

Association between smoking and psychosis may be mediated by maternal smoking during pregnancy

David Curtis

d.curtis@ucl.ac.uk

UCL Genetics Institute, UCL, Darwin Building, Gower Street London WC1E 6BT.
Centre for Psychiatry, Barts and the London School of Medicine and Dentistry.

Financial support: none

Word count: 267

The report demonstrating a cross-sectional association between current smoking and psychotic experiences (Bhavsar *et al.* 2017) is interesting but the authors fail to consider what I believe may be an important and plausible potential causal mechanism. Smoking during pregnancy has been shown to be a risk factor for the subsequent development of schizophrenia (Niemel *et al.* 2016). Since similar effects are seen with maternal malnutrition and infections during pregnancy (Xu *et al.* 2009; Khandaker *et al.* 2013), it seems reasonable to conclude that a broad range of insults during pregnancy, including maternal smoking, may impact on neurodevelopment in a way which increases susceptibility to subsequent psychosis. Smoking behaviour is familial, with both genetic and cultural effects (Bares *et al.* 2016). Thus, research subjects who currently smoke are more likely to have mothers who smoked during pregnancy and this would be expected to increase their risk of schizophrenia and/or psychotic experiences. This mechanism might also explain, at least in part, the apparent genetic association between smoking behaviour and schizophrenia (Hartz *et al.* 2017). The effect might be mediated indirectly through maternal smoking behaviour, which would lead to the observation that genetic variants associated with smoking (in the mother) were also associated with schizophrenia (in the offspring).

Of course, the mechanism I propose does not explain the observed association between smoking and schizophrenia in discordant monozygotic twins (Lyons *et al.* 2002), nor the correlation in individuals between smoking behaviour and psychotic experiences over time (Taylor *et al.* 2014). Nevertheless the effect of maternal smoking does seem to be potentially relevant and worthy of consideration in studies of this nature.

Conflict of interest

The author declares he has no conflict of interest.

References

Bares CB, Kendler KS, Maes HHM (2016). Racial differences in heritability of cigarette smoking in adolescents and young adults. *Drug and Alcohol Dependence* **166**, 75–84.

Bhavsar V, Jauhar S, Murray RM, Hotopf M, Hatch SL, McNeill A, Boydell J, MacCabe JH (2017). Tobacco smoking is associated with psychotic experiences in the general population of South London. *Psychological Medicine*, 1–9.

Hartz SM, Horton AC, Hancock DB, Baker TB, Caporaso NE, Chen L-S, Hokanson JE, Lutz SM,

Marazita ML, McNeil DW, Pato CN, Pato MT, Johnson EO, Bierut LJ (2017). Genetic correlation between smoking behaviors and schizophrenia. *Schizophrenia Research*

Khandaker GM, Zimbron J, Lewis G, Jones PB (2013). Prenatal maternal infection, neurodevelopment and adult schizophrenia: a systematic review of population-based studies. *Psychological Medicine* **43**, 239–257.

Lyons MJ, Bar JL, Kremen WS, Toomey R, Eisen SA, Goldberg J, Faraone S V, Tsuang M (2002). Nicotine and familial vulnerability to schizophrenia: a discordant twin study. *Journal of abnormal psychology* **111**, 687–93.

Niemel S, Sourander A, Surcel H-M, Hinkka-Yli-Salomäki S, McKeague IW, Cheslack-Postava K, Brown AS (2016). Prenatal Nicotine Exposure and Risk of Schizophrenia Among Offspring in a National Birth Cohort. *American Journal of Psychiatry* **173**, 799–806.

Taylor G, McNeill A, Girling A, Farley A, Lindson-Hawley N, Aveyard P (2014). Change in mental health after smoking cessation: systematic review and meta-analysis. *BMJ (Clinical research ed.)* **348**, g1151.

Xu M-Q, Sun W-S, Liu B-X, Feng G-Y, Yu L, Yang L, He G, Sham P, Susser E, St. Clair D, He L (2009). Prenatal Malnutrition and Adult Schizophrenia: Further Evidence From the 1959-1961 Chinese Famine. *Schizophrenia Bulletin* **35**, 568–576.