

Accepted Manuscript

A randomised controlled trial of guided internet-based cognitive behavioural therapy for perfectionism: Effects on psychopathology and transdiagnostic processes.

Radha Kothari, Chris Barker, Nancy Pistrang, Alexander Rozental, Sarah Egan, Tracey Wade, Hannah Allcott-Watson, Gerhard Andersson, Roz Shafran



PII: S0005-7916(18)30055-7

DOI: <https://doi.org/10.1016/j.jbtep.2019.03.007>

Reference: BTEP 1472

To appear in: *Journal of Behavior Therapy and Experimental Psychiatry*

Received Date: 28 February 2018

Revised Date: 19 March 2019

Accepted Date: 26 March 2019

Please cite this article as: Kothari, R., Barker, C., Pistrang, N., Rozental, A., Egan, S., Wade, T., Allcott-Watson, H., Andersson, G., Shafran, R., A randomised controlled trial of guided internet-based cognitive behavioural therapy for perfectionism: Effects on psychopathology and transdiagnostic processes., *Journal of Behavior Therapy and Experimental Psychiatry* (2019), doi: <https://doi.org/10.1016/j.jbtep.2019.03.007>.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

A Randomised Controlled Trial of Guided Internet-Based Cognitive Behavioural Therapy for Perfectionism: Effects on psychopathology and transdiagnostic processes.

Radha Kothari^{a, b}, Chris Barker^a, Nancy Pistrang^a, Alexander Rozental^{b, c}, Sarah Egan^d, Tracey Wade^e, Hannah Allcott-Watson^b, Gerhard Andersson^{f, g}, Roz Shafran^b.

^a Division of Psychology and Language Sciences, University College London, London, 1 - 19 Torrington Place, London, WC1E 7HB, United Kingdom

^b Institute of Child Health Population, Policy and Practice Programme, University College London, 30 Guilford Street, London, WC1N 1EH, United Kingdom

^c Department of Clinical Neuroscience, Karolinska Institutet, Norra Stationsgatan 69, 113 64, Stockholm, Sweden

^d School of Psychology, Faculty of Health Sciences, Curtin University, Kent St, Bentley, WA 6102, Perth, Australia

^e School of Psychology, Flinders University, Adelaide, Sturt Rd, Bedford Park, SA 5042, Australia

^f Department of Behavioural Sciences and Learning, Linköping University, 581 83 Linköping, Sweden

^g Department of Clinical Neuroscience, Karolinska Institute, Solnavägen 1, 171 77 Solna, Stockholm, Sweden

Corresponding author: Radha Kothari, Radha.kothari.10@ucl.ac.uk

c.barker@ucl.ac.uk

n.pistrang@ucl.ac.uk

alexander.rozentel@ki.se

s.egan@curtin.edu.au

tracey.wade@flinders.edu.au

hannah.allcott-watson.14@ucl.ac.uk

gerhard.andersson@liu.se

r.shafran@ucl.ac.uk

**A Randomised Controlled Trial of Guided Internet-Based Cognitive Behavioural
Therapy for Perfectionism: Effects on psychopathology and transdiagnostic
processes.**

ACCEPTED MANUSCRIPT

Abstract

Background and Objectives: Perfectionism is a transdiagnostic process that has been associated with a range of psychopathology and also with other transdiagnostic processes. We have previously shown that guided internet-based cognitive behavioural therapy (ICBT) can reduce symptoms of dysfunctional perfectionism, however, no impact was observed on symptoms of depression and anxiety. Here we explore the impact of guided ICBT for perfectionism on symptoms of other associated psychopathology, specifically obsessive-compulsive disorder (OCD) and eating disorders, and also on other associated transdiagnostic processes (self-esteem, intolerance of uncertainty, and self-compassion).

Methods: Participants who presented with clinical levels of perfectionism were randomised to an experimental group that received the intervention ($n = 62$), or a wait list control group ($n = 58$). Questionnaires assessing symptoms of OCD, eating disorders, self-esteem, intolerance of uncertainty, and fear of self-compassion were completed pre-intervention, post-intervention (12 weeks), and at follow-up (24 weeks). Between group effect sizes are reported.

Results: The intervention led to significant decreases in symptoms of OCD ($d = -0.9$; CI: -1.4, -0.4) and eating disorders ($d = -0.6$; CI: -1.0, -0.1), and had an impact on other transdiagnostic processes resulting in increased self-esteem ($d = 0.7$; CI: 0.2, 1.2), decreases in intolerance of uncertainty ($d = -0.9$; CI: -1.4, -0.4), and fear of self-compassion ($d = -0.8$; CI: -1.3, -0.3). At follow-up changes were maintained in symptoms of OCD ($d = -1.3$; CI: -1.8, -0.8), disordered eating ($d = -0.7$; CI: -1.2, -0.2), intolerance of uncertainty ($d = -0.8$; CI: -1.2, -0.3), and fear of self-compassion ($d = -1.0$; CI: -1.5, -0.5).

Conclusions: Guided ICBT for perfectionism improves associated psychopathology and transdiagnostic processes. *ClinicalTrials.gov registration no. NCT02756871.*

Keywords: perfectionism, cognitive behavioural therapy, online intervention, guided self-help, transdiagnostic, randomised controlled trial, self-esteem; intolerance of uncertainty, self-compassion, obsessive-compulsive disorder, eating disorder.

Highlights

- Internet based cognitive behavioural therapy for perfectionism led to significant decreases in symptoms of associated psychopathology, specifically obsessive-compulsive disorder and eating disorders.
- Internet based cognitive behavioural therapy for perfectionism had an impact on other associated transdiagnostic processes, specifically increased self-esteem, and decreases in intolerance of uncertainty and fear of self-compassion.
- Changes in symptoms of obsessive-compulsive disorder and eating disorders, and in intolerance of uncertainty and fear of self-compassion were maintained at 24-week follow-up.

1. Introduction

Perfectionism has, for some time now, been considered to present in both adaptive and maladaptive forms, and the numerous theoretical accounts of perfectionism tend to agree that it comprises of two factors: one, having high standards; and two, engaging in intense self-criticism in response to these standards not being met (Burns, 1980; Frost, Marten, Lahart, & Rosenblate, 1990; Hamachek, 1978; Hewitt & Flett, 1991; Shafran, Cooper, & Fairburn, 2002; Slaney, Rice, & Ashby, 2002). This distinction is supported by factor analytic studies of the most commonly used measures of perfectionism which have resulted in a two-factor solution: perfectionistic strivings and perfectionistic concerns, respectively (Stoeber & Otto, 2006).

Both dimensions of perfectionism have been associated with generalised anxiety disorder (GAD), depression, eating disorders, obsessive compulsive disorder (OCD), and panic disorder; however, a recent meta-analysis found that most psychopathology was more strongly associated with perfectionistic concerns, except for eating disorders which were found to be strongly associated with both (Limburg, Watson, Hagger, & Egan, 2016; Smith et al., 2017). Perfectionism has been identified as a transdiagnostic process, meaning that it increases risk for, and contributes to maintenance of, a range of psychopathology and mental health disorders (Coughtrey, Shafran, Bennett, Kothari, & Wade, 2017; Egan, Wade, & Shafran, 2011; Radomsky, Rachman, Shafran, Coughtrey, & Barber, 2014; Shafran & Mansell, 2001; Stoeber & Otto, 2006). Maladaptive perfectionism can also be clinically problematic in its own right, causing both practical and emotional difficulties, and commonly resulting in a narrowing of interests and social isolation (Shafran, Coughtrey, & Kothari, 2016).

Cognitive behavioural therapy (CBT) for perfectionism is efficacious in reducing symptoms of perfectionism in individual, group and guided self-help formats (Egan, van Noort, et al., 2014; Lloyd, Schmidt, Khondoker, & Tchanturia, 2015), and in reducing symptoms of anxiety, depression, and eating disorders (Egan & Hine, 2008; Egan, van Noort, et al., 2014; Glover, Brown, Fairburn, & Shafran, 2007; Lloyd et al., 2015; Riley, Lee,

Cooper, Fairburn, & Shafran, 2007; Steele & Wade, 2008; Steele et al., 2013). More recently, such benefits have also been found with internet-based CBT (ICBT) interventions for perfectionism (Egan, van Noort, et al., 2014; Rozental et al., 2017; Shafran et al., 2017). Internet-based interventions, where patients work with or without support, are able to fill the gap between need and availability and have been associated with a number of advantages such as increased convenience, patient anonymity, lower cost to health care providers, and improved mental well-being (Andersson & Cuijpers, 2009; Cuijpers, Van Straten, & Andersson, 2008; Lauder, Chester, & Berk, 2007; Mitchell, Stanimirovic, Klein, & Vella-Brodrick, 2009; Musiat, Goldstone, & Tarrrier, 2014; Powell et al., 2013; Wagner, Horn, & Maercker).

In a recent publication we provided evidence that guided ICBT is effective at reducing symptoms of maladaptive perfectionism, as measured by the concern over mistakes subscale of the Frost Multidimensional Perfectionism Scale (FMPS; Frost et al., 1990) and the Clinical Perfectionism Questionnaire (CPQ; Fairburn et al., 2003). No impact was found on symptoms of depression and anxiety however. In this paper we explore the impact of this guided ICBT intervention on other forms of psychopathology that have been associated with perfectionism, specifically symptoms of OCD and eating disorders. In addition to this, and based on previous evidence that perfectionism is associated with other constructs also hypothesised to be transdiagnostic in nature, we also explore in this paper the impact of guided ICBT for perfectionism on self-esteem, intolerance of uncertainty, and self-compassion. We have previously reported on correlations between variables in this sample which show that perfectionism, as measured by the CPQ and subscales of the FMPS, is negatively associated with self-esteem and positively associated with intolerance of uncertainty and fear of self-compassion with small to medium effect sizes (Coughtrey et al., 2017; Kothari, 2017).

Self-esteem, defined as an individual's subjective appraisal of the self at an affective and evaluative level (Rosenberg, 1965), has been associated with risk for and maintenance of a number of mental health disorders and may be negatively associated with perfectionism

(Coopersmith, 1967; Fairburn, Cooper, & Shafran, 2003; Kothari, 2017; Krabbendam et al., 2002; Orth, Robins, Trzesniewski, Maes, & Schmitt, 2009; Rosenberg, 1965). It has been hypothesised that extreme perfectionism may result in comparison between the real and ideal self and between perfectionistic goals and actual performance, resulting in low self-esteem; or that low self-esteem may result from individuals with clinical perfectionism basing their self-worth on their ability to achieve unattainable standards and being highly self-critical when standards are not met (Beck, 1976; Burns & Beck, 1978).

Intolerance of uncertainty, initially defined as a negative and fearful style of responding at a cognitive, emotional, and behavioural level to uncertain situations (Freeston, Rhéaume, Letarte, Dugas, & Ladouceur, 1994), is also hypothesised to be a transdiagnostic process, contributing to the development and maintenance of most anxiety disorders, depression, and OCD (Gentes & Ruscio, 2011; Mahoney & McEvoy, 2012). Intolerance of uncertainty has been defined as having two factors: desiring and actively seeking predictability and certainty, and being cognitively and behaviourally paralysed in the face of uncertainty (Birrell, Meares, Wilkinson, & Freeston, 2011). The Obsessive Compulsive Cognitions Working Group (OCCWG; 2003) have suggested that among individuals with OCD perfectionism acts in conjunction with intolerance of uncertainty and that the need to achieve perfection is an attempt to make the future more certain, particularly in domains that are experienced as uncertain or distressing. Empirical research supports the theorised relationship between perfectionism and intolerance of uncertainty, with most studies finding a positive correlation between the two constructs in the general population, and among eating disordered and socially anxious samples (Kothari, 2017).

Self-compassion, the ability to be kind, empathic, gentle, and warm towards oneself, even in times of difficulty, may also be a transdiagnostic process, though more research is required to show this empirically (Gilbert, McEwan, Matos, & Rivis, 2011). Self-compassion is negatively associated with maladaptive facets of perfectionism which are representative of self-criticism, key to theoretical accounts of perfectionism (Frost et al., 1990; Neff, 2003; Shafran et al., 2002). Fear of self-compassion, theorised to occur when compassion is

associated with negative emotions such as grief or threat, is positively associated with self-criticism (Gilbert & Procter, 2006). Given the growing evidence of a negative association between self-compassion and perfectionism, it is possible to hypothesise that fear of self-compassion may be directly positively associated with perfectionism (Egan, Wade, Shafran, & Antony, 2014). Compassion towards others and the self requires an understanding that imperfection and the making of mistakes is a normal part of the human condition, and individuals high in perfectionism may fear that self-compassion might reduce motivation to meet high standards without making errors (Gilbert, 2009; Gilbert et al., 2011).

Though there is growing evidence for relationships between perfectionism and self-esteem, intolerance of uncertainty, and self-compassion, the directions of causality, and indeed whether associations are direct, have not yet been determined.

Aims

We have previously shown that guided ICBT for perfectionism results in significant decreases in symptoms of perfectionism, both post-intervention and at six month follow-up, with medium to large effect sizes (Kothari, 2017; Shafran et al., 2017). In contrast to our hypotheses, no impact was observed on symptoms of depression or anxiety. Here we explore the impact of the intervention on symptoms of other psychopathology associated with perfectionism, specifically OCD and eating disorders. We also explore the impact on other transdiagnostic processes, associated with and theoretically relevant to the maintenance of perfectionism: self-esteem, intolerance of uncertainty, and self-compassion.

Hypotheses

Theoretical models indicate that perfectionism contributes to the development and/or maintenance of OCD and eating disorders. Experimental evidence supports this theoretical perspective, and interventions targeting perfectionism appear to result in a reduction in symptoms of both disorders. Based on this we hypothesised that this guided ICBT

intervention for perfectionism would also result in reduced symptoms of OCD and eating disorders (Egan et al., 2011; Handley, Egan, Kane, & Rees, 2015; Obsessive Compulsive Cognitions Working, 1997, 2003; Pleva & Wade, 2007; Shafran et al., 2002; Steele & Wade, 2008).

Based upon current evidence of the relationship between perfectionism and other transdiagnostic processes reviewed above, it was hypothesised that guided ICBT for perfectionism would also result in increased self-esteem and decreases in intolerance of uncertainty and fear of self-compassion.

2. Method

2.1 Protocol, ethics and trial registration

We report here on secondary findings from a randomised controlled trial of a guided ICBT intervention for clinical perfectionism called Overcoming Perfectionism. Ethical approval was granted by the University College London (UCL) Research Ethics Committee (Project ID: 6222:001). The full protocol has previously been published (Kothari, Egan, Wade, Andersson, & Shafran, 2016) and the study was registered as a clinical trial on ClinicalTrials.gov (NCT02756871).

2.2 Setting and Intervention

This version of the treatment was adapted from the Cognitive Behavioural Treatment of Perfectionism (Shafran, Egan, & Wade, 2010). For the internet-based version the content was made briefer, video was used, and worksheets were adapted to be interactive. The intervention was divided into eight modules designed to be completed weekly (Figure 1); however, participants were provided with guidance and support over 12 weeks to allow for breaks such as illness and holidays. Participants continued to have access to the intervention after 12 weeks, but ongoing guidance was not provided. Post-intervention measures were collected 12 weeks after participants were randomly allocated to the

experimental or control group (T_2) and follow-up measures were collected 24 weeks after (T_3). As a token of thanks participants were given a ten pound voucher after completing measures at T_2 and another when completing measures at T_3 .

Psychoeducation and examples were provided in each module, followed by an interactive section requiring participants to answer questions and complete worksheets. In this way participants were able to create an idiosyncratic model (or formulation) of their own unhelpful perfectionism, challenge and restructure unhelpful cognitions, and also design surveys and behavioural experiments.

2.3 Guidance and Feedback

Guidance and feedback were provided by 12 guides who were psychology undergraduates, Masters students, PhD students/graduates, or trainee clinical psychologists. Each participant was allocated a guide who was able to view submitted worksheets and responses. Guides provided feedback and suggestions to the participant in the form of asynchronous internet-based written communication. Participants received feedback and guidance as they completed each module and submitted the relevant worksheets. The average length of feedback for each worksheet was one to two paragraphs. Participants were also able to ask questions and respond to feedback and they received guidance if they specifically requested help with understanding or completing modules and between-session work. Guides were provided with regular supervision by a clinical psychologist and research psychologist to discuss guidance and ensure consistency.

2.4 Participants

An a priori power calculation was conducted using a tool designed by Hedeker and colleagues which is appropriate for determining power for longitudinal designs (Hedeker, Gibbons, & Waternaux, 1999). A two-tailed alpha of .05, three assessment points (pre, post and follow-up), a pre-post correlation for the primary outcome measure (concern over mistakes subscale, see below) of 0.61 and attrition rates of 50%, were used. Both the pre-

post correlation and expected attrition rate were based upon a similar RCT of a web-based intervention for perfectionism (Egan, van Noort, et al., 2014). A sample size of 40 participants per group, with 20 participants completing per group, was found to provide 80% power at a two-tailed $p < .05$ to detect a large effect size (0.80) difference between the control and intervention groups. This use of a large effect size was also based upon the previous RCT (Egan, van Noort, et al., 2014).

Full details of participant recruitment and consent have previously been published (Kothari, 2017; Shafran et al., 2017). Participants completed questionnaires online which included screening measures to determine their eligibility. To be eligible for inclusion participants had to be 18 or over, with no upper age limit, score one standard deviation above published norms on the 'concern over mistakes' subscale of the Frost Multidimensional Perfectionism Scale (Frost et al., 1990), a score of ≥ 29 (Suddarth & Slaney, 2001), and be fluent in English. Participants were excluded if they reported suicidal thoughts or intent, either current or in the past, at any point over the duration of the intervention. Due to the established co-occurrence between a range of psychopathology and perfectionism, participants reporting symptoms of co-occurring mental health disorders (depression, anxiety, OCD and eating disorders) were not excluded from the study. If eligible, participants were randomly allocated to the experimental group to complete the intervention, or the control group (no intervention). Randomisation of participants was performed by a third party, unconnected to the study, who created a randomization schedule using a Web-based randomizer ("Sealed Envelope,").

A total of 156 participants registered for participation and completed the screening measures, of whom 35 (22.4%) participants were excluded as they did not meet the inclusion criteria of ≥ 29 on the FMPS concern over mistakes subscale (Frost et al., 1990) and one participant refused to be randomised, resulting in a total of 120 participants that were randomised into the experimental ($n = 62$) and control ($n = 58$) groups (Figure 2). Participants allocated to the experimental group were paired with guides after randomisation. Participants who did not meet criteria for inclusion in the study were sent a copy of

Overcoming Perfectionism: A Self-Help Guide Using Cognitive Behavioural Techniques

(Shafran et al., 2010) and were signposted to other services.

2.5 Measures

Participants completed self-report questionnaires at three time points: (i) prior to any intervention at baseline (T_1), (ii) 12 weeks after the participant was randomised to the experimental or control group to assess change post-intervention (T_2), and (iii) 24 weeks after the participant was randomised to assess whether change was maintained at follow-up (T_3). Findings pertaining to the effect of the intervention on symptoms of perfectionism, depression and anxiety have previously been published (Shafran et al., 2017). This study reports on a different set of measures to Shafran et al. (2017) which assess symptoms of other psychopathology (specifically symptoms of OCD and eating disorders) and other transdiagnostic processes (self-esteem, intolerance of uncertainty, and self-compassion).

Measures of Psychopathology

The Obsessive-Compulsive Inventory – Revised (OCI-R; Foa et al., 2002), a shortened version of the Obsessive-Compulsive Inventory, assesses symptoms of OCD. The measure consists of 18 items (e.g. “I frequently have nasty thoughts and have difficulty in getting rid of them”). On a five-point scale, respondents rate how distressed or bothered they have been in the past month by the symptom described, with responses ranging from “Not at all” to “Extremely.” It has been found to have good validity and reliability in both clinical and non-clinical samples (Hajcak, Huppert, Simons, & Foa, 2004; Huppert et al., 2007). Cronbach’s alpha in the current study was 0.92.

The Eating Disorder Examination-Questionnaire (EDE-Q; Fairburn, 1994) assesses for symptoms of the eating disorders: Anorexia Nervosa (AN), Bulimia Nervosa (BN), Binge Eating Disorder (BED) and sub-threshold variants. The measure consists of 28 items rated on a seven point scale appropriate to the item: number of days a symptom has been experienced over the past month, ranging from no days to every day; and for remaining

questions (e.g. “How dissatisfied have you been with your weight?”) responses ranging from “Not at all” to “Markedly.” It has been found to have good reliability and has been validated for use among clinical and community samples (Aardoom, Dingemans, Op’t Landt, & Van Furth, 2012; Reas, Grilo, & Masheb, 2006). Cronbach’s alpha in the current study was 0.92.

Measures of Transdiagnostic Processes

The Rosenberg Self-esteem Scale (RSES; Morris Rosenberg, 1965) is a self-report measure consisting of 10 items (e.g. “On the whole I am satisfied with myself” and “I wish I could have more respect for myself”) and is rated on a four point scale ranging from 1 = “strongly disagree” to 4 = “strongly agree”. It has been found to be reliable and has been validated for use among clinical and community samples (Bagley & Mallick, 2001; Schmitt & Allik, 2005). Cronbach’s alpha in the present study was .87.

The Intolerance of Uncertainty Scale (IUS; Freeston et al., 1994) is a self-report measure consisting of 27 items (e.g. “Uncertainty stops me from having a firm opinion” and “It’s unfair not having any guarantees in life”) and is rated on a five point scale ranging from 1 = “Not at all characteristic of me” to 5 = “Entirely characteristic of me.” It has been found to be reliable and has been validated for use among clinical and community samples (Buhr & Dugas, 2006; Jacoby, Fabricant, Leonard, Riemann, & Abramowitz, 2013). The scale was highly reliable in the present study, Cronbach’s alpha .94.

The Fear of Compassion Scales (FCS; Gilbert et al., 2011) consist of three scales which assess fear of compassion for self (e.g. “I worry that if I start to develop compassion for myself I will become dependent on it”), fear of compassion from others (e.g. “I try to keep my distance from others, even if I know they are kind”) and fear of compassion for others (e.g. “Being too compassionate makes people soft and easy to take advantage of”). The FCS is made up of 38 items in total which are rated on a four-point scale ranging from “Don’t agree at all” to “Completely agree.” It has been shown to be reliable and valid for use among clinical and community samples (Gilbert et al., 2012; Gilbert et al., 2011). In the current study

only the first scale was used (fear of compassion for the self), for which Cronbach's alpha was 0.95.

2.6 Statistical Analysis

Logistic regression analyses were conducted to determine whether demographic and baseline characteristics were predictive of missing data at T_2 and T_3 . Completer and intention-to-treat (ITT) analyses were conducted using SPSS version 24. Multiple imputation was used to manage missing data for ITT analysis as this has been shown to be superior to last observation carried forward (Elobeid et al., 2009). Demographic data and outcome data from all three time points were included in the multiple imputation model and ten datasets were imputed.

To compare change over time between experimental and control groups ANCOVA analyses were conducted using the residualised gains procedure. Observations at T_2 (post-intervention and primary endpoint, 12 weeks after randomisation) were adjusted for observations at T_1 (baseline, pre-intervention). The follow-up effect of the intervention was investigated in the same way, using observations at T_3 (follow-up, 24 weeks after randomisation) as the outcome variable, adjusted for observations at T_1 . In line with recent recommendations regarding the reporting of the results of statistical analysis we have adopted an open science approach by reporting Cohen's d between group effect sizes and 95% confidence intervals to indicate significance rather than providing significance or p values (Cumming, 2013; Cumming & Calin-Jageman, 2016). Given our inclusion of baseline observations as a covariate, we interpret a significant difference between groups as being an effect size that does not cross zero. Criteria for reliable change were used to determine whether the difference on all outcomes was reliable and due to the intervention, rather than due to measurement error. A Reliable Change Index (RCI) was computed using the formula $SE_{diff} = SD_1 \sqrt{2\sqrt{1-r}}$, where SD_1 is the standard deviation at baseline and r is the Cronbach's alpha reliability coefficient of the measure (Jacobson & Truax, 1991). Change scores are required to exceed 1.96 times the SE_{diff} to show improvement (Evans, Margison,

& Barkham, 1998). A negative change score exceeding the RCI was used to determine deterioration (Rozental et al., 2014).

3. Results

3.1 Descriptive Statistics

The mean age of the 120 participants was 28.9 years (SD = 8), the majority were female (n = 98; 82%) and just over half were studying for a University degree (n = 62; 52%). A notable proportion of participants had previously received (n = 47; 40%) or were currently receiving (n = 34; 28%) treatment for a mental health disorder (Table 1).

3.2 Missing Data Analysis

At T₂ (post-intervention at 12 weeks) data was available on 71 participants (experimental = 30, control = 41) and at T₃ (follow-up at 24 weeks) data was available on 66 participants (experimental = 27, control = 39). The flow of participants through the trial is shown in Figure 2. Logistic regression analysis conducted to identify predictors of missing data showed that having an educational qualification certificate was associated with lower odds of having missing data at T₂, while being in the experimental group was associated with having higher odds of having missing data at T₂ and T₃. Additionally, higher scores on the EDE-Q were associated with having higher odds of having missing data at T₂ and higher scores on the OCI-R were associated with having higher odds of missing data at T₃ (Table 1).

3.3 Completion of Modules and Measures

Of participants in the experimental group, 17 (27.4%) completed no modules, 36 (58.1%) completed one to four modules (half or less) and nine (14.5%) completed five to eight modules. The mean number of modules completed was 2.48 (SD = 2.37).

3.4 Change in Psychopathology

Completer and ITT analysis showed significant group differences on the OCI-R post intervention (T_2) and at follow-up (T_3), and participants in the experimental group had significantly higher odds than those in the control group of meeting criteria for reliable change in the expected direction (Tables 2 and 3). Completer but not ITT analysis also revealed significant between group differences on the EDE-Q at T_2 and T_3 ; however, participants in the experimental group did not have significantly higher odds of meeting criteria for reliable change.

3.5 Change in other Transdiagnostic Processes

Completer and ITT analysis revealed significant between group differences on the IUS and the FCS at T_2 and T_3 , and on the RSES at T_2 but not T_3 . Participants in the experimental group also had significantly higher odds than those in the control group of meeting criteria for reliable change on the FCS at T_2 .

4. Discussion

We have previously shown that guided ICBT for perfectionism can be effective at reducing perfectionism as measured by the clinical perfectionism questionnaire and the concern over mistakes subscale of the Frost Multidimensional Perfectionism Scale (Frost et al., 1990), post intervention and at follow-up. Here we show that the same intervention reduced symptoms of other psychopathology, specifically OCD and eating disorders, and also had an impact on other transdiagnostic processes associated with perfectionism, with participants showing decreased intolerance of uncertainty and fear of compassion post intervention and at follow-up. Increased self-esteem was also observed post intervention, but not at follow-up.

Our findings contribute to the literature suggesting that that an intervention aimed at decreasing perfectionism can also lead to a reduction in associated psychopathology, specifically symptoms of OCD and eating disorders (Handley et al., 2015; Pleva & Wade,

2007; Steele & Wade, 2008). In comparison to the control group, participants in the experimental group showed decreases in symptoms of OCD with medium to large effect sizes, post-intervention and at follow-up. Between group differences in symptoms of OCD were statistically significant when conducting completer and intent to treat analysis, and participants in the experimental group had over twice the odds of meeting criteria for reliable change in the hypothesised direction (i.e. decreased scores on the OCI-R). Between group effect sizes in ED symptoms were also medium to large, both post-intervention and at follow-up, but differences only met criteria for statistical significance when analysing data from completers so this finding must be interpreted with caution. Additionally, participants in the experimental group were not significantly more likely to meet criteria for reliable change; however, scores on the EDE-Q for the current sample were low to begin with, meaning only limited change would be expected.

A novel finding is that guided ICBT for perfectionism may impact on other associated transdiagnostic processes. The experimental group showed comparative increases in self-esteem with medium to large effect sizes post-intervention which were statistically significant when conducting completer and intent to treat analysis. This improvement was not maintained at follow-up, however. One could hypothesise that engagement with treatment generally, rather than treatment for perfectionism specifically, is what led to an increase in self-esteem, potentially explaining why self-esteem scores decreased again once engagement ceased. This might be an interesting line of enquiry for future research, particularly if it was possible to maintain this increase.

In comparison to the control group, participants in the experimental group showed a decrease in intolerance of uncertainty and in fear of self-compassion, post-intervention and at follow-up with medium to large effect sizes. Between group differences were statistically significant for both completer and intent to treat analysis. It is possible that the impact on intolerance of uncertainty and fear of self-compassion is due to aspects of the intervention acting directly on these processes. While modules one to four focus specifically on understanding and challenging perfectionistic thoughts and beliefs, modules five to eight are

broader. For example, modules seven and eight target self-worth and self-compassion, meaning a potentially direct impact on fear of self-compassion. Module five also teaches problem solving skills, which may increase an individual's confidence in their ability to deal with uncertain situations and unexpected events, potentially leading to a decrease in intolerance of uncertainty. It is worth bearing in mind, however, that fewer than 15% of participants completed more than four modules, meaning that the majority of participants only completed modules that focus specifically on perfectionism. Given this, the findings might be indication of causal relationships between perfectionism and intolerance of uncertainty, and perfectionism and self-compassion. Results of completer analysis may also be taken in support of this as effect sizes were found to be larger for intolerance of uncertainty and fear of self-compassion amongst completers; however, it is not possible to determine causality from the design of the current study.

To our knowledge this is the first study to show that CBT for perfectionism can lead to improvement in other associated transdiagnostic processes, specifically intolerance of uncertainty and self-compassion. Although the odds of meeting criteria for reliable change in self-esteem and intolerance of uncertainty were not significantly higher for participants in the experimental group, the current sample were not observed to have extreme scores on these processes, meaning only small changes might be expected. Given this, more research is warranted and should also investigate whether any causal relationships are direct or are due to other unknown variables.

Research investigating the causal relationship between perfectionism and OCD, and perfectionism and the transdiagnostic processes of intolerance of uncertainty and self-compassion, may be particularly fruitful given the strong evidence that guided ICBT for perfectionism can impact on these. It was not possible here to investigate whether change in perfectionism was a potential mediator for changes in psychopathology and other associated transdiagnostic processes due to limitations inherent in the design of this study, and limited evidence regarding the direction of causality between variables. Future studies would benefit from including multiple time-points of measurement during the period of treatment so that

change can be tracked over time and analysis of potential mediators of change is possible. This would also allow exploration of causal direction, i.e. does change in perfectionism mediate change in associated psychopathology and other transdiagnostic processes, or vice versa.

This research derives many of its strengths from its design. A randomised controlled trial allows for causal inferences to be made, which means that changes in psychopathology and transdiagnostic processes can be attributed (either directly or indirectly) to the guided ICBT intervention. Because participants were randomly allocated, differences between groups and potential confounding factors were minimised. This was confirmed in an exploratory analysis which showed no significant differences between experimental and control groups in demographics or baseline scores of outcome variables. In addition, the randomisation schedule was created by a third party uninvolved with the research, reducing the likelihood of randomisation bias.

Considerable drop-out was observed, with over a quarter of participants in the experimental group completing no modules after being allocated and fewer than 15% completing more than half the intervention. This is a limitation common to studies of online interventions. A systematic review found that drop-out for online psychological interventions ranged from 2% to 83% and average drop-out was 35% (Melville, Casey, & Kavanagh, 2010). The review also found that the majority of individuals drop out before commencing treatment, which is in line with the current findings. It has been suggested that reminders might prompt participants to persevere (Donkin & Glozier, 2012). In the current research guides were directed to prompt participants who were inactive for a week or more. Though drop-out was still high, it was not as high as a previous RCT evaluating ICBT for perfectionism without guidance, so it is possible that the addition of guidance is effective in increasing retention (Egan, van Noort, et al., 2014). It should also be noted that the pattern of missing data was somewhat impacted by education, and symptoms of disordered eating and obsessionality, which may challenge some assumptions of the statistical testing used. An RCT evaluating the effectiveness of the current intervention was run in Sweden at the

same time as the current RCT and much lower drop-out rates were observed (approximately 80% retention; Rozental et al., 2017). It is likely that important procedural differences in the design of the trials might have contributed to this difference in drop-out. In particular, participants in the Swedish trial were contacted via telephone by their therapists prior to commencing the intervention to confirm their intention to participate. It is likely that this meeting not only boosted retention through checking that participants still wanted to participate after being randomised, but also that it laid a better foundation for the building of a good therapeutic relationship, which in turn may have increased motivation to participate (Shafran et al., 2017).

Finally, multiple comparisons are subject to statistical analysis and though this is typical in the reporting of randomised controlled trials, the risk of experiment-wise error rate must be considered. We have refrained from adjusting for multiple comparisons as such an approach assumes that all null hypotheses are true simultaneously and can therefore increase the likelihood of type II errors, meaning truly important differences are deemed non-significant. Instead we have taken the approach of clearly describing the statistical tests that have been performed (Perneger, 1998).

5. Conclusions

The current study contributes to existing literature showing that treating perfectionism can have a positive impact on symptoms of associated psychopathology, specifically OCD and eating disorders. It also provides preliminary evidence for the hypothesis that effecting change in one transdiagnostic process can lead to change in other associated transdiagnostic processes. Exploring the nature of the relationships observed between these transdiagnostic processes and how they interact within different mental health disorders could provide new insights into how to identify those at high risk and the best and most effective method of treating these disorders. With further understanding it is possible that CBT and ICBT interventions focussing on a cluster of transdiagnostic processes could

become integral to the treatment of mental health disorders and provide an effective early intervention strategy.

Acknowledgements

We would like to thank Julia Bowen, Tuhina Bhattacharyya, Harriet Clarkson, Sophie Cripps, Jovita Leung, Lauren Robinson, Jean Stafford, Martha von Werthern and Chloe Shu. All research at Great Ormond Street Hospital NHS Foundation Trust and UCL Great Ormond Street Institute of Child Health is made possible by the NIHR Great Ormond Street Hospital Biomedical Research Centre. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

Conflicts of Interest

None declared.

References

- Aardoom, J. J., Dingemans, A. E., Op't Landt, M. C. S., & Van Furth, E. F. (2012). Norms and discriminative validity of the Eating Disorder Examination Questionnaire (EDE-Q). *Eating behaviors*, *13*(4), 305-309.
- Andersson, G., & Cuijpers, P. (2009). Internet-based and other computerized psychological treatments for adult depression: a meta-analysis. *Cognitive Behaviour Therapy*, *38*(4), 196-205.
- Bagley, C., & Mallick, K. (2001). Normative data and mental health construct validity for the Rosenberg Self-Esteem Scale in British Adolescents. *International Journal of Adolescence and Youth*, *9*(2-3), 117-126.
- Beck, A. T. (1976). *Cognitive Therapy and the Emotional Disorders*. New York: International Universities Press.
- Birrell, J., Meares, K., Wilkinson, A., & Freeston, M. (2011). Toward a definition of intolerance of uncertainty: A review of factor analytical studies of the Intolerance of Uncertainty Scale. *Clinical psychology review*, *31*(7), 1198-1208.
- Buhr, K., & Dugas, M. J. (2006). Investigating the construct validity of intolerance of uncertainty and its unique relationship with worry. *Journal of anxiety disorders*, *20*(2), 222-236. doi:https://doi.org/10.1016/j.janxdis.2004.12.004
- Burns, D. D. (1980). The perfectionist's script for self-defeat. *Psychology today*, *14*(6), 34-52.
- Burns, D. D., & Beck, A. T. (1978). Cognitive behaviour modification of mood disorders. In E. Foreyt & D. P. Rathjen (Eds.), *Cognitive Behaviour Therapy: Research and Application* (pp. 109-134). New York: Plenum.
- Coopersmith, S. (1967). *The antecedents of self-esteem*. San Francisco: Freeman.
- Coughtrey, A., Shafran, R., Bennett, S., Kothari, R., & Wade, T. (2017). Mental contamination: Relationship with psychopathology and transdiagnostic processes. *Journal of Obsessive-Compulsive and Related Disorders*. doi:https://doi.org/10.1016/j.jocrd.2017.08.009
- Cuijpers, P., Van Straten, A., & Andersson, G. (2008). Internet-administered cognitive behavior therapy for health problems: a systematic review. *Journal of behavioral medicine*, *31*(2), 169-177.
- Donkin, L., & Glozier, N. (2012). Motivators and motivations to persist with online psychological interventions: a qualitative study of treatment completers. *Journal of medical Internet research*, *14*(3), e91.
- Egan, S. J., & Hine, P. (2008). Cognitive behavioural treatment of perfectionism: A single case experimental design series. *Behaviour Change*, *25*(04), 245-258.
- Egan, S. J., van Noort, E., Chee, A., Kane, R. T., Hoiles, K. J., Shafran, R., & Wade, T. D. (2014). A randomised controlled trial of face to face versus pure online self-help cognitive behavioural treatment for perfectionism. *Behaviour research and therapy*, *63*, 107-113.
- Egan, S. J., Wade, T. D., & Shafran, R. (2011). Perfectionism as a transdiagnostic process: A clinical review. *Clinical psychology review*, *31*(2), 203-212.
- Egan, S. J., Wade, T. D., Shafran, R., & Antony, M. M. (2014). *Cognitive Behavioural Treatment of Perfectionism*. New York: Guilford Press.
- Elobeid, M. A., Padilla, M. A., McVie, T., Thomas, O., Brock, D. W., Musser, B., . . . St-Onge, M.-P. (2009). Missing data in randomized clinical trials for weight loss: scope of the problem, state of the field, and performance of statistical methods. *PloS one*, *4*(8), e6624.

- Evans, C. (1998). Reliable Change Criterion Calculator. Retrieved from <http://www.psych.org/stats/rcsc1.htm>
- Evans, C., Margison, F., & Barkham, M. (1998). The contribution of reliable and clinically significant change methods to evidence-based mental health. *Evidence-Based Mental Health, 1*(3), 70-72. doi:10.1136/ebmh.1.3.70
- Fairburn, C. G. (1994). Assessment of eating disorders: Interview or self-report questionnaire? *The International journal of eating disorders, 16*(4), 363.
- Fairburn, C. G., Cooper, Z., & Shafran, R. (2003). Cognitive behaviour therapy for eating disorders: a "transdiagnostic" theory and treatment. *Behav Res Ther, 41*(5), 509-528. doi:S0005796702000888 [pii]
- Foa, E. B., Huppert, J. D., Leiberg, S., Langner, R., Kichic, R., Hajcak, G., & Salkovskis, P. M. (2002). The Obsessive-Compulsive Inventory: development and validation of a short version. *Psychological assessment, 14*(4), 485.
- Freeston, M. H., Rhéaume, J., Letarte, H., Dugas, M. J., & Ladouceur, R. (1994). Why do people worry? *Personality and Individual Differences, 17*(6), 791-802.
- Frost, R. O., Marten, P., Lahart, C., & Rosenblate, R. (1990). The dimensions of perfectionism. *Cognitive therapy and research, 14*(5), 449-468.
- Gentes, E. L., & Ruscio, A. M. (2011). A meta-analysis of the relation of intolerance of uncertainty to symptoms of generalized anxiety disorder, major depressive disorder, and obsessive-compulsive disorder. *Clinical psychology review, 31*(6), 923-933.
- Gilbert, P. (2009). Introducing compassion-focused therapy. *Advances in Psychiatric Treatment, 15*(3), 199-208. doi:10.1192/apt.bp.107.005264
- Gilbert, P., McEwan, K., Gibbons, L., Chotai, S., Duarte, J., & Matos, M. (2012). Fears of compassion and happiness in relation to alexithymia, mindfulness, and self-criticism. *Psychology and Psychotherapy: Theory, Research and Practice, 85*(4), 374-390.
- Gilbert, P., McEwan, K., Matos, M., & Rivis, A. (2011). Fears of compassion: Development of three self-report measures. *Psychology and Psychotherapy: Theory, Research and Practice, 84*(3), 239-255.
- Gilbert, P., & Procter, S. (2006). Compassionate mind training for people with high shame and self-criticism: Overview and pilot study of a group therapy approach. *Clinical Psychology & Psychotherapy: An International Journal of Theory & Practice, 13*(6), 353-379.
- Glover, D. S., Brown, G. P., Fairburn, C. G., & Shafran, R. (2007). A preliminary evaluation of cognitive-behaviour therapy for clinical perfectionism: A case series. *British Journal of Clinical Psychology, 46*(1), 85-94.
- Hajcak, G., Huppert, J. D., Simons, R. F., & Foa, E. B. (2004). Psychometric properties of the OCI-R in a college sample. *Behaviour research and therapy, 42*(1), 115-123.
- Hamachek, D. E. (1978). Psychodynamics of normal and neurotic perfectionism. *Psychology: A Journal of Human Behavior.*
- Handley, A. K., Egan, S. J., Kane, R. T., & Rees, C. S. (2015). A randomised controlled trial of group cognitive behavioural therapy for perfectionism. *Behaviour research and therapy, 68*, 37-47.
- Hedeker, D., Gibbons, R. D., & Waternaux, C. (1999). Sample size estimation for longitudinal designs with attrition: comparing time-related contrasts between two groups. *Journal of Educational and Behavioral Statistics, 24*(1), 70-93.

- Hewitt, P. L., & Flett, G. L. (1991). Perfectionism in the self and social contexts: Conceptualization, assessment, and association with psychopathology. *Journal of personality and social psychology*, 60(3), 456 - 470.
- Huppert, J. D., Walther, M. R., Hajcak, G., Yadin, E., Foa, E. B., Simpson, H. B., & Liebowitz, M. R. (2007). The OCI-R: validation of the subscales in a clinical sample. *Journal of anxiety disorders*, 21(3), 394-406.
- Jacobson, N. S., & Truax, P. (1991). Clinical significance: a statistical approach to defining meaningful change in psychotherapy research. *Journal of consulting and clinical psychology*, 59(1), 12.
- Jacoby, R. J., Fabricant, L. E., Leonard, R. C., Riemann, B. C., & Abramowitz, J. S. (2013). Just to be certain: Confirming the factor structure of the Intolerance of Uncertainty Scale in patients with obsessive-compulsive disorder. *Journal of anxiety disorders*, 27(5), 535-542.
- Kothari, R. (2017). *Guided Internet-based Cognitive Behavioural Therapy for Perfectionism, and its Impact on Self-esteem and Intolerance of Uncertainty: A Randomised Controlled Trial*. (Doctorate in Clinical Psychology Doctoral Thesis), University College London, UK.
- Kothari, R., Egan, S., Wade, T., Andersson, G., & Shafran, R. (2016). Overcoming Perfectionism: Protocol of a Randomized Controlled Trial of an Internet-Based Guided Self-Help Cognitive Behavioral Therapy Intervention. *JMIR Research Protocols*, 5(4).
- Krabbendam, L., Janssen, I., Bak, M., Bijl, R. V., de Graaf, R., & van Os, J. (2002). Neuroticism and low self-esteem as risk factors for psychosis. *Social psychiatry and psychiatric epidemiology*, 37(1), 1-6.
- Lauder, S., Chester, A., & Berk, M. (2007). Net-effect? Online psychological interventions. *Acta Neuropsychiatrica*, 19(6), 386-388.
- Limburg, K., Watson, H. J., Hagger, M. S., & Egan, S. J. (2016). The Relationship Between Perfectionism and Psychopathology: A Meta-Analysis. *Journal of Clinical Psychology*.
- Lloyd, S., Schmidt, U., Khondoker, M., & Tchanturia, K. (2015). Can psychological interventions reduce perfectionism? A systematic review and meta-analysis. *Behavioural and cognitive psychotherapy*, 43(06), 705-731.
- Mahoney, A. E., & McEvoy, P. M. (2012). A transdiagnostic examination of intolerance of uncertainty across anxiety and depressive disorders. *Cognitive Behaviour Therapy*, 41(3), 212-222.
- Melville, K. M., Casey, L. M., & Kavanagh, D. J. (2010). Dropout from Internet-based treatment for psychological disorders. *British Journal of Clinical Psychology*, 49(4), 455-471.
- Mitchell, J., Stanimirovic, R., Klein, B., & Vella-Brodrick, D. (2009). A randomised controlled trial of a self-guided internet intervention promoting well-being. *Computers in Human Behavior*, 25(3), 749-760.
doi:<https://doi.org/10.1016/j.chb.2009.02.003>
- Musiat, P., Goldstone, P., & Tarrier, N. (2014). Understanding the acceptability of e-mental health-attitudes and expectations towards computerised self-help treatments for mental health problems. *BMC psychiatry*, 14(1), 109.
- Neff, K. D. (2003). The Development and Validation of a Scale to Measure Self-Compassion. *Self and Identity*, 2(3), 223-250. doi:10.1080/15298860309027
- Obsessive Compulsive Cognitions Working, G. (1997). Cognitive assessment of obsessive-compulsive disorder. *Behaviour research and therapy*, 35(7), 667-681. doi:[https://doi.org/10.1016/S0005-7967\(97\)00017-X](https://doi.org/10.1016/S0005-7967(97)00017-X)

- Obsessive Compulsive Cognitions Working, G. (2003). Psychometric validation of the Obsessive Beliefs Questionnaire and the Interpretation of Intrusions Inventory: Part I. *Behaviour research and therapy*, 41(8), 863-878. doi:[https://doi.org/10.1016/S0005-7967\(02\)00099-2](https://doi.org/10.1016/S0005-7967(02)00099-2)
- Orth, U., Robins, R. W., Trzesniewski, K. H., Maes, J., & Schmitt, M. (2009). Low self-esteem is a risk factor for depressive symptoms from young adulthood to old age. *Journal of abnormal psychology*, 118(3), 472.
- Pleva, J., & Wade, T. D. (2007). Guided self-help versus pure self-help for perfectionism: A randomised controlled trial. *Behaviour research and therapy*, 45(5), 849-861. doi:<https://doi.org/10.1016/j.brat.2006.08.009>
- Powell, J., Hamborg, T., Stallard, N., Burls, A., McSorley, J., Bennett, K., . . . Christensen, H. (2013). Effectiveness of a Web-Based Cognitive-Behavioral Tool to Improve Mental Well-Being in the General Population: Randomized Controlled Trial. *Journal of medical Internet research*, 15(1), e2. doi:10.2196/jmir.2240
- Radomsky, A. S., Rachman, S., Shafran, R., Coughtrey, A. E., & Barber, K. C. (2014). The nature and assessment of mental contamination: A psychometric analysis. *Journal of Obsessive-Compulsive and Related Disorders*, 3(2), 181-187. doi:<https://doi.org/10.1016/j.jocrd.2013.08.003>
- Reas, D. L., Grilo, C. M., & Masheb, R. M. (2006). Reliability of the Eating Disorder Examination-Questionnaire in patients with binge eating disorder. *Behaviour Research and Therapy*, 44(1), 43-51.
- Riley, C., Lee, M., Cooper, Z., Fairburn, C. G., & Shafran, R. (2007). A randomised controlled trial of cognitive-behaviour therapy for clinical perfectionism: A preliminary study. *Behaviour Research and Therapy*, 45(9), 2221-2231.
- Rosenberg, M. (1965). *Society and the adolescent child*: Princeton, NJ: Princeton University Press.
- Rosenberg, M. (1965). *Society and the adolescent self-image*. Princeton, NJ: Princeton University Press.
- Rozental, A., Andersson, G., Boettcher, J., Ebert, D. D., Cuijpers, P., Knaevelsrud, C., . . . Carlbring, P. (2014). Consensus statement on defining and measuring negative effects of Internet interventions. *Internet Interventions*, 1(1), 12-19. doi:10.1016/j.invent.2014.02.001
- Rozental, A., Shafran, R., Wade, T., Egan, S., Nordgren, L. B., Carlbring, P., . . . Andersson, G. (2017). A randomized controlled trial of Internet-Based Cognitive Behavior Therapy for perfectionism including an investigation of outcome predictors. *Behaviour research and therapy*. doi:<https://doi.org/10.1016/j.brat.2017.05.015>
- Rozental, A., Shafran, R., Wade, T., Nordgren, L. B., Carlbring, P., Landstrom, A., . . . Andersson, A. (2017). A Randomized Controlled Trial of Internet-Based Cognitive Behavior Therapy for Perfectionism Including an Investigation of Outcome Predictors. *Behaviour research and therapy*.
- Schmitt, D. P., & Allik, J. (2005). Simultaneous administration of the Rosenberg Self-Esteem Scale in 53 nations: exploring the universal and culture-specific features of global self-esteem. *Journal of personality and social psychology*, 89(4), 623.
- Sealed Envelope. Retrieved from <http://www.sealedenvelope.com/>
- Shafran, R., Cooper, Z., & Fairburn, C. G. (2002). Clinical perfectionism: A cognitive-behavioural analysis. *Behaviour research and therapy*, 40(7), 773-791.

- Shafran, R., Coughtrey, A., & Kothari, R. (2016). New frontiers in the treatment of perfectionism. *International Journal of Cognitive Therapy*, 9(2), 156-170.
- Shafran, R., Egan, S., & Wade, T. (2010). *Overcoming perfectionism: A self-help guide using cognitive behavioural techniques*: Robinson Publishing.
- Shafran, R., & Mansell, W. (2001). Perfectionism and psychopathology: A review of research and treatment. *Clinical psychology review*, 21(6), 879-906.
- Shafran, R., Wade, T., Egan, S., Kothari, R., Allcott-Watson, H., Carlbring, P., . . . Andersson, G. (2017). Is the devil in the detail? A randomized controlled trial of guided internet-based CBT for perfectionism. *Behaviour research and therapy*.
- Slaney, R. B., Rice, K. G., & Ashby, J. S. (2002). A programmatic approach to measuring perfectionism: The Almost Perfect Scales.
- Smith, M. M., Sherry, S. B., Chen, S., Saklofske, D. H., Mushquash, C., Flett, G. L., & Hewitt, P. L. (2017). The perniciousness of perfectionism: A meta-analytic review of the perfectionism–suicide relationship. *Journal of Personality*, n/a-n/a. doi:10.1111/jopy.12333
- Steele, A. L., & Wade, T. D. (2008). A randomised trial investigating guided self-help to reduce perfectionism and its impact on bulimia nervosa: A pilot study. *Behaviour research and therapy*, 46(12), 1316-1323.
- Steele, A. L., Waite, S., Egan, S. J., Finnigan, J., Handley, A., & Wade, T. D. (2013). Psycho-education and group cognitive-behavioural therapy for clinical perfectionism: a case-series evaluation. *Behavioural and cognitive psychotherapy*, 41(02), 129-143.
- Stoeber, J., & Otto, K. (2006). Positive conceptions of perfectionism: Approaches, evidence, challenges. *Personality and social psychology review*, 10(4), 295-319.
- Suddarth, B. H., & Slaney, R. B. (2001). An investigation of the dimensions of perfectionism in college students. *Measurement and evaluation in counseling and development*, 34(3), 157.
- Wagner, B., Horn, A. B., & Maercker, A. Internet-based versus face-to-face cognitive-behavioral intervention for depression: A randomized controlled non-inferiority trial. *Journal of Affective Disorders*, 152, 113-121. doi:10.1016/j.jad.2013.06.032

Figure 1 Modules and components of Overcoming Perfectionism

Module	Module Components
1. Understanding Perfectionism	1.1. What is unhelpful perfectionism? 1.2. Why perfectionism continues 1.3. Fact or fiction? 1.4. "The harder you work, the better you'll do" - Fact or fiction? 1.5. Facts about perfectionism and performance 1.6. Preparing for change 1.7. Key take away 1.8. Between module work
2. Your Perfectionism Cycle	2.1. Between-module work 2.2. A reminder 2.3. The first steps 2.4. Drawing your own diagram 2.5. Between-module work 2.6. Take-home message
3. Surveys and Experiments	3.1. Between-module work 3.2. Perfectionism behaviours 3.3. Surveys 3.4. Reflect on the responses 3.5. Behavioural experiments 3.6. Different forms of behavioural experiments 3.7. An added benefit 3.8. Between-module work 3.9. Take home message
4. New Ways of Thinking	4.1. Between-module work 4.2. Changing thinking 4.3. Imagining vivid future positive outcomes 4.4. From all or nothing thinking to flexibility and freedom 4.5. "Rules break, guidelines bend": Turning rigid rules into guidelines 4.6. Changing thinking styles 4.7. Between module work 4.8. Key take away
5. Useful Skills for Managing Unhelpful Perfectionism	5.1. Procrastination 5.2. Problem solving 5.3. Pleasant events 5.4. Take home message 5.5. Before the next module
6. Self-criticism or Self-compassion	6.1. How to respond 6.2. Take home message 6.3. Before the next module
7. Re-examining the way we Examine our Self-worth	7.1. Your self-worth 7.2. Step 1. Recognizing that your self-worth can be independent of your achievements 7.3. Step 2. Encouraging flexible and realistic goals 7.4. Step 3. Spreading your self-worth across as many areas of your life as possible 7.5. Step 4. Develop more balance in what you pay attention to daily 7.6. Take home message 7.7. Before the next module
8. Staying Well – Managing Unhelpful Perfectionism in the Long-term	8.1. Improve your sense of self-worth 8.2. Questions 8.3. Thank you!


Figure 2 Flow of participants through trial.

Table 1 Baseline characteristics of overall sample and logistic regression analysis of whether baseline variables predict missing data (post-intervention, follow-up).

Characteristics (N=120)	Frequency (%)	Post-intervention OR (95% CI), p-value	Follow-up OR (95% CI)
<i>Gender</i>			
Female	98 (81.7)	Ref.	Ref.
Male	22 (18.3)	1.26 (0.5 – 3.2), 0.63	1.6 (0.63 – 4.05)
<i>Age (years)</i>	28.93 (7.98)	0.96 (0.92 – 1.01), 0.15	0.98 (0.93 – 1.02)
<i>Marital Status</i>			
Single (never married)	88 (73.3)	Ref.	Ref.
Married/domestic Partnership	27 (22.5)	3.04 (0.33 – 28.32), 0.33	1.14 (0.18 – 7.17)
Divorced/separated/ Widowed	5 (4.1)	2.35 (0.23 – 24.1), 0.47	1.62 (0.23 – 11.26)
<i>Educational Qualification Certificate</i>			
No	11 (9.2)	Ref.	Ref.
Yes	109 (90.8)	0.23 (0.06 – 0.9), 0.04*	0.27 (0.07 – 1.09)
<i>Professional Vocational Certificate</i>			
No	63 (52.5)	Ref.	Ref.
Yes	57 (47.5)	0.73 (0.35 – 1.52), 0.4	0.92 (0.45 – 1.88)
<i>Currently studying for degree level qualification</i>			
No	58 (48.3)	Ref.	Ref.
Yes	62 (51.7)	0.96 (0.46 – 1.98), 0.91	0.68 (0.33 – 1.39)
<i>Ethnicity</i>			
White British	52 (44.1)	Ref.	Ref.
Other Ethnicity	66 (55.9)	1.48 (0.7 – 3.14), 0.31	1.14 (0.55 – 2.37)
<i>Currently receiving treatment for a mental health problem</i>			
No	86 (71.7)	Ref.	Ref.
Yes	34 (28.3)	1.21 (0.54 – 2.7), 0.65	1.32 (0.6 – 2.94)
<i>Previously received treatment for a mental health problem</i>			
No	73 (60.8)	Ref.	Ref.
Yes	47 (39.2)	1.12 (0.53 – 2.37), 0.76	1.3 (0.62 – 2.71)
<i>Group</i>			
Control	58 (48.3)	Ref.	Ref.
Experimental	62 (51.7)	2.57 (1.21 – 5.47), 0.01*	2.66 (1.27 – 5.6)
	Mean (SD)	Post-intervention OR (95% CI), p-value	Follow-up OR (95% CI)
<i>Psychopathology</i>			
OCI-R	27.88 (13.71)	1.02 (1.0 – 1.05), 0.08	1.03 (1.00 – 1.06)
EDE-Q	2.43 (1.63)	1.29 (1.02 – 1.62), 0.03	1.2 (0.96 – 1.5)
<i>Transdiagnostic Processes</i>			
RSES	11.85 (5.3)	0.98 (0.92 – 1.05), 0.61	0.96 (0.9 – 1.03)
US	91.38 (20.95)	1 (0.98 – 1.01), 0.73	1 (0.98 – 1.01)
FSC-1	28.98 (14.8)	1.01 (0.99 – 1.04), 0.36	1.01 (0.99 – 1.04)

Note: OR = odds ratio; 95% CI = 95% confidence intervals; OCI-R = Obsessive Compulsive Inventory – Revised; EDE-Q = Eating Disorder Examination Questionnaire; RSES = Rosenberg Self-Esteem Scale; IUS = Intolerance of Uncertainty Scale; FCS = Fear of Self Compassion Scale 1 (fear of self-compassion).

Table 2 ANCOVA analysis comparing control and experimental groups on scores at 12 and 24 weeks, adjusted for pre-intervention scores

	Baseline covariate	Control, M (SE) n = 58 (42 Completers)	Experimental, M (SE) n = 62 (31 Completers)	Between group effect size d, 95% C.I.
Post intervention (T₂)				
<i>Psychopathology</i>				
OCI-R	26.07 (13.27) <i>27.88 (13.71)</i>	22.64 (1.46) <i>21.69 (1.22)</i>	14.13 (1.7) <i>16.51 (1.21)</i>	-0.91 (-1.4 – -0.42) <i>-0.55 (-0.92 – -0.19)</i>
EDE-Q	2.16 (1.46) <i>2.43 (1.63)</i>	2.15 (0.17) <i>2.03 (0.18)</i>	1.54 (0.2) <i>1.77 (0.19)</i>	-0.56 (-1.03 – -0.09) <i>-0.18 (-0.54 – 0.18)</i>
<i>Transdiagnostic Processes</i>				
RSES	12.06 (5.53) <i>11.85 (5.3)</i>	13.1 (0.54) <i>12.75 (0.75)</i>	15.6 (0.63) <i>15.25 (0.78)</i>	0.72 (0.24 – 1.2) <i>0.42 (0.06 – 0.79)</i>
IUS	91.93 (21.71) <i>91.38 (20.95)</i>	89.95 (3.06) <i>87.38 (2.21)</i>	71.71 (3.58) <i>77.4 (2.14)</i>	-0.93 (-1.42 – -0.44) <i>-0.6 (-0.96 – -0.23)</i>
FCS	27.96 (14.68) <i>28.98 (14.8)</i>	26.35 (1.85) <i>24.78 (1.43)</i>	16.72 (2.16) <i>20.06 (1.41)</i>	-0.81 (-1.3 – -0.33) <i>-0.43 (-0.79 – -0.07)</i>
Follow-up (T₃)				
<i>Psychopathology</i>				
OCI-R	25.53 (12.62) <i>27.88 (13.71)</i>	21.19 (1.23) <i>19.75 (1.12)</i>	11.49 (1.42) <i>14.68 (1.07)</i>	-1.27 (-1.78 – -0.75) <i>-0.6 (-0.97 – -0.24)</i>
EDE-Q	2.22 (1.51) <i>2.43 (1.63)</i>	2.15 (0.15) <i>1.99 (1.19)</i>	1.52 (0.18) <i>1.76 (0.17)</i>	-0.66 (-1.15 – -0.18) <i>-0.04 (-0.39 – 0.32)</i>
<i>Transdiagnostic Processes</i>				
RSES	12.33 (5.29) <i>11.85 (5.3)</i>	13.68 (0.83) <i>13.15 (1.49)</i>	15.09 (1) <i>14.59 (2.78)</i>	0.27 (-0.21 – 0.74) <i>0.08 (-0.28 – 0.44)</i>
IUS	92.15 (19.78) <i>91.38 (20.95)</i>	83.65 (3.18) <i>81.47 (2.13)</i>	68.44 (3.83) <i>73.72 (2.1)</i>	-0.75 (-1.24 – -0.26) <i>-0.48 (-0.84 – -0.11)</i>
FCS	27.76 (14.72) <i>28.98 (14.8)</i>	26.41 (1.91) <i>24.83 (1.47)</i>	14.64 (2.29) <i>18.37 (1.47)</i>	-0.97 (-1.47 – -0.47) <i>-0.57 (-0.94 – -0.21)</i>

Note: Results of intent to treat analysis presented in italics; d = effect size; 95% CI = 95% confidence intervals; OCI-R = Obsessive Compulsive Inventory – Revised; EDE-Q = Eating Disorder Examination Questionnaire; RSES = Rosenberg Self-Esteem Scale; IUS = Intolerance of Uncertainty Scale; FCS = Fear of Self Compassion Scale (fear of self-compassion).

Table 3 Frequency and odds of participants in the experimental group (vs. control group) achieving reliable change on all outcomes, post-treatment.

	Cronbach's Alpha	Change Criterion	Control, n (%) Improvement	No Change	Det.	Experimental, n (%) Improvement	No Change	Det.	OR (95% CI)
Post-treatment (T₂)									
<i>Psychopathology</i>									
OCI-R	0.92	± 10.75	16 (27.6)	40 (69.0)	2 (3.4)	29 (46.8)	30 (48.4)	3 (4.8)	2.42 (1.04 – 5.66)
EDE-Q	0.92	± 1.28	11 (19.0)	41 (70.7)	6 (10.3)	21 (33.9)	31 (50)	10 (16.1)	2.48 (0.96 – 6.37)
<i>Transdiagnostic Processes</i>									
RSES	0.87	± 5.3	9 (15.5)	45 (75.9)	5 (8.6)	18 (29)	41 (64.5)	4 (6.5)	2.32 (0.78 – 6.92)
IUS	0.94	± 14.22	20 (34.5)	28 (48.3)	10 (17.2)	26 (41.9)	29 (46.8)	7 (11.3)	1.35 (0.62 – 2.95)
FCS-1	0.95	± 9.17	22 (37.9)	31 (53.4)	5 (8.6)	35 (56.5)	17 (27.4)	10 (16.1)	2.98 (1.29 – 6.87)
Follow-up (T₃)									
<i>Psychopathology</i>									
OCI-R	0.92	± 10.75	21 (36.2)	36 (62.1)	1 (1.7)	35 (56.5)	25 (40.3)	2 (3.2)	2.42 (1.09 – 5.41)
EDE-Q	0.92	± 1.28	12 (20.7)	41 (70.7)	5 (8.6)	18 (29.0)	38 (61.3)	6 (9.7)	1.60 (0.6 – 4.29)
<i>Transdiagnostic Processes</i>									
RSES	0.87	± 5.3	13 (22.4)	37 (63.8)	8 (13.8)	23 (37.1)	26 (41.9)	13 (21)	1.98 (0.74 – 5.26)
IUS	0.94	± 14.22	28 (48.3)	23 (39.6)	7 (12.1)	29 (46.8)	26 (41.9)	7 (11.3)	0.95 (0.45 – 1.99)
FCS-1	0.95	± 9.17	21 (36.2)	27 (46.6)	10 (17.2)	31 (50.0)	23 (37.1)	8 (12.9)	1.76 (0.77 – 4.03)

1. Reliable change criterion calculated using *Reliable Change Criterion Calculator* (Chris Evans, 1998).

2. Det. = Deterioration.

3. Table shows results of logistic regression analysis (OR, 95% CI).

4. OCI-R = Obsessive Compulsive Inventory – Revised; EDE-Q = Eating Disorder Examination Questionnaire; RSES = Rosenberg Self-Esteem Scale; IUS = Intolerance of Uncertainty Scale; FCS = Fear of Self Compassion Scale 1 (fear of self-compassion).

The authors declare no conflicts of interest. The research did not receive any specific grant from funding agencies in the public, commercial, or not-for-profit sectors, however, all research at Great Ormond Street Hospital NHS Foundation Trust and UCL Great Ormond Street Institute of Child Health is made possible by the NIHR Great Ormond Street Hospital Biomedical Research Centre. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.