

LETTER

Concomitant fragile X-associated tremor ataxia syndrome and Parkinson's disease: a clinicopathological report of two cases

INTRODUCTION

Parkinsonism is common in fragile X-associated tremor/ataxia syndrome (FXTAS) but its underlying pathophysiology remains unknown. Our group reported a patient with FXTAS and parkinsonism, and *in vivo* evidence of presynaptic and postsynaptic nigrostriatal dysfunction.¹ We report the histological findings of this case (case 1) and another case with dual pathologies of FXTAS and Lewy body Parkinson's disease (PD).

METHODS

Genetic testing for FXTAS was requested in seven cases from the Queen Square Brain Bank for Neurological Disorders (QSBB) based on the characteristic radiological 'middle cerebellar peduncle' (MCP) sign (case 1, diagnosis confirmed in life) and histological findings of round, eosinophilic p62-positive neuronal intranuclear inclusions in the hippocampus or unexplained cerebellar degeneration (6 postmortem cases were tested). Two cases were positive for *FMR1* premutation and both showed histological features of FXTAS and PD. The QSBB brain donor programme was approved by a Research Ethics Committee and written consent for research was obtained from all cases.

CASE REPORTS

Case 1

A 68-year-old man presented with a 10-year history of slowly progressive right-hand tremor. There was no family history of neurological disorders or mental retardation. Examination showed hypomimia, asymmetric rest and kinetic tremor, rigidity, mild bradykinesia and inability to tandem walk. The patient was diagnosed with tremor-predominant PD but did not respond to L-dopa (600 mg/day). Over the next few years he developed mild cognitive impairment, gait ataxia and intention tremor. Brain MRI demonstrated the MCP sign and diffuse atrophy. Genetic analysis confirmed *FMR1* premutation with 87 CGG expansion. ¹²³I-FP-CIT SPECT revealed asymmetric reduced tracer uptake with the right side more affected. D₂ receptor (123I) IBZM-SPECT (IBZM) scan showed

slightly reduced tracer uptake bilaterally. There was clinical and neurophysiological evidence of large fibre axonal peripheral neuropathy. The patient's University of Pennsylvania Smell Identification Test (UPSIT) score was 25/40 (severe hyposmia: 19–25). His condition gradually deteriorated and he died of pneumonia at age 72.

Case 2

A 21-year-old man was diagnosed with schizophrenia and his symptoms responded well to long-term fluphenazine. At the age of 35, he developed a right-hand tremor initially attributed to the neuroleptic treatment. The tremor persisted despite switching fluphenazine to olanzapine 5 mg/day and administering anticholinergics (orphenadrine and procyclidine). His tremor remained stable and only began to deteriorate at age 51 along with the onset of slurred speech and dragging of right leg. Examination showed mild hypomimia, hypophonia, restricted downgaze, slow vertical saccades, jaw and tongue tremor, intermittent right-sided predominant hand tremor, bradykinesia and cogwheel rigidity. Olanzapine was replaced by quetiapine 200 mg/day and L-dopa (300 mg/day) was administered without any improvement. Brain MRI showed global atrophy without the MCP sign. At age 58, his symptoms rapidly deteriorated with severe hand tremor and frequent falls requiring a wheelchair. ¹²³I-FP-CIT SPECT demonstrated asymmetrical reduction of tracer uptake in putamen and left caudate. A clinical diagnosis of progressive supranuclear palsy (PSP) was made due to the rapid motor deterioration, postural instability, restriction of downgaze and absence of L-dopa response. He died of pulmonary embolism at age 59. Neuropathological findings prompted genetic testing and the identification of *FMR1* gene premutation with 100 CGG expansions.

NEUROPATHOLOGICAL EXAMINATION

In both cases, histological examination demonstrated widespread p62-positive and ubiquitin-positive and 1C2-negative neuronal and glial intranuclear inclusions with mild Purkinje cell depletion consistent with FXTAS. There was loss of pigmented neurons in the substantia nigra with α -synuclein-positive Lewy bodies, in keeping with PD. Lewy body pathology conformed to Braak stage 5 in case 1 and stage 3 in case 2. Age-related neurofibrillary tangle pathology was also noted in both cases (Braak and Braak stage II in case 1 and stage I in case 2). An acute left

middle cerebral artery territory infarction was observed in case 2.

DISCUSSION

In the literature, only 13 of 29 (45%) cases with *FMR1* premutation and parkinsonism had abnormal dopamine transporter scans (see online supplementary table) and it is likely that these 13 cases may have had concomitant PD pathology akin to the two cases in this series. On the other hand, IBZM scans were only available in four cases with *FMR1* premutation including our case and all had abnormal findings (see online supplementary table). We propose that postsynaptic dopaminergic striatal pathway dysfunction is responsible for parkinsonism in individuals with FXTAS. *FMR1* premutation carriers with an abnormal dopamine transporter scan probably signifies the presence of concomitant PD pathology and may have a rapidly progressive disease course due to the dual pathologies. These patients may benefit from L-dopa therapy.

Of the 22 FXTAS cases with postmortem data (table 1), concomitant pathologies were identified in nine cases (41%): four had Alzheimer's disease pathology,² one had multiple sclerosis (MS)² and four had PD pathology including the two cases presented here.^{2,3}

It is difficult to ascertain the exact contribution of each pathology to the clinical course of parkinsonism in our cases. In case 2, the long-term use of neuroleptics complicated the clinical picture, thus the longstanding hand tremor could have been caused by a combination of FXTAS and neuroleptics. However, at age 51, the deterioration of hand tremor, relentless progression of parkinsonism and the onset of typical features of PD including jaw and tongue tremor collectively marked the onset of an additional underlying neurodegenerative process of PD. Both cases had a relatively benign course in the first decade of symptom onset followed by rapidly progressive parkinsonism and motor disability leading to death. This rapid deterioration is atypical for either FXTAS or PD. Indeed, the clinical diagnosis for case 2 was revised to PSP. Among the four patients with dual FXTAS and PD pathology in the literature,^{2,3} the time from onset to use of walking aid (median: 7.5 years) and life expectancy (median: 11 years) are much shorter than in other patients with isolated FXTAS, in whom the median time to walking aid is 15 years and median survival is 21 years.⁴ These data suggest that the synergistic effect of more than one neurodegenerative condition might have

Table 1 Features of 22 reported postmortem FXTAS cases

	Duration from onset (years)		CCG repeats	MCP sign	Additional neuropathological findings
	Wheelchair	Death			
Greco <i>et al</i> ³ ; 11 males ⁶	6	10	113	–	–
	–	13	80	–	–
	NA	5	80	Y	–
	13	14	71	–	PD
	12	19	105	–	–
	7	13	77	Y	–
	5	6	88	Y	–
	NA	3	92	–	–
	13	16	88	Y	–
	4	6	93	Y	–
	NA	1	65	–	–
Tassone <i>et al</i> ² ; 8 females	NA	NA	70	–	–
	–	10	80	N	AD (B&B V-VI)
	–	6	87	–	–
	–	14	75	Y	MS
	–	7	65	–	AD (B&B V-VI)
	4	7	–	N	AD (B&B V-VI)
	4	6	59	N	PD
Louis <i>et al</i> ; 1 male ⁷	7	9	90	–	–
	13	14	72	Y	PD (Braak 5)
	7	8	100	N	PD (Braak 3)
Present series: De Pablo-Fernandez <i>et al</i> ; 2 males	7	8	100	N	PD (Braak 3)
Total: 8 females; 14 males	8	9		Y	
				64% (7/11)	

AD, Alzheimer disease; B&B, Braak and Braak stage for Alzheimer; Braak, Braak stage for Parkinson's disease; FXTAS, fragile X-associated tremor/ataxia syndrome; MCP, middle cerebellar peduncle; MS, multiple sclerosis; N, no; NA, not applicable; PD, Parkinson's disease; Y, yes.

Provenance and peer review Not commissioned; externally peer reviewed.



OPEN ACCESS

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/jnnp-2014-309460>).

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

To cite De Pablo-Fernandez E, Doherty K M, Holton JL, *et al*. *J Neurol Neurosurg Psychiatry* Published Online First: [please include Day Month Year] doi:10.1136/jnnp-2014-309460

Received 10 September 2014

Revised 7 November 2014

Accepted 13 November 2014

J Neurol Neurosurg Psychiatry 2014;**0**:1–2.

doi:10.1136/jnnp-2014-309460

contributed to an accelerated disease course. Carriers of *FMR1* premutation with concomitant Alzheimer's disease,² heterozygous *Parkin* mutation⁵ or MS² have been reported and these secondary pathological processes have been postulated to modulate the effect of *FMR1* premutation.

Eduardo De Pablo-Fernandez^{1,2}
Karen M Doherty^{1,2} **Janice L Holton**^{1,2,3}
Tamas Revesz^{1,2} **Atbin Djamshidian**¹
Patricia Limousin^{3,4} **Kailash P Bhatia**^{3,4}
Thomas T Warner^{1,2,3} **Andrew J Lees**^{1,2}
Helen Ling^{1,2}

¹Reta Lila Weston Institute of Neurological Studies, UCL Institute of Neurology, London, UK

²Queen Square Brain Bank for Neurological Disorders, UCL Institute of Neurology, London, UK

³National Hospital for Neurology and Neurosurgery, Queen Square, UCL Hospitals NHS Foundation Trust, London, UK

⁴Sobell Department of Motor Neuroscience, UCL Institute of Neurology, London, UK

Correspondence to Dr Helen Ling, Reta Lila Weston

Institute of Neurological Studies, UCL Institute of Neurology, 1 Wakefield Street, London WC1N 1PJ, UK; h.ling@ucl.ac.uk

Acknowledgements The authors thank Dr Thomasin Andrews for providing clinical information regarding case 2.

Contributors ED-PF contributed to project conception and organisation, and data collection, and wrote the first draft; KMD contributed to project conception and organisation, and data collection, and critically reviewed the manuscript; JLH and TR performed histological analysis, contributed to pathology content and critically reviewed the manuscript; AD, PL, KPB, TTW and AJL contributed to clinical assessment, diagnostic process and critically reviewed the manuscript; HL contributed to project conception and organisation, and critically reviewed the manuscript.

Competing interests None.

Patient consent Obtained.

Ethics approval The brain donor programme of the QSBB was approved by a London Multi-Centre Research Ethics Committee and tissue is stored under a license from the Human Tissue Authority. Consent for use of tissue for research was obtained from donors and/or their next of kin.

REFERENCES

- 1 Healy DG, Bressman S, Dickson J, *et al*. Evidence for pre and postsynaptic nigrostriatal dysfunction in the fragile X tremor-ataxia syndrome. *Mov Disord* 2009;**24**:1245–7.
- 2 Tassone F, Greco CM, Hunsaker MR, *et al*. Neuropathological, clinical and molecular pathology in female fragile X premutation carriers with and without FXTAS. *Genes Brain Behav* 2012;**11**:577–85.
- 3 Greco CM, Berman RF, Martin RM, *et al*. Neuropathology of fragile X-associated tremor/ataxia syndrome (FXTAS). *Brain* 2006;**129**(Pt 1):243–55.
- 4 Leehey MA, Berry-Kravis E, Min SJ, *et al*. Progression of tremor and ataxia in male carriers of the *FMR1* premutation. *Mov Disord* 2007;**22**:203–6.
- 5 Hedrich K, Pramstaller PP, Stubke K, *et al*. Premutations in the *FMR1* gene as a modifying factor in *Parkin*-associated Parkinson's disease? *Mov Disord* 2005;**20**:1060–2.
- 6 Greco CM, Hagerman RJ, Tassone F, *et al*. Neuronal intranuclear inclusions in a new cerebellar tremor/ataxia syndrome among fragile X carriers. *Brain* 2002;**125**:1760–71.
- 7 Louis E, Moskowitz C, Friez M, *et al*. Parkinsonism, dysautonomia, and intranuclear inclusions in a fragile X carrier: a clinical-pathological study. *Mov Disord* 2006;**21**:420–5.



Concomitant fragile X-associated tremor ataxia syndrome and Parkinson's disease: a clinicopathological report of two cases

Eduardo De Pablo-Fernandez, Karen M Doherty, Janice L Holton, Tamas Revesz, Atbin Djamshidian, Patricia Limousin, Kailash P Bhatia, Thomas T Warner, Andrew J Lees and Helen Ling

J Neurol Neurosurg Psychiatry published online December 4, 2014

Updated information and services can be found at:

<http://jnp.bmj.com/content/early/2014/12/04/jnp-2014-309460>

These include:

Supplementary Material

Supplementary material can be found at:

<http://jnp.bmj.com/content/suppl/2014/12/04/jnp-2014-309460.DC1.html>

References

This article cites 5 articles, 1 of which you can access for free at:

<http://jnp.bmj.com/content/early/2014/12/04/jnp-2014-309460#BIBL>

Open Access

This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

Email alerting service

Receive free email alerts when new articles cite this article. Sign up in the box at the top right corner of the online article.

Topic Collections

Articles on similar topics can be found in the following collections

[Open access](#) (157)

Notes

To request permissions go to:

<http://group.bmj.com/group/rights-licensing/permissions>

To order reprints go to:

<http://journals.bmj.com/cgi/reprintform>

To subscribe to BMJ go to:

<http://group.bmj.com/subscribe/>