

Airway smooth muscle inflammation is regulated by microRNA-145 in COPD

Lawrence O'Leary^{1,2}, Kenan Sevinç^{1,2}, Ilektra M. Papazoglou^{1,2}, Bernadett Tildy^{1,2}, Karen Detillieux³, Andrew J. Halayko³, Kian Fan Chung^{1,2} and Mark M. Perry⁴

1 Airways Disease, National Heart and Lung Institute, Imperial College, London, UK

2 Royal Brompton NIHR Biomedical Research Unit, London, UK

3 Departments of Internal Medicine & Physiology, Respiratory Hospital, Winnipeg, MB, Canada

4 Molecular Neurosciences, The Dubowitz Neuromuscular Centre, UCL Institute of Child Health, London, UK

Correspondence

M. M. Perry, Molecular Neurosciences, The Dubowitz Neuromuscular Centre, UCL Institute of Child Health, 30 Guildford Street, London WC1N 1EH, UK
Fax: 0207 905 2832
Tel: 0207 905 2136
E-mail: mark.perry@ucl.ac.uk

(Received 8 February 2016, revised 30 March 2016, accepted 30 March 2016)

doi:10.1002/1873-3468.12168

Edited by Tamas Dalmay

Chronic obstructive pulmonary disease (COPD) is a common, highly debilitating disease of the airways, primarily caused by smoking. Chronic inflammation and structural remodelling are key pathological features of this disease, in part caused by the aberrant function of airway smooth muscle (ASM) cells under the regulation of transforming growth factor (TGF)- β . miRNA are short, noncoding gene transcripts involved in the negative regulation of specific target genes, through their interactions with mRNA. Previous studies have proposed that mRNA-145 (*miR-145*) may interact with *SMAD3*, an important downstream signalling molecule of the TGF- β pathway. TGF- β was used to stimulate primary human ASM cells isolated from healthy non-smokers, healthy smokers and COPD patients. This resulted in a TGF- β -dependent increase in CXCL8 and IL-6 release, most notably in the cells from COPD patients. TGF- β stimulation increased *SMAD3* expression, only in cells from COPD patients, with a concurrent increased *miR-145* expression. Regulation of *miR-145* was found to be negatively controlled by pathways involving the MAP kinases, MEK-1/2 and p38 MAPK. Subsequent, overexpression of *miR-145* (using synthetic mimics) in ASM cells from patients with COPD suppressed IL-6 and CXCL8 release, to levels comparable to the nonsmoker controls. Therefore, this study suggests that *miR-145* negatively regulates pro-inflammatory cytokine release from ASM cells in COPD by targeting *SMAD3*.

Keywords: COPD; inflammation; microRNA

Chronic obstructive pulmonary disease (COPD) is a chronic inflammatory condition of the lung of high global prevalence [1] and is associated with high morbidity, mortality and socioeconomic cost [2]; moreover, its contribution to deaths worldwide is predicted to increase over the course of both the current and next decade [3].

Abbreviations

ASM, airway smooth muscle; COPD, chronic obstructive pulmonary disease; CXCL8, CXC chemokine ligand 8; HSFs, hypertrophic scar fibroblasts; IL-6, interleukin 6; MAPKs, mitogen-activated protein kinases; *miR-145*, microRNA-145; NF- κ B, nuclear factor- κ B; SSC, systemic sclerosis; TGF, transforming growth factor; VSMCs, vascular smooth muscle cells; WNT, wntless/integrase-1.

Chronic obstructive pulmonary disease is a heterogeneous condition, primarily affecting the lung, but often with significant systemic features, which can augment the morbidity of the disorder and hamper its management [2]. The pulmonary component of the disease refers to a progressive and largely irreversible obstruction of airflow, due to a pathological combination of

narrowing of the small airways, parenchymal destruction and structural remodelling [4]. The aberrant remodelling occurs in addition, or in response, to the ongoing and worsening inflammation [5]. It is associated with architectural alterations to the bronchi [6], small airways and parenchymal tissue and affects the epithelium, its underlying extracellular matrix and the surrounding smooth muscle and these changes have been correlated with disease severity [7]. Although the primary cause of COPD in more economically developed countries is cigarette smoking, the majority of heavy smokers do not develop the disease [8] and indicators for determining those that will be affected remain elusive.

An increased ASM mass has been observed in both the large and small airways in COPD [6], which correlates with disease severity [7]. Whether increased ASM mass in COPD is due to either hypertrophy or hyperproliferation, or a combination of both has not been definitively determined. ASM cells have also been shown to produce a number of cytokines, chemokines and growth factors in response to a variety of inflammatory stimuli, which may contribute towards the inflammatory process in COPD [9]. Among these secreted cytokines are CXC chemokine ligand 8 (CXCL8) and interleukin 6 (IL-6) both of which have been found to be elevated in the sputum of patients with COPD; even more so during exacerbations [10].

Transforming growth factor (TGF)- β is a pleiotropic cytokine that stimulates ASM proliferation and pro-inflammatory cytokine production [11,12], and elevated TGF- β secretion by pulmonary epithelial cells from patients with COPD has been observed in comparison to healthy controls [13]. TGF- β signalling is typically through the Smad-dependent pathway [14] although other signalling pathways can also be involved, including the nuclear factor- κ B (NF- κ B) pathway, pathways involving mitogen-activated protein kinases (MAPKs) and the Wingless/integrase-1 (WNT) pathway [15].

In addition to the transcription of mRNA, which forms the first step of the 'central dogma of molecular biology', noncoding RNA are transcribed from DNA [16]. miRNA bind, in a complementary manner, to the 3' end of their target mRNA and instigate their suppression/degradation [16]. Cell proliferation has been shown to be regulated by microRNA-221 (*miR-221*) in ASM cells from patients with asthma [12]; in a human epithelial cell line [17]; and in murine vascular smooth muscle cells (VSMCs) [18]. IL-6 and CXCL8 secretion has also been shown to be attenuated by miRNA; by *miR-221* in ASM cells from patients with asthma [12]; and by *miR-146a* and *miR-146b* in human alveolar

epithelial cells [19,20]. Our previous studies have helped to identify *miR-145* as a potential key regulator of airway smooth muscle function in COPD. Firstly, it is highly expressed in the healthy lung [21] and in healthy ASM cells specifically [22]. It has also been shown to be overexpressed in the airways of patients with cystic fibrosis, and to correlate with a decrease in *SMAD3* expression [23]. A number of human and animal models have linked *miR-145* to mechanisms that could also contribute towards the development of COPD [24,25]. Smooth muscle cell proliferation correlated inversely with expression levels of *miR-145* in murine [26–28], leporine [29] and human [28] vasculatures. Moreover, exposure to cigarette smoke has been shown to affect expression levels of *miR-145* in the lungs of rats [30].

We hypothesized that increased IL-6 and CXCL8 release from the ASM cells of COPD patients is mediated by the TGF- β -induced expression of *miR-145*. We examined the effects of TGF- β upon ASM IL-6 and CXCL8 release from patients with COPD, and in healthy nonsmokers and healthy smokers. We then examined the regulation of *miR-145* with specific kinase inhibitors. Finally, we examined the effects of modulating the expression levels of *miR-145* in these cells on cytokine release and on the phosphorylation of SMAD3. *miR-145* controls the excessive cytokine release observed in ASM cells from patients with COPD, by reducing SMAD3 phosphorylation.

Materials and methods

Primary human ASM cell culture

Primary human ASM cells were previously dissected from the lungs of healthy nonsmokers, healthy smokers and patients with COPD; disease and smoking status were defined according to guidelines produced by the American Thoracic Society [31]. Healthy smokers had a smoking history of at least 10 pack years. There were significant differences between FEV₁ in litres, FEV₁ percent predicted, and FEV₁/FVC ratio between smokers and patients with COPD compared with nonsmokers but matched for age and smoking history (Table 1).

ASM cells were cultured and plated as previously described [11,12,22,32]. ASM cells were plated onto 96-well plates for the measurement of cytokine release, and six well plates for RNA and protein extraction. Confluent cells were growth-arrested by FCS deprivation for 24 h in Dulbecco's Modified Eagle's Medium supplemented with sodium pyruvate (1 mM), L-glutamine (2 mM), nonessential amino acids (1 : 100), penicillin (100 U·mL⁻¹)/streptomycin (100 mg·mL⁻¹), amphotericin B (1.5 mg·mL⁻¹) and BSA (0.1%). Passages 3–4 from nine different donors were used.

Table 1. Patient characteristics.

	Nonsmokers	Smokers	COPD
<i>n</i>	9	9	9
Age (years)	66.4 ± 12.72	59.2 ± 7.6	65.4 ± 6.6
Sex (♂ – ♀)	7 – 2	4 – 5	5 – 5
Pack years smoking	N/A	29.25 ± 3.3	38.32 ± 26.92
FEV ₁ (L)	4.02 ± 0.48	3.12 ± 0.78	1.76 ± 0.45
FEV ₁ (% Predicted)	104.23 ± 7.28	101.5 ± 4.51	77 ± 21.97
FEV ₁ /FVC (%)	78.89 ± 5.98	77.57 ± 3.32	38.88 ± 15.75
PC ₂₀ (mg·mL ⁻¹)	> 16	> 16	Too severe

FEV₁, forced expiratory volume in 1 s; FVC, forced vital capacity; PC₂₀, provocative concentration of methacholine causing a 20% fall in FEV₁. Data shown as mean ± SEM.

Cells were stimulated in triplicate ± TGF-β at the indicated concentrations.

Alternatively, ASM cells were cultured for 1 h in the presence or absence of the indicated concentrations of TPCA-1 (an IKK-2 inhibitor), PD098059 (a MEK-1/2 inhibitor), SP600125 (a JNK-1/2 inhibitor) and SB 203580 (a p38 MAP kinase inhibitor) and then stimulated with 1 ng·mL⁻¹ of TGF-β for 24 h. All inhibitors were obtained from Calbiochem. The supernatants were removed, and IL-6 and CXCL8 levels were determined by DuoSet ELISA (R&D Systems, Abingdon, UK).

miRNA and mRNA Expression

The human (hsa)-*miR-145* and *SMAD3* expression levels were measured as previously described [11,12,22].

Transfection with *miR-145* mimics and controls

ASM cells were transfected as previously described [11,12]. A mimic for *miR-145* and controls were obtained from Ambion/Applied Biosystems, Ltd. (Paisley, UK). Transfected cells were plated into 96-well or 6-well plates, and left to adhere overnight before being serum starved for 6 h before stimulation with 1 ng·mL⁻¹ TGF-β for the indicated times.

Western blotting

Proteins were measured as previously described [12,32,33]. Antibodies against human phospho-S423-S425-Smad3 and total Smad3 were purchased from AbCam (Cambridge, UK).

Data analysis

Data were analysed using GRAPHPAD PRISM, version 5.03 (GraphPad Software, San Diego, CA). Data were not

normally distributed (as assessed by the Kolmogorov–Smirnov test), and therefore groups were compared using the Dunn nonparametric test. All data are expressed as means ± SEMs. Significance was defined as a *P* value of less than 0.05.

Results

The effect of TGF-β stimulation on CXCL8 and IL-6 release and *SMAD3* and *miR-145* expression by ASM cells after 24 h

ASM cells were stimulated with 2.5% FCS and TGF-β at the indicated concentrations (0.001–10 ng·mL⁻¹) for 24 h. TGF-β induced a concentration-dependent increase in CXCL8 and IL-6 release from ASM cells which plateaued at 1 ng·mL⁻¹ in the nonsmokers (*P* < 0.05), smokers (*P* < 0.01) and COPD (*P* < 0.001) cells (Fig. 1A,B). A significant increase in CXCL8 release was observed in the COPD ASM cells compared to the nonsmokers (*P* < 0.01) when the ASM cells were stimulated with 1 ng·mL⁻¹ of TGF-β (Fig. 1A). Furthermore, there was significant increase in IL-6 release between the nonsmokers and smokers (*P* < 0.01), nonsmokers and COPDs (*P* < 0.001), and the smoker and COPD ASM cells (*P* < 0.05) (Fig. 1B).

TGF-β (0.01 ng·mL⁻¹) induced an increase in *SMAD3* expression in ASM cells from COPD patients ~60-fold higher than baseline (*P* < 0.001; Fig. 1C). Relatively little change to *SMAD3* expression was seen in the nonsmokers and smokers compared to unstimulated cells. *miR-145* expression in ASM cells from COPD patients exhibited a concentration dependent increase which plateaued at 1 ng·mL⁻¹ (*P* < 0.01) (Fig. 1D). A significant increase in expression was observed in the COPD ASM cells compared to the nonsmokers and smokers (both *P* < 0.01) (Fig. 1D).

The effects of specific kinase inhibitors on CXCL8 and IL-6 release by ASM cells stimulated with FCS and TGF-β after 24 h

In previous studies, we and others have demonstrated that cytokines can induce activation of IKK2/NF-κB and the MAP kinases, ERK-1/2, JNK-1/2 and p38 MAP kinase in ASM cells and that these are inhibited in the presence of the selective pharmacological inhibitors of these [34–43]. We therefore used the biological active concentrations of these inhibitors to examine the role of the NF-κB and MAP kinases pathways during *miR-145* expression.




Fig. 1. Effect of increasing concentrations of transforming growth factor- β (TGF- β) on airway smooth muscle (ASM) CXCL8 (A) and IL-6 release (B), *SMAD3* (C) and *miR-145* (D) expression from the ASM cells of non-smokers, smokers and patients with COPD at 24 h. Points represent the means \pm SEMs from nine ASM donors in each group. */\$/# $P < 0.05$; **/\$\$/## $P < 0.01$; ***/###/\$\$\$\$ $P < 0.001$. Asterisks indicate comparison with no TGF- β control. Hash signs indicate COPD vs. nonsmoker ASM cells. Dollar signs indicate COPD vs. smoker ASM cells. Plus signs indicate smoker vs. nonsmoker ASM cells.

Following 1 h pre-treatment with inhibitors, ASM cells were stimulated with TGF- β (1 ng-mL $^{-1}$) and the generation of IL-6 (Fig. 2A,D,G,J), CXCL8 (Fig. 2B,E,H,K) and *miR-145* (Fig. 2C,F,I,L) were determined at 24 h. Exposure to TPCA-1 completely inhibited production of IL-6 and CXCL8 in the non-smokers at 10 μ M, and a significant reduction was observed in the COPD ASM cells (both $P < 0.05$) (Fig. 2A,B). No effect was observed upon *miR-145* expression (Fig. 2C). The MEK-1/2 inhibitor (10 μ M) also attenuated IL-6 and CXCL8 production (both $P < 0.05$) (Fig. 2D,E). Interestingly, a significant increase in *miR-145* expression was observed in the COPD ASM cells ($P < 0.05$) (Fig. 2F). Inhibition of the JNK-1/2 kinase demonstrated no effect upon either cytokine release or *miR-145* expression (Fig. 2G,H,I). In contrast, inhibition of the p38 MAP kinase had differential actions upon cytokine and *miR-145* production. Blocking p38 MAP kinase inhibited CXCL8 but not IL-6 in both the non-smoker and COPD ASM cells (Fig. 2J,K), and a significant increase in *miR-145* expression was observed in the COPD ASM cells (Fig. 2L). Overall, pharmacological studies indicate that TGF- β -induced *miR-*

145 expression is regulated via an MEK-1/2- and p38-dependent pathway.

The effect upon TGF- β -stimulated ASM cells of *miR-145* overexpression on CXCL8 and IL-6 release

To clarify the role of *miR-145*, we examined the effect of overexpressing *miR-145* on TGF- β -induced CXCL8 and IL-6 release. Transfection using Amaxa electroporation (Lonza, Slough, UK) showed that *miR-145* mimics (100 nM) inhibited CXCL8 release by approximately 47% ($P < 0.01$) and IL-6 release by approximately 49% ($P < 0.001$), to levels comparably seen in the healthy smokers (Fig. 3A,B). Altering the endogenous levels of *miR-145* exerted no effect in either healthy or smoker ASM cells or those from patients with severe asthma. To confirm efficient transfection, we undertook parallel studies that examined the effects of a small, interfering RNA (100 nM) targeted to *IL-6*. As demonstrated previously [12,41], we showed a reduction in IL-6 release induced by TGF- β stimulation in ASM cells (data not shown), with no effect upon cell viability (data not shown).




Fig. 2. Effect of inhibitors of IKK2 and MAP kinases upon TGF- β -induced IL-6 and CXCL8 release, and *miR-145* expression from the ASM cells of nonsmokers and patients with COPD at 24 h. ASM cells were pretreated for 60 min with the indicated concentrations of the inhibitors of IKK-2 (TPCA-1), MEK-1/2 (PD098059), JNK-1/2 (SP600125) and p38 MAP kinase (SB203580). Following exposure to vehicle control or TGF- β (1 ng·mL⁻¹) for 24 h, the release of IL-6 and CXCL8 was determined by ELISA. *miR-145* expression was measured by RT-PCR. Points represent the means \pm SEM of nine ASM donors in each group. *[#] $P < 0.05$; **[#] $P < 0.01$; ***[#] $P < 0.001$. Asterisks indicate stimulated nonsmoker comparison with no TGF- β control. Hash signs indicate stimulated COPD comparison with no TGF- β control.




Fig. 3. Effects of the overexpression of *miR-145* in the ASM cells of nonsmokers, smokers and patients with COPD at 24 h. ASM cells were electroporated in the presence of buffer, control mimic or *miR-145* mimic. Cells were then exposed to vehicle control or 1 ng·mL⁻¹ TGF-β and the release of CXCL8 (A) and IL-6 (B) was measured by ELISA at 24 h. Furthermore, p-SMAD3 was measured by western blotting (C). Points represent the means ± SEMs from nine ASM donors in each group. **P* < 0.05; **/###/+*P* < 0.01; ***/###/+*P* < 0.001.

Effects of *miR-145* on SMAD3

We next determined whether *miR-145* could regulate SMAD3 phosphorylation. As previously demonstrated [33], TGF-β (1 ng·mL⁻¹) increased SMAD3 phosphorylation in the nonsmoker ASM cells at 24 h (*P* < 0.05) (Fig. 3C). For the first time, we demonstrate that there is a slight increase in phosphorylation in the non-smokers, and an even greater degree of phosphorylation in the ASM cells from COPD patients induced by TGF-β (*P* < 0.001) (Fig. 3C). The *miR-145* mimic (100 nM) decreased the TGF-β-induced phosphorylation of SMAD3 in the COPD ASM cells by approximately 50% (*P* < 0.05) (Fig. 3C), and demonstrated no effect upon either the nonsmoker or smoker ASM cells.

Discussion

We have made several important observations regarding the behaviour of ASM cells from patients with COPD. First, we showed that TGF-β increased both ASM IL-6 and CXCL8 release in the COPD patients to a greater degree than those from the nonsmoker subjects. We also observed a concurrent increase in the expression of both *SMAD3* and *miR-145* in the ASM cells from the COPD patients. We next investigated the mechanisms that regulate the expression of *miR-145*. We showed that expression of *miR-145* is mediated, at least in part, through activation of MEK-1/2 and p38 in ASM cells from COPD patients. Examination of the effect of these MAP kinase inhibitors upon generation of inflammatory mediators showed that

IL-6 release was mediated via IKK2 and MEK-1/2 while CXCL8 release was mediated via IKK2, MEK-1/2 and p38 in both the ASM cohorts, from nonsmokers and those with COPD. Finally, we found that *miR-145* regulates the enhanced IL-6 and CXCL8 release seen in the ASM cells from patients with COPD, that could be partly through the control of *SMAD3* (Summarized in Fig. 4).

Previous observations have described the effect of inducing hyperproliferation of ASM cells with TGF- β in both asthma [11,12] and COPD [44]. This is the first time that TGF- β has been shown to induce both IL-6 and CXCL8 release from primary ASM cells isolated from individuals with COPD. The regulation of smooth muscle cell phenotype has previously been shown to be correlated with expression of *miR-145*. Specifically, *miR-145* has been demonstrated to regulate smooth muscle cell fate [26,45], the contractile phenotype of VSMCs [27,46], and acts a novel VSMC phenotypic marker in murine models [28], and prevent vein graft disease in rabbits [29]. Interestingly, exposure to cigarette smoke has been shown to affect expression levels of *miR-145* in the lungs of rats [30], and has been found to be differentially expressed in lung homogenates in rats with COPD [47]. Furthermore, *miR-145* (along with others) has recently been proposed to be a promising

plasma based biomarker for the diagnosis of COPD [48]. This is the first time that a role for *miR-145* in the ASM cells from COPD patients has been reported. We have shown that expression of *miR-145* is through activation of both MEK-1/2 and p38. Interestingly, Hu *et al.* [49], have reported that activation of MEK-1/2 suppresses *miR-145* expression in VSMCs, and Kent *et al.* [50], demonstrate that *miR-145* expression is inhibited through activation of the MAPK and JNK pathways in colorectal cancer. Furthermore, in cardiomyocytes, the protective activity of *miR-145* is associated with modulation of both MEK-1/2 and JNK [51] and in gastric mucosal epithelial cell regulation of *miR-145* involves JNK [52]. Similar to our results, p38 has previously been shown to be linked to *miR-145* induction in VSMCs [53] and Hong *et al.* [54], suggest that the p38 MAPK signalling pathway promotes miRNA biogenesis by facilitating the nuclear localization of p68. Clearly, the regulation of *miR-145* is, unsurprisingly, cell type specific.

Finally, we examined the effect of increasing *miR-145* expression in ASM cells from COPD patients upon IL-6 and CXCL8 release. Although studies have suggested a correlation between *miR-145* expression and IL-6 & CXCL8 release [55–58], we show for the first time that increasing the expression of




Fig. 4. SMAD3 dependent regulation of IL-6 & CXCL8 release by TGF- β -induced *miR-145* expression. In response to TGF- β stimulation, MEK-1/2 and p38 activation results in increased *miR-145* expression. The concurrent increase in expression and phosphorylation of SMAD3, is regulated by *miR-145* to prevent further generation and release of IL-6 & CXCL8.

miR-145 can reduce the levels of IL-6 & CXCL8 release from the COPD ASM cells to levels comparable to that of the nonsmoker ASM cells. *miR-145* is proposed to target *SMAD3* in systemic sclerosis (SSc) [59], cystic fibrosis [23], cartilage dysfunction [60], nasopharyngeal cancer [61] and in hypertrophic scar fibroblasts (HSFBs) [62], we show here that this may also be the case in ASM cells from patients with COPD.

Interestingly, *miR-145* is also likely to be important in regulating ASM cells in asthma, as it is highly expressed in the healthy lung [21] and in healthy ASM cells specifically [22], and inhibition of *miR-145* inhibits eosinophilic inflammation, mucus hypersecretion, T_H2 cytokine production and airway hyperresponsiveness in house dust mite-induced allergic mouse airways [63].

In conclusion, *miR-145* is vital in controlling the increased inflammatory response of human ASM cells in patients with COPD. This finding may open a new avenue in COPD therapeutics by targeting of *miRNA-145* and diagnosis by its detection.

Acknowledgements

This work was supported by a fellowship from Imperial College to MMP and a grant from The Wellcome Trust (085935) to KFC. It was also supported by the NIHR Respiratory Disease Biomedical Research Unit at the Royal Brompton NHS Foundation Trust and Imperial College London. KFC is a Senior Investigator of NIHR, UK. MMP and KFC are members of Interuniversity Attraction Poles Program-Belgian State-Belgian Science Policy- project P7/30.

Author contributions

MP and LL were responsible for preparation of the manuscript; LL, KS, IP and BT conducted *in-vitro* experiments; KD, AH and KFC provided the primary ASM cells; MP designed the study.

References

- Buist AS, McBurnie MA, Vollmer WM, Gillespie S, Burney P, Mannino DM, Menezes AM, Sullivan SD, Lee TA, Weiss KB *et al.* (2007) International variation in the prevalence of COPD (the BOLD Study): a population-based prevalence study. *Lancet* **370**, 741–750.
- Vestbo J, Hurd SS, Agusti AG, Jones PW, Vogelmeier C, Anzueto A, Barnes PJ, Fabbri LM, Martinez FJ, Nishimura M *et al.* (2013) Global strategy for the diagnosis, management, and prevention of chronic obstructive pulmonary disease: GOLD executive summary. *Am J Respir Crit Care Med* **187**, 347–365.
- Mathers CD and Loncar D (2006) Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med* **3**, e442.
- Barnes PJ (2014) Cellular and molecular mechanisms of chronic obstructive pulmonary disease. *Clin Chest Med* **35**, 71–86.
- Jeffery PK (2004) Remodeling and inflammation of bronchi in asthma and chronic obstructive pulmonary disease. *Proc Am Thorac Soc* **1**, 176–183.
- Trian T, Benard G, Begueret H, Rossignol R, Girodet PO, Ghosh D, Ousova O, Vernejoux JM, Marthan R, Tunon-de-Lara JM *et al.* (2007) Bronchial smooth muscle remodeling involves calcium-dependent enhanced mitochondrial biogenesis in asthma. *J Exp Med* **204**, 3173–3181.
- Hogg JC, Chu F, Utokaparch S, Woods R, Elliott WM, Buzatu L, Cherniack RM, Rogers RM, Sciurba FC, Coxson HO *et al.* (2004) The nature of small-airway obstruction in chronic obstructive pulmonary disease. *N Engl J Med* **350**, 2645–2653.
- Fletcher C and Peto R (1977) The natural history of chronic airflow obstruction. *Br Med J* **1**, 1645–1648.
- Prakash YS (2013) Airway smooth muscle in airway reactivity and remodeling: what have we learned? *Am J Physiol Lung Cell Mol Physiol* **305**, L912–L933.
- Bhowmik A, Seemungal TA, Sapsford RJ and Wedzicha JA (2000) Relation of sputum inflammatory markers to symptoms and lung function changes in COPD exacerbations. *Thorax* **55**, 114–120.
- Perry MM, Durham AL, Austin PJ, Adcock IM and Chung KF (2015) BET bromodomains regulate transforming growth factor-beta-induced proliferation and cytokine release in asthmatic airway smooth muscle. *J Biol Chem* **290**, 9111–9121.
- Perry MM, Baker JE, Gibeon DS, Adcock IM and Chung KF (2013) Airway smooth muscle hyperproliferation is regulated by microRNA-221 in severe asthma. *Am J Respir Cell Mol Biol* **50**, 7–17.
- Takizawa H, Tanaka M, Takami K, Ohtoshi T, Ito K, Satoh M, Okada Y, Yamasawa F, Nakahara K and Umeda A (2001) Increased expression of transforming growth factor-beta1 in small airway epithelium from tobacco smokers and patients with chronic obstructive pulmonary disease (COPD). *Am J Respir Crit Care Med* **163**, 1476–1483.
- Yeganeh B, Mukherjee S, Moir LM, Kumawat K, Kashani HH, Bagchi RA, Baarsma HA, Gosens R and Ghavami S (2013) Novel non-canonical TGF-beta signaling networks: emerging roles in airway smooth muscle phenotype and function. *Pulm Pharmacol Ther* **26**, 50–63.
- Massague J, Seoane J and Wotton D (2005) Smad transcription factors. *Genes Dev* **19**, 2783–2810.

- 16 Bartel DP (2009) MicroRNAs: target recognition and regulatory functions. *Cell* **136**, 215–233.
- 17 Zhang C, Zhang J, Zhang A, Wang Y, Han L, You Y, Pu P and Kang C (2010) PUMA is a novel target of miR-221/222 in human epithelial cancers. *Int J Oncol* **37**, 1621–1626.
- 18 Liu X, Cheng Y, Zhang S, Lin Y, Yang J and Zhang C (2009) A necessary role of miR-221 and miR-222 in vascular smooth muscle cell proliferation and neointimal hyperplasia. *Circ Res* **104**, 476–487.
- 19 Perry MM, Moschos SA, Williams AE, Shepherd NJ, Larner-Svensson HM and Lindsay MA (2008) Rapid changes in microRNA-146a expression negatively regulate the IL-1beta-induced inflammatory response in human lung alveolar epithelial cells. *J Immunol* **180**, 5689–5698.
- 20 Perry MM, Williams AE, Tsitsiou E, Larner-Svensson HM and Lindsay MA (2009) Divergent intracellular pathways regulate interleukin-1beta-induced miR-146a and miR-146b expression and chemokine release in human alveolar epithelial cells. *FEBS Lett* **583**, 3349–3355.
- 21 Williams AE, Larner-Svensson H, Perry MM, Campbell GA, Herrick SE, Adcock IM, Erjefalt JS, Chung KF and Lindsay MA (2009) MicroRNA expression profiling in mild asthmatic human airways and effect of corticosteroid therapy. *PLoS One* **4**, e5889.
- 22 Perry M, Tsitsiou E, Austin P, Lindsay M, Gibeon D, Adcock I and Chung K (2014) Role of non-coding RNAs in maintaining primary airway smooth muscle cells. *Respir Res* **15**, 58.
- 23 Megiorni F, Cialfi S, Cimino G, De Biase RV, Dominici C, Quattrucci S and Pizzuti A (2013) Elevated levels of miR-145 correlate with SMAD3 down-regulation in cystic fibrosis patients. *J Cyst Fibros* **12**, 797–802.
- 24 Perry MM, Adcock IM and Chung KF (2015) Role of microRNAs in allergic asthma: present and future. *Curr Opin Allergy Clin Immunol* **15**, 156–162.
- 25 Brook PO, Perry MM, Adcock IM and Durham AL (2015) Epigenome-modifying tools in asthma. *Epigenomics* **7**, 1017–1032.
- 26 Cordes KR, Sheehy NT, White MP, Berry EC, Morton SU, Muth AN, Lee TH, Miano JM, Ivey KN and Srivastava D (2009) miR-145 and miR-143 regulate smooth muscle cell fate and plasticity. *Nature* **460**, 705–710.
- 27 Boettger T, Beetz N, Kostin S, Schneider J, Kruger M, Hein L and Braun T (2009) Acquisition of the contractile phenotype by murine arterial smooth muscle cells depends on the Mir143/145 gene cluster. *J Clin Invest* **119**, 2634–2647.
- 28 Cheng Y, Liu X, Yang J, Lin Y, Xu DZ, Lu Q, Deitch EA, Huo Y, Delphin ES and Zhang C (2009) MicroRNA-145, a novel smooth muscle cell phenotypic marker and modulator, controls vascular neointimal lesion formation. *Circ Res* **105**, 158–166.
- 29 Ohnaka M, Marui A, Yamahara K, Minakata K, Yamazaki K, Kumagai M, Masumoto H, Tanaka S, Ikeda T and Sakata R (2014) Effect of microRNA-145 to prevent vein graft disease in rabbits by regulation of smooth muscle cell phenotype. *J Thorac Cardiovasc Surg* **148**, 676–682.
- 30 Izzotti A, Calin GA, Arrigo P, Steele VE, Croce CM and De FS (2009) Downregulation of microRNA expression in the lungs of rats exposed to cigarette smoke. *FASEB J* **23**, 806–812.
- 31 Standards for the diagnosis and care of patients with chronic obstructive pulmonary disease. American Thoracic Society. *Am J Respir Crit Care Med* **152**, S77–S121.
- 32 Perry MM, Hui CK, Whiteman M, Wood ME, Adcock I, Kirkham P, Michaeloudes C and Chung KF (2011) Hydrogen sulfide inhibits proliferation and release of IL-8 from human airway smooth muscle cells. *Am J Respir Cell Mol Biol* **45**, 746–752.
- 33 Michaeloudes C, Sukkar MB, Khorasani NM, Bhavsar PK and Chung KF (2010) TGF-beta regulates Nox4, MnSOD and catalase expression and IL-6 release in airway smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* **300**, L295–L304.
- 34 Wuyts WA, Vanaudenaerde BM, Dupont LJ, Demedts MG and Verleden GM (2003) Involvement of p38 MAPK, JNK, p42/p44 ERK and NF-kappaB in IL-1beta-induced chemokine release in human airway smooth muscle cells. *Respir Med* **97**, 811–817.
- 35 Oltmanns U, Issa R, Sukkar MB, John M and Chung KF (2003) Role of c-jun N-terminal kinase in the induced release of GM-CSF, RANTES and IL-8 from human airway smooth muscle cells. *Br J Pharmacol* **139**, 1228–1234.
- 36 Wang CC, Lin WN, Lee CW, Lin CC, Luo SF, Wang JS and Yang CM (2005) Involvement of p42/p44 MAPK, p38 MAPK, JNK, and NF-kappaB in IL-1beta-induced VCAM-1 expression in human tracheal smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* **288**, L227–L237.
- 37 Laporte JD, Moore PE, Lahiri T, Schwartzman IN, Panettieri RA Jr and Shore SA (2000) p38 MAP kinase regulates IL-1 beta responses in cultured airway smooth muscle cells. *Am J Physiol Lung Cell Mol Physiol* **279**, L932–L941.
- 38 Laporte JD, Moore PE, Abraham JH, Maksym GN, Fabry B, Panettieri RA Jr and Shore SA (1999) Role of ERK MAP kinases in responses of cultured human airway smooth muscle cells to IL-1beta. *Am J Physiol* **277**, L943–L951.
- 39 Hedges JC, Singer CA and Gerthoffer WT (2000) Mitogen-activated protein kinases regulate cytokine

- gene expression in human airway myocytes. *Am J Respir Cell Mol Biol* **23**, 86–94.
- 40 Hallsworth MP, Moir LM, Lai D and Hirst SJ (2001) Inhibitors of mitogen-activated protein kinases differentially regulate eosinophil-activating cytokine release from human airway smooth muscle. *Am J Respir Crit Care Med* **164**, 688–697.
 - 41 Larner-Svensson HM, Williams AE, Tsitsiou E, Perry MM, Jiang X, Chung KF and Lindsay MA (2010) Pharmacological studies of the mechanism and function of interleukin-1beta-induced miRNA-146a expression in primary human airway smooth muscle. *Respir Res* **11**, 68.
 - 42 Issa R, Xie S, Lee KY, Stanbridge RD, Bhavsar P, Sukkar MB and Chung KF (2006) GRO-alpha regulation in airway smooth muscle by IL-1beta and TNF-alpha: role of NF-kappaB and MAP kinases. *Am J Physiol Lung Cell Mol Physiol* **291**, L66–L74.
 - 43 Shan L, Redhu NS, Saleh A, Halayko AJ, Chakir J and Gounni AS (2010) Thymic stromal lymphopoietin receptor-mediated IL-6 and CC/CXC chemokines expression in human airway smooth muscle cells: role of MAPKs (ERK1/2, p38, and JNK) and STAT3 pathways. *J Immunol* **184**, 7134–7143.
 - 44 Wiegman CH, Michaeloudes C, Haji G, Narang P, Clarke CJ, Russell KE, Bao W, Pavlidis S, Barnes PJ, Kanerva J *et al.* (2015) Oxidative stress-induced mitochondrial dysfunction drives inflammation and airway smooth muscle remodeling in patients with chronic obstructive pulmonary disease. *J Allergy Clin Immunol* **136**, 769–780.
 - 45 Yamaguchi S, Yamahara K, Homma K, Suzuki S, Fujii S, Morizane R, Monkawa T, Matsuzaki Y, Kangawa K and Itoh H (2011) The role of microRNA-145 in human embryonic stem cell differentiation into vascular cells. *Atherosclerosis* **219**, 468–474.
 - 46 Hutcheson R, Terry R, Chaplin J, Smith E, Musiyenko A, Russell JC, Lincoln T and Rocic P (2013) MicroRNA-145 restores contractile vascular smooth muscle phenotype and coronary collateral growth in the metabolic syndrome. *Arterioscler Thromb Vasc Biol* **33**, 727–736.
 - 47 Li B, Zhou X, Chen L, Feng C and Li T (2014) Expression of microRNAs in lung homogenates in rats with chronic obstructive pulmonary disease. *Zhonghua Wei Zhong Bing Ji Jiu Yi Xue* **26**, 905–909.
 - 48 Wang M, Huang Y, Liang Z, Liu D, Lu Y, Dai Y, Feng G and Wang C (2016) Plasma miRNAs might be promising biomarkers of chronic obstructive pulmonary disease. *Clin Respir J* **10**, 104–111.
 - 49 Hu B, Song JT, Qu HY, Bi CL, Huang XZ, Liu XX and Zhang M (2014) Mechanical stretch suppresses microRNA-145 expression by activating extracellular signal-regulated kinase 1/2 and upregulating angiotensin-converting enzyme to alter vascular smooth muscle cell phenotype. *PLoS One* **9**, e96338.
 - 50 Kent OA, Fox-Talbot K and Halushka MK (2013) RREB1 repressed miR-143/145 modulates KRAS signaling through downregulation of multiple targets. *Oncogene* **32**, 2576–2585.
 - 51 Li R, Yan G, Zhang Q, Jiang Y, Sun H, Hu Y, Sun J and Xu B (2013) miR-145 inhibits isoproterenol-induced cardiomyocyte hypertrophy by targeting the expression and localization of GATA6. *FEBS Lett* **587**, 1754–1761.
 - 52 Luo XJ, Liu B, Dai Z, Li TB, Li NS, Zhang XJ, Yang ZC, Li YJ and Peng J (2013) Expression of apoptosis-associated microRNAs in ethanol-induced acute gastric mucosal injury via JNK pathway. *Alcohol* **47**, 481–493.
 - 53 Blumensatt M, Wronkowitz N, Wiza C, Cramer A, Mueller H, Rabelink MJ, Hoeber RC, Eckel J, Sell H and Ouwens DM (2014) Adipocyte-derived factors impair insulin signaling in differentiated human vascular smooth muscle cells via the upregulation of miR-143. *Biochim Biophys Acta* **1842**, 275–283.
 - 54 Hong S, Noh H, Chen H, Padia R, Pan ZK, Su SB, Jing Q, Ding HF and Huang S (2013) Signaling by p38 MAPK stimulates nuclear localization of the microprocessor component p68 for processing of selected primary microRNAs. *Sci Signal* **6**, ra16.
 - 55 Oliva EN, Cuzzola M, Aloe Spiriti MA, Poloni A, Lagana C, Rigolino C, Morabito F, Galimberti S, Ghio R, Cortelezzi A *et al.* (2013) Biological activity of lenalidomide in myelodysplastic syndromes with del5q: results of gene expression profiling from a multicenter phase II study. *Ann Hematol* **92**, 25–32.
 - 56 Yu CC, Tsai LL, Wang ML, Yu CH, Lo WL, Chang YC, Chiou GY, Chou MY and Chiou SH (2013) miR145 targets the SOX9/ADAM17 axis to inhibit tumor-initiating cells and IL-6-mediated paracrine effects in head and neck cancer. *Cancer Res* **73**, 3425–3440.
 - 57 Li W, Ma K, Zhang S, Zhang H, Liu J, Wang X and Li S (2015) Pulmonary microRNA expression profiling in an immature piglet model of cardiopulmonary bypass-induced acute lung injury. *Artif Organs* **39**, 327–335.
 - 58 Zhou J, Chaudhry H, Zhong Y, Ali MM, Perkins LA, Owens WB, Morales JE, McGuire FR, Zumbun EE, Zhang J *et al.* (2015) Dysregulation in microRNA expression in peripheral blood mononuclear cells of sepsis patients is associated with immunopathology. *Cytokine* **71**, 89–100.
 - 59 Zhu H, Li Y, Qu S, Luo H, Zhou Y, Wang Y, Zhao H, You Y, Xiao X and Zuo X (2012) MicroRNA expression abnormalities in limited cutaneous scleroderma and diffuse cutaneous scleroderma. *J Clin Immunol* **32**, 514–522.
 - 60 Yang B, Kang X, Xing Y, Dou C, Kang F, Li J, Quan Y and Dong S (2014) Effect of microRNA-145 on IL-1beta-induced cartilage degradation in human chondrocytes. *FEBS Lett* **588**, 2344–2352.

- 61 Huang H, Sun P, Lei Z, Li M, Wang Y, Zhang HT and Liu J (2015) miR-145 inhibits invasion and metastasis by directly targeting Smad3 in nasopharyngeal cancer. *Tumour Biol* **36**, 4123–4131.
- 62 Zhu HY, Li C, Zheng Z, Zhou Q, Guan H, Su LL, Han JT, Zhu XX, Wang SY, Li J *et al.* (2015) Peroxisome proliferator-activated receptor-gamma (PPAR-gamma) agonist inhibits collagen synthesis in human hypertrophic scar fibroblasts by targeting Smad3 via miR-145. *Biochem Biophys Res Commun* **459**, 49–53.
- 63 Collison A, Mattes J, Plank M and Foster PS (2011) Inhibition of house dust mite-induced allergic airways disease by antagonism of microRNA-145 is comparable to glucocorticoid treatment. *J Allergy Clin Immunol* **128**, 160–167.