

The impact of phylogenetic dating method on interpreting trait evolution: a case study of  
Cretaceous-Palaeogene eutherian body size evolution

Halliday, T.J.D\*<sup>1</sup>, and Goswami, A.<sup>1,2</sup>

1 – Department of Genetics, Evolution, and Environment, UCL, Gower Street, London,  
WC1E 6BT

2 – Department of Earth Sciences, UCL, Gower Street, London, WC1E 6BT

\*TJDH is the corresponding author – [thomas.halliday.11@ucl.ac.uk](mailto:thomas.halliday.11@ucl.ac.uk)

## **Abstract**

The fossil record of the earliest Cenozoic contains the first large-bodied placental mammals. Several evolutionary models have been invoked to explain the transition from small to large body sizes, but methods for determining evolutionary mode of trait change depend on input from tree topology and divergence dates. Different dating methods may therefore affect inference of evolutionary model. Here, we fit models of body mass evolution onto dated phylogenies of Cretaceous and Palaeogene mammals, comparing the effect of dating method on interpretation of evolutionary model. Among traditional palaeontological dating approaches, an OU model with high alpha parameters is recovered as best-fitting when minimum-age dating is used, while branch-sharing methods are highly sensitive to topology. Release or release-radiate models are preferred when Bayesian fossilised birth-death methods are used, but when using stochastic cal3 dating of trees, a model of increased evolutionary rate without a release in constraint at the K-Pg boundary has highest support. These results demonstrate unambiguously that choice of dating method is critical for interpretation of continuous trait evolution, and that care must therefore be taken to consider these effects in macroevolutionary studies.

## **Keywords**

Mammalia, Body Size, Evolutionary Model, Tempo, Mode

## **Background**

Ancestral trait reconstruction and the fitting of models of trait evolution to phylogenies is commonplace in evolutionary biology (1). To understand how clades developed key traits or responded to environmental changes, such approaches are extremely useful. However, as most methods for reconstructing trait evolution require a dated phylogeny (2, 3), conclusions

may be heavily influenced by variations in tree topology, taxonomic sample, or reconstructed divergence dates.

Mammalian body mass has received much attention in macroevolutionary studies, particularly across the Cretaceous-Palaeogene (K-Pg) boundary (4-7). The fossil record suggests that mean body mass of eutherians (7, 8) and overall rate of morphological evolution (9) increased over the K-Pg boundary. However, body mass reconstructions onto different eutherian phylogenies have found either stasis (5) or a reduction in rate of body size evolution combined with a release of constraints (6) at 66 Ma.

Several methods exist for dating divergences in fossil-based phylogenies. Fossil ages provide minimum estimates for dating clades including that taxon, but some analyses have directly used fossil ages as divergence dates (10), a potentially problematic approach (11). Alternative “branch-sharing” methods divide ancestral branch lengths proportionally to the amount of morphological change (12) or equally (13) among descendant branches. Finally, the rate-calibrated cal3 approach (14), which under some conditions better approximates true dates in simulations (15), and Bayesian approaches (e.g. 16) use parametric models of diversification and sampling to derive divergence dates. If dating methods systematically result in different patterns of divergence, this would be expected to impact inference of models of continuous trait evolution due to differences in phylogenetic variance-covariance structure across the tree. We assess the impact of different methods for estimating divergence dates on support for a range of evolutionary models of placental body size evolution.

## **Methods**

### *Mass Estimation*

Body masses were estimated using dental, cranial, and postcranial proxies for 177 eutherian genera (Fig. S1, File S1) sampled in a recent phylogenetic analysis of Cretaceous-Palaeogene taxa (17), sampling 76% of known Cretaceous and 66% of Paleocene eutherian families, densely sampling eutherian diversity around the K-Pg boundary. Standard deviation on body size was assumed to be 0.15, following previously published estimates (18).

### *Dating*

Four methods were used to date six previously published sets of phylogenies derived from different topological constraints (17). The first method used fossil occurrences as minimum clade ages. Second, the “equal” branch-sharing method (13) was applied. This approach uses minimum-age dating, but, where zero-length branches exist, “shares” the duration of the ancestral branch with the daughter branches, preventing simultaneous occurrence of nested branching events. Third, trees dated using “cal3” – a stochastic method incorporating estimates of sampling, speciation, and extinction rate (14) – were used. Fourth, a fossilised birth-death (FBD) model was used to date the phylogeny in MRBAYES v.3.2.6 (19) in which the topology was fixed, but allowing free resolution of polytomies. Branch rates were drawn from a gamma distribution, with tip ages calibrated based on fossil occurrences (Files S2-4).

Taxon first occurrences were taken from the literature and [www.paleobiodb.org](http://www.paleobiodb.org) (accessed 03/11/15) and divided into stage and North American Land Mammal Age time bins. Tip-dates were assigned randomly and uniformly within the earliest bin in which that taxon occurred for minimum-age and branch-sharing dating; for the FBD, the same uniform distributions formed tip date priors. Models were fitted to a sample of 50 resolved, dated trees from each pattern of constraint and dating method; these were the 50 trees with the highest posterior probability in the FBD, and a random sample of trees for other methods.

## *Modelling Evolution*

We used the *fitContinuous\_paleo* function (6) to fit seven macroevolutionary models to comparative datasets spanning all dating approaches. Akaike weights (20) were used to assess relative support, in each case comparing between Brownian motion (BM), Ornstein-Uhlenbeck (OU), white noise, and trend models, as well as three models incorporating changes in parameters at a fixed point in time – here, 66 million years ago. These three models are the “rate-shift”, which assumes BM before and after the shift point, with rates free to vary before and after, “release”, which assumes an OU model before the shift point, and BM with an identical rate parameter after, and “release-radiate” model, which is the “release” model with rates free to vary before and after the shift point (6).

## **Results**

The minimum-age method produced multiple zero-length branches, and dating using cal3 resulted in multiple short, non-zero-length branches concentrated near the K-Pg boundary. The FBD approach resulted in trees with considerably older divergence dates, with long branches at the tips. Branch-sharing spread internal nodes more evenly through time than other methods (Fig. 1, Table S1).

Support for each evolutionary model was strongly influenced by the dating method (Fig. 2, File S5). OU models received greatest support under minimum-age dating, while the best-fitting models for topologies dated using the “equal” method depended on the topology, and included rate-shift, release-radiate, and OU. Similarly, FBD dating resulted in greatest support for a release model, but also release-radiate and OU models. The most strongly

supported model for cal3-dated trees was a tenfold increase in rate at the K-Pg boundary (Table S2).

## Discussion

Different evolutionary models affect our understanding of the effect of the end-Cretaceous mass extinction on eutherian evolution. Models without a shift component like the OU model here best supported by the data when dating trees with minimum-age approaches imply no change in body size evolution at the K-Pg boundary. Similar results have previously been interpreted to show little effect of the mass extinction on mammal evolution (5, 21). The rate-shift and release-radiate models preferred by cal3, FBD, or branch-sharing dating methods imply a change in tempo, and perhaps mode, in body size evolution, and thus a significant effect of the extinction on subsequent mammal evolution.

These patterns may not hold for other datasets, and simulations would establish which dating method produces trees that better fit real evolutionary scenarios. Some discussion of the merits of each is possible in this specific example.

Some authors have noted a tendency to erroneously recover strong support for OU models in simulated data where the true model is unbounded BM (22). However, these effects are strongest in small trees, and are indicated by low alpha parameters, neither of which occurs here. The OU models preferred by minimum-age dating in this study have high alpha values, indicating a strong pull towards the optimum – here, the root state, reconstructed as approximately the size of the tree-shrew *Tupaia* – and, as a result, low variance. Highly-nested early Paleocene taxa such as *Periptychus* in the dataset, the placental diversification is reconstructed as the simultaneous emergence of several lineages across the majority of the

diversity of Placentalia; most lineages include taxa of approximately this size. Moreover, the sampling of extant taxa is limited to small afrotheres, xenarthrans, euarchontans (including *Tupaia*) and eulipotyphlans, most of which are small in size, and which might appear as an ultimate optimum size. Testing with an expanded sample of extant taxa might alter this observed pattern.

Preference between evolutionary models for “equal” dated trees is strongly influenced by topology, resulting in well-supported but contradictory models. The problem of early but highly-nested taxa manifests itself differently here, as the duration of an ancestral branch must be shared among more zero-length descendant branches than if phylogenetic position and stratigraphy were better correlated. This branch-sharing results in older divergence dates for large clades, affecting the timescales over which character evolution is reconstructed. Differences in topology, especially for key early taxa, have a relatively strong impact on choice of different models under branch-sharing dating compared with other dating methods. To some extent, the models best fitting the data when using FBD dating were also varied, but most often the model with greatest support was one of a change in mode of evolution, with or without a shift in rates.

The rate-shift models that had greatest support when using cal3-dated trees reconstruct a large increase in evolutionary rate at the K-Pg boundary, contrary to the findings of Slater (6) of a reduction in rate but a shift in mode. This difference is perhaps a result of the focus in this dataset on Palaeogene taxa. With a large number of taxa that are close in time to one another and passing through multiple time bins, the estimated sampling and speciation rates are high, leading to a concentration of short branches around the K-Pg boundary. With shorter branches, similar changes in body mass must occur over shorter timescales, resulting

in increased rates around the end-Cretaceous mass extinction. As our sampling of the Cretaceous and Paleocene fossil record is relatively even and far more complete than previous studies, it allows the reconstruction of hypothesised rate changes during this interval, but the caveat must be introduced that as the dataset still substantially undersamples the Oligocene and Neogene, and if a change in mode did occur at the end-Cretaceous mass extinction, it is conceivable that the Palaeogene represents only the initial phase of the new evolutionary model. Increased sampling of later Cenozoic taxa might help place any observed transitions in greater context, and would help to identify any later transitions.

The dates reconstructed by the different dating methods vary considerably and imply very divergent evolutionary histories for mammals based on the same observed pattern. The assumptions about the quality of the fossil record and the nature of speciation that underlie each choice of dating method colour our subsequent interpretation of trait evolution.

Describing the history of mammal evolution in the context of a release in constraint or a shift in rates (or both) at the end-Cretaceous mass extinction tells very different stories. The former suggests some form of ecological interaction between mammals and some clade which became extinct at 66 Ma; the latter alone some intrinsic key adaptation that allowed larger body sizes. Interpreting a single-optimum OU model for the whole of eutherian evolution requires alternative explanations for increased body size in the early Palaeogene. The results of this study strongly demonstrate the importance of being explicit with one's assumptions when dating phylogenies, and the need for continued effort in the development of robust methods for dating trees of fossil organisms.

### **Data accessibility**

All data from this study is deposited at Dryad at the following address:

<http://dx.doi.org/10.5061/dryad.7tp7f>.

### **Competing interests**

We have no competing interests.

### **Author contributions**

TJDH collected the data and conducted the analyses. TJDH and AG conceived the study, wrote the manuscript, approved the publication, and agree to be accountable for the work herein.

### **Acknowledgements**

We thank Drs Bapst, Lloyd, Matzke, and Wright for inviting this contribution. We thank Dr Slater and two anonymous reviewers for their comments, and members of UCL ADaPTiVE group and UCL-NHM-IC palaeobiology journal club for discussions.

### **Funding**

Funding was provided by Leverhulme Grant 2014-364 to AG and TJDH.

### **References**

1. Goswami A, Mannion PD, Benton MJ. Radiation and extinction: investigating clade dynamics in deep time. *Biological Journal of the Linnean Society*. 2016.
2. O'Meara BC, Ane C, Sanderson MJ, Wainwright PC. Testing for different rates of continuous trait evolution using likelihood. *Evolution*. 2006;60(5):922-33.
3. Slater GJ, Harmon LJ, Wegmann D, Joyce P, Revell LJ, Alfaro ME. FITTING MODELS OF CONTINUOUS TRAIT EVOLUTION TO INCOMPLETELY SAMPLED COMPARATIVE DATA USING APPROXIMATE BAYESIAN COMPUTATION. *Evolution*. 2012;66(3):752-62.
4. Alroy J. Cope's rule and the dynamics of body mass evolution in North American fossil mammals. *Science*. 1998;280(5364):731-4.

5. Venditti C, Meade A, Pagel M. Multiple routes to mammalian diversity. *Nature*. 2011;479(7373):393-6.
6. Slater GJ. Phylogenetic evidence for a shift in the mode of mammalian body size evolution at the Cretaceous-Palaeogene boundary. *Methods in Ecology and Evolution*. 2013;4(8):734-44.
7. Alroy J. The fossil record of North American mammals: Evidence for a Paleocene evolutionary radiation. *Systematic Biology*. 1999;48(1):107-18.
8. Smith FA, Boyer AG, Brown JH, Costa DP, Dayan T, Ernest SKM, et al. The Evolution of Maximum Body Size of Terrestrial Mammals. *Science*. 2010;330(6008):1216-9.
9. Halliday TJD, Upchurch P, Goswami A. Eutherians experienced elevated evolutionary rates in the immediate aftermath of the K-Pg mass extinction. *Proceedings of the Royal Society B*. review.
10. O'Leary MA, Bloch JI, Flynn JJ, Gaudin TJ, Giallombardo A, Giannini NP, et al. The Placental Mammal Ancestor and the Post-K-Pg Radiation of Placentals. *Science*. 2013;339(6120):662-7.
11. dos Reis M, Donoghue PCJ, Yang Z. Neither phylogenomic nor palaeontological data support a Palaeogene origin of placental mammals. *Biology Letters*. 2014;10(1):20131003.
12. Ruta M, Wagner PJ, Coates MI. Evolutionary patterns in early tetrapods. I. Rapid initial diversification followed by decrease in rates of character change. *Proceedings of the Royal Society B-Biological Sciences*. 2006;273(1598):2107-11.
13. Brusatte SL, Benton MJ, Ruta M, Lloyd GT. Superiority, competition, and opportunism in the evolutionary radiation of dinosaurs. *Science*. 2008;321(5895):1485-8.
14. Bapst DW. A stochastic rate-calibrated method for time-scaling phylogenies of fossil taxa. *Methods in Ecology and Evolution*. 2013;4(8):724-33.
15. Bapst DW. Assessing the effect of time-scaling methods on phylogeny-based analyses in the fossil record. *Paleobiology*. 2014;40(3):331-51.
16. Fanti F, Cau A, Cantelli L, Hassine M, Audatore M. New Information on *Tataouinea hannibal* from the Early Cretaceous of Tunisia and Implications for the Tempo and Mode of Rebbachisaurid Sauropod Evolution. *Plos One*. 2015;10(4).
17. Halliday TJD, Upchurch P, Goswami A. Resolving the relationships of Paleocene placental mammals. *Biological Reviews*. 2015.
18. Evans AR, Jones D, Boyer AG, Brown JH, Costa DP, Ernest SKM, et al. The maximum rate of mammal evolution. *Proceedings of the National Academy of Sciences of the United States of America*. 2012;109(11):4187-90.
19. Ronquist F, Huelsenbeck JP. MRBAYES 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics*. 2003;19:1572-4.
20. Akaike H. A new look at the statistical model identification. *IEEE Transactions on Automatic Control*. 1974;19:716-23.
21. Bininda-Emonds ORP, Cardillo M, Jones KE, MacPhee RDE, Beck RMD, Grenyer R, et al. The delayed rise of present-day mammals. *Nature*. 2007;446(7135):507-12.
22. Cooper N, Thomas GH, Venditti C, Meade A, Freckleton RP. A cautionary note on the use of Ornstein Uhlenbeck models in macroevolutionary studies. *Biological Journal of the Linnean Society*. 2015.

## Figure Captions

Figure 1 – Exemplar dated phylogenies from each of four dating methods. In each case, the topology is derived from the discretised dataset of Halliday *et al.* (17) under the “full”

constraint. A: Minimum-age dating contains multiple polytomies due to taxa that are both early and deeply nested. B: 'equal' branch-sharing causes internal nodes to be much earlier, and divides the long temporal gap between the ancestors of the early Cretaceous stem-eutherians and the crown group evenly. C: Dating using cal3 results in ages intermediate between A and B, including several very short (but non-zero) branches near the diversification of major clades. D: FBD dating gives relatively long terminal branches, with most diversification occurring earlier in the tree.

Figure 2 – Akaike weights for models of body mass evolution reconstructed on trees dated using minimum-age, branch-sharing, cal3, and Bayesian methods. Each barplot shows the Akaike weights for models reconstructed on all trees, divided into sets based on their constraints (see Halliday *et al.* 2015 for details of those constraints). When using minimum-age dating, OU models best fit the data; under branch-sharing methods, release-radiate models have greatest support, with rate-shift models also fitting the data well. Under cal3 dating, rate-shift models have greatest support, while the best fitted models under FBD dating are trend models.