

PEDIATRICS®

Short malnourished children do not gain excessive fat with food supplementation

Journal:	<i>Pediatrics</i>
Manuscript ID	2018-0679.R1
Article Type:	Regular Article
Date Submitted by the Author:	n/a
Complete List of Authors:	<p>Fabiansen, Christian; Kobenhavns Universitet Det Natur- og Biovidenskabelige Fakultet, Department of Nutrition, Sports and Exercise Phelan, Kevin; ALIMA, ALIMA, Route de l'Aéroport, Rue NG 96 BP: 15530. Dakar, Sénégal</p> <p>Cichon, Bernardette; Kobenhavns Universitet Det Natur- og Biovidenskabelige Fakultet, Department of Nutrition, Sports and Exercise Yaméogo, Charles; Département Biomédical et Santé Publique, Institut de Recherche en Sciences de la Santé, 03 BP 7047 Ouagadougou 03, Burkina Faso , Département Biomédical et Santé Publique</p> <p>Iuel-Brockdorff, Ann-Sophie; Kobenhavns Universitet Det Natur- og Biovidenskabelige Fakultet, Department of Nutrition, Sports and Exercise Kurpad, Anura; St John's Research Institute, Of Nutrition</p> <p>Wells, Jonathan; MRC Childhood Nutrition Research Centre, Nutrition Ritz, Christian; Kobenhavns Universitet, Department of Nutrition, Exercise and Sports</p> <p>Filteau, Suzanne; London School of Hygiene and Tropical Medicine, Epidemiology and Population Health</p> <p>Briend, Andre; Kobenhavns Universitet Det Natur- og Biovidenskabelige Fakultet, Department of Nutrition, Sports and Exercise</p> <p>Christensen, Vibeke; MSF Denmark, Msf Denmark</p> <p>Michaelsen, Kim; University of Copenhagen, Department of Nutrition, Exercise and Sports</p> <p>Shepherd, Susan; ALIMA, ALIMA, Route de l'Aéroport, Rue NG 96 BP: 15530. Dakar, Sénégal</p> <p>Friis, Henrik; University of Copenhagen, Department of Nutrition, Exercise and Sports</p>
Keyword/Topic:	Nutrition

Short malnourished children do not gain excessive fat with food supplementation

Christian Fabiansen^{a,b}, MD, PhD, Kevin P.Q. Phelan^c, MSc, Bernardette Cichon^{a,b}, PhD, Charles W Yaméogo^{a,d}, MSc, Ann-Sophie Iuel-Brockdorff^{a,b}, PhD, Anura Kurpad^e, MD, PhD, Jonathan C Wells^f, PhD, Christian Ritz^a, PhD, Suzanne Filteau^g, PhD, André Briend^{a,h}, MD, PhD, Vibeke B Christensen^{a,i}, MD, DMSc, Kim F Michaelsen^a, MD, DMSc, Susan Shepherd^c, MD, Henrik Friis^a, MD, PhD

Affiliations:

^aDepartment of Nutrition, Exercise and Sports, University of Copenhagen, Rolighedsvej 30, DK-1958 Frederiksberg C, Denmark

^bMédecins Sans Frontières - Denmark, Dronningensgade 68, 3, 1420 Copenhagen, Denmark

^cALIMA, Route de l'Aéroport, Rue NG 96 BP: 15530. Dakar, Sénégal

^dDépartement Biomédical et Santé Publique, Institut de Recherche en Sciences de la Santé, 03 BP 7047 Ouagadougou 03, Burkina Faso

^eDivision of Nutrition, St John's Research Institute, Opp. BDA Complex, Koramangala, Bangalore – 560 034

^fChildhood Nutrition Research Centre, UCL Great Ormond Street Institute of Child Health, 30 Guilford Street, London WC1N 1EH, UK

^gLondon School of Hygiene and Tropical Medicine, Faculty of Epidemiology and Population Health; Keppel Street, London, WC1E 7HT

^hCenter for Child Health Research, University of Tampere School of Medicine and Tampere University Hospital, Lääkärintie 1, 33014 University of Tampere, Finland

ⁱDepartment of Paediatrics, Rigshospitalet, Blegdamsvej 9, 2100, Copenhagen, Denmark

Address correspondence to:

Dr. Christian Fabiansen
Department of Nutrition, Sports and Exercise
University of Copenhagen,
Rolighedsvej 25,
1958 Frederiksberg C, Denmark
Email chfa@nexs.ku.dk
Reprints will not be available from the author

Short title: Short malnourished children do not gain excessive fat

Funding Source: The study was funded by Danish International Development Assistance (09-097LIFE) (KFM); Médecins Sans Frontières (Denmark, Norway); Arvid Nilsson's Foundation; The World Food Program, which was part of a donation to the World Food Program from the American people through the support of the US Agency for International Development's Office of Food for

1
2
3
4 Peace; the Alliance for International Medical Action; and the European Union's humanitarian aid
5 funds, in partnership with Action Contre la Faim. This document covers humanitarian aid activities
6 implemented with the financial assistance of the European Union. The views expressed herein
7 should not be taken, in any way, to reflect the official opinion of the European Union, and the
8 European Commission is not responsible for any use that can be made of the information it
9 contains. The funders had no role in study design, data collection and analysis, decision to publish,
10 or preparation of the manuscript.
11

12 **Financial Disclosure:** The authors of this manuscript have the following competing interests: KFM
13 has received research grants from US Dairy Export Council and the Danish Dairy Research
14 Foundation, and also has research collaboration with Nutriset, a producer of LNS products, and
15 patent owner; HF has received research grants from ARLA Food for Health Centre, and also has
16 research collaboration with Nutriset, a producer of LNS products, and patent owner; AB was the
17 inventor of LNS, for which Nutriset has the patent, but abandoned claims to royalties in 2003. Other
18 authors declare no financial relationships with any organisations that might have an interest in the
19 submitted work in the previous three years, and declare no other relationships or activities that
20 could appear to have influenced the submitted work.
21

22 **Potential Conflict of Interest:** The authors have no conflicts of interest relevant to this article to
23 disclose
24

25 **Clinical Trial Registration:** The trial is registered at www.controlled-trials.com
26 (ISRCTN42569496)
27

28 **Abbreviations:** The Alliance for International Medical Action (ALIMA); (CHW); corn-soy blend
29 (CSB); children recruited by MUAC and ≥ 67 cm (LONG); children recruited by MUAC and < 67
30 cm (SHORT); lipid-based nutrient supplements (LNS); Mid-upper arm circumference (MUAC);
31 moderate acute malnutrition (MAM); ready-to-use therapeutic food (RUTF); severe acute
32 malnutrition (SAM); weight-for-height z-score (WHZ)
33
34

35 **Table of Contents Summary**

36 We assess if it is justified to exclude shorter children with moderate acute malnutrition from
37 treatment out of fear they will accumulate too much fat
38

39 **What's Known on This Subject**

40 It is policy for national malnutrition programs in many African countries to exclude shorter children
41 with moderate acute malnutrition (MAM) from treatment due to concerns they will accumulate too
42 much fat thereby putting them at later risk of non-communicable diseases
43

44 **What This Study Adds**

45 Our study shows that shorter children with low MUAC do not gain excessive fat during
46 supplementation and thus should be offered treatment. The WHO should integrate this evidence into
47 its recently updated Integrated Management of childhood Illness (IMCI) recommendations
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 **Contributors' Statement Page**

5
6 Dr Fabiansen conceptualized the research, conducted the research, acquired the data, performed
7 statistical analysis and wrote the first draft of the manuscript. Mr Phelan, Drs Kurpard, Wells,
8 Filteau, Briend, Christensen and Drs Cichon, Iuel-Brockdorff and Mr Yaméogo conducted the
9 research and acquired the data. Dr Ritz conceptualized the research and design and performed
10 statistical analysis and interpretation of data. Dr Shepherd conceptualized the research and
11 contributed to data interpretation. Drs Michaelsen and Friis conceptualized the research and design
12 and contributed to the interpretation of data. All authors critically revised the article for important
13 intellectual content, and approved the final manuscript as submitted and agree to be accountable for
14 all aspects of the work.
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Review Copy

ABSTRACT

Background: In moderate acute malnutrition (MAM) programs, it is common practice not to measure mid-upper arm circumference (MUAC) of children whose length is <67 cm. This is based on expert opinion that supplementation of shorter children with low MUAC and weight-for-height z-score (WHZ) ≥ -2 may increase risk of excessive fat accumulation. Our aim was to assess if shorter children gain more fat than taller children when treated for MAM diagnosed by low MUAC alone.

Methods: This observational study included children aged 6-23 months with MUAC between 115-125 mm and WHZ ≥ -2 . Based on length at admission, children were categorized as SHORT if <67 cm and LONG if ≥ 67 cm. Linear mixed-effects models were used to assess body composition based on deuterium dilution and skinfold-thickness.

Results: Following 12 weeks of supplementation, there was no difference in change in fat mass index (-0.038kg/m^2 , 95%CI $-0.257;0.181$, $p=0.74$) or fat-free mass index (0.061kg/m^2 , 95%CI $-0.150;0.271$; $p=0.57$) in SHORT vs LONG. In absolute terms, the SHORT children gained both less fat-free mass (-230g , 95%CI: $-355;-106$, $P<0.001$) and fat mass (-97g , 95%CI $-205;10$, $p=0.076$). There were no difference in changes in absolute subscapular and triceps skinfold-thickness and z-scores ($p>0.5$).

Conclusions: SHORT children with low MUAC do not gain excessive fat during supplementation. These data support a recommendation for policy change to include all children ≥ 6 months with low MUAC in supplementary feeding programs, regardless of length. The use of length as a criterion for measuring MUAC to determine treatment eligibility should be discontinued in policy and practice.

INTRODUCTION

Childhood malnutrition contributes to almost half the mortality in children under five years¹. Moderate acute malnutrition (MAM) is currently defined as weight-for-height z-score (WHZ) between -3 and -2 (i.e. moderate wasting), and/or mid-upper arm circumference (MUAC) between 115 and 125 mm². While the prevalence of MAM is unknown, moderate wasting alone affects 33 million children at any time³ and is associated with a three-fold increased risk of death⁴.

Protocols for management of acute malnutrition in many African countries (including Cameroon, Central African Republic, Chad, Guinea, Ivory Coast, Mali, Mauritania, Senegal, and Togo) instruct health personnel to measure MUAC only of children with length ≥ 67 cm when assessing eligibility for MAM or severe acute malnutrition (SAM) treatment programs⁵⁻¹³. In Ethiopia, admission by MUAC alone for SAM treatment is restricted to children with lengths >65 cm¹⁴. These shorter children are enrolled in treatment only if they meet WHZ criterion, which is less closely linked to the risk of death and misses high-risk children who would have been identified by MUAC. Hence, current practice excludes short children who would benefit from nutritional support.

The practice of restricting treatment admission by MUAC alone to children of length ≥ 67 cm is not supported by data and has never been formally endorsed by the World Health Organization (WHO). Two reasons can be identified to explain this exclusion. First, the inclusion of a length restriction for MUAC is likely a holdover from older versions of emergency nutrition guidelines that used length as a proxy for age (<6 months) because it can sometimes be difficult to ascertain age in a fast moving emergency. The WHO, for example, used length of 60 cm as a proxy for age of 6 months in publications from 1995 and 2000^{15,16} while length of 65 cm was used in older versions of guidelines from non-governmental humanitarian organizations like Médecins Sans Frontières (MSF) in 1995 and Valid International in 2006^{17,18}. It is important to identify children <6 months

1
2
3
4 of age as they should be exclusively breastfed, and treated as in-patients if in need of treatment.
5
6 However, length is a poor proxy for age, and most often a caretaker can recall the month of birth for
7
8 an infant.
9

10
11 Second, WHO now recommends against routine food supplementation of children with MAM due
12
13 to a concern it may promote obesity and increase the risk of non-communicable diseases later in life
14
15 ¹⁹. Some experts worry that shorter children above 6 months of age who only meet the low-MUAC
16
17 criteria for MAM are at most at risk of excessive fat accumulation during treatment ². Although
18
19 only based on expert opinion, this seems to be the reason why many national malnutrition protocols
20
21 exclude shorter children from having their MUAC measured, thereby making children ineligible for
22
23 treatment, unless they meet WHZ criteria. This concern led WHO to call for research to identify
24
25 appropriate MUAC admission and discharge criteria for children <67 cm and ≥ 6 months to
26
27 supplementary feeding programs ².
28
29

30
31 We have previously shown, that when supplemented, ponderal growth rates are similar in short and
32
33 long children ≥ 6 months with a low MUAC ²⁰. Niger dropped this length restriction in 2016, and its
34
35 revised protocols now call for measuring MUAC in all children ≥ 6 months, regardless of length ²¹.
36
37

38
39 To what extent there is a difference in fat accumulation during treatment has not been assessed. The
40
41 objective of this paper is to assess if short children (<67 cm) gain more fat than long children (≥ 67
42
43 cm) when treated for MAM diagnosed by low MUAC (115-125 mm).
44
45
46
47

48 49 **SUBJECTS AND METHODS**

50
51 As previously reported ²⁰, a prospective cohort study was nested in a randomized nutrition trial
52
53 (Treatfood) ²² investigating the effectiveness of 500 kcal/day supplement either as corn-soy blend
54
55
56
57
58
59
60

1
2
3
4 (CSB) porridge or ready-to-use lipid-based nutrient supplements (LNS). The LNS supplements
5 provided almost three times more energy as fat than the CSB supplements (~57% vs ~21%).
6

7
8 Assignment to one of the 12 supplements (6 CSB and 6 LNS) followed randomisation stratified by
9
10 site.
11

12 13 **Participants**

14
15
16 Data were collected in the Province du Passoré in the Northern region of Burkina Faso at 5 research
17 sites located at different governmental health centers and staffed by the non-governmental
18 organization Alliance for International Medical Action (ALIMA, Dakar, Senegal).
19

20
21 Children were screened in villages either by community health workers using MUAC tapes or by
22 designated screening teams with the use of both MUAC and WHZ. Moreover, children could be
23 referred from a health centre or could present at site on caretaker's initiative. At the sites, the final
24 assessment of eligibility for inclusion was performed.
25
26
27
28
29

30
31
32 For the main Treatfood trial, children aged 6–23 months with MAM (defined as MUAC between
33 115-124 and/or a WHZ between –3 and –2), resident in the catchment area and whose caretaker
34 consented to participate, were included. Children were not included if treated for severe acute
35 malnutrition (SAM) or hospitalized within the past two months, if already in a nutritional program
36 or if they presented medical complications requiring hospitalization. Likewise, children with a
37 severe disability limiting the possibility of investigations and children with suspected allergy to
38 milk, peanuts, corn or soy were not included.
39
40
41
42
43
44
45
46
47
48

49
50 Of the 1609 children included in the trial 468 had MUAC between 115-125mm and WHZ \geq -2, and
51 were included in the current study. In many settings these children are excluded from treatment if
52 they are also short (i.e. <65 or 67 cm). Accordingly, children in this study were categorized, based
53
54
55
56
57
58
59
60

1
2
3
4 on baseline length, as SHORT (<67 cm) or LONG (\geq 67 cm) to assess if SHORT children gain more
5 fat than LONG children when treated for MAM diagnosed by low MUAC.
6
7
8
9
10

11 **Procedures and study visits**

12
13
14 We previously described clinic visits, standard anthropometric measurements and age determination
15 ²⁰. In the present paper we additionally report on indices of body composition and skinfolds
16 assessed at baseline and at end-intervention at 12 weeks. Total body water (TBW) was assessed
17 using the deuterium dilution technique (D2O) to enable calculation of fat-free mass (FFM) and fat
18 mass (FM). The method involved giving an oral dose of 5 g deuterium oxide (D2O) (99.8%,
19 Cambridge Isotope Laboratories Inc., Andover, USA). The isotope was diluted in 5 g of bottled
20 water (LAFI, Burkina Faso), with the dosing bottle weighed with 0.01 g precision (Adam
21 equipment: model CQT 202, United Kingdom) before and after administration of the dose. Pre-dose
22 saliva samples were obtained to assess background isotope levels in body fluids, and post-dose
23 saliva samples were collected after a three-hour equilibration period as established during the pilot
24 study ²³. For each assessment, D2O enrichment was measured in duplicate in the pre- and post-dose
25 saliva samples and in a diluted sample of the dose, using Fourier Transform Infrared Spectrometry
26 (FTIR; Agilent Technologies, CA, USA) ²⁴ at St. John's Research Centre, Bangalore, India.
27 Saliva samples required at least 60 μ l saliva for analysis. Deuterium dilution space was calculated
28 as described previously ²⁵, and converted to total body water (TBW) using a factor of 1.044 to
29 adjust for proton exchange ²⁶. FFM was calculated as TBW/hydration, using age- and sex-specific
30 hydration coefficients ²⁷. FM was calculated as weight minus FFM. Data were cleaned for
31 typographical errors and implausible TBW values, based on the association of TBW with length
32 and a cut-off for FM of <0.1 and >2.4 kg.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 Skinfold thickness was measured in duplicate by a Harpenden caliper. The mean of the duplicate
5
6 measurements were taken for analysis. Weight was measured in duplicate to the nearest 100 g using
7
8 electronic scales (Seca model 881 1021659) with double weighing function. Length was measured
9
10 in duplicate with a wooden length board to the nearest 1 mm. WHZ was determined at sites using
11
12 WHO field tables, and this value was used for recruitment. MUAC was measured in duplicate to the
13
14 nearest 1 mm, at the midpoint between the olecranon and the acromion process using a standard
15
16 measuring tape. Anthropometric measurements were undertaken by trained staff, after
17
18 standardization sessions.
19

20
21
22 In later analyses, WHZ, length-for-age z-score (LAZ) and weight-for-age z-score were calculated
23
24 using the package “zscore06” in STATA 12 (College Station, Texas, USA). Skinfold-for-age z-
25
26 scores were calculated using WHO's Anthro Plus software (version 3.2.2, 2011, World Health
27
28 Organization, Geneva, Switzerland). All z-scores were calculated using the 2006 WHO child
29
30 growth standards.
31

32 33 **Outcomes**

34
35
36 Changes were evaluated in FM, FFM, weight, length, and skinfold thickness (both raw values and
37
38 skinfold thickness-for-age z-scores). By dividing FFM and FM by length squared the indices are
39
40 expressed as (kg/m^2) independent of length, i.e. FFM index (FFMI) and FM index (FMI). In
41
42 addition, fat was calculated as a percentage of total body weight.
43
44
45

46 47 **Statistical analysis**

48
49
50 Data were double entered in Epidata 3.1 (Epidata Association, Odense, Denmark) and double entry
51
52 checks were performed on a daily basis. All statistical analyses were carried out using the statistical
53
54 software package Stata version 12 (StataCorp, College station, Texas, USA). Baseline and endline
55
56
57
58
59
60

1
2
3
4 characteristics of children were summarized as mean (SD) or percentages, and categories were
5
6 compared at baseline using two-sample t-tests for continuous variables and chi-square tests for
7
8 categorical variables.
9

10
11 To evaluate differences in outcomes between groups during the 12-week supplementation
12
13 intervention, a linear mixed analysis of covariance model was considered as the main analysis,
14
15 resulting in an estimated mean difference at the end of the study adjusted for differences at baseline.
16
17 Specifically, the analyses included adjustment for food intervention as well as a number of baseline
18
19 characteristics: baseline measure of the outcome, age, sex, and month of admission (as fixed
20
21 effects). Random intercepts were included in the model to adjust for variation between sites. In the
22
23 adjusted analysis, weight was derived by adding FFM and FM. Additionally, for direct outcomes of
24
25 fat (FM and FMI) we evaluated the two-way interaction between length group and product group
26
27 (CSB compared with LNS) in order to assess effect modification by type of product. Moreover,
28
29 possible effect modification was also evaluated for stunting at admission in a) two categories: above
30
31 and below LAZ of <-2 , or b) or three categories: severe stunting (LAZ <-3), moderate stunting
32
33 (LAZ ≥-3 to <-2) or absence of stunting (LAZ ≥-2). The same adjustments were applied in these
34
35 additional analyses as in the main analysis. Model checking was based on residuals and normal
36
37 probability plots. A significance level of 0.05 was applied.
38
39
40
41
42
43
44
45

46 **RESULTS**

47
48
49 A total of 1609 children, of which 55% were girls, were randomized to 12 weeks of supplementary
50
51 feeding in the Treatfood trial²². Of these, 50% (804) were included by both WHZ and MUAC, 21%
52
53 (337) by WHZ alone and 29% (468) by MUAC alone. Among the 468 children recruited by MUAC
54
55
56
57
58
59
60

1
2
3
4 only, on whom this this paper is based, 230 (49%) were <67 cm (SHORT) and 238 (51%) \geq 67 cm
5 (LONG)²⁰. During the intervention, 3 SHORT and 11 LONG children dropped out prior to
6 completion of supplementation at week 12. Body composition based on D2O dilution both at
7 baseline and after 12 weeks were successfully determined in 195 (85%) of SHORT children and
8 195 (82 %) of LONG children (**Figure 1**). At baseline, there were substantial differences in age and
9 sex distributions, as well as in standard anthropometry between SHORT and LONG, whereas there
10 were no differences in allocation to the different experimental diets and season of inclusion (**Table**
11 **1**). Mean values of body composition variables determined at baseline and after supplementation are
12 presented in **Table 2**. At baseline, the weight of SHORT children was 1.474 (95%CI: 1.355; 1.593)
13 kg less than that of LONG children, which reflected a 0.120 (0.047; 0.193) kg lower FM and a
14 1.354 (1.219; 1.489) kg lower FFM. Adjusted for length squared this corresponded to a 0.325
15 (0.163; 0.487) kg/m² higher FMI in SHORT compared to LONG children, but no difference (-0.056
16 kg/m², -0.230; 0.118) in FFMI. Body fat was 1.9 (0.8; 3.0) percentage points higher in SHORT
17 compared to LONG children.

18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35 Adjusted differences from baseline to endline are presented in **Table 3**. Following the 12 weeks of
36 supplementation, there was no difference of change in FMI (-0.038 kg/m², 95%CI -0.257; 0.181,
37 p=0.74) or FFMI (0.061 kg/m², 95%CI -0.150; 0.271; p=0.57) in SHORT vs LONG children. The
38 SHORT children actually gained both less FFM (-230 g, 95%CI: -355; -106, p<0.001) as well as
39 FM (-97 g, 95%CI -205; 10) although the latter was not significant (p=0.076). The differences in
40 FM and FFM summed up to a 328 g (95%CI: 199; 456, p <0.001) lower weight gain in SHORT
41 compared to LONG children. There were no differences in subscapular and triceps skinfold
42 thickness and z-scores (all p>0.5).

43
44
45
46
47
48 For FM and FMI there was no effect modification of product type, (CSB or LNS) (p>0.30).

49
50
51
52
53
54
55 Likewise, there was no effect modification by stunting at admission (present or non-present)

1
2
3
4 (p>0.50) or stunting at admission stratified by severity (severe stunting, moderate stunting or non-
5
6 present) (p>0.45).
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Review Copy

DISCUSSION

In response to WHO's call for research to identify appropriate MUAC admission and discharge criteria for children <67 cm and ≥ 6 months to supplementary foods², we show here that short children with MUAC 115-125 mm but WHZ >-2 do not gain excessive fat during supplementation.

Our results therefore challenge current policy and practice in several African countries of measuring MUAC only in children more than 6 months of age above a certain length threshold, thus excluding shorter children with only low MUAC from eligibility for MAM or SAM treatment programs⁵⁻¹⁴.

The exclusion of short children is based solely on expert opinion, and reflects a concern that these children are just stunted rather than wasted, have slow catch-up growth, and may accumulate excessive fat mass following supplementation, thereby putting them at risk of non-communicable diseases later in life. We previously reported that ponderal growth rates were similar in short and long children who received supplementation for MAM determined only by low MUAC²⁰, and a similar finding was reported in children with SAM²⁸.

The current study is the first to report direct evidence on fat accumulation following treatment for such children, and our data show there is no excessive fat gain in the SHORT children compared to LONG children. In fact, weight gain in both groups overwhelmingly came in the form of FFM.

These findings were independent of children having received LNS or CSB supplement. Because LNS contains a high level of fat, some experts were concerned that this food could lead to higher fat deposition in the malnourished child. Earlier we showed that LNS did not lead to high fat accumulation in treatment of children for MAM²². We now show that this concern is unwarranted even in shorter children enrolled in treatment solely by MUAC and, furthermore, is not modified by stunting at admission.

1
2
3
4 By using the state-of-the-art deuterium dilution technique seconded by skinfold thickness
5
6 measurements, we present comprehensive data on body fat following supplementation in MAM.
7
8 Since the gold standard for body composition is cadaver dissection, all *in vivo* techniques are
9
10 necessarily imperfect. Here, we used two different techniques in which their error is uncorrelated,
11
12 but their results are similar, making our findings more robust.
13
14

15 At baseline, the SHORT compared to the LONG children were younger, contained more girls, had
16
17 lower MUAC, and had higher prevalence of stunting and underweight, but were less wasted²⁰.
18
19 More than 90% of the 1.5 kg higher weight in LONG children at baseline was FFM, while SHORT
20
21 children had higher baseline FMI and fat%, and this was supported by higher baseline skinfold
22
23 thicknesses.
24
25

26
27 One possible explanation for this baseline difference is that, in the process of growing poorly,
28
29 SHORT children invested relatively more in fat reserves than LONG children, at a cost to their
30
31 functional fat-free tissue. However, a simpler alternative explanation is that even in children with
32
33 optimal growth conditions, such as those used to establish WHO 2006 growth standards, both
34
35 triceps and subscapular skinfold thicknesses tend to decrease in thickness between 6 months and 2
36
37 years, and more so in boys than girls. Hence, the fact that the SHORT children were also ~6 months
38
39 younger at baseline might explain the higher FMI at baseline. Supporting this, baseline skinfold
40
41 thicknesses were thicker in SHORT children, but when expressed as z-scores relative to WHO
42
43 reference data, SHORT children actually had lower triceps values than LONG children, while their
44
45 subscapular skinfold thickness was only 0.2 z-score greater than in LONG children.
46
47
48

49 At the end of the intervention, the average skinfolds thickness in this study were below -1 z-score in
50
51 both groups suggesting no excessive fat gain in either group.
52
53
54
55
56
57
58
59
60

1
2
3
4 At the end of the supplementation relative measures of FMI, FFMI and percentage fat were similar
5
6 between the two groups and likewise there was no difference between groups in skinfold thickness
7
8 changes, whether expressed in absolute terms or as z-scores. The SHORT children gained less
9
10 weight and FFM and also showed a trend towards less FM gain.
11

12
13 A recent study found that children with MAM diagnosed solely on low MUAC experienced high
14
15 rates of deterioration to SAM if left untreated²⁹. We are not aware of data on mortality data
16
17 specifically for short children with MAM based only on low MUAC. However, the low MUAC
18
19 SHORT children in our cohort exhibited high prevalence of additional anthropometric deficits. Half
20
21 of the SHORT children were stunted and SHORT children were more underweight compared to
22
23 LONG children. Stunted and underweight children have a higher mortality risk and may therefore
24
25 benefit most from nutritional interventions³⁰. MUAC is a simple way to target them for treatment.
26
27
28

29 The WHO recently updated its Integrated Management of Childhood Illness (IMCI), and now
30
31 recommends against routinely supplementing children with MAM due to a concern for obesity. We
32
33 previously showed that there was no excess fat accumulation in children with MAM supplemented
34
35 with LNS or CSB²². Our findings here show that even the shorter children with MAM determined
36
37 solely by MUAC, who were considered by some experts to be at the greatest risk for obesity from
38
39 food supplementation, do not become obese. Such children should no longer be excluded from
40
41 treatment eligibility.
42
43
44

45 We are not aware of any data on fat accumulation following treatment in short children with SAM
46
47 by MUAC alone. Indeed, currently only data on skinfolds could be generated in large field studies
48
49 in children with SAM, as current techniques available for body composition assessment are not
50
51 suitable for use in SAM. However, we see no rationale in maintaining a length restriction for
52
53 MUAC assessment in SAM.
54
55
56
57
58
59
60

1
2
3
4 In conclusion, SHORT children with low MUAC do not gain excessive fat during supplementation.
5
6 Admission by the same MUAC criteria (i.e. <115 mm for SAM and 115-125 mm for MAM) should
7
8 apply to any child 6-59 months of age, regardless of their length, and irrespective of their WHZ.
9
10 The use of length as criterion for measuring MUAC should be discontinued in policy and practice
11
12 wherever such restrictions exist. It will also be important for the WHO to integrate this evidence
13
14 showing that children with MAM are not put at risk for obesity following food supplementation into
15
16 its recently updated IMCI recommendations.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Review Copy

DECLARATIONS**Ethics approval and consent to participate**

The trial was approved by the Ethics Committee for Health Research in Burkina Faso (2012-8-059) and consultative approval was obtained from the Danish National Committee on Biomedical Research Ethics (1208204). Consent was obtained from caregivers, prior to inclusion, verbally and in writing (signature or fingerprints). All children were treated free of charge at the study sites, irrespective of study participation. The trial is registered at www.controlled-trials.com (ISRCTN42569496).

Consent for publication

Not applicable

Availability of data and material

The dataset used during the current study is available from the corresponding author on reasonable request.

Acknowledgements

Not applicable


References

1. WHO | Children: reducing mortality. WHO. <http://www.who.int/mediacentre/factsheets/fs178/en/>. Accessed September 12, 2016.
2. WHO, UNICEF, WFP and UNHCR Consultation on the Programmatic Aspects of the Management of Moderate Acute Malnutrition in Children under five years of age. Geneva. WHO. 2010.
3. Black RE, Victora CG, Walker SP, et al. Maternal and child undernutrition and overweight in low-income and middle-income countries. *The Lancet*. 2013;382(9890):427–451.
4. Olofin I, McDonald CM, Ezzati M, et al. Associations of suboptimal growth with all-cause and cause-specific mortality in children under five years: a pooled analysis of ten prospective studies. *PloS One*. 2013;8(5):e64636. doi:10.1371/journal.pone.0064636
5. *Cameroon Ministry of Public Health. National Protocol for the Integrated Management of Acute Malnutrition. 2013: Pp 20, 29, 119. [in French].*
6. *Central African Republic Ministry of Public Health and Population. Protocol for the Integrated Management of Acute Malnutrition. December 2014: Pp 40, 57, 168. [in French].*
7. *Republic of Chad Ministry of Public Health, Social Action and National Solidarity. Protocol for the Integrated Management of Acute Malnutrition. May 2014: P 166. [in French].*
8. *Guinea Ministry of Health and Public Hygiene. National Protocol for the Integrated Management of Acute Malnutrition. 2012: Pp 36, 52, 149. [in French].*
9. *Ivory Coast Ministry of Public Health and Hygiene. National Protocol for the Integrated Management of Acute Malnutrition. 2015: Pp 56, 83, 188. [in French].*
10. *Islamic Republic of Mauritania Ministry of Health. Protocol for the Integrated Management of Acute Malnutrition. December 2011: Pp 57, 146 [in French].*
11. *Republic of Senegal Ministry of Health and Social Action. Protocol for the Integrated Management of Acute Malnutrition. November 2013: Pp 26, 40, 57, 156 [in French].*
12. *Togo Ministry of Health. National Protocol for the Integrated Management of Acute Malnutrition. July 2013: Pp 33, 123 [in French].*
13. *Mali Ministry of Health. Protocol for the Integrated Management of Acute Malnutrition in Mali. 2011: Pp 31, 138 [in French].*
14. *Ethiopia Federal Ministry of Health. Protocol for the Management of Severe Acute Malnutrition. March 2007: P 10.*
15. WHO | The management of nutrition in major emergencies. Geneva. WHO. 2000.

16. WHO | WHO Regional Office for the Eastern Mediterranean. *Field Guide on Rapid Nutritional Assessment in Emergencies*. Alexandria (Egypt): WHO; 1995.
17. MSF | Nutrition Guidelines. Médecins Sans Frontières. 1995.
18. VALID | Community-based Therapeutic Care (CTC) A field manual, Valid International. 2006.
19. WHO | Assessing and managing children at primary health-care facilities to prevent overweight and obesity in the context of the double burden of malnutrition. WHO. <http://www.who.int/nutrition/publications/guidelines/children-primaryhealthcare-obesity-dbm/en/>. Accessed November 1, 2017.
20. Fabiansen C, Phelan KP, Cichon B, et al. Short children with a low midupper arm circumference respond to food supplementation: an observational study from Burkina Faso. *Am J Clin Nutr*. 2016;103(2):415-421. doi:10.3945/ajcn.115.124644
21. Niger Ministry of Public Health. Protocol for the integrated management of acute malnutrition. March 2016: p 127 [in French].
22. Fabiansen C, Yaméogo CW, Iuel-Brockdorf A-S, et al. Effectiveness of food supplements in increasing fat-free tissue accretion in children with moderate acute malnutrition: A randomised 2 × 2 × 3 factorial trial in Burkina Faso. *PLoS Med*. 2017;14(9):e1002387. doi:10.1371/journal.pmed.1002387
23. Fabiansen C, Yaméogo CW, Devi S, Friis H, Kurpad A, Wells JC. Deuterium dilution technique for body composition assessment: resolving methodological issues in children with moderate acute malnutrition. *Isotopes Environ Health Stud*. 2017;0(0):1-12. doi:10.1080/10256016.2017.1295043
24. International Atomic Energy Agency. *Introduction to Body Composition Assessment Using the Deuterium Dilution Technique with Analysis of Saliva Samples by Fourier Transform Infrared Spectrometry*. Vienna: International Atomic Energy Agency; 2011.
25. Davies PS, Wells JC. Calculation of total body water in infancy. *Eur J Clin Nutr*. 1994;48(7):490-495.
26. Racette SB, Schoeller DA, Luke AH, Shay K, Hnilicka J, Kushner RF. Relative dilution spaces of 2H-and 18O-labeled water in humans. *Am J Physiol-Endocrinol Metab*. 1994;267(4):E585-E590.
27. Lohman T.G. "Estimating body composition in children and the elderly", *Advances in Body Composition Assessment (Current Issues in Exercise Science, Monograph 3)* (LOHMAN, T.G., Ed.), Human Kinetics, Champaign, IL (1992) 65-77.
28. Dale NM, Myatt M, Prudhon C, Briend A. Using Mid-Upper Arm Circumference to End Treatment of Severe Acute Malnutrition Leads to Higher Weight Gains in the Most Malnourished Children. *PLoS ONE*. 2013;8(2):e55404. doi:10.1371/journal.pone.0055404

- 1
2
3
4 29. James P, Sadler K, Wondafrash M, et al. Children with Moderate Acute Malnutrition with No
5 Access to Supplementary Feeding Programmes Experience High Rates of Deterioration and
6 No Improvement: Results from a Prospective Cohort Study in Rural Ethiopia. *PLOS ONE*.
7 2016;11(4):e0153530. doi:10.1371/journal.pone.0153530
8
- 9 30. McDonald CM, Olofin I, Flaxman S, et al. The effect of multiple anthropometric deficits on
10 child mortality: meta-analysis of individual data in 10 prospective studies from developing
11 countries. *Am J Clin Nutr*. 2013;97(4):896-901. doi:10.3945/ajcn.112.047639
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Review Copy

Figure 1 participant flow chart^a

^a Treatfood participants included in the cohort study²². ^b D2O = Body composition assessment by deuterium dilution technique

Table 1. Baseline characteristics of children with moderate acute malnutrition included by MUAC only with complete D2O data (n=390), by length category^a

	^b SHORT (n=195)	^c LONG (n=195)	P-value
Girls, % (n)	84% (163)	72% (140)	0.005
Age, months	7.7 (1.6)	14.0 (4.3)	<0.001
Weight, kg	6.012 (0.4)	7.487 (0.7)	<0.001
Length, cm	63.9 (2.1)	72.0 (3.9)	<0.001
Mid upper arm circumference, mm	121.0 (2.6)	122.2 (1.9)	<0.001
Weight for height z-score	-1.5 (0.4)	-1.7 (0.3)	<0.001
Weight for age z	-2.4 (0.6)	-2.1 (0.6)	<0.001
Length for age z	-2.0 (1.0)	-1.7 (1.0)	0.002
Length for age z, <-2 % (n)	49% (95)	35% (68)	0.006
Season at time of inclusion, % (n)			0.17
Dry season	61% (118)	67% (131)	
Rainy season	40% (77)	33% (64)	
Food supplement, % (n)			0.84
CSB	48% (94)	47% (92)	
LNS	52% (101)	53% (103)	
Site, % (n)			0.009
0	12% (24)	27% (52)	
1	15% (30)	11% (21)	
2	21% (41)	18% (35)	
3	31% (60)	26% (50)	
4	21% (40)	19% (37)	
Breastfeeding, % (n)	100% (194)	91% (178)	<0.001

^aData are mean (\pm SD) unless otherwise indicated. ^bSHORT <67 cm, ^cLONG \geq 67 cm.

Table 2. Body composition among short and long children with moderate acute malnutrition at baseline and endline, without adjustments^a

	Baseline			Endline		
	^b SHORT (n=195)	^c LONG (n=195)	^d P Δ	SHORT (n=195)	LONG (n=195)	^e P Δ
Fat mass, kg	1.098 (0.4)	1.218 (0.4)	0.001	1.123 (0.4)	1.246 (0.4)	0.003
Fat-free mass, kg	4.914 (0.5)	6.268 (0.8)	<0.001	5.673 (0.5)	7.096 (0.9)	<0.001
%Fat, (Fat-mass/weight*100)	18.3 (5.8)	16.4 (5.3)	<0.001	16.3 (5.7)	15.0 (4.9)	0.015
Fat mass index, kg/m ²	2.689 (0.9)	2.364 (0.8)	<0.001	2.499 (0.9)	2.259 (0.8)	0.006
Fat-free mass index, kg/m ²	12.000 (0.9)	12.056 (0.9)	0.53	12.640 (0.8)	12.723 (0.9)	0.36
Triceps skinfold, mm	6.6 (1.1)	6.3 (1.0)	0.001	6.8 (1.1)	6.8 (1.2)	0.85
Triceps skinfold-for-age z	-1.5 (0.8)	-1.3 (0.8)	0.04	-1.1 (0.8)	- 0.8 (0.9)	0.007
Subscapular skinfold, mm	5.6 (0.9)	5.1 (0.8)	<0.001	5.8 (1.0)	5.4 (0.8)	<0.001
Subscapular skinfold-for-age z	-1.3 (0.9)	-1.5 (0.9)	0.03	-0.9 (1.0)	-0.9 (0.9)	0.96

^aData are mean (±SD) unless otherwise indicated. ^bSHORT <67 cm, ^cLONG ≥67 cm. ^dP value for difference at baseline between SHORT and LONG children. ^eP value for difference at endline between SHORT and LONG children.

Table 3. Changes in outcomes during supplementation^a

	^b SHORT (n=195)	^c LONG (n=195)	^d Difference in change	^e P Δ
Fat mass, kg	-0.022 (-0.088; 0.043)	0.075 (0.010; 0.140)	-0.097 (-0.205; 0.010)	0.076
Fat-free mass, kg	0.678 (0.604; 0.752)	0.908 (0.834; 0.982)	-0.230 (-0.355; -0.106)	<0.001
%Fat (Fat-mass/weight*100)	-1.8 (-2.6; -1.0)	-1.5 (-2.3; -0.7)	-0.3 (-1.6; 1.1)	0.70
Fat mass index, kg /m ²	-0.166 (-0.309; -0.234)	-0.128 (-0.271; 0.015)	-0.038 (-0.257; 0.181)	0.74
Fat-free mass index, kg/m ²	0.684 (0.556; 0.813)	0.624 (0.495; 0.752)	0.061 (-0.150; 0.271)	0.57
Triceps skinfold, mm	0.39 (0.04; 0.73)	0.35 (-0.002; 0.70)	0.04 (-0.24; 0.32)	0.77
Triceps skinfold-for-age z	0.48 (0.25; 0.71)	0.50 (0.26; 0.73)	-0.02 (-0.22; 1.85)	0.88
Subscapular skinfold, mm	0.32 (0.17; 0.47)	0.24 (0.09; 0.40)	0.08 (-0.15; 0.30)	0.52
Subscapular skinfold-for-age z	0.52 (0.36; 0.67)	0.49 (0.34; 0.65)	0.02 (-0.21; 0.26)	0.84

^aData are mean and 95% CI calculated using linear mixed models adjusted for food intervention, baseline measure of the outcome, age, sex, month of admission and site (random effects). ^bSHORT <67 cm, ^cLONG ≥67 cm. ^dDifference in change of SHORT vs LONG. ^eP value for difference in change from baseline to endline for SHORT and LONG children.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Review Copy

List word counts below (do not paste the text here). Please see the Decision Letter Attachment for allowances as they pertain to your manuscript type.

of words in Abstract: **247** (250 words allowed)

of words in Manuscript Body: **3155** (3000 allowed for Regular Articles/Quality Reports; 4000 Reviews/Special Articles; 800 Commentaries; 1200 Perspectives)

of characters in Main Title: **79** characters (97 characters allowed, including spaces)

of characters in Short Title: **55** (55 characters allowed, including spaces)

of words in "Table of Contents Summary": **25** (25 words allowed; this section appears in all articles with abstracts)

of words in "What's Known on this Subject": **40** (40 words allowed; this section appears in Regular Articles only)

of words in "What this Study Adds": **39** (40 words allowed; this section appears in Regular Articles only)

2016-XXXX.RX – Short malnourished children do not gain excessive fat with food supplementation -- by Fabiansen et al.

EDITOR/REVIEWER COMMENTS <i>Paste each of the editor and reviewer queries here.</i>	AUTHOR'S RESPONSE <i>Paste your answer to the editor and reviewer queries here. If you alter your manuscript to address this query, you MUST paste the relevant altered text here – verbatim as it appears in the manuscript.</i>	REFERENCE PAGE <i>State where* the change now appears in your newly revised manuscript.</i>	CHANGE APPROVED ? FOR EDITORIAL USE ONLY
Editors Comments	<p>We thank you for provisionally accepting our manuscript.</p> <p>Our current title is: Short malnourished children with a low MUAC do not gain excessive fat with food supplementation</p> <p>You suggest we change it into: Short malnourished children and fat accumulation with food supplementation</p> <p>We fully acknowledge that our original title was rather long. However, we would like the title to give an idea of the main message of the paper, and your suggestion may indicate fat accumulation. We therefore kindly ask to change the title into:</p> <p>“Short malnourished children do not gain excessive fat with food supplementation”</p> <p>Likewise, we hope it is acceptable that your suggested short title Short malnourished children and fat accumulation</p>		

<p>1</p> <p>2</p> <p>3</p> <p>4</p> <p>5</p> <p>6</p>	<p>Is changed into:</p> <p>“Short malnourished children do not gain excessive fat”</p>		
<p>7</p> <p>8</p> <p>9</p> <p>10</p> <p>11</p> <p>12</p> <p>13</p> <p>14</p> <p>15</p> <p>16</p> <p>17</p> <p>18</p> <p>19</p> <p>20</p> <p>21</p> <p>22</p> <p>23</p> <p>24</p> <p>25</p> <p>26</p> <p>27</p> <p>28</p> <p>29</p> <p>30</p> <p>31</p> <p>32</p> <p>33</p> <p>34</p> <p>35</p> <p>36</p> <p>37</p>	<p>Reviewer 1’s comment</p> <p>We thank reviewer 1 for the very positive review.</p> <ol style="list-style-type: none"> 1. Yes thanks, this is now avoided in the new title. 2. Thanks for pointing this out. Sample size calculation was made for the main trial. We now made it clearer which children were included in current study. The text now reads: <p>“Of the 1609 children included in the trial 468 had MUAC between 115-125mm and WHZ \geq-2, and were included in the current study. In many settings these children are excluded from treatment if they are also short (i.e. <65 or 67 cm). Accordingly, children in this study were categorized, based on baseline length, as SHORT (<67 cm) or LONG (\geq67 cm) to assess if SHORT children gain more fat than LONG children when treated for MAM diagnosed by low MUAC .”</p> <ol style="list-style-type: none"> 3. Thanks for pointing this out. The problem is that “²²” is a reference in “Pediatrics” format. To improve readability we now changed table superscripts to letters. 4. Thanks. MAM is now written as moderate acute malnutrition. 	<ol style="list-style-type: none"> 1. Title 2. In <i>statistical analysis</i> 3. In <i>Participants</i> 4. Tab 1 +2 	
<p>39</p> <p>40</p> <p>41</p>	<p>Reviewer 2’ comment</p> <p>We thank reviewer 2 for the kind comments on our study and for us testing BOGSAT opinion</p>		
<p>43</p>	<p>Reviewer 3’s comment</p> <p>We thank reviewer 3 for the very positive review</p>	<p>1 In <i>statistical</i></p>	

<p>1 1. Methodology: Were the mixed 2 models “random intercept” or 3 “random intercept” and “random 4 slope” models? 5 6 2. Results: How were the results 7 (mean difference, 95% CI and P- 8 value) for weight gain obtained? For 9 me is not clear: why as the weight 10 difference between end of 11 intervention and baseline not used? 12 13 3. Discussion: Are there potential 14 bias that could limit the 15 generalization of the results? 16 17 4. Are other studies needed to 18 validate the results from this study? 19 20 5. Is it possible that these 21 supplemented children develop 22 obesity later in childhood or 23 adolescence? 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42</p>	<p>1. We’ve now clarified that the linear mixed models only included random intercepts corresponding to the sites. The text now reads:</p> <p>“To evaluate differences in outcomes between groups during the 12-week supplementation intervention, a linear mixed analysis of covariance model was considered as the main analysis, resulting in an estimated mean difference at the end of the study adjusted for differences at baseline. Specifically, the analyses included adjustment for food intervention as well as a number of baseline characteristics: baseline measure of the outcome, age, sex, and month of admission (as fixed effects). Random intercepts were included in the model to adjust for variation between sites. In the adjusted analysis, weight was derived by adding FFM and FM. Additionally, for direct outcomes of fat (FM and FMI) we evaluated the two-way interaction between length group and product group (CSB compared with LNS) in order to assess effect modification by type of product. Moreover, possible effect modification was also evaluated for stunting at admission in a) two categories: above and below LAZ of <-2, or b) or three categories: severe stunting (LAZ <-3), moderate stunting (LAZ ≥-3 to <-2) or absence of stunting (LAZ ≥-2). The same adjustments were applied in these additional analyses as in the main analysis. Model checking was based on residuals and normal probability plots. A significance level of 0.05 was applied.”</p> <p>2. Thanks for the question. Weight gain was estimated using the weight measurements at the end of the study but adjusted for baseline weight (analysis of covariance). We’ve now spelled out that we used an analysis of covariance type of linear mixed model. In this way, confidence intervals and p-values were obtained directly from the model fits (This is included in text in the paper as seen in the answer to your first question)</p> <p>3. We don’t consider bias to be a problem in this study. First, selection bias is unlikely, since we included a large proportion of</p>	<p><i>analysis</i></p>	
---	---	------------------------	--

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

those with MAM, based on MUAC, in the study, and the follow-up rate was very high, and not different between the two groups being compared. Hence, it is unlikely that selection could have led to a bias because of, say, lower probability of inclusion/higher attrition of children more prone to accumulate fat in the short compared to the long children. Second, information bias is unlikely since the outcome was assessed in the lab by staff who were unaware of the exposure of interest. Theoretically, a source of information bias could be that short children were more or less likely to spit when given the deuterium. For example, if short (younger) children were more likely to spit, then the concentration of deuterium would be underestimated, and the total body water overestimated, and therefore, the fat mass would be underestimated. However, since this would then happen base at baseline and endline, it is not likely to affect the change in fat mass, which is our outcome of interest. In general, generalizability is likely to be lower in nutrition compared to other intervention trials, since the effect of an intervention depends on the background nutritional intake and status. However, in this substudy, we are not testing the effect of a nutrition intervention, but if short stature affects changes in body composition. Certainly, since we used strict inclusion criteria we think that our findings can be generalized to other West African populations of young children with MAM, diagnosed using MUAC, at least. Yet, we cannot exclude the possibility that different fetal and early life trajectories in growth and body composition, as well as differences in background diets, may modify the difference in change in body composition during supplementation.

This paper specifically asks for a policy change in the remaining African countries where length restriction is applied to MUAC measurements, these countries are located in West and Central Africa where morbidity/mortality and malnutrition patterns (very high incidence of acute malnutrition accompanied by high levels of stunting) are similar to that seen in Burkina. We would welcome further studies. It is probably worthwhile to repeat such body

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

composition studies in other malnutrition contexts such as high prevalence of kwashiorkor in DRC/East Africa and in South Asia. However, we feel that policy must be changed now as no evidence support fat accumulation in this group of children. Given the fact that we are already at the full word limit we hope it is acceptable that we make no further changes in the manuscript text.

4. This is a very good question. Malnutrition in itself might dispose to later non-communicable diseases (NCD) including increased susceptibility to fat accumulation, but firm evidence is lacking. However, if children with MAM are not treated, they are at a 3 times higher risk of dying here and now. The fact that they do not accumulate excess fat directly following 12 weeks treatment indicates, we believe, that the risk of later obesity following modern supplementation is, if at all present, low. The longer term implications of nutritional supplementation on NCD risk would be more likely to emerge in studies of young children supplemented over longer periods, e.g. the 9-18 month. However, the only way to find out would be doing long-term follow-up and compare our present cohort with another cohort who received different or no supplementation. Also, while getting fat later increases NCD risk, so does losing lean mass early, ie the thrifty phenotype. So, it makes sense to try to address the lean deficit when it is still amenable to nutritional therapy (early critical window). Given the fact that we are already at the full word limit we hope it is acceptable that we make no further changes in the manuscript text.

Instructions:

Please use this table format to answer the questions posed by the editors and reviewers of your paper. Copy and paste the editor/reviewer's question in the "Comments" column and your answer to that question in the corresponding "Response" column. Be sure to ALSO paste the corrected text along with your response. For minor copyediting changes such as spelling and grammar corrections, you may simply state that the error was corrected, without pasting the altered text.

* Use the page/line numbers from your revised .doc, .rtf, or .txt file; do *not* use the page/line numbers from the submission system's auto-generated PDF.

For clarity, use one row per question. Make sure to list the page and line reference where your change can be found. If no change was made, please make sure to note that in your response in addition to your reasoning. You may delete the sample row and insert rows to this table as needed.

- 1
- 2
- 3
- 4
- 5
- 6
- 7
- 8
- 9
- 10
- 11
- 12
- 13
- 14
- 15
- 16
- 17
- 18
- 19
- 20
- 21
- 22
- 23
- 24
- 25
- 26
- 27
- 28
- 29
- 30
- 31
- 32
- 33
- 34
- 35
- 36
- 37
- 38
- 39
- 40
- 41
- 42
- 43
- 44
- 45
- 46
- 47

Review Copy