Nondestructive adult age at death estimation: visualizing cementum annulations in a known age historical human assemblage using synchrotron X-ray microtomography

Adeline Le Cabec^{1,2,*}, Nancy K. Tang^{3,*}, Valentin Ruano Rubio⁴, Simon Hillson⁵

¹Department of Human Evolution, Max Planck Institute for Evolutionary Anthropology, Deutscher Platz 6, D- 04103 Leipzig, Germany. adeline.lecabec@eva.mpg.de

²ID19 Beamline, Structure of Materials Group, European Synchrotron Radiation Facility, 71 avenue des Martyrs, CS-40220, F-38043 Grenoble cedex 09, France.

³Unaffiliated, nktang@gmail.com

⁴Data Science Data Engineering, Broad Institute, 415 Main St., Cambridge, Massachusetts, USA, 02142. valentin@broadinstitute.org

⁵Institute of Archaeology, University College London, 31-34 Gordon Square, London, WC1H 0PY, United Kingdom simon.hillson.ucl@goolemail.com

*: Adeline Le Cabec and Nancy K. Tang should be considered joint first authors.

Numbers: 20 text pages plus 15 pages of bibliography, 11 figures, 4 tables, 1 graph, 8 Supplementary files.

Abbreviated title: Adult age estimation by synchrotron imaging of cementum

Corresponding authors:

Dr. Adeline Le Cabec
adeline.lecabec@eva.mpg.de
Department of Human Evolution
Max Planck Institute for Evolutionary Anthropology
Deutscher Platz 6
D- 04103 Leipzig,
Germany.

Dr. Nancy K. Tang
nktang@gmail.com

Grant sponsorship: ESRF, MPI-EVA

Conflict of Interest Statement

The authors declare no conflict of interest. All authors have read, commented on and approved the final version of this paper.

Abstract

Objectives

Adult age at death estimation continues to challenge physical anthropologists. One estimation method involves counting tooth cementum annulations (TCA). Non-destructively accessing TCA is a critical step to approaching fossil teeth of unknown age and to verifying life history profiles of human ancestors. This pilot study aims to (1) non-destructively image TCA in teeth from a known age archeological human population by propagation phase contrast X-ray synchrotron μ CT (PPC-SR- μ CT) (2) test the correlation between real and estimated ages, and the accuracy, precision and bias of age estimates.

Materials and Methods

We examine 20 permanent human canines (aged 20-81 years), from a 18th to 19th century known age collection from St. Luke’s Church (London, England). We scanned transverse segments of acellular cementum in the apical portion of the middle root third using PPC-SR- μ CT. We generated virtual transverse sections on which two observers perform two sessions of blind TCA counts. We calculate the estimated ages at death by adding 10 years to the TCA counts.

Results

A moderately strong positive linear relationship exists between real and estimated ages ($r=0.76$, $p<0.001$), with an average inaccuracy of 16.1 years and an average bias towards underestimation of 15.7 years. This difference is lower in individuals <50 years (6.8 and 6.5 years, respectively, $n=10$) compared to those >50 years (24.9 years, $n=10$).

Discussion

We reliably imaged and identified TCA in individuals <50 years from a known-age archeological sample. Scanning refinement will yield a promising alternative to current destructive methods of TCA analyses and to aid access to life history events in adult fossil hominins.

Keywords: cementochronology, adult age at death, virtual histology, propagation phase contrast

1 | INTRODUCTION

Age at death estimation is an important part of the information derived from skeletal and dental remains. On the one hand, in forensic anthropology, aging contributes to building a biological profile for the identification of missing persons (Ritz-Timme et al., 2000). On the other hand, in archeology, age estimation is more concerned with understanding the demography of past populations (see Chamberlain, 2006). Especially because there are not always written documents existing to provide this information—even in historical periods—, recovering the age at death of individuals may serve reconstructing mortality profiles at the population level in a community. Several variables can subsequently be explored such as life expectancies, fertility, growth and composition, and patterns of pathologies and epidemics. When investigating the primate fossil record, recovering the age at death of individuals allows us to better understand how the pattern and timing of dental development evolved and how this correlates with skeletal maturation within a taxon and along a lineage (e.g., Dean, 2010; Dean & Liversidge, 2015; Le Cabec, Dean, & Begun, 2017; T. M. Smith et al., 2015).

1.1 | The challenge of estimating age at death in adults

When constructing methods to track age-related changes, researchers may rely on four general states of the aging process: initiation, growth, maintenance, and degeneration (Angel, Suchey, İşcan, & Zimmerman, 1986). Estimating age at death in a developing skeleton can be based on relatively standard developmental and growth processes, such as the formation of enamel microstructure or the eruption sequence of the developing dentition (e.g., AlQahtani, Hector, & Liversidge, 2010; Dean & Beynon, 1991). After growth has stopped, adult aging methods often rely on physiological manifestations of degenerative processes, such as dental attrition and bony changes in articular surfaces, thus yielding what could be called a “biological age”. These processes are influenced by genetic, epigenetic, and environmental factors within and across populations, between sexes, and even within a given individual (Belsky et al., 2015; reviewed in Couoh, 2017; Meindl & Russell, 1998; Prince & Konigsberg, 2008; Tang, Antoine, & Hillson, 2014). The goal of adult age at death estimation is to bypass biological age, i.e., as manifested relatively and physiologically on the skeleton or teeth by the degenerative processes described above, and arrive at an absolute estimate as close as possible to the chronological (calendar) age of the individual. In spite of the efforts made to refine techniques (e.g., Boyd, Villa, & Lynnerup, 2015; Calce, 2012; Cameriere, De Luca, Alemán, Ferrante, & Cingolani, 2012; Cox, 2000; Crowder, Heinrich, & Stout, 2012; Crowder et al., 2012; Falys & Prangle, 2014; Garvin & Passalacqua, 2012; Kampan, Sinthubua, & Mahakkanukrauh, 2017; Rissech, Estabrook, Cunha,

& Malgosa, 2007; Tayles & Halcrow, 2015; Whittaker, 2000) adult age at death estimation remains the "Achilles' tendon of anthropology" (Cunha et al., 2009, p. 2). Indeed, Bocquet-Appel and Masset (1982) have suggested that more information than possible is demanded of current aging methods given the limitations of age indicators and that existing methods could at best estimate human remains into the youngest and the oldest age classes.

Because of their highly mineralized state and relatively small size, teeth are considerably resistant to decomposition processes that more rapidly affect the rest of the skeleton. Unlike bone, which undergoes regular remodeling during life, dental tissues – enamel, dentin, and cementum – are considerably more stable and grow in periodical increments which are recorded directly in the dental microstructure (Hillson & Antoine, 2003).

1.2 | Cementum: types, location and incremental growth

Cementum is composed of an inorganic component that is mainly hydroxyapatite (65% by weight or 45% by volume), an organic component that is mostly type I collagen fibers (23% by weight or 33% by volume), and water (12% by weight or 22% by volume) (Berkovitz, Holland, & Moxham, 2002). Cementum can be classified based on composition, location, and function (See reviews in Bosshardt & Schroeder, 1996; Selvig, 1964, 1965; Yamamoto, Hasegawa, Yamamoto, Hongo, & Amizuka, 2016). The two main types are primary and secondary cementum. Acellular extrinsic fibers cementum (AEFC), or primary cementum, is an avascular connective hard tissue that is incrementally deposited over time onto the dentin root surface. AEFC does not contain cementocytes, i.e., cementoblasts that get entrapped in the cementum matrix and eventually die and deteriorate leaving empty lacunae (see Yamamoto et al., 2016 for a review). In humans, it is mainly found in the cervical and middle thirds of the root, and it anchors the root surface to the periodontal ligaments and thus within the alveolar cavity (Bosshardt & Schroeder, 1991). Deposition occurs at least partially in response to providing new attachments for the periodontal ligaments, which secure the tooth in its bony socket as it is repositioned for effective occlusion during life (Foster, Nociti Jr, & Somerman, 2014; Ho et al., 2010; Ho, Marshall, Ryder, & Marshall, 2007). Acellular cementum—as to a lesser extent dentin—has no capability for repair and regeneration—and indeed, the amount of remodeling in dentin is considered negligible in biochemical studies (e.g., Beaumont, Gledhill, Lee-Thorp, & Montgomery, 2013).

Acellular cementum is most often, though not always (Huffman & Antoine, 2010; Klevezal, 1996; Klevezal, Kirillova, Shishlina, Sokolov, & Trunova, 2006), used for estimating age. Acellular extrinsic fiber cementum shows a relatively slow, continuous, and regular growth compared to secondary cementum, i.e., cellular mixed stratified cementum (CMSC) and cellular

intrinsic fibers cementum (CIFC) (Bosshardt, Luder, & Schroeder, 1989). Secondary cementum has areas containing cementocytes, tends to be less well-mineralized than primary cementum (Nanci & Bosshardt, 2006), and is found mostly where mechanical loads are highest, such as at the root apex and underneath furcations. It is generally a reactive tissue with periodic, irregular, and fast appositional growth (Schroeder, 1993). For example, cellular cementum has been shown to be responsive to extrinsic factors such as growth hormone and periodontal disease (Smid et al., 2004) whereas acellular cementum was generally unaffected.

Because AEFC grows incrementally throughout the life of the tooth (with some exceptions, see e.g., Foster et al., 2014)—i.e., also during the individual's adulthood—, it represents an invaluable quantitative adult age at death indicator over enamel and dentin. In fact, enamel ceases growth once the tooth crown is fully formed, while dentin ceases to be deposited once the pulp cavity is filled (Lieberman & Meadow, 1992). While the cyclical growth of dental hard tissues is not well understood, it has been postulated that enamel and dentin grow under the control of a master clock in the suprachiasmatic nucleus of the hypothalamus that synchronizes these rhythms to external light-dark cycles (e.g., Bromage, Hogg, Lacruz, & Hou, 2012; Bromage et al., 2016; Dean, 2017; Lacruz et al., 2012; Mahoney, Miskiewicz, Pitfield, Deter, & Guatelli-Steinberg, 2017; Papagerakis et al., 2014; Zheng et al., 2014). It may be that a biological timing mechanism involving clock genes similar to that proposed to regulate metabolic and genomic oscillations affecting circadian growth rhythms in enamel also may influence AEFC formation (see Zheng et al., 2014; Zheng, Papagerakis, Schnell, Hoogerwerf, & Papagerakis, 2011).

1.3 | The challenge of counting tooth cementum annulations (TCA) for estimating adult age at death

The layers of acellular cementum have traditionally been studied by visual inspection of physical thin sections under the microscope (methods reviewed in Naji et al., 2016 for paleodemography in human populations; in Stutz, 2002 for faunal archeological remains). With this method, cementum annulations appear as alternating light and dark bands. Based on ecological studies in marine and terrestrial wildlife (e.g., polar bears in Calvert & Ramsay, 1998; elk in Hamlin, Pac, Sime, DeSimone, & Dusek, 2000; grey seals in Hewer, 1964; and macaques in Kay & Cant, 2005; see review in Geusa et al., 1999), these bands are thought to form in a periodic rhythm resulting from seasonal changes in mineralization (Colard et al., 2016; Dean, Le Cabec, Spiers, Zhang, & Garrevoet, 2018; Großkopf & McGlynn, 2011; reviewed in Grue & Jensen, 1979; Klevezal, 1996; Klevezal & Kleinenberg, 1969; Lieberman, Deacon, & Meadow, 1990). When viewed under transmitted, cross-polarized light, the dark bands can be relatively thin. Through observations of density using Scanning Electron Microscopy (see Lieberman, 1994),

these dark bands have been attributed to hypermineralization, which in turn has been linked to a slowing down of cementoblast activity during formation in late autumn, winter, and early spring. The lighter and wider bands are thought to result from a less well-organized collagen network combined with hypomineralization that may occur during seasons of rapid tissue growth in spring and summer (Colard et al., 2016). Together, each dark and light pair is thought to represent one calendar year (Klevezal & Kleinenberg, 1969).

The biological nature of cementum-forming cells and the factors that regulate their formation are not well-known. Aging using cementum increments has been called a black-box technique from the poor understanding of their formation process (Bosshardt & Schroeder, 1996; Lieberman et al., 1990). Nonetheless, cementum annulations in non-human mammals have been successfully used in age at death estimations (Calvert & Ramsay, 1998; Hamlin et al., 2000; Hewer, 1964; Kay & Cant, 2005; Klevezal & Pucek, 1987; Pike-Tay, Cabrera Valdés, & Bernaldo de Quirós, 1999; Scheffer, 1950; Wada, Ohtaishi, & Hachiya, 1978), and in identifying events that influence the calcium and phosphate biochemistry of an organism (Großkopf & McGlynn, 2011; reviewed in Lieberman, 1994) such as hormonal cycles (Coy & Garshelis, 1992; Grue & Jensen, 1979; Kagerer & Grupe, 2001; Klevezal, 1996; Klevezal & Kleinenberg, 1969; Klevezal & Pucek, 1987; Klevezal & Stewart, 1994; Medill, Derocher, Stirling, & Lunn, 2010; Saxon & Higham, 1968), diet (Lieberman, 1993), and seasonality (Grue & Jensen, 1979; Klevezal, 1996; Klevezal et al., 2006; Klevezal & Pucek, 1987; Lieberman & Meadow, 1992; Lubinski & O'Brien, 2001).

In 1950, Gustafson included cementum thickness in his six criteria dental aging method for humans based on a qualitative scoring system (Gustafson, 1950). Since then, most studies have transitioned to counting cementum annulations, a quantitative adaptation to cementum apposition. The accuracy of using cementum annulations to estimate age at death in humans has been tested on a substantial number of known age contemporary human samples from museum collections, dentist's offices, and morgues (Avadhani, Tupkari, Khambaty, & Sardar, 2009; Charles, Condon, Cheverud, & Buikstra, 1986; Czermak, Ernst, & Grupe, 2006; Großkopf, 1990; Naylor, Miller, Stokes, & Stott, 2005; Stott, Sis, & Levy, 1982). While cementum annulations have been correlated significantly to chronological/calendar age in humans (Condon, Charles, Cheverud, & Buikstra, 1986; Couoh, 2017; Meinel et al., 2008; Obertová & Francken, 2009; Stein & Corcoran, 1994; Wittwer-Backofen & Buba, 2002; Wittwer-Backofen, Gampe, & Vaupel, 2004), their applicability to archeological material is questionable because of potential taphonomic changes resulting from the postmortem environment, such as abrasion by the alveolus or erosive effects of soil resulting in tissue removal (Charles, Condon, Cheverud, & Buikstra, 1989). Indeed, collagen leaching and apatite recrystallization can obscure or mimic true bands of cementum growth (Klevezal & Shishlina, 2001; Stutz, 2002). Only a handful of studies have

examined cementum annulations in archeological material and all of them have used teeth with unknown age (Bertrand, Robbins Schug, Polet, Naji, & Colard, 2016; Gauthier & Schutkowski, 2013; Großkopf, 1990; Huffman & Antoine, 2010; Klevezal et al., 2006; Klevezal & Shishlina, 2001; Lieberman, 1994; Robbins Schug, Brandt, & Lukacs, 2012; Roksandic, Vlak, Schillaci, & Voicu, 2009; Stutz, 2002; Wittwer-Backofen et al., 2008). Given the potential impact of diagenesis on archeological cementum annulation counts, samples should have minimal taphonomic alterations, especially in counting areas, by macroscopic and microscopic inspections. This becomes even more crucial when considering any fossil material.

Although protocols have been published for analyzing cementum increments in human teeth (e.g., Colard et al., 2016; Geusa et al., 1999; Huffman & Antoine, 2010; Wittwer-Backofen, 2012), the methodology employed across studies is not yet fully standardized. For example, studies vary in their selection of tooth type, and there is inconsistent information on whether teeth from the same individual differ in their number of annulations (Charles et al., 1986; Wittwer-Backofen et al., 2004). Keeping in mind that each tooth locus has its own developmental history—initiation, timing and pace of mineralization and eruption, loading regime and tooth use once the tooth is in functional occlusion—, examining several teeth from a given individual may contribute to constrain the reported range of age estimation. This could be applicable in the cases where identifiable stress events are visible on several teeth belonging to a single individual, and that the counts in cementum annulations in between two stress events finally match across tooth loci. However in archeological and paleoanthropological settings, investigators are limited by what is preserved and by the technique employed (e.g., destructive for classical histology vs. acquisition time, computer power and data storage needed to handle huge datasets, and sufficient access to the facilities for nondestructive techniques). Because cementum types are distributed differentially along the tooth root surface and according to tooth type, development and morphology (e.g., single- vs. multi-rooted teeth; see 1.2), the number of annulations in a given tooth has been observed to vary depending on location of sampling (Lippitsch & Grupe, 2007; Renz & Radlanski, 2006). Therefore, location of analysis along the tooth root is also a crucial consideration.

At present the only way to clearly visualize cementum annulations is through destructive techniques, which vary across investigations (reviewed in Großkopf & McGlynn, 2011). Most studies use cross-sections but some use longitudinal sections or a combination of both either singularly or in series (e.g., Aggarwal, Saxena, & Bansal, 2008; Kvaal, Kolltveit, Thomsen, & Solheim, 1995; Stein & Corcoran, 1994). Section thicknesses have ranged from 7 to 500 μm ; too thin sections may eliminate critical microstructural information whereas too thick sections may create artefacts from overlapping planes. Sections are sometimes decalcified and stained to better view the cementum growth layers (Lipsinic, Paunovich, Houston, & Robison,

1986). Analysis of the sectioned teeth can be done directly under the microscope or through digital images that allow the microstructure to be enhanced. Different microscopic techniques, such as the use of reflected or transmitted light, have led to confusion in the description of dark and light bands (reviewed in Lieberman, 1994). Various attempts at semi-automated methods of visualizing and counting annulations have not been widely adopted (Czermak et al., 2006; Lieberman et al., 1990). Investigations caution that accurate analysis of cementum annulations involves specific preparatory steps, as mentioned above, deviation from which can result in low reliability with high inter-observer variation (Colard, Bertrand, Naji, Delannoy, & Bécart, 2015; Lipsinic et al., 1986; Maat, Gerretsen, & Aarents, 2006; Renz & Radlanski, 2006).

1.4 | Towards non-destructively imaging TCA?

In fossil and subfossil teeth, sectioning may not be an option as conservation often requires that these valuable specimens remain intact (Le Cabec & Toussaint, 2017; Tafforeau et al., 2006). So far, only two techniques have attempted to image non-destructively cementum annulations: OCT (Optical Coherence Tomography) and Terahertz imaging (Leiss-Holzinger et al., 2015). Although both techniques provide 2D and 3D data, OCT has a limited spatial resolution (axial resolution at best of 1.7 μm by 4.4 μm) and a penetration depth of at most 3 mm (Leiss-Holzinger et al., 2015).

Propagation phase contrast synchrotron X-ray microtomography (PPC-SR- μCT) has established itself over the last decade as the leading imaging technique to non-destructively visualize microstructures that remain otherwise invisible with conventional microtomographic absorption techniques (Le Cabec, Tang, & Tafforeau, 2015; Sanchez, Ahlberg, Trinajstić, Mirone, & Tafforeau, 2012; Sanchez et al., 2013; Tafforeau & Smith, 2008; Zabler, 2008; Zabler, Riesemeier, Fratzl, & Zaslansky, 2006). Indeed, synchrotron X-ray imaging has been employed for the study of the developing dentition of juvenile fossil remains for age at death calculations and inferences on the evolution of primate dental development, primarily through the visualization of Retzius lines and perikymata in enamel and matching stress events in enamel and dentin in 2D and 3D images (Le Cabec et al., 2017, 2015; T. M. Smith et al., 2015, 2007, 2010; Tafforeau, Bentalab, Jaeger, & Martin, 2007; Tafforeau & Smith, 2008). Virtual histology by PPC-SR- μCT can be considered as structurally non-destructive and allows greater freedom in choosing the plane of section and section thickness, image enhancement, all of which increase flexibility in the examination of cementum annulations.

Here we extend the use of synchrotron imaging to adult age at death estimation in an attempt to non-destructively visualize cementum annulations in known age archeological modern human permanent canines. More specifically, we aim (1) to image, visualize and

correctly identify cementum increments in acellular cementum; (2) to calculate an age at death estimate from counting these increments and correlate these estimates to known age; (3) to test whether these estimates are accurate, precise, and reproducible when obtained from two observers; (4) to assess whether any observed variation between real and estimated age could be linked with the age category of these individuals. We hypothesize that we will observe a statistically significant positive correlation between the calculated age at death from annulations visualized with PPC-SR μ CT, and the actual age of individuals belonging to our British archeological population. To our knowledge, this is the first study that non-destructively examines cementum annulations for age estimation in a known age archeological population using synchrotron μ CT imaging.

2 | MATERIALS AND METHODS

2.1 | Sample

The excavation of the northern and southern burial ground and crypt at St. Luke's Church in London (England) by Oxford Archaeology in 2000 unveiled a relatively large named sample (*Departum* plate inscriptions displayed name, age, and sex of the deceased) comprising 241 individuals interred between 1760 and 1850. Of these 241 named individuals, 219 were adults, 165 of which could be associated with a known age at death ranging from 17 to 90 years. Sex of the individuals is noted based on first names but the groups are not large enough in the current study to draw sex-based conclusions. The majority of named individuals were inferred to be from the middle and upper classes (see Boyle, Boston, & Witkin, 2005 for more information).

Twenty permanent canines (maxillary and mandibular) from 20 different individuals ranging in age from 20-81 years were selected for this pilot study (Table 1). Canines were chosen because of their robustness (survivability) and tendency to have a single straight root. Teeth were selected to create a broad age range for statistical analysis. Exclusion criteria included overt macroscopic pathologies (e.g., hypercementosis) and visible postmortem change or damage to the root (e.g., presence of major cracks). Roots with minor carious lesions or evidence of periodontitis that did not inhibit measuring annulations in the middle root third were included (Couoh, 2017; de Broucker, Colard, Penel, Blondiaux, & Naji, 2016).

2.2 | Synchrotron data acquisition

The canine teeth were scanned at the European Synchrotron Radiation Facility (ESRF, Grenoble, France) on the beamline BM05 with a detector system giving a 0.615 μ m isotropic voxel size. The beamtime available within consecutive days was a major constraint which

limited the number of teeth we could scan. Therefore 20 teeth could be considered to represent a decent sample for a study based on synchrotron data. The white beam spectrum of the bending magnet was filtered using 2 mm of aluminum and 0.3 mm of tungsten (see supplementary SI1 for details on filters and detectors). We scanned transverse segments of AEFC in the apical region of the middle third of the buccal aspect of the canine root.

To define the region of interest (ROI) to scan, we divided the root transversely into six equal regions and took two scans that were vertically contiguous, each scan took about 46 minutes and covered 0.55 mm of root at the midpoint of the third region from the apex. We used an exposure time of one second per frame and 2499 projections over 360 degrees per scan. Scans were reconstructed using single distance phase retrieval process (Paganin, Mayo, Gureyev, Miller, & Wilkins, 2002) implemented in the PyHST2 (Mirone, Brun, Gouillart, Tafforeau, & Kieffer, 2014) software for application of filtered back-projection algorithm, followed by the application of a 3D unsharp mask (Sanchez et al., 2012). The resulting reconstructions were stacks of TIFF slices. Residual ring artefacts were corrected using a customized algorithm developed at the ESRF (Lyckegaard, Johnson, & Tafforeau, 2011). Scans were made anonymous through assignment of random ID numbers.

Of the two scans per tooth, one scan was chosen based on image quality from which to cut the virtual thin sections. Image stacks were imported into VGStudio Max 2.2.3 (Volume Graphics GmbH, Heidelberg, Germany). Virtual transverse sections were cut from the randomized scans. The precise alignment of the whole volume along the incremental lines is critical to improve annulation visibility while reducing the rest of the structures (Tafforeau et al., 2007). One can optimize the visualization of TCA for a given ROI in the scan, knowing that this configuration will not be applicable to the whole volume. Thus the cementum microstructure will appear blurry around this ROI. Locally the TCA can be considered as planes which are parallel between each other in 3D, but also to the cementum-dentin junction (CDJ) and to the root surface. First, one would set the center of rotation of the volume in an area where the increments are best visible. Second, the volume is orientated (by fine rotation in the three planes of space) so that, in 2D sections, the CDJ, TCA and root surface appear horizontal in the transverse axis, and roughly vertical in the sagittal axis. The frontal plane should consequently be parallel to the root surface and to the increments (see the position of each plane in 3D on Figure S1). Third, several contiguous μ CT slices (each being 0.615 μ m-thick) may be overlaid to reach a slab thickness from 4 to 15 μ m in our study (15 μ m in Figure S1). This will enhance the visibility of the TCA (Fig. S2). To note that with PPC-SR μ CT, the optimal slice thickness is much smaller than in classical histology ($\sim$ 100 μ m), as PPC-SR μ CT deals with tissue density. In addition, minute rotations in the three planes of space and a limited translation in depth (i.e., scrolling through the scan after the optimal thickness has been

determined) will help in finding the position where the increments are best visible. The scans, from which the images we counted were produced, are freely available on the ESRF Paleontological Database repository: <http://paleo.esrf.eu/index.php?/category/1613>.

2.3 | Identifying and counting cementum annulations

Cementum annulation counts were made on the randomized virtual thin sections by two observers (NKT = obs1, ALC = obs2), in Photoshop CS6 (Adobe Systems, San Jose, California, USA). These two observers each made two sets of counts per scan at least 24 hours apart.

Because the reconstruction uses single distance phase retrieval (Paganin et al., 2002), the images display fine modulations in density. This process uses a reconstruction parameter called " δ/β " which is proper to each material assumed to be quasi-homogeneous. To note that δ and β are the estimated X-ray optical properties of the sample (decrement of the X-ray refractive index, and absorption coefficient, respectively) (Paganin et al., 2002; Weitkamp, Haas, Wegrzynek, & Rack, 2011, 2013). This explains that the large and saturated white band at the surface of the root (Fig. 1) is actually a phase contrast fringe which was not retrieve during the reconstruction, because the δ/β value chosen was not adapted to account for the very different material properties existing between air and dental tissues. In the end, regarding cementum, this means that the fine dark and light pattern we see are fine variations in density in the cementum and not fringes of phase contrast.

The first crucial point is to harmonize the recognition of cementum increments on the virtual slices among observers. This was done by having both observers training together for a few counts on test images, by drawing the increments as they identified them on the section using individual layers in Photoshop. In case of discrepancies, this allowed superimposing the layers, discussing the potential reasons (e.g., cracks may affect some areas of the slice, thick phase contrast fringes are often present at the root surface), and harmonizing the definition of the spacing between TCA. Most importantly, accentuated lines should not be mistaken for TCA, as they are much fatter in appearance, and do not occur periodically nor on every slice investigated. Whether those accentuated lines could correspond to stress events remains unclear; their etiology is certainly complex, invoking multiple factors. Cipriano (2002) observed some accentuated lines in the AEFC of captive great apes kept in the northern hemisphere, and these lines were interpreted as corresponding to cold winter seasons, thus recording a thermic stress. Kagerer and Grupe (2001) observed accentuated lines in the AEFC of modern humans known to have experience pregnancies, skeletal trauma or renal diseases, which are all known to impact the calcium metabolism and could thus result in hypomineralized lines. Figure 1 illustrates the difference between fat, episodic accentuated lines (blue arrows) and regular thinner TCA (fine green brackets on the right hand side close-up). One of the challenges, in

common with classical histology, lies in identifying clear increments throughout the whole thickness of cementum from the CDJ to the root surface, potentially by counting in different places where the increments are clearest and joining those counts by following one accentuated line that runs parallel to the CDJ.

One of the goals of this pilot study is to assess whether PPC-SR μ CT may successfully resolve TCA. This would optimally involve a validation step involving physically sectioning the teeth to count the TCA using classical microscopy, and then comparing this thin section with the corresponding virtual slice of the same ROI. However, due to the uniqueness of this known age sample, it was not authorized to perform destructive analyses after the PPC-SR μ CT scanning. We will appreciate the validity of the identified increments by (1) checking the spacing between increments, which should be regular and consistent with what has been documented in other published studies (any abnormally large or small spacing should call caution, as for instance regarding the accentuated lines observed in Fig. 1), and (2) assessing the likelihood of correct identification of the TCA by detecting any systematic and consistent large offset in the age estimate in comparison to the real ages, for all age categories. For example, an old adult will have likely accumulated a more considerable thickness of cementum during their life than a very young individual—and thus the AEFC should display many more yearly increments.

2.4 | Age at death calculation

2.4.1 | Debate around the constant to add to TCA counts: eruption vs. emergence time

Perhaps one of the least explored areas of cementum annulations analysis is the constant that is added to the TCA count to achieve a final age at death estimate. Studies by convention add a value representing how long the individual existed prior to the first annulation. In general, tooth formation is thought to be a more reliable indicator than tooth emergence (Ubelaker, 1978). Therefore, the age at root completion has been occasionally used (Hillson & Antoine, 2003; Kvaal & Solheim, 1995) for calculating age at death from TCA counts. However, this proxy is imprecise as cementum begins to form on the root before root completion (Grue & Jensen, 1979). The value that is used most often is age at “eruption” of the particular tooth type, however, often without further specification (Gauthier & Schutkowski, 2013; Kasetty, Rammanohar, & Raju Ragavendra, 2010; Naylor et al., 2005). The age at eruption is complicated by various definitions of eruption in the literature (see Hurme, 1948), and by the dependence on environmental factors (i.e., malocclusion: Helm, 1968; space within the oral cavity: Liversidge, 2015b; illnesses such as HIV: Psoter et al., 2016). Tooth eruption describes the continuous process of a tooth moving from within the jaw, through the alveolar bone (alveolar emergence) and gingiva, to a functional position in the oral cavity, and subsequent

movement thereafter. Eruption 'age' therefore is inevitably an 'age range' that is marked by a brief event, clinical emergence, when the tooth emerges from the oral mucosa into the oral cavity (Liversidge, 2015a, 2015b). Eruption is most often actually referring to the age at "gingival emergence" (for living individuals or the study of orthopantomograms) or "alveolar emergence" (especially for archeological and prehistorical material: Harris, 2015, p. 143; Olze, Van Niekerk, Schulz, & Schmeling, 2007, fig. 1). Smith et al. (1994, p. 183) and Ubelaker (1978, p. 64) defined 'emergence' as any part of a cusp or crown piercing the gingiva. Grøn (1962, p. 574) saw 'emergence' more precisely as a tooth that had just pierced the gingiva but was no more than 3 mm above the gingival level. AlQahtani et al. (2010) and T.D. Smith et al. (2015) marked eruption as emergence from the alveolar bone whereas Haaviko (1970) stated that alveolar eruption occurs when the alveolar bone is completely resorbed on the occlusal side of the tooth (and thus has completely vanished).

2.4.2 | Age at AEFC initiation

Bosshardt and Schroeder (1991, 1996) documented the early stages of AEFC growth in premolars when they were at 50-60% of their final root length (also see Yamamoto et al., 2016 for a review). They further observed that the development of AEFC coincides roughly with the time of emergence of premolars in the oral cavity. This observation is relevant because many TCA studies choose mid-root height to make their counts. Sequeira et al. (1992) further demonstrated that cervical AEFC grows in thickness by 2.9 μm per year in maxillary first premolars and by ~ 1.5 μm per year in mandibular second premolars. Bosshardt and Schroeder (1996) reported an early AEFC growth rate for human premolars ranging from 1.8 to 3.4 μm per year. From their Table 2, by averaging their growth rate for levels 3 and 5 (mm apical to the CEJ) one gets an average growth rate of AEFC in the middle root third of 2.75 μm per year for upper premolars and 2.35 μm per year for lower premolars. The root of human premolars takes 5 to 7 years to form (AlQahtani et al., 2010). Coming back to the half-formed tooth root, Bosshardt and Schroeder (1991, 1996) suggest that the cusp tips will just be emerging through the gums and at that point there is 7 to 9 μm thickness of AEFC already near the cervix after 3 to 4 years of root development. As the crown erupts into occlusion over the next months and perhaps years, the root is still growing and the periodontal ligament attachment is developing. It really is not clear when the apex is complete but taking their assumptions further, it would be 2 to 3 years after this point. Thus it is common practice, though perhaps not ideal –as evidenced by the growth studies above–, to assume that the first annulation of AEFC is formed at around mean eruption age (Stott et al., 1982; Wittwer-Backofen et al., 2004). Rösing and Kvaal (1998) stressed the importance of using an ontogenetic method that adds the developmental age of the exact section position rather than fixed values for emergence age; however this has not been tested. The

years added onto the annulation count do not vary significantly depending on the value chosen (see Table 2a-b and SI 2), however the convention of using emergence ages should not be done without knowledge of its origins.

Since we are taking measurements at approximately two thirds of the root length measured from the cervix, we are more interested in developmental ages when the root is half complete and three-quarters complete. There is indeed no existing quantitative data on when the root is two-third complete. Three studies cite ages for these stages in both maxillary and mandibular canines (AlQahtani et al., 2010; Bengston, 1935; Haavikko, 1970). We used root formation times from a London-based reference population (AlQahtani et al., 2010) even though there does not seem to be overt population differences in dental maturity of permanent teeth (Braga, Heuze, Chabadel, Sonan, & Gueramy, 2005; Liversidge, 2009, 2011; Maber, Liversidge, & Hector, 2006) and thus population-specific reference data do not appear to improve estimates of dental age (Liversidge, 2015a). Specifically, we used the midpoint (10 years of age) between when the canine tooth is half complete and three-quarters complete, although we acknowledge that root extension in primates is a non-linear process; it reaches a peak and flattens off towards root completion (e.g., Le Cabec et al., 2017; Macchiarelli et al., 2006). Further we acknowledge that the London Atlas is based on radiographic data, which do not show the thinnest parts of the root and thus are known to underestimate root extension. AlQahtani et al. (2010, p. 483) justified pooling data from males and females because females preceded males in development between 6 and 14 years but usually only by one stage and not in all teeth, and this was not consistent. In general it is observed that girls are in advance of boys. This has been particularly pronounced in canines (Demirjian & Levesque, 1980). The Haavikko (1970) data for canines suggest that this sex-difference in canines does not exceed a little over a year (Table 2a-b and SI 2).

2.5 | Statistical analyses

We quantify the level of agreement between repeated measurements performed by each observer (intraobserver variation) and averages between observers (interobserver variation) using intraclass correlation coefficient (ICC) (Shrout & Fleiss, 1979). We report ICC estimates and their 95% confidence intervals and p-values calculated using a 2-way random effects model for two raters/measurements ($k=2$): ICC(2, k) or mean-rating for quantifying inter-observer absolute agreement; and a 2-way mixed effects model: ICC(3, k) or mean-rating for quantifying intra-observer consistency (Koo & Li, 2016). Both ICC(2,2) and ICC(3,2) allow to evaluate the reliability of the average measure across $k=2$ raters. For interobserver error, ICC(2,2) deals with a random set of two raters supposedly picked from a larger population (e.g., other researchers

rating cementum annulations), while ICC(3,2), for intra-observer consistency, considers both counting sessions of each rater as a fixed set of observers, and cannot be seen as a random two-observer sample from a larger population. This method has been used in published studies in this context (Ferrante & Cameriere, 2009; Meinel et al., 2008; Wong, Peng, Wen, & McGrath, 2016). We use Pearson correlation coefficient ("Pearson's r ") to assess the strength of the linear relationship between real age and estimated age (i.e., the sum of TCA counts and the estimated age of two-thirds of canine root formation at 10 years), and within observers (Kvaal & Solheim, 1995; Wittwer-Backofen et al., 2008). Accuracy of the method was assessed as the mean absolute difference between real age and estimated age (average deviation, Lucy, Pollard, & Roberts, 1995) and bias as the difference between estimated age and real age, with a negative value indicating underestimation. In order to assess the significance of any bias and the inter-rater reliability, we took the difference between average counts of both observers and we used the Wilcoxon signed paired rank test to verify whether these differences distributed around zero, indicating that the disagreement between both observer is about random. All the analyses were run using R statistical software (R Development Core Team, 2017).

3 | RESULTS

The raw data from this pilot study are summarized in Table 1.

3.1 | Repeatability in TCA counts

Within each observer, there was high consistency of the two annulation counts and thus strong precision. Intraobserver variation showed a mean absolute difference of 2.3 annulations for observer 1 (obs1; ICC(3,2) $r=0.99$; 95% confidence interval: [0.97, 0.995], $p<0.001$) and 3.3 annulations for observer 2 (obs2; ICC(3,2) $r=0.96$; 95% confidence interval: [0.90, 0.98], $p<0.001$). For both observers, these variations were lower in those under 50 years ($n=10$; 1.2 annulations for obs1 and obs2) compared to those over 50 years ($n=10$; 3.3 annulations for obs1; 5.3 annulations for obs2). Pairwise Pearson correlation comparisons showed strong linear relationships between counts for each observer ($r=0.97$ for obs1; $r=0.93$ for obs2, $p<0.001$). In two teeth, obs2 demonstrated variations in count greater than or equal to 10 annulations; when these were removed, the average absolute difference between the two sets of counts for obs1 and obs2 were nearly identical (1.9 annulations and 2.1 annulations, respectively). The lowest intraobserver variation for both observers was 0 annulation (six teeth for obs1 and two teeth for obs2) and the highest was 8 annulations for obs1 (Fig. 1) and 17 annulations for obs2 (Fig. 2).

The two counts for each observer were averaged and compared. Interobserver variation showed a mean absolute difference of 3.5 annulations (ICC(2,2) $r=0.98$; 95% confidence

interval: [0.60, 0.99, $p < 0.001$], indicating that the counts were highly correlated, and precision strong. The average difference in deviation between those under 50 years (3 annulations) and those over 50 years (3.9 annulations) was moderate. There was a strong linear relationship between observers' annulation counts ($r = 0.98$, $p < 0.001$). Interobserver variation was greatest (8 annulations) in a 58-year old male. There was perfect agreement between observers for one individual (Fig. 3), which was underestimated by 18 years. The Wilcoxon test run between the average counts of obs1 and obs2 yields a significant result ($p < 0.001$), coming from obs2 underestimating her counts as compared to obs1.

3.2 | Correlation between estimated age and real age

We have added 10 years to each annulation count representing the proxy for median age at two-thirds of root formation (Table 2a-b and SI 2; AlQahtani et al., 2010). Assuming that each annulation (i.e., a pair of dark and light bands) represents one year of life, we can establish an estimate for the age at death of each individual. Pearson correlation coefficients described the linear relationship between real and estimated ages as positive and moderately strong for both observers ($r = 0.75$ for obs1; $r = 0.77$ for obs2; $p < 0.001$). The overall average Pearson correlation coefficient was $r = 0.76$, $p < 0.001$ (Fig. 4). A linear regression model yielded a determination coefficient of $r^2 = 0.58$, which indicates that TCA counts reflect age more than would be expected by chance ($r^2 > 0.50$) but that they are not strong enough to predict real age accurately. The average correlation was higher for those under 50 years ($n = 10$, $r = 0.69$, $p = 0.03$) than those over 50 years ($n = 10$, $r = 0.50$, $p = 0.14$; the lack of significance may result from the small sample size). Indeed, there was a smaller tendency to underestimate in the under-50 age group (2-set Wilcoxon test, $p = 0.02$). On average, those under 50 years were underestimated by $18.40\% \pm 16.32\%$ and those over 50 years were underestimated by $37.53\% \pm 16.65\%$ of their actual calendar age. A larger sample size in both age groups would be necessary to confirm the validity of this observation.

3.3 | Accuracy and bias

Although a linear relationship between real and estimated ages was observed, the accuracy of estimations was low. The average absolute difference between real age and estimated age was considerably high for both observers (14.9 years for obs1; 17.3 years for obs2), resulting in an overall average inaccuracy of 16.1 years. Estimated age was closer to real age in individuals under 50 years (6.8 years) compared to those over 50 years (24.9 years) (Table 3).

Obs1 underestimated age by an average of 14.1 years (underestimating age in all but three individuals) and obs2 underestimated age by an average of 17.3 years (underestimating

age in all individuals), resulting in an overall average bias towards underestimation of 15.7 years. This difference was again smaller in the under 50 years group (6.5 years) compared to the over 50 years group (24.9 years). The lowest difference between real and estimated age was an underestimation of 0.5 years for obs1 (Fig. 5a) and an underestimation of 1.5 years for obs2 (Fig. 5b-c). The largest bias occurred in the same individual for both observers, a 77 year-old male who was underestimated by 41.5 years (obs1) and 45.5 years (obs2) (Fig. 6). To note, in the archeological report by Boyle et al. (2005), this individual was age estimated at 40-49 years using the auricular surface (Lovejoy, Meindl, Pryzbeck, & Mensforth, 1985), 46 years using the Brooks and Suchey (1990) public symphysis, 45-50 years using Todd (1920, 1921) public symphysis, and 43-55 years using sternal rib end (İşcan, Loth, & Wright, 1984, 1985) – all underestimated which does not exclude a possibility of mislabeling during excavation.

The grey plain line in Figure 4 represents the 1-to-1 match between estimated (Y-axis) and real age (X-axis), where each box represents one tooth and each dot is the average of four counts (two by each observer). The graph clearly shows that we tended to underestimate the age of the person based on the repeat measurements of two observers for each tooth. Interestingly a number of teeth seem to follow a 1-to-1 ratio between estimated and actual age: these data points falling on the grey plain line mostly involve individuals younger than 50 years of age. For those individuals, no outstanding characteristics could be elucidated in these virtual sections. The rest of the under-estimated ages seem to follow a linear model with a slope different than 1 (black dotted line on Fig. 4). A simple linear regression (Fig. 4) does not provide an adequate fit ($r^2=0.58$) due to the data following these two regimes.

4 | DISCUSSION

4.1 | Non-destructively imaging cementum annulations using synchrotron microtomography

In this pilot study, we have shown that PPC-SR- μ CT may be successfully used to non-destructively image yearly increments in acellular cementum of modern human teeth ($r = 0.76$, $p < 0.001$, $n = 20$). This preliminary result has major implications for studying fossil remains or historical specimens for which any destructive sampling is not allowed.

Some fine-tuning may intervene in the post-scanning processing steps, such as during the reconstruction of the tomographic data, during which Paganin's algorithm (2002) may already significantly improve the contrast of the structures of interest. Nonetheless, the low accuracy and general underestimation we presently obtained suggests that, while the

annulations were generally visible, many may have been unresolved or absent either within the cementum thickness or at the outer edges.

In archeological and paleoanthropological teeth, diagenetic processes can lead to confusion through false interpretations of annulation counts. For example, collagen dissolution reduces the number of bands making them look thicker, and microbes can remove outer increments giving the edge a scalloped appearance (Lieberman, 1994). In this context, a potential drawback of PPC-SR- μ CT thus far concerns the pre-screening for potential areas of interest prior to scanning. In classical histology different magnifications may be used to screen an area of interest on the whole surface of the section. Here, the volume scanned at $\sim 0.6 \mu\text{m}$ is very small, and thus may well be affected entirely by a crack or taphonomic damage. Unless a middle resolution scan ($\sim 5 \mu\text{m}$) is performed on the portion of interest of the root prior to scanning at high resolution ($\sim 0.6 \mu\text{m}$), identifying the ROI exempt of fine cracks or alterations is random. The major alterations can however be detected in radiography mode when aligning the sample.

In addition to the non-destructive nature of synchrotron scans and virtual histology, it is possible to scroll through the depth of a scan and work at variable section thicknesses to locate areas where annulations are clearest (Figs. S2-6). Furthermore, it is possible to optimize the structure visibility by orientation control (i.e., translation and rotation). Figure 7 shows an example on the canine of a 78-year old man (also see Fig. S3), in which scrolling through the scan allows to avoid damages (pits, irregularities) and optimize the visibility of the annulations for counting. Areas affected by microbial damage or mechanical loss through excavation and handling (e.g., wavy and broken edges on Fig. 8) may be circumvented either by counting in another area of the scan, or by excluding these individuals from the sample. Areas of opacity and cracks penetrating the thickness of the cementum were also observed, obscuring cementum lines in some areas (Fig. 5a). In certain cases (e.g., Fig. 9), separation or cracking of cementum from the dentin at the CDJ made it difficult to accurately count the number of lines that were within the detached zone. In some cases, root morphology may induce a wavy appearance of cementum annulations (“wavy lines”, e.g., Huffman & Antoine, 2010; Roksandic et al., 2009) at the CDJ (Fig. 10) or at the outer cementum edge (Fig. 11).

Future extensions of this exploratory work, including a validation with physical sectioning, will fine-tune the scanning setup to increase the spatial and contrast resolution at the ESRF and at other synchrotron facilities (e.g., at SLS, see Newham, Robson-Brown, Gill, & Corfe, 2017). If fossil teeth potentially containing ancient DNA were to be scanned for TCA imaging, particular attention should be made that the X-ray dose delivered remains compatible

with potential future genetic studies (see dose calculator available in SI Table 5 in Immel et al., 2016).

4.2 | Intra- and inter-observer variation

The level of experience necessary to count cementum annulations can influence how accurately age at death is estimated. One of the major shortcomings of paleodemography and forensic science is the absence of highly repeatable methods of age estimation (Charles et al., 1989). Intra- and interobserver variations in this pilot study suggest that cementum annulations from virtual thin sections are on the whole clearly presented with high ICC and Pearson correlation coefficients for both intra- and inter-observer counts. This finding contrasts results from Witter-Backofen et al. (2008) who examined increments in thin sections from an early medieval cemetery; three observers counted cementum increments and Pearson correlation coefficients of pairwise comparisons were statistically significant though relatively low ($r=0.58, 0.61, 0.72$). While experience in section preparation is required, Charles et al. (1989) argue that counting annulations produce accurate and repeatable results for those with no prior experience and a 15-minute tutorial. However, Colard et al. (2015) point out that a lack of experience, as well as teamwork (i.e., several observers training together) may impact the accuracy and precision of a counting technique, and thus the conclusions about its efficiency.

4.3 | Linear relationship between real age and estimated age

One of the benefits of cementum annulation analysis is that regression equations and thus reference populations are not needed, with the exception of deriving a constant for the first cementum deposition age (reviewed in Buckberry, 2015). Age at death can essentially be calculated from the tooth itself, even if isolated, and therefore results are not influenced by the statistical treatment of reference samples, such as the use of inverse calibration methods that contribute to overestimation of age in younger individuals and underestimation of age in older individuals (Aykroyd, Lucy, Pollard, & Solheim, 1997; Lucy, Aykroyd, & Pollard, 2002; Lucy et al., 1995).

Condon et al. (1986) suggested that TCA have a correlation strength ($r=0.78$, using 73 clinically extracted premolars) equivalent to established skeletal aging methods (Lovejoy, Meindl, Mensforth, & Barton, 1985). Meindl et al. (2008) compared three dental aging methods (Bang & Ramm, 1970; Lamendin et al., 1992, and cementum annulations) in 67 teeth. They noticed that the mean absolute difference between real age and estimated age using cementum annulations was about half that of the other two methods. The authors report a high determination coefficient, $r^2=0.85$, when regressing true age against estimated age from TCA.

Yet, overall, validations studies on modern human teeth have reported high correlations (Pearson's $r > 0.80$; see review in Naji et al., 2016) in known age collections. Finally, Wittwer-Backofen et al. (2004) obtained a very strong correlation between real age and estimated age using TCA counts for both males and females, in a sample of 363 modern freshly-extracted incisors, canines, and premolars (males: $r=0.97$, females: $r=0.98$; Wittwer-Backofen et al., 2004). This strong sex-unspecific correlation was later confirmed by Couoh (2017) ($r=0.99$).

It is however worth noting that the correlation between real and estimated ages is not consistent across cementum increment studies, and this may be explained by several factors related to study design. The first factor is using a root with a damaged surface that may be missing the last increments of life. This seems to have at least partially affected the results presented by Kasetty et al. (2010) who examined 200 formalin-preserved roots, some of which had resorbed root surfaces. They found a low correlation between real and estimated ages ($r=0.42$), also mentioning the difficulty in clearly identifying increments close to the CDJ. The second main cause for explaining the variability in correlation between real and estimated age in published results lies in the thickness of the slice generated from the tooth of interest. In a sample of 100 freshly extracted teeth, Miller et al. (1988) used 350- μm -thick cementum sections and found that 29% had obscured bands of annulations. Of the 71 remaining sections with visible TCA for counting, only 15% could be correctly placed within 10 years of real age. In addition to the thick slices used, the fact that they averaged the counts of four observers probably also account for a large part to this poor result. Last, the profile of the sample itself may affect the overall strength of the correlation. Kvaal and Solheim (1995) observed a relatively strong correlation, $r=0.84$ ($n=95$), but suggested that most of this strength came from individuals under 30 years ($r=0.93$, $n=14$), with older individuals having a greater incidence of dental disease thought to affect the age indicator.

Given the moderately strong Pearson correlation coefficient between real and estimated ages ($r=0.76$, $p<0.001$) in the present study, we may be confident that the lines visualized in these virtual thin sections are annulations rather than imaging artefacts related to μCT data acquisition or reconstruction. This linear relationship was only somewhat higher than that obtained using simpler measures of root dentin translucency on a larger number of anterior teeth from the same collection ($r=0.69$, $n=150$, Tang, 2011). Comparison against studies using tooth sections (Naji et al., 2016) suggests that the correlation observed in the present study still lies at the lower end of those obtained from non-archeological teeth. Bocquet-Appel and Masset (1982) have stated that it is not possible to estimate the death structure of a skeletal population if the correlation coefficient between real and estimated ages is lower than $r=0.90$. In this respect it may be that the data quality from the scans is insufficient (resolution) for all

annulations to be resolved. As already stated in the methods, a full validation would require a direct comparison between scans and thin section analysis on the same teeth.

4.4 | Accuracy and bias

A concerning issue in the present study is the low accuracy of estimates (large deviations between real and estimated ages) despite sufficient precision and a moderately strong linear relationship. This pattern is particularly evident in individuals over 50 years where estimates differed on average by more than two decades from real age. While individuals under 50 years had an average deviation of less than a decade, this is still not any more accurate than the broad age ranges of skeletal age estimation methods.

The trend we observe in the present study, of decreasing accuracy with increasing age, aligns to studies in modern teeth (e.g., Aggarwal et al., 2008; Kvaal & Solheim, 1995; Lipsinic et al., 1986). It has been suggested that growth layers in older individuals may get thinner (show minimal increase) or skipped altogether making them more difficult to discern (Charles et al., 1989; Gauthier & Schutkowski, 2013; Grue & Jensen, 1979; Johannson, 1971; Klevezal & Shishlina, 2001; Lipsinic et al., 1986; Rösing & Kvaal, 1998; Solheim & Sundnes, 1980; Stein & Corcoran, 1994). Kargerer and Grupe (2001) concluded that teeth of older individuals may also be more subject to pathological conditions (e.g., hypercementosis causing an increase in TCA or periodontal disease causing a decrease in TCA) and systemic disturbances to calcium metabolism (e.g., renal diseases, skeletal trauma, pregnancies causing hypomineralized lines) than those of younger individuals. Periodontal disease did not appear to affect the archeological material examined by Bertrand et al. (2016). However, in the current study, teeth with potential pathologies were excluded from analysis. In paleodemographic applications, age at death estimates for the oldest individuals in the population are particularly important in the determination of life expectancy and calculations of life history parameters such as survivorship (Wood, Holman, O'Connor, & Ferrell, 2002). Being able to accurately estimate age in older individuals is challenging due to environmental influences that increasingly affect aging features.

The quality of the tomographic data itself may at least be partially responsible for the low accuracy. One minor source of ambiguity, which could account for a difference of a few increments, lies in how to identify the first annulation. Due to the noisy aspect of the images in this area, the CDJ (Tomes' granular layer) sometimes appeared thick and unsharp (i.e., with poorly defined borders and containing frequent and closely spaced variations in grey values). Similarly, at the cementum-air border the outer increments may have been obscured by a prominent white fringe (related to the phase contrast and the δ/β value chosen for revealing material interfaces inside the tooth, and not at its surface) that was present with varying

degrees of expression in all of the scans. As it would have affected all teeth in a similar way, it could explain a regular bias with a negative offset toward the lower annulation counts. This phase fringe certainly prevented analysis of season of death, which has previously been observed in human teeth (Wedel, 2007).

4.5 | Prior assignation of age at emergence

As exposed in the Method section 2.5, various differing datasets exist on emergence times (Table 2a-b and SI 2) and their use in cementum annulation analysis is not standardized (Kagerer & Grupe, 2001; Lipsinic et al., 1986; Meinl et al., 2008; Miller et al., 1988; Obertová & Francken, 2009; Robbins Schug et al., 2012; Stein & Corcoran, 1994; Stott et al., 1982; Wittwer-Backofen et al., 2004). It should be noted however that using mean or median variation from a given study does not present a significant contribution to differences in final estimated age. However, the range of emergence can vary significantly within a population, for example in a large study of a mostly low socioeconomic group of healthy schoolchildren, mandibular canine emergence ranged broadly from 8.3 to 13.9 years in males and 7.4 to 12.3 years in females (Grøn, 1962). Emergence is also markedly influenced by physical factors such as tooth positioning within the alveolar bone, craniofacial complex remodeling, loss of deciduous teeth, malocclusion/lack of space in the dental arch, and health-related factors such as birth weight and illness (Liversidge, 2015b; Moorrees, Fanning, & Hunt Jr., 1963; Psoter et al., 2016; Wong et al., 2016). In general, and as mentioned in the methods, it is thought that tooth formation is superior to tooth emergence for assessing dental maturation. Demirjian et al. (1973, p. 212) conceded that visible emergence usually occurs when root formation is about three quarters complete, but “quite large departures from this rule have been observed.”

To calculate age at death accurately, the number of annulations should be added to the age at which the cementum has begun forming on the area of the root where the counts are made, in the case of this study when approximately two thirds of the root is complete. Root formation is a continuous process that may not occur at the same speed throughout (Dean & Cole, 2013). Thus categorizing formation into discrete stages (e.g., half complete, two-thirds complete, three-quarters complete) means that the majority of individuals do not fit any stage precisely; most are somewhere in between two stages (one is unable to extrapolate whether a tooth in one stage has just entered that stage or is approaching the next stage) and age assignment must be interpolated (Grøn, 1962; Hillson, 1996). Our own approximation at 10 years of age—i.e., the age midway between root half-complete to root three-quarters complete (see method section for details and Table 2a,b, SI2)—appears reasonable given that formation during this period is not very long (greatest duration for maxillary and mandibular canines is 1.62 years and 2.16 years, respectively, reported by Bengston, 1935, see Table 2a-b and SI2 in

present study). Fluctuations in age estimates based on these different values would be small, especially regarding archeological teeth, where the most significant deviations inevitably occur in the counting of annulations.

5 | CONCLUSION

In this pilot study, we investigated the potential of counting cementum annulations in a known age historical assemblage imaged non-destructively using phase contrast synchrotron X-ray microtomography. Because estimated ages showed a moderate positive significant linear correlation to real age ($r=0.76$, $p<0.001$), it may be inferred that virtual sections are able to identify true cementum annulations. Increment analysis showed sufficient precision but low accuracy. Therefore, in its present state of development, this method cannot be adopted for aging fossil material of unknown age but nonetheless represents a promising step in non-destructive adult age at death estimation. Future technical developments to improve imaging protocols will be beneficial in revealing the unexplored adult life history of fossil hominins.

ACKNOWLEDGEMENTS

We are thankful to Paul Tafforeau (ID 19, ESRF, France) for his support and access to in-house beamtime on beamline BM05, as well as for his suggestions on a previous version of this manuscript. We are obliged to Alexander Rack (ID 19, ESRF) for his support and advice, as well as to the ID19 and BM05 staff members who contributed to develop the technical facilities on BM05. Christopher Dean (UCL, UK) and Thomas Colard (University of Lille, France) are acknowledged for constructive discussions. Our gratitude also goes to the Max Planck Society. We are grateful to Peter Ellison, the Associate Editor, and two anonymous reviewers for their constructive comments which improved significantly this manuscript.

ORCID

Nancy K. Tang: [0000-0002-0523-2754](https://orcid.org/0000-0002-0523-2754)

Adeline Le Cabec: [0000-0001-6948-4726](https://orcid.org/0000-0001-6948-4726)

(Valentin Ruano Rubio: no ORCID)

Simon Hillson: [0000-0002-8557-0111](https://orcid.org/0000-0002-8557-0111)

REFERENCES

Aggarwal, P., Saxena, S., & Bansal, P. (2008). Incremental lines in root cementum of human teeth: An approach to their role in age estimation using polarizing microscopy. *Indian Journal of Dental Research*, 19(4), 326–330. Retrieved from <http://www.ijdr.in/text.asp?2008/19/4/326/44536>

AlQahtani, S. J., Hector, M. P., & Liversidge, H. M. (2010). Brief communication: the London atlas of human tooth development and eruption. *American Journal of Physical Anthropology*, 142(3), 481–490. doi:10.1002/ajpa.21258

Angel, J. L., Suchey, J. M., İşcan, M. Y., & Zimmerman, M. R. (1986). Age at death estimated from the skeleton and viscera. In M. R. Zimmerman & J. L. Angel (Eds.), *Dating and Age Determination of Biological Materials*. Croom Helm, London (pp. 179–220). Kent: Croom Helm Ltd.

Avadhani, A., Tupkari, J., Khambaty, A., & Sardar, M. (2009). Cementum annulations and age determination. *Journal of Forensic Dental Sciences*, 1(2), 73–76. Retrieved from <http://www.jfds.org/text.asp?2009/1/2/73/60377>

Aykroyd, R. G., Lucy, D., Pollard, A. M., & Solheim, T. (1997). Technical note: Regression analysis in adult age estimation. *American Journal of Physical Anthropology*, 104(2), 259–265. doi:10.1002/(SICI)1096-8644(199710)104:2<259::AID-AJPA11>3.0.CO;2-Z

Bang, G., & Ramm, E. (1970). Determination of age in humans from root dentin transparency. *Acta Odontologica Scandinavica*, 28(1), 3–35.

Beaumont, J., Gledhill, A., Lee-Thorp, J., & Montgomery, J. (2013). Childhood diet: a closer examination of the evidence from dental tissues using stable isotope analysis of incremental human dentine. *Archaeometry*, 55(2), 277–295. doi:10.1111/j.1475-4754.2012.00682.x

Belsky, D. W., Caspi, A., Houts, R., Cohen, H. J., Corcoran, D. L., Danese, A., ... Moffitt, T. E. (2015). Quantification of biological aging in young adults. *Proceedings of the National Academy of Sciences*, 112(30), E4104. doi:10.1073/pnas.1506264112

Bengston, R. G. (1935). A study of the time of eruption and root development of the permanent teeth between six and thirteen years. *Northwestern University Bulletin*, 35, 3–9.

Berkovitz, B. K. B., Holland, G. R., & Moxham, B. J. (Eds.). (2002). *Oral anatomy, histology and embryology*. (3rd edition). Edinburgh: New York Mosby.

Bertrand, B., Robbins Schug, G., Polet, C., Naji, S., & Colard, T. (2016). Age-at-death estimation of pathological individuals: A complementary approach using teeth cementum annulations. *International Journal of Paleopathology*, 15, 120–127. doi:10.1016/j.ijpp.2014.04.001

Bocquet-Appel, J.-P., & Masset, C. (1982). Farewell to paleodemography. *Journal of Human Evolution*, 11(4), 321–333. doi:10.1016/S0047-2484(82)80023-7

- 1
- 2
- 3 Bosshardt, D. D., Luder, H. U., & Schroeder, H. E. (1989). Rate and growth pattern of
- 4 cementum apposition as compared to dentine and root formation in a fluorochrome-
- 5 labelled monkey (*Macaca fascicularis*). *Journal de Biologie Buccale*, 17(1), 3–13.
- 6
- 7 Bosshardt, D. D., & Schroeder, H. E. (1991). Initiation of acellular extrinsic fiber cementum
- 8 on human teeth. *Cell and Tissue Research*, 263(2), 311–324.
- 9 doi:10.1007/BF00318773
- 10
- 11 Bosshardt, D. D., & Schroeder, H. E. (1996). Cementogenesis reviewed: A comparison
- 12 between human premolars and rodent molars. *The Anatomical Record*, 245(2), 267–
- 13 292. doi:10.1002/(SICI)1097-0185(199606)245:2<267::AID-AR12>3.0.CO;2-N
- 14
- 15 Boyd, K. L., Villa, C., & Lynnerup, N. (2015). The Use of CT Scans in Estimating Age at
- 16 Death by Examining the Extent of Ectocranial Suture Closure. *Journal of Forensic*
- 17 *Sciences*, 60(2), 363–369. doi:10.1111/1556-4029.12683
- 18
- 19 Boyle, A., Boston, C., & Witkin, A. (2005). *The archaeological experience at St Luke's*
- 20 *church, Old Street, Islington* ((Unpublished Report)). Oxford: Oxford Archaeology.
- 21
- 22 Braga, J., Heuze, Y., Chabadel, O., Sonan, N. K., & Gueramy, A. (2005). Non-adult dental
- 23 age assessment: correspondence analysis and linear regression versus Bayesian
- 24 predictions. *International Journal of Legal Medicine*, 119(5), 260–274.
- 25 doi:10.1007/s00414-004-0494-8
- 26
- 27 Bromage, T. G., Hogg, R. T., Lacruz, R. S., & Hou, C. (2012). Primate enamel evinces long
- 28 period biological timing and regulation of life history. *Journal of Theoretical Biology*,
- 29 305(0), 131–144. doi:10.1016/j.jtbi.2012.04.007
- 30
- 31 Bromage, T. G., Idaghdour, Y., Lacruz, R. S., Crenshaw, T. D., Ovsy, O., Rotter, B., ...
- 32 Schrenk, F. (2016). The Swine Plasma Metabolome Chronicles “Many Days”
- 33 Biological Timing and Functions Linked to Growth. *PLOS ONE*, 11(1), e0145919.
- 34 doi:10.1371/journal.pone.0145919
- 35
- 36 Brooks, S., & Suchey, J. M. (1990). Skeletal age determination based on the os pubis: A
- 37 comparison of the Acsádi-Nemeskéri and Suchey-Brooks methods. *Human Evolution*,
- 38 5(3), 227–238. doi:10.1007/BF02437238
- 39
- 40 Buckberry, J. (2015). The (mis)use of adult age estimates in osteology. *Annals of Human*
- 41 *Biology*, 42(4), 323–331. doi:10.3109/03014460.2015.1046926
- 42
- 43 Calce, S. E. (2012). A new method to estimate adult age at death using the acetabulum.
- 44 *American Journal of Physical Anthropology*, 148(1), 11–23. doi:10.1002/ajpa.22026
- 45
- 46 Calvert, W., & Ramsay, M. A. (1998). Evaluation of age determination of polar bears by
- 47 counts of cementum growth layer groups. *Ursus*, 449–453.
- 48
- 49 Cameriere, R., De Luca, S., Alemán, I., Ferrante, L., & Cingolani, M. (2012). Age estimation
- 50 by pulp/tooth ratio in lower premolars by orthopantomography. *Forensic Science*
- 51 *International*, 214(1), 105–112. doi:10.1016/j.forsciint.2011.07.028
- 52
- 53 Chamberlain, A. T. (2006). *Demography in Archaeology*. New York: Cambridge University
- 54 Press.
- 55
- 56
- 57
- 58
- 59
- 60

- Charles, D. K., Condon, K., Cheverud, J. M., & Buikstra, J. E. (1986). Cementum annulation and age determination in *Homo sapiens*. I. Tooth variability and observer error. *American Journal of Physical Anthropology*, 71(3), 311–320. doi:10.1002/ajpa.1330710306
- Charles, D. K., Condon, K., Cheverud, J. M., & Buikstra, J. E. (1989). Estimating age at death from growth layer groups in cementum. In M. Y. İşcan (Ed.), *Age Markers in the Human Skeleton*. (pp. 277–302). Springfield: Charles C. Thomas.
- Cipriano, A. (2002). Cold Stress in Captive Great Apes Recorded in Incremental Lines of Dental Cementum. *Folia Primatologica*, 73(1), 21–31. doi:10.1159/000060416
- Colard, T., Bertrand, B., Naji, S., Delannoy, Y., & Bécart, A. (2015). Toward the adoption of cementochronology in forensic context. *International Journal of Legal Medicine*, 1–8.
- Colard, T., Falgayrac, G., Bertrand, B., Naji, S., Devos, O., Balsack, C., ... Penel, G. (2016). New Insights on the Composition and the Structure of the Acellular Extrinsic Fiber Cementum by Raman Analysis. *PLoS ONE*, 11(12), e0167316. doi:10.1371/journal.pone.0167316
- Condon, K., Charles, D. K., Cheverud, J. M., & Buikstra, J. E. (1986). Cementum annulation and age determination in *Homo sapiens*. II. Estimates and accuracy. *American Journal of Physical Anthropology*, 71(3), 321–330. doi:10.1002/ajpa.1330710307
- Couoh, L. R. (2017). Differences between biological and chronological age at death in human skeletal remains: A change of perspective. *American Journal of Physical Anthropology*, 163(4), 671–695. doi:10.1002/ajpa.23236
- Cox, M. (2000). Ageing adults from the skeleton. In M. Cox & S. Mays (Eds.), *Human osteology in archaeology and forensic science* (pp. 61–82). New York: Cambridge University Press.
- Coy, P. L., & Garshelis, D. L. (1992). Reconstructing reproductive histories of black bears from the incremental layering in dental cementum. *Canadian Journal of Zoology*, 70(11), 2150–2160. doi:10.1139/z92-290
- Crowder, C., Heinrich, J., & Stout, S. D. (2012). Rib Histomorphometry for Adult Age Estimation. In L. S. Bell (Ed.), *Forensic Microscopy for Skeletal Tissues: Methods and Protocols* (pp. 109–127). Totowa, NJ: Humana Press. doi:10.1007/978-1-61779-977-8_7
- Cunha, E., Baccino, E., Martrille, L., Ramsthaler, F., Prieto, J., Schuliar, Y., ... Cattaneo, C. (2009). The problem of aging human remains and living individuals: A review. *Forensic Science International*, 193(1), 1–13. doi:10.1016/j.forsciint.2009.09.008
- Czermak, A., Ernst, H., & Grupe, G. (2006). A new method for the automated age-at-death evaluation by tooth-cementum annulation (TCA). *Anthropologischer Anzeiger*, 64(1), 25–40.
- de Broucker, A., Colard, T., Penel, G., Blondiaux, J., & Naji, S. (2016). The impact of periodontal disease on cementochronology age estimation. *International Journal of Paleopathology*, 15, 128–133. doi:10.1016/j.ijpp.2015.09.004

- Dean, M. C. (2010). Retrieving chronological age from dental remains of early fossil hominins to reconstruct human growth in the past. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 365(1556), 3397–3410. doi:10.1098/rstb.2010.0052
- Dean, M. C. (2017). How the microstructure of dentine can contribute to reconstructing developing dentitions and the lives of hominoids and hominins. *Comptes Rendus Palevol*, 16(5-6.), 557–571. doi:10.1016/j.crpv.2016.10.006
- Dean, M. C., & Beynon, A. D. (1991). Histological reconstruction of crown formation times and initial root formation times in a modern human child. *American Journal of Physical Anthropology*, 86(2), 215–228. doi:10.1002/ajpa.1330860210
- Dean, M. C., & Cole, T. J. (2013). Human Life History Evolution Explains Dissociation between the Timing of Tooth Eruption and Peak Rates of Root Growth. *PLoS ONE*, 8(1), e54534. doi:10.1371/journal.pone.0054534
- Dean, M. C., Le Cabec, A., Spiers, K., Zhang, Y., & Garrevoet, J. (2018). Incremental distribution of strontium and zinc in great ape and fossil hominin cementum using synchrotron X-ray fluorescence mapping. *Journal of The Royal Society Interface*, 15(138). doi:10.1098/rsif.2017.0626
- Dean, M. C., & Liversidge, H. M. (2015). Age estimation in fossil hominins: comparing dental development in early *Homo* with modern humans. *Annals of Human Biology*, 42(4), 415–429. doi:10.3109/03014460.2015.1046488
- Demirjian, A., Goldstein, H., & Tanner, J. M. (1973). A new system of dental age assessment. *Human Biology*, 45(2), 211–227.
- Demirjian, A., & Levesque, G.-Y. (1980). Sexual Differences in Dental Development and Prediction of Emergence. *Journal of Dental Research*, 59(7), 1110–1122. doi:10.1177/00220345800590070301
- Falys, C. G., & Prangle, D. (2014). Estimating age of mature adults from the degeneration of the sternal end of the clavicle. *American Journal of Physical Anthropology*, 156(2), 203–214. doi:10.1002/ajpa.22639
- Ferrante, L., & Cameriere, R. (2009). Statistical methods to assess the reliability of measurements in the procedures for forensic age estimation. *International Journal of Legal Medicine*, 123(4), 277–283. doi:10.1007/s00414-009-0349-4
- Foster, B. L., Nociti Jr, F. H., & Somerman, M. J. (2014). The rachitic tooth. *Endocrine Reviews*, 35(1), 1–34. doi:10.1210/er.2013-1009
- Garvin, H. M., & Passalacqua, N. V. (2012). Current Practices by Forensic Anthropologists in Adult Skeletal Age Estimation. *Journal of Forensic Sciences*, 57(2), 427–433. doi:10.1111/j.1556-4029.2011.01979.x
- Gauthier, J., & Schutkowski, H. (2013). Assessing the application of tooth cementum annulation relative to macroscopic aging techniques in an archeological sample. *HOMO - Journal of Comparative Human Biology*, 64(1), 42–57. doi:10.1016/j.jchb.2012.11.001

- Geusa, G., Bondioli, L., Capucci, E., Cipriano, A., Grupe, G., Savoré, C., & Macchiarelli, R. (1999). Dental cementum annulations and age at death estimates. Osteodental biology of people of Portus Romae (Necropolis of Isola Sacra, 2nd–3rd c. AD). In L. Bondioli & R. Macchiarelli (Eds.), *Osteodental biology of people of Portus Romae (Necropolis of Isola Sacra, 2nd–3rd c. AD)* (Vol. 2). Roma: Digital Archives of Human Paleobiology.
- Grøn, A.-M. (1962). Prediction of tooth emergence. *Journal of Dental Research*, 41(3), 573–585.
- Großkopf, B. (1990). Individualaltersbestimmung mit Hilfe von Zuwachsringen im Zement bodengelagerter menschlicher Zähne. *Zeitschrift Für Rechtsmedizin*, 103(5), 351–359. doi:10.1007/BF00204456
- Großkopf, B., & McGlynn, G. (2011). Age diagnosis based on incremental lines in dental cementum: A critical reflection. *Anthropologischer Anzeiger*, 68(3), 275–289. doi:10.1127/0003-5548/2011/0004
- Grue, H., & Jensen, B. (1979). Review of the formation of incremental lines in tooth cementum of terrestrial mammals [age determination, game animal, variation, sex, reproductive cycle, climate, region, condition of the animal]. *Danish Review of Game Biology (Denmark)*, 11, 1–48.
- Gustafson, G. (1950). Age Determinations on Teeth. *The Journal of the American Dental Association*, 41(1), 45–54. doi:10.14219/jada.archive.1950.0132
- Haavikko, K. (1970). The formation and the alveolar and clinical eruption of the permanent teeth. An orthopantomographic study. *Proceedings of the Finnish Dental Society*, 66(3), 103.
- Hamlin, K. L., Pac, D. F., Sime, C. A., DeSimone, R. M., & Dusek, G. L. (2000). Evaluating the accuracy of ages obtained by two methods for Montana ungulates. *The Journal of Wildlife Management*, 64, 441–449. doi:10.2307/3803242
- Harris, E. F. (2015). Odontogenesis. In J. D. Irish & G. R. Scott (Eds.), *A Companion to Dental Anthropology* (pp. 142–158). Chichester, UK: John Wiley & Sons, Inc. doi:10.1002/9781118845486.ch11
- Helm, S. (1968). Malocclusion in Danish children with adolescent dentition: An epidemiologic study. *American Journal of Orthodontics*, 54(5), 352–366. doi:10.1016/0002-9416(68)90304-7
- Hewer, H. R. (1964). The determination of age, sexual maturity, longevity and a life-table in the grey seal (*Halichoerus grypus*). *Proceedings of the Zoological Society of London*, 142, 593–624. doi:10.1111/j.1469-7998.1964.tb04631.x
- Hillson, S. (1996). *Dental Anthropology* (3rd edition). Cambridge: Cambridge University Press.
- Hillson, S., & Antoine, D. M. (2003). Ancient bones and teeth on the microstructural level. In G. Grupe & J. Peters (Eds.), *Deciphering ancient bones: the research potential of bioarchaeological collections*. (pp. 141–157). Leidorf: Rahden/Westf.: Verlag Marie Leidorf GmbH.

- 1
- 2
- 3 Ho, S. P., Kurylo, M. P., Fong, T. K., Lee, S. S. J., Wagner, H. D., Ryder, M. I., & Marshall,
- 4 G. W. (2010). The biomechanical characteristics of the bone-periodontal ligament-
- 5 cementum complex. *Biomaterials*, 31(25), 6635–6646.
- 6 doi:10.1016/j.biomaterials.2010.05.024
- 7
- 8 Ho, S. P., Marshall, S. J., Ryder, M. I., & Marshall, G. W. (2007). The tooth attachment
- 9 mechanism defined by structure, chemical composition and mechanical properties of
- 10 collagen fibers in the periodontium. *Biomaterials*, 28(35), 5238–5245.
- 11 doi:10.1016/j.biomaterials.2007.08.031
- 12
- 13 Huffman, M., & Antoine, D. (2010). Analysis of cementum layers in archaeological material.
- 14 *Dental Anthropology*, 23, 67–73.
- 15
- 16 Hurme, V. O. (1948). Standards of variation in the eruption of the first six permanent teeth.
- 17 *Child Development*, 19(4), 213–231. doi:10.2307/1125669
- 18
- 19 Immel, A., Le Cabec, A., Bonazzi, M., Herbig, A., Temming, H., Schuenemann, V. J., ...
- 20 Krause, J. (2016). Effect of X-ray irradiation on ancient DNA in sub-fossil bones –
- 21 Guidelines for safe X-ray imaging. *Scientific Reports*, 6, 32969.
- 22 doi:10.1038/srep32969
- 23
- 24 Işcan, M. Y., Loth, S. R., & Wright, R. K. (1984). Age estimation from the rib by phase
- 25 analysis: white males. *Journal of Forensic Science*, 29(4), 1094–1104.
- 26 doi:10.1520/JFS11776J
- 27
- 28 Işcan, M. Y., Loth, S. R., & Wright, R. K. (1985). Age estimation from the rib by phase
- 29 analysis: white females. *Journal of Forensic Science*, 30(3), 853–863.
- 30 doi:10.1520/JFS11018J
- 31
- 32 Johansson, G. (1971). Age determination from human teeth, a critical evaluation with special
- 33 consideration of change after 14 years of age. *Odontologisk Revy*, 21, 1–126.
- 34
- 35 Kagerer, P., & Grupe, G. (2001). Age-at-death diagnosis and determination of life-history
- 36 parameters by incremental lines in human dental cementum as an identification aid.
- 37 *Forensic Science International*, 118(1), 75–82. doi:10.1016/S0379-0738(00)00382-0
- 38
- 39 Kampan, N., Sinthubua, A., & Mahakkanukrauh, P. (2017). A New Method for Age
- 40 Estimation from Ectocranial Suture Closure in a Thai Population. *Siriraj Medical*
- 41 *Journal*, 66(3), 5. Retrieved from
- 42 <http://www.smj.si.mahidol.ac.th/sirirajmedj/index.php/smj/article/view/256>
- 43
- 44 Kasetty, S., Rammanohar, M., & Raju Ragavendra, T. (2010). Dental Cementum in Age
- 45 Estimation: A Polarized Light and Stereomicroscopic Study. *Journal of Forensic*
- 46 *Sciences*, 55(3), 779–783. doi:10.1111/j.1556-4029.2010.01363.x
- 47
- 48 Kay, R. F., & Cant, J. G. H. (2005). Age assessment using cementum annulus counts and
- 49 tooth wear in a free-ranging population of *Macaca mulatta*. *American Journal of*
- 50 *Primatology*, 15(1), 1–15. doi:10.1002/ajp.1350150103
- 51
- 52 Klevezal, G. A. (1996). *Recording structures of mammals. Determination of age and*
- 53 *reconstruction of life history*. Rotterdam: Brookfield VT, Balkema AA.
- 54
- 55
- 56
- 57
- 58
- 59
- 60

- 1
- 2
- 3 Klevezal, G. A., Kirillova, I. V., Shishlina, N. I., Sokolov, A. A., & Trunova, Y. E. (2006).
4 Growth layers in tooth dentin and cementum: problems and perspectives of their use
5 in the study of fossil and subfossil mammal remains including humans. *Doc.*
6 *Archaeobiol.*, 4, 113–124.
7
- 8 Klevezal, G. A., & Kleinenberg, S. E. (1969). Age determination of mammals by layered
9 structure in teeth and bone. In D. R. Sergeant (Ed.), *Fisheries Research Board of*
10 *Canada, Arctic Biological Station*. (pp. 1–142). Ste Anne de Bellevue: Fisheries
11 Research Board of Canada, Arctic Biological Station.
12
- 13 Klevezal, G. A., & Pucek, Z. (1987). Bisoniana XCIII. Growth layers in tooth cement and
14 dentine of European bison and its hybrids with domestic cattle. *Acta Theriologica*,
15 32(9), 115–128.
16
- 17 Klevezal, G. A., & Shishlina, N. I. (2001). Assessment of the season of death of ancient
18 human from cementum annual layers. *Journal of Archaeological Science*, 28(5), 481–
19 486. doi:10.1006/jasc.2000.0585
20
- 21 Klevezal, G. A., & Stewart, B. S. (1994). Patterns and calibration of layering in tooth
22 cementum of female northern elephant seals, *Mirounga angustirostris*. *Journal of*
23 *Mammalogy*, 75(2), 483–487. doi:10.2307/1382571
24
- 25 Koo, T. K., & Li, M. Y. (2016). A Guideline of Selecting and Reporting Intraclass
26 Correlation Coefficients for Reliability Research. *Journal of Chiropractic Medicine*,
27 15(2), 155–163. doi:10.1016/j.jcm.2016.02.012
28
- 29 Kvaal, S. I., Kolltveit, K. M., Thomsen, I. O., & Solheim, T. (1995). Age estimation of adults
30 from dental radiographs. *Forensic Sci. Int.*, 74, 175–185. Retrieved from
31 [http://www.sciencedirect.com/science/article/B6T6W-4002FF7-](http://www.sciencedirect.com/science/article/B6T6W-4002FF7-2/2/e063d5466d00a479ee37b43b5a4d93c6)
32 [2/2/e063d5466d00a479ee37b43b5a4d93c6](http://www.sciencedirect.com/science/article/B6T6W-4002FF7-2/2/e063d5466d00a479ee37b43b5a4d93c6)
33
34
- 35 Kvaal, S. I., & Solheim, T. (1995). Incremental lines in human dental cementum in relation to
36 age. *European Journal of Oral Sciences*, 103(4), 225–230. doi:10.1111/j.1600-
37 0722.1995.tb00164.x
38
- 39 Lacruz, R. S., Hacia, J. G., Bromage, T. G., Boyde, A., Lei, Y., Xu, Y., ... Snead, M. L.
40 (2012). The Circadian Clock Modulates Enamel Development. *Journal of Biological*
41 *Rhythms*, 27(3), 237–245. doi:10.1177/0748730412442830
42
- 43 Lamendin, H., Baccino, E., Humbert, J. F., Tavernier, J. C., Nossintchouk, R. M., & Zerilli,
44 A. (1992). A simple technique for age estimation in adult corpses: the two criteria
45 dental method. *Journal of Forensic Science*, 37(5), 1373–1379.
46 doi:10.1520/JFS13327J
47
- 48 Le Cabec, A., Dean, M. C., & Begun, D. R. (2017). Dental development and age at death of
49 the holotype of *Anapithecus hernyaki* (RUD 9) using synchrotron virtual histology.
50 *Journal of Human Evolution*, 108, 161–175. doi:10.1016/j.jhevol.2017.03.007
51
- 52 Le Cabec, A., Tang, N. K., & Tafforeau, P. (2015). Accessing developmental information of
53 fossil hominin teeth using new synchrotron microtomography-based visualization
54 techniques of dental surfaces and interfaces. *PLoS ONE*, 10(4), e0123019.
55 doi:10.1371/journal.pone.0123019
56
57
58
59
60

- Le Cabec, A., & Toussaint, M. (2017). Impacts of curatorial and research practices on the preservation of fossil hominid remains. *Journal of Anthropological Sciences*, 95, 1–28. doi:10.4436/jass.95002
- Leiss-Holzinger, E., Wiesauer, K., Stephani, H., Heise, B., Stifter, D., Kriechbaumer, B., ... Withalm, G. (2015). Imaging of the inner structure of cave bear teeth by novel non-destructive techniques. *Palaeontologia Electronica*, 18(1), 1–15.
- Lieberman, D. E. (1993). Life history variables preserved in dental cementum microstructure. *Science*, 261(5125), 1162–1164. doi:10.1126/science.8356448
- Lieberman, D. E. (1994). The biological basis for seasonal increments in dental cementum and their application to archaeological research. *Journal of Archaeological Science*, 21(4), 525–539. doi:10.1006/jasc.1994.1052
- Lieberman, D. E., Deacon, T. W., & Meadow, R. H. (1990). Computer image enhancement and analysis of cementum increments as applied to teeth of *Gazella gazella*. *Journal of Archaeological Science*, 17(5), 519–533. doi:10.1016/0305-4403(90)90033-2
- Lieberman, D. E., & Meadow, R. H. (1992). The biology of cementum increments (with an archaeological application). *Mammal Review*, 22(2), 57–77. doi:10.1111/j.1365-2907.1992.tb00120.x
- Lippitsch, A., & Grupe, G. (2007). Variability of the apposition of the acellular, extrinsic fibre cementum and its influence on the tooth cementum annulation technique in humans: The influence of physical demands and functional morphology. *Doc. Archaeobiol.*, 5, 87–112.
- Lipsinic, F. E., Paunovich, E., Houston, G. D., & Robison, S. F. (1986). Correlation of age and incremental lines in the cementum of human teeth. *Journal of Forensic Science*, 31(3), 982–989. doi:10.1520/JFS11106J
- Liversidge, H. M. (2009). Permanent tooth formation as a method of estimating age. In T. Koppe, G. Meyer, K. W. Alt, A. S. Brooks, M. C. Dean, I. Kjaer, ... M. F. Teaford (Eds.), *Comparative Dental Morphology* (Vol. 13, pp. 153–157). Karger Publishers.
- Liversidge, H. M. (2011). Similarity in dental maturation in two ethnic groups of London children. *Annals of Human Biology*, 38(6), 702–715. doi:10.3109/03014460.2011.609565
- Liversidge, H. M. (2015a). Controversies in age estimation from developing teeth. *Annals of Human Biology*, 42(4), 397–406.
- Liversidge, H. M. (2015b). Tooth Eruption and Timing. In J. D. Irish & G. R. Scott (Eds.), *A Companion to Dental Anthropology* (pp. 159–171). Chichester, UK: John Wiley & Sons, Inc. Retrieved from <https://doi.org/10.1002/9781118845486.ch12>
- Lovejoy, C. O., Meindl, R. S., Mensforth, R. P., & Barton, T. J. (1985). Multifactorial determination of skeletal age at death: A method and blind tests of its accuracy. *American Journal of Physical Anthropology*, 68(1), 1–14. doi:10.1002/ajpa.1330680102

- 1
- 2
- 3 Lovejoy, C. O., Meindl, R. S., Pryzbeck, T. R., & Mensforth, R. P. (1985). Chronological
- 4 metamorphosis of the auricular surface of the ilium: A new method for the
- 5 determination of adult skeletal age at death. *American Journal of Physical*
- 6 *Anthropology*, 68(1), 15–28. doi:10.1002/ajpa.1330680103
- 7
- 8 Lubinski, P. M., & O'Brien, C. J. (2001). Observations on Seasonality and Mortality from a
- 9 Recent Catastrophic Death Assemblage. *Journal of Archaeological Science*, 28(8),
- 10 833–842. doi:10.1006/jasc.2000.0603
- 11
- 12 Lucy, D., Aykroyd, R. G., & Pollard, A. M. (2002). Nonparametric calibration for age
- 13 estimation. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*,
- 14 51(2), 183–196. doi:10.1111/1467-9876.00262
- 15
- 16 Lucy, D., Pollard, A. M., & Roberts, C. A. (1995). A Comparison of Three Dental
- 17 Techniques for Estimating Age at Death in Humans. *Journal of Archaeological*
- 18 *Science*, 22(3), 417–428. doi:10.1006/jasc.1995.0041
- 19
- 20 Lyckegaard, A., Johnson, G., & Tafforeau, P. (2011). Correction of ring artifacts in X-ray
- 21 tomographic images. *International Journal of Tomography & Statistics*, 18(F11), 1–9.
- 22
- 23 Maat, G. J. R., Gerretsen, R. R. R., & Aarents, M. J. (2006). Improving the visibility of tooth
- 24 cementum annulations by adjustment of the cutting angle of microscopic sections.
- 25 *Forensic Science International*, 159, S95–S99. doi:10.1016/j.forsciint.2006.02.038
- 26
- 27 Maber, M., Liversidge, H. M., & Hector, M. P. (2006). Accuracy of age estimation of
- 28 radiographic methods using developing teeth. *Forensic Science International*, 159,
- 29 S68–S73. doi:10.1016/j.forsciint.2006.02.019
- 30
- 31 Macchiarelli, R., Bondioli, L., Debénath, A., Mazurier, A., Tournepiche, J.-F., Birch, W., &
- 32 Dean, M. C. (2006). How Neanderthal molar teeth grew. *Nature*, 444, 748.
- 33 doi:10.1038/nature05314
- 34
- 35 Mahoney, P., Miskiewicz, J. J., Pitfield, R., Deter, C., & Guatelli Steinberg, D. (2017).
- 36 Enamel biorhythms of humans and great apes: the Havers Halberg Oscillation
- 37 hypothesis reconsidered. *Journal of Anatomy*, 230(2), 272–281.
- 38 doi:10.1111/joa.12551
- 39
- 40 Medill, S., Derocher, A. E., Stirling, I., & Lunn, N. (2010). Reconstructing the reproductive
- 41 history of female polar bears using cementum patterns of premolar teeth. *Polar*
- 42 *Biology*, 33(1), 115–124. doi:10.1007/s00300-009-0689-z
- 43
- 44 Meindl, R. S., & Russell, K. F. (1998). Recent advances in method and theory in
- 45 paleodemography. *Annual Review of Anthropology*, 27(1), 375–399.
- 46 doi:10.1146/annurev.anthro.27.1.375
- 47
- 48 Meinl, A., Huber, C. D., Tangl, S., Gruber, G. M., Teschler-Nicola, M., & Watzek, G.
- 49 (2008). Comparison of the validity of three dental methods for the estimation of age at
- 50 death. *Forensic Science International*, 178(2), 96–105.
- 51 doi:10.1016/j.forsciint.2008.02.008
- 52
- 53 Miller, C. S., Dove, S. B., & Cottone, J. A. (1988). Failure of use of cemental annulations in
- 54 teeth to determine the age of humans. *Journal of Forensic Science*, 33(1), 137–143.
- 55 doi:10.1520/JFS12444J
- 56
- 57
- 58
- 59
- 60

- 1
- 2
- 3 Mirone, A., Brun, E., Gouillart, E., Tafforeau, P., & Kieffer, J. (2014). The PyHST2 hybrid
- 4 distributed code for high speed tomographic reconstruction with iterative
- 5 reconstruction and a priori knowledge capabilities. *Nuclear Instruments and Methods*
- 6 *in Physics Research Section B: Beam Interactions with Materials and Atoms*, 324,
- 7 41–48. doi:10.1016/j.nimb.2013.09.030
- 8
- 9 Moorrees, C. F. A., Fanning, E. A., & Hunt Jr., E. E. (1963). Age Variation of Formation
- 10 Stages for Ten Permanent Teeth. *Journal of Dental Research*, 42(6), 1490–1502.
- 11 doi:10.1177/00220345630420062701
- 12
- 13 Naji, S., Colard, T., Blondiaux, J., Bertrand, B., d’Incau, E., & Bocquet-Appel, J.-P. (2016).
- 14 Cementochronology, to cut or not to cut? *International Journal of Paleopathology*,
- 15 15, 113–119. doi:10.1016/j.ijpp.2014.05.003
- 16
- 17 Nanci, A., & Bosshardt, D. D. (2006). Structure of periodontal tissues in health and disease.
- 18 *Periodontology 2000*, 40(1), 11–28. doi:10.1111/j.1600-0757.2005.00141.x
- 19
- 20 Naylor, J. W., Miller, W. G., Stokes, G. N., & Stott, G. G. (2005). Cemental annulation
- 21 enhancement: A technique for age determination in man. *American Journal of*
- 22 *Physical Anthropology*, 68(2), 197–200. doi:10.1002/ajpa.1330680206
- 23
- 24 Newham, E., Robson-Brown, K., Gill, P., & Corfe, I. (2017). Sexual dimorphism in primate
- 25 dental cementum microstructure (p. 166). Presented at the 17th International
- 26 Symposium on Dental Morphology & 2nd congress of International Association for
- 27 Paleodontology, Bordeaux, France. Retrieved from [https://isdmiapo-](https://isdmiapo-2017.sciencesconf.org/data/ISDM_IAPO_2017_book_of_abstracts.pdf)
- 28 [2017.sciencesconf.org/data/ISDM_IAPO_2017_book_of_abstracts.pdf](https://isdmiapo-2017.sciencesconf.org/data/ISDM_IAPO_2017_book_of_abstracts.pdf)
- 29
- 30 Obertová, Z., & Francken, M. (2009). Tooth cementum annulation method: accuracy and
- 31 applicability. In T. Koppe, G. Meyer, K. W. Alt, A. S. Brooks, M. C. Dean, I. Kjaer,
- 32 ... M. F. Teaford (Eds.), *Comparative Dental Morphology* (Vol. 13, pp. 184–189).
- 33 Chichester, UK: Karger Publishers.
- 34
- 35 Olze, A., Van Niekerk, P., Schulz, R., & Schmeling, A. (2007). Studies of the Chronological
- 36 Course of Wisdom Tooth Eruption in a Black African Population. *Journal of Forensic*
- 37 *Sciences*, 52(5), 1161–1163. doi:10.1111/j.1556-4029.2007.00534.x
- 38
- 39 Paganin, D., Mayo, S. C., Gureyev, T. E., Miller, P. R., & Wilkins, S. W. (2002).
- 40 Simultaneous phase and amplitude extraction from a single defocused image of a
- 41 homogeneous object. *Journal of Microscopy*, 206(1), 33–40. doi:10.1046/j.1365-
- 42 2818.2002.01010.x
- 43
- 44 Papagerakis, S., Zheng, L., Schnell, S., Sartor, M. A., Somers, E., Marder, W., ...
- 45 Papagerakis, P. (2014). The Circadian Clock in Oral Health and Diseases. *Journal of*
- 46 *Dental Research*, 93(1), 27–35. doi:10.1177/0022034513505768
- 47
- 48 Pike-Tay, A., Cabrera Valdés, V., & Bernaldo de Quirós, F. (1999). Seasonal variations of
- 49 the Middle–Upper Paleolithic transition at El Castillo, Cueva Morín and El Pendo
- 50 (Cantabria, Spain). *Journal of Human Evolution*, 36(3), 283–317.
- 51 doi:10.1006/jhev.1998.0271
- 52
- 53
- 54
- 55
- 56
- 57
- 58
- 59
- 60

Prince, D. A., & Konigsberg, L. W. (2008). New Formulae for Estimating Age at Death in the Balkans Utilizing Lamendin's Dental Technique and Bayesian Analysis*. *Journal of Forensic Sciences*, 53(3), 578–587. doi:10.1111/j.1556-4029.2008.00713.x

Psoter, W. J., Nelson, E. M., Psoter, K. J., Dearing, B. A., Katz, R. V., & Catalanotto, F. (2016). Delayed tooth emergence in children infected with human immunodeficiency virus. *Oral Surgery, Oral Medicine, Oral Pathology and Oral Radiology*, 122(4), 442–447. doi:10.1016/j.oooo.2016.06.010

R Development Core Team. (2017). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. Retrieved from <http://www.R-project.org/>

Renz, H., & Radlanski, R. J. (2006). Incremental lines in root cementum of human teeth —A reliable age marker? *HOMO - Journal of Comparative Human Biology*, 57(1), 29–50. doi:10.1016/j.jchb.2005.09.002

Rissech, C., Estabrook, G. F., Cunha, E., & Malgosa, A. (2007). Estimation of Age at Death for Adult Males Using the Acetabulum, Applied to Four Western European Populations. *Journal of Forensic Sciences*, 52(4), 774–778. doi:10.1111/j.1556-4029.2007.00486.x

Ritz-Timme, S., Cattaneo, C., Collins, M. J., Waite, E. R., Schütz, H. W., Kaatsch, H.-J., & Borrman, H. I. M. (2000). Age estimation: The state of the art in relation to the specific demands of forensic practise. *International Journal of Legal Medicine*, 113(3), 129–136. doi:10.1007/s004140050283

Robbins Schug, G., Brandt, E. T., & Lukacs, J. R. (2012). Cementum annulations, age estimation, and demographic dynamics in Mid-Holocene foragers of North India. *HOMO - Journal of Comparative Human Biology*, 63(2), 94–109. doi:10.1016/j.jchb.2012.01.002

Roksandic, M., Vlak, D., Schillaci, M. A., & Voicu, D. (2009). Technical note: Applicability of tooth cementum annulation to an archaeological population. *American Journal of Physical Anthropology*, 140(3), 583–588. doi:10.1002/ajpa.21136

Rösing, F. W., & Kvaal, S. I. (1998). Dental age in adults—A review of estimation methods. In K. W. Alt, F. W. Rösing, & M. Teschler-Nicola (Eds.), *Dental Anthropology* (pp. 443–468). Wien: Springer-Verlag. Retrieved from https://doi.org/10.1007/978-3-7091-7496-8_22

Sanchez, S., Ahlberg, P. E., Trinajstic, K. M., Mirone, A., & Tafforeau, P. (2012). Three-dimensional synchrotron virtual paleohistology: a new insight into the world of fossil bone microstructures. *Microscopy and Microanalysis*, 18(05), 1095–1105. doi:10.1017/S1431927612001079

Sanchez, S., Dupret, V., Tafforeau, P., Trinajstic, K. M., Ryll, B., Gouttenoire, P.-J., ... Ahlberg, P. E. (2013). 3D microstructural architecture of muscle attachments in extant and fossil vertebrates revealed by synchrotron microtomography. *PLoS One*, 8(2), e56992. doi:10.1371/journal.pone.0056992

- 1
- 2
- 3 Saxon, A., & Higham, C. F. W. (1968). Identification and interpretation of growth rings in
- 4 the secondary dental cementum of *Ovis aries* L. *Nature*, 219, 634.
- 5 doi:10.1038/219634a0
- 6
- 7 Scheffer, V. B. (1950). Growth Layers on the Teeth of Pinnipedia as an Indication of Age.
- 8 *Science*, 112(2907), 309–311. Retrieved from <http://www.jstor.org/stable/1677822>
- 9
- 10 Schroeder, H. E. (1993). Human cellular mixed stratified cementum: a tissue with alternating
- 11 layers of acellular extrinsic-and cellular intrinsic fiber cementum. *Schweizer*
- 12 *Monatsschrift Fur Zahnmedizin= Revue Mensuelle Suisse d'odonto-Stomatologie=*
- 13 *Rivista Mensile Svizzera Di Odontologia e Stomatologia/SSO*, 103(5), 550–560.
- 14
- 15 Selvig, K. A. (1964). An ultrastructural study of cementum formation. *Acta Odontologica*
- 16 *Scandinavica*, 22(1), 105–120.
- 17
- 18 Selvig, K. A. (1965). The fine structure of human cementum. *Acta Odontologica*
- 19 *Scandinavica*, 23(4), 423–441.
- 20
- 21 Sequeira, P., Bosshardt, D. D., & Schroeder, H. E. (1992). Growth of acellular extrinsic fiber
- 22 cementum (AEFC) and density of inserting fibers in human premolars of adolescents.
- 23 *Journal of Periodontal Research*, 27(2), 134–142. doi:10.1111/j.1600-
- 24 0765.1992.tb01815.x
- 25
- 26 Shrout, P. E., & Fleiss, J. L. (1979). Intraclass correlations: uses in assessing rater reliability.
- 27 *Psychological Bulletin*, 86(2), 420–428. doi:10.1037/0033-2909.86.2.420
- 28
- 29 Smid, J. R., Rowland, J. E., Young, W. G., Daley, T. J., Coschigano, K. T., Kopchick, J. J., &
- 30 Waters, M. J. (2004). Mouse Cellular Cementum is Highly Dependent on Growth
- 31 Hormone Status. *Journal of Dental Research*, 83(1), 35–39.
- 32 doi:10.1177/154405910408300107
- 33
- 34 Smith, B. H., Crummett, T. L., & Brandt, K. L. (1994). Ages of eruption of primate teeth: a
- 35 compendium for aging individuals and comparing life histories. *American Journal of*
- 36 *Physical Anthropology*, 37(S19), 177–231. doi:10.1002/ajpa.1330370608
- 37
- 38 Smith, T. D., Muchlinski, M. N., Jankord, K. D., Progar, A. J., Bonar, C. J., Evans, S., ...
- 39 Deleon, V. B. (2015). Dental Maturation, Eruption, and Gingival Emergence in the
- 40 Upper Jaw of Newborn Primates. *The Anatomical Record*, 298(12), 2098–2131.
- 41 doi:10.1002/ar.23273
- 42
- 43 Smith, T. M., Tafforeau, P., Le Cabec, A., Bonnin, A., Houssaye, A., Pouech, J., ... Menter,
- 44 C. G. (2015). Dental Ontogeny in Pliocene and Early Pleistocene Hominins. *PLoS*
- 45 *ONE*, 10(2), e0118118. doi:10.1371/journal.pone.0118118
- 46
- 47 Smith, T. M., Tafforeau, P., Reid, D. J., Grün, R., Eggins, S., Boutakiout, M., & Hublin, J.-J.
- 48 (2007). Earliest evidence of modern human life history in North African early *Homo*
- 49 *sapiens*. *Proceedings of the National Academy of Sciences*, 104(15), 6128–6133.
- 50 doi:10.1073/pnas.0700747104
- 51
- 52 Smith, T. M., Tafforeau, P., Reid, D. J., Pouech, J., Lazzari, V., Zermeno, J. P., ... Hublin, J.-
- 53 J. (2010). Dental evidence for ontogenetic differences between modern humans and
- 54 Neanderthals. *Proceedings of the National Academy of Sciences of the United States*
- 55 *of America*, 107, 20923–20928. doi:10.1073/pnas.1010906107
- 56
- 57
- 58
- 59
- 60

- Solheim, T., & Sundnes, P. K. (1980). Dental age estimation of Norwegian adults — A comparison of different methods. *Forensic Science International*, 16(1), 7–17. doi:10.1016/0379-0738(80)90174-7
- Stein, T. J., & Corcoran, J. F. (1994). Pararadicular cementum deposition as a criterion for age estimation in human beings. *Oral Surgery, Oral Medicine, Oral Pathology*, 77(3), 266–270. doi:10.1016/0030-4220(94)90297-6
- Stott, G. G., Sis, R. F., & Levy, B. M. (1982). Cemental Annulation as an Age Criterion in Forensic Dentistry. *Journal of Dental Research*, 61(6), 814–817. doi:10.1177/00220345820610063401
- Stutz, A. J. (2002). Polarizing microscopy identification of chemical diagenesis in archaeological cementum. *Journal of Archaeological Science*, 29(11), 1327–1347. doi:10.1006/jasc.2001.0805
- Tafforeau, P., Bentaleb, I., Jaeger, J.-J., & Martin, C. (2007). Nature of laminations and mineralization in rhinoceros enamel using histology and X-ray synchrotron microtomography: potential implications for palaeoenvironmental isotopic studies. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 246(2), 206–227. doi:10.1016/j.palaeo.2006.10.001
- Tafforeau, P., Boistel, R., Boller, E., Bravin, A., Brunet, M., Chaimanee, Y., ... Jaeger, J. J. (2006). Applications of X-ray synchrotron microtomography for non-destructive 3D studies of paleontological specimens. *Applied Physics A: Materials Science & Processing*, 83(2), 195–202. doi:10.1007/s00339-006-3507-2
- Tafforeau, P., & Smith, T. M. (2008). Nondestructive imaging of hominoid dental microstructure using phase contrast X-ray synchrotron microtomography. *Journal of Human Evolution*, 54(2), 272–278. doi:10.1016/j.jhevol.2007.09.018
- Tang, N. K. (2011). *Root dentine translucency and its correlation to chronological age at death in archaeological human remains* (Unpublished PhD dissertation). University College London, London.
- Tang, N. K., Antoine, D., & Hillson, S. (2014). Application of the Bang and Ramm age at death estimation method to two known-age archaeological assemblages. *American Journal of Physical Anthropology*, 155(3), 332–351. doi:10.1002/ajpa.22566
- Tayles, N., & Halcrow, S. E. (2015). Age at death estimation in a sample of prehistoric Southeast Asian adolescents and adults. In M. Oxenham & H. Buckley (Eds.), *The Routledge Handbook of Bioarchaeology in Southeast Asia and the Pacific Islands* (pp. 220–238). London.
- Todd, T. W. (1920). Age changes in the pubic bone. I. The male white pubis. *American Journal of Physical Anthropology*, 3(3), 285–334. doi:10.1002/ajpa.1330030301
- Todd, T. W. (1921). Age changes in the pubic bone. *American Journal of Physical Anthropology*, 4(1), 1–70. doi:10.1002/ajpa.1330040102
- Ubelaker, D. H. (1978). *Human skeletal remains. Excavation, analysis, interpretation* (3rd edition). Washington: Taraxacum.

- 1
- 2
- 3 Wada, K., Ohtaishi, N., & Hachiya, N. (1978). Determination of age in the Japanese monkey
- 4 from growth layers in the dental cementum. *Primates*, 19(4), 775–784.
- 5 doi:10.1007/BF02373645
- 6
- 7 Wedel, V. L. (2007). Determination of Season at Death Using Dental Cementum Increment
- 8 Analysis. *Journal of Forensic Sciences*, 52(6), 1334–1337. doi:10.1111/j.1556-
- 9 4029.2007.00546.x
- 10
- 11 Weitkamp, T., Haas, D., Wegrzynek, D., & Rack, A. (2011). ANKAphase: software for
- 12 single-distance phase retrieval from inline X-ray phase-contrast radiographs. *Journal*
- 13 *of Synchrotron Radiation*, 18(4), 617–629. doi:10.1107/S0909049511002895
- 14
- 15 Weitkamp, T., Haas, D., Wegrzynek, D., & Rack, A. (2013). ANKAphase: software for
- 16 single-distance phase retrieval from inline X-ray phase-contrast radiographs. Erratum.
- 17 *Journal of Synchrotron Radiation*, 20(1), 205. doi:10.1107/S0909049512044871
- 18
- 19 Whittaker, D. K. (2000). Ageing from the dentition. In M. Cox & S. Mays (Eds.), *Human*
- 20 *osteology in archaeology and forensic science* (pp. 83–99). New York: Cambridge
- 21 University Press.
- 22
- 23 Wittwer-Backofen, U. (2012). Age Estimation Using Tooth Cementum Annulation. In L. S.
- 24 Bell (Ed.), *Forensic Microscopy for Skeletal Tissues: Methods and Protocols* (pp.
- 25 129–143). Totowa, NJ: Humana Press. doi:10.1007/978-1-61779-977-8_8
- 26
- 27 Wittwer-Backofen, U., & Buba, H. (2002). Age estimation by tooth cementum annulation:
- 28 perspectives of a new validation study. In R. D. Hoppa & J. W. Vaupel (Eds.),
- 29 *Paleodemography, Age Distributions from Skeletal Samples* (pp. 107–128).
- 30 Cambridge: Cambridge University Press.
- 31
- 32 Wittwer□Backofen, U., Buckberry, J., Czarnetzki, A., Doppler, S., Grupe, G., Hotz, G., ...
- 33 Weise, S. (2008). Basics in paleodemography: A comparison of age indicators applied
- 34 to the early medieval skeletal sample of Lauchheim. *American Journal of Physical*
- 35 *Anthropology*, 137(4), 384–396. doi:10.1002/ajpa.20881
- 36
- 37 Wittwer-Backofen, U., Gampe, J., & Vaupel, J. W. (2004). Tooth cementum annulation for
- 38 age estimation: Results from a large known-age validation study. *American Journal of*
- 39 *Physical Anthropology*, 123(2), 119–129. doi:10.1002/ajpa.10303
- 40
- 41 Wong, H. M., Peng, S.-M., Wen, Y. F., & McGrath, C. P. J. (2016). A population survey of
- 42 early factors associated with permanent tooth emergence: findings from a prospective
- 43 cohort study. *Community Dentistry and Oral Epidemiology*, 44(5), 467–475.
- 44 doi:10.1111/cdoe.12236
- 45
- 46
- 47 Wood, J. W., Holman, D. J., O'Connor, K. A., & Ferrell, R. J. (2002). Mortality models for
- 48 paleodemography. In R. D. Hoppa & J. W. Vaupel (Eds.), *Paleodemography. Age*
- 49 *distributions from skeletal samples. Cambridge Studies in Biological and*
- 50 *Evolutionary Anthropology* (Vol. 31, pp. 129–168). Cambridge: Cambridge
- 51 University Press.
- 52
- 53 Yamamoto, T., Hasegawa, T., Yamamoto, T., Hongo, H., & Amizuka, N. (2016). Histology
- 54 of human cementum: Its structure, function, and development. *Japanese Dental*
- 55 *Science Review*, 52(3), 63–74. doi:10.1016/j.jdsr.2016.04.002
- 56
- 57
- 58
- 59
- 60

Zabler, S. A. (2008). *X-ray Imaging by Partially Coherent Synchrotron Light: Application to Metallic Alloys, Tooth Dentin, and Natural Rock* (PhD Dissertation (Dr. rer. nat.)). Fakultät III - Prozesswissenschaften der Technischen Universität Berlin, Berlin, Germany. Retrieved from <https://depositonce.tu-berlin.de/handle/11303/2061>

Zabler, S. A., Riesemeier, H., Fratzl, P., & Zaslansky, P. (2006). Fresnel-propagated imaging for the study of human tooth dentin by partially coherent x-ray tomography. *Optics Express*, 14(19), 8584–8597.

Zheng, L., Ehardt, L., McAlpin, B., About, I., Kim, D., Papagerakis, S., & Papagerakis, P. (2014). The tick tock of odontogenesis. *Experimental Cell Research*, 325(2), 83–89. doi:10.1016/j.yexcr.2014.02.007

Zheng, L., Papagerakis, S., Schnell, S. D., Hoogerwerf, W. A., & Papagerakis, P. (2011). Expression of clock proteins in developing tooth. *Gene Expression Patterns*, 11(3), 202–206. doi:10.1016/j.gep.2010.12.002

FIGURE LEGENDS

Figure 1. Thomas Sowter, age at death 76 years. This individual had the highest intra-observer variation for observer two with a difference of 17 annulations between two counts. Dotted rectangle represents ROI where annulations were most clearly preserved.

Figure 2. Peter Clarke, age at death 78 years. This individual had the highest intra-observer variation for observer one with a difference of eight annulations between two counts. Dotted rectangle represents ROI for counting. Refer to Figure 1 for labeling of dental features.

Figure 3. William Wood, age at death 46 years. This individual had perfect agreement in counts between observer one and observer two, and was underestimated by 18 years. Dotted rectangle represents ROI for counting. Refer to Figure 1 for labeling of dental features.

Figure 4. Bivariate plot illustrating the moderately significant positive linear relationship between real age and estimated age. Whisker plot shows mean, maximum, and minimum age measured by the two observers. The background is shaded in light blue for individuals younger than 50 years of age, and in a darker blue otherwise. The grey plain line figures the 1-to-1 ratio, and the black dotted line the best-fit line of the data, showing an underestimation of the estimated ages in comparison with the real ages.

Figure 5. (a) John Pugh, age at death 22 years. This individual showed the lowest inaccuracy for observer one, with an average deviation between real and estimated age of 0.5 years

underestimation; (b) Thomas Giles, age at death 20 years and (c) Thomas Lan, age at death 42 years. These individuals showed the lowest inaccuracy for observer two, with an average deviation between real and estimated age of 1.5 years underestimation. The ROI for counting (dotted rectangle) was minimally affected by the ring artefacts present in (c). Refer to Figure 1 for labeling of dental features.

Figure 6. Individual (William Farmer, 77 years) showing the greatest inaccuracy for both observers, with an age at death underestimated by 41.5 years for observer one and 45.5 years for observer two. Dotted rectangle represents ROI for counting. Refer to Figure 1 for labeling of dental features.

Figure 7. Illustration of how taphonomic damage can be avoided with virtual thin sections by scrolling through the scan: (a) virtual section located 27.5 μm before the virtual section used for the TCA counting (b). Virtual sections (c) and (d) are respectively 97.2 μm and 173.9 μm beyond (b). (Peter Clarke, 78 years; red arrows point at various defects).

Figure 8. Destruction of the outer edge of cementum, producing a scalloped appearance. In this case, a more optimal plane was used for counting annulations. (William Rone, 59 years).

Figure 9. Separation of cementum from dentin at the cementum-dentin junction (red arrow). (Jane Colchett, 81 years).

Figure 10. Marked wavy appearance of the cementum-dentin junction making it difficult to determine the first cementum annulation. (Susannah Smith, 39 years).

Figure 11. Marked wavy appearance of the outer edge of the cementum (Andrew Duff, 56 years).

TABLES

Table 1. Raw data of cementum annulations counts by two observers, used for the present study from which all calculations are based.

Table 2a. Literature review of crown emergence and root development times for permanent maxillary human canines.

Table 2b. Literature review of crown emergence and root development times for permanent mandibular human canines.

Table 3. Inaccuracy and bias in age predictions for St. Luke's individuals.

SUPPLEMENTARY ONLINE INFORMATION

SI 1. Technical details of the setup used on BM05.

SI 2. Methods used in references listed in Tables 2 a and b.

Figure S1. Protocol for aligning the scan to reveal the TCA, using a 15 μm thick slab. Fine-tuning of the orientation by minute rotation and translation in the sagittal (a; red), frontal (b; green), and transverse (c; blue) planes optimizes the visibility of TCA in 3D space (d). See Section 2.2 for further details. D: dentin, CDJ: cemento-dentin junction, AEFC: acellular extrinsic fiber cementum, RS: root surface. (Peter Clarke, 78 years).

Figure S2. Impact of the slice thickness ("thick slab"; thickness values are indicated above the images, e.g., 0.615 μm corresponds to one single slice, while 12.3 μm is the superimposing of 20 slices) on the visualization of TCA. See Section 2.2 for further details. D: dentin, CDJ: cemento-dentin junction, AEFC: acellular extrinsic fiber cementum, RS: root surface. (Peter Clarke, 78 years).

Figure S3. Virtual 15 μm -thick sections (a) used for TCA counting, and (b) illustrating taphonomic damages (e.g., cracking, wavy lines, pits) 175.66 μm beyond (a). (Peter Clarke, 78 years).

Figure S4. Virtual 4 μm -thick sections (a) used for TCA counting, and (b) illustrating taphonomic damages (e.g., cracking, fractures running from the outer root surface to the CDJ) 75.26 μm beyond (a). (William Farmer, 77 years).

Figure S5. Virtual 11 μm -thick sections (a) used for TCA counting, and (b) illustrating taphonomic damages (e.g., cracking, fractures running from the outer root surface to the CDJ) 71.32 μm before (a). (William Wood, 46 years).

Figure S6. Virtual 5 μm -thick sections (a) used for TCA counting, and (b) illustrating taphonomic damages (e.g., large fracture running from the outer root surface to the CDJ) 97.67 μm beyond (a). (Jane Colchett, 81 years).

Table 1. Raw data used for the present study from which all calculations are based.

First Name	Surname	Skeletal ID	Tooth type	Sex ^a	Age at death ^b (years)	Annulation counts ^c			
						Observer 1 Count 1	Observer 1 Count 2	Observer 2 Count 1	Observer 2 Count 2
Thomas	Giles	538	LLC	M	20	10	12	9	8
John	Pugh	467	LLC	M	22	12	11	7	7
Henry	Stevens	338	LRC	M	23	8	8	7	6
Matilda	Gibson	970	ULC	F	24	11	13	9	11
Samuel	Corney	722	LRC	M	29	22	23	17	17
Susannah	Joselyn	468	LLC	F	35	16	17	13	14
Rouland	Wilkes	1223	ULC	M	38	16	20	15	14
Susannah	Smith	537	LRC	F	39	15	15	12	13
Thomas	Lan	1146	LRC	M	42	35	36	32	29
William	Wood	494	LRC	M	46	18	18	17	19
John	Lowe	497	LRC	M	53	40	44	41	31
Andrew	Duff	1071	LLC	M	56	30	30	26	30
George	Scott	1008	LLC	M	58	23	23	12	18
William	Rone	910	LLC	M	59	18	22	15	16
Ann	Fisher	1088	LRC	F	61	22	23	20	21
Thomas	Willet	699	LRC	M	61	25	25	28	24
Thomas	Sowter	361	LRC	M	76	62	56	46	63
William	Farmer	1203	LLC	M	77	23	28	21	22
Peter	Clarke	1157	LLC	M	78	35	43	35	33
Jane	Colchett	968	LLC	F	81	39	44	40	47

^ainferred from first name^bfrom departum plate inscriptions^cEstimated age at death can be calculated by adding 10 years (age at emergence of the permanent mandibular canine, see methods) to the number of annulation counted.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Table 2a. Literature review of crown emergence and root development times (years) for permanent maxillary human canines.

Study*	Crown complete		Crown emergence**		Root initiation		Root 1/4 complete		Root 1/2 complete		Root 2/3 complete		Root 3/4 complete		Root complete***			
	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M		
Bengston, 1935			11.78 ^a				8.11		9.75				11.37					
Schour and Massler, 1941	6 to 7		11 ^b		6 to 7		8 to 9		10 to 11		12 to 15		11 to 12		12 to 15			
Thoma and Goldman, 1960	6 to 7		11 to 12 ^c														13 to 15	
Halikis, 1961			10.8 ± 1.03 ^d	11.6 ± 1.29 ^d														
Haavikko, 1970	4.5	4.6	10.6 ^e	12.1 ^e			6.3	7.0	7.7	8.4			9.0	9.8	12.7	13.6		
Helm and Steidler, 1974			10.84 ± 1.25 ^f	11.45 ± 1.21 ^f														
Anderson, Thompson, & Popovich, 1976	4.1 ± 0.49	4.9 ± 0.53			4.9 ± 0.57	5.8 ± 0.61	5.9 ± 0.63	7.0 ± 0.67	7.0 ± 0.72	8.2 ± 0.76			8.2 ± 0.76	9.6 ± 0.82	11.9 ± 1.04	13.7 ± 1.28		
Ubelaker, 1978	6 to 7 ± 2		11 ± 2.5								9 ± 2				10 ± 2.5			
AlQahtani, Hector, & Liversidge, 2010	5.5		11.5 ^g		6.5		7.5 to 8.5		9.5				10.5 to 11.5		15.5 to 16.5			
Liversidge, 2016			10.9 ± 0.98 ^h														10.9 ± 0.98	

* brief explanation of study in supplementary SI2; ** variously defined; *** apex closed
^a tooth positioned at line of occlusion (alveolar eruption 10.20 years)
^b presumably gingival emergence
^c definitions of emergence/root completion unspecified
^d emergence of any part of crown into oral cavity
^e very first appearance of crown through gums (alveolar eruption: 9.3 years (females); 11.2 years (males))
^f gingival emergence
^g full eruption 12.5 years
^h alveolar eruption

Table 2b. Literature review of crown emergence and root development times (years) for permanent mandibular human canines.

Study*	Crown completion (years)		Crown emergence** (years)		Root initiation (years)		Root 1/4 complete (years)		Root 1/2 complete (years)		Root 2/3 complete (years)		Root 3/4 complete (years)		Root completion*** (years)			
	F	M	F	M	F	M	F	M	F	M	F	M	F	M	F	M		
Bengston, 1935			10.49 ^a				7.65		8.73				10.89		12.38			
Schour and Massler, 1941	6 to 7		11 ^b		6 to 7		8 to 10		10 to 11		12 to 15		11 to 12		12 to 15			
Thoma and Goldman, 1960	6 to 7		9 to 10													12 to 14 ^c		
Halikis, 1961			9.9 ± 0.97 ^d	10.7 ± 1.17 ^d														
Grøn, 1962			9.67 ± 0.82 ^e	10.91 ± 1.18 ^e					9.16	10.23			9.52	10.82	10.27 ^f	12.25 ^f		
Moorrees, Fanning, & Hunt, 1963	3.9 ± 0.5	4.0 ± 0.5					5.4 ± 0.7	5.7 ± 0.6	7.1 ± 0.8	8.1 ± 0.8			8.3 ± 0.9	9.7 ± 0.9	11.3 ± 1.2	13.1 ± 1.3		
Haavikko, 1970	4.1	4.3	9.2 ^g	10.4 ^g			6.3	6.9	7.0	8.2			8.7	9.6	11.5	13.2		
Fanning and Brown, 1971	4.39	4.53							7.94	8.58					11.18	12.67		
Helm and Steidler, 1974			9.64 ± 1.05 ^h	10.54 ± 1.13 ^h														
Anderson, Thompson, & Popovich, 1976	4.1 ± 0.49	4.8 ± 0.59			4.8 ± 0.52	5.7 ± 0.68	5.9 ± 0.60	6.9 ± 0.74	6.9 ± 0.68	8.1 ± 0.73	8.1 ± 0.79	9.4 ± 0.80			11.4 ± 0.92	13.5 ± 1.22		
Ubelaker, 1978	3 to 4 ± 2		10 ± 2.5								9 ± 2				12 ± 2.5			
Demirjian and Levesque, 1980	2.9	3.3	9.6 ^b	10.5 ^b													12.2	13.4
Smith, 1991	4.3	4.4			5.0	5.2	6.2	6.9	7.7	8.8			8.6	9.9	11.3	13.0		
Liversidge, 2009	5.61				6.38		7.47		8.63				10.03		12.23 ± 1.42			
AlQahtani, Hector, & Liversidge, 2010	5.5		9.5 ⁱ		6.5		7.5 to 8.5		9.5				10.5 to 11.5		14.5 to 16.5			
Liversidge, 2016			10.14 ± 1.14 ^j										10.14 ± 1.14					

* brief explanation of study in supplementary SI2; ** variously defined; *** apex closed

^a tooth positioned at line of occlusion (alveolar eruption 9.44 years)

^b presumably gingival emergence

^c definitions of emergence/root completion unspecified

^d emergence of any part of crown into oral cavity

^e just pierced the gingiva but is no more than 3 mm above the gingival level, estimated from the tip of the cusp or from the incisal margin

^f root complete but apex open

^g very first appearance of crown through gums (alveolar eruption: 8.8 years (females); 9.8 years (males))

^h gingival emergence

ⁱ full eruption 11.5 years

^j alveolar eruption

Table 3. Inaccuracy and bias in age predictions for St. Luke’s Individuals

Age Class	n	Inaccuracy ^a	Bias ^b
20-29	5	2.5	-2.2
30-49	5	11.2	-10.8
50-59	4	20.3	-20.3
60-89	6	28.0	-28.0

^a $\Sigma(|\text{estimated age} - \text{real age}|)/n$

^b $\Sigma(\text{estimated age} - \text{real age})/n$

Figure 1 - Thomas Sowter, age at death 76 years. This individual had the highest intra-observer variation for observer two with a difference of 17 annulations between two counts. Dotted rectangle represents ROI where annulations were most clearly preserved.

58x22mm (300 x 300 DPI)

Figure 2 - Peter Clarke, age at death 78 years. This individual had the highest intra-observer variation for observer one with a difference of eight annulations between two counts. Dotted rectangle represents ROI for counting. Refer to Figure 1 for labeling of dental features.

55x18mm (300 x 300 DPI)

Figure 3 - William Wood, age at death 46 years. This individual had perfect agreement in counts between observer one and observer two, and was underestimated by 18 years. Dotted rectangle represents ROI for counting. Refer to Figure 1 for labeling of dental features.

918x213mm (96 x 96 DPI)

Figure 4 - Bivariate plot illustrating the moderately significant positive linear relationship between real age and estimated age. Whisker plot shows mean, maximum, and minimum age measured by the two observers. The background is shaded in light blue for individuals younger than 50 years of age, and in a darker blue otherwise. The grey plain line figures the 1-to-1 ratio, and the black dotted line the best-fit line of the data, showing an underestimation of the estimated ages in comparison with the real ages.

167x152mm (300 x 300 DPI)

Figure 5 - (a) John Pugh, age at death 22 years. This individual showed the lowest inaccuracy for observer one, with an average deviation between real and estimated age of 0.5 years underestimation; (b) Thomas Giles, age at death 20 years and (c) Thomas Lan, age at death 42 years. These individuals showed the lowest inaccuracy for observer two, with an average deviation between real and estimated age of 1.5 years underestimation. The ROI for counting (dotted rectangle) was minimally affected by the ring artefacts present in (c). Refer to Figure 1 for labeling of dental features.

254x150mm (300 x 300 DPI)

Figure 6 - Individual (William Farmer, 77 years) showing the greatest inaccuracy for both observers, with an age at death underestimated by 41.5 years for observer one and 45.5 years for observer two. Dotted rectangle represents ROI for counting. Refer to Figure 1 for labeling of dental features.

254x66mm (300 x 300 DPI)

Figure 7 - Illustration of how taphonomical damage can be avoided with virtual thin sections by scrolling through the scan: (a) virtual section located 27.5 μm before the virtual section used for the TCA counting (b). Virtual sections (c) and (d) are respectively 97.2 μm and 173.9 μm beyond (b). (Peter Clarke, 78 years; red arrows point at various defects).

149x201mm (300 x 300 DPI)

Figure 8 - Destruction of the outer edge of cementum, producing a scalloped appearance. In this case, a more optimal plane was used for counting annulations. (William Rone, 59 years).

254x59mm (300 x 300 DPI)

Figure 9 - Separation of cementum from dentin at the cementum-dentin junction (red arrow). (Jane Colchett, 81 years).

67x22mm (300 x 300 DPI)

Figure 10 - Marked wavy appearance of the cementum-dentin junction making it difficult to determine the first cementum annulation. (Susannah Smith, 39 years).

254x44mm (300 x 300 DPI)

Figure 11 - Marked wavy appearance of the outer edge of the cementum (Andrew Duff, 56 years).

254x63mm (300 x 300 DPI)

Figure S1 - Protocol for aligning the scan to reveal the TCA, using a 0.01 mm thick slab. Fine-tuning of the orientation by minute rotation and translation in the sagittal (a; red), frontal (b; green), and transverse (c; blue) planes optimizes the visibility of TCA in 3D space (d). See Section 2.2 for further details. D: dentin, CDJ: cemento-dentin junction, AEFC: acellular extrinsic fiber cementum, RS: root surface. (Peter Clarke, 78 years).

194x188mm (300 x 300 DPI)

Figure S2. Impact of the slice thickness ("thick slab"; various value indicated above the image, e.g., 0.615 μm corresponds to one single slice, while 12.3 μm is the superimposing of 20 slices) on the visualization of TCA. See Section 2.2 for further details. D: dentin, CDJ: cemento-dentin junction, AEFC: acellular extrinsic fiber cementum, RS: root surface. (Peter Clarke, 78 years).

241x292mm (300 x 300 DPI)

Figure S3 - Virtual 15 μm -thick sections (a) used for TCA counting, and (b) illustrating taphonomical damages (e.g., cracking, wavy lines, pits) 175.66 μm beyond (a). (Peter Clarke, 78 years).

135x92mm (300 x 300 DPI)

Figure S4 - Virtual 4 μm -thick sections (a) used for TCA counting, and (b) illustrating taphonomical damages (e.g., cracking, fractures running from the outer root surface to the CDJ) 75.26 μm beyond (a). (William Farmer, 77 years).

126x80mm (300 x 300 DPI)

Figure S5 - Virtual 11 µm-thick sections (a) used for TCA counting, and (b) illustrating taphonomical damages (e.g., cracking, fractures running from the outer root surface to the CDJ) 71.32 µm before (a). (William Wood, 46 years).

99x49mm (300 x 300 DPI)

Figure S6 - Virtual 5 μm -thick sections (a) used for TCA counting, and (b) illustrating taphonomical damages (e.g., large fracture running from the outer root surface to the CDJ) 97.67 μm beyond (a). (Jane Colchett, 81 years).

139x97mm (300 x 300 DPI)