Brain size and organization in the Middle Pleistocene hominins from Sima de los Huesos.

Inferences from endocranial variation

Eva María Poza-Rey^{a,b}, Aida Gómez-Robles^{c,d,e,*}, Juan Luis Arsuaga^{a,b}

^a Centro Mixto UCM-ISCIH de Evolución y Comportamiento Humanos, Instituto de Salud

Carlos III, 28029 Madrid, Spain

^b Departamento de Paleontología, Facultad de Ciencias Geológicas, Universidad Complutense

de Madrid, 28040 Madrid, Spain

^c Department of Anthropology, University College London, London WC1E 0BW, UK.

^d Department of Genetics, Evolution and Environment, University College London, London

WC1E 6BT, UK.

^e Department of Life Sciences, Natural History Museum, London SW7 5BD, UK.

* Corresponding author: a.gomez-robles@ucl.ac.uk (A. Gómez-Robles)

Acknowledgements

We are indebted to all the members of the Sima de los Huesos excavation team for the invaluable work they have carried out for many years to recover the SH hominin remains. We are also grateful to Ana Gracia and María Cruz Ortega for the time and energy spent in preserving and restoring the SH collection. Laura Rodríguez and Elena Santos graciously scanned most of the SH cranial collection. Henry de Lumley, Dominique Grimaud-Hervé, Philippe Menecier, Aurelie Fort, Odile Romain, Amelie Vialet and Anna Echassoux facilitated access to the comparative endocranial collections deposited in Paris and Nice. Mark

Grabowski generously shared with us unpublished data on individual encephalization quotients, and Emiliano Bruner provided images of *H. erectus* and Neanderthal endocasts. We are grateful to Mike Plavcan, the associate editor, Simon Neubauer, Antoine Balzeau and one anonymous reviewer for their constructive feedback on our manuscript. This research has been funded by the Dirección General de Investigación of Ministerio de Economía y Competitividad of Spain (CGL-2009-12703-C03-01, CGL2012-38434-C03-01, CGL2012-38434-C03-03) and the Comunidad de Madrid (S2010/BMD-2330 I+D project). E.M.P.-R. has been funded by Fundación Atapuerca. A.G.-R. is funded by the UCL-Excellence program.

Brain size and organization in the Middle Pleistocene hominins from Sima de los Huesos.

Inferences from endocranial variation

Abstract

The Sima de los Huesos (SH) endocranial sample includes 16 complete or partial endocasts corresponding to European Middle Pleistocene hominins. Different anatomical and molecular studies have demonstrated that these hominins are phylogenetically related to Neanderthals, thus making them the earliest unquestionable representatives of the Neanderthal lineage. The description of endocranial variation in this population is fundamental to shedding light on the evolution of the Neanderthal brain. In this contribution, we analyze and describe endocranial variation in this sample, including aspects related to brain size (endocranial volume and encephalization) and brain organization (through qualitative descriptions and quantitative analyses). Our results indicate that the SH hominins show a transitional state between a primitive hominin endocranial configuration (which is found in *Homo erectus* and non-SH Middle Pleistocene *Homo*) and the derived configurations found in Neanderthals and modern humans, without a clear anticipation of classic Neanderthal endocranial traits. In comparison with other cranial and postcranial traits that show a fully Neanderthal or clear pre-Neanderthal condition in the SH collection, endocranial variation in these hominins is surprisingly primitive and shows no Neanderthal affinity. These results and the comparison with other cranial traits confirm that Neanderthals evolved in a mosaic fashion. Traits related to mastication (dental, facial and mandibular anatomy) led the Neanderthalization process, whereas neurocranial anatomy must have acquired a fully Neanderthal condition considerably later.

Introduction

26

27 Studying endocranial morphology is fundamental to inferring key aspects of human brain
28 evolution. Similarities in brain size and general brain proportions between chimpanzees and
29 early hominins (as inferred from endocranial anatomy) indicate that the brain of the last
30 common ancestor of the chimpanzees and humans was more similar to the chimpanzee brain.
31 Therefore, a dramatic modification in brain size and organization must have happened in the
32 human lineage after its separation from the chimpanzee lineage, which is also inferred from
33 endocranial variation in fossil hominins. Detailed study of the hominin fossil record has
34 allowed researchers to describe in some detail endocranial variation in australopiths (Falk,
35 1980, 1983, 2014; Holloway, 1981a, 1983; Holloway et al., 2004a, 2004b; Carlson et al., 2011;
36 Beaudet et al., 2018; Falk et al., 2018), early *Homo* (Tobias, 1987; Spoor et al., 2015), *Homo*
37 *erectus* and *Homo ergaster* (Grimaud-Hervé, 1997; Coqueugniot et al., 2004; Balzeau et al.,
38 2005; Wu et al., 2010; Bruner et al., 2015; Beaudet and Bruner, 2017), *Homo floresiensis* (Falk
39 et al., 2005; Kubo et al., 2013), classic Neanderthals (Holloway, 1981b; Grimaud-Hervé, 1997;
40 Bruner et al., 2003; Bruner and Manzi, 2008; Gunz et al., 2010; Ponce de León et al., 2016)
41 and early and recent modern humans (Hublin et al., 2017; Neubauer et al., 2018), and to make
42 some inferences regarding their brain anatomy and their behavior.

43

44 Other periods of human evolution, however, remain poorly understood. Among them, patterns
45 of variation in Middle Pleistocene hominins, usually referred to as *H. heidelbergensis*, remain
46 particularly elusive. This is due, at least in part, to the lack of consensus on the relationships
47 and the putative evolutionary cohesion (or lack thereof) of different Middle Pleistocene groups
48 (Tattersall, 2007; Mounier et al., 2009; Stringer, 2012; Gómez-Robles et al., 2013a). This lack
49 of consensus primarily concerns the relationships between European and African fossils, as
50 different lines of evidence indicate that these European and African populations may be already

part of the Neanderthal and modern human lineages, respectively (see below). In addition, it has been argued that European Pleistocene hominins may include two different evolutionary lineages, one of which would be phylogenetically more closely related to Neanderthals than the other (Tattersall, 2007; Roksandic et al., 2018). A key sample that has a fundamental role to inform this debate is the Middle Pleistocene collection from the Sima de los Huesos site (Sierra de Atapuerca, Burgos, Spain).

The Sima de los Huesos (SH) hominin sample comprises a minimum of 28 individuals as determined through dental associations (Bermúdez de Castro and Nicolás, 1997; Bermudez de Castro et al., 2004; Arsuaga et al., 2014). Sixteen individuals preserve complete or fragmentary cranial remains and form the SH endocranial (and cranial) collection (Arsuaga et al., 2014; Poza-Rey, 2015). Reliable associations have been established between six dental and cranial individuals, whereas the remaining dental individuals are not associated with their cranial counterparts (Arsuaga et al., 2014; Poza-Rey et al., 2017). All these hominins were found in a single pit site and in the same stratigraphic level (Arsuaga et al., 2014; Aranburu et al., 2017), suggesting that they belonged to the same biological population. The most recent analyses of the site indicate that the fossil hominins are dated to 430 ka (Arnold et al., 2014). This age is younger than the ca. 600 ka indicated by earlier analyses (Bischoff et al., 2007), but older than the 350 ka suggested initially for this sample (Bischoff et al., 2003). Recent molecular studies of a bone fragment from this sample have shown that the SH hominins are phylogenetically related to Neanderthals (Meyer et al., 2014, 2016), a result that confirms the Neanderthal affinity indicated by a multitude of anatomical studies of this sample published for more than 25 years (e.g., Arsuaga et al., 1993, 1997). Indeed, studies of the cranial and postcranial anatomy of these hominins show a transitional morphology between a more primitive hominin anatomy and a classic Neanderthal condition in most traits (Arsuaga et al., 1993, 1997, 2014,

2015; Carretero et al., 1997; Martínez and Arsuaga, 1997; Rosas, 1997; Gómez-Olivencia et al., 2007; Bonmatí et al., 2010; Rodríguez et al., 2016; Pablos et al., 2017). Studies of the SH teeth, however, have shown a surprisingly derived dental anatomy that is virtually indistinguishable from a classic Neanderthal condition for discrete traits (Martínón-Torres et al., 2012) and even more derived than the classic Neanderthal condition in terms of dental size (Bermudez de Castro and Nicolas, 1995) and structural reduction in the number and size of individual cusps (Gómez-Robles et al., 2015a). The eclectic combination of fully Neanderthal and transitional traits of this population indicates that the process of acquiring a classic Neanderthal anatomy has occurred in a mosaic fashion (Arsuaga et al., 2014), with different traits evolving at different rates in response to different selective pressures (Gómez-Robles et al., 2017).

Given the role that cognitive advancements must have had during Neanderthal evolution, it is particularly important to describe the evolution of the brain in the early representatives of this lineage as inferred from endocranial variation. The SH sample offers not only an incomparable glimpse into the evolution of the early Neanderthal brain, but also documents intrapopulation variation because of its large sample size. Recently, endocranial asymmetry of the SH hominins has been assessed in association with behavioral lateralization as inferred from labial striations on the anterior dentition (Poza-Rey et al., 2017). In addition, some particular aspects of endocranial variation in some individuals of the sample have been described in previous studies (Bookstein et al., 2000; Bruner et al., 2003; Arsuaga et al., 2014). However, a comprehensive descriptive analysis of this collection has not yet been published. The purpose of the present contribution is to provide such a study, including descriptions of major anatomical traits, size and morphometric variation of the SH endocranial collection and comparisons with other European, African and Asian hominins.

It has to be noted that inferences about brain anatomy based on endocranial variation are subject to different interpretations and lead to occasional controversy (e.g., Holloway, 1981a; Falk, 1983). Brain sulci do not leave unequivocal marks on the endocranial surface, particularly in older individuals, thus making the interpretation of sulcal imprints particularly challenging and open to debate. Behavioral and cognitive inferences based on endocranial variation are even more problematic and very often impossible to test. Other aspects of brain variation, such as size, general proportions and the relative expansion or reduction of certain areas, are much less problematic and can be inferred from endocranial size, from different sets of linear metrics, or from geometric morphometric configurations of landmarks. Therefore, the present study focuses on the anatomical description of the SH endocranial collection with special emphasis on the variation of endocranial size and shape while acknowledging that these are but imperfect proxies for the evolution of brain size and organization.

Material and methods

Material

The SH endocranial sample comprises 16 endocasts obtained from the preserved crania in the sample. Some of these crania are incomplete and fragmentary, so the completeness and preservation of endocasts differs widely. These endocasts were given the same number as their corresponding crania, from cranium 2 to cranium 17. Crania 4 and 5 were scanned in 1999 with a CT Scanner Toshiba Xpress/GX, using a slice thickness of 1 mm, slice interval of 1 mm and pixel size of 0.468 mm. All the other crania from the Sima de los Huesos sample were scanned

in 2006 with an industrial CT Scanner MU 2000-CT YXLON International. For these crania, slice thickness was 0.5 mm, slice interval 1 mm and pixel size 0.217 mm (Poza-Rey, 2015).

This sample was compared with a sample of other Middle and Late Pleistocene hominin endocasts which, according to a splitting taxonomy (Wood and Boyle, 2016), belong to *H. erectus*, *Homo heidelbergensis* and *Homo rhodesiensis* (here together referred to as Middle Pleistocene *Homo*), *Homo neanderthalensis* and *Homo sapiens*. Details of the specimens included in the quantitative analyses are provided in Table 1, but additional specimens are mentioned in the qualitative comparisons. Some of those comparisons come from personal observations, whereas others come from published studies.

Methods

General qualitative descriptions of the Sima de los Huesos endocasts are provided, which include the description of the completeness of the specimens, an assessment of Broca's cap asymmetry and of petalial patterns (LeMay, 1976; LeMay et al., 1982), a description of the projection of the encephalic rostrum (Grimaud-Hervé, 1997) and of the shape of the occipital lobe (Fig. 1). The assessment of endocranial asymmetries is based on previous studies (Poza-Rey, 2015; Poza-Rey et al., 2017). As described in further detail in Poza-Rey et al. (2017), the assessment of petalial patterns relies on linear metrics, particularly on the hemispheric differences in the length and width of the frontal and occipital poles measured at the points marking 10% of the total length (Fig. 2). The asymmetry of Broca's cap is based on the comparison of left and right areas as measured following the sulcal boundaries of the pars triangularis (anterior horizontal and ascending rami of the Sylvian fissure and inferior frontal sulcus) and of the pars opercularis (anterior vertical ramus of the Sylvian fissure, inferior

segment of the precentral sulcus and inferior frontal sulcus). Additional methodological details and quantifications of endocranial asymmetries are provided in Poza-Rey et al. (2017). It has to be noted, however, that the human Broca's cap also includes the pars orbitalis (the inferior frontal cortex laying inferior and anteriorly to the horizontal ramus of the Sylvian fissure), which is not included here. In addition, because sulcal boundaries on the inferior frontal region do not reproduce well in hominin endocasts (Balzeau et al., 2014; Falk, 2014), the assessment of Broca's cap asymmetry remains tentative.

Endocranial volumes (ECV) were calculated using the approach described in previous studies, which consisted of scaling complete endocasts (SH4 and SH5) to the size of incomplete endocasts using 15 endocranial landmarks and an affine transformation (Arsuaga et al., 2014; Poza-Rey, 2015). The accuracy of this approach was tested using the very complete cranium 6 (SH6), for which the empirically measured ECV could be compared with the estimated ECV using the previously explained approach. The empirical ECV of SH6 is 1220 cc, whereas its estimated ECV is 1225 cc (Supplementary Information in Arsuaga et al., 2014). The estimated value is, therefore, less than 0.5% higher than the empirical value.

Endocranial volumes were used to calculate encephalization quotients (EQ) based on an average body mass estimate of 69.1 kg for the SH hominins (Arsuaga et al., 2015). SH's EQ was estimated using the recent formula proposed by Grabowski and colleagues ($EQ = ECV / e^{(0.60 * \log(\text{Body Mass}) - 1.40)}$, Grabowski et al., 2016), which provides updated results versus those based on Jerison's (1973) and Martin's (1981) earlier approaches. Those values were compared with data for other species that were calculated using the same approach (Grabowski et al., 2016). Those comparisons use individual ECVs, but, in most cases, species-specific body masses, which can result in misleading values because interindividual variation

in ECV is considered, but the same variation in body mass is not accounted for. Therefore, a range of possible EQs for the SH sample was calculated based on the range of body mass estimates from 59.1 to 79.0 kg (Arsuaga et al., 2015). This approach, however, can still yield unrealistically high variation because it does not account for sexual dimorphism in brain size and body mass. This source of variation was further explored by assuming an equal distribution of sexes in the SH sample, as has been suggested by previous studies (Bermúdez de Castro and Nicolás, 1997; Bermudez de Castro et al., 2004), and assuming that individuals with smaller ECVs are females and that individuals with larger ECVs are males. It should be noted, however, that this way of classifying males and females does not perfectly match previous sex assignments of the individuals in this population based on mandibular size and shape and on dental size (Bermúdez de Castro and Nicolás, 1997; Bermudez de Castro et al., 2004). EQs in the SH individuals were then additionally calculated using female- and male-specific body mass estimates (females: mean 57.6 kg, range 52.9-62.3; males: mean 76.7 kg, range 70.7-82.9). These EQ calculations can potentially be influenced by the immature state of some SH individuals (Arsuaga et al., 1997; Gracia et al., 2009), who may not have attained their adult ECV and body mass. However, the paleodemographic analysis of this population shows that most individuals within the collection are adolescents or young adults who would have attained their adult brain size (Arsuaga et al., 1997; Bermúdez de Castro and Nicolás, 1997; Bermudez de Castro et al., 2004), so an immature state is unlikely to affect most EQ calculations.

Detailed morphometric analyses were based on 17 linear metrics between specific endocranial landmarks measured by E.M.P.-R. (Tables 2 and 3; Fig. 3). Correlations between those variables within the SH sample were computed for those pairs of variables that are present in more than five individuals. A principal components analysis (PCA) of size-adjusted metrics (divided by the geometric mean of all the employed variables) was carried out to assess the

major patterns of shape variation of the SH sample with respect to other hominins. This analysis included only seven variables that were present in the largest possible number of SH specimens, and only those fossil specimens without missing data for those variables. This analysis was based on 29 fossil specimens (which includes eight specimens from the SH sample) and 25 modern humans (Table 1). Comparative data for fossil specimens were compiled from different sources (Grimaud-Hervé, 1997; Falk et al., 2000; Holloway et al., 2004a). Comparative data for the modern human sample were measured by E.M.P.-R.

Results

General anatomical descriptions

A summary of these descriptions is provided in Table 4.

SH2 (Fig. 4) This endocast is rather incomplete. A small part of the posterior frontal region is preserved. The right parietal lobe is almost complete, but the left parietal and occipital lobes are represented only by a few fragments. Due to the fragmentary nature of this cranium, its endocast lacks the anterior region of the temporal lobes and the whole base. Broca's cap is not preserved. The occipital lobes are rounded and slightly projected. The endocranial volume of SH2 was estimated at 1333 cc.

SH3 (Fig. 5) This endocast is also fragmentary and moderately incomplete. A large part of the frontal lobe and the base of the endocast are lost. Occipital and parietal lobes are quite complete, and the left temporal lobe is also preserved. Broca's cap is preserved in both hemispheres, although the right one is not complete. Like SH2, the occipital lobes of SH3 are

rounded and moderately projected. The endocranial volume of this individual was estimated at 1230 cc.

SH4 (Fig. 6) This is one of the most complete endocasts from the SH collection, where all of the endocranial areas are preserved and undistorted. Broca's cap is preserved in both hemispheres, but there is no clear right or left asymmetry. Encephalic rostrum development is intermediate between typology 1 and 2 (Grimaud-Hervé, 1997), which means that the rostrum is pointed and considerably projecting, as observed in most *H. erectus* (type 1) and Neanderthals and Middle Pleistocene fossils (type 2). This endocast shows a left-frontal and left-occipital length petalia, and a right-frontal and right-occipital width petalia. This means that the right hemisphere is wider and shorter than the left. The occipital lobes are rounded and highly projected. The endocranial volume of SH4 was measured to be 1360 cc.

SH5 (Fig. 7) This is also one of the most complete endocasts from the SH collection, which lacks only a fragment of the anterior region of the left parietal lobe. Broca's cap is preserved in both hemispheres and it is larger on the left side. The degree of development of the encephalic rostrum corresponds to typology 1-2 of Grimaud-Hervé (1997). This endocast shows a right-frontal and left-occipital petalial pattern in both length and width. The occipital lobes are rounded and moderately projecting. The endocranial volume of SH5 is 1092 cc.

SH6 (Fig. 8) This endocast is not as complete as SH4 and SH5, but it is better preserved than other endocasts in the collection. Most of the frontal lobes are preserved and the temporal and occipital lobes are quite complete. However, SH6 lacks most of the anterior parietal region of both lobes and the base of the endocast. Broca's cap is preserved in both hemispheres, but there is no clear left or right asymmetry. This endocast shows no petalia in the frontal lobes, and it

shows a left-occipital petalial pattern in both length and width. The occipital lobes are rounded and slightly projecting. The endocranial volume of this individual was estimated at 1225 cc.

SH7 (Fig. 9) This endocast preserves most of the right parietal lobe and a large part of the left one. It lacks a large part of the base involving frontal, temporal and occipital lobes. Broca's cap is preserved on both hemispheres, although it is incomplete on the right one. This endocast shows a left-frontal and left-occipital length petalia, with no frontal and a left-occipital width petalia. The occipital lobes are rounded and slightly projected. The endocranial volume was estimated at 1143 cc.

SH8 (Fig. 10) This is a fragmentary specimen that preserves only the posterior part of the right parietal and temporal lobes. Some middle meningeal vessels can be observed. Based on the preserved endocranial regions, the endocranial volume of this individual was estimated at 1313 cc.

SH9 (Fig. 11) This endocast lacks the base, as well as the cerebellar lobes and the ventral region of the frontal lobe. The temporal lobes are incomplete, but both parietal and occipital lobes are almost complete. Only the left Broca's cap is preserved. This endocast shows a left-frontal and left-occipital width petalia, and a right-frontal and left-occipital length petalia. The occipital lobes are rounded and highly projected. The endocranial volume of this individual was estimated at 1201 cc.

SH10 (Fig. 12) This endocast is rather incomplete. It lacks the entire base, almost all of the occipital lobes, the prefrontal region, and most of the right temporal and parietal lobes. A large part of the frontal lobes, the temporal poles and the left parietal lobe are preserved. Broca's cap

is present in both hemispheres and it appears to be larger on the left side. The endocranial volume of SH10 was estimated at 1218 cc.

SH11 (Fig. 13) This endocast is incomplete. It lacks the base, including the ventral regions of the frontal and temporal lobes. The complete right temporal and parietal lobes are absent. The occipital and cerebellar lobes are quite well preserved, together with the left parietal lobe. Based on the preserved parts, the encephalic rostrum shows a 1-2 typology of Grimaud-Hervé (1997). This endocast shows a right-frontal width petalia and a left-occipital length petalia. The occipital lobes are rounded and slightly projected. The endocranial volume was estimated at 1057 cc.

SH12 (Fig. 14) This endocast preserves only the right parietal lobe, a large part of the left temporal lobe, and the prefrontal area. A large part of Broca's cap is preserved in the right hemisphere. The endocranial volume of this individuals was estimated at 1227 cc.

SH13 (Fig. 15) This endocast is quite complete on its right side, but very incomplete on its left side. It preserves the right frontal, parietal and temporal lobes, whereas the left hemisphere preserves only a minor part of the temporal and inferior frontal region. Broca's cap is preserved on both sides and it is larger on the right hemisphere. Based on the anatomy of the preserved parts, the encephalic rostrum shows a 1-2 typology (Grimaud-Hervé, 1997). The endocranial volume of SH13 was estimated at 1436 cc. This value makes this the biggest endocast in the SH sample, with an endocranial volume that falls comfortably within the range of variation observed for classic Neanderthals (Holloway et al., 2004a).

SH14 (Fig. 16) This endocast is almost complete. It lacks only a part of the base involving the cerebellar area and the left temporal lobe and frontal region. Broca's cap is preserved on both sides and it shows a left asymmetry. The degree of development of the encephalic rostrum is between typology 1 and 2 of Grimaud-Hervé (1997). This endocast shows a right-frontal width petalia and a right-frontal and left-occipital length petalia. The occipital lobes are rounded and slightly projecting. This endocast shows a marked asymmetry that is particularly apparent in posterior view. However, the endocranial traits of this individual cannot be considered representative of the normal range of variation of this population, as SH14 has been previously suggested to suffer from a pathological condition termed lambdoid single suture craniosynostosis (Gracia et al., 2009). According to the previously published diagnosis, this condition would be associated with ecto- and endocranial deformities in this individual (Gracia et al., 2009). The endocranial volume of this individual was estimated at 1224 cc.

SH15 (Fig. 17) This endocast preserves most of the left parietal and temporal lobes, but only a small part of the right parietal and temporal lobes. It lacks portions of the frontal, occipital and cerebellar lobes. Only the left Broca's cap is preserved. The endocranial volume of SH15 was estimated at 1283 cc.

SH16 (Fig. 18) This endocast is rather incomplete. It preserves only a small part of the frontal lobes on both sides, the right temporal lobe and a part of the right parietal lobe; the occipital lobe is almost complete. Broca's cap is preserved in both hemispheres and it shows a right asymmetry. Encephalic rostrum development shows a typology 1-2 (Grimaud-Hervé, 1997). There is no length petalia in the frontal lobes and a very minor right-occipital length petalia (considered no petalia). Comparison of the preserved regions indicates that this endocast shows no frontal width petalia, and it was not possible to assess the occipital width petalia. The

occipital lobes are rounded and slightly projected. The endocranial volume of this individual was estimated at 1236 cc.

SH17 (Fig. 19) The left side of this endocast is very well preserved, but the right side is rather incomplete. The frontal lobe is almost completely preserved. The left temporal and parietal lobes are quite complete, and so are the occipital and cerebellar lobes. However, SH17 lacks the central basicranial region and the right parietal and temporal lobes. Broca's cap is present in both hemispheres and it is larger on the right side. The encephalic rostrum shows a 1-2 typology (Grimaud-Hervé, 1997). This endocast shows a right-frontal petalial pattern for both width and length, but it was not possible to determine the occipital width petalia. Both hemispheres are very similar with almost no length asymmetry. The occipital lobes are rounded and moderately projected. The endocranial volume of this individual was estimated at 1218 cc.

Anatomical comparison with other endocasts

The maximum endocranial width in SH endocasts tends to be located at the level of the temporal lobes, around the mid-posterior region of the second temporal convolution. This pattern is also observed in Ceprano (Bruner and Manzi, 2005), Hexian (Wu et al., 2006), Zhoukoudian (Wu et al., 2009, 2010), the Ngandong collection and Swanscombe. In Neanderthals, the maximum endocranial width tends to be found more posteriorly, at the end of the first temporal convolution and close to the parietal lobe. This same position is observed in some fossil *H. sapiens*, such as those from the Předmostí collection, but other fossil *H. sapiens* (including Brno 3, Cro-Magnon 3, and Skhul 1) show an even more posterior location of the maximum width, at the level of the mid-parietal region.

Frontal organization in the SH collection has been mostly assessed based on the asymmetry of Broca's cap, the endocranial remnant of Broca's area. Broca's cap in SH hominins is well developed on both sides, but it tends to be slightly larger and more prominent on the left side (Poza-Rey, 2015; Poza-Rey et al., 2017). This trait anticipates the pattern seen in modern humans and other later hominins, including Neanderthals, where Broca's cap is generally asymmetrically enlarged on the left side (de Sousa and Cunha, 2012). Indeed, a number of later *Homo* specimens show this asymmetry in the Broca's cap, including *H. erectus* from Zhoukoudian (Wu et al., 2009, 2010) and Hexian (Wu et al., 2006), Middle Pleistocene *Homo* (Kabwe) and Neanderthals (Teshik Tash, Feldhofer and La Quina 5; Holloway et al., 2004a). Some other specimens, including some from the SH collection, show the opposite pattern consisting of an enlarged Broca's cap on the right side (La Ferrassie 1 and La Chapelle-aux-Saints; Holloway et al., 2004a) or an unclear one (Bodo, Arago and Cro-Magnon 3).

Superiorly, the parietal lobes of the SH endocasts are depressed para-sagittally and many of them show a keel in the upper part. This gives rise to a "roof-like" anatomy that is observed in other Middle Pleistocene specimens, such as Swanscombe, Arago, or Reilingen (Fig. 20). This morphology differs from the "tent-like" ("en tente") configuration observed in *H. erectus* specimens, such as those from Hexian (Wu et al., 2006) and Zhoukoudian (Wu et al., 2010). Juvenile SH specimens show a less marked para-sagittal depression and, therefore, a more rounded appearance. Rounded parietals (termed "en bombe") can be observed in La Ferrassie 1, La Quina 5, Teshik Tash and other Neanderthals such as Saccopastore 1 (Bruner and Manzi, 2008). Fossil *H. sapiens*, such as Cro-Magnon 3, Brno 3, the Předmostí collection, Skhul 1, Dolní Věstonice 1, Combe Capelle and Singa, show rounded parietal lobes and vertical temporal lobes, which result in a "house-like" ("en maison") anatomy.

A qualitative assessment shows that the temporal lobes of the SH endocasts are narrower than those of modern humans, but similar to Neanderthals' (including Teshik Tash, Gibraltar 1, Monte Circeo, La Chapelle-aux-Saints and La Ferrassie 1). The temporal lobes of Zhoukoudian hominins have been described as narrow and slender (Wu et al., 2010), whereas those from Hexian, Sambungmacan and Trinil are fuller and broader.

Most of the SH specimens show slight or moderate posterior projection of the occipital lobes (except SH4 and SH9, which have a stronger posterior projection than the others). This is quite different from the pattern observed in Zhoukoudian and Hexian specimens, which are dorso-ventrally flattened and have strong posterior projections (Wu et al., 2006, 2010). Some Middle Pleistocene specimens, such as Petralona and Kabwe, have been described as having a strong projection of the occipital lobes (Seidler et al., 1997; Holloway et al., 2004a), but we consider this a moderate projection that is similar to that observed in the SH endocasts. Neanderthal endocasts show a strong posterior protrusion that can be related to the occipital "chignon" (Holloway et al., 2004a), which is not observed in SH hominins (Arsuaga et al., 2014). Fossil modern humans (including those from Jebel Irhoud, Dolní Věstonice, Předmostí, Brno, Combe Capelle and Cro-Magnon) and recent modern humans show rounded occipital lobes and very slight posterior projections.

The petalial patterns of the SH endocasts are relatively variable when considering both width and length projections. A left occipital length petalia is clearly predominant in the SH hominins (seven out of nine individuals where this petalia can be assessed), but this is not always associated with a right frontal petalia (see Table 4). This left-occipital right-frontal (LORF) petalia is the predominant condition in modern humans (Balzeau et al., 2012), and it seems to become more and more frequent during the evolution of the genus *Homo* (de Sousa and Cunha,

2012). Consistent with this observation, numerous hominin endocasts show a LORF petalia (Saccopastore 1, Reilingen, Sambungmacan 3, Ngandong 1, 6 and 7, Trinil 2, Amud 1, Teshik Tash, Brno 3, Combe Capelle, Cro-Magnon 3, Feldhofer, Gibraltar 1, La Quina 5, Monte Circeo, Předmostí 4 and 10, Spy 1 and 2, La Ferrassie 1, Salé, Jebel Irhoud 1). Others, however, show less common patterns, including a right-occipital right-frontal petalia in Hexian (Wu et al., 2006) and a right-occipital left-frontal petalia in Brno 2, Le Moustier 1 and Předmostí 3 (Holloway et al., 2004a). Substantial variability is also observed in the Zhoukoudian collection (Wu et al., 2010).

Endocranial volume and encephalization

Endocranial volume in the Sima de los Huesos sample ranges from 1057 to 1436 cc (Fig. 21, Table 5). This range is slightly lower than that observed for classic Neanderthals (1172-1740 cc) and similar to that observed in non-SH Middle Pleistocene *Homo* (1165-1325 cc), although the latter can be affected by the lack of a clear definition for this group (see below). In addition, the lowest ECV observed in the SH sample corresponds to one of the subadult individuals whose adult ECV may be slightly higher (SH11), so the lowest ECV corresponding to an SH adult individual is 1092 cc.

The mean encephalization quotient of the SH sample, obtained from the average ECV and estimated mean body mass for this population, is 6.29. This EQ is slightly higher than the Middle Pleistocene *Homo* value (6.14) and smaller than that observed in classic Neanderthals (6.86), obtained from data in Grabowski et al. (2016). The range of variation of EQs observed in SH hominins is similar to that found in Middle Pleistocene *Homo* (referred to as *H. heidelbergensis* in Grabowski et al., 2016) and lower than in Neanderthals (Fig. 21). EQs

within the SH sample range between 5.36 and 7.28, with a potential range of variation of 5.11-8.01 when variation in body mass is also considered (Table 5). This range of variation, however, is substantially reduced when potential dimorphism in brain size and body mass is accounted for, giving rise to largely overlapping ranges of variation for putative females and males (6.02-6.99 for females and of 5.85-6.83 for males).

Endocranial morphometrics

The analysis of the endocranial metrics of SH hominins (Table 6) reveals some expected patterns. Most of the correlations across variables are not significant (Table 7). Although this lack of significant correlations can result from the relative independence of these variables within the SH sample, it is also likely related to the small sample sizes, which are the result of the fragmentary nature of most SH endocasts. Some variables, however, still show significant correlations due to their general relationship with endocranial volume (e.g., endocranial length and maximum width) or with general endocranial proportions (e.g., maximum width, frontal width, and width at Broca's cap).

The principal components analysis of size-adjusted linear metrics shows a clear separation between modern humans and all the other groups, including the SH hominins (Fig. 22). Unsurprisingly, variable loadings show that the variable driving this separation is the increase in the parietal chord in modern humans (Fig. 22, Table 8). Simple parietal linear metrics have been shown to be a reliable predictor of parietal expansion in modern humans and to separate them from earlier hominins (Bruner et al., 2011; Gómez-Robles et al., 2017), so this appears to be the anatomical trait underlying this distribution of individuals. The SH sample fully overlaps with the Neanderthal sample, thus pointing to negligible differences between early

and classic Neanderthals in these endocranial metrics. *H. erectus* and Middle Pleistocene *Homo* (MP *Homo*) also overlap with classic Neanderthals and SH hominins to a large extent. Some *H. erectus*, however, have negative scores for PC1 overlapping with modern humans. This indicates that these *H. erectus* specimens have a relatively long parietal chord, but their low PC1 scores do not indicate the high degree of parietal expansion observed in modern humans. The ranges of variation of all species overlap along PC2, which is mostly associated with the relative length of the complete endocast and of the frontal chord. Within the modern human sample, clear differences exist between the fossil and the recent samples, with a much higher degree of parietal expansion in the latter (Neubauer et al., 2018). The simple linear metrics used in our study, however, do not clearly separate different groups of fossil modern humans according to their geological age and level of parietal expansion.

An additional PCA was carried out only on the fossil sample to get rid of the strong effect that the upper parietal expansion observed in recent modern humans has on the ordination of individuals. That PCA shows extensive overlap of species along PC1. The variables with the highest —and opposite— loadings on PC1 are the frontal chord and the parietal chord, which means that the ordination of individuals along PC1 is driven by the relative size of the frontal and the parietal lobes (Fig. 22, Table 8). *H. erectus*, MP *Homo* and fossil *H. sapiens* overlap to a large extent along PC2. There is, however, certain separation between SH hominins and classic Neanderthals along PC2, which is associated in its positive values with maximum and occipital width (that is, with the width of the caudal half of the brain) and in its negative values with the frontal length. Therefore, SH hominins tend to have broader brains than classic Neanderthals, particularly in the temporo-parietal and occipital regions, whereas Neanderthals tend to have longer brains, which is particularly true for the frontal lobe. The Ehringsdorf

endocast, also considered an early Neanderthal (although substantially younger than SH hominins), plots with classic Neanderthals and is separated from SH individuals in this PCA.

Discussion

The analysis of the SH endocranial collection provides some fundamental insights into the evolution of the Neanderthal brain configuration. A number of anatomical (Arsuaga et al., 1997; Martínón-Torres et al., 2012; Arsuaga et al., 2014, 2015; Gómez-Robles et al., 2015a) and recent molecular studies (Meyer et al., 2014, 2016) demonstrate that the SH hominins are early representatives of the Neanderthal lineage. Therefore, their endocranial anatomy provides information on the brain phenotype that was typical of the early stages of the Neanderthalization process. Brain size and, to a lesser extent, the level of encephalization are similar in classic Neanderthals and in fossil modern humans (Holloway et al., 2004a; Grabowski et al., 2016). Brain organization as inferred from endocranial morphology also shows similarities between classic Neanderthals and modern humans in some aspects of frontal and occipital anatomy (reviewed in de Sousa and Cunha, 2012), but also important differences that are mostly related to the expansion of the parietal cortex and to the level of globularization (Bruner et al., 2003, 2011; Gunz et al., 2010; Neubauer and Gunz, 2018). The analysis of the SH collection shows that some of those similarities between classic Neanderthals and modern humans were already present in early Neanderthals, thus suggesting a common origin of those traits. Other traits that are shared by modern humans and classic Neanderthals and that are not observed in the SH sample point to a parallel acquisition of those traits in both lineages. One of the traits that seems to have evolved in parallel in both species is a very large brain size, so it is possible that other parallel-evolving traits are allometrically associated with brain size increase. The evolution of brain size and brain organization in these groups is discussed below.

Brain size and encephalization

Brain size in SH hominins has a mean value of 1241 cc, which is close to the MP *Homo* mean value (1227 cc) and lower than the Neanderthal mean value (1387 cc). The range of variation of brain size in SH hominins largely overlaps with that seen in MP *Homo*, but it falls in the lower end of the range of variation seen in Neanderthals (Fig. 21). This indicates a gradual increase in brain size from earlier to later Neanderthals. Other individuals that are considered early Neanderthals show similarly variable endocranial volumes, with some specimens showing SH-like values (1200 cc in Steinheim and 1325 cc in Swanscombe) and others showing more Neanderthal-like values (1430 cc in Reilingen and 1450 cc in Ehringsdorf). Interestingly, brain size in the earliest representatives of the modern human lineage already shows values that are typical of later modern humans (Neubauer et al., 2018). Indeed, cranial capacities in the Jebel Irhoud specimens, now considered to be the earliest known hominins that can be clearly identified as part of the modern human lineage (Hublin et al., 2017), are 1375 and 1467 cc (Neubauer et al., 2018), very close to the mean brain size of fossil modern humans (approx. 1485 cc). The Jebel Irhoud remains are younger than those from Sima de los Huesos (315 ka versus 430 ka; Arnold et al., 2014; Richter et al., 2017), so it is possible that pre-modern humans predating those from Jebel Irhoud had smaller brains that increased in size later in the evolution of the modern human lineage. However, considering the current hypodigms of both species and their ancestral groups, a gradual increase in brain size is more clearly observed in the Neanderthal lineage than in the modern human lineage.

As for encephalization, the SH sample shows a transitional state between that seen in more primitive groups (*H. erectus* and MP *Homo*) and in classic Neanderthals. The EQ of the SH

sample is 6.3, which is closer to the values observed in MP *Homo* (6.1) than in Neanderthals (6.9). All these values are substantially lower than the ones observed in modern humans (7.7). The comparison of ranges of variation across species and of individual values is hampered by the scarcity of data regarding individual body masses. Therefore, EQs have been estimated with respect to species-specific (in some cases, group-specific) body masses (Ruff et al., 1997; Grabowski et al., 2016). This scenario is far from ideal because it accounts for interindividual differences in brain size, but not in body mass, thus making large-brained specimens look highly encephalized. However, this can be simply the result of their large body size. Studies of contemporary humans show that brain size (as measured from MRI brain scans) and body mass have a significant but moderate correlation of about 0.4 (A.G.-R., unpublished data). Although this correlation is not extremely high, it does show a moderate association between both variables within species whereby larger individuals tend to have larger brains. This implies that the actual range of variation in EQs within each species would be probably narrower were interindividual variation in body mass included in these calculations. Despite this methodological challenge, the evaluation of species- or group-specific values indicates that the SH hominins show both an absolute brain size and a level of encephalization that are intermediate between those seen in earlier and more primitive hominins (*H. erectus* and non-SH MP *Homo*) and later classic Neanderthals.

Brain organization

Brain organization has been evaluated through quantitative analysis and qualitative descriptions. Quantitative analysis shows a clear separation between recent modern humans and all other hominin groups that is driven by the length of the parietal chord. All other hominin groups, however, including *H. erectus*, MP *Homo*, SH hominins, Neanderthals and fossil

modern humans, are virtually indistinguishable on the basis of their level of parietal expansion. This difference between recent modern humans and other later hominins has been already noted by other authors (Bruner et al., 2003; Neubauer et al., 2018). Unsurprisingly, the SH endocranial collection aligns with other non-recent modern humans in showing no parietal expansion. Whether parietal expansion is linked to functional differences related to modern human behavior (Gunz et al., 2010; Bruner and Iriki, 2016) or simply the result of general craniofacial integration (Martínez-Abadías et al., 2012; Zollikofer et al., 2017) is a matter of some debate.

The second principal component analysis, which includes only fossil specimens, shows some intriguing differences between SH hominins and later Neanderthals. SH hominins seem to have broader brains, particularly in their posterior (temporo-parietal and occipital) regions. Later Neanderthals, however, have relatively longer brains, particularly in the anterior (frontal) region. The fact that other hominin groups do not show any trend with respect to the posterior endocranial width (measured in the temporo-parietal and occipital regions) and frontal length makes it unlikely that they are related to functional/behavioral differences between early and late Neanderthals. However, certain behavioral differences may have existed and they may be related to the already mentioned differences in brain size and encephalization or on other aspects not included in our quantitative analyses (see below).

The qualitative comparison of SH and Neanderthal endocranial anatomy shows some additional differences. The encephalic rostrum of SH hominins tends to be pointed and considerably projecting, showing a typology that is intermediate between that seen in *H. erectus* (long and narrow encephalic rostrum with a marked invagination at the interorbital region) and Neanderthals (shorter and broader rostrum as a result of the expansion of the

anterior frontal convolutions; Grimaud-Hervé, 1997). This intermediate typology of the encephalic rostrum is indicative of a moderate expansion of the prefrontal region. This expansion, particularly the one involving the inferior frontal region, has occurred asymmetrically during hominin evolution, giving rise to an asymmetry of Broca's language area in modern humans whereby the left region is larger than its right counterpart. This size-asymmetry can be associated with microstructural asymmetries of the inferior frontal region (Amunts et al., 1999; Keller et al., 2009), some of which are also present in chimpanzees (Schenker et al., 2010). Structural asymmetries of the inferior frontal region that can be identified on endocasts are also found in Neanderthals and other later hominins (de Sousa and Cunha, 2012). Because of the large proportion of modern humans that exhibit a left hemisphere language dominance (Springer et al., 1999), it is thought that the anatomical asymmetry of Broca's area can be related to functional lateralization for language production (but see Keller et al., 2009). The small number of SH individuals where inferior frontal asymmetry can be assessed makes it hard to infer a population-specific pattern. Out of eight SH endocasts that preserve Broca's cap on both sides, three of them have a larger left area, three of them have a larger right area, and two individuals do not show a clear pattern. This observation may indicate a transitional state with respect to that observed in later hominins, or it may simply indicate that evolutionary trends in Broca's cap asymmetry are not as clear-cut as previously assumed (Balzeau et al., 2014). In addition, the analysis of evolutionary trends in endocranial and brain asymmetry is unavoidably affected by the different methodological approaches used in different studies, which can lead to apparently contradictory results. For example, asymmetries in surface areas assessed in hominin endocasts may not accurately reflect the evolutionary trends indicated by volumetric and microstructural asymmetries measured in MRI scans and postmortem brain tissue of extant primates.

Whole-hemispheric asymmetries, usually referred to as petalias, also show a transitional state in SH hominins. These hominins tend to show a left-occipital petalia that is typical of modern humans and other later hominins. In modern humans, the left-occipital petalia tends to be associated with a right-frontal petalia. This pattern, however, is not so clearly observed in the SH hominins. The left-occipital right-frontal (LORF) petalia has been classically considered to correlate with right-handedness in humans (LeMay and Kido, 1978), but later analyses have not confirmed that association (Good et al., 2001). Indeed, even if this petalial pattern is not as consistent in SH hominins as it is in modern humans, studies of the orientation of labial striations on the surface of the incisors, which are formed when individuals hold something between their anterior teeth and use a stone tool to cut it, indicate that SH hominins were predominantly right-handed (Lozano et al., 2009) in a proportion that is similar to that observed in modern humans (9:1). Likewise, studies of dental striations in Neanderthal populations show that classic Neanderthals were also preferentially right-handed in a similar proportion (Fruyer et al., 2010; Estalrich and Rosas, 2013). The presence of a population-level right-handedness in SH hominins is not surprising considering that a preferential right-handedness has been suggested even for early *Homo* based on the morphology of stone tools (Toth, 1985) and on the orientation of labial striations on the anterior teeth (Fruyer et al., 2016). Indeed, a preferential right-handedness, at least for some tasks, has even been suggested for chimpanzees (Hopkins et al., 2007), which indicates that this aspect of behavioral lateralization may predate the separation of the hominin and panin clades and may have been present in the earliest hominins.

As in other MP *Homo*, the SH hominins tend to show para-sagittally depressed parietal lobes, with the maximum width located quite inferior and posteriorly. This configuration is different from that observed in *H. erectus* (which tends to show a tent-like profile in posterior view),

623 classic Neanderthals (with a rounded or “en bombe” profile), and modern humans (with a
624 house-like or “en maison” profile). Perhaps surprisingly, the SH endocasts do not show any
625 clear differentiation towards the Neanderthal condition (Bruner et al., 2003), but a pure
626 intermediate state between the most primitive tent-like anatomy and the two derived states (“en
627 bombe” and “en maison”). This intermediate condition of the SH sample makes it difficult to
628 identify even incipient Neanderthal traits in the endocasts and neurocrania of early
629 Neanderthals (Arsuaga et al., 1997). However, the affinity of SH hominins with classic
630 Neanderthals indicates that the rounded, laterally expanded parietal lobes typical of classic
631 Neanderthals must have evolved from the flatter parietal lobes that are typical of SH hominins
632 and other early Neanderthals. These shape changes result not only from the level of upper
633 parietal expansion, but also from the level of inferior parietal expansion and from the
634 relationships across different parietal domains and with temporal areas. Although upper
635 parietal expansion in fossil hominins has been extensively discussed in the literature (Bruner
636 et al., 2003, 2011; Gunz et al., 2010; Ponce de León et al., 2016; Neubauer et al., 2018),
637 comparatively little has been discussed about inferior parietal variation. Quantitative studies of
638 chimpanzee and human brains, however, have shown that the inferior parietal region is a hot
639 spot of variation and evolutionary change in humans in terms of symmetric and asymmetric
640 variation (Gómez-Robles et al., 2013b), and a particularly plastic target for environmental
641 influences (Gómez-Robles et al., 2015b). These observations are in line with the range of
642 complex cognitive tasks in which the inferior parietal region is involved, including language,
643 attention and memory (Sherwood et al., 2008). The clear differences in parietal configuration
644 between SH hominins, classic Neanderthals and modern humans, which seem to involve both
645 superior and inferior parietal regions, might point to behavioral differences in higher-order
646 tasks that are controlled by these domains. As far as it can be inferred from endocranial
647 anatomy, however, the temporal lobes of SH hominins are similar to those of Neanderthals but

narrower than those of modern humans, confirming the results of previous quantitative analyses (Bastir et al., 2011). These differences in temporal lobe development have been suggested to correlate with differences in visual memory, language and theory of mind that might be mediated by higher olfactory functions (Bastir et al., 2011).

The occipital lobes of the SH hominins tend to be slightly projected, but substantially less so than the occipital lobes of classic Neanderthals. The size and shape of the occipital lobes can simply reflect other ectocranial traits, such as the presence of an occipital bun in Neanderthals, but they can be also associated with the size of the visual cortex, which is located in the occipital region. The size of the visual cortex (as inferred from orbital size rather than from endocranial anatomy) has been controversially linked to differences in social cognition and social organization between Neanderthals and modern humans (Pearce et al., 2013; but see Traynor et al., 2015, among others). Given the methodological difficulties associated with these inferences, however, interpretations about sociality in the SH sample are more safely based on demographic aspects of this group. The SH collection is formed by at least 28 individuals (Arsuaga et al., 2014) where infants are underrepresented (Bermúdez de Castro and Nicolás, 1997). Given the excellent preservation of all the skeletal parts in the SH sample, which includes even auditory ossicles, it is unlikely that the absence of infants in the sample is the result of taphonomic factors. This underrepresentation of infants and children is more likely to have resulted from differential accumulation related to social or cultural factors (Bermúdez de Castro and Nicolás, 1997) or from ecological factors (Bocquet-Appel and Arsuaga, 1999). Among them, a demographic crisis caused by severe environmental fluctuation has been suggested to explain the age distribution of the SH sample and of other fossil hominin samples, such as that from Krapina (Bocquet-Appel and Arsuaga, 1999). In any case, the absence of very young individuals in the SH population indicates that the potential size of the social group

to which these individuals belonged was probably larger (perhaps substantially larger) than the number of individuals in the SH assemblage. In addition, the study of the nearby site of Gran Dolina TD10-2, which is roughly contemporary to Sima de los Huesos and probably associated with the same or similar pre-Neanderthal populations, shows evidence of communal hunting activities that involve complex technological, cognitive and social skills (Rodríguez-Hidalgo et al., 2017).

SH behavior and evolutionary relationships

Although SH hominins have a more primitive endocranial morphology than that of classic Neanderthals, this may not be informative of their behavioral capabilities. Indeed, most behavioral inferences about the SH collection come from the study of the associated paleontological and archaeological evidence rather than from the study of endocranial anatomy per se. Based on that associated evidence, the SH hominins have been suggested to show most of the behavioral innovations that were also present in later Neanderthals, including a right-hand preference and other signs of behavioral lateralization (Lozano et al., 2009; Poza-Rey et al., 2017), conspecific care (Gracia et al., 2009), lethal interpersonal conflict (Sala et al., 2015), the capacity to hunt cooperatively (Rodríguez-Hidalgo et al., 2017), symbolic behavior (Carbonell and Mosquera, 2006) and perhaps a spoken language (Martínez et al., 2004). Some of these behavioral traits, however, particularly those related to behavioral lateralization and lethal aggression, may largely predate the origin of the Neanderthal lineage and may have been inherited from much earlier hominins or even from the chimpanzee-human last common ancestor (Hopkins et al., 2007; Gómez et al., 2016). These observations indicate that the association between particular endocranial evolutionary changes and the evolution of human behavioral innovation can be rather loose. Behavioral inferences should therefore be based on

a comprehensive and holistic study of all the available evidence from the fossil and archaeological record.

The analysis of endocranial variation in the SH sample, however, is important because it allows us to make inferences about the timing of the Neanderthalization process. The SH endocranial collection shows, in general, a transitional state between that found in more primitive hominins (*H. erectus* and non-SH Middle Pleistocene *Homo*) and the derived states found in classic Neanderthals and modern humans, without any differentiation towards the Neanderthal condition. Indeed, SH endocranial anatomy differs markedly from the endocranial anatomy of classic Neanderthals in some traits, such as the absence of a strong occipital projection and the relationship between upper parietal, lower parietal and temporal domains. These distinct endocranial configurations give rise to very different profiles in posterior view that have been extensively discussed at the level of external cranial anatomy (e.g., Hublin, 1982; Arsuaga et al., 1997) and that are also reproduced endocranially. In most traits included in our study, SH hominins also differ from the conditions observed in earlier and later hominins, showing a squarely intermediate state between those found in *H. erectus*, Neanderthals and modern humans. Future quantitative studies should assess whether finer-grained brain organization still shows a primitive condition or whether it shows some incipient affinities with Neanderthals.

SH primitive endocranial anatomy can be compared with a large number of features where SH hominins show an intermediate morphology between that observed in earlier hominins and that observed in classic Neanderthals (e.g., Arsuaga et al., 1993, 1997; Carretero et al., 1997; Martínez and Arsuaga, 1997; Rosas, 1997; Gómez-Olivencia et al., 2007; Bonmatí et al., 2010; Arsuaga et al., 2014, 2015; Quam et al., 2016; Rodríguez et al., 2016; Pablos et al., 2017). Those traits are differentially developed along the primitive-to-classic Neanderthal continuum,

with some of them showing a fully Neanderthal or quasi-Neanderthal condition, whereas others show a more primitive configuration. Indeed, SH primitive endocranial anatomy stands out in stark contrast with SH dental anatomy, which shows a surprisingly derived Neanderthal condition that is not found in any other traits of this population (Bermúdez de Castro and Nicolas, 1995; Martínón-Torres et al., 2012; Gómez-Robles et al., 2015a). The different evolutionary trends observed for different traits indicate that classic Neanderthal anatomy evolved in a mosaic fashion whereby masticatory traits (dental, mandibular and facial anatomy) acquired a fully Neanderthal condition substantially earlier than non-masticatory traits (Arsuaga et al., 2014). Different selective pressures related to mastication - and perhaps also to the para-masticatory use of anterior teeth (Lozano et al., 2008) - may have driven the early evolution of the Neanderthal dentition and face (Arsuaga et al., 2014). Therefore, neurocranial anatomy and the anatomy of other non-masticatory regions with very diverse functional roles seem to have lagged behind masticatory features, which apparently led the Neanderthalization process as it can be inferred from SH hominins.

References

- Amunts, K., Schleicher, A., Burgel, U., Mohlberg, H., Uylings, H.B., Zilles, K., 1999. Broca's region revisited: Cytoarchitecture and intersubject variability. *Journal of Comparative Neurology* 412, 319–341.
- Aranburu, A., Arsuaga, J.L., Sala, N., 2017. The stratigraphy of the Sima de los Huesos (Atapuerca, Spain) and implications for the origin of the fossil hominin accumulation. *Quaternary International* 433, 5–21.
- Arnold, L.J., Demuro, M., Parés, J.M., Arsuaga, J.L., Aranburu, A., Bermúdez de Castro, J.M., Carbonell, E., 2014. Luminescence dating and palaeomagnetic age constraint on

748 hominins from Sima de los Huesos, Atapuerca, Spain. *Journal of Human Evolution* 67,
749 85–107.

750 Arsuaga, J.-L., Martínez, I., Gracia, A., Carretero, J.-M., Carbonell, E., 1993. Three new human
751 skulls from the Sima de los Huesos Middle Pleistocene site in Sierra de Atapuerca,
752 Spain. *Nature* 362, 534–537.

753 Arsuaga, J.L., Martínez, I., Gracia, A., Lorenzo, C., 1997. The Sima de los Huesos crania
754 (Sierra de Atapuerca, Spain). A comparative study. *Journal of Human Evolution* 33,
755 219–281.

756 Arsuaga, J.L., Martínez, I., Arnold, L.J., Aranburu, A., Gracia-Téllez, A., Sharp, W.D., Quam,
757 R.M., Falguères, C., Pantoja-Pérez, A., Bischoff, J., Poza-Rey, E., Parés, J.M.,
758 Carretero, J.M., Demuro, M., Lorenzo, C., Sala, N., Martínón-Torres, M., García, N.,
759 Alcázar de Velasco, A., Cuenca-Bescós, G., Gómez-Olivencia, A., Moreno, D., Pablos,
760 A., Shen, C.-C., Rodríguez, L., Ortega, A.I., García, R., Bonmatí, A., Bermúdez de
761 Castro, J.M., Carbonell, E., 2014. Neandertal roots: Cranial and chronological evidence
762 from Sima de los Huesos. *Science* 344, 1358–1363.

763 Arsuaga, J.L., Carretero, J.-M., Lorenzo, C., Gómez-Olivencia, A., Pablos, A., Rodríguez, L.,
764 García-González, R., Bonmatí, A., Quam, R.M., Pantoja-Pérez, A., Martínez, I.,
765 Aranburu, A., Gracia-Téllez, A., Poza-Rey, E., Sala, N., García, N., Alcázar de
766 Velasco, A., Cuenca-Bescós, G., Bermúdez de Castro, J.M., Carbonell, E., 2015.
767 Postcranial morphology of the middle Pleistocene humans from Sima de los Huesos,
768 Spain. *Proceedings of the National Academy of Sciences USA* 112, 11524–11529.

769 Balzeau, A., Grimaud-Hervé, D., Jacob, T., 2005. Internal cranial features of the Mojokerto
770 child fossil (East Java, Indonesia). *Journal of Human Evolution* 48, 535–553.

771 Balzeau, A., Gilissen, E., Grimaud-Hervé, D., 2012. Shared pattern of endocranial shape
 772 asymmetries among great apes, anatomically modern humans, and fossil hominins.
 773 PLoS ONE 7, e29581.

774 Balzeau, A., Gilissen, E., Holloway, R.L., Prima, S., Grimaud-Hervé, D., 2014. Variations in
 775 size, shape and asymmetries of the third frontal convolution in hominids:
 776 Paleoneurological implications for hominin evolution and the origin of language.
 777 Journal of Human Evolution 76, 116–128.

778 Bastir, M., Rosas, A., Gunz, P., Peña-Melian, A., Manzi, G., Harvati, K., Kruszynski, R.,
 779 Stringer, C., Hublin, J.-J., 2011. Evolution of the base of the brain in highly
 780 encephalized human species. Nature Communications 2, 588.

781 Beaudet, A., Bruner, E., 2017. A frontal lobe surface analysis in three archaic African human
 782 fossils: OH 9, Buia, and Bodo. Comptes Rendus Palevol 16, 499–507.

783 Beaudet, A., Dumoncel, J., Beer, F., Durrleman, S., Gilissen, E., Oettlé, A., Subsol, G.,
 784 Thackeray, J.F., Braga, J., 2018. The endocranial shape of *Australopithecus africanus*:
 785 Surface analysis of the endocasts of Sts 5 and Sts 60. Journal of Anatomy 232, 296–
 786 303.

787 Bermúdez de Castro, J.M., Nicolás, M.E., 1995. Posterior dental size reduction in hominids:
 788 The Atapuerca evidence. American Journal of Physical Anthropology 96, 335–356.

789 Bermúdez de Castro, J.M., Nicolás, M.E., 1997. Palaeodemography of the Atapuerca-SH
 790 Middle Pleistocene hominid sample. Journal of Human Evolution 33, 333–355.

791 Bermúdez de Castro, J.M., Martínón-Torres, M., Lozano, M., Sarmiento, S., Muela, A., 2004.
 792 Paleodemography of the Atapuerca-Sima de los Huesos hominin sample: A revision
 793 and new approaches to the paleodemography of the European Middle Pleistocene
 794 population. Journal of Anthropological Research 60, 5–26.

795 Bischoff, J.L., Shamp, D.D., Aramburu, A., Arsuaga, J.L., Carbonell, E., Bermudez de Castro,
796 J.M., 2003. The Sima de los Huesos hominids date to beyond U/Th equilibrium (>350
797 kyr) and perhaps to 400–500 kyr: New radiometric dates. *Journal of Archaeological*
798 *Science* 30, 275–280.

799 Bischoff, J.L., Williams, R.W., Rosenbauer, R.J., Aramburu, A., Arsuaga, J.L., García, N.,
800 Cuenca-Bescós, G., 2007. High-resolution U-series dates from the Sima de los Huesos
801 hominids yields 600–66+ ∞ kyrs: Implications for the evolution of the early
802 Neanderthal lineage. *Journal of Archaeological Science* 34, 763–770.

803 Bocquet-Appel, J.P., Arsuaga, J.L., 1999. Age distributions of hominid samples at Atapuerca
804 (SH) and Krapina could indicate accumulation by catastrophe. *Journal of*
805 *Archaeological Science* 26, 327–338.

806 Bonmatí, A., Gómez-Olivencia, A., Arsuaga, J.-L., Carretero, J.M., Gracia, A., Martínez, I.,
807 Lorenzo, C., Bermudez de Castro, J.M., Carbonell, E., 2010. Middle Pleistocene lower
808 back and pelvis from an aged human individual from the Sima de los Huesos site, Spain.
809 *Proceedings of the National Academy of Sciences USA* 107, 18386.

810 Bookstein, F., Schäfer, K., Prossinger, H., Seidler, H., Fieder, M., Stringer, C., Weber, G.W.,
811 Arsuaga, J.L., Slice, D.E., Rohlf, F.J., Recheis, W., Mariam, A.J., Marcus, L.F., 2000.
812 Comparing frontal cranial profiles in archaic and modern *Homo* by morphometric
813 analysis. *Anatomical Record* 257, 217–224.

814 Bruner, E., Manzi, G., 2005. CT-based description and phyletic evaluation of the archaic
815 human calvarium from Ceprano, Italy. *Anatomical Record* 285A, 643–657.

816 Bruner, E., Manzi, G., 2008. Paleoneurology of an “early” Neandertal: Endocranial size, shape,
817 and features of Saccopastore 1. *Journal of Human Evolution* 54, 729–742.

818 Bruner, E., Iriki, A., 2016. Extending mind, visuospatial integration, and the evolution of the
819 parietal lobes in the human genus. *Quaternary International* 405, 98–110.

820 Bruner, E., Manzi, G., Arsuaga, J.L., 2003. Encephalization and allometric trajectories in the
821 genus *Homo*: evidence from the Neandertal and modern lineages. Proceedings of the
822 National Academy of Sciences USA 100, 15335–15340.

823 Bruner, E., de la Cuetara, J.M., Holloway, R., 2011. A bivariate approach to the variation of
824 the parietal curvature in the genus *Homo*. Anatomical Record 294, 1548–1556.

825 Bruner, E., Grimaud-Hervé, D., Wu, X., de la Cuétara, J.M., Holloway, R., 2015. A
826 paleoneurological survey of *Homo erectus* endocranial metrics. Quaternary
827 International 368, 80–87.

828 Carbonell, E., Mosquera, M., 2006. The emergence of a symbolic behaviour: The sepulchral
829 pit of Sima de los Huesos, Sierra de Atapuerca, Burgos, Spain. Comptes Rendus
830 Palevol 5, 155–160.

831 Carlson, K.J., Stout, D., Jashashvili, T., de Ruiter, D.J., Tafforeau, P., Carlson, K., Berger,
832 L.R., 2011. The endocast of MH1, *Australopithecus sediba*. Science 333, 1402–1407.

833 Carretero, J.M., Arsuaga, J.L., Lorenzo, C., 1997. Clavicles, scapulae and humeri from the
834 Sima de los Huesos site (Sierra de Atapuerca, Spain). Journal of Human Evolution 33,
835 357–408.

836 Coqueugniot, H., Hublin, J.-J., Veillon, F., Houët, F., Jacob, T., 2004. Early brain growth in
837 *Homo erectus* and implications for cognitive ability. Nature 431, 299–302.

838 de Sousa, A., Cunha, E., 2012. Hominins and the emergence of the modern human brain. In:
839 Hofman, M.A, Falk, D. (Eds.), Evolution of the Primate Brain. Elsevier, Amsterdam,
840 pp. 293–322.

841 Estalrich, A., Rosas, A., 2013. Handedness in Neandertals from the El Sidrón (Asturias,
842 Spain): Evidence from instrumental striations with ontogenetic inferences. PLOS ONE
843 8, e62797.

844 Falk, D., 1980. A reanalysis of the South African australopithecine natural endocasts. American
845 Journal of Physical Anthropology 53, 525–539.

846 Falk, D., 1983. The Taung endocast: A reply to Holloway. American Journal of Physical
847 Anthropology 60, 479–489.

848 Falk, D., 2014. Interpreting sulci on hominin endocasts: Old hypotheses and new findings.
849 Frontiers in Human Neuroscience 8, 134.

850 Falk, D., Redmond Jr, J.C., Guyer, J., Conroy, C., Recheis, W., Weber, G.W., Seidler, H., 2000.
851 Early hominid brain evolution: A new look at old endocasts. Journal of Human
852 Evolution 38, 695–717.

853 Falk, D., Hildebolt, C., Smith, K., Morwood, M.J., Sutikna, T., Brown, P., Jatmiko, Saptomo,
854 E.W., Brunnsden, B., Prior, F., 2005. The brain of LB1, *Homo floresiensis*. Science 308,
855 242–245.

856 Falk, D., Zollikofer, C.P., de León, M.P., Semendeferi, K., Warren, J.L.A., Hopkins, W.D.,
857 2018. Identification of in vivo sulci on the external surface of eight adult chimpanzee
858 brains: Implications for interpreting early hominin endocasts. Brain, Behavior and
859 Evolution 91, 45–58.

860 Frayer, D.W., Fiore, I., Lalueza-Fox, C., Radovcic, J., Bondioli, L., 2010. Right handed
861 Neandertals: Vindija and beyond. Journal of Anthropological Sciences 88, 113–127.

862 Frayer, D.W., Clarke, R.J., Fiore, I., Blumenschine, R.J., Pérez-Pérez, A., Martinez, L.M.,
863 Estebananz, F., Holloway, R., Bondioli, L., 2016. OH-65: The earliest evidence for
864 right-handedness in the fossil record. Journal of Human Evolution 100, 65–72.

865 Gómez, J.M., Verdú, M., González-Megías, A., Méndez, M., 2016. The phylogenetic roots of
866 human lethal violence. Nature 538, 233.

867 Gómez-Olivencia, A., Carretero, J.M., Arsuaga, J.L., Rodríguez-García, L., García-González,
868 R., Martínez, I., 2007. Metric and morphological study of the upper cervical spine from

869 the Sima de los Huesos site (Sierra de Atapuerca, Burgos, Spain). *Journal of Human*
870 *Evolution* 53, 6–25.

871 Gómez-Robles, A., Bermúdez de Castro, J.M., Arsuaga, J.-L., Carbonell, E., Polly, P.D.,
872 2013a. No known hominin species matches the expected dental morphology of the last
873 common ancestor of Neanderthals and modern humans. *Proceedings of the National*
874 *Academy of Sciences USA* 110, 18196–18201.

875 Gómez-Robles, A., Hopkins, W.D., Sherwood, C.C., 2013b. Increased morphological
876 asymmetry, evolvability and plasticity in human brain evolution. *Proceedings of the*
877 *Royal Society B* 280, 20130575.

878 Gómez-Robles, A., Bermúdez de Castro, J.M., Martínón-Torres, M., Prado-Simón, L.,
879 Arsuaga, J.L., 2015a. A geometric morphometric analysis of hominin lower molars:
880 Evolutionary implications and overview of postcanine dental variation. *Journal of*
881 *Human Evolution* 82, 34–50.

882 Gómez-Robles, A., Hopkins, W.D., Schapiro, S.J., Sherwood, C.C., 2015b. Relaxed genetic
883 control of cortical organization in human brains compared with chimpanzees.
884 *Proceedings of the National Academy of Sciences USA* 112, 14799–14804.

885 Gómez-Robles, A., Smaers, J.B., Holloway, R.L., Polly, P.D., Wood, B.A., 2017. Brain
886 enlargement and dental reduction were not linked in hominin evolution. *Proceedings of*
887 *the National Academy of Sciences USA* 114, 468–473.

888 Good, C.D., Johnsrude, I., Ashburner, J., Henson, R.N.A., Friston, K.J., Frackowiak, R.S.J.,
889 2001. Cerebral asymmetry and the effects of sex and handedness on brain structure: A
890 voxel-based morphometric analysis of 465 normal adult human brains. *NeuroImage* 14,
891 685–700.

892 Grabowski, M., Voje, K.L., Hansen, T.F., 2016. Evolutionary modeling and correcting for
893 observation error support a 3/5 brain-body allometry for primates. *Journal of Human*
894 *Evolution* 94, 106–116.

895 Gracia, A., Arsuaga, J.L., Martínez, I., Lorenzo, C., Carretero, J.M., Bermúdez de Castro, J.M.,
896 Carbonell, E., 2009. Craniosynostosis in the Middle Pleistocene human Cranium 14
897 from the Sima de los Huesos, Atapuerca, Spain. *Proceedings of the National Academy*
898 *of Sciences USA* 106, 6573.

899 Grimaud-Hervé, D., 1997. L'évolution de l'encéphale chez l'*Homo erectus* et l'*Homo sapiens*,
900 Les cahiers de Paléanthropologie. CNRS Editions, Paris.

901 Gunz, P., Neubauer, S., Maureille, B., Hublin, J.-J., 2010. Brain development after birth differs
902 between Neanderthals and modern humans. *Current Biology* 20, R921–R922.

903 Holloway, R.L., 1981a. Revisiting the South African Taung australopithecine endocast: The
904 position of the lunate sulcus as determined by the stereoplotting technique. *American*
905 *Journal of Physical Anthropology* 56, 43–58.

906 Holloway, R.L., 1981b. Volumetric and asymmetry determinations on recent hominid
907 endocasts: Spy I and II, Djebel Ihroud I, and the Salè *Homo erectus* specimens, with
908 some notes on Neandertal brain size. *American Journal of Physical Anthropology* 55,
909 385–393.

910 Holloway, R.L., 1983. Cerebral brain endocast pattern of *Australopithecus afarensis* hominid.
911 *Nature* 303, 420.

912 Holloway, R.L., Broadfield, D.C., Yuan, M.S., Schwartz, J.H., Tattersall, I., 2004a. The human
913 fossil record. Brain endocasts—The paleoneurological evidence (Vol. 3). Wiley-Liss,
914 New York.

915 Holloway, R.L., Clarke, R.J., Tobias, P.V., 2004b. Posterior lunate sulcus in *Australopithecus*
916 *africanus*: Was Dart right? *Comptes Rendus Palevol* 3, 287–293.

917 Hopkins, W.D., Russell, J.L., Lambeth, S., Schapiro, S.J., 2007. Handedness and
 918 neuroanatomical asymmetries in captive chimpanzees: A summary of 15 years of
 919 research. In: Hopkins, W.D. (Ed.), *The Evolution of Hemispheric Specialization in*
 920 *Primates*. Elsevier, Oxford, pp. 146–181.

921 Hublin, J.-J., 1982. Les anténéandertaliens: Presapiens ou prénéandertaliens. *Geobios* 15, 345–
 922 357.

923 Hublin, J.-J., Ben-Ncer, A., Bailey, S.E., Freidline, S.E., Neubauer, S., Skinner, M.M.,
 924 Bergmann, I., Le Cabec, A., Benazzi, S., Harvati, K., Gunz, P., 2017. New fossils from
 925 Jebel Irhoud, Morocco and the pan-African origin of *Homo sapiens*. *Nature* 546, 289–
 926 292.

927 Jerison, H., 1973. *Evolution of the Brain and Intelligence*. Academic Press, New York.

928 Keller, S.S., Crow, T., Foundas, A., Amunts, K., Roberts, N., 2009. Broca's area:
 929 Nomenclature, anatomy, typology and asymmetry. *Brain and Language* 109, 29–48.

930 Kubo, D., Kono, R.T., Kaifu, Y., 2013. Brain size of *Homo floresiensis* and its evolutionary
 931 implications. *Proceedings of the Royal Society of London B* 280, 20130338.

932 LeMay, M., 1976. Morphological cerebral asymmetries of modern man, fossil man, and
 933 nonhuman primate. *Annals of the New York Academy of Sciences* 280, 349–366.

934 LeMay, M., Kido, D.K., 1978. Asymmetries of the cerebral hemispheres on computed
 935 tomograms. *Journal of Computer Assisted Tomography* 2, 471–476.

936 LeMay, M., Billig, M.S., Geschwind, N., 1982. Asymmetries of the brains and skulls of
 937 nonhuman primates. In: Armstrong, E., Falk, D. (Eds.), *Primate Brain Evolution:*
 938 *Methods and Concepts*. Springer, Boston, MA, pp. 263–277.

939 Lozano, M., Bermúdez de Castro, J.M., Carbonell, E., Arsuaga, J.L., 2008. Non-masticatory
 940 uses of anterior teeth of Sima de los Huesos individuals (Sierra de Atapuerca, Spain).
 941 *Journal of Human Evolution* 55, 713–728.

942 Lozano, M., Mosquera, M., Bermúdez de Castro, J.M., Arsuaga, J.L., Carbonell, E., 2009.
 943 Right handedness of *Homo heidelbergensis* from Sima de los Huesos (Atapuerca,
 944 Spain) 500,000 years ago. *Evolution and Human Behavior* 30, 369–376.
 945 Martin, R.D., 1981. Relative brain size and basal metabolic rate in terrestrial vertebrates.
 946 *Nature* 293, 57–60.
 947 Martínez, I., Arsuaga, J.L., 1997. The temporal bones from Sima de los Huesos Middle
 948 Pleistocene site (Sierra de Atapuerca, Spain). A phylogenetic approach. *Journal of*
 949 *Human Evolution* 33, 283–318.
 950 Martínez, I., Rosa, M., Arsuaga, J.-L., Jarabo, P., Quam, R., Lorenzo, C., Gracia, A., Carretero,
 951 J.-M., Bermúdez de Castro, J.M., Carbonell, E., 2004. Auditory capacities in Middle
 952 Pleistocene humans from the Sierra de Atapuerca in Spain. *Proceedings of the National*
 953 *Academy of Sciences USA* 101, 9976–9981.
 954 Martínez-Abadías, N., Esparza, M., Sjøvold, T., González-José, R., Santos, M., Hernández,
 955 M., Klingenberg, C.P., 2012. Pervasive genetic integration directs the evolution of
 956 human skull shape. *Evolution* 66, 1010–1023.
 957 Martínón-Torres, M., Bermúdez de Castro, J.M., Gómez-Robles, A., Prado-Simón, L.,
 958 Arsuaga, J.L., 2012. Morphological description and comparison of the dental remains
 959 from Atapuerca-Sima de los Huesos site (Spain). *Journal of Human Evolution* 62, 7–
 960 58.
 961 Meyer, M., Fu, Q., Aximu-Petri, A., Glocke, I., Nickel, B., Arsuaga, J.-L., Martínez, I., Gracia,
 962 A., Bermudez de Castro, J.M., Carbonell, E., 2014. A mitochondrial genome sequence
 963 of a hominin from Sima de los Huesos. *Nature* 505, 403–406.
 964 Meyer, M., Arsuaga, J.-L., de Filippo, C., Nagel, S., Aximu-Petri, A., Nickel, B., Martínez, I.,
 965 Gracia, A., Bermudez de Castro, J.M., Carbonell, E., Viola, B., Kelso, J., Prüfer, K.,

966 Pääbo, S., 2016. Nuclear DNA sequences from the Middle Pleistocene Sima de los
967 Huesos hominins. *Nature* 531, 504–507.

968 Mounier, A., Marchal, F., Condemi, S., 2009. Is *Homo heidelbergensis* a distinct species? New
969 insight on the Mauer mandible. *Journal of Human Evolution* 56, 219–246.

970 Neubauer, S., Gunz, P., 2018. Endocasts and the Evo-Devo approach to study human brain
971 evolution. In: Bruner, E., Ogihara, N., Tanabe, H.C. (Eds.), *Digital Endocasts: From*
972 *Skulls to Brains*. Springer, Tokyo, pp. 173–190.

973 Neubauer, S., Hublin, J.-J., Gunz, P., 2018. The evolution of modern human brain shape.
974 *Science Advances* 4, eaao5961.

975 Pablos, A., Pantoja-Pérez, A., Martínez, I., Lorenzo, C., Arsuaga, J.L., 2017. Metric and
976 morphological analysis of the foot in the Middle Pleistocene sample of Sima de los
977 Huesos (Sierra de Atapuerca, Burgos, Spain). *Quaternary International* 433, 103–113.

978 Pearce, E., Stringer, C., Dunbar, R.I.M., 2013. New insights into differences in brain
979 organization between Neanderthals and anatomically modern humans. *Proceedings of*
980 *the Royal Society B* 280, 20130168.

981 Ponce de León, M.S., Bienvenu, T., Akazawa, T., Zollikofer, C.P.E., 2016. Brain development
982 is similar in Neanderthals and modern humans. *Current Biology* 26, R665–R666.

983 Poza-Rey, E., 2015. Estudio del encéfalo de los homínidos del yacimiento de La Sima de los
984 Huesos (Atapuerca, Burgos), a partir del análisis de sus moldes endocraneales. Ph.D.
985 Dissertation, Complutense University of Madrid.

986 Poza-Rey, E.M., Lozano, M., Arsuaga, J.L., 2017. Brain asymmetries and handedness in the
987 specimens from the Sima de los Huesos site (Atapuerca, Spain). *Quaternary*
988 *International* 433, 32–44.

989 Quam, R., Lorenzo, C., Martínez, I., Gracia-Téllez, A., Arsuaga, J.L., 2016. The bony labyrinth
990 of the middle Pleistocene Sima de los Huesos hominins (Sierra de Atapuerca, Spain).
991 Journal of Human Evolution 90, 1–15.

992 Richter, D., Grün, R., Joannes-Boyau, R., Steele, T.E., Amani, F., Rué, M., Fernandes, P.,
993 Raynal, J.-P., Geraads, D., Ben-Ncer, A., Hublin, J.-J., McPherron, S.P., 2017. The age
994 of the hominin fossils from Jebel Irhoud, Morocco, and the origins of the Middle Stone
995 Age. Nature 546, 293–296.

996 Rodríguez, L., Carretero, J.M., García-González, R., Lorenzo, C., Gómez-Olivencia, A.,
997 Quam, R., Martínez, I., Gracia-Téllez, A., Arsuaga, J.L., 2016. Fossil hominin radii
998 from the Sima de los Huesos Middle Pleistocene site (Sierra de Atapuerca, Spain).
999 Journal of Human Evolution 90, 55–73.

1000 Rodríguez-Hidalgo, A., Saladié, P., Ollé, A., Arsuaga, J.L., Bermúdez de Castro, J.M.,
1001 Carbonell, E., 2017. Human predatory behavior and the social implications of
1002 communal hunting based on evidence from the TD10.2 bison bone bed at Gran Dolina
1003 (Atapuerca, Spain). Journal of Human Evolution 105, 89–122.

1004 Roksandic, M., Radović, P., Lindal, J., 2018. Revising the hypodigm of *Homo heidelbergensis*:
1005 A view from the Eastern Mediterranean. Quaternary International 466, 66–81.

1006 Rosas, A., 1997. A gradient of size and shape for the Atapuerca sample and Middle Pleistocene
1007 hominid variability. Journal of Human Evolution 33, 319–331.

1008 Ruff, C.B., Trinkaus, E., Holliday, T.W., 1997. Body mass and encephalization in Pleistocene
1009 *Homo*. Nature 387, 173.

1010 Sala, N., Arsuaga, J.L., Pantoja-Pérez, A., Pablos, A., Martínez, I., Quam, R.M., Gómez-
1011 Olivencia, A., Bermúdez de Castro, J.M., Carbonell, E., 2015. Lethal interpersonal
1012 violence in the Middle Pleistocene. PLOS ONE 10, e0126589.

1013 Schenker, N.M., Hopkins, W.D., Spocter, M.A., Garrison, A.R., Stimpson, C.D., Erwin, J.M.,
1014 Hof, P.R., Sherwood, C.C., 2010. Broca's area homologue in chimpanzees (*Pan*
1015 *trogloodytes*): Probabilistic mapping, asymmetry, and comparison to humans. *Cerebral*
1016 *Cortex* 20, 730–742.

1017 Seidler, H., Falk, D., Stringer, C., Wilfing, H., Müller, G.B., zur Nedden, D., Weber, G.W.,
1018 Reicheis, W., Arsuaga, J.-L., 1997. A comparative study of stereolithographically
1019 modelled skulls of Petralona and Broken Hill: Implications for future studies of Middle
1020 Pleistocene hominid evolution. *Journal of Human Evolution* 33, 691–703.

1021 Sherwood, C.C., Subiaul, F., Zawidzki, T.W., 2008. A natural history of the human mind:
1022 Tracing evolutionary changes in brain and cognition. *Journal of Anatomy* 212, 426–
1023 454.

1024 Spoor, F., Gunz, P., Neubauer, S., Stelzer, S., Scott, N., Kwekason, A., Dean, M.C., 2015.
1025 Reconstructed *Homo habilis* type OH 7 suggests deep-rooted species diversity in early
1026 *Homo*. *Nature* 519, 83–86.

1027 Springer, J.A., Binder, J.R., Hammeke, T.A., Swanson, S.J., Frost, J.A., Bellgowan, P.S.,
1028 Brewer, C.C., Perry, H.M., Morris, G.L., Mueller, W.M., 1999. Language dominance
1029 in neurologically normal and epilepsy subjects: A functional MRI study. *Brain* 122,
1030 2033–2046.

1031 Stringer, C., 2012. The status of *Homo heidelbergensis* (Schoetensack 1908). *Evolutionary*
1032 *Anthropology* 21, 101–107.

1033 Tattersall, I., 2007. Neanderthals, *Homo sapiens*, and the question of species in
1034 paleoanthropology. *Journal of Anthropological Sciences* 85, 139–146.

1035 Tobias, P.V., 1987. The brain of *Homo habilis*: A new level of organization in cerebral
1036 evolution. *Journal of Human Evolution* 16, 741–761.

1037 Toth, N., 1985. Archaeological evidence for preferential right-handedness in the Lower and
 1038 Middle Pleistocene, and its possible implications. *Journal of Human Evolution* 14, 607–
 1039 614.

1040 Traynor, S., Gurtov, A.N., Senjem, J.H., Hawks, J., 2015. Assessing eye orbits as predictors of
 1041 Neandertal group size. *American Journal of Physical Anthropology* 157, 680–683.

1042 Wood, B., Boyle, E.K., 2016. Hominin taxic diversity: Fact or fantasy? *American Journal of*
 1043 *Physical Anthropology* 159, 37–78.

1044 Wu, X., Schepartz, L.A., Falk, D., Liu, W., 2006. Endocranial cast of Hexian *Homo erectus*
 1045 from South China. *American Journal of Physical Anthropology* 130, 445–454.

1046 Wu, X., Schepartz, L.A., Liu, W., 2009. A new *Homo erectus* (Zhoukoudian V) brain endocast
 1047 from China. *Proceedings of the Royal Society B* 277, 337–344.

1048 Wu, X., Schepartz, L.A., Norton, C.J., 2010. Morphological and morphometric analysis of
 1049 variation in the Zhoukoudian *Homo erectus* brain endocasts. *Quaternary International*
 1050 211, 4–13.

1051 Zollikofer, C.P.E., Bienvenu, T., Ponce de León, M.S., 2017. Effects of cranial integration on
 1052 hominid endocranial shape. *Journal of Anatomy* 230, 85–105.

1053

Figure legends

Figure 1. Anatomy of the encephalic rostrum (top row) showing configurations observed in *H. erectus* (type 1 as shown by Zhoukoudian 12), SH hominins (type 1-2 as shown by SH4) and *H. sapiens* (type 3 as shown by a recent modern human; Grimaud-Hervé, 1997). Degree of projection of the occipital lobes (bottom row) showing slightly projecting (SH6), moderately projecting (SH3, mirror-imaged) and highly projecting (SH4) configurations. Broca's cap region is highlighted in red in the lateral view of SH4. Endocasts are not to scale.

Figure 2. Analysis of petalial patterns. Left-right differences are calculated for the length and width of the frontal and occipital poles as measured at the points marking 10% of total endocranial length. Further details and quantifications are provided in Poza-Rey et al. (2017).

Figure 3. Linear metrics used in the quantitative analysis of the SH sample and in the comparisons with other endocasts. Linear measurements and landmarks are defined in Tables 2 and 3.

Figure 4. Endocast SH2. Six different views are provided (top row: left lateral, superior, right lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black. Scale bar is 10 cm.

Figure 5. Endocast SH3. Six different views are provided (top row: left lateral, superior, right lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black. Scale bar is 10 cm.

1078

1079 **Figure 6.** Endocast SH4. Six different views are provided (top row: left lateral, superior, right
1080 lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black.
1081 Scale bar is 10 cm.

1082

1083 **Figure 7.** Endocast SH5. Six different views are provided (top row: left lateral, superior, right
1084 lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black.
1085 Scale bar is 10 cm.

1086

1087 **Figure 8.** Endocast SH6. Six different views are provided (top row: left lateral, superior, right
1088 lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black.
1089 Scale bar is 10 cm.

1090

1091 **Figure 9.** Endocast SH7. Six different views are provided (top row: left lateral, superior, right
1092 lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black.
1093 Scale bar is 10 cm.

1094

1095 **Figure 10.** Endocast SH8. Six different views are provided (top row: left lateral, superior, right
1096 lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black.
1097 Scale bar is 10 cm.

1098

1099 **Figure 11.** Endocast SH9. Six different views are provided (top row: left lateral, superior, right
1100 lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black.
1101 Scale bar is 10 cm.

1102

Figure 12. Endocast SH10. Six different views are provided (top row: left lateral, superior, right lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black. Scale bar is 10 cm.

Figure 13. Endocast SH11. Six different views are provided (top row: left lateral, superior, right lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black. Scale bar is 10 cm.

Figure 14. Endocast SH12. Six different views are provided (top row: left lateral, superior, right lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black. Scale bar is 10 cm.

Figure 15. Endocast SH13. Six different views are provided (top row: left lateral, superior, right lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black. Scale bar is 10 cm.

Figure 16. Endocast SH14. Six different views are provided (top row: left lateral, superior, right lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black. Scale bar is 10 cm.

Figure 17. Endocast SH15. Six different views are provided (top row: left lateral, superior, right lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black. Scale bar is 10 cm.

Figure 18. Endocast SH16. Six different views are provided (top row: left lateral, superior, right lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black. Scale bar is 10 cm.

Figure 19. Endocast SH17. Six different views are provided (top row: left lateral, superior, right lateral; bottom row: anterior, inferior and posterior). Missing parts are represented in black. Scale bar is 10 cm.

Figure 20. Posterior view of four hominin endocasts showing the configurations observed in *H. erectus* (tent-like or “en tente”, Zhoukoudian 12), SH-hominins (roof-like, SH4), *H. neanderthalensis* (rounded or “en bombe”, Saccopastore 1) and *H. sapiens* (house-like or “en maison”, recent modern human). Endocasts are not to scale.

Figure 21. Variation in brain size (top left) and encephalization quotients (top right) across hominins. EQ data for all the groups except Sima de los Huesos are based on unpublished data from Grabowski et al. (2016) with some minor taxonomic reallocations. The *H. sapiens* sample does not include recent individuals, but only fossil *H. sapiens*. Bottom panels show the possible range of variation in EQ of all SH individuals when considering the range of variation in body mass for the complete sample (bottom left) and when separating male and female body masses (bottom right). SH individuals in the bottom panels are sorted according to their increasing endocranial volumes.

Figure 22. Principal components analysis of endocranial shape variation based on seven size-adjusted linear metrics (graphically represented in the insets on the right). The two plots on the

1151 top of the figure correspond to the analysis of all groups, including recent modern humans. The
1152 two plots on the bottom of the figure include only fossil specimens to better ascertain the
1153 relationships between non-recent specimens. Plots on the right show the same analyses as those
1154 on the left without specimen labels, but with variable loadings on PC1 and PC2.

1155

H. erectus

SH

H. sapiens

SH6

SH3

SH4

SH2

SH3

SH5

SH6

SH7

SH8

SH9

SH11

SH12

SH14

SH15

SH16

SH17

H. erectus

SH

H. neanderthalensis

H. sapiens

Table 1. Comparative sample used for metric analyses

Individual	Abbreviation	Taxon
Ngandong 6	Ng6	<i>H. erectus</i>
Ngandong 7	Ng7	<i>H. erectus</i>
Ngandong 11	Ng11	<i>H. erectus</i>
Ngandong 12	Ng12	<i>H. erectus</i>
Sambungmacan 3	Sb3	<i>H. erectus</i>
Trinil 2	Tr2	<i>H. erectus</i>
Zhoukoudian 3	Zk3	<i>H. erectus</i>
Zhoukoudian 12	Zk12	<i>H. erectus</i>
Arago 21-47	A21-47	Middle Pleistocene <i>Homo</i>
Kabwe	Kb	Middle Pleistocene <i>Homo</i>
Petralona	Ptr	Middle Pleistocene <i>Homo</i>
Ehringsdorf 9	Eh9	<i>H. neanderthalensis</i>
Feldhofer	Fdh	<i>H. neanderthalensis</i>
La Chapelle-aux-Saints	LCh	<i>H. neanderthalensis</i>
La Ferrassie 1	LF	<i>H. neanderthalensis</i>
La Quina 5	LQ5	<i>H. neanderthalensis</i>
Teshik Tash	TT	<i>H. neanderthalensis</i>
Cro-Magnon 3	CM3	Fossil <i>H. sapiens</i>
Jebel Irhoud 1	Jl1	Fossil <i>H. sapiens</i>
Předmostí 10	Pr10	Fossil <i>H. sapiens</i>
Singa	Sng	Fossil <i>H. sapiens</i>
25 recent modern humans*		Recent <i>H. sapiens</i>

*Recent modern humans are non-pathological adults from the anthropological collections of the Musée de l'Homme, Paris. Most individuals are from recent historical times and their geographic origin is diverse.

Table 2. List of landmarks used to obtain endocranial metrics.

Landmark abbreviation	Definition
fp	Frontal pole: the most anterior point of the frontal lobe.
fv	The most ventral point of the frontal lobe.
tv	The most ventral point of temporal lobe.
cv	The most ventral point of the cerebellar hemisphere.
op	Occipital pole: the most posterior point of the occipital lobe.
o	The most laterally projected point of the occipital lobe.
ev	Endovortex: the most superior point of the endocast, perpendicular to the hemispheric length (L).
rf/sg	Point where the central sulcus (Roland fissure) and the sagittal sinus intersect.
pf/sg	Point where the perpendicular fissure and the sagittal sinus intersect.
eat	Point where the perpendicular fissure and the lateral sinus intersect.
eba	Endobasion: middle point on the anterior margin of the foramen magnum.
bat	The most anterior point on the temporal lobe in basal view.
mat	The most lateral point on the endocast at the level of “bat” point in basal view.
mbat	Middle point on the line connecting the two “bat” points.
fl	The most laterally projected point of the frontal lobe.
lat	The most laterally projected point of the endocast.
Br	The most laterally projected point of the Broca’s region.
rof	The most anteriorly projected point on the orbital surfaces of the frontal lobes in basal view.
bpc	The most posteriorly projected point of the occipital lobes in basal view.

Table 3. List of linear metrics measured in the SH sample and in the comparative sample.

Measurements	
Lateral view	
L	Total length; average value of both hemispheres (fp-op).
H	Total height (ev-cv).
CH	Cerebral height (ev-tv).
FC	Frontal chord (fp-rs/sg).
PC	Parietal chord (rf/sg- pf/sg).
FH	Frontal height (rf/sg- fv).
rf/sg-eba	Distance between the intersection point between the Roland fissure with the sagittal sinus and the endobasion point.
rf/sg-cv	Distance between the intersection point between the Roland fissure with the sagittal sinus and the most ventral point of the cerebellum.
rf/sg-eat	Distance between the intersection point between the Roland fissure with the sagittal sinus and the intersection point between the perpendicular fissure and the lateral sinus.
P	Distance between op and cv.
Basal view	
bat-bat	Distance between the most anterior points of the temporal lobes.
mat-mat	Width of the frontal lobes at the level of bat.
mbat-rof	The shortest distance between the mid-point of the line connecting the two bats and the tangent to the most rostral point on the orbital surfaces of the frontal lobes.
BW	Width of the endocast at the level of Broca's cap.
Dorsal view	
FW	Width of the frontal lobe.
MW	Maximum width of the brain endocast.
OW	Width of the occipital lobe.

Table 4. Summary of the anatomical descriptions of SH endocasts

Specimen	Preservation	Broca's cap asymmetry	Petalias	Rostrum development	Occipital lobes
SH2	Rather incomplete	--	--	--	Slightly projected
SH3	Moderately incomplete	--	--	--	Moderately projected
SH4	Complete	$L \approx R$	L-frontal, L-occipital (length) R-frontal, R-occipital (width)	1-2	Highly projected
SH5	Almost complete	$L \square R$	R-frontal, L-occipital (length) R-frontal, L-occipital (width)	1-2	Moderately projected
SH6	Moderately incomplete	$L \approx R$	No frontal, L-occipital (length) No frontal, L-occipital (width)	--	Slightly projected
SH7	Moderately incomplete	--	L-frontal, L-occipital (length) No frontal, L-occipital (width)	--	Slightly projected
SH8	Very incomplete	--	--	--	--
SH9	Moderately incomplete	--	R-frontal, L-occipital (length) L-frontal, L-occipital (width)	--	Highly projected
SH10	Rather incomplete	$L \square R$	--	--	--
SH11	Rather incomplete	--	No frontal, L-occipital (length) R-frontal, ?-occipital (width)	1-2	Slightly projected

SH12	Rather incomplete	--	--	--	--
SH13	Almost complete (R) and very incomplete (L)	L□R	--	1-2	--
SH14	Almost complete	L□R	R-frontal, L-occipital (length) R-frontal, No occipital (width)	1-2	Slightly projected
SH15	Rather incomplete	--	--	--	--
SH16	Rather incomplete	L□R	No frontal, No occipital (length) No frontal, ?-occipital (width)	1-2	Slightly projected
SH17	Almost complete (L) and rather incomplete (R)	L□R	R-frontal, No occipital (length) R-frontal, ?-occipital (width)	1-2	Moderately projected

Table 5. Endocranial volume and encephalization quotients in SH individuals

Specimen	Endocranial volume (cc)	EQ mean (range)	Assigned sex	EQ as male or female
SH2	1333.5 $\pm$ 3.5	6.79 (6.24-7.43)	Male	6.34 (6.06-6.65)
SH3*	1230 $\pm$ 8	6.27 (5.76-6.86)	Male	5.85 (5.59-6.14)
SH4	1360**	6.93 (6.37-7.58)	Male	6.47 (6.18-6.79)
SH5	1092**	5.57 (5.11-6.09)	Female	6.22 (5.97-6.49)
SH6	1225 $\pm$ 21	6.24 (5.73-6.83)	Female	6.98 (6.70-7.28)
SH7	1143.5 $\pm$ 26.5	5.83 (5.35-6.37)	Female	6.51 (6.25-6.80)
SH8	1313 $\pm$ 31	6.69 (6.16-7.32)	Male	6.25 (5.97-6.55)
SH9	1201 $\pm$ 9.5	6.12 (5.62-6.70)	Female	6.84 (6.56-7.14)
SH10	1218 $\pm$ 5	6.21 (5.70-6.79)	Female	6.94 (6.66-7.24)
SH11*	1057 $\pm$ 23.5	5.38 (4.95-5.89)	Female	6.02 (5.78-6.29)
SH12	1227.5 $\pm$ 0.5	6.25 (5.74-6.84)	Female	6.99 (6.71-7.30)
SH13	1436.5 $\pm$ 29.5	7.32 (6.72-8.01)	Male	6.83 (6.52-7.17)
SH14*	1224 $\pm$ 8	6.24 (5.73-6.83)	Female	6.98 (6.69-7.28)
SH15	1283.5 $\pm$ 0.5	6.54 (6.00-7.15)	Male	6.11 (5.83-6.40)
SH16	1236 $\pm$ 10	6.30 (5.78-6.89)	Male	5.88 (5.61-6.17)
SH17	1218 $\pm$ 7	6.53 (6.00-7.14)	Male	6.10 (5.82-6.39)

* Immature individuals whose adult ECV may have not been attained

** Endocranial volumes measured in complete endocasts. All the other values have been estimated by scaling the complete endocasts (SH4 and SH5) to the size of incomplete endocasts using 15 endocranial landmarks as described in Arsuaga et al. (2014) and Poza-Rey (2015).

Table 6. Linear metrics in the SH sample. Values are provided in mm.

Ind	L	H	CH	FC	PC	FH	rs/sg-eba	rs/sg-cv	rs/sg-eat	P	bat-bat	mat-mat	mbat-rof	BW	FW	MW	OW
SH2					52.21										122.28*	136.28*	97.97
SH3					54.98				101.83		61.22*			96.3	107.9	134.98	98.32
SH4	174.35	124.4	113.95	119.26	50.53	115.4	113.48	119.82	107.22	76.37	73.47	113.15	47.15	101.61	115.98	146.62	113.62
SH5	159.86	118.63	107.19	110.72	47.8	105.27	110.02	114.56	98.35	62.43	64.82	98.16	51.14	92.15	106.54	129.22	105.97
SH6	167.57									69.34	62.28	101.6	47.44	99.73	106.23	129.72	97.21
SH7	161.8	113.22		114.43	57.92			112.45		71.25	62.53*			88.70*	104.34	126.76*	98.09*
SH8																	
SH9	172.24			115.55	47.47				94.08					92.88*	113.58	134.93	102.41
SH10											68.08			95.72	109.9*	131.49*	
SH11	158.76			118.03	51.75		109.23	114.72	94.28	60.92				98.18	113.53*	129.34*	93.76
SH12											68.32*			100.32	116.61	132.61*	
SH13								123.3			73.08*			101.4	114.45	141.78	
SH14	161.83		110.52	114.59	57.81				99.64		67.68			94.38	113.56	132.43	91.14
SH15	167.24			117.48	54.65				104.32		91.18*			103.26*	114.01*	140.36*	95.69*

SH16	167.88									84.11	68.85*			97	103.84*	131.39*	97.52*
SH17	168.18	128.05		111.74	59.2	108.54		121.8	100.71	70.93	64.5			93.47	108.99*	132.86*	102.04*

* Values that have been estimated in incomplete endocasts by mirror-imaging preserved parts.

Table 7. Correlations between linear metrics in the SH sample. Sample sizes for each comparison are provided above the diagonal, and correlation coefficients and *p*-values (in parentheses) under the diagonal.

	L	H	CH	FC	PC	FH	rs/sg-eba	rs/sg-cv	rs/sg-eat	P	bat-bat	mat-mat	mbat-rof	BW	FW	MW	OW	
L		4	3	8	8	3	3	5	7	7	8	3	3	10	10	10	10	L
H			2	4	4	3	2	4	3	4	4	2	2	4	4	4	4	H
CH				3	3	2	2	2	3	2	3	2	2	3	3	3	3	CH
FC	0.36 (0.397)				8	3	3	5	7	5	6	2	2	8	8	8	8	FC
PC	-0.20 (0.653)			-0.17 (0.700)		3	3	5	8	5	7	2	2	9	10	10	10	PC
FH							2	3	3	3	3	2	2	3	3	3	3	FH
rs/sg-eba						2		3	3	3	2	2	2	3	3	3	3	rs/sg-eba
rs/sg-cv									4	5	5	2	2	6	6	6	5	rs/sg-cv
rs/sg-eat	0.47 (0.308)			0.28 (0.565)	0.32 (0.464)					4	6	2	2	8	8	8	8	rs/sg-eat
P	0.72 (0.071)										6	3	3	7	7	7	7	P
bat-bat	0.33 (0.450)			0.65 (0.179)	-0.15 (0.769)				0.56 (0.272)	0.59 (0.244)		3	3	12	12	12	9	bat-bat
mat-mat													3	3	3	3	3	mat-mat
mbat-rof														3	3	3	3	mbat-rof
BW	0.39 (0.502)			0.74 (0.035)	-0.15 (0.715)			0.64 (0.185)	0.63 (0.097)	0.21 (0.667)	0.66 (0.011)				14	14	11	BW

FW	0.29 (0.430)			0.75 (0.029)	-0.28 (0.454)			0.60 (0.225)	0.17 (0.707)	-0.24 (0.631)	0.55 (0.040)			0.56 (0.034)		15	12	FW
MW	0.73 (0.015)			0.62 (0.108)	-0.21 (0.565)			0.75 (0.094)	0.80 (0.015)	0.39 (0.405)	0.64 (0.066)			0.66 (0.008)	0.58 (0.023)		12	MW
OW	0.56 (0.094)			0.01 (0.981)	-0.49 (0.161)				0.40 (0.343)	0.18 (0.713)	-0.03 (0.806)			0.04 (0.905)	0.05 (0.885)	0.51 (0.091)		OW
	L	H	CH	FC	PC	FH	rs/sg-eba	rs/sg-cv	rs/sg-eat	P	bat-bat	mat-mat	mbat-rof	BW	FW	MW	OW	

Bold: comparisons involving more than five individuals (above the diagonal) and significant correlations (below the diagonal). Correlations have been calculated only when pairs of variables have been measured in more than five individuals.

Table 8. Variable loadings corresponding to the principal components analysis of all groups and of all the fossil groups (excluding recent modern humans).

	All groups		Only fossils	
Variable	PC1 loading	PC2 loading	PC1 loading	PC2 loading
Length (L)	0.398	0.499	-0.266	0.150
Frontal chord (FC)	0.232	0.643	-0.747	-0.448
Parietal chord (PC)	-0.701	0.114	0.529	-0.072
Width at Broca's cap (BW)	-0.013	-0.201	0.104	-0.311
Frontal width (FW)	0.001	-0.195	0.087	-0.265
Maximum width (MW)	0.380	-0.328	-0.100	0.490
Occipital width (OW)	0.391	-0.371	-0.252	-0.604