[Click here to view linked References](#)

1 Martin H. Trauth^{*1}, Asfawossen Asrat², Walter Duesing¹, Verena Foerster³, K. Hauke Kraemer^{1,4}, Norbert
2 Marwan⁴, Mark A. Maslin⁵, Frank Schaebitz³

3

4 Classifying past climate change in the Chew Bahir basin, 5 southern Ethiopia, using recurrence quantification analysis

6

7

8 ¹ University of Potsdam, Institute of Earth and Environmental Science, Potsdam, Germany

9 ² Addis Ababa University, School of Earth Sciences, Addis Ababa, Ethiopia

10 ³ University of Cologne, Institute of Geography Education, Cologne, Germany

11 ⁴ Potsdam Institute for Climate Impact Research, Potsdam, Germany

12 ⁵ Department of Geography, University College London, London, UK

13

14 *Corresponding Author: email trauth@uni-potsdam.de, Tel. +49 331 977 5810

15

16 Acknowledgements

17 Our project was funded by the DFG priority program SPP 1006 ICDP to M.H.T. and F.S. as well as the CRC
18 806 "Our Way to Europe" to F.S. at the University of Cologne. The project also received funds from a DFG
19 grant to N.M. and M.H.T. to improve the recurrence plots/recurrence quantification analysis methods. This
20 paper is publication number ### of the Hominin Sites and Paleolakes Drilling Project (HSPDP).

21 Abstract

22 The Chew Bahir Drilling Project (CBDP) aims to test possible linkages between climate and evolution in Africa
23 through the analysis of sediment cores that have recorded environmental changes in the Chew Bahir basin.
24 In this statistical project we consider the Chew Bahir palaeolake to be a dynamical system consisting of
25 interactions between its different components, such as the waterbody, the sediment beneath lake, and the
26 organisms living within and around the lake. Recurrence is a common feature of such dynamical systems, with
27 recurring patterns in the state of the system reflecting typical influences. Identifying and defining these
28 influences contributes significantly to our understanding of the dynamics of the system. Different recurring
29 changes in precipitation, evaporation, and wind speed in the Chew Bahir basin could result in similar (but not
30 identical) conditions in the lake (e.g., depth and area of the lake, alkalinity and salinity of the lake water, species
31 assemblages in the water body, and diagenesis in the sediments). Recurrence plots (RPs) are graphic displays
32 of such recurring states within a system. Measures of complexity were subsequently introduced to complement
33 the visual inspection of recurrence plots, and provide quantitative descriptions for use in recurrence
34 quantification analysis (RQA). We present and discuss herein results from an RQA on the environmental
35 record from six short (<17 m) sediment cores collected during the CBDP, spanning the last 45 kyrs. The
36 different types of variability and transitions in these records were classified to improve our understanding of
37 the response of the biosphere to climate change, and especially the response of humans in the area.

38 Keywords

39 Paleoclimate dynamics, eastern Africa, Pleistocene, Holocene, time-series analysis, recurrence plot

40 Introduction

41 The development by humans of settlement systems, artistic representations, and hunting strategies during the
42 last 50–40 kyrs marks a "human revolution" and the emergence of behaviorally modern humans in Africa
43 ([Renfrew, 2009](#); [Richter et al., 2012](#)). These behaviorally modern humans may have adapted to environmental

44 changes and extreme climatic oscillations through technological, behavioural, cultural and cognitive innovation
45 rather than through physical adaptation ([Klein, 1995](#); [Klein and Steele, 2013](#); [Clarke et al., 2016](#)). Determining
46 the nature and pace of changes in the environment of early modern humans is crucial to understanding the
47 factors that influenced this human revolution. For example, different types of climate variability would have
48 resulted in different types of climatic stress and changes to environmental boundaries ([Hildebrand and Grillo,](#)
49 [2012](#); [Vogelsang and Keding, 2013](#); [Foerster et al., 2015](#)).

50
51 There are currently ongoing discussions concerning global and regional climate fluctuations that had an effect
52 on human habitats (e.g. [Trauth et al., 2007](#); [Foerster et al., 2015](#); [Lamb et al., 2018](#); [Ivory and Russell, 2018](#);
53 [Maley et al., 2018](#); [Garcin et al., 2018](#); [Trauth et al., 2018](#)) and which did not, either because these fluctuations
54 had little or no effect on the African climate (e.g. [Timmerman and Friedrich, 2016](#)) or because their effect was
55 buffered by the environmental system (e.g. [Cuthbert et al., 2017](#)). Among the most debated episodes of African
56 climate (including their onset, termination, and internal variability) are the Dansgaard-Oeschger (DO) cycles
57 and Heinrich Events (HEs) (~110–12 kyr BP, e.g. [Brown et al., 2007](#); [Lamb et al., 2007](#); [Garcin, 2008](#); [Tierney](#)
58 [and deMenocal, 2013](#); [Berke et al., 2014](#); [Timmerman and Friedrich, 2016](#)), the marine isotope stage 4 (MIS
59 4) aridification (~71 kyr BP, e.g. [Timmerman and Friedrich, 2016](#)), the Last Glacial Maximum (LGM, 23.5–18
60 kyr BP, e.g. [Gasse, 2000](#); [Shakun and Carlson, 2010](#); [Tierney and deMenocal, 2013](#)), and the African Humid
61 Period (AHP, ~15–5 kyr BP, e.g. [deMenocal et al., 2000](#); [Tierney and deMenocal, 2013](#); [Shanahan et al.,](#)
62 [2015](#); [Tierney et al., 2017](#)). Such global (DO cycles, HEs, LGM) and regional (AHP) episodes may have
63 affected the availability of water and food, spatial retreats and shelter, and migration corridors, over variable
64 periods of time (e.g. [Ambrose et al., 1998](#); [Carto et al., 2009](#); [Castañeda et al., 2009](#); [Brandt et al., 2012](#);
65 [Foerster et al., 2015](#); [Flohr et al., 2016](#); [Marchant et al., 2018](#)).

66
67 Time-series analysis provides a number of tools with which to characterize past climate change, which can be
68 random, clustered, cyclic, or chaotic (e.g. [Marwan et al., 2007](#); [Mudelsee, 2014](#); [Trauth, 2015](#)). The most
69 popular methods for characterizing variations are based on Fourier or wavelet transforms, decomposing time
70 series into a linear combination of sinusoids (e.g. [Trauth, 2015](#) and references therein). Past climate change

71 is, however, often nonlinear (i.e. there is no simple proportional relation between cause and effect) and
72 techniques to describe nonlinear behavior have therefore become increasingly popular in recent decades,
73 defining the scaling laws and fractal dimensions of natural processes ([Turcotte 1997](#); [Kantz and Schreiber,](#)
74 [1997](#); [Tsonis, 2016](#)) and detecting nonlinear interrelationships using methods such as transfer entropies,
75 graphic models, and recurrence plots ([Zbilut and Webber, 1992](#); [Marwan et al., 2003](#); [Rodó and Rodríguez-](#)
76 [Arias, 2006](#); [Feldhoff et al., 2012](#); [Goswami et al., 2013](#); [Runge et al., 2012, 2014](#); [Builes-Jaramillo et al., 2018](#);
77 [Ramos et al., 2018](#)). The availability of long time series in the Earth sciences in recent times (for example from
78 multi-sensor core logger and micro X-ray fluorescence results) facilitates the use of these methods and
79 increases the reliability of the results obtained.

80
81 In this paper we present a classification of past climate variability in the Chew Bahir basin of southern Ethiopia
82 over approximately the last 45 kyrs using recurrence plots, which provide a graphic display of recurring states
83 in the environmental system ([Eckmann et al., 1987](#); [Marwan et al., 2007](#)). Quantitative descriptions (measures
84 of complexity) have been developed to complement visual inspection of recurrence plots (RPs) and for
85 recurrence quantification analysis (RQA) (e.g. [Zbilut & Webber, 1992](#); [Marwan et al., 2007](#); [Marwan, 2008](#)).
86 We previously used recurrence plots to identify past climate transitions during the Plio-Pleistocene in Africa,
87 the Middle East, and East Asia (e.g., [Donges et al., 2011](#); [Eroglu et al., 2016](#)). Such plots enable us to detect
88 nonlinear patterns in past climate change, helping to improve our understanding of the underlying process of
89 climate transitions in the Chew Bahir basin by statistically describing the dynamical characteristics of the
90 environment ([Marwan et al., 2007](#); [Donges et al., 2011](#); [Marwan et al., 2013](#); [Eroglu et al., 2014](#); [Marwan and](#)
91 [Kurths, 2015](#)). We first used the method on prototypical data in order to assess its performance with typical
92 palaeoclimate transitions. We then performed an RQA on the Late Quaternary climate record from Chew Bahir
93 cores CB01–06 because it provides one of the most detailed records of climate change available from the
94 vicinity of important sites for modern human fossil ([Foerster et al., 2012, 2015, 2018](#); [Trauth et al., 2015, 2018](#)).

95 Regional setting

96 The sediment cores described herein were recovered from the Chew Bahir basin in the southern Ethiopian
97 Rift (4.1–6.3°N, 36.5–38.1°E; [Fig. 1](#)) (see details in [Foerster et al., 2012](#)). Chew Bahir is a closed basin,
98 separated from the Turkana Basin to the west by the Hammar Range. The western part of the 32,400 km²
99 catchment, drained by the perennial Segen and Weyto rivers, is mostly formed by Late Proterozoic gneisses,
100 whereas the eastern part is dominated by Miocene basalts ([Moore and Davidson, 1978](#); [Davidson, 1983](#)).
101 Rainfall in the area is associated with the seasonal migration of the Intertropical Convergence Zone (ITCZ),
102 resulting in two rainy seasons in March–May and October–November ([Nicholson, 2017](#)). Rainfall intensity
103 strongly depends on Atlantic and Indian Ocean sea-surface temperature (SST) variations caused by the Indian
104 Ocean Dipole (IOD) and the El Niño-Southern Oscillation (ENSO), also explaining the recent reduction of
105 rainfall intensities over the last decades ([Saji et al., 1999](#); [Seleshi and Zanke, 2004](#); [Cheung et al., 2008](#);
106 [Segele et al., 2009](#); [Nicholson, 2017](#)).

107 Methods

108 *The Late Pleistocene–Holocene record of the Chew Bahir basin*

109 We reconstructed climate fluctuations in the Chew Bahir from six short (<17 m) sediment cores (CB01–06),
110 collected along a ~20 km transect from Northwest to Southeast in the basin ([Foerster et al., 2012, 2015](#); [Trauth](#)
111 [et al., in 2018](#)) ([Fig. 1](#)). These six cores were collected in a pilot study for the Hominin Sites and Paleolakes
112 Drilling Project (HSPDP; HSPDP-CHB deep coring site in [Fig. 1](#)) ([Cohen et al., 2016](#); [Campisano et al., 2017](#))
113 and the Collaborative Research Center CRC-806 at the University of Cologne, and were described in detail in
114 earlier publications (e.g., [Foerster et al., 2012, 2015](#); [Trauth et al., 2015](#)). We used the potassium (K)
115 concentrations of the sediment, determined by micro X-ray fluorescence (μXRF) scanning, as a proxy of aridity
116 ([Foerster et al., 2012](#); [Trauth et al., 2015](#); [Foerster et al., 2018](#)). Dynamic time warping (DTW) was employed
117 for aligning the K records from the six cores CB01–06 ([Trauth et al., 2018](#)). The composite age model of [Trauth](#)
118 [et al. \(2015\)](#), based on 32 AMS ¹⁴C ages derived from biogenic carbonate, fossilized charcoal and organic

119 sediment, resulting in a very solid chronology for lake record spanning the last ~45 kyr, was then used to
120 convert composite core depths into ages.

121

122 A principal component analysis (PCA) helped us to separate the mixed regional and local environmental
123 signals in the K records from the six aligned cores. The first principal component (PC1) contains more than
124 94% of the variance of the data and was therefore interpreted to best represent regional climate. The temporal
125 resolution of the climate proxy record in CB01, with 2,812 original measurements, has a calculated mean
126 spacing of ~16 yrs, ranging from ~4 yrs in the upper part of the core to almost 2 kyrs in the deeper part of the
127 core ([Foerster et al., 2012, 2014; Trauth et al., 2015](#)). The K record (following DTW alignment of cores CB01
128 to CB06 and PCA-based unmixing) runs from 45.358 to 0 kyr BP with a mean resolution of 8 yrs (ranging from
129 2.6 to 30.6 yrs) and the record has therefore been interpolated to an evenly-spaced time axis running from
130 45.358 to 0 kyr BP at 10 year intervals, which is close to the mean intervals of the original data (~16 yrs,
131 ranging from ~4 yrs to 2 kyrs) and in the aligned and unmixed data (~8 yrs, ranging from 2.6 to 30.6 yrs)
132 ([Trauth et al., 2018](#)).

133

134 *Principles of recurrence plots (RPs) and recurrence quantification analysis (RQA)*

135 We consider the Chew Bahir palaeolake to represent a complex system of interacting components, such as
136 the waterbody, the sediment beneath the lake, and the organisms living within and around the lake. Systems
137 with properties that change over time, such as the Chew Bahir palaeolake with its slowly changing
138 geomorphologic features, much more rapidly varying climatic factors, and the possible (very recent) influence
139 of human activities, are known as dynamical systems. The Chew Bahir multi-dimensional palaeolake system
140 is affected by a number of factors (known as *state variables* of the system) such as precipitation (with higher
141 rainfall resulting in increased weathering and erosion within the catchment, and hence more potassium washed
142 into the lake), evaporation (with increased evaporation producing more extreme hydrochemical conditions that
143 enhance potassium fixation in the sediment through authigenic clay-mineral alterations, [Foerster et al., 2012,](#)

2018) and wind speed (with higher wind speeds and reduced vegetation cover resulting in more potassium-rich particles being blown into the lake).

Analysis of temporal variations in the state variables of the Chew Bahir palaeolake requires a record of the variables influencing those state variables over a relevant time period. Since the state variables of the Chew Bahir system (e.g., precipitation, evaporation and wind speed) cannot be measured directly, we use indirect indicators (known as climate proxies) measured from natural archives of environmental change such as the sediments of the Chew Bahir basin, sampled by coring. A proxy record of a multi-dimensional dynamical system obtained by sampling a single variable is equivalent to projecting the dynamics of a complex system onto a single axis (Iwanski and Bradley, 1998). In our case the sampled variable is the series $x(t)$ of potassium concentrations x along a lake sediment core, which provides a natural archive of past influences on the Chew Bahir lake system, with sediment depth d converted into time t using the age model from Trauth et al. (2015).

One way to untangle the dynamics of a multi-dimensional system from a one-dimensional time series $x(t)$ is by time-delay embedding, which preserves the dynamic characteristics of the system (Packard et al., 1980) (Fig. 2). The reason why an entire system needs to be reconstructed from a single variable is that information about the system and the factors affecting its state variables is contained in a one-dimensional time series. In other words, since the potassium concentration in the Chew Bahir sediments is the result of a complex interplay between unknown amplitudes of different environmental (or state-) variables (such as precipitation, evaporation, and wind velocity), an analysis of the temporal variations in this environmental proxy will help us to understand the state variables of the Chew Bahir palaeolake and the time-varying interactions between its different components. The embedding of the time series $x(t)$ in a three-dimensional ($m=3$) coordinate system (a phase space), for example, means that three successive values $x(t)$, $x(t+\tau)$, and $x(t+2\tau)$ with a temporal separation of τ are represented by a single point within the phase space (Iwanski and Bradley, 1998; Webber and Zbilut, 2005; Marwan et al., 2007) (Fig. 2). The geometric representation of the embedded time series of observations as trajectories $\vec{s}(t)=[x(t), x(t+\tau), x(t+2\tau)]$ within the phase space is called a phase portrait. The

170 reconstructed phase space is not exactly the same as the original phase space, but its topological properties
 171 are preserved provided that the embedding dimension is sufficiently large (Packard et al., 1980, Takens, 1981).
 172

173 A common feature of dynamical systems is the property of recurrence (Webber and Zbilut, 2005). Recurring
 174 patterns in the state of a system are a reflection of typical characteristics of the dynamical system. Defining
 175 these patterns can contribute significantly to our understanding of the system's dynamics. Changes in
 176 environmental (or state-) variables (such as precipitation, evaporation, and wind velocity) often follow
 177 characteristic courses (represented as trajectories $\vec{s}$ in phase space) that could lead to similar (but not
 178 identical) lake characteristics (e.g. depth and area of the lake, alkalinity and salinity of the lake, species
 179 assemblage in the waterbody, or formation of authigenic minerals in the sediment). Recurrence plots (RPs),
 180 first introduced by Eckmann et al. (1987), are graphic displays of such recurring states within a system,
 181 calculated from the distance (e.g. the Euclidean distance) between all pairs of phase space vectors $\vec{s}(t_1)$ and
 182 $\vec{s}(t_2)$, below a threshold value e (Marwan et al., 2007) (Fig. 2+3).
 183

184 Measures of complexity were subsequently introduced to complement the visual inspection of recurrence plots,
 185 providing quantitative descriptions for use in recurrence quantification analysis (RQA) (e.g. Zbilut & Webber,
 186 1992; Marwan et al., 2007; Marwan, 2008). Among these, a selection of recurrence characteristics are very
 187 useful to summarize the appearance of the recurrence plots, ultimately helping to describe the behavior of the
 188 Chew Bahir lake system. An example of such a characteristic is the *recurrence rate* (RR), which is the density
 189 of black dots within the recurrence plot. This measure simply describes the probability of states of the system
 190 recurring within a particular time period.
 191

192 Diagonal lines typically occur in recurrence plots when one segment of the trajectory runs almost parallel to
 193 another segment (e.g. within a given tolerance) representing an earlier episode in the history of the system
 194 within the phase space. Diagonal lines in recurrence plots are therefore diagnostic of cyclicity in time series;
 195 such cyclicity detected using recurrence plots is not restricted to sinusoidal structures, in contrast to cyclicity

196 in Fourier-based time-series analysis. Cyclicity can be used to predict future conditions from present and past
 197 conditions using the ratio of the recurrence points that form diagonal patterns (of a certain length) to all other
 198 recurrence points, as a measure of the *determinism* (*DET*) or predictability of the system.
 199

200 Misleading results can be obtained during analysis of the diagonal and vertical lines in recurrence plots due to
 201 what is known as tangential motion (Marwan et al., 2007). Tangential motion occurs when calculating points
 202 of recurrence if there are points in the phase space below the threshold value ϵ that define the same trajectory
 203 segment, which is very typical of smooth or high resolution data (Marwan et al., 2007). This can lead to
 204 misleading RQA values, especially *DET* values. There are a number of methods that can be used to suppress
 205 this effect: (1) using a Theiler window defining the minimum time separation of nearest neighbors, (2) trailing
 206 data points on the phase space trajectory, or (3) by increasing the minimum length of the lines used to calculate
 207 the *DET* (Marwan et al., 2007). The latter method defines a lower limit for the length of the diagonal lines in
 208 order to exclude points on the same trajectory from *DET* calculations (Marwan et al., 2007; Marwan, 2011).
 209

210 RP analysis is further complicated by trends in the mean and variance (i.e. nonstationarity and long-term
 211 variations), which have a marked effect on the recurrence rate (*RR*), i.e. on the density of dots in RPs. If the
 212 long-scale variations are not of interest, the RQA characteristics can be used to reveal undesired fluctuations
 213 that do not reflect the more rapid variations in the dynamics of the system. We therefore use an adaptive
 214 threshold instead of a fixed value ϵ , which is determined in such a way that all points in the phase space have
 215 the same number of neighbors. The *RR* is thus corrected for any changes in the density of points within the
 216 phase space. The resulting constant *RRs* (and hence more homogeneous RPs) make it easier to detect rapid
 217 changes of the system dynamics at a particular point, while long-term changes can be observed at other points.
 218 Note that the *RR* obtained by using an adaptive threshold is constant with respect to the whole RP and also
 219 to each column within the RP, since one column represents one point in time, hence one point in the
 220 (reconstructed) phase space. This means, we expect a time-dependent behavior of *RR*, when performing a
 221 windowed analysis over the entire RP and a constant *RR* with respect to the whole RP (see description below).

222 One other possible problem in analyzing palaeoclimate records, which typically have very low signal-to-noise
223 ratios, is the disruption of patterns in the RP (such as diagonal lines) by noise, gaps and other disturbances.

224

225 The RQA can be carried out using moving windows in order to detect changes in the system dynamics,
226 represented by a change in the RQA measures, (1) by developing a single RP and calculating the RQA
227 measures for windows moving along this RP, or (2) by developing multiple RPs of individual windows and
228 calculating the RQA measures for these RPs. If nonstationarities (e.g. trends) are not the main focus of the
229 analysis, then approach (2) makes it easier to find transitions while ignoring any nonstationarities. However, if
230 the detection of overall changes is of interest (e.g. to test for nonstationarity), the recurrence conditions should
231 be kept constant over time (thus taking into account the RP of the entire time series) and approach (1) will be
232 more appropriate (Marwan, 2011). A window size needs to be chosen that is small enough to ensure good
233 temporal resolution but large enough to cover typical variations (e.g. the number of cycles) in order to be able
234 to detect recurrences. Since our data show a very dominant long-term trend, we first perform the RQA on the
235 original data and then on a high-pass filtered (and hence detrended) version of the data.

236

237 *Synthetic examples of a recurrence quantification analysis*

238 The investigation of synthetic data using RPs and the RQA measures described above, which are then used
239 to analyse real data, has proven to be particularly advantageous when the methods are complex and the
240 results not immediately obvious (e.g. Marwan et al., 2007; Trauth, 2015) (Fig. 3). The use of conceptual models
241 that mimic typical system changes helps us to understand the typical changes seen in RPs and to assign them
242 to one or other of those changes. The first example investigated was of normally-distributed (Gaussian) noise,
243 for which an RP and RQA measures were derived (Fig. 3A). The RP shows randomly distributed points, each
244 representing times when the system randomly returned to a similar state. Similar states frequently recur in
245 random noise but without any regularity except the states represented by the main diagonal line, the line of
246 identity. The *RR* is therefore more or less constant with very low values. Since there is no systematic pattern
247 (e.g. cyclically recurring states), the RP does not show any linear patterns and the *DET* is therefore very low.

248

249 The second example investigated was a composite signal of two sine waves, for which an RP and RQA
250 measures were again derived. In the interval between $-1,400$ and -600 there is a positive trend in the mean
251 (Fig. 3B). The RP shows long diagonal lines, diagnostic of cyclicity in time series, with shorter lines in between.
252 The horizontal distances between these lines corresponds to the periods of the two sine waves ($T_1=50$ and
253 $T_2=200$). Since the higher frequency is a harmonic of the lower frequency, the corresponding diagonal lines
254 appear thicker in the RP. The RP clearly shows the effect of the trend in the mean as the diagonal lines
255 disappear towards the upper left and lower right corners of the RP due to the trend-induced increase in the
256 distance, leaving a blocky pattern in the middle of the RP. Because the dynamic itself does not change, neither
257 do the DET values; instead they persist at their maximum values, which are unaffected by the trend. The RR
258 is however affected, as shown by the lower density of black dots in the RP, indicating cyclic variations with a
259 period of the same order as the dominant period in the signal ($T_2=200$).

260

261 The synthetic data in the third example comprised a sine wave and Gaussian noise with a signal-to-noise ratio
262 that decreases from left to right (Fig. 3C). As a result the continuity of the diagonal lines decreases to the right,
263 as do both the DET and the RR values. The synthetic data in the fourth example investigated comprised a
264 composite signal from two sine waves with distinct trends in their frequencies (Fig. 3D). The distances between
265 diagonal lines decrease as a result of increasing signal compression with time. The convergence of the lines
266 and their degree of curvature depend on the function describing the signal compression. The increase in
267 frequency with time results in reductions in the DET because the higher-frequency cycles seen on the right of
268 the plot are no longer adequately resolved. Recurrence points in between diagonal lines are scarce due to the
269 fact that the time delay τ chosen for the plot no longer suits the increased frequency. A higher sampling rate
270 eliminates this phenomenon.

271

272 The results of the fifth example revealed how RPs and RQAs respond to an abrupt transition from a composite
273 signal consisting of two sine waves ($T_1=300$, $T_2=50$) to a signal with only one sine wave ($T_3=60$) (Fig. 3E). As
274 before, the oscillations of the signal produce diagonal lines in the RP, with horizontal distances between the

lines corresponding to the periods of the signals. Two sets of diagonal lines in the lower-left corner of the RP correspond to the two frequencies of the sine waves, while there is only one set of diagonals in the upper right corner. During the transition we also let the amplitude of the signal decrease. Both the change from two sine waves to a single sine wave and the decrease in variance during the transition are clearly visible in the RP, as well as in the RQA measures. The single period oscillation has more recurrences than the two-period signal and therefore a higher *RR*, peaking at the transition because the lower signal variance produces a blocky pattern in the RP.

Because of the two different time scales of the two-period signal it has a more complicated phase-space trajectory on the left-hand side of the RP than the one-period signal. Trajectory segments are therefore only parallel at particular times, resulting in long diagonals for the long period signal (because *T2* is a harmonic of *T1*, as in the second example above), but interrupted diagonals (and lines that are curved at times) for the short period signal. This results in slightly lower *DET* values before the transition to a one-period signal than after it. An effect similar to that seen in our third example is observed in the interval between $t=-1200$ and $t=-1050$ (where the actual transition from a two period signal to a one period signal occurs), which is why the *DET* decreases here before again increasing due to the blocky pattern in the RP, which also causes the *RR* to increase, as mentioned previously.

The synthetic data in the sixth example investigated was of Gaussian noise with a stepwise transition in the mean and a change in the autocorrelation prior to this transition (Fig. 3F). It is important to note that neither the mean nor the variances change in the pre-transition section, so this change cannot be recognized using conventional methods. The two blocky features in the RP before and after the transition look very similar to those in the RP of the first example above. However, the interval between $-1,200$ and $-1,000$ clearly shows distinctive patterns within the RP that are different from those typically occurring in RPs of pure noise. White vertical lines help to define blocky features that represent episodes with different dynamics. The RQA characteristics look similar to those in the Gaussian noise example except for a section with higher autocorrelation, which is reflected in the higher density of black dots, clear diagonal lines, and a change in the

302 dynamics of the system. This fact can be used to detect such a change in autocorrelation as a precursor to a
303 tipping point and to ultimately predict the tipping point itself.

304 Results

305 We have used recurrence plots (RPs), complemented by a recurrence quantification analysis (RQA), to
306 characterize past climate change in the Chew Bahir basin over approximately the last 45 kyrs. We selected
307 the *RR* and *DET* measures because they describe fundamental properties of the complex Chew Bahir system
308 dynamics, while keeping the theoretical complexity within reasonable limits to facilitate interpretations. The RP
309 and RQA approach was applied to the record of K concentrations in the sediment cores (following DTW-based
310 alignment and linear unmixing using a PCA), which has previously been shown to be a reliable proxy for aridity
311 in the Chew Bahir basin (Foerster et al., 2012, 2018; Trauth et al., 2015, 2018). The K record was embedded
312 in phase spaces with dimensions varying from $m=5$ to $m=6$ and temporal distances varying from $\tau=3$ to $\tau=10$
313 data points, equivalent to 3×10 yrs=30 yrs and 10×10 yrs=100 yrs, where 10 yrs is the resolution of the time
314 series following interpolation. The RPs were calculated using a Euclidean norm, a Theiler window of $thei=1$
315 data point, a minimum length of $lmin=3$ for the lines used to calculate *DET*, and an adaptive threshold value e
316 rather than a fixed value. The size w and the step size ws of the moving window depend on the period of time
317 investigated and the number of data points contained therein.

318
319 In order to compare different climatic conditions we first looked at the RP of the complete, unfiltered time series
320 documenting the long-term variations in the Chew Bahir system (Fig. 4). This RP reveals a clear division of
321 the time series into sections of different lengths, indicated by square blocky features in the RP separated by
322 white vertical lines (see also Figs 3B, E and F). The first cluster of recurrence points occurs between 45.35
323 and 37 kyr BP, comprising both connected and isolated points. This interval is characterized by both vertical
324 and horizontal lines, representing episodes of stability (both wet and dry) interrupted by a series of extremely
325 wet events, indicated by white vertical lines in the RP. We observe low but gradually increasing *DET* values in
326 this episode, suggesting increasing predictability in the system.

327

328 A second, very obvious, cluster of recurrence points occurs between 37 and 20 kyr BP, which includes the
329 time intervals in which the Dansgaard-Oeschger (DO) cycles (~110–23 kyr BP), the Heinrich Events (HE, ~60–
330 12 kyr BP) and the Last Glacial Maximum (LGM, 23.5–18 kyr BP) affected the climate further to the north. The
331 RQA reveals consistently high *DET* values until about 15 kyr BP, exceeding those of the previously described
332 cluster and suggesting a much higher predictability in the system during that time. The recurrence plot for the
333 time interval from 20 to 0 kyr BP includes the African Humid Period (AHP, ~15–5 kyr BP). This interval is
334 characterized by three large clusters of recurrence points, interrupted by both white vertical and horizontal
335 lines, together with fluctuating *RR* values and a long-term trend towards lower *DET* values. The white vertical
336 lines again help to define blocky features marking episodes with different dynamics.

337

338 To analyze the dynamics of these individual sections, the time series was high-pass filtered with a cutoff
339 frequency of 0.001 yrs⁻¹ in order to remove any long-term trends (Fig. 5 to 7). An RP was constructed and an
340 RQA performed using a sliding window (*w*=100, *ws*=10) over the entire RP, assuming stationarity within the
341 intervals. The interval between 35 and 20 kyr BP is remarkable for numerous short, slightly curved diagonal
342 lines, suggesting a cyclic recurrence of wetter episodes in the Chew Bahir basin within a period that had a
343 generally dry climate (Fig. 5). The oscillating climatic conditions are reflected in higher *DET* values, while the
344 low *RR* values suggest a low probability of recurring states occurring within the system over a particular time
345 period (Marwan et al., 2007).

346

347 The recurrence plot for the time interval from 20 to 1 kyr BP includes the AHP (~15–5 kyr BP) (Fig. 6). As
348 before, using moving windows with a length of *w*=100 (1,000 yrs) and a step size of *ws*=10 (100 yrs) reveals
349 a series of blocky recurrence point patterns, interrupted by a series of white vertical lines. These patterns
350 suggest distinct episodes of relative stability, both wet and dry, separated by abrupt transitions at ~13.2 kyr
351 BP, ~11.8 kyr BP, ~7.5 kyr BP, ~5.2 kyr BP, and ~4.5 kyr BP. The overall appearance of the RP (and of the
352 time series itself) reflects dynamics characterized by a period of higher variability followed by a period of low
353 variability between 9.5 and 8 kyr BP, and dynamics dominated by a high-frequency cyclicity between 8 and 1

kyr BP, which is roughly similar to the fifth synthetic example above (see Fig. 3E). In addition to these patterns, the RP for the interval from 20 to 1 kyr BP also shows numerous short diagonal lines, suggesting a weak cyclicity. The diagonal lines, however, are very different from each other in width and in the number present. The *RR* values are very low except for an interval of relative stability between 10 and 8 kyr BP, which has a high probability of recurring states. The *DET* values document a general trend towards lower predictability in the system dynamics, but this decline exhibits a very complicated and somewhat cyclical pattern, rather than a simple linear trend.

In contrast to the RP for the first half of the time interval from 20 to 1 kyr BP, the RP for the period between 8 and 1 kyr BP contains numerous spotty diagonals following two blocky features at about 9 and 7 kyr BP (Fig. 7). These blocky features indicate a slowing-down of the system dynamics and therefore a higher predictability, as also indicated by high *DET* values (similar to the synthetic example shown in Figs 3E+F). The *RR* values are very low, except for one interval of relative stability. The distinctive diagonal lines after about 6.8 kyr BP suggest a marked cyclic recurrence of droughts approximately every 100–150 yrs, which lasted until stable conditions returned following the termination of the AHP. The relatively low *DET* values (~0.5) and therefore low predictability during this interval, however, reflects the discontinuity in the diagonal lines and the dispersion of cycles (i.e. the variability in distances between the diagonal lines, similar to the synthetic example shown in Figs 3C+D), suggesting that a stochastic process is superimposed on the cyclicity. The *DET* values during the wet-dry transition at the end of the AHP remain moderately high in the high-pass filtered time series until stable dry conditions are established. The interval between 4 and 2 kyr BP with very low *DET* values (<0.5) reflects a predominantly stochastic process.

Discussion

A section-by-section analysis of the RPs of the time series together with an examination of the temporal course of the RQA measures allows us to identify and eventually classify different types of variability and transitions (Table 1). The classification of variability and transitions can help to improve our understanding of the response

379 of the biosphere (including humans) to climate changes (e.g. [Donges et al., 2011](#); [Kennett et al., 2012](#); [Foerster](#)
380 [et al., 2015, 2016](#); [Trauth et al., 2015, 2018](#)). There is a general consensus amongst anthropologists that both
381 long-term trends and severe, abrupt changes resulted in significant changes to the social and socio-economic
382 behavior of early humans ([Gatto and Zerboni, 2015](#); [Clarke et al., 2016](#); [Lahr, 2016](#); [Flohr et al., 2016](#)). The
383 response to a changing habitat, both subtle or dramatic, accompanied by changes in essential resources such
384 as food and water, could be either expansion, decrease, migration, or adaption (e.g. [Foerster et al., 2015,](#)
385 [2016](#)). In this process, the nature of the adaptation is a function of the adaptability and the time scale (e.g.
386 [Foerster et al., 2015, 2016](#)). Of particular interest is how the adaptability of humans has enabled them to deal
387 with such diverse and profound environmental changes since 45 kyrs BP through behavioral changes, and
388 what level of environmental change met the limit of resilience (e.g. [Marshall and Hildebrandt, 2002](#); [Foerster](#)
389 [et al., 2015, 2016](#); [Trauth et al., 2015, 2018](#)). Furthermore, the different types of variability and transitions, and
390 the corresponding response of the biosphere (including humans), will help to detect similar types of changes
391 in the long (~280 m) sediment cores recently collected in the Chew Bahir basin within the Hominin Sites and
392 Paleolakes Drilling Project (HSPDP) and to investigate whether or not these types are typical for and exclusive
393 to the basin, as well as the response of the biosphere to these changes.

394
395 Our analysis clearly shows a number of different types of variability in the K record, separated by either gradual
396 or rapid transitions. The first type of variability occurs between the beginning of the record (45 kyr BP) and
397 about 35 kyr BP. Within this interval we observe a relatively low but gradually increasing predictability during
398 times of relative stability (both wet and dry), interrupted by a series of extremely wet events. There is no
399 cyclicity in this interval but rather an irregular pattern of different types of variability. Both the extreme events
400 and the rapid transitions between episodes of relative (wet and dry) stability will certainly have had an impact
401 on humans in the area, leaving them with a range of possible responses (adapt, migrate, starve) to a
402 dramatically changing environment (e.g. [Foerster et al., 2015, 2016](#); [Trauth et al., 2015, 2018](#)).

403
404 The second type of variability occurs between 35 and 20 kyr BP, with slightly different dynamics before and
405 after 25 kyr BP. This interval is characterized by a millennial-scale climate variations during the last glacial

406 cycle, which includes the time intervals in which the Dansgaard-Oeschger (DO) cycles (~110–23 kyr BP), the
407 Heinrich Events (HE, ~60–12 kyr BP) and the Last Glacial Maximum (LGM, 23.5–18 kyr BP) affected the
408 climate further to the north. This millennial-scale climate fluctuations are cyclic with minor variations in the
409 period, as indicated by the slight curvature of the diagonals in the RP (see synthetic example shown in [Fig.](#)
410 [3D](#)), although this curvature could also suggest inaccuracies in the age model, rather than real variations in
411 the cyclicity. The pronounced cyclicity is reflected in the RP and in the RQA measures by a very high
412 predictability, but on time scales that are certainly not relevant to humans, because it is orders of magnitude
413 longer than the lifetime of individual humans. However, the transitions between the dry and wet episodes were
414 very rapid which has probably caused significant stress to human populations.

415
416 The third type of variability occurs between 16 and 10 kyr BP, including parts of the AHP (~15–5 kyr BP).
417 During this interval we find episodes of relative stability, both wet and dry, separated by abrupt transitions at
418 ~13.2 kyr BP and ~11.8 kyr BP. There is also evidence of a weak cyclicity with a general (but complicated)
419 trend towards lower predictability. The onset of the AHP in the Chew Bahir area was relatively rapid (covering
420 ~240 yrs, [Trauth et al., 2018](#)), which is in agreement with similar records from elsewhere, as a result of large-
421 scale deglacial forcings (i.e. changes of Atlantic sea-surface temperatures causing meridional shifts of the
422 African easterly jet and the monsoon belt, and changes in the atmospheric concentration of greenhouse gases
423 causing changes in atmospheric temperatures, [Shanahan et al., 2015](#)). The onset and termination of the dry
424 episode during the Younger Dryas (YD) were also rapid transitions (over less than 100 yrs, [Trauth et al., 2018](#)),
425 again very similar to other sites in N and NE Africa (e.g. [Shanahan et al., 2015](#); [Trauth et al., 2018](#)).

426
427 The climate variability within the AHP and the long-term transition that it represents clearly affected human
428 communities and has fueled massive changes in the population size and structure such as the profound socio-
429 economic transformations that have been documented for N and NE Africa (e.g. [Marshall and Hildebrand,](#)
430 [2002](#); [Brooks, 2006](#); [Clarke et al., 2016](#); [Lahr, 2016](#); [Marchant et al., 2018](#)). A well-studied example is provided
431 by demographic reconstructions that have been made for the Saharan Holocene. These reconstructions show
432 a temporal delay between the onset of humid conditions (based on sedimentary dust flux records) and human

433 reoccupation of former desert areas, with associated societal changes seen as a response to the
434 environmental changes ([Manning and Timpson, 2014](#); [Gatto and Zerboni, 2015](#); [Clarke et al., 2016](#)).

435

436 Having adapted to the wet climate of the AHP, humans certainly had to cope with the very rapid transition
437 towards extreme dryness at the onset of the YD dry episode. Highly mobile groups of hunter-gatherers
438 responded to short-term arid pulses by vertical migration as documented by the settlement patterns in what
439 are assumed to have been refuge areas such as the SW Ethiopian Highlands (e.g. [Foerster et al., 2015](#)). The
440 YD is followed by a short interval of relative stability between 10 and 8 kyr BP, followed by a dry episode
441 centered around 7.5 kyr BP. This event, which is synchronous with the prolonged pause in the Green Sahara
442 conditions 8 kyrs ago (within the uncertainty of our age model, see [Trauth et al., 2015](#)), coincides with a
443 temporary abandonment of sites previously occupied by Neolithic humans ([Tierney et al., 2017](#)).

444

445 The fourth type of variability occurs between 8 and 4 kyr BP. This interval includes the transition from the
446 humid climate of the AHP to the subsequent dry climate. The termination of the AHP was a result of weaker,
447 insolation-driven forcing (and hence more complex and time-transgressive responses) than that which
448 produced the DO cycles, the onset of the AHP, or the onset/termination of the YD dry episode ([Shanahan et al., 2015](#)). This was nevertheless compared to those produced by more subtle changes in orbital forcing, but
449 certainly not abrupt compared to human time scales, i.e. human lifespan or a little more, as it continued for
450 approximately 990 years ([Trauth et al., 2018](#)). The termination of the AHP occurred at different times in other
451 areas, suggesting a strong influence of Indian Ocean SSTs on the East African climate ([Shanahan et al., 2015](#)). Most of the transition at the end of the AHP is characterized by wet conditions, interrupted by at ~14
452 dry events that have recurred every 160 ± 40 years and lasted 20–80 years ([Trauth et al., 2015](#)). Compared to
453 the low-frequency cyclicity of climate variability during the DO cycles, this type of cyclicity occurs on time scales
454 equivalent to a few human generations. In other words, it is very likely (albeit speculative) that people were
455 conscious of these changes and adapted their lifestyles to the consequent changes in water and food
456 availability ([Marshall and Hildebrand, 2002](#); [Clarke et al., 2016](#)).

459

460 An interesting aspect of this variability is the nature of the transitions close to the threshold in the system
461 response, and how the environment switches from one stable mode to another. A rapid change of climate in
462 response to a relatively modest change in forcing appears to be typical of tipping points in complex systems
463 such as the Chew Bahir basin ([Lenton et al., 2008](#); [Ditlevsen and Johnsen, 2010](#)). If this is the case then the
464 14 dry events at the end of the AHP could represent precursors of an imminent tipping point that would have
465 allowed a prediction of climate change in the Chew Bahir basin at that time. A deeper analysis of our data is
466 however required to understand whether the wet-dry climate transition in the area was due to a saddle-node
467 bifurcation in the structural stability of the climate, or whether it was induced by a stochastic fluctuation ([Lenton
468 et al., 2008](#); [Ditlevsen and Johnsen, 2010](#)). The time interval after the termination of the AHP (<4 kyr BP) leads
469 into present-day conditions in the Chew Bahir basin.

470 Conclusions

471 We have used a recurrence quantification analysis (RQA) on environmental records from short cores collected
472 during a pilot study within the Chew Bahir basin to characterize the Chew Bahir palaeolake as a dynamical
473 system composed of interacting components. The different types of variability and transitions in these records
474 were classified to improve our understanding of the response of the biosphere to climate change, and
475 especially the response of humans in the area. This classification and the corresponding responses of the
476 biosphere will enable the detection of similar types of variability and transitions in the long (~280 m) ICDP core
477 collected in the Chew Bahir basin within the Hominin Sites and Paleolakes Drilling Project (HSPDP) and allow
478 us to investigate whether or not these types are typical for and exclusive to the basin.

479 References

- 480 Ambrose SH (1998) Late Pleistocene human population bottlenecks, volcanic winter, and the differentiation of
481 modern humans. *Journal of Human Evolution* 35:115-118. <https://doi.org/10.1006/jhev.1998.0219>
482 Brandt SA, Fisher EC, Hildebrand EA, Vogelsang R, Ambrose SH, Lesur J, Wang H (2012) Early MIS 3
483 occupation of Mochena Borago Rockshelter, Southwest Ethiopian Highlands: Implications for Late

484 Pleistocene archaeology, paleoenvironments and modern human dispersals. *Quaternary International*
485 274:38-54. <https://doi.org/10.1016/j.quaint.2012.03.047>

486 Brooks N (2006) Cultural responses to aridity in the Middle Holocene and increased social complexity.
487 *Quaternary International* 151:29-49. <https://doi.org/10.1016/j.quaint.2006.01.013>

488 Brown ET, Johnson TC, Scholz CA, Cohen AS, King JW (2007) Abrupt change in tropical African climate
489 linked to the bipolar seesaw over the past 55,000 years. *Geophysical Research Letters* 34:L20702.
490 <https://doi.org/10.1029/2007GL031240>

491 Builes-Jaramillo A, Marwan N, Poveda G, Kurths J (2018) Nonlinear interactions between the Amazon River
492 basin and the Tropical North Atlantic at interannual timescales. *Climate Dynamics* 50:2951-2969.
493 <https://doi.org/10.1007/s00382-017-3785-8>

494

495

496 Campisano CJ, Cohen AS, Arrowsmith JR, Asrat A, Behrensmeyer AK, Brown ET, Deino AL, Deocampo DM,
497 Feibel CS, Kingston JD, Lamb HF, Lowenstein TK, Noren A, Olago DO, Owen RB, Pelletier JD, Potts R,
498 Reed KE, Renaut RW, Russell JM, Russell JL, Schäbitz F, Stone JR, Trauth MH, Wynn JG (2017) The
499 Hominin Sites and Paleolakes Drilling Project: high-resolution paleoclimate records from the East African
500 Rift System and their implications for understanding the environmental context of hominin evolution.
501 *PaleoAnthropology* 2017:1-43. <https://doi.org/10.4207/PA.2017.ART104>

502 Carto SL, Weaver AJ, Hetherington R, Lam Y, Wiebe EC (2009) Out of Africa and into an ice age: On the role
503 of global climate change in the late Pleistocene expansion of early modern humans out of Africa. *Journal*
504 *of Human Evolution* 56:139-151. <https://doi.org/10.1016/j.jhevol.2008.09.004>

505 Castañeda IS, Mulitza S, Schefuß E, Santos RAL, Damste JSS, Schouten S (2009) Wet phases in the
506 Sahara/Sahel region and human expansion patterns in North Africa. *Proceedings of the National Academy*
507 *of Sciences* 106:1-5. <https://doi.org/10.1073/pnas.0905771106>

508 Cheung WH, Senay GB, Singh A (2008) Trends and spatial distribution of annual and seasonal rainfall in
509 Ethiopia. *International Journal of Climatology* 28:1723-1734. <https://doi.org/10.1002/joc.1623>

510 Clark, J, Brooks N, Banning EB, Bar-Matthews M, Campbell S, Clare L, Cremaschi M, di Lernia S, Drake N,
 511 Gallinaro M, Manning S, Nicoll K, Philip G, Rosen S, Schoop UD, Tafuri MA, Weninger B, Zerboni A (2016)
 512 Climatic changes and social transformations in the Near East and North Africa during the 'long' 4th
 513 millennium BC: A comparative study of environmental and archaeological evidence. *Quaternary Science*
 514 *Reviews* 136:96-121. <https://doi.org/10.1016/j.quascirev.2015.10.003>
 515 Cohen A, Campisano C, Arrowsmith R, Asrat A, Behrensmeyer AK, Deino A, Feibel C, Hill A, Johnson R,
 516 Kingson J, Lamb H, Lowenstein T, Noren A, Olago D, Owen RB, Potts R, Reed K, Renaut R, Schäbitz F,
 517 Tiercelin JJ, Trauth MH, Wynn J, Ivory S, Brady K, O'Grady R, Rodysill J, Githiri J, Russell J, Foerster V,
 518 Dommain R, Rucina S, Deocampo D, Russell J, Billingsley A, Beck C, Dorenbeck G, Dullo L, Feary D,
 519 Garelo D, Gromig R, Johnson T, Junginger A, Karanja M, Kimburi E, Mbuthia A, McCartney T, McNulty E,
 520 Muiruri V, Nambiro E, Negash EW, Njagi D, Wilson JN, Rabideaux N, Raub T, Sier MJ, Smith P, Urban J,
 521 Warren M, Yadeta M, Yost C, Zinaye B (2016) The Hominin Sites and Paleolakes Drilling Project: Inferring
 522 the Environmental Context of Human Evolution from Eastern African Rift Lake Deposits. *Scientific Drilling*
 523 21:1-16. <https://doi.org/10.5194/sd-21-1-2016>
 524 Davidson A (1983) The Omo River project: reconnaissance geology and geochemistry of parts of Ilubabor,
 525 Kefa, Gemu Gofa and Sidamo. *Ethiopian Institute of Geological Surveys Bulletin* 2:1-89.
 526 deMenocal P, Ortiz J, Guilderson T, Adkins J, Sarnthein M, Baker L, Yarusinsky M (2000) Abrupt onset and
 527 termination of the African Humid Period: rapid climate responses to gradual insolation forcing. *Quaternary*
 528 *Science Reviews* 19:347-361. [https://doi.org/10.1016/S0277-3791\(99\)00081-5](https://doi.org/10.1016/S0277-3791(99)00081-5)
 529 Ditlevsen PD, Johnsen SJ (2010) Tipping points: Early warning and wishful thinking. *Geophysical Research*
 530 *Letters* 37:L19703. <https://doi.org/10.1029/2010GL044486>
 531 Donges JF, Donner RV, Trauth MH, Marwan N, Schellnhuber HJ, Kurths J (2011) Nonlinear detection of
 532 paleoclimate-variability transitions possibly related to human evolution. *Proceedings of the National*
 533 *Academy of Sciences* 108:20423-20427. <https://doi.org/10.1073/pnas.1117052108>
 534 Donner RV, Heitzig J, Donges JF, Zou Y, Marwan N, Kurths J (2011) The Geometry of Chaotic Dynamics - A
 535 Complex Network Perspective, *European Physical Journal B* 84:653-672.
 536 <https://doi.org/10.1140/epjb/e2011-10899-1>

537 Eckmann JP, Kamphorst SO, Ruelle D (1987) Recurrence Plots of Dynamical Systems. *Europhysics Letters*,
538 5, 973-977. <https://doi.org/10.1209/0295-5075/4/9/004>

539 Egerer S, Claussen M, Reick C, Stanelle T (2017) Could gradual changes in Holocene Saharan landscape
540 have caused the observed abrupt shift in North Atlantic dust deposition? *Earth and Planetary Science*
541 *Letters* 473:104-112. <https://doi.org/10.1016/j.epsl.2017.06.010>

542 Eroglu D, McRobie FH, Ozken I, Stemler T, Wyrwoll KH, Breitenbach SFM, Marwan N, Kurths J (2016) See-
543 saw relationship of the Holocene East Asian-Australian summer monsoon. *Nature Communications* 7.
544 <https://doi.org/10.1038/ncomms12929>

545 Feldhoff JH, Donner RV, Donges JF, Marwan N, Kurths J (2013) Geometric signature of complex
546 synchronisation scenarios. *Europhysics Letters* 102:30007. <https://doi.org/10.1209/0295-5075/102/30007>

547 Flohr P, Fleitmann D, Matthews R, Matthews W, Black S (2016) Evidence of resilience to past climate change
548 in Southwest Asia: Early farming communities and the 9.2 and 8.2 ka events. *Quaternary Science Reviews*
549 136:23-39. <https://doi.org/10.1016/j.quascirev.2015.06.022>

550 Foerster V, Junginger A, Langkamp O, Gebru T, Asrat A, Umer M, Lamb H, Wennrich V, Rethemeyer J,
551 Nowaczyk N, Trauth MH, Schäbitz F (2012) Climatic change recorded in the sediments of the Chew Bahir
552 basin, southern Ethiopia, during the last 45,000 yr. *Quaternary International* 274:25-37.
553 <https://doi.org/10.1016/j.quaint.2012.06.028>

554 Foerster V, Vogelsang R, Junginger A, Asrat A, Lamb HF, Schaebitz F, Trauth MH (2015) Environmental
555 Change and Human Occupation of Southern Ethiopia and Northern Kenya during the last 20,000
556 yr. *Quaternary Science Reviews* 129:333-340. <https://doi.org/10.1016/j.quascirev.2015.10.026>

557 Foerster V, Deocampo DM, Asrat A, Günter C, Junginger A, Kraemer H, Stroncik NA, Trauth MH (2018)
558 Towards an understanding of climate proxy formation in the Chew Bahir basin, southern Ethiopian Rift.
559 *Palaeogeography, Palaeoclimatology, Palaeoecology* 501:111-123.
560 <https://doi.org/10.1016/j.palaeo.2018.04.009>

561 Fritsch FN, Carlson RE (1980) Monotone Piecewise Cubic Interpolation. *SIAM J. Numerical Analysis* 17:238-
562 246. <https://doi.org/10.1137/0717021>

563 Garcin Y (2008) Comment on "Abrupt change in tropical African climate linked to the bipolar seesaw over the
 564 past 55,000 yr" by Brown, E. T, Johnson, T. C, Scholz, C. A, Cohen, A. S, and King, J. W. (2007).
 565 Geophysical Research Letters 35:L04701. <https://doi.org/10.1029/2007GL032399>
 566 Garcin Y, Deschamps P, Ménot G, de Saulieu G, Schefuß E, Sebag D, Dupont LM, Oslisly R, Brademann B,
 567 Mbusnum KG, Onana JM, Ako AA, Epp LS, Tjallingii R, Strecker MR, Brauer A, Sachse D (2018) Early
 568 anthropogenic impact on Western Central African rainforests 2,600 y ago. Proceedings of the National
 569 Academy of Sciences 115:3261-3266. <https://doi.org/10.1073/pnas.1715336115>
 570 Gasse F (2000) Hydrological changes in the African tropics since the Last Glacial Maximum. Quaternary
 571 Science Reviews 19:189-211. [https://doi.org/10.1016/S0277-3791\(99\)00061-X](https://doi.org/10.1016/S0277-3791(99)00061-X)
 572 Gatto MC, Zerboni A (2015) Holocene Supra-Regional Environmental Changes as Trigger for Major Socio-
 573 Cultural Processes in Northeastern Africa and the Sahara. African Archaeological Review 32:301-333.
 574 <https://doi.org/10.1007/s10437-015-9191-x>
 575 Goswami B, Marwan N, Feulner G, Kurths J (2013) How do global temperature drivers influence each other?
 576 A network perspective using recurrences. European Physical Journal, Special Topics 222:861-873.
 577 <https://doi.org/10.1140/epjst/e2013-01889-8>
 578 Hildebrand E, Grillo K (2012) Early herders and monumental sites in eastern Africa: New radiocarbon dates,
 579 Antiquity 86:338-352. <https://doi.org/10.1017/S0003598X00062803>
 580 Ivory SJ, Russell J (2017) Lowland forest collapse and early human impacts at the end of the African Humid
 581 Period at Lake Edward, equatorial East Africa. Quaternary Research 89:7-20.
 582 <https://doi.org/10.1017/qua.2017.48>
 583 Iwanski J, Bradley E (1998) Recurrence plot analysis: To embed or not to embed? Chaos 8:861-871.
 584 <https://doi.org/10.1063/1.166372>
 585 Junginger A, Roller S, Trauth MH (2014) The effect of solar irradiation changes on water levels in the paleo-
 586 Lake Suguta, Northern Kenya Rift, during the late Pleistocene African Humid Period (15-5 ka
 587 BP). Palaeogeography, Palaeoclimatology, Palaeoecology 396:1-16.
 588 <https://doi.org/10.1016/j.gloplacha.2013.09.005>

589 Kantz H, Schreiber T (1997) Nonlinear Time Series Analysis. Cambridge University Press, Cambridge.
590 <https://doi.org/10.1017/CBO9780511755798>

591 Klein RG (1995) Anatomy, behavior, and modern human origins. *Journal of World Prehistory* 9:167-198.
592 <https://doi.org/10.1007/BF02221838>

593 Klein RG, Steele TE (2013) Archaeological shellfish size and later human evolution in Africa. *Proceedings of*
594 *the National Academy of Sciences* 110:10910-10915. <https://doi.org/10.1073/pnas.1304750110>

595 Lahr M (2016) The shaping of human diversity: filters, boundaries and transitions. *Philosophical Transactions*
596 *of the Royal Society B* 371:20150241. <https://doi.org/10.1098/rstb.2015.0241>

597 Lamb HF, Bates CR, Bryant CL, Davies SJ, Huws DG, Marshall MH, Roberts HM (2018) 150,000-year
598 palaeoclimate record from northern Ethiopia supports early, multiple dispersals of modern humans from
599 Africa. *Scientific Reports* 8:1077. <https://doi.org/10.1038/s41598-018-19601-w>

600 Lenton TM, Held H, Kriegler E, Hall JW, Lucht W, Rahmstorf S, Schellnhuber HJ (2008) Tipping elements in
601 the Earth's climate system. *Proceedings of the National Academy of Sciences* 105:1786-1796.
602 <https://doi.org/10.1073/pnas.0705414105>

603 Maley J, Doumenge C, Giresse P, Mahé G, Philippon N, Hubau W, Lokonda MO, Tshibamba JM, Chepstow-
604 Lusty A (2018) Late Holocene forest contraction and fragmentation in central Africa. *Quaternary Research*
605 89:43-59. <https://doi.org/10.1017/qua.2017.97>

606 Manning K, Timpson A (2014) The demographic response to Holocene climate change in the Sahara.
607 *Quaternary Science Reviews* 101:28-35. <https://doi.org/10.1016/j.quascirev.2014.07.003>

608 Maraun D, Rust HW, Timmer J (2004) Tempting long-memory - on the interpretation of DFA results. *Nonlinear*
609 *Processes in Geophysics*. <https://doi.org/10.5194/npg-11-495-2004>

610 Marchant R, Richer S, Boles O, Capitani C, Courtney-Mustaphi CJ, Lane PJ et al. (2018) Drivers and
611 trajectories of land cover change in East Africa: Human and environmental interactions from 6000 years
612 ago to present. *Earth-Science Reviews* 178:322-378. <https://doi.org/10.1016/j.earscirev.2017.12.010>

613 Marshall F, Hildebrand E (2002) Cattle before the crops: the beginnings of food production in Africa. *Journal*
614 *of World Prehistory* 16:99-143. <https://doi.org/10.1023/A:10199549>

615 Marwan N, Trauth MH, Vuille M, Kurths J (2003) Comparing modern and Pleistocene ENSO-like influences in
 616 NW Argentina using nonlinear time series analysis methods. *Climate Dynamics* 21:317-326.
 617 <https://doi.org/10.1007/s00382-003-0335-3>

618 Marwan N, Kurths J (2005) Line structures in recurrence plots. *Physics Letters A* 336:349-357.
 619 <https://doi.org/10.1016/j.physleta.2004.12.056>

620 Marwan N, Romano MC, Thiel M, Kurths J (2007) Recurrence Plots for the Analysis of Complex Systems.
 621 *Physics Reports* 438:237-329. <https://doi.org/10.1016/j.physrep.2006.11.001>

622 Marwan N (2008) A Historical Review of Recurrence Plots. *European Physical Journal, Special Topics* 164:3-
 623 12. <https://doi.org/10.1140/epjst/e2008-00829-1>

624 Marwan N (2011) How to avoid potential pitfalls in recurrence plot based data analysis. *Journal of Biofurcation*
 625 *and Chaos* 21:1003-1017. <https://doi.org/10.1142/S0218127411029008>

626 Marwan N, Schinkel S, Kurths J (2013) Recurrence plots 25 years later - Gaining confidence in dynamical
 627 transitions. *Europhysics Letters* 101:20007. <https://doi.org/10.1209/0295-5075/101/20007>

628 Marwan N, Kurths J (2015) Complex network based techniques to identify extreme events and (sudden)
 629 transitions in spatio-temporal systems. *Chaos* 25:097609. <https://doi.org/10.1063/1.4916924>

630 Maslin MA (2004) Ecological verses climatic thresholds. *Science* 306:2197-2198.
 631 <https://doi.org/10.1126/science.1107481>

632 Moore JM, Davidson A (1978) Rift structure in southern Ethiopia. *Tectonophysics* 46:159-173.
 633 [https://doi.org/10.1016/0040-1951\(78\)90111-7](https://doi.org/10.1016/0040-1951(78)90111-7)

634 Mudelsee M (2014) *Climate Time Series Analysis: Classical Statistical and Bootstrap Methods*, Second
 635 Edition. Springer. <https://doi.org/10.1007/978-3-319-04450-7>

636 Nicholson SE (2017) Climate and climatic variability of rainfall over eastern Africa. *Reviews of Geophysics*
 637 55:590-635. <https://doi.org/10.1002/2016RG000544>

638 Packard NH, Crutchfield JP, Farmer JD, Shaw RS (1980) Geometry from a time series. *Physical Review*
 639 *Letters* 45:712-716. <https://doi.org/10.1103/PhysRevLett.45.712>

640 de Ramos AMT, Zou Y, de Oliveira, GS, Kurths J, Macau EEN (2018) Unveiling non-stationary coupling
641 between Amazon and ocean during recent extreme events. *Climate Dynamics* 50:767-776.
642 <https://doi.org/10.1007/s00382-017-3640-y>

643 Renfrew C (2009) Situating the creative explosion: universal or local? In: Renfrew, C, Morley, I. (Eds.),
644 *Becoming Human: Innovation in Prehistoric Material and Spiritual Culture*. Cambridge University Press,
645 Cambridge, pp. 74-92. ISBN-13: 978-0521734660

646 Richter J, Hauck T, Vogersang R, Widlock T, Le Tensorer JM, Schmid P (2012) "Contextual areas" of early
647 *Homo sapiens* and their significance for human dispersal from Africa into Eurasia between 200 ka and 70
648 ka. *Quaternary International* 274:5-24. <https://doi.org/10.1016/j.quaint.2012.04.017>

649 Rodó X, Rodríguez-Arias MA (2006) A new method to detect transitory signatures and local time/space
650 variability structures in the climate system: the scale-dependent correlation analysis. *Climate Dynamics*
651 27:441-458. <https://doi.org/10.1007/s00382-005-0106-4>

652 Runge J, Heitzig J, Marwan N, Kurths J (2012) Quantifying causal coupling strength: A lag-specific measure
653 for multivariate time series related to transfer entropy. *Physical Review E* 86:061121.
654 <https://doi.org/10.1103/PhysRevE.86.061121>

655 Runge J, Petoukhov V, Kurths J (2014) Quantifying the strength and delay of climatic interactions: The
656 ambiguities of cross correlation and a novel measure based on graphic models. *Journal of Climate*, 27,
657 720-739. <https://doi.org/10.1175/JCLI-D-13-00159.1>

658 Saji NH, Goswami BN, Vinayachandran PN, Yamagata T (1999) A dipole mode in the tropical Indian Ocean.
659 *Nature* 401:360-363. <https://doi.org/10.1038/43854>

660 Segele ZT, Lamb PJ, Leslie LM (2009) Seasonal-to-Interannual Variability of Ethiopia/Horn of Africa Monsoon.
661 Part I: Associations of Wavelet-Filtered Large-Scale Atmospheric Circulation and Global Sea Surface
662 Temperature. *Journal of Climate* 22:3396-3421. <https://doi.org/10.1175/2008JCLI2859.1>

663 Seleshi Y, Zanke U (2004) Recent changes in rainfall and rainy days in Ethiopia. *International Journal of*
664 *Climatology* 24:973-983. <https://doi.org/10.1002/joc.1052>

665 Shakun JD, Carlson AE (2010) A global perspective on Last Glacial Maximum to Holocene climate change.
666 *Quaternary Science Reviews* 29:1801-1816. <https://doi.org/10.1016/j.quascirev.2010.03.016>

667 Shanahan TM, McKay NP, Hughen KA, Overpeck JT, Otto-Bliesner B, Heil CW, King J, Scholz CA, Peck J
 668 (2015) The time-transgressive termination of the African Humid Period. *Nature Geoscience* 8:140-144.
 669 <https://doi.org/10.1038/NGEO2329>

670 Takens F (1981) Detecting strange attractors in turbulence. *Lecture Notes in Mathematics* 898, Springer, Berlin
 671 Heidelberg New York, pp. 366-381. <https://doi.org/10.1007/BFb0091924>

672 Tierney JE, deMenocal PB (2013) Abrupt shifts in Horn of Africa hydroclimate since the last glacial maximum.
 673 *Science* 342:843-846. <https://doi.org/10.1126/science.1240411>

674 Tierney JE, Pausata FSR, deMenocal PB (2017) Rainfall regimes of the Green Sahara. *Science Advances*, 3,
 675 e1601503. <https://doi.org/10.1126/sciadv.1601503>

676 Timmerman A, Friedrich T (2016) Late Pleistocene climate drivers of early human migration. *Nature* 538:92-
 677 95. <http://dx.doi.org/10.1038/nature19365>

678 Tsonis AA (2017) *Advances in Nonlinear Geosciences*. Springer, 707 p, ISBN: 978-3-319-58894-0

679 Trauth MH, Maslin MA, Deino A, Strecker MR, Bergner AGN, Dühnforth M (2007) High- and low-latitude forcing
 680 of Plio-Pleistocene African climate and human evolution. *Journal of Human Evolution* 53:475-486.
 681 <https://doi.org/10.1016/j.jhevol.2006.12.009>

682 Trauth MH (2015) *MATLAB Recipes for Earth Sciences - 4th Edition*. Springer, 427 p, ISBN: 978-3-662-46244-
 683 7

684 Trauth MH, Bergner AGN, Foerster V, Junginger A, Maslin MA, Schaebitz F (2015) Episodes of environmental
 685 stability and instability in late Cenozoic lake records of Eastern Africa. *Journal of Human Evolution* 87:21-
 686 31. <https://doi.org/10.1016/j.jhevol.2015.03.011>

687 Trauth MH, Foerster V, Junginger A, Asrat A, Lamb HF, Schaebitz F (2018) Abrupt or gradual? Change point
 688 analysis of the late Pleistocene-Holocene climate record from Chew Bahir, southern Ethiopia. *Quaternary*
 689 *Research* 90:321-330. <https://doi.org/10.1017/qua.2018.30>

690 Turcotte DL (1992) *Fractals and Chaos in Geology and Geophysics*. Cambridge University Press, Cambridge.
 691 <https://doi.org/10.1002/gj.3350280216>

692 Vogelsang R, Keding B (2013) Climate, culture, and change: From hunters to herders in northeastern and
 693 southwestern Africa, in: *Comparative Archaeology and Paleoclimatology - Socio-cultural responses to a*

694 changing world, Baldia MO, Perttula TK, Frink DS, eds.), BAR International Series, 2456, Archaeopress,
695 Oxford, UK, pp. 43-62.

696 Webber CL, Zbilut JP (2005) Recurrence Quantification Analysis of Nonlinear Dynamical Systems, in: from:
697 Riley, M.A, Van Orden, G.C, Tutorials in contemporary nonlinear methods for the behavioral sciences,
698 available online at <https://www.nsf.gov/sbe/bcs/pac/nmbs/nmbs.jsp>. [https://doi.org/10.1007/978-3-319-](https://doi.org/10.1007/978-3-319-07155-8)
699 07155-8

700 Widlok T, Aufgebauer A, Bradtmöller M, Dikau R, Hoffmann T, Kretschmer I, Panagiotopoulos K, Pastoors A,
701 Peters R., Schäbitz F, Schlummer M, Solich M, Wagner B, Weniger GC, Zimmermann A (2012) Towards
702 a theoretical framework for analyzing integrated socio-environmental systems. Quaternary International
703 274:259-272. <https://doi.org/10.1016/j.quaint.2012.01.020>

704 Zbilut JP, Webber Jr CL (1992) Embeddings and delays as derived from quantification of recurrence plots.
705 Physics Letters A 171:199-203. [https://doi.org/10.1016/0375-9601\(92\)90426-M](https://doi.org/10.1016/0375-9601(92)90426-M)

706

707 **Table 1** Compilation of the most important time periods in the Chew Bahir sediment cores, classified according
708 to main climate, environmental conditions, recurrence plot appearance, recurrence rate RR, determinism DET,
709 occurrence of extreme events and human habitat.

Age (kyr)	45–37	37–20	20–16	16–10	10–8	8–4	4–now
Main climate	intermediate	dry	intermediate	wet, with YD dry event	wet	wet	dry
Environmental conditions	wet-dry trend	1500 yrs cycles	Long-term sinusoidal dry-wet-dry trend due to 20 kyr precession cycle			stable with 160 yrs cycles	stable, wet at the end
RP appearance	irregular	regular	regular	irregular	regular	regular	regular
Recurrence rate (RR)	low	intermediate	intermediate	high, with low YD event	high	high	intermediate
Determinism (DET)	low	high	high	intermediate	intermediate	intermediate	low
Extreme events	random wet	wet-dry	none	YD dry event	possibly 8.2 kyr dry event	~14 dry events, 20–80 yrs long	very arid, possibly wet after 2 kyr BP
Human habitat	extreme random wet events	wet-dry cycles	stable	threshold	stable, except for 8.2 kyr event	threshold and cycles	aridity

710

711

Figure 1 (A) Topographic map of the Chew Bahir basin, showing the outline of the catchment, the drainage network, the locations of the short cores in the pilot study (2009, 2010), and the 2014 HSPDP-CHB drill site. (B) Geologic map of the Chew Bahir basin, showing the three generalized rock types: Cenozoic rift sediments, Cenozoic rift volcanics, and Proterozoic basement. Compilation based on Omo River Project Map ([Davidson, 1983](#)), Geology of the Sabarei Area ([Key, 1987](#)), Geology of the Yabello Area ([Hailemeskel and Fekadu, 2004](#)), and Geology of the Agere Maryam Area ([Hassen et al., 1997](#)). Maps are modified versions of the ones previously published in [Trauth et al. \(2018\)](#) and [Foerster et al. \(2018\)](#).

Figure 2 Principle of recurrence plot. To untangle the dynamics of a multi-dimensional system from a one-dimensional time series $x(t)$ (black line, in panel A) is by time-delay embedding. The embedding of the time series $x(t)$ in a three-dimensional ($m=3$) coordinate system (a phase space, shown in panel B), for example, means that three successive values $x(t)$, $x(t+\tau)$, and $x(t+2\tau)$ with a temporal separation of τ are represented as a single point $\vec{s}(t)$ within the phase space (Iwanski and Bradley, 1998; Webber and Zbilut, 2005; Marwan et al., 2007). Recurrence plots (RPs, panel D), first introduced by Eckmann et al. (1987), are graphic displays of such recurring states within a system, calculated from the distance (e.g. the Euclidean distance d , shown in panel C) between all pairs of phase space vectors $\vec{s}(t_1)$ and $\vec{s}(t_2)$, below a threshold value e (also shown in panel C) (Marwan et al., 2007).

A **B**

Figure 3 A+B Recurrence plots (RPs) and recurrence quantification analysis (RQA) measures for synthetic data representing common types of dynamic behavior: (A) normally-distributed (Gaussian) noise, (B) composite signal comprising two sine waves and a positive trend in the mean. The examples display the time series (upper panel), the recurrence plot (middle panel) and the RQA measures (lower panel). Embedding parameters m =embedding dimension, τ =time delay, e =threshold, w =window size, ws =window moving steps, $norm$ =vector norm, $thei$ =size of Theiler window, $lmin$ =minimum line length, RQA measures RR =recurrence rate and DET =determinism. Please read the methods section for a detailed description of the embedding parameters and RQA measures.

C **D**

Figure 3 C+D Recurrence plots (RPs) and recurrence quantification analysis (RQA) measures for synthetic data representing common types of dynamic behavior: (C) composite signal comprising a sine wave and Gaussian noise with decreasing signal-to-noise ratio from left to right, (D) composite signal comprising two sine waves and a trend in the frequencies. The examples display the time series (upper panel), the recurrence plot (middle panel) and the RQA measures (lower panel). See previous figure for the meaning of the abbreviations.

E **F**

Figure 3 E+F Recurrence plots (RPs) and recurrence quantification analysis (RQA) measures for synthetic data representing common types of dynamic behavior: (E) abrupt transition from a composite signal comprising two sine waves to a signal with only one sine wave, (F) abrupt transition in the amplitudes of two sine waves, and (F) normally-distributed (Gaussian) noise with a stepwise transition in the mean and a change in the autocorrelation prior to this transition. The examples display the time series (upper panel), the recurrence plot (middle panel) and the RQA measures (lower panel). See previous figure for the meaning of the abbreviations.

A **B**

Figure 4 Recurrence plot (RP) and recurrence quantification analysis (RQA) measures of the complete record (–45,358 to 0 yrs BP) from the Chew Bahir basin: time series (upper panel), the recurrence plot (middle panel) and the RQA measures (lower panel) of moving windows determined either by (A) calculating the RQA measures for windows moving along a single (global) RP and (B) by calculating individual RPs for windows moving along the entire time series. See previous figure for the meaning of the abbreviations.

Figure 5 Recurrence plot (RP) and recurrence quantification analysis (RQA) measures for the Chew Bahir basin covering of the time interval between 35 and 20 kyr BP: time series (upper panel), the recurrence plot (middle panel) and the RQA measures (lower panel) of moving windows determined by calculating individual RPs for windows moving along the entire time series. See previous figure for the meaning of the abbreviations.

Figure 6 Recurrence plot (RP) and recurrence quantification analysis (RQA) measures for the Chew Bahir basin covering of the time interval between 20 and 1 kyr BP: time series (upper panel), the recurrence plot (middle panel) and the RQA measures (lower panel) of moving windows determined by calculating individual RPs for windows moving along the entire time series. See previous figure for the meaning of the abbreviations.

Figure 7 Recurrence plot (RP) and recurrence quantification analysis (RQA) measures for the Chew Bahir basin of the time interval between 9 and 1 kyr BP: time series (upper panel), the recurrence plot (middle panel) and the RQA measures (lower panel) of moving windows determined by calculating individual RPs for windows moving along the time series. See previous figure for the meaning of the abbreviations.