

## BOUNDARY ELEMENT METHODS WITH WEAKLY IMPOSED BOUNDARY CONDITIONS\*

TIMO BETCKE<sup>†</sup>, ERIK BURMAN<sup>†</sup>, AND MATTHEW W. SCROGGS<sup>†</sup>

**Abstract.** We consider boundary element methods where the Calderón projector is used for the system matrix and boundary conditions are weakly imposed using a particular variational boundary operator designed using techniques from augmented Lagrangian methods. Regardless of the boundary conditions, both the primal trace variable and the flux are approximated. We focus on the imposition of Dirichlet, mixed Dirichlet–Neumann, and Robin conditions. A salient feature of the Robin condition is that the conditioning of the system is robust also for stiff boundary conditions. The theory is illustrated by a series of numerical examples.

**Key words.** boundary element method, weak boundary conditions, mixed boundary conditions, Robin conditions, Calderon projection

**AMS subject classifications.** 65N38, 65R20

**DOI.** 10.1137/18M119625X

**1. Introduction.** Weak imposition of boundary conditions has been very successful in the context of finite element methods. In particular, Nitsche’s method [19] has recently received increased interest in the scientific computation community. Our aim in this paper is to discuss how the idea behind this type of method can be applied in the context of boundary element methods to impose different types of boundary condition in a unified framework.

Weak imposition of boundary conditions here means that neither the Dirichlet trace nor the Neumann trace is imposed exactly; instead an  $h$ -dependent boundary condition is imposed that is weighted in such a way that optimal error estimates may be derived and the exact boundary condition is recovered in the asymptotic limit. Methods based on Nitsche’s method have been successfully utilized for boundary element method domain decomposition problems, where they have been used to impose interface conditions at one-dimensional (1D) interfaces between segments of 2D screens embedded in 3D space [13, 10]. Our approach instead focuses on imposing boundary conditions on the 2D boundary of a single domain problem through the addition of penalty terms to a general formulation written in terms of the multitrace operator, in a similar vein to the method discussed in [1] for the finite element method.

The use of systems of boundary integral equations for problems with mixed boundary conditions is quite classical [11, 25, 26, 27]. While these papers require the assembly of boundary operators on subsets of the boundary mesh, the penalty method proposed in this paper requires only the addition of sparse mass matrices to the multitrace operator assembled on the entire mesh. In addition to the greater simplicity of the resulting formulation, this method has the advantage that the sparse penalty terms only affect the entries in the matrix for near interactions: this gives the resulting

---

\*Submitted to the journal’s Methods and Algorithms for Scientific Computing section June 25, 2018; accepted for publication (in revised form) January 25, 2019; published electronically May 2, 2019.

<http://www.siam.org/journals/sisc/41-3/M119625.html>

**Funding:** The work of the first and second authors was supported by EPSRC grants EP/P01576X/1 and EP/P012434/1.

<sup>†</sup>Department of Mathematics, University College London, London, WC1E 6BT, UK (t.betcke@ucl.ac.uk, e.burman@ucl.ac.uk, matthew.scroggs.14@ucl.ac.uk, <http://www.mscroggs.co.uk>).

system a structure that can be utilized when designing effective preconditioners.

This approach may not be competitive in the simple case of pure Dirichlet or Neumann conditions due to the increase in the number of unknowns. Therefore the main focus of this work is on more complex situations. We will discuss the following four model cases:

1. nonhomogeneous Dirichlet conditions,
2. nonhomogeneous Neumann conditions,
3. mixed Dirichlet–Neumann boundary conditions,
4. generalized Robin conditions.

We consider the Laplace equation: Find  $u$  such that

$$\begin{aligned}
 (1.1a) \quad & -\Delta u = 0 && \text{in } \Omega, \\
 (1.1b) \quad & u = g_D && \text{on } \Gamma_D, \\
 (1.1c) \quad & \frac{\partial u}{\partial \boldsymbol{\nu}} = g_N && \text{on } \Gamma_N, \\
 (1.1d) \quad & \frac{\partial u}{\partial \boldsymbol{\nu}} = \frac{1}{\varepsilon}(g_D - u) + g_N && \text{on } \Gamma_R.
 \end{aligned}$$

Here  $\Omega \subset \mathbb{R}^3$  denotes a polyhedral domain with outward pointing normal  $\boldsymbol{\nu}$  and boundary  $\Gamma := \Gamma_D \cup \Gamma_N \cup \Gamma_R$ . We assume for simplicity that the boundaries between  $\Gamma_D$ ,  $\Gamma_N$ , and  $\Gamma_R$  coincide with edges between the faces of  $\Gamma$ . Whenever it is ambiguous, we will write  $\boldsymbol{\nu}_{\boldsymbol{x}}$  for the outward pointing normal at the point  $\boldsymbol{x}$ . We assume that  $g_D \in H^{1/2}(\Gamma_D \cup \Gamma_R)$  and  $g_N \in L^2(\Gamma_N \cup \Gamma_R)$ . Observe that, by the Lax–Milgram lemma, there exists a unique solution to (1.1). We assume that  $u \in H^{3/2+\epsilon}(\Omega)$  for some  $\epsilon > 0$ .

For the Robin boundary condition, we will use the ideas of Juntunen and Stenberg [16]. A salient feature of this type of imposition of the Robin condition is that it is robust under singular perturbations. Indeed regardless of the Robin coefficient, the conditioning of the resulting system matrix is no worse than for the Neumann or the Dirichlet problem.

The proposed framework is flexible and allows for the design of a range of different methods depending on the choice of weights and residuals. We will present a sample of possible methods with the ambition of showing the versatility of the framework rather than claiming that for each case the choices are optimal.

An outline of the paper is as follows. First, we review some of the basic elements of the theory of boundary operators in section 2. Then, in section 3 we discuss the design of formulations for the linear model problems in a formal setting. We propose the corresponding boundary element methods in section 4 and give an abstract analysis. The boundary elements obtained using the formulations from section 3 are then shown to satisfy the assumptions of the abstract theory. Finally, we show some computational examples in section 5.

While the present paper focuses on weak imposition of boundary conditions through Nitsche type coupling for BEM, ultimately the goal is to develop a framework for complex BEM/BEM and FEM/BEM multiphysics coupling situations. Existing approaches here are often built upon FETI and BETI type methods [17, 18]. While BETI is usually formulated in terms of Steklov–Poincaré operators, the framework proposed in this paper builds directly upon Calderón projectors of the subdomains.

For the method proposed in the present work the multidomain coupling will take a form similar to that using Nitsche’s method in the FEM/FEM coupling setting of [5]; see also the FEM/BEM coupling of [9], where a Nitsche’s method for the coupling was proposed, using the Steklov–Poincaré operator for the BEM system.

An important application area for the presented weak imposition of boundary conditions is inverse problems with unknown boundary conditions. Since the boundary condition only enters through a sparse operator this can be easily updated in each step of a solver iteration, while the boundary integral operators only need to be computed once. In particular, for reconstruction of the coefficient in a Robin condition (see, e.g., [15] for a finite element approach and [3] for a detailed analysis of the stability of this problem), the robustness with respect to the coefficient of the present method is an advantage.

**2. Boundary operators.** We define the Green's function for the Laplace operator in  $\mathbb{R}^3$  by

$$(2.1) \quad G(\mathbf{x}, \mathbf{y}) = \frac{1}{4\pi|\mathbf{x} - \mathbf{y}|}.$$

In this paper, we focus on the problem in  $\mathbb{R}^3$ . A similar analysis can be used for problems in  $\mathbb{R}^2$ , in which case this definition should be replaced by  $G(\mathbf{x}, \mathbf{y}) = -\log|\mathbf{x} - \mathbf{y}|/2\pi$ .

In the standard fashion (see, e.g., [23, Chapter 6]), we define the single layer potential operator,  $\mathcal{V} : H^{-1/2}(\Gamma) \rightarrow H^1(\Omega)$ , and the double layer potential operator,  $\mathcal{K} : H^{1/2}(\Gamma) \rightarrow H^1(\Omega)$ , for  $v \in H^{1/2}(\Gamma)$ ,  $\mu \in H^{-1/2}(\Gamma)$ , and  $\mathbf{x} \in \Omega \setminus \Gamma$  by

$$(2.2) \quad (\mathcal{V}\mu)(\mathbf{x}) := \int_{\Gamma} G(\mathbf{x}, \mathbf{y})\mu(\mathbf{y}) \, d\mathbf{y},$$

$$(2.3) \quad (\mathcal{K}v)(\mathbf{x}) := \int_{\Gamma} \frac{\partial G(\mathbf{x}, \mathbf{y})}{\partial \nu_{\mathbf{y}}} v(\mathbf{y}) \, d\mathbf{y}.$$

We define the space  $H^1(\Delta, \Omega) := \{v \in H^1(\Omega) : \Delta v \in L^2(\Omega)\}$ , and then we define the Dirichlet and Neumann traces,  $\gamma_{\text{D}} : H^1(\Omega) \rightarrow H^{1/2}(\Gamma)$  and  $\gamma_{\text{N}} : H^1(\Delta, \Omega) \rightarrow H^{-1/2}(\Gamma)$ , by

$$(2.4) \quad \gamma_{\text{D}}f(\mathbf{x}) := \lim_{\Omega \ni \mathbf{y} \rightarrow \mathbf{x} \in \Gamma} f(\mathbf{y}),$$

$$(2.5) \quad \gamma_{\text{N}}f(\mathbf{x}) := \lim_{\Omega \ni \mathbf{y} \rightarrow \mathbf{x} \in \Gamma} \nu_{\mathbf{x}} \cdot \nabla f(\mathbf{y}).$$

We recall that if the Dirichlet and Neumann traces of a harmonic function are known, then the potentials (2.2) and (2.3) may be used to reconstruct the function in  $\Omega$  using the following relation:

$$(2.6) \quad u = -\mathcal{K}(\gamma_{\text{D}}u) + \mathcal{V}(\gamma_{\text{N}}u).$$

It is also known [23, Lemma 6.6] that  $\forall \mu \in H^{-1/2}(\Gamma)$ , the function

$$(2.7) \quad u_{\mu}^{\mathcal{V}} := \mathcal{V}\mu$$

satisfies  $-\Delta u_{\mu}^{\mathcal{V}} = 0$  and

$$(2.8) \quad \|u_{\mu}^{\mathcal{V}}\|_{H^1(\Omega)} \leq c\|\mu\|_{H^{-1/2}(\Gamma)}.$$

Similarly, for the double layer potential there holds [23, Lemma 6.10] that  $\forall v \in H^{1/2}(\Gamma)$ , the function

$$(2.9) \quad u_v^{\mathcal{K}} := \mathcal{K}v$$

satisfies  $-\Delta u_v^{\mathcal{K}} = 0$  and

$$(2.10) \quad \|u_v^{\mathcal{K}}\|_{H^1(\Omega)} \leq c\|v\|_{H^{1/2}(\Gamma)}.$$

We define  $\{\gamma_{\mathcal{D}}f\}_{\Gamma}$  and  $\{\gamma_{\mathcal{N}}f\}_{\Gamma}$  to be the averages of the interior and exterior Dirichlet and Neumann traces of  $f$ . We define the single layer, double layer, adjoint double layer, and hypersingular boundary integral operators,  $\mathcal{V} : H^{-1/2}(\Gamma) \rightarrow H^{1/2}(\Gamma)$ ,  $\mathcal{K} : H^{1/2}(\Gamma) \rightarrow H^{1/2}(\Gamma)$ ,  $\mathcal{K}' : H^{-1/2}(\Gamma) \rightarrow H^{-1/2}(\Gamma)$ , and  $\mathcal{W} : H^{1/2}(\Gamma) \rightarrow H^{-1/2}(\Gamma)$ , by

$$(2.11a) \quad (\mathcal{K}v)(\mathbf{x}) := \{\gamma_{\mathcal{D}}\mathcal{K}v\}_{\Gamma}(\mathbf{x}), \quad (\mathcal{V}\mu)(\mathbf{x}) := \{\gamma_{\mathcal{D}}\mathcal{V}\mu\}_{\Gamma}(\mathbf{x}),$$

$$(2.11b) \quad (\mathcal{W}v)(\mathbf{x}) := -\{\gamma_{\mathcal{N}}\mathcal{K}v\}_{\Gamma}(\mathbf{x}), \quad (\mathcal{K}'\mu)(\mathbf{x}) := \{\gamma_{\mathcal{N}}\mathcal{V}\mu\}_{\Gamma}(\mathbf{x}),$$

where  $\mathbf{x} \in \Gamma$ ,  $v \in H^{1/2}(\Gamma)$ , and  $\mu \in H^{-1/2}(\Gamma)$  [23, Chapter 6].

The following coercivity results are known for the single layer and hypersingular operators in  $\mathbb{R}^3$ , where  $\langle \cdot, \cdot \rangle_{\Gamma}$  denotes the  $H^{1/2}(\Gamma)$ – $H^{-1/2}(\Gamma)$  duality pairing.

LEMMA 2.1 (coercivity of  $\mathcal{V}$ ). *There exists  $\alpha_{\mathcal{V}} > 0$  such that*

$$\alpha_{\mathcal{V}}\|\mu\|_{H^{-1/2}(\Gamma)}^2 \leq \langle \mathcal{V}\mu, \mu \rangle_{\Gamma} \quad \forall \mu \in H^{-1/2}(\Gamma).$$

*Proof.* See [23, Theorem 6.22] for the proof.  $\square$

LEMMA 2.2 (coercivity of  $\mathcal{W}$ ). *There exists  $\alpha_{\mathcal{W}} > 0$  such that*

$$\alpha_{\mathcal{W}}\|v\|_{H^{1/2}(\Gamma)}^2 \leq \langle \mathcal{W}v, v \rangle_{\Gamma}, \quad \forall v \in H_*^{1/2}(\Gamma),$$

where  $H_*^{1/2}(\Gamma)$  denotes the set of functions  $v \in H^{1/2}(\Gamma)$  such that  $\bar{v} = 0$ , where  $\bar{v} := \frac{\langle v, 1 \rangle_{\Gamma}}{\langle 1, 1 \rangle_{\Gamma}}$  is the average value of  $v$ . From this it follows that

$$\alpha_{\mathcal{W}}|v|_{H_*^{1/2}(\Gamma)}^2 \leq \langle \mathcal{W}v, v \rangle_{\Gamma} \quad \forall v \in H^{1/2}(\Gamma),$$

where  $|\cdot|_{H_*^{1/2}(\Gamma)}$  is defined, for  $v \in H^{1/2}(\Gamma)$ , by  $|v|_{H_*^{1/2}(\Gamma)} := \|v - \bar{v}\|_{H^{1/2}(\Gamma)}$ .

*Proof.* See [23, Theorem 6.24] for the proof.  $\square$

The following boundedness results are also known.

LEMMA 2.3 (boundedness). *There exist  $C_{\mathcal{V}}, C_{\mathcal{K}}, C_{\mathcal{K}'}, C_{\mathcal{W}} > 0$  such that*

$$(i) \quad \|\mathcal{V}\mu\|_{H^{1/2}(\Gamma)} \leq C_{\mathcal{V}}\|\mu\|_{H^{-1/2}(\Gamma)} \quad \forall \mu \in H^{-1/2}(\Gamma),$$

$$(ii) \quad \|\mathcal{K}v\|_{H^{1/2}(\Gamma)} \leq C_{\mathcal{K}}\|v\|_{H^{1/2}(\Gamma)} \quad \forall v \in H^{1/2}(\Gamma),$$

$$(iii) \quad \|\mathcal{K}'\mu\|_{H^{-1/2}(\Gamma)} \leq C_{\mathcal{K}'}\|\mu\|_{H^{-1/2}(\Gamma)} \quad \forall \mu \in H^{-1/2}(\Gamma),$$

$$(iv) \quad \|\mathcal{W}v\|_{H^{-1/2}(\Gamma)} \leq C_{\mathcal{W}}\|v\|_{H^{1/2}(\Gamma)} \quad \forall v \in H^{1/2}(\Gamma).$$

*Proof.* See [23, sections 6.2–6.5] for the proof.  $\square$

We define the Calderón projector by

$$(2.12) \quad \mathcal{C} := \begin{pmatrix} (1 - \sigma)\text{Id} - \mathcal{K} & \mathcal{V} \\ \mathcal{W} & \sigma\text{Id} + \mathcal{K}' \end{pmatrix},$$

where  $\sigma$  is defined as in [23, equation (6.11)], and recall that if  $u$  is a solution of (1.1), then it satisfies

$$(2.13) \quad \mathcal{C} \begin{pmatrix} \gamma_{\mathcal{D}}u \\ \gamma_{\mathcal{N}}u \end{pmatrix} = \begin{pmatrix} \gamma_{\mathcal{D}}u \\ \gamma_{\mathcal{N}}u \end{pmatrix}.$$

Taking the product of (2.13) with two test functions, and using the fact that  $\sigma = \frac{1}{2}$  almost everywhere, we arrive at the following equations:

$$(2.14) \quad \langle \gamma_{\mathbb{D}}u, \mu \rangle_{\Gamma} = \langle (\frac{1}{2}\text{Id} - \mathbf{K})\gamma_{\mathbb{D}}u, \mu \rangle_{\Gamma} + \langle \mathbf{V}\gamma_{\mathbb{N}}u, \mu \rangle_{\Gamma} \quad \forall \mu \in H^{-1/2}(\Gamma),$$

$$(2.15) \quad \langle \gamma_{\mathbb{N}}u, v \rangle_{\Gamma} = \langle (\frac{1}{2}\text{Id} + \mathbf{K}')\gamma_{\mathbb{N}}u, v \rangle_{\Gamma} + \langle \mathbf{W}\gamma_{\mathbb{D}}u, v \rangle_{\Gamma} \quad \forall v \in H^{1/2}(\Gamma).$$

For a more compact notation, we introduce  $\lambda = \gamma_{\mathbb{N}}u$  and  $u = \gamma_{\mathbb{D}}u$  and the Calderón form

$$(2.16) \quad \mathcal{C}[(u, \lambda), (v, \mu)] := \langle (\frac{1}{2}\text{Id} - \mathbf{K})u, \mu \rangle_{\Gamma} + \langle \mathbf{V}\lambda, \mu \rangle_{\Gamma} + \langle (\frac{1}{2}\text{Id} + \mathbf{K}')\lambda, v \rangle_{\Gamma} + \langle \mathbf{W}u, v \rangle_{\Gamma}.$$

We may then rewrite (2.14) and (2.15) as

$$(2.17) \quad \mathcal{C}[(u, \lambda), (v, \mu)] = \langle u, \mu \rangle_{\Gamma} + \langle \lambda, v \rangle_{\Gamma}.$$

We will also frequently use the multitrace form, defined by

$$(2.18) \quad \mathcal{A}[(u, \lambda), (v, \mu)] := -\langle \mathbf{K}u, \mu \rangle_{\Gamma} + \langle \mathbf{V}\lambda, \mu \rangle_{\Gamma} + \langle \mathbf{K}'\lambda, v \rangle_{\Gamma} + \langle \mathbf{W}u, v \rangle_{\Gamma}.$$

Using this, we may rewrite (2.17) as

$$(2.19) \quad \mathcal{A}[(u, \lambda), (v, \mu)] = \frac{1}{2} \langle u, \mu \rangle_{\Gamma} + \frac{1}{2} \langle \lambda, v \rangle_{\Gamma}.$$

To quantify the two traces we introduce the product space

$$\mathbb{V} := \begin{cases} H^{1/2}(\Gamma) \times H^{-1/2}(\Gamma) & \text{if } \Gamma_{\mathbb{N}} \cup \Gamma_{\mathbb{R}} = \emptyset, \\ H^{1/2}(\Gamma) \times L^2(\Gamma) & \text{otherwise.} \end{cases}$$

The additional regularity on the flux variable is required later when imposing Neumann and Robin conditions. We also introduce the associated norm

$$\|(v, \mu)\|_{\mathbb{V}} := \|v\|_{H^{1/2}(\Gamma)} + \|\mu\|_{H^{-1/2}(\Gamma)}.$$

Using the results in Lemmas 2.1 to 2.3, we obtain the continuity and coercivity of  $\mathcal{A}$ .

LEMMA 2.4 (continuity). *There exists  $C > 0$  such that*

$$|\mathcal{A}[(w, \eta), (v, \mu)]| \leq C\|(w, \eta)\|_{\mathbb{V}}\|(v, \mu)\|_{\mathbb{V}} \quad \forall (w, \eta), (v, \mu) \in \mathbb{V}.$$

*Proof.* Use the stability results from Lemma 2.3. □

LEMMA 2.5 (coercivity). *There exists  $\alpha > 0$  such that*

$$\alpha \left( \|v\|_{H^{1/2}(\Gamma)}^2 + \|\mu\|_{H^{-1/2}(\Gamma)}^2 \right) \leq \mathcal{A}[(v, \mu), (v, \mu)] \quad \forall (v, \mu) \in \mathbb{V}.$$

*Proof.* Use the coercivity of  $\mathbf{V}$  and  $\mathbf{W}$  from Lemmas 2.1 and 2.2 and let  $\alpha = \min(\alpha_{\mathbf{W}}, \alpha_{\mathbf{V}})$ . □

**3. Weak imposition of boundary conditions.** In this section, we will derive boundary integral formulations of the problem (1.1), that we will then use for our boundary element formulations. We assume that the boundary condition may be written as

$$(3.1) \quad R_\Gamma(u, \lambda) = 0.$$

The idea that we will exploit in the following is simply to add a suitable weighted weak form of this constraint to the Calderón form (2.17). Formally, this leads to an expression of the form

$$(3.2) \quad \mathcal{C}[(u, \lambda), (v, \mu)] = \langle u, \mu \rangle_\Gamma + \langle \lambda, v \rangle_\Gamma + \langle R_\Gamma(u, \lambda), \beta_1 v + \beta_2 \mu \rangle_\Gamma,$$

or equivalently

$$(3.3) \quad \mathcal{A}[(u, \lambda), (v, \mu)] = \frac{1}{2} \langle u, \mu \rangle_\Gamma + \frac{1}{2} \langle \lambda, v \rangle_\Gamma + \langle R_\Gamma(u, \lambda), \beta_1 v + \beta_2 \mu \rangle_\Gamma,$$

where  $\beta_1$  and  $\beta_2$  are problem-dependent scaling operators that will be chosen as a function of the physical parameters in order to obtain robustness of the method.

**3.1. Dirichlet boundary condition.** In this section, we assume that  $\Gamma_D \equiv \Gamma$  and consider the resulting Dirichlet problem. We choose  $\beta_1 = \beta_D^{1/2}$ ,  $\beta_2 = \beta_D^{-1/2}$ , where  $\beta_D$  will be identified with a mesh-dependent penalty parameter, and

$$(3.4) \quad R_{\Gamma_D}(u, \lambda) := \beta_D^{1/2}(g_D - u),$$

where  $g_D \in H^{1/2}(\Gamma)$  is the Dirichlet data.

Inserting this into (3.3), we obtain the formulation

$$(3.5) \quad \mathcal{A}[(u, \lambda), (v, \mu)] - \frac{1}{2} \langle \lambda, v \rangle_{\Gamma_D} + \frac{1}{2} \langle u, \mu \rangle_{\Gamma_D} + \langle \beta_D u, v \rangle_{\Gamma_D} = \langle g_D, \beta_D v + \mu \rangle_{\Gamma_D}.$$

One can compare the method with the classical (nonsymmetric) Nitsche's method by formally identifying  $\lambda$  with  $\partial_\nu u$  and  $\mu$  with  $\partial_\nu v$  (up to the multiplicative factor  $\frac{1}{2}$ ).

For a more compact notation, we introduce the boundary operator associated with the nonhomogeneous Dirichlet condition

$$(3.6) \quad \mathcal{B}_D[(u, \lambda), (v, \mu)] := -\frac{1}{2} \langle \lambda, v \rangle_{\Gamma_D} + \frac{1}{2} \langle u, \mu \rangle_{\Gamma_D} + \langle \beta_D u, v \rangle_{\Gamma_D}$$

and the operator associated with the right-hand side

$$(3.7) \quad \mathcal{L}_D(v, \mu) := \langle g_D, \beta_D v + \mu \rangle_{\Gamma_D}.$$

Using these and (3.5), we arrive at the following problem: Find  $(u, \lambda) \in \mathbb{V}$  such that

$$(3.8) \quad \mathcal{A}[(u, \lambda), (v, \mu)] + \mathcal{B}_D[(u, \lambda), (v, \mu)] = \mathcal{L}_D(v, \mu) \quad \forall (v, \mu) \in \mathbb{V}.$$

If we set  $\beta_D = 0$  in (3.6) and (3.7), we obtain a penalty-free formulation for the Dirichlet problem.

**3.2. Neumann boundary condition.** In this section, we assume that  $\Gamma_N \equiv \Gamma$  and consider the resulting Neumann problem. We choose  $\beta_1 = \beta_N^{-1/2}$ ,  $\beta_2 = \beta_N^{1/2}$ , and define

$$(3.9) \quad R_{\Gamma_N}(u, \lambda) := \beta_N^{1/2}(g_N - \lambda),$$

where  $g_N \in L^2(\Gamma_N)$ , with  $\int_\Gamma g_N = 0$ , is the Neumann data.

Proceeding as in the Dirichlet case, we obtain the formulation

$$(3.10) \quad \mathcal{A}[(u, \lambda), (v, \mu)] - \frac{1}{2} \langle u, \mu \rangle_{\Gamma_N} + \frac{1}{2} \langle \lambda, v \rangle_{\Gamma_N} + \langle \beta_N \lambda, \mu \rangle_{\Gamma_N} = \langle g_N, \beta_N \mu + v \rangle_{\Gamma_N}.$$

Defining

$$(3.11) \quad \mathcal{B}_N[(u, \lambda), (v, \mu)] := -\frac{1}{2} \langle u, \mu \rangle_{\Gamma_N} + \frac{1}{2} \langle \lambda, v \rangle_{\Gamma_N} + \langle \beta_N \lambda, \mu \rangle_{\Gamma_N},$$

$$(3.12) \quad \mathcal{L}_N(v, \mu) := \langle g_N, \beta_N \mu + v \rangle_{\Gamma_N},$$

we may write this as the variational problem: Find  $(u, \lambda) \in \mathbb{V}^*$  such that

$$(3.13) \quad \mathcal{A}[(u, \lambda), (v, \mu)] + \mathcal{B}_N[(u, \lambda), (v, \mu)] = \mathcal{L}_N(v, \mu) \quad \forall (v, \mu) \in \mathbb{V}^*.$$

Here, we use the space  $\mathbb{V}^* := H_*^{1/2}(\Gamma_N) \times L^2(\Gamma_N)$ , as the solution to the Neumann problem can only be determined up to a constant, so we include the extra condition that  $\bar{u} = 0$ .

If we set  $\beta_N = 0$  in (3.11) and (3.12), we obtain a penalty-free formulation for the Neumann problem. In this case, we may take  $\mathbb{V}^* = H_*^{1/2}(\Gamma_N) \times H^{-1/2}(\Gamma_N)$  and  $g_N \in H^{-1/2}(\Gamma_N)$ .

When  $\beta_N > 0$ , observe that for the terms imposing the Neumann condition to be well defined, we need  $\lambda \in L^2(\Gamma_N)$ . This can be avoided by replacing  $\beta_N$  with a regularizing operator  $R : H^{-1/2}(\Gamma_N) \rightarrow H^{1/2}(\Gamma_N)$ . For example, we could take  $R = \beta_V \mathbf{V}$ , where  $\beta_V \in \mathbb{R}$  and  $\mathbf{V}$  is the single layer boundary operator on  $\Gamma_N$ . This formulation with the operator  $R$  is given in [24, (3.10) and (3.11)], where it was derived using a domain decomposition approach where a Robin condition was used to weakly impose a Neumann condition.

The resulting formulations using  $\beta_N$  are in general easier to analyze, since they give control of  $\lambda$  on the Neumann boundary in the natural norm  $\|\lambda\|_{H^{-1/2}(\Gamma_N)}$ .

**3.3. Mixed Dirichlet–Neumann boundary condition.** We now consider the case of mixed Dirichlet–Neumann boundary conditions, when  $\Gamma = \Gamma_D \cup \Gamma_N$ . We note that in this case, and in the Robin case, we take  $\mathbb{V} = H^{1/2}(\Gamma) \times L^2(\Gamma)$ .

Let  $R_{\Gamma_D}$  and  $R_{\Gamma_N}$  be defined by (3.4) and (3.9). Using the abstract form (3.3), we obtain

$$(3.14) \quad \mathcal{A}[(u, \lambda), (v, \mu)] = \frac{1}{2} \langle u, \mu \rangle_{\Gamma} + \frac{1}{2} \langle \lambda, v \rangle_{\Gamma} + \left\langle R_{\Gamma_D}(u, \lambda), \beta_D^{1/2} v + \beta_D^{-1/2} \mu \right\rangle_{\Gamma_D} + \left\langle R_{\Gamma_N}(u, \lambda), \beta_N^{-1/2} v + \beta_N^{1/2} \mu \right\rangle_{\Gamma_N}.$$

Developing (3.14), and defining

$$(3.15) \quad \mathcal{B}_{ND}[(u, \lambda), (v, \mu)] := \frac{1}{2} \langle u, \mu \rangle_{\Gamma_D} - \frac{1}{2} \langle \lambda, v \rangle_{\Gamma_D} + \langle \beta_D u, v \rangle_{\Gamma_D} + \frac{1}{2} \langle \lambda, v \rangle_{\Gamma_N} - \frac{1}{2} \langle u, \mu \rangle_{\Gamma_N} + \langle \beta_N \lambda, \mu \rangle_{\Gamma_N},$$

$$(3.16) \quad \mathcal{L}_{ND}(v, \mu) := \langle g_D, \beta_D v + \mu \rangle_{\Gamma_D} + \langle g_N, \beta_N \mu + v \rangle_{\Gamma_N},$$

we arrive the variational formulation: Find  $(u, \lambda) \in \mathbb{V}$  such that

$$(3.17) \quad \mathcal{A}[(u, \lambda), (v, \mu)] + \mathcal{B}_{ND}[(u, \lambda), (v, \mu)] = \mathcal{L}_{ND}(v, \mu) \quad \forall (v, \mu) \in \mathbb{V}.$$

If we set  $\beta_D = 0$  and  $\beta_N = 0$  in (3.15) and (3.16), we obtain a penalty-free formulation for the mixed Dirichlet–Neumann problem. By taking  $\Gamma_N = \emptyset$  or  $\Gamma_D = \emptyset$ , formulations for both Dirichlet and Neumann problems can be obtained from (3.17).

**3.4. Robin conditions.** For simplicity, we consider the case where  $\Gamma = \Gamma_R$ . Considering the Robin condition (1.1d), we may write, for some  $\varepsilon > 0$ ,

$$(3.18) \quad R_{\Gamma_R}(u, \lambda) := \beta_R^{1/2} \left( \varepsilon^{1/2}(g_N - \lambda) + \varepsilon^{-1/2}(g_D - u) \right).$$

This function is a linear combination of the Dirichlet and the Neumann conditions.

$$(3.19) \quad R_{\Gamma_R}(u, \lambda) = \alpha_D R_{\Gamma_D}(u, \lambda) + \alpha_N R_{\Gamma_N}(u, \lambda),$$

where  $\alpha_N = \beta_R^{1/2} \beta_N^{-1/2} \varepsilon^{1/2}$  and  $\alpha_D = \beta_R^{1/2} \beta_D^{-1/2} \varepsilon^{-1/2}$ .

We take  $\beta_1 = \beta_R^{1/2}$  and  $\beta_2 = \beta_R^{-1/2}$  and look for a term of the form

$$(3.20) \quad \left\langle \phi R_{\Gamma_R}(u, \lambda), \beta_R^{1/2} v + \beta_R^{-1/2} \mu \right\rangle_{\Gamma_R},$$

where the  $\phi$  and  $\beta_R$  must have the following properties to ensure that the formulation degenerates into the formulation for the Dirichlet and Neumann problems as  $\varepsilon \rightarrow 0$  and  $\varepsilon \rightarrow \infty$ :

$$\begin{array}{llll} \beta_R \rightarrow \beta_D, & \alpha_D \phi \rightarrow 1, & \text{and} & \alpha_N \phi \rightarrow 0 & \text{as } \varepsilon \rightarrow 0, \\ \beta_R \rightarrow \beta_N^{-1}, & \alpha_N \phi \rightarrow 1, & \text{and} & \alpha_D \phi \rightarrow 0 & \text{as } \varepsilon \rightarrow \infty. \end{array}$$

It is straightforward to verify that these conditions are satisfied for the choices

$$(3.21) \quad \phi := \frac{\varepsilon^{1/2}}{\varepsilon \beta_R + 1},$$

$$(3.22) \quad \beta_R := \frac{\varepsilon \beta_N^{-1} + \beta_D}{\varepsilon + 1}.$$

Later, we will use  $\beta_D = \beta h^{-1}$  and  $\beta_N = \beta h$ , where  $\beta$  is a constant, as in the mixed Dirichlet–Neumann case.

Collecting the above considerations, we arrive at the formulation

$$(3.23) \quad \mathcal{A}[(u, \lambda), (v, \mu)] = \frac{1}{2} \langle u, \mu \rangle_{\Gamma} + \frac{1}{2} \langle \lambda, v \rangle_{\Gamma} + \left\langle \varepsilon(g_N - \lambda) + (g_D - u), \frac{\beta_R}{\varepsilon \beta_R + 1} v + \frac{1}{\varepsilon \beta_R + 1} \mu \right\rangle_{\Gamma_R}.$$

Taking  $\varepsilon \rightarrow 0$ , we recover the Dirichlet formulation (3.5), and taking  $\varepsilon \rightarrow \infty$  results in the Neumann formulation (3.10).

By introducing

$$\begin{aligned} \mathcal{B}_R[(u, \lambda), (v, \mu)] &:= \frac{1}{2} \left\langle \frac{\varepsilon \beta_R - 1}{\varepsilon \beta_R + 1} \lambda, v \right\rangle_{\Gamma_R} - \frac{1}{2} \left\langle \frac{\varepsilon \beta_R - 1}{\varepsilon \beta_R + 1} u, \mu \right\rangle_{\Gamma_R} \\ &\quad + \left\langle \frac{\varepsilon}{\varepsilon \beta_R + 1} \lambda, \mu \right\rangle_{\Gamma_R} + \left\langle \frac{\beta_R}{\varepsilon \beta_R + 1} u, v \right\rangle_{\Gamma_R} \end{aligned}$$

and

$$\mathcal{L}_R(v, \mu) := \left\langle g_D + \varepsilon g_N, \frac{\beta_R}{\varepsilon \beta_R + 1} v + \frac{1}{\varepsilon \beta_R + 1} \mu \right\rangle_{\Gamma_R},$$

we may write this as the variational problem: Find  $(u, \lambda) \in \mathbb{V}$  such that

$$(3.24) \quad \mathcal{A}[(u, \lambda), (v, \mu)] + \mathcal{B}_R[(u, \lambda), (v, \mu)] = \mathcal{L}_R(v, \mu) \quad \forall (v, \mu) \in \mathbb{V}.$$



**4. Boundary element method for the single domain problem.** All the methods introduced above are written as the sum of the multitrace operator  $\mathcal{A}$  and a boundary condition operator  $\mathcal{B}$ . We write this generally as follows: Find  $(u, \lambda) \in \mathbb{V}$  such that

$$(4.1) \quad \mathcal{A}[(u, \lambda), (v, \mu)] + \mathcal{B}[(u, \lambda), (v, \mu)] = \mathcal{L}(v, \mu) \quad \forall (v, \mu) \in \mathbb{V}.$$

In this section, we analyze this general problem, then show that the analysis is applicable to the boundary conditions discussed in section 3.

For the sake of example and to fix the ideas, we introduce a family of conforming, shape regular triangulations of  $\Gamma$ ,  $\{\mathcal{T}_h\}_{h>0}$ , indexed by the largest element diameter of the mesh,  $h$ . We assume that the triangulations are fitted to the different boundary sets  $\Gamma_D, \Gamma_R$ , and  $\Gamma_N$ . We then consider the following finite element spaces:

$$\begin{aligned} V_h^k &:= \{v_h \in C^0(\Gamma) : v_h|_T \in \mathbb{P}_k(T) \text{ for every } T \in \mathcal{T}_h\}, \\ \Lambda_h^l &:= \{v_h \in L^2(\Gamma) : v_h|_T \in \mathbb{P}_l(T) \text{ for every } T \in \mathcal{T}_h\}, \\ \tilde{\Lambda}_h^l &:= \{v_h \in \Lambda_h^l : v_h|_{\Gamma_i} \in C^0(\Gamma_i) \text{ for } i = 1, \dots, M\}, \end{aligned}$$

where  $\mathbb{P}_k(T)$  denotes the space of polynomials of order less than or equal to  $k$ , and  $\{\Gamma_i\}_{i=1}^M$  are the polygonal faces of  $\Gamma$ .

We observe that  $V_h^k \subset H^{1/2}(\Gamma)$ ,  $\Lambda_h^l \subset L^2(\Gamma)$ , and  $\tilde{\Lambda}_h^l \subset L^2(\Gamma)$ . We now introduce the discrete product space  $\mathbb{V}_h := V_h^k \times \Lambda_h^l$ . The space  $\tilde{\Lambda}_h^l$  may be used in the place of  $\Lambda_h^l$  without any modifications of the arguments below.

The boundary element formulation of the generic problem (4.1) then takes the following form: Find  $(u_h, \lambda_h) \in \mathbb{V}_h$  such that

$$(4.2) \quad \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] + \mathcal{B}[(u_h, \lambda_h), (v_h, \mu_h)] = \mathcal{L}(v_h, \mu_h) \quad \forall (v_h, \mu_h) \in \mathbb{V}_h.$$

If we assume that  $(u, \lambda) \in \mathbb{V}$  and  $(u_h, \lambda_h) \in \mathbb{V}_h$  satisfy (4.1) and (4.2), it immediately follows that the following Galerkin orthogonality relation holds:

$$(4.3) \quad \mathcal{A}[(u - u_h, \lambda - \lambda_h), (v_h, \mu_h)] + \mathcal{B}[(u - u_h, \lambda - \lambda_h), (v_h, \mu_h)] = 0 \quad \forall (v_h, \mu_h) \in \mathbb{V}_h.$$

We also get the following representation formula for the approximation in the bulk using (2.6):

$$(4.4) \quad \tilde{u}_h = -\mathcal{K}u_h + \mathcal{V}\lambda_h.$$

We will now proceed to derive some estimates for the solution of (4.2) and the reconstruction (4.4).

Let  $\mathbb{W}$  be a product Hilbert space for the primal and flux variables, such that  $\mathbb{V}_h \subset \mathbb{W} \subset \mathbb{V}$ . Let  $\|\cdot\|_{\mathcal{B}}$  be a norm defined on  $\mathbb{W}$ , such that  $\forall (v, \mu) \in \mathbb{W}, \|(v, \mu)\|_{\mathcal{B}} \geq \|(v, \mu)\|_{\mathbb{V}}$ .

To reduce the number of constants that appear, especially when proving that Assumption 4.4 holds, we introduce the following notation:

- If  $\exists C > 0$ , independent of  $h$ , such that  $a \leq Cb$ , then we write  $a \lesssim b$ .
- If  $a \lesssim b$  and  $b \lesssim a$ , then we write  $a \approx b$ .

For the abstract analysis, we will make use of the following standard assumptions.

*Assumption 4.1* (weak coercivity). There exists  $\alpha > 0$  such that  $\forall (v, \mu) \in \mathbb{W}$

$$\alpha \|(v, \mu)\|_{\mathcal{B}} \leq \sup_{(w, \eta) \in \mathbb{W} \setminus \{0\}} \frac{\mathcal{A}[(v, \mu), (w, \eta)] + \mathcal{B}[(v, \mu), (w, \eta)]}{\|(w, \eta)\|_{\mathcal{B}}},$$

and  $\forall (w, \eta) \in \mathbb{W} \setminus \{0\}$

$$\sup_{(v, \mu) \in \mathbb{W}} |\mathcal{A}[(v, \mu), (w, \eta)] + \mathcal{B}[(v, \mu), (w, \eta)]| > 0.$$

*Assumption 4.2* (discrete coercivity). There exists  $\alpha > 0$  such that  $\forall (v_h, \mu_h) \in \mathbb{V}_h$

$$\alpha \|(v_h, \mu_h)\|_{\mathcal{B}} \leq \sup_{(w_h, \eta_h) \in \mathbb{V}_h \setminus \{0\}} \frac{\mathcal{A}[(v_h, \mu_h), (w_h, \eta_h)] + \mathcal{B}[(v_h, \mu_h), (w_h, \eta_h)]}{\|(w_h, \eta_h)\|_{\mathcal{B}}},$$

and  $\forall (w_h, \eta_h) \in \mathbb{V}_h \setminus \{0\}$

$$\sup_{(v_h, \mu_h) \in \mathbb{V}_h} |\mathcal{A}[(v_h, \mu_h), (w_h, \eta_h)] + \mathcal{B}[(v_h, \mu_h), (w_h, \eta_h)]| > 0.$$

*Assumption 4.3* (continuity). There exists an auxiliary norm  $\|(v, \mu)\|_*$  defined on  $\mathbb{W}$ , and there exists  $M > 0$  such that  $\forall (w, \eta), (v, \mu) \in \mathbb{W}$

$$|\mathcal{A}[(w, \eta), (v, \mu)] + \mathcal{B}[(w, \eta), (v, \mu)]| \leq M \|(w, \eta)\|_* \|(v, \mu)\|_{\mathcal{B}}.$$

*Assumption 4.4* (approximation). For all  $(v, \mu) \in H^s(\Gamma) \times H^r(\Gamma)$ ,

$$\inf_{(w_h, \eta_h) \in \mathbb{V}_h} \|(v - w_h, \mu - \eta_h)\|_* \lesssim h^{\zeta-1/2} |v|_{H^{\zeta}(\Gamma)} + h^{\xi+1/2} |\mu|_{H^{\xi}(\Gamma)},$$

where  $\zeta = \min(k + 1, s)$ ,  $\xi = \min(l + 1, r)$ ,  $s \geq \frac{1}{2}$  and  $r \geq -\frac{1}{2}$ .

*Remark 4.5.* In the right-hand side of the bound of Assumption 4.4, the Sobolev norm in the second term should be interpreted as the broken norm over the faces of the polyhedral boundary  $\Gamma$  when  $\xi > 0$ , i.e.,  $|\mu|_{H^{\xi}(\Gamma)} := \sum_{i=1}^M |\mu|_{H^{\xi}(\Gamma_i)}$ , since  $\mu$  may not be globally smooth. Likewise, below we will write  $\lambda \in H^r(\Gamma)$  as short form for  $\lambda \in H^{-\frac{1}{2}}(\Gamma)$  and  $\lambda|_{\Gamma_i} \in H^r(\Gamma_i)$ ,  $i = 1, \dots, M$ , when  $r > 0$ .

Typically, we use approximation spaces with  $k = l + 1$ , where the polynomial spaces used for  $\lambda$  are one order lower than those for  $u$ , or spaces with  $k = l$ , where equal order spaces are used for both variables.

We note that if the form  $\mathcal{A} + \mathcal{B}$  is coercive, that is, there exists  $\alpha > 0$  such that  $\forall (v, \mu) \in \mathbb{W}$

$$\alpha \|(v, \mu)\|_{\mathcal{B}}^2 \leq \mathcal{A}[(v, \mu), (v, \mu)] + \mathcal{B}[(v, \mu), (v, \mu)],$$

then Assumptions 4.1 and 4.2 hold.

We now proceed to prove some results about the abstract problem.

**PROPOSITION 4.6.** *Assume that Assumption 4.1 holds; then the linear system defined by (4.2) is invertible. If, in addition, we assume that*

- *Assumption 4.3 holds,*
- *there exists  $L > 0$  such that  $\mathcal{L}(w, \eta) \leq L \|(w, \eta)\|_{\mathcal{B}} \forall (w, \eta) \in \mathbb{W}$ ,*
- *and  $\|\cdot\|_*$  is equivalent to  $\|\cdot\|_{\mathcal{B}}$ ,*

*then the formulation (4.1) admits a unique solution in  $\mathbb{W}$ .*

*Proof.* Note that Assumption 4.1 implies the inf-sup condition,

$$(4.5) \quad \inf_{(v, \mu) \in \mathbb{W} \setminus \{0\}} \sup_{(w, \eta) \in \mathbb{W} \setminus \{0\}} \frac{\mathcal{A}[(v, \mu), (w, \eta)] + \mathcal{B}[(v, \mu), (w, \eta)]}{\|(v, \mu)\|_{\mathcal{B}} \|(w, \eta)\|_{\mathcal{B}}} > 0.$$

Therefore we may apply the Babuška–Lax–Milgram theorem [2, Theorem 5.2.1].  $\square$

PROPOSITION 4.7. Assume that  $(u, \lambda) \in \mathbb{V}$  is the solution to a boundary value problem of the form (1.1) satisfying the abstract form (4.1). Let  $(u_h, \lambda_h) \in \mathbb{V}_h$  be the solution of (4.2). If Assumptions 4.2 and 4.3 are satisfied, then

$$(4.6) \quad \|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}} \leq \frac{M}{\alpha} \inf_{(v_h, \mu_h) \in \mathbb{V}_h} \|(u - v_h, \lambda - \mu_h)\|_*$$

*Proof.* See [28, Theorem 2]. □

COROLLARY 4.8. Let  $(u, \lambda) \in H^s(\Gamma) \times H^r(\Gamma)$ , for some  $s \geq \frac{1}{2}$  and  $r \geq -\frac{1}{2}$ , satisfy the abstract form (4.1). Under the assumptions of Proposition 4.7 and Assumption 4.4,

$$\|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}} \lesssim h^{\zeta-1/2} |u|_{H^\zeta(\Gamma)} + h^{\xi+1/2} |\lambda|_{H^\xi(\Gamma)},$$

where  $\zeta = \min(k + 1, s)$  and  $\xi = \min(l + 1, r)$ .

*Proof.* Apply Assumption 4.4 to the right-hand side of (4.6). □

PROPOSITION 4.9. Assume that  $(u, \lambda) \in \mathbb{V}$  is the solution to a boundary value problem of the form (1.1) satisfying the abstract form (4.1) and that the assumptions of Proposition 4.7 are satisfied. Let  $(u_h, \lambda_h) \in \mathbb{V}_h$ . Let  $\tilde{u} : \Omega \rightarrow \mathbb{R}$  be the reconstruction obtained using (2.6), with  $\gamma_{\mathbb{N}} u = \lambda$  and  $\gamma_{\mathbb{D}} u = u$ , and let  $\tilde{u}_h : \Omega \rightarrow \mathbb{R}$  be the reconstruction obtained using (4.4). Then there holds

$$\|\tilde{u} - \tilde{u}_h\|_{H^1(\Omega)} \lesssim \frac{M}{\alpha} \inf_{v_h, \mu_h \in \mathbb{V}_h} \|(u - v_h, \lambda - \mu_h)\|_*$$

*Proof.* Using (2.7) and (2.9), we may write

$$\tilde{u} - \tilde{u}_h = (u_\lambda^{\mathcal{V}} - u_{\lambda_h}^{\mathcal{V}}) + (u_u^{\mathcal{K}} - u_{u_h}^{\mathcal{K}}).$$

Using the triangle inequality, we have

$$(4.7) \quad \|\tilde{u} - \tilde{u}_h\|_{H^1(\Omega)} \leq \|u_\lambda^{\mathcal{V}} - u_{\lambda_h}^{\mathcal{V}}\|_{H^1(\Omega)} + \|u_u^{\mathcal{K}} - u_{u_h}^{\mathcal{K}}\|_{H^1(\Omega)}.$$

By (2.8) and (2.10), there exist  $c_1, c_2 > 0$  such that

$$(4.8) \quad \|u_\lambda^{\mathcal{V}} - u_{\lambda_h}^{\mathcal{V}}\|_{H^1(\Omega)} \leq c_1 \|\lambda - \lambda_h\|_{H^{-1/2}(\Gamma)},$$

$$(4.9) \quad \|u_u^{\mathcal{K}} - u_{u_h}^{\mathcal{K}}\|_{H^1(\Omega)} \leq c_2 \|u - u_h\|_{H^{1/2}(\Gamma)}.$$

Collecting (4.7)–(4.9), we see that there exists  $C > 0$  such that

$$(4.10) \quad \|\tilde{u} - \tilde{u}_h\|_{H^1(\Omega)} \leq C \|(\lambda - \lambda_h, u - u_h)\|_{\mathbb{V}} \leq C \|(\lambda - \lambda_h, u - u_h)\|_{\mathcal{B}}.$$

The statement now follows from Proposition 4.7. □

COROLLARY 4.10. Under the same assumptions of Proposition 4.9 and Assumption 4.4,

$$\|\tilde{u} - \tilde{u}_h\|_{H^1(\Omega)} \lesssim h^{\zeta-1/2} |u|_{H^\zeta(\Gamma)} + h^{\xi+1/2} |\lambda|_{H^\xi(\Gamma)},$$

where  $\zeta = \min(k + 1, s)$  and  $\xi = \min(l + 1, r)$ .

*Proof.* Apply Assumption 4.4 to (4.10) in the proof of Proposition 4.9. □

**4.1. Application of the theory to the Dirichlet problem.** For the finite element spaces defined above, the Dirichlet problem takes the following form: Find  $(u_h, \lambda_h) \in \mathbb{V}_h$  such that

$$(4.11) \quad \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] + \mathcal{B}_D[(u_h, \lambda_h), (v_h, \mu_h)] = \mathcal{L}_D(v_h, \mu_h) \quad \forall (v_h, \mu_h) \in \mathbb{V}_h.$$

We introduce the following  $\mathcal{B}_D$ -norm.

$$\|(v, \mu)\|_{\mathcal{B}_D} := \|(v, \mu)\|_{\mathbb{V}} + \beta_D^{1/2} \|v\|_{L^2(\Gamma_D)},$$

we let  $\|\cdot\|_* = \|\cdot\|_{\mathcal{B}_D}$ , and we let  $\mathbb{W} = \mathbb{V}$ . We now proceed to verify that Assumptions 4.1 to 4.4 hold.

PROPOSITION 4.11 (coercivity). *Assumptions 4.1 and 4.2 are satisfied for the Dirichlet problem if  $\exists \beta_{\min} > 0$ , independent of  $h$ , such that  $\beta_D > \beta_{\min}$ .*

*Proof.* Using the fact that  $|v|_{H^{1/2}(\Gamma_D)}^2 + \|\bar{v}\|_{L^2(\Gamma_D)}^2 \gtrsim \|v\|_{H^{1/2}(\Gamma_D)}^2$ , we deduce from Lemma 2.5 that for every positive  $\alpha' \leq \alpha$ ,

$$\alpha' \|(v, \mu)\|_{\mathbb{V}}^2 - \alpha' \|\bar{v}\|_{L^2(\Gamma_D)}^2 \leq \mathcal{A}[(v, \mu), (v, \mu)] \quad \forall (v, \mu) \in \mathbb{W}.$$

Using the definition of  $\mathcal{B}_D$ , we see that

$$\mathcal{B}_D[(v, \mu), (v, \mu)] = \beta_D \langle v, v \rangle_{\Gamma_D} = \beta_D \|v\|_{L^2(\Gamma_D)}^2.$$

Taking  $\alpha' = \min(\alpha, \beta_{\min}/2)$ , we see that

$$\begin{aligned} \mathcal{A}[(v, \mu), (v, \mu)] + \mathcal{B}_D[(v, \mu), (v, \mu)] &\geq \alpha' \|(v, \mu)\|_{\mathbb{V}}^2 + \left(1 - \frac{\alpha'}{\beta_{\min}}\right) \beta_D \|v\|_{L^2(\Gamma_D)}^2 \\ &\geq \alpha'' \|(v, \mu)\|_{\mathcal{B}_D}^2 \end{aligned}$$

for some  $\alpha'' > 0$ . Therefore, in this case the form  $\mathcal{A} + \mathcal{B}_D$  is coercive, and so Assumptions 4.1 and 4.2 hold.  $\square$

PROPOSITION 4.12 (weak coercivity). *Assumptions 4.1 and 4.2 are satisfied for the Dirichlet problem with  $\beta_D = 0$ .*

*Proof.* Taking  $w = v$  and  $\eta = \mu + c\bar{v}$ , for some  $c \in \mathbb{R}$  to be fixed, we obtain

$$(4.12) \quad \begin{aligned} L &:= \mathcal{A}[(v, \mu), (w, \eta)] + \mathcal{B}_D[(v, \mu), (w, \eta)] \\ &= \langle \mathbf{V}\mu, \mu \rangle_{\Gamma} + c \langle \mathbf{V}\mu, \bar{v} \rangle_{\Gamma} - c \langle \mathbf{K}v, \bar{v} \rangle_{\Gamma} + \langle \mathbf{W}v, v \rangle_{\Gamma} + \frac{c}{2} \langle v, \bar{v} \rangle_{\Gamma}. \end{aligned}$$

By Lemmas 2.1 and 2.2, we know that

$$(4.13) \quad \langle \mathbf{V}\mu, \mu \rangle_{\Gamma} + \langle \mathbf{W}v, v \rangle_{\Gamma} \geq \alpha_V \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \alpha_W |v|_{H^{1/2}(\Gamma)}^2.$$

By Lemma 2.3, we see that

$$\begin{aligned} c |\langle \mathbf{V}\mu, \bar{v} \rangle_{\Gamma}| &\leq c \|\mathbf{V}\mu\|_{H^{1/2}(\Gamma)} \|\bar{v}\|_{H^{-1/2}(\Gamma)} \\ &\leq c C_V \|\mu\|_{H^{-1/2}(\Gamma)} \|\bar{v}\|_{H^{-1/2}(\Gamma)} \\ &= c C_V \|\mu\|_{H^{-1/2}(\Gamma)} \|\bar{v}\|_{L^2(\Gamma)}. \end{aligned}$$

Using the fact that for  $a, b \geq 0$ ,  $ab \leq (a^2 + b^2)/2$ , we obtain

$$(4.14) \quad c |\langle \mathbf{V}\mu, \bar{v} \rangle_{\Gamma}| \leq \frac{c^2 C_V^2}{2\alpha_V} \|\bar{v}\|_{L^2(\Gamma)}^2 + \frac{\alpha_V}{2} \|\mu\|_{H^{-1/2}(\Gamma)}^2.$$

We note that  $u = \bar{v}$  is a solution to (1.1),  $\gamma_D \bar{v} = \bar{v}$  and  $\gamma_N \bar{v} = 0$ . Using this and applying (2.14), we see that  $\forall \mu \in H^{-1/2}(\Gamma)$ ,  $\langle \mathbf{K}\bar{v}, \mu \rangle_\Gamma = -\frac{1}{2} \langle \bar{v}, \mu \rangle_\Gamma$ . Therefore, using  $\mu = \bar{v}$ ,

$$\begin{aligned} c \langle \mathbf{K}v, \bar{v} \rangle_\Gamma &= c \langle \mathbf{K}(v - \bar{v}), \bar{v} \rangle_\Gamma + c \langle \mathbf{K}\bar{v}, \bar{v} \rangle_\Gamma \\ &= c \langle \mathbf{K}(v - \bar{v}), \bar{v} \rangle_\Gamma - \frac{c}{2} \langle \bar{v}, \bar{v} \rangle_\Gamma. \end{aligned}$$

Using the fact that  $\|v - \bar{v}\|_{H^{1/2}(\Gamma)} = |v|_{H_*^{1/2}(\Gamma)}$ , and proceeding in the same way as we did for the single layer term above, we obtain

$$(4.15) \quad c \langle \mathbf{K}v, \bar{v} \rangle_\Gamma \leq \frac{\alpha_W}{2} |v|_{H_*^{1/2}(\Gamma)}^2 + \frac{C_K^2 c^2}{2\alpha_W} \|\bar{v}\|_{L^2(\Gamma)}^2 - \frac{c}{2} \|\bar{v}\|_{L^2(\Gamma)}^2.$$

We also have that

$$(4.16) \quad \frac{c}{2} \langle v, \bar{v} \rangle = \frac{c}{2} \|\bar{v}\|_{L^2(\Gamma)}^2.$$

Taking  $\alpha = \min(\alpha_V, \alpha_K)$  and  $C = \max(C_V, C_K)$ , and putting (4.13)–(4.16) together, we obtain

$$L \geq \frac{\alpha}{2} \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \frac{\alpha}{2} |v|_{H_*^{1/2}(\Gamma)}^2 + \left( c - \frac{c^2 C^2}{\alpha} \right) \|\bar{v}\|_{L^2(\Gamma)}^2.$$

Letting  $c = \frac{\alpha}{2C^2}$  gives

$$\begin{aligned} L &\geq \frac{\alpha}{2} \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \frac{\alpha}{2} |v|_{H_*^{1/2}(\Gamma)}^2 + \frac{\alpha}{4C^2} \|\bar{v}\|_{L^2(\Gamma)}^2 \\ &\gtrsim \|\mu\|_{H^{-1/2}(\Gamma)}^2 + |v|_{H_*^{1/2}(\Gamma)}^2 + \|\bar{v}\|_{L^2(\Gamma)}^2. \end{aligned}$$

Finally, we show that

$$\begin{aligned} \|(v, \mu)\|_{\mathbb{V}} &= \|v\|_{H^{1/2}(\Gamma)} + \|\mu\|_{H^{-1/2}(\Gamma)} \\ &\leq \|v - \bar{v}\|_{H^{1/2}(\Gamma)} + \|\bar{v}\|_{H^{1/2}(\Gamma)} + \|\mu\|_{H^{-1/2}(\Gamma)} \\ &= |v|_{H_*^{1/2}(\Gamma)} + \|\bar{v}\|_{L^2(\Gamma)} + \|\mu\|_{H^{-1/2}(\Gamma)}, \\ \|(w, \eta)\|_{\mathbb{V}} &\leq |v|_{H_*^{1/2}(\Gamma)} + \|\bar{v}\|_{L^2(\Gamma)} + \|\mu + c\bar{v}\|_{H^{-1/2}(\Gamma)} \\ &\leq |v|_{H_*^{1/2}(\Gamma)} + \|\bar{v}\|_{L^2(\Gamma)} + \|\mu\|_{H^{-1/2}(\Gamma)} + c\|\bar{v}\|_{H^{-1/2}(\Gamma)} \\ &\lesssim |v|_{H_*^{1/2}(\Gamma)} + \|\bar{v}\|_{L^2(\Gamma)} + \|\mu\|_{H^{-1/2}(\Gamma)}. \end{aligned}$$

Therefore

$$\begin{aligned} \|(v, \mu)\|_{\mathbb{V}} \|(w, \eta)\|_{\mathbb{V}} &\lesssim \|\mu\|_{H^{-1/2}(\Gamma)}^2 + |v|_{H_*^{1/2}(\Gamma)}^2 + \|\bar{v}\|_{L^2(\Gamma)}^2 \\ &\lesssim L. \end{aligned}$$

We obtain the first part of Assumption 4.1 by dividing through by  $\|(w, \eta)\|_{\mathbb{V}}$  and taking the supremum.

To show the second part of Assumption 4.1, we let  $(w, \eta) \in \mathbb{W} \setminus \{0\}$  and proceed as follows:

$$\begin{aligned} L &:= \sup_{(v, \mu) \in \mathbb{W}} |\mathcal{A}[(v, \mu), (w, \eta)] + \mathcal{B}_D[(v, \mu), (w, \eta)]| \\ &\geq \mathcal{A}[(w, \eta - \bar{w}), (w, \eta)] + \mathcal{B}_D[(w, \eta - \bar{w}), (w, \eta)] \\ &= -\langle \mathbf{K}'\bar{w}, w \rangle_\Gamma + \langle \mathbf{V}\eta, \eta \rangle_\Gamma - \langle \mathbf{V}\bar{w}, \eta \rangle_\Gamma + \langle \mathbf{W}w, w \rangle_\Gamma + \frac{1}{2} \langle \bar{w}, w \rangle_\Gamma. \end{aligned}$$

This is of the same form as (4.12), so we proceed as above to obtain

$$L \gtrsim \|(v, \mu)\|_{\mathbb{V}} \|(w, \eta)\|_{\mathbb{V}}.$$

This is greater than zero  $\forall (w, \eta) \neq 0$ , and so we have proven the second part of Assumption 4.1.

Assumption 4.2 can be proven in the same way as above using the discrete space  $\mathbb{V}_h$  in the place of  $\mathbb{W}$ .  $\square$

**PROPOSITION 4.13** (continuity). *Assumption 4.3 is satisfied for the Dirichlet problem.*

*Proof.* Applying Lemma 2.4, the relation

$$\langle \eta, v \rangle_{\Gamma} \leq \|\eta\|_{H^{-1/2}(\Gamma)} \|v\|_{H^{1/2}(\Gamma)},$$

and the Cauchy–Schwarz inequality,

$$\beta_{\mathbb{D}} \langle w, v \rangle_{\Gamma} \leq \beta_{\mathbb{D}}^{1/2} \|w\|_{L^2(\Gamma)} \beta_{\mathbb{D}}^{1/2} \|v\|_{L^2(\Gamma)},$$

to the form  $\mathcal{A} + \mathcal{B}_{\mathbb{D}}$  yields the desired continuity result.  $\square$

**PROPOSITION 4.14** (approximation). *Assumption 4.4 is satisfied for the Dirichlet problem if  $0 \leq \beta_{\mathbb{D}} \lesssim h^{-1}$ .*

*Proof.* Using standard approximation results (see, e.g., [23, Theorems 10.4 and 10.9]), we see that

$$\begin{aligned} \inf_{(w_h, \eta_h) \in \mathbb{V}_h} \|(v - w_h, \mu - \eta_h)\|_{\mathbb{V}} &= \inf_{w_h \in \mathbb{V}_h^k} \|v - w_h\|_{H^{1/2}(\Gamma)} + \inf_{\eta_h \in \Lambda_h^l} \|\mu - \eta_h\|_{H^{-1/2}(\Gamma)} \\ &\lesssim h^{\zeta-1/2} |v|_{H^{\zeta}(\Gamma)} + h^{\xi+1/2} |\mu|_{H^{\xi}(\Gamma)}, \\ \inf_{w_h \in \mathbb{V}_h^k} \|v - w_h\|_{L^2(\Gamma_{\mathbb{D}})} &\lesssim h^{\zeta} |v|_{H^{\zeta}(\Gamma)}. \end{aligned}$$

Applying these to the definition of  $\|\cdot\|_*$  gives

$$\inf_{(w_h, \eta_h) \in \mathbb{V}_h} \|(v - w_h, \mu - \eta_h)\|_* \lesssim h^{\zeta-1/2} |v|_{H^{\zeta}(\Gamma)} + h^{\xi+1/2} |\mu|_{H^{\xi}(\Gamma)} + \beta_{\mathbb{D}}^{1/2} h^{\zeta} |v|_{H^{\zeta}(\Gamma)}.$$

If  $\beta_{\mathbb{D}} = 0$ , Assumption 4.4 holds. If  $0 < \beta_{\mathbb{D}} \lesssim h^{-1}$ , then  $\beta_{\mathbb{D}}^{1/2} h^{\zeta} \lesssim h^{\zeta-1/2}$ , and so Assumption 4.4 holds.  $\square$

We have shown that Assumptions 4.1 to 4.4 are satisfied. Additionally the extra assumptions in Proposition 4.6 are satisfied, so we conclude that the results of Propositions 4.6, 4.7, and 4.9 and Corollaries 4.8 and 4.10 apply to the Dirichlet problem. This is summarized in the following result.

**THEOREM 4.15.** *The Dirichlet problem (3.8) has a unique solution  $(u, \lambda) \in H^s(\Gamma) \times H^r(\Gamma)$ , for some  $s \geq \frac{1}{2}$  and  $r \geq -\frac{1}{2}$ . The discrete Dirichlet problem (4.11) is invertible. If  $\exists \beta_{\min} > 0$  such that  $\beta_{\min} < \beta_{\mathbb{D}} \lesssim h^{-1}$  or  $\beta_{\mathbb{D}} = 0$ , its solution  $(u_h, \lambda_h) \in \mathbb{V}_h^k \times \Lambda_h^l$  satisfies*

$$\|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}_{\mathbb{D}}} \lesssim h^{\zeta-1/2} |u|_{H^{\zeta}(\Gamma)} + h^{\xi+1/2} |\lambda|_{H^{\xi}(\Gamma)},$$

where  $\zeta = \min(k + 1, s)$  and  $\xi = \min(l + 1, r)$ . Additionally,

$$\|\tilde{u} - \tilde{u}_h\|_{H^1(\Omega)} \lesssim h^{\zeta-1/2} |u|_{H^{\zeta}(\Gamma)} + h^{\xi+1/2} |\lambda|_{H^{\xi}(\Gamma)},$$

where  $\tilde{u}$  and  $\tilde{u}_h$  are the solutions in  $\Omega$  computed using (2.6).

**4.2. Application of the theory to the Neumann problem.** The Neumann problem takes the following form: Find  $(u_h, \lambda_h) \in \mathbb{V}_h^*$  such that

$$(4.17) \quad \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] + \mathcal{B}_N[(u_h, \lambda_h), (v_h, \mu_h)] = \mathcal{L}_N(v_h, \mu_h) \quad \forall (v_h, \mu_h) \in \mathbb{V}_h^*.$$

Here  $\mathbb{V}_h^* := \mathbb{V}_h^{*k}(\Gamma) \times \Lambda_h^l(\Gamma)$  and  $\mathbb{V}_h^{*k}(\Gamma) := \{v \in \mathbb{V}_h^k : \bar{v} = 0\}$ .

We introduce the  $\mathcal{B}_N$ -norm

$$\|(v, \mu)\|_{\mathcal{B}_N} := \|(v, \mu)\|_{\mathbb{V}} + \beta_N^{1/2} \|\mu\|_{L^2(\Gamma_N)},$$

we let  $\|\cdot\|_* = \|\cdot\|_{\mathcal{B}_N}$ , and we let  $\mathbb{W} = \mathbb{V}^*$ .

We now proceed to verify that Assumptions 4.1 to 4.4 hold.

**PROPOSITION 4.16** (coercivity). *Assumptions 4.1 and 4.2 are satisfied for the Neumann problem with  $\beta_N \geq 0$ .*

*Proof.* As  $v \in H_*^{1/2}(\Gamma_N)$ , we may immediately apply Lemmas 2.1 and 2.2 to show that the form is coercive. □

**PROPOSITION 4.17** (continuity). *Assumption 4.3 is satisfied for the Neumann problem.*

*Proof.* The proof is the same as in the Dirichlet case. □

**PROPOSITION 4.18** (approximation). *Assumption 4.4 is satisfied for the Neumann problem if  $0 \leq \beta_N \lesssim h$ .*

*Proof.* The proof is the same as in the Dirichlet case. □

As in the Dirichlet case, the extra assumptions in Proposition 4.6 are satisfied. We therefore conclude with the following result.

**THEOREM 4.19.** *The Neumann problem (3.13) has a unique solution  $(u, \lambda) \in H_*^s(\Gamma) \times H^r(\Gamma)$ , for some  $s \geq \frac{1}{2}$  and  $r \geq 0$  if  $\beta_N > 0$ . If  $\beta_N = 0$ , this holds for some  $r \geq -\frac{1}{2}$ . The discrete Neumann problem (4.17) is invertible. If  $0 \leq \beta_N \lesssim h$ , its solution  $(u_h, \lambda_h) \in \mathbb{V}_h^{*k} \times \Lambda_h^l$  satisfies*

$$\|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}_N} \lesssim h^{\zeta-1/2} |u|_{H^\zeta(\Gamma)} + h^{\xi+1/2} |\lambda|_{H^\xi(\Gamma)},$$

where  $\zeta = \min(k + 1, s)$  and  $\xi = \min(l + 1, r)$ . Additionally,

$$\|\tilde{u} - \tilde{u}_h\|_{H^1(\Omega)} \lesssim h^{\zeta-1/2} |u|_{H^\zeta(\Gamma)} + h^{\xi+1/2} |\lambda|_{H^\xi(\Gamma)},$$

where  $\tilde{u}$  and  $\tilde{u}_h$  are the solutions in  $\Omega$  computed using (2.6).

**4.3. Application of the theory to the mixed Dirichlet–Neumann problem.** For the mixed problem, the boundary element method takes the following form: Find  $(u_h, \lambda_h) \in \mathbb{V}_h$  such that

$$(4.18) \quad \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] + \mathcal{B}_{ND}[(u_h, \lambda_h), (v_h, \mu_h)] = \mathcal{L}_{ND}(v_h, \mu_h) \quad \forall (v_h, \mu_h) \in \mathbb{V}_h.$$

We now show that the assumptions for the abstract error estimate are satisfied for the formulation (4.18). First, we introduce the following norms:

$$\begin{aligned} \|(v, \mu)\|_{\mathcal{B}_{ND}} &:= \|(v, \mu)\|_{\mathbb{V}} + \beta_D^{1/2} \|v\|_{L^2(\Gamma_D)} + \beta_N^{1/2} \|\mu\|_{L^2(\Gamma_N)}, \\ \|(v, \mu)\|_* &:= \|(v, \mu)\|_{\mathbb{V}} + \beta_D^{1/2} \|v\|_{L^2(\Gamma)} + \beta_N^{1/2} \|\mu\|_{L^2(\Gamma)}. \end{aligned}$$

We let  $\mathbb{W} = H^{1/2}(\Gamma) \times L^2(\Gamma)$ .

Observe that in this case the two norms are not the same, nor are they equivalent, so the results below cannot be used to prove existence of a unique solution to (3.17). Nevertheless, it is easy to verify that if the exact solution to the mixed Dirichlet–Neumann problem is in  $\mathbb{V}$ , then it satisfies (3.17).

PROPOSITION 4.20 (coercivity). *Assumptions 4.1 and 4.2 are satisfied for the mixed Dirichlet–Neumann problem if  $\exists \beta_{\min} > 0$ , independent of  $h$ , such that  $\beta_D > \beta_{\min}$ .*

*Proof.* We obtain using Lemma 2.5 that for  $(v, \mu) \in \mathbb{W}$ ,

$$\begin{aligned} L &:= \mathcal{A}[(v, \mu), (v, \mu)] + \mathcal{B}_{\text{ND}}[(v, \mu), (v, \mu)] \\ &\geq \alpha \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \alpha |v|_{H_*^{1/2}(\Gamma)}^2 + \beta_D \|v\|_{L^2(\Gamma_D)}^2 + \beta_N \|\mu\|_{L^2(\Gamma_N)}^2. \end{aligned}$$

Taking  $\alpha' = \min(\alpha, \beta_{\min}/2)$ , we get

$$\begin{aligned} L &\geq \alpha' \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \alpha' \left( |v|_{H_*^{1/2}(\Gamma)}^2 + \|v\|_{L^2(\Gamma_D)}^2 \right) \\ &\quad + (\beta_D - \alpha') \|v\|_{L^2(\Gamma_D)}^2 + \beta_N \|\mu\|_{L^2(\Gamma_N)}^2. \end{aligned}$$

By [23, Theorem 2.6],  $(|\cdot|_{H_*^{1/2}(\Gamma)}^2 + \|\cdot\|_{L^2(\Gamma_D)}^2)^{1/2}$  is an equivalent norm to  $\|\cdot\|_{H^{1/2}(\Gamma)}$ . Therefore

$$\begin{aligned} L &\geq \alpha' \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \alpha' \|v\|_{H^{1/2}(\Gamma)}^2 + \beta_D \left( 1 - \frac{\alpha'}{\beta_{\min}} \right) \|v\|_{L^2(\Gamma_D)}^2 + \beta_N \|\mu\|_{L^2(\Gamma_N)}^2 \\ &\gtrsim \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \|v\|_{H^{1/2}(\Gamma)}^2 + \beta_D \|v\|_{L^2(\Gamma_D)}^2 + \beta_N \|\mu\|_{L^2(\Gamma_N)}^2. \end{aligned}$$

Coercivity follows using the definition of  $\|\cdot\|_{\mathcal{B}_{\text{ND}}}$ . □

PROPOSITION 4.21 (continuity). *Assumption 4.3 is satisfied for the mixed Dirichlet–Neumann problem if  $\exists \beta_{\min} > 0$ , independent of  $h$ , such that  $\beta_D^{1/2} \beta_N^{1/2} > \beta_{\min}$ .*

*Proof.* Using the fact that  $\langle v, \mu \rangle_{\Gamma} = \langle v, \mu \rangle_{\Gamma_D} + \langle v, \mu \rangle_{\Gamma_N}$ , we see that

$$\begin{aligned} \mathcal{B}_{\text{ND}}[(w, \eta), (v, \mu)] &= \frac{1}{2} \langle w, \mu \rangle_{\Gamma_D} - \frac{1}{2} \langle \eta, v \rangle_{\Gamma_D} + \beta_D \langle w, v \rangle_{\Gamma_D} \\ &\quad + \frac{1}{2} \langle \eta, v \rangle_{\Gamma_N} - \frac{1}{2} \langle w, \mu \rangle_{\Gamma_N} + \beta_N \langle \eta, \mu \rangle_{\Gamma_N} \\ &= \frac{1}{2} \langle w, \mu \rangle_{\Gamma} - \langle \eta, v \rangle_{\Gamma_D} + \beta_D \langle w, v \rangle_{\Gamma_D} \\ &\quad + \frac{1}{2} \langle \eta, v \rangle_{\Gamma} - \langle w, \mu \rangle_{\Gamma_N} + \beta_N \langle \eta, \mu \rangle_{\Gamma_N} \\ &= \frac{1}{2} \langle w, \mu \rangle_{\Gamma} - \beta_D^{1/2} \beta_D^{-1/2} \langle \eta, v \rangle_{\Gamma_D} + \beta_D \langle w, v \rangle_{\Gamma_D} \\ &\quad + \frac{1}{2} \langle \eta, v \rangle_{\Gamma} - \beta_N^{-1/2} \beta_N^{1/2} \langle w, \mu \rangle_{\Gamma_N} + \beta_N \langle \eta, \mu \rangle_{\Gamma_N}. \end{aligned}$$

Proceeding as in Proposition 4.13 leads to the desired result. □

PROPOSITION 4.22 (approximation). *Assumption 4.4 is satisfied for the mixed Dirichlet–Neumann problem if  $0 < \beta_D \lesssim h^{-1}$  and  $0 < \beta_N \lesssim h$ .*

*Proof.* Proceeding as in the Dirichlet case, we see that

$$\begin{aligned} \inf_{(w_h, \eta_h) \in \mathbb{V}_h} \|(v - w_h, \mu - \eta_h)\|_* &\lesssim h^{\zeta-1/2} |v|_{H^{\zeta}(\Gamma)} + h^{\xi+1/2} |\mu|_{H^{\xi}(\Gamma)} \\ &\quad + \beta_D^{1/2} h^{\zeta} |v|_{H^{\zeta}(\Gamma)} + \beta_N^{1/2} h^{\xi} |\mu|_{H^{\xi}(\Gamma)}. \end{aligned}$$

If  $0 < \beta_D \lesssim h^{-1}$  and  $0 < \beta_N \lesssim h$ , then



$$\beta_D^{1/2} h^\zeta |v|_{H^\zeta(\Gamma)} + \beta_N^{1/2} h^\xi |\mu|_{H^\xi(\Gamma)} \lesssim h^{\zeta-1/2} |v|_{H^\zeta(\Gamma)} + h^{\xi+1/2} |\mu|_{H^\xi(\Gamma)},$$

and so Assumption 4.4 holds. □

Motivated by the bounds on  $\beta_D$  and  $\beta_N$  in this proposition, we will later take  $\beta_D = \beta h^{-1}$  and  $\beta_N = \beta h$ , where  $\beta$  is a constant.

If  $k = l$ ,  $\beta_N \lesssim h^{-1}$ , and the solution is smooth enough, then

$$\beta_N^{1/2} h^\xi = \beta_N^{1/2} h^\zeta \lesssim h^{\zeta-1/2}.$$

Therefore the same order of convergence will be observed when the bounds on  $\beta_N$  here and in the theorem below may be replaced by  $\beta_N \lesssim h^{-1}$  without loss of convergence. In this case, both  $\beta_N$  and  $\beta_D$  may be taken to be constants independent of  $h$ .

We conclude that the best approximation result of Proposition 4.7 and the error estimate of Corollary 4.8 hold for the discrete solutions of (4.18), as given in the following theorem.

**THEOREM 4.23.** *Let  $(u, \lambda) \in H^s(\Gamma) \times H^r(\Gamma)$ , for some  $s \geq \frac{1}{2}$  and  $r \geq 0$ , be the unique solution to the mixed Dirichlet–Neumann problem. This solution satisfies (3.17). Let  $(u_h, \lambda_h) \in V_h^k \times \Lambda_h^l$  be the solution of (4.18). If  $0 < \beta_D \lesssim h^{-1}$ ,  $0 < \beta_N \lesssim h$ , and  $\exists \beta_{\min} > 0$  such that  $\beta_D^{1/2} \beta_N^{1/2} > \beta_{\min}$  and  $\beta_D > \beta_{\min}$ , then*

$$\|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}_{ND}} \lesssim h^{\zeta-1/2} |u|_{H^\zeta(\Gamma)} + h^{\xi+1/2} |\lambda|_{H^\xi(\Gamma)},$$

where  $\zeta = \min(k + 1, s)$  and  $\xi = \min(l + 1, r)$ .

If we set  $\beta_D = 0$  and  $\beta_N = 0$ , we arrive at a penalty-free formulation for the mixed Dirichlet–Neumann problem. We conjecture based on numerical experiments that this result also holds for the penalty-free formulation. The analysis for this case would take a similar form as in the Dirichlet and Neumann penalty-free cases.

**4.4. Application of the theory to the Robin problem.** The formulation for Robin conditions was proposed in (3.24). To simplify the notation we introduce a function  $\omega : \Gamma \rightarrow \mathbb{R}_+$  defined by

$$\omega(\mathbf{x}) := \frac{1}{\varepsilon(\mathbf{x})\beta_R(\mathbf{x}) + 1},$$

and we assume that  $\varepsilon$  and  $\beta_R$  are sufficiently regular so that

$$(4.19) \quad \omega \in W^{1,2}(\Gamma) \cap L^\infty(\Gamma).$$

This will be true if the mesh has some local quasi uniformity and  $\varepsilon$  is smooth enough. Noting that

$$\omega - \frac{1}{2} = \frac{2 - (\varepsilon\beta_R + 1)}{2(\varepsilon\beta_R + 1)} = -\frac{1}{2} \frac{\varepsilon\beta_R - 1}{\varepsilon\beta_R + 1},$$

we may then write the operators  $\mathcal{B}_R$  and  $\mathcal{L}_R$  as

$$(4.20)$$

$$\mathcal{B}_R[(u, \lambda), (v, \mu)] = \langle (\omega - \frac{1}{2})u, \mu \rangle_{\Gamma_R} - \langle (\omega - \frac{1}{2})\lambda, v \rangle_{\Gamma_R} + \langle \omega\beta_R u, v \rangle_{\Gamma_R} + \langle \omega\varepsilon\lambda, \mu \rangle_{\Gamma_R},$$

$$(4.21) \quad \mathcal{L}_R[(v, \mu)] = \langle (g_D + \varepsilon g_N)\omega, \beta_R v + \mu \rangle_{\Gamma_R}.$$

The boundary element method for the Robin problem reads as follows: Find  $(u_h, \lambda_h) \in \mathbb{V}_h$  such that

$$(4.22) \quad \mathcal{A}[(u_h, \lambda_h), (v_h, \mu_h)] + \mathcal{B}_R[(u_h, \lambda_h), (v_h, \mu_h)] = \mathcal{L}_R[(v_h, \mu_h)] \quad \forall (v_h, \mu_h) \in \mathbb{V}_h.$$

For the analysis the following technical lemmas will be useful.

LEMMA 4.24. *If  $\varphi \in W^{1,2}(\Gamma) \cap L^\infty(\Gamma)$  and  $f \in H^{1/2}(\Gamma)$ , then  $\varphi f \in H^{1/2}(\Gamma)$  and*

$$\|\varphi f\|_{H^{1/2}(\Gamma)} \leq C (\|\varphi\|_{L^\infty(\Gamma)} + \|\varphi\|_{W^{1,2}(\Gamma)}) \|f\|_{H^{1/2}(\Gamma)}.$$

*Proof.* The proof is a consequence of [7, Lemma 6] which shows that

$$(4.23) \quad \|\varphi f\|_{H^{1/2}(\Gamma)} \leq C \left( \|\varphi\|_{L^\infty(\Gamma)} \|f\|_{H^{1/2}(\Gamma)} + \|f\|_{L^4(\Gamma)} \|\varphi\|_{W^{1,2}(\Gamma)}^{1/2} \|\varphi\|_{L^\infty(\Gamma)}^{1/2} \right).$$

We then recall the Sobolev injection  $\|f\|_{L^4(\Gamma)} \leq C \|f\|_{H^{1/2}(\Gamma)}$  from [12, Theorem 6.7] and conclude using this result and an arithmetic-geometric inequality of the right-hand side of (4.23).  $\square$

LEMMA 4.25. *If  $\varphi, f \in L^2(\Gamma)$  and  $\varphi(\mathbf{x}) > 0 \forall \mathbf{x} \in \Gamma$ , then there exists  $C > 0$  such that*

$$\|\varphi f\|_{L^2(\Gamma)}^2 \geq C \|f\|_{L^2(\Gamma)}^2.$$

*Proof.* Let  $a = \inf_{\mathbf{x} \in \Gamma} \varphi(\mathbf{x})$ . Since  $\Gamma$  is closed, there exists  $\mathbf{y} \in \Gamma$  such that  $\varphi(\mathbf{y}) = a$ . Therefore  $a > 0$ . We now see that

$$\begin{aligned} \|\varphi f\|_{L^2(\Gamma)}^2 &= \int_{\Gamma} \varphi^2 f^2 \\ &\geq a^2 \int_{\Gamma} f^2 \\ &= C \|f\|_{L^2(\Gamma)}^2, \end{aligned}$$

where  $C = a^2$ .  $\square$

We introduce the norm

$$\|(v, \mu)\|_{\mathcal{B}_R} := \|(v, \mu)\|_{\mathbb{V}} + \|(\varepsilon\omega)^{1/2}\mu\|_{L^2(\Gamma)} + \|(\omega\beta_R)^{1/2}v\|_{L^2(\Gamma)},$$

we let  $\|\cdot\|_* = \|\cdot\|_{\mathcal{B}_R}$ , and we let  $\mathbb{W} = H^{1/2}(\Gamma) \times L^2(\Gamma)$ . We note that if  $\varepsilon \rightarrow 0$  or  $\varepsilon \rightarrow \infty$ , then  $\|\cdot\|_{\mathcal{B}_R}$  converges to  $\|\cdot\|_{\mathcal{B}_D}$  or  $\|\cdot\|_{\mathcal{B}_N}$ , respectively. We now proceed to show that Assumptions 4.1 to 4.4 hold.

PROPOSITION 4.26 (coercivity). *Assumptions 4.1 and 4.2 are satisfied for the Robin problem.*

*Proof.* Let  $(v, \mu) \in \mathbb{W}$ , and let  $L := \mathcal{A}[(v, \mu), (v, \mu)] + \mathcal{B}_R[(v, \mu), (v, \mu)]$ . Using Lemma 2.5, we see that

$$\begin{aligned} L &\geq \alpha \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \alpha \|v\|_{H^{1/2}(\Gamma)}^2 - \alpha \|v\|_{L^2(\Gamma)}^2 \\ &\quad + \|(\varepsilon\omega)^{1/2}\mu\|_{L^2(\Gamma)}^2 + \|(\omega\beta_R)^{1/2}v\|_{L^2(\Gamma)}^2 \end{aligned}$$

for any  $\alpha \leq \min(\alpha_V, \alpha_W)$ .

By Lemma 4.25, we have

$$(4.24) \quad -\alpha \|v\|_{L^2(\Gamma)}^2 \geq -\frac{\alpha}{C} \|(\omega\beta_R)^{1/2}v\|_{L^2(\Gamma)}^2.$$

Taking  $\alpha = \min(\alpha_V, \alpha_W, C/2)$ , we obtain

$$L \geq \alpha \|\mu\|_{H^{-1/2}(\Gamma)}^2 + \alpha \|v\|_{H^{1/2}(\Gamma)}^2 + \|(\varepsilon\omega)^{1/2}\mu\|_{L^2(\Gamma)}^2 + \frac{1}{2} \|(\omega\beta_R)^{1/2}v\|_{L^2(\Gamma)}^2.$$

Using the definition of  $\|\cdot\|_{\mathcal{B}_R}$ , we see that the form is coercive.  $\square$

PROPOSITION 4.27 (continuity). *Assumption 4.3 is satisfied for the Robin problem if  $\exists \beta_{\min} > 0$ , independent of  $h$ , such that  $\beta_{\mathbb{R}} > \beta_{\min}$ .*

*Proof.* Using Lemma 4.24 and the fact that  $\omega \in W^{1,2}(\Gamma) \cap L^\infty(\Gamma)$ , we see that for  $g \in H^{-1/2}(\Gamma)$  and  $f \in H^{1/2}(\Gamma)$ ,

$$\langle \omega g, f \rangle_\Gamma \leq C (\|\omega\|_{L^\infty(\Gamma)} + \|\omega\|_{W^{1,2}(\Gamma)}) \|g\|_{H^{-1/2}(\Gamma)} \|f\|_{H^{1/2}(\Gamma)}.$$

Let  $\varepsilon_{\min} := \inf_{\mathbf{x} \in \Gamma} \varepsilon(\mathbf{x})$ . As in the proof of Lemma 4.25, we see that  $\varepsilon_{\min} > 0$ . Hence,

$$-\frac{1}{2} < \omega - \frac{1}{2} < \frac{1}{\beta_{\min} \varepsilon_{\min} + 1},$$

and so

$$\|\omega - \frac{1}{2}\|_{L^\infty(\Gamma)} + \|\omega - \frac{1}{2}\|_{W^{1,2}(\Gamma)} < \max\left(\frac{1}{2}, \frac{1}{\beta_{\min} \varepsilon_{\min} + 1}\right) (\|1\|_{L^\infty(\Gamma)} + \|1\|_{W^{1,2}(\Gamma)}).$$

Applying these two results to the first two boundary terms in  $\mathcal{B}_{\mathbb{R}}[(w, \eta), (v, \mu)]$ , we obtain

$$\langle (\omega - \frac{1}{2})w, \mu \rangle_\Gamma - \langle (\omega - \frac{1}{2})v, \eta \rangle_\Gamma \leq C \|(w, \eta)\|_{\mathbb{V}} \|(v, \mu)\|_{\mathbb{V}}.$$

By the Cauchy–Schwarz inequality, we obtain for the remaining terms

$$\begin{aligned} & \langle \omega \varepsilon \eta, \mu \rangle_\Gamma + \langle \omega \beta_{\mathbb{R}} w, v \rangle_\Gamma \\ & \leq \|(\omega \varepsilon)^{1/2} \eta\|_{L^2(\Gamma)} \|(\omega \varepsilon)^{1/2} \mu\|_{L^2(\Gamma)} + \|(\omega \beta_{\mathbb{R}})^{1/2} w\|_{L^2(\Gamma)} \|(\omega \beta_{\mathbb{R}})^{1/2} v\|_{L^2(\Gamma)}. \end{aligned}$$

Collecting the terms, we then have

$$\mathcal{B}_{\mathbb{R}}[(w, \eta), (v, \mu)] \lesssim \|(w, \eta)\|_{\mathcal{B}_{\mathbb{R}}} \|(v, \mu)\|_{\mathcal{B}_{\mathbb{R}}}. \quad \square$$

PROPOSITION 4.28 (approximation). *Assumption 4.4 is satisfied for the Robin problem if  $\beta_{\mathbb{R}} \approx h^{-1}$ .*

*Proof.* First note that  $\omega < 1$  and

$$\omega \varepsilon = \frac{\varepsilon}{\varepsilon \beta_{\mathbb{R}} + 1} = \frac{1}{\beta_{\mathbb{R}} + \frac{1}{\varepsilon}} < \frac{1}{\beta_{\mathbb{R}}}.$$

Therefore,

$$(4.25) \quad \|(\omega \beta_{\mathbb{R}})^{1/2} v\|_{L^2(\Gamma)} \leq \beta_{\mathbb{R}}^{1/2} \|v\|_{L^2(\Gamma)} \quad \text{and} \quad \|(\omega \varepsilon)^{1/2} \mu\|_{L^2(\Gamma)} \leq \beta_{\mathbb{R}}^{-1/2} \|\mu\|_{L^2(\Gamma)}.$$

If  $\beta_{\mathbb{R}} \approx h^{-1}$ , then Assumption 4.4 can be shown to hold. □

When using equal order approximation, the same order of convergence will be observed when the bounds on  $\beta_{\mathbb{R}}$  here and in the theorem below may be replaced by  $\beta_{\min} \lesssim \beta_{\mathbb{R}} \lesssim h^{-1}$  for sufficiently smooth solutions.

PROPOSITION 4.29. *The extra assumptions in Proposition 4.6 are satisfied for the Robin problem.*

*Proof.* As a consequence of the coercivity and continuity above and observing that by the Cauchy–Schwarz inequality and the definition of  $\omega$ , there exists  $C$  such that

$$\langle \omega(g_{\mathbb{D}} + \varepsilon g_{\mathbb{N}}), \beta_{\mathbb{R}} v + \mu \rangle_\Gamma \leq C (\|g_{\mathbb{D}}\|_{L^2(\Gamma)} + \|g_{\mathbb{N}}\|_{L^2(\Gamma)}) \|(v, \mu)\|_{\mathcal{B}_{\mathbb{R}}}. \quad \square$$

We conclude that Propositions 4.6 and 4.7 and Corollaries 4.8 and 4.10 hold for the Robin problem. This is summarized in the following result.

**THEOREM 4.30.** *The Robin problem (3.24) has a unique solution  $(u, \lambda) \in H^s(\Gamma) \times H^r(\Gamma)$  for some  $s \geq \frac{1}{2}$  and  $r \geq 0$ . The discrete Robin problem (4.22) is invertible. If  $\beta_R \approx h^{-1}$ , its solution  $(u_h, \lambda_h) \in V_h^k \times \Lambda_h^l$  satisfies*

$$\|(u - u_h, \lambda - \lambda_h)\|_{\mathcal{B}_R} \leq C \left( h^{\zeta-1/2} |u|_{H^\zeta(\Gamma)} + h^{\xi+1/2} |\lambda|_{H^\xi(\Gamma)} \right)$$

for some  $C > 0$ , where  $\zeta = \min(k+1, s)$  and  $\xi = \min(l+1, r)$ . Additionally,

$$\|\tilde{u} - \tilde{u}_h\|_{H^1(\Omega)} \leq C \left( h^{\zeta-1/2} |u|_{H^\zeta(\Gamma)} + h^{\xi+1/2} |\lambda|_{H^\xi(\Gamma)} \right),$$

where  $\tilde{u}$  and  $\tilde{u}_h$  are the solutions in  $\Omega$  computed using (2.6).

Again, we could set  $\beta_R = 0$  to arrive at a penalty-free formulation for Robin problems. In this case, our numerical experiments show large errors for some values of the parameter  $\varepsilon$ , which leads us to conclude that this result does not hold for the penalty-free formulation.

As  $\varepsilon \rightarrow 0$  and  $\varepsilon \rightarrow \infty$ , we obtain the Dirichlet and Neumann formulations analyzed in subsections 4.1 and 4.2. We expect the condition number of the discrete system for the Robin problem to be no worse than in either extreme case, and we observe this in subsection 5.3.

**5. Numerical results.** Drawing inspiration from [16], we define

$$\begin{aligned} u(x, y, z) &= \sin(\pi x) \sin(\pi y) \sinh(\sqrt{2}\pi z), \\ g_D(x, y, z) &= \sin(\pi x) \sin(\pi y) \sinh(\sqrt{2}\pi z), \\ g_N(x, y, z) &= \begin{pmatrix} \pi \cos(\pi x) \sin(\pi y) \sinh(\sqrt{2}\pi z) \\ \pi \sin(\pi x) \cos(\pi y) \sinh(\sqrt{2}\pi z) \\ \sqrt{2}\pi \sin(\pi x) \sin(\pi y) \cosh(\sqrt{2}\pi z) \end{pmatrix} \cdot \boldsymbol{\nu}. \end{aligned}$$

It is easy to check that for any bounded domain  $\Omega \subset \mathbb{R}^3$  with boundary  $\Gamma = \Gamma_D \cup \Gamma_N \cup \Gamma_R$  and any fixed  $\varepsilon \in \mathbb{R}$ ,  $u$  is the solution of

$$\begin{aligned} (5.1a) \quad & -\Delta u = 0 && \text{in } \Omega, \\ (5.1b) \quad & u = g_D && \text{on } \Gamma_D, \\ (5.1c) \quad & \frac{\partial u}{\partial \boldsymbol{\nu}} = g_N && \text{on } \Gamma_N, \\ (5.1d) \quad & \frac{\partial u}{\partial \boldsymbol{\nu}} = \frac{1}{\varepsilon}(u - g_D) + g_N && \text{on } \Gamma_R. \end{aligned}$$

In the examples presented here, we let  $\Omega$  be the unit sphere and  $\Gamma$  its boundary. In the computations presented, a series of approximations of the sphere by plane triangles are used. The results in this section were computed using the boundary element library Bempp [22], an open source boundary element library developed by the authors of this paper. All examples in this paper were computed with version 3.3.2 of the Bempp library. Jupyter notebooks demonstrating the functionality used in this paper will be made available at [www.bempp.com](http://www.bempp.com).

**5.1. Dirichlet boundary conditions.** First, we look at the case where  $\Gamma = \Gamma_D$ , in which the problem reduces to the Dirichlet problem:

$$(5.2a) \quad -\Delta u = 0 \quad \text{in } \Omega,$$

$$(5.2b) \quad u = g_D \quad \text{on } \Gamma.$$

For this problem, we compare the penalty method proposed in this paper (4.11) to the standard single layer formulation: Find  $\lambda \in \Lambda_h$  such that

$$(5.3) \quad \langle \nabla \lambda, \mu \rangle = \langle (\frac{1}{2} \text{Id} + \mathbf{K}) g_D, \mu \rangle \quad \forall \mu \in \Lambda_h.$$

Figure 1 shows the convergence and iteration counts when  $\beta_D = 0.1$  and  $k = l = 1$ , and so we look for  $(u_h, \lambda_h) \in V_h^1 \times \tilde{\Lambda}_h^1$ . We note that as  $h$  decreases,  $h^{-1}$  increases, so  $0.1 \lesssim h^{-1}$ . In this case,  $\Gamma$  is smooth, and so  $V_h^1 = \tilde{\Lambda}_h^1$ . The iteration count plot (right) shows the number of iterations taken to solve the nonpreconditioned system (red diamonds), compared with the system with mass matrix preconditioning applied blockwise from the left (red circles), as described in [6]. Mass matrix preconditioning greatly reduces the number of iterations required, so for the remainder of this paper, we precondition all linear systems using mass matrix preconditioning.

For larger and more complex geometries, however, more specialized preconditioners are required. With systems of boundary element equations, it is common to use operator preconditioning or Calderón preconditioning [8], where properties of the boundary operators at the continuous level are used to derive a preconditioned equation of a form known to be well conditioned. In our case, it is not clear how to apply this approach, although further investigation of this warrants future work.

An alternative avenue of investigation leads to hierarchical LU based preconditioners, or even direct solvers of this type [4]. The penalty terms in this paper are all sparse matrices that have nonzero entries only for neighboring triangles, and so adding these terms only affects the entries in the matrix arising from near interactions; the far interactions—which are exactly those that are approximated in a hierarchical




FIG. 1. The convergence (left) and GMRES iteration counts (right) of the penalty method with  $\beta_D = 0.1$  (red circles) compared to the standard single layer method (5.3) (gray triangles), for the Dirichlet problem on the unit sphere, with  $k = l = 1$ . The iteration count plot shows the number of iterations taken to solve the mass matrix preconditioned system (red circles) and the nonpreconditioned system (red diamonds). The dashed line shows order 2 convergence.




FIG. 2. The dependence of the error (left) and iteration count (right) on the value of  $\beta_D$  for  $h = 2^{-2}$  (red triangles),  $h = 2^{-3.5}$  (red diamonds), and  $h = 2^{-5}$  (red pentagons), for the Dirichlet problem on the unit sphere, with  $k = l = 1$ .

matrix compression—are not affected by these terms. Therefore H-matrix methods can be applied to this method with few algorithmic changes required.

Figure 2 shows the dependence of the error and iteration count on the chosen value of  $\beta_D$ , for a range of values of  $h$ . It can be seen that the number of iterations increases when  $\beta_D$  is above around 0.1, and the error increases when  $\beta_D$  is above 100. This motivates our earlier choice of 0.1 as the value of  $\beta_D$ , although anything smaller than this appears to be a good choice of  $\beta_D$ .

In Figure 1, it can be seen that the penalty method proposed here gives comparable convergence to the standard method in a similar number of iterations. However, the system in the penalty method contains around twice the number of unknowns, and so each iteration will be more expensive.

Additionally, the discrete systems for the penalty method are nonsymmetric, so are solved using GMRES [21]. The discrete systems for the standard method (5.3) are symmetric, so CG [14] or MINRES [20] could be used: these methods are typically less expensive than GMRES, so this is a further disadvantage of the penalty method for pure Dirichlet and Neumann problems and justifies our focus on more complex boundary conditions.

**5.2. Mixed Dirichlet–Neumann boundary conditions.** We now consider the case where  $\Gamma = \Gamma_D \cup \Gamma_N$  and the problem reduces to a mixed Dirichlet–Neumann problem:

$$(5.4a) \quad -\Delta u = 0 \quad \text{in } \Omega,$$

$$(5.4b) \quad u = g_D \quad \text{on } \Gamma_D,$$

$$(5.4c) \quad \frac{\partial u}{\partial \nu} = g_N \quad \text{on } \Gamma_N.$$

Let  $\Gamma_N := \{(x, y, z) \in \Gamma : x > 0\}$  and  $\Gamma_D := \Gamma \setminus \Gamma_N$ . We use the same  $g_D$  and  $g_N$  as above.

We compare the method proposed in this paper with the standard method for mixed Dirichlet–Neumann problems [25, equation (3.2)]: Find  $(u, \lambda) \in \hat{H}^{1/2}(\Gamma_N) \times \hat{H}^{-1/2}(\Gamma_D)$  such that




FIG. 3. The dependence of the error (left) and iteration count (right) on the value of  $\beta$  for  $h = 2^{-2}$  (blue triangles),  $h = 2^{-3.5}$  (blue diamonds), and  $h = 2^{-5}$  (blue pentagons), for the mixed Dirichlet–Neumann problem on the unit sphere, with  $k = l + 1 = 1$ . Here we use  $\beta_D = \beta h^{-1}$  and  $\beta_N = \beta h$ .

$$\begin{aligned}
 (5.5) \quad & \langle W_{NN}u, v \rangle + \langle K'_{DN}, v \rangle - \langle K_{ND}u, \mu \rangle + \langle V_{DD}\lambda, \mu \rangle \\
 & = - \langle W_{DN}g_D, v \rangle + \langle (\tfrac{1}{2}\text{Id} - K'_{NN})g_N, v \rangle + \langle (\tfrac{1}{2}\text{Id} + K_{DD})g_D, \mu \rangle - \langle V_{ND}, \mu \rangle \\
 & \quad \forall (v, \mu) \in \tilde{H}^{1/2}(\Gamma_N) \times \tilde{H}^{-1/2}(\Gamma_D),
 \end{aligned}$$

where for a given boundary operator  $B$ ,  $B_{ij}$  is the corresponding boundary operator with the integral taken over  $\Gamma_i$  and the point  $\mathbf{x} \in \Gamma_j$ . For example,  $V_{ND}$  is defined by

$$(5.6) \quad [V_{ND}f](\mathbf{x}) := \int_{\Gamma_N} f(\mathbf{y})G(\mathbf{x}, \mathbf{y}) \, d\mathbf{y} \quad \text{for } \mathbf{x} \in \Gamma_D.$$

We first let  $k = l + 1 = 1$  and so look for  $(u_h, \lambda_h) \in V_h^1 \times \Lambda_h^0$ . As motivated above by Proposition 4.22, we set  $\beta_D = \beta h^{-1}$  and  $\beta_N = \beta h$ , where  $\beta$  is a constant. The dependence of the error and iteration count on  $\beta$  is shown in Figure 3. We observe that  $\beta = 0.01$  is a good choice, as this gives a small error and iteration count.

Theorem 4.23 says that taking  $\beta$  to be constant and  $\beta_D = \beta h^{-1}$  and  $\beta_N = \beta h$  will lead to convergence. The results in Figure 3 appear to contradict this, as they show that for smaller values of  $h$ , smaller values of  $\beta$  are required to obtain a small error. The fast increase in the number of iterations required to solve the linear system suggests that this discrepancy is in fact due to the increased ill-conditioning of the system as  $\beta$  is increased, with the ill-conditioning becoming apparent at lower values of  $\beta$  as  $h$  is reduced. This is confirmed by our experiments in which we ran the same problem as in Figure 3 but with a smaller GMRES tolerance: this decreased the error for  $h = 2^{-5}$  and  $\beta = 10$  from 144 to 0.909.

The convergence of the error as we reduce  $h$  is shown in Figure 4. Here we observe order 1.5 convergence, and the same rate of convergence as the standard method (5.5), with a marginally lower error in the standard method. The iteration count for the penalty method increases more gradually than the standard method, although this issue could be removed through better preconditioning of the standard method.

We next consider the case where  $k = l = 1$ . In this case, as remarked in subsection 4.3, we may replace the bound on  $\beta_N$  by  $\beta_N \lesssim h^{-1}$ , and so we may take both




FIG. 4. The convergence (left) and iterations counts (right) of the penalty method with  $\beta = 0.01$  (blue squares) compared to the standard method (5.5) (gray triangles), for the mixed Dirichlet–Neumann problem on the unit sphere, with  $k = l + 1 = 1$ . The dashed line shows order 1.5 convergence. Here we use  $\beta_{\text{D}} = \beta h^{-1}$  and  $\beta_{\text{N}} = \beta h$ .




FIG. 5. The dependence of the error (left) and iteration count (right) on the value of  $\beta$  for  $h = 2^{-2}$  (blue triangles),  $h = 2^{-3.5}$  (blue diamonds), and  $h = 2^{-5}$  (blue pentagons), for the mixed Dirichlet–Neumann problem on the unit sphere, with  $k = l = 1$ . Here we use  $\beta_{\text{D}} = \beta_{\text{N}} = \beta$ .

$\beta_{\text{D}}$  and  $\beta_{\text{N}}$  to be constant: we set  $\beta_{\text{D}} = \beta_{\text{N}} = \beta$ . The dependence of the error and iteration count on  $\beta$  for this choice of parameters is shown in Figure 5.

The convergence to the solution when  $\beta = 0.01$  is shown in Figure 6. It can be seen here that order 2 convergence is observed, higher than the expected order 1.5 convergence. In this case, the standard method (5.5) only achieves order 1 convergence, with a much higher iteration count than the penalty method. For this choice of discrete spaces, we also compared our method with the formulation given in [11, equation (1.19)]: this formulation is better conditioned than (5.5) but still achieves only order 1 convergence.

In Figures 3 and 6, the error and iteration count remain steady as  $\beta \rightarrow 0$ . In numerical experiments on a sphere and cube with  $\beta = 0$ , we see convergence similar to that observed in this section. This leads us to conjecture that Theorem 4.23 will hold for the penalty-free formulation, when  $\beta = 0$ .

**5.3. Robin problem.** We now consider the case where  $\Gamma = \Gamma_{\text{R}}$  and the problem reduces to a Robin problem:






FIG. 6. The convergence (left) and iterations counts (right) of the penalty method with  $\beta = 0.01$  (blue circles) compared to the standard method (5.5) (gray triangles) and the method given in [11, equation (1.19)] (gray diamonds), for the mixed Dirichlet–Neumann problem on the unit sphere, with  $k = l = 1$ . The dashed lines show order 2 and order 1 convergence. Here we use  $\beta_D = \beta_N = \beta$ .

$$(5.7a) \quad -\Delta u = 0 \quad \text{in } \Omega,$$

$$(5.7b) \quad \frac{\partial u}{\partial \nu} = \frac{1}{\varepsilon}(u - g_D) + g_N \quad \text{on } \Gamma,$$

for some  $\varepsilon \in \mathbb{R}$ .

In this section, we compare the method proposed in this paper with the following method: Find  $u \in H^{1/2}(\Gamma)$  such that

$$(5.8) \quad \langle Wu, v \rangle + \left\langle \frac{1}{\varepsilon} \left( \frac{1}{2} \text{Id} - K' \right) u, v \right\rangle = \left\langle \left( \frac{1}{2} \text{Id} - K' \right) \left( \frac{1}{\varepsilon} g_D + g_N \right), v \right\rangle \quad \forall v \in H^{1/2}(\Gamma).$$

Again, we begin letting  $k = l + 1 = 1$ . Here we use

$$\beta_R := \frac{\varepsilon \beta_N + \beta_D}{\varepsilon + 1},$$

where  $\beta_D = \beta h^{-1}$  and  $\beta_N = \beta h$ , for some constant  $\beta$ , as in the mixed Dirichlet–Neumann case.

The dependence of the error and iteration count on both  $\varepsilon$  and  $\beta$ , on a grid with  $h = 0.1$ , is shown in Figure 7. The convergence as  $h$  is reduced for  $\varepsilon = \frac{1}{300}$ ,  $\varepsilon = 1$ , and  $\varepsilon = 300$ , and using  $\beta = 0.01$ , is shown in Figure 8. In this case, order 1.5 convergence is observed.

As in the mixed Dirichlet–Neumann case, when  $k = l = 1$ , we may replace the bound on  $\beta_N$  with  $\beta_N \lesssim h^{-1}$ . Again, we take  $\beta_D = \beta_N = \beta$  for some constant  $\beta$ . The dependence of the error and iteration count on both  $\beta$  and  $\varepsilon$  is shown in Figure 9. As in the previous case,  $\beta = 0.01$  looks to be a suitable choice for the parameter.

The convergence as we reduce  $h$  for  $\varepsilon = \frac{1}{300}$ ,  $\varepsilon = 1$ , and  $\varepsilon = 300$ , and using  $\beta = 0.01$ , is shown in Figure 10. In this case, order 2 convergence is observed. For the method (5.8), the same order of convergence and errors of almost identical size are observed. For the method (5.8), the number of iterations required to solve the system is higher for smaller values of  $\varepsilon$ ; for the penalty method, the number of iterations is less affected by the value of  $\varepsilon$ , leading to lower iteration counts than the method (5.8) for small values of  $\varepsilon$ .




FIG. 7. The dependence of the error on  $\epsilon$  and  $\beta$  for the Robin problem on the unit sphere with  $h = 0.1$ , with  $k = l + 1 = 1$ . Here we use  $\beta_D = \beta h^{-1}$  and  $\beta_N = \beta h$ .




FIG. 8. The convergence (left) and iteration counts (right) of the penalty method for the Robin problem with  $\epsilon = 300$  (green triangles),  $\epsilon = 1$  (green diamonds) and  $\epsilon = 1/300$  (green pentagons) on the unit sphere, using  $k = l + 1 = 1$  and  $\beta = 0.01$ . The dashed line shows order 1.5 convergence. Here we use  $\beta_D = \beta h^{-1}$  and  $\beta_N = \beta h$ .




FIG. 9. The dependence of the error on  $\epsilon$  and  $\beta$  for the Robin problem on the unit sphere with  $h = 0.1$ , with  $k = l = 1$ . Here we use  $\beta_D = \beta_N = \beta$ .




FIG. 10. The convergence (left) and iteration counts (right) of the penalty method (green) compared to the method (5.8) (gray dashed), for the Robin problem with  $\varepsilon = 300$  (triangles),  $\varepsilon = 1$  (diamonds), and  $\varepsilon = 1/300$  (pentagons) on the unit sphere, using  $k = l = 1$  and  $\beta = 0.01$ . The dashed line shows order 2 convergence. Here we use  $\beta_D = \beta_N = \beta$ .

Again, we could consider the penalty-free formulation for the Robin problem. However, Figures 7 and 9 suggest that as  $\beta \rightarrow 0$ , the error increases for some values of  $\varepsilon$ . This increased error can also be observed in the numerical experiments we have run with  $\beta = 0$ . Hence in the Robin case, the penalty term is necessary and Theorem 4.30 does not hold for  $\beta_R = 0$ .

**6. Conclusions.** We have analyzed and demonstrated the effectiveness of Nitsche type coupling methods for boundary element formulations. In particular, for Robin and mixed Neumann/Dirichlet boundary conditions these are simpler than the strong imposition of boundary conditions since the boundary condition only enters the equations through a sparse operator.

An open problem is preconditioning. While the iteration counts in the presented examples were already practically useful, for large and complex structures preconditioning is still essential. The hope is to use the properties of the Calderón projector to build effective operator preconditioning techniques for the presented Nitsche type frameworks.

An extension of the presented method to FEM/BEM formulations is currently in preparation. Other directions are the Helmholtz and Maxwell problems. Although the analysis for these cases is more involved, we expect that their implementation will be structurally similar to the presented Laplace case.

#### REFERENCES

- [1] I. BABUŠKA, *The finite element method with penalty*, Math. Comp., 27 (1973), pp. 221–228.
- [2] I. BABUŠKA AND A. K. AZIZ, *Survey lectures on the mathematical foundations of the finite element method*, in The Mathematical Foundations of the Finite Element Method with Applications to Partial Differential Equations, Academic Press, New York, 1972, pp. 1–359.
- [3] L. BARATCHART, L. BOURGEOIS, AND J. LEBLOND, *Uniqueness results for inverse Robin problems with bounded coefficient*, J. Funct. Anal., 270 (2016), pp. 2508–2542.
- [4] M. BEBENDORF, *Hierarchical LU decomposition-based preconditioners for BEM*, Computing, 74 (2005), pp. 225–247.
- [5] R. BECKER, P. HANSBO, AND R. STENBERG, *A finite element method for domain decomposition with non-matching grids*, Math. Model. Numer. Anal., 37 (2003), pp. 209–225.

- [6] T. BETCKE, M. SCROGGS, AND W. SMIGAJ, *Product algebras for Galerkin discretizations of boundary integral operators and their applications*, preprint, arXiv:1711.10607, 2017, <https://ui.adsabs.harvard.edu/abs/2017arXiv171110607B>.
- [7] H. BREZIS AND P. MIRONESCU, *Gagliardo-Nirenberg, composition and products in fractional Sobolev spaces*, *J. Evol. Equ.*, 1 (2001), pp. 387–404.
- [8] A. BUFFA AND S. H. CHRISTIANSEN, *A dual finite element complex on the barycentric refinement*, *C. R. Math.*, 340 (2005), pp. 461–464.
- [9] A. CHERNOV AND P. HANSBO, *Nonconforming hp-FE/BE coupling for interface problems based on Nitsche’s method*, unpublished manuscript.
- [10] F. CHOULY AND N. HEUER, *A Nitsche-based domain decomposition method for hypersingular integral equations*, *Numer. Math.*, 121 (2012), pp. 705–729.
- [11] M. COSTABEL AND E. STEPHAN, *Boundary integral equations for mixed boundary value problems in polygonal domains and Galerkin approximation*, *Banach Center Publ.*, 15 (1985), pp. 175–251.
- [12] E. DI NEZZA, G. PALATUCCI, AND E. VALDINOCI, *Hitchhiker’s guide to the fractional Sobolev spaces*, *Bull. Sci. Math.*, 136 (2012), pp. 521–573.
- [13] G. GATICA, M. HEALEY, AND N. HEUER, *The boundary element method with Lagrangian multipliers*, *Numer. Methods Partial Differential Equations*, 25 (2009), pp. 1303–1319.
- [14] M. R. HESTENES AND E. STIEFEL, *Methods of conjugate gradients for solving linear systems*, *J. Res. National Bureau of Standards*, 49 (1952), pp. 409–436.
- [15] B. JIN AND J. ZOU, *Numerical estimation of the Robin coefficient in a stationary diffusion equation*, *IMA J. Numer. Anal.*, 30 (2010), pp. 677–701.
- [16] M. JUNTUNEN AND R. STENBERG, *Nitsche’s method for general boundary conditions*, *Math. Comp.*, 78 (2009), pp. 1353–1374.
- [17] U. LANGER AND O. STEINBACH, *Boundary element tearing and interconnecting methods*, *Computing*, 71 (2003), pp. 205–228.
- [18] U. LANGER AND O. STEINBACH, *Coupled boundary and finite element tearing and interconnecting methods*, in *Domain Decomposition Methods in Science and Engineering*, *Lect. Notes Comput. Sci. Eng.*, Springer, Berlin, 40 (2005), pp. 83–97.
- [19] J. NITSCHKE, *Über ein Variationsprinzip zur Lösung von Dirichlet-Problemen bei Verwendung von Teilräumen, die keinen Randbedingungen unterworfen sind*, *Abh. Math. Seminar Univ. Hamb.*, 36 (1971), pp. 9–15.
- [20] C. PAIGE AND M. SAUNDERS, *Solution of sparse indefinite systems of linear equations*, *SIAM J. Numer. Anal.*, 12 (1975), pp. 617–629.
- [21] Y. SAAD AND M. H. SCHULTZ, *GMRES: A generalized minimal residual algorithm for solving nonsymmetric linear systems*, *SIAM J. Sci. Statist. Comput.*, 7 (1986), pp. 856–869.
- [22] W. ŚMIGAJ, T. BETCKE, S. ARRIDGE, J. PHILLIPS, AND M. SCHWEIGER, *Solving boundary integral problems with BEM++*, *ACM Trans. Math. Software*, 41 (2015).
- [23] O. STEINBACH, *Numerical Approximation Methods for Elliptic Boundary Value Problems: Finite and Boundary Elements*, Springer, New York, 2008.
- [24] O. STEINBACH AND M. WINDISCH, *Stable boundary element domain decomposition methods for the Helmholtz equation*, *Numer. Math.*, 118 (2011), pp. 171–195.
- [25] E. STEPHAN, *Boundary integral equations for mixed boundary value problems in  $\mathbb{R}^3$* , *Math. Nachr.*, 134 (1987), pp. 21–53.
- [26] T. VON PETERSDORFF AND R. LEIS, *Boundary integral equations for mixed Dirichlet, Neumann and transmission problems*, *Math. Methods Appl. Sci.*, 11 (1989), pp. 185–213.
- [27] T. VON PETERSDORFF AND E. P. STEPHAN, *Regularity of mixed boundary value problems in  $\mathbb{R}^3$  and boundary element methods on graded meshes*, *Math. Methods Appl. Sci.*, 12 (1990), pp. 229–249.
- [28] J. XU AND L. ZIKATANOV, *Some observations on Babuška and Brezzi theories*, *Numer. Math.*, 94 (2003), pp. 195–202.