JGR Space Physics

RESEARCH ARTICLE

10.1029/2018JA026169

Key Points:

- A technique to resolve mass-per-charge ambiguity of O^+ and S^{2+} for ESA-TOF instruments using carbon foil effects
- An analysis tool to extract ion distributions from in situ measurements of Juno JADE-I
- Ion bulk properties derived from the forward model with convected kappa distributions in the jovian plasma sheet

Correspondence to:

T. K. Kim,
thomas.kim@contractor.swri.org

Citation:

Kim, T. K., Ebert, R. W., Valek, P. W., Allegrini, F., McComas, D. J., Bagenal, F., et al. (2020). Method to derive ion properties from Juno JADE including abundance estimates for O^+ and S^{2+} . *Journal of Geophysical Research: Space Physics*, 125, e2018JA026169. <https://doi.org/10.1029/2018JA026169>

Received 12 OCT 2018

Accepted 29 JUL 2019

Accepted article online 4 AUG 2019

Method to Derive Ion Properties From Juno JADE Including Abundance Estimates for O^+ and S^{2+}

Thomas K. Kim^{1,2}, R. W. Ebert^{2,1}, P. W. Valek², F. Allegrini^{2,1}, D. J. McComas^{2,3}, F. Bagenal⁴, K. Chae², G. Livadiotis², C. E. Loeffler², C. Pollock⁵, D. A. Ranquist⁴, M. F. Thomsen⁶, R. J. Wilson⁴, G. Clark⁷, P. Kollmann⁷, B. H. Mauk⁷, S. Bolton², S. Levin⁸, and G. Nicolaou^{2,9}

¹Department of Physics and Astronomy, University of Texas at San Antonio, TX, USA, ²Space Science and Engineering Division, Southwest Research Institute, San Antonio, TX, USA, ³Department of Astrophysical Sciences, Princeton University, Princeton, NJ, USA, ⁴Laboratory for Atmospheric and Space Physics, University of Colorado Boulder, CO, USA, ⁵Denali Scientific, Healy, AK, USA, ⁶Planetary Science Institute, Tucson, AZ, USA, ⁷The Johns Hopkins University Applied Physics Laboratory, Laurel, MD, USA, ⁸Jet Propulsion Laboratory, Pasadena, CA, USA, ⁹Department of Space and Climate Physics, Mullard Space Science Laboratory, University College London, Dorking, UK

Abstract The Jovian Auroral Distributions Experiment Ion sensor (JADE-I) on Juno is a plasma instrument that measures the energy-per-charge (E/Q) distribution of 0.01 to 46.2 keV/q ions over a mass-per-charge (M/Q) range of 1–64 amu/q. However, distinguishing O^+ and S^{2+} from JADE-I's measurements is a challenging task due to similarities in their M/Q (~ 16 amu/q). Because of this, O^+ and S^{2+} have not been fully resolved in the in situ measurements made by plasma instruments at Jupiter (e.g., Voyager PLS and Galileo PLS) and their relative ratios has been studied using physical chemistry models and ultraviolet remote observations. To resolve this ambiguity, a ray tracing simulation combined with carbon foil effects is developed and used to obtain instrument response functions for H^+ , O^+ , O^{2+} , O^{3+} , Na^+ , S^+ , S^{2+} , and S^{3+} . The simulation results indicate that JADE-I can resolve the M/Q ambiguity between O^+ and S^{2+} due to a significant difference in their charge state modification process and a presence of a large electric potential difference (~ 8 kV) between its carbon foils and MCPs. A forward model based on instrument response functions and convected kappa distributions is then used to obtain ion properties at the equatorial plasma sheet ($\sim 36 R_J$) in the predawn sector of magnetosphere. The number density ratio between O^+ and S^{2+} for the selected plasma sheet crossings ranges from 0.2 to 0.7 (0.37 ± 0.12) and the number density ratio between total oxygen ions to total sulfur ions ranges from 0.2 to 0.6 (0.41 ± 0.09).

1. Introduction

Jupiter's magnetosphere is the largest cohesive structure in the heliosphere. The dynamics of Jupiter's magnetosphere is mainly driven by the planet's strong magnetic field, fast rotation period, and an immense internal plasma source from Io. Volcanic activity on Io's surface releases neutral SO_2 , which is dissociated and ejected from Io with subsequent ionization by electron impact at a rate of about 1 ton/s (see review by Thomas et al., 2004, and references therein).

The magnetospheric ions at Jupiter are largely composed of dissociation products of iogenic SO_2 , 80% of which comprise ions with similar mass-per-charge (M/Q) (e.g., $O^+ \approx S^{2+} \approx 16$ amu/q). Newborn iogenic ions are then picked up by the ambient magnetic field and accelerated up to the bulk speed (~ 57 km/s relative to Keplerian orbital speed of Io) of the surrounding plasma at the orbit of Io. These relatively cold dense plasmas move radially outward via centrifugally driven flux tube interchange instability forming Jupiter's plasma sheet (Hill & Michel, 1976).

Jovian magnetospheric ions have been observed by instruments onboard several previous missions (e.g., Bridge et al., 1977; Frank et al., 1992; Gloeckler et al., 1992; Krimigis et al., 1977; McComas et al., 2008; Williams et al., 1992). However, the in situ relative abundance of O^+ and S^{2+} in the plasma sheet are not well constrained due to their similarities in M/Q (e.g., Bagenal & Sullivan, 1981; Bridge et al., 1979; McNutt et al., 1981). It is known that the magnetospheric plasmas are corotating at Jupiter's ~ 10 -hr rotation period within a distance of $\sim 10 R_J$ ($1 R_J = 71,492$ km) from the planet. Beyond $\sim 10 R_J$, plasmas are under the corotation

speed (i.e., subcorotating) but still share a common flow speed (Bagenal et al., 2017; McNutt et al., 1981). O^+ and S^{2+} have a similar energy-per-charge (E/Q) under these circumstances. These similarities in E/Q and M/Q make it difficult to resolve these ions in measurements that employ electrostatic analyzer (ESA), or ESA and time-of-flight (TOF) techniques. A mass resolution ($M/\Delta M$) of ~ 500 is required to resolve the M/Q ambiguity between O^+ and S^{2+} , but typical plasma instruments have $M/\Delta M < 10$ (e.g., Galileo PLS had mass resolution ranging from $2 \leq M/\Delta M \leq 4.1$, see Frank et al., 1992).

Instruments comprised of an ESA and a carbon-foil based TOF mass spectrometer are commonly used to measure plasma in space environments (see Wüest, 1998, and references therein). An ESA selects incident ions by their E/Q , while a TOF mass spectrometer can determine the speed (v) of the ions so that the M/Q can be derived from its kinetic energy ($E/Q = 1/2 M/Q \cdot v^2$). The speed of the ions is estimated by measuring their TOF through a known path length in the instrument, the TOF being determined by the time difference between start signals generated by secondary electrons (SE) and stop signals produced by the ions. These SEs are generated when the ions transmit through thin carbon foils. In addition to the SE generation, there are other effects on the transmitted ions (e.g., Allegrini et al., 2003; McComas et al., 2004; Ritzau & Baragiola, 1998; Wüest, 1998). These include charge state modification (e.g., Allegrini et al., 2014; Allegrini, Ebert & Funsten 2016; Gonin et al., 1994; Kallenbach et al., 1993, 1995), angular scattering (e.g., Ebert et al., 2014; Funsten et al., 1993), and energy straggling (e.g., Allegrini et al., 2006, 2015). Such effects, which we refer to collectively as ‘carbon foil effects’ hereafter, are the consequences of nuclear and electronic interactions between the incident ions and carbon foil materials. The count-TOF distribution (i.e., TOF measurements at a given energy step) is strongly affected by carbon foil effects since they can alter incoming ion properties (e.g., incident energy, angle, and charge states with respect to the carbon foil) and can change the electrodynamics of ions in the TOF instrument. The shape of the count-TOF distribution for ions with similar M/Q can be different if they have distinct carbon foil effects. Thus, by characterizing these effects for an individual ion species per energy step of the instrument, better species determination of the ions with M/Q ambiguity can be achieved for instruments using ESAs with non-field-free TOF mass spectrometers.

The Jovian Auroral Distributions Experiment (JADE) on NASA’s Juno mission to Jupiter (McComas et al., 2017) is an instrument suite composed of two identical electron sensors (JADE-E) and an ion sensor (JADE-I). JADE-I measures the E/Q (~ 0.01 – 46.2 keV/q) and M/Q (~ 1 – 64 amu/q) of ions while JADE-E measures electron energy from ~ 0.05 – 100 keV/q. JADE-I is composed of a top-hat ESA, carbon foils, and a TOF mass spectrometer. It is designed to measure ion energy distributions, angular distributions, and ion composition in the polar regions of the jovian magnetosphere and is one of the key instruments for addressing the magnetospheric science questions discussed in Bagenal et al. (2017). One goal of a plasma instrument is to derive macroscopic plasma parameters (i.e., velocity moments) from the measured distributions. To achieve this, a distribution for each individual ion species is necessary for obtaining proper bulk parameters in a multi ion plasma environment (e.g., the jovian magnetosphere). However, the mass resolution of JADE-I (i.e., $2.5 \leq M/\Delta M \leq 11$) is not sufficient to directly resolve the M/Q between O^+ and S^{2+} .

On the other hand, the TOF section of JADE-I has a huge electric potential difference (~ 8 kV) between its carbon foils and microchannel plates (MCP). This potential difference generates nonlinear electric fields that accelerate electrons and negative ions or decelerate positively charged ions toward MCPs (see Figure 1). While most of the ions have a neutral charge state after transiting the carbon foils, a fraction exits with non-neutral charge states. The difference in exit (or residual) charge states for individual ion species produce a distinct count-TOF distribution under the strong nonlinear electric field in the JADE-I’s TOF section. In this study, we demonstrate how the M/Q ambiguity between O^+ and S^{2+} can be resolved in JADE-I by detailed modeling of the instrument response using carbon foil effects. Our simulation results show that they have distinct count-TOF distribution due to the significant difference in their charge state modification process (especially on the non-neutral charge states) and the presence of a strong electrostatic potential (~ 8 kV) in the TOF section of the instrument.

This study also presents bulk ion properties in the predawn sector (local time of $\sim 04:16$) of the jovian plasma sheet observed by Juno JADE-I. A kappa distribution is used in our analysis to account for wide E/Q measurements from JADE-I. They were first used on analysis of electron data from IMP-1, IMP-2, OGO-1, and OGO-3 (e.g., Olbert, 1968; Vasyliunas, 1968) and it is an appropriate distribution function that can describe both low energy (typically Maxwell-Boltzmann) and energetic (power law) ion/electron populations. We combine a forward model using kappa distributions and the instrument response functions from the JADE-I

Figure 1. The JADE-I SIMION model. Each colored line shows possible trajectories of incident H^+ (1 keV/q) for different residual charge states (blue: $H^{(+)}$, green: $H^{(0)}$, and red: $H^{(-)}$) after transit through the carbon foil. The trajectory for different charge states are different due to large electric potential differences from carbon foils to MCPs. The orange line is a trajectory of secondary electrons released from the carbon foil.

Simulation Tool (JIST). Section 2 describes the data from JADE and MAG (Connerney et al., 2017) that are used in this study. Section 3 explains the forward model. Section 4 discusses ion parameters obtained from the forward model and comparisons with previous observations. An appendix presents JIST, comparisons with laboratory measurements, background subtraction method, and the updated geometric factor $\times$ detection efficiency.

2. Data

JADE-I measures the distribution of ions every 2 s. It measures the $4\pi \text{ sr}$ distribution every $\sim 30\text{-s}$ spacecraft spin period (McComas et al., 2017). The JADE-I data used in this study are provided with three different formats, LOG, TOF, and SPECIES data products. LOG data have 25 logical counters including counts from 13 anodes (12 anodes + background anode; Wilson, 2017). We use background anode counts for subtracting backgrounds in TOF data (see Appendix D). The TOF data product has high mass resolution (93 TOF bands) summed over all look directions, and the SPECIES data product has high angular resolution with coarse mass resolution. SPECIES data product has 12 anodes (only the data obtained from eight anodes are used for Low Rate Science), and TOF data product has 248 separate TOF channels (or bins) onboard that are collapsed down to 93 TOF bands for transmission. Both TOF and SPECIES data sets have the same 64 logarithmically spaced energy steps covering from ~ 0.01 to 46.2 keV/q . This study uses JADE-I TOF data to derive ion properties in the Jovian plasma sheet and the JADE-I TOF data sets presented in this study are all Low-Rate Science (LRS) data with 30-s time resolution.

In July 2016, Juno arrived at Jupiter and initiated its prime science investigation (Bolton, Lunine, et al. 2017; Bolton, Adriani, et al. 2017; Connerney et al., 2017). Juno has a polar orbit with a period of $\sim 53 \text{ days}$. Perijove (PJ) occurs very close to the planet (3,000–5,000 km above the 1-bar level of the atmosphere) while apojove is at $\sim 110 R_J$. Through this orbit, Juno explores Jupiter's polar magnetosphere, plasma sheet, and outer magnetosphere. Figure 2 shows total JADE-I count rates and the spacecraft trajectory in a magnetic coordinate system. The magnetic field lines (magenta lines) were obtained from VIP4+CAN magnetic field model (Connerney et al., 1981, 1998). The wiggly nature of the spacecraft trajectory in a magnetic coordinate system is due to the tilt angle between Jupiter's rotation axis and magnetic dipole axis. As shown in Figure 2,

Figure 2. Count rates of JADE-I in a magnetic coordinate system from 2017-188T18:00:00 to 2017-191T12:00:00 (from left to right). The magnetic field lines (magenta) are from the VIP4+CAN magnetic field model (Connerney et al., 1981, 1998) and the spacecraft trajectory for Juno orbit 7 is shown by the white line. Start and end time of the data sets presented in this study are highlighted with white boxes.

Figure 3. Count rates measured by JADE-I anode 4–11 from 2017-189T16:42:20 to 2017-189T20:40:50. The plasma sheet ions are mostly observed in anodes 8–11. R is the radial distance in spherical coordinate system and ρ is the radial distance in cylindrical coordinate system (z axis for both coordinate systems are aligned with Jupiter's rotation axis). Both R and ρ are in units of the jovian radii (R_J). ρ is used to compute a rigid corotation speed.

Figure 4. E/Q -time spectra of ion (top) and electron (middle) JADE data from 2017-189T16:42:20 to 2017-189T20:40:50 (highlighted with white boxes in Figure 2). Bottom panel shows magnetic field in spherical coordinates. Equatorial plasma sheet crossings are identified by the reversal of the magnetic field vector radial component. Fourteen intervals are selected to obtain the ion properties of the plasma sheet.

the spacecraft encounters multiple plasma sheet crossings in both the equatorial and high-latitude region of the magnetosphere during the inbound trajectory prior to PJ 7.

We selected the plasma sheet data sets using the following criteria: (1) the count rates are focused in the magnetic equator (i.e., $z_{\text{MAG}} \approx 0$); (2) the radial component of the magnetic field vector is near zero (i.e., equatorial plasma sheet crossing); and (3) the plasma sheet flow is mainly observed in anodes 8–11 of JADE-I. Criteria (3) is important to avoid oversampling the distribution (anodes 0–3 and 4–7 measures same part of the sky over a spacecraft spin) considering that the JADE-I TOF data essentially are the sum of counts measured from anodes 0–11. Selected data ranges are highlighted with white boxes in Figure 2. Figure 3 shows the JADE-I count rates for all anodes during this period, and it shows that the counts are mainly observed at the 8–11. Anodes 0–3 are not shown because in LRS modes, just anodes 4–11 are returned to provide a 4π sr view of the sky.

A summary plot showing the E/Q -time spectra for ions (JADE-I), electrons (JADE-E), and magnetic field (MAG) data during this interval is shown in Figure 4. The ions are mainly observed at >1 keV/q where the energy centroid appears at ~ 10 keV/q. The electrons are distributed over a much wider energy range (occupying energy from ~ 0.1 keV to ~ 40 keV) compared to the ions. Both ions and electrons have fine structures, possibly due to dynamic events during this interval. This is also well represented in the magnetic field data. The radial component of the magnetic field data shows that the spacecraft crossed the magnetic equator twice indicated by when the radial component of the magnetic field vector (B_r) is near zero.

It is clear from the E/Q -time spectra that the ion and electron energy distributions are very dynamic over time. There are times when a significant portion of the ion energy distribution extends above the upper limit

Table 1
Time Intervals of This Study During 2017-189

Interval	UTC Start	UTC End	Total Accumulation Time
A	17:31:54	17:33:54	00:02:00 (120 s)
B	17:37:54	17:40:24	00:02:30 (150 s)
C	17:47:24	17:49:24	00:02:00 (120 s)
D	18:01:24	18:06:54	00:04:30 (270 s)
E	18:09:24	18:13:24	00:04:00 (240 s)
F	18:23:24	18:28:24	00:05:00 (300 s)
G	18:31:54	18:34:24	00:02:30 (150 s)
H	18:45:54	18:50:54	00:05:00 (300 s)
I	18:52:54	18:58:54	00:06:00 (360 s)
J	19:13:54	19:16:54	00:03:00 (180 s)
K	19:22:24	19:24:24	00:02:00 (120 s)
L	19:30:24	19:33:54	00:02:30 (150 s)
M	19:40:54	19:44:54	00:04:00 (240 s)
N	19:50:54	19:57:24	00:06:30 (390 s)

of JADE-I's E/Q range. Because of this, we further divide the data sets into subintervals by choosing time periods where (1) the variation in energy distribution (both ion and electron) and the total magnetic field strength is small over time; and (2) the peak of the energy distribution is resolved in the JADE-I data. These selected intervals are indicated in Figure 4 and are listed on Table 1.

The TOF data can be presented in E/Q -time spectra (summed over all TOF bands for each energy step) as shown in the top panel of Figure 5 and in E/Q -TOF spectra (summed over each record or time steps) for detailed mass analysis as shown in the middle panel of Figure 5. The mass tracer lines shown in the E/Q -TOF spectra (middle panel) are the estimated peak position of the given ion species based on equation (B2) that was obtained during in-flight calibration (see Appendix B1). As shown in the middle panel of Figure 5, O^+ and S^{2+} are the dominant ion species in the data. Because they have similar M/Q (~ 16 amu/ q), their count-TOF distributions (i.e., horizontal slice of the E/Q -TOF spectra shown in the middle panel of Figure 5) are nearly superposed to each other (see mass tracer lines for O^+ and S^{2+}) so that it is difficult to determine their relative ratios.

During the plasma sheet crossings we have also identified the presence of high-energy ion populations which typically extend to the top of JADE-I's E/Q range (i.e., 46.2 keV/ q). They are not the major component of the plasma sheet ion distributions and are hard to characterize due to the limited information provided by JADE-I. Because we only use a single distribution function for individual ions species, we remove them from in situ E/Q -TOF spectra when the forward model results have kappa values less than 1.6. The removal process adopts equation (B2) to account for corotational flow of the plasma sheet. Under a common flow speed, ion species with higher M/Q will appear at higher E/Q , whereas ion species with lower M/Q appears at lower E/Q . Equation (B2) provides an estimate of E/Q value for ions with different M/Q . For this study, we crudely determine seven boundaries where the flow speeds for empirical removal boundaries are given by plugging energies (derived from empirical corotation speed that corresponds to O^+ energies at 15, 20, 25, 30, 35, and 40 keV/ q + post-acceleration -10 keV) to E_T (total ion energy at the carbon foil), set of mass arrays (ranging from 0.01 to 50 amu) to M , and $Q=1$ in equation (B2). The bottom panel shows an example of boundaries obtained with O^+ at 20 keV/ q (solid and dashed white lines from bottom left to upper right). We preserve proton data in the E/Q -TOF spectra by giving another boundary obtained by plugging $M=2$ and $Q=1$ into equation (B2). This boundary is shown next to the proton tracer line in the bottom panel of Figure 5. Then the upper area covered by both corotational flow line and the proton boundary (white solid lines above ~ 2 keV/ q) are removed from the E/Q -TOF spectra for the forward model calculations.

To resolve the M/Q ambiguity between O^+ and S^{2+} , we developed JIST for characterizing count-TOF distributions for individual ion species. JIST is a ray tracing simulation tool capable of calculating count-TOF distributions for individual ions using carbon foil effects (see Appendix A1 for details). The sum of multiple instrument response functions (i.e., normalized count-TOF distributions over JADE-I's E/Q range for a

Figure 5. E/Q -time spectra (top) and E/Q -TOF (time-of-flight) spectra (middle) of interval A (2 min of data from 2017-189T17:31:54 to 2017-189T17:33:54). Colored lines in the E/Q -TOF spectra are the mass tracer lines that indicate expected peak position of each ion species. Protons have two tracer lines due to the charge modification process in the carbon foil. The bottom panel shows an example of data removal boundary at 20 keV/q of O^+ cutoff energy. The data enclosed by white solid lines are removed from E/Q -TOF spectra for the forward model calculation. An additional measurement error of 10% are applied in the data where TOF is greater than the magenta lines shown in the bottom panel.

given ion species) shown in Figure 6 gives E/Q -TOF spectra that will be used to characterize JADE-I TOF measurements. The top panel shows E/Q -TOF spectra and bottom two panels show normalized count-TOF distributions at 1.07 and 9.37 keV/q, respectively. We combine the instrument response functions with convected kappa distributions to obtain ion properties for the periods identified in Table 1.

3. Forward Model

This section describes the forward model for obtaining plasma parameters from the JADE-I measurements. The model uses convected kappa distributions, the instrument response functions obtained from JIST for eight ion species (H^+ , O^+ , O^{2+} , O^{3+} , Na^+ , S^+ , S^{2+} , and S^{3+}), and chi-square minimization to reproduce in situ E/Q -TOF spectra.

Figure 6. E/Q -TOF spectra (top) of the normalized instrument response functions from JIST. Bottom two panels show count-TOF distributions at $E/Q=1.07$ keV/q (middle) and $E/Q=9.97$ keV/q (bottom). The black line is the total distribution of all ions and each colored line represent individual ion species (shown in gray box around the top panel). Each ion species has different shaped distributions at each energy step due to the carbon foil effects and incident energy.

We start with the standard form of the convected kappa distribution (Livadiotis, 2017, Chapter 4) with common isotropic temperature for individual ions and a common flow velocity $\vec{v}_f$ (i.e., corotating plasma),

$$f(E, \delta) = n \cdot \left[\pi \left(\kappa - \frac{3}{2} \right) \eta^2 \right]^{-\frac{3}{2}} \cdot \frac{\Gamma(\kappa + 1)}{\Gamma(\kappa - \frac{1}{2})} \cdot \left[1 + \frac{E + E_f - 2\sqrt{E}\sqrt{E_f} \cos \delta}{(\kappa - \frac{3}{2})k_B T} \right]^{-\kappa-1}, \quad (1)$$

where $\eta = \sqrt{\frac{2k_B T}{m}}$ is the thermal velocity, $E_f = 1/2 \cdot m v_f^2$ is the flow energy where v_f is the flow speed, and δ is the angle between the individual ion velocity $\vec{v}$ and the corotational flow vector $\vec{v}_f$ representing the thermal spread of the velocity. The spacecraft speed v_{sc} is not considered here since the TOF data product does not have any directional information and it is relatively small (~ 8 km/s) compared to the flow speed (see Tables 2 and 3) for the intervals shown in Table 1.

The level 3 TOF data product is the sum of count rates observed by 12 JADE-I anodes covering an instantaneous field of view of $\sim 270^\circ \times 15^\circ$. Converting count rates to counts c (by multiplying total accumulation time T_{acc} (see Table 1) and the accumulation time per energy step for JADE-I, $\Delta t [s] = \frac{1}{32} - 2 \times 10^{-3}$), the total measured counts C at energy level E is

$$C(E) = \sum_{l,m} c(E, \theta_l, \phi_m), \quad (2)$$

where l designates the anode number and m represents the azimuth (accumulation) number.

The counts are related to flux j and phase space density f such that equation (2) can be rewritten as

$$C(E) = \sum_{l,m} E G \epsilon(E) \Delta t \cdot j(E, \theta_l, \phi_m) = \sum_{l,m} \frac{2E^2 G \epsilon(E) \Delta t}{m^2} \cdot f(E, \theta_l, \phi_m), \quad (3)$$

where $G \epsilon [m^2 \cdot eV/eV \cdot sr] = 6.84 \times 10^{-5} + 2.52 \times 10^{-6} \cdot \log_{10}(E/Q [eV/q]) + 6.84 \times 10^{-5} \cdot E/Q [eV/q]^{-0.28}$ is the geometric factor $G \times$ detection efficiency ϵ (see Appendix E) and $\Delta t [s] = \frac{1}{32} - 2 \times 10^{-3}$ is the accumulation

time per energy step for JADE-I, after accounting for the hemispherical voltage settling time (the 2×10^{-3} term).

Total accumulation time (T_{acc}) of the LRS data sets are approximately multiples of the spacecraft spin period (~ 30 s), the total counts C is roughly equivalent to multiple omni-directional summations of the true distribution function. Then the average counts per sector (one anode, one azimuth) is

$$\begin{aligned} \langle c(E) \rangle &= \frac{C(E)}{12 \cdot \frac{T_{\text{acc}}}{2}} = \frac{2E^2 G \epsilon(E) \Delta t}{m^2} \cdot \langle f(E, \theta_l, \phi_m) \rangle \\ &= \frac{2E^2 G \epsilon(E) \Delta t}{m^2} \cdot \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi f(E, \theta_l, \phi_m) \sin \theta d\theta d\phi, \end{aligned} \quad (4)$$

where 12 represents the number of JADE-I anodes and $\frac{T_{\text{acc}}}{2}$ is the number of accumulation (or azimuth) including the duty cycle of the instrument (2 s). Thus, the analytical form of a one-dimensional count distribution for each JADE-I energy step over all look directions for accumulation N_{acc} is

$$C(E) = \left(12 \cdot \frac{T_{\text{acc}}}{2} \right) \cdot \frac{2E^2 G \epsilon(E) \Delta t}{m^2} \cdot \frac{1}{4\pi} \int_0^{2\pi} \int_0^\pi f(E, \theta_l, \phi_m) \sin \theta d\theta d\phi, \quad (5)$$

Plugging equation (1) into (5) and choosing a coordinate system such that $\theta = \delta$ (i.e., $f = f(E, \theta_l) = f(E, \delta)$, f is now independent of ϕ) yields an analytical form of the convected kappa distribution in a JADE-I frame of reference,

$$\begin{aligned} C(E) &= \left(12 \cdot \frac{T_{\text{acc}}}{2} \right) \cdot \frac{n}{8} \sqrt{\frac{2}{m}} \cdot \left[\pi^3 E_f k_B T \left(\kappa - \frac{3}{2} \right) \right]^{-\frac{1}{2}} \cdot \frac{\Gamma(\kappa)}{\Gamma(\kappa - \frac{1}{2})} \\ &\cdot \left\{ \left[1 + \frac{E + E_f - 2\sqrt{E}\sqrt{E_f}}{(\kappa - \frac{3}{2})k_B T} \right]^{-\kappa} - \left[1 + \frac{E + E_f + 2\sqrt{E}\sqrt{E_f}}{(\kappa - \frac{3}{2})k_B T} \right]^{-\kappa} \right\} \cdot E^{\frac{3}{2}} G \epsilon(E) \Delta t. \end{aligned} \quad (6)$$

For a given interval (see Table 1), we use equation (6) to generate eight convected kappa distributions (count- E/Q distributions each representing H^+ , O^+ , O^{2+} , O^{3+} , Na^+ , S^+ , S^{2+} , and S^{3+}). Without any further physical assumptions, this analysis has 25 free parameters (i.e., a common flow speed, 8 densities, 8 temperatures, and 8 kappa values) which requires significant computation time to find a global minimum (i.e., absolute minimum where the difference between the data and the model is smallest in chi-square space). In order to reduce the number of free parameters in the analysis, we use a common kappa value κ_{HEAVY} and temperature T_{HEAVY} for all heavy ions (i.e., O^+ , O^{2+} , O^{3+} , Na^+ , S^+ , S^{2+} , and S^{3+}) which cuts the number of free parameters from 25 down to 13. This is a reasonable approximation since the heavy ions will be in thermal equilibrium by the time they are transported from Io plasma torus to the plasma sheet as the transport timescale is much greater than the Coulomb equilibration timescale for these ions (Delamere et al., 2005).

Once the kappa distributions are generated, a normalized (such that the total of each count-TOF distribution in each energy step is equal to 1) instrument response function S (shown in Figure 6) obtained from JIST needs to be multiplied to each point (i.e., JADE-I energy step) of the convected kappa distribution functions to produce the forward model in E/Q -TOF spectra,

$$\text{Model}(E, \tau) = \sum_{i=1}^8 C_i(E) \cdot S_i(E, \tau), \quad (7)$$

where subscript i represents each ion species and τ is each bin of 93 TOF bands. Instrument response function S is a crucial part of the forward model because it converts the kappa distributions (i.e., count- E/Q distribution) to E/Q -TOF spectra and provides constraints to resolve the M/Q ambiguity between O^+ and S^{2+} . Most of the previous studies used the forward model only in energy or E/Q space due to the absence of M/Q measurements or the inability to model M/Q (or TOF) measurements. However, O^+ and S^{2+} are the dominant species in Jupiter's magnetosphere (indicated at the bottom panel of Figure 5) and they have similar M/Q and E/Q if they share common flow speed. We use the TOF information and consider carbon foil effects, which are different for each ion species, to resolve this ambiguity.

Figure 7. Forward model results for interval A. The top two panels show E/Q -TOF spectra of JADE-I measurements (top) and forward model results (middle). The bottom panel shows a comparison between the in situ measurement and the forward model results (the color-coded lines represent each ion and the magenta dots represent the total of the forward model) at $E/Q=8.58$ keV/q in count-TOF space.

The best fit parameters of the forward model that represent the JADE-I E/Q -TOF spectra can be found by minimizing the chi-square value defined as

$$\chi^2 = \sum_{j=0}^{63} \sum_{k=0}^{92} \frac{[\text{Data}(E_j, \tau_k) - \text{Model}(E_j, \tau_k)]^2}{\sigma_{\text{TOTAL}}^2(E_j, \tau_k)}, \quad (8)$$

where Data is the background subtracted in situ data (see Appendix D for background subtraction) and σ_{TOTAL} is the uncertainty for the background-subtracted data expressed as

$$\sigma_{\text{TOTAL}}(E, \tau) = \sqrt{\sigma_{\text{RD}}^2(E, \tau) + \sigma_{\text{BG}}^2(E, \tau)}, \quad (9)$$

assuming that σ_{RD} (error of the raw data with no background subtraction) and σ_{BG} are independent to each other (Bevington & Robinson, 2003). We use the Powell minimization method presented in Seiler and Seiler (1989). Thus, the forward model finds 13 parameters v_f , $n(\text{H}^+)$, $n(\text{O}^+)$, $n(\text{O}^{2+})$, $n(\text{O}^{3+})$, $n(\text{Na}^+)$, $n(\text{S}^+)$, $n(\text{S}^{2+})$, $n(\text{S}^{3+})$, T_{PROTON} , T_{HEAVY} , κ_{PROTON} , and κ_{HEAVY} that can best represent the plasma sheet data obtained by JADE-I.

However, we decided to fix the proton kappa value to 3.5 ($\kappa_{\text{PROTON}}=3.5$) based on Collier and Hamilton (1995) due to the presence of the high-energy populations (described in section 6.8). Leaving the proton kappa as a free parameter often resulted in a very small kappa (~ 1.5) and very large temperature ($\gg 100$ keV) for the proton distributions, which does not make any physical sense considering that JADE-I can only measure up to 46.2 keV/q. κ_{HEAVY} also had same issues for some intervals. To obtain reasonable temperature and kappa value for heavy ions, we use the boundaries introduced in the data section (one of the boundaries is shown as solid white lines in the bottom panel of Figure 5) to trim the data for the forward model calculations. We start with the raw data and proceed with trimmed data sets provided by corotational tracer lines for O^+ at 15, 20, 25, 30, 35, and 40 keV/q until the kappa converged to a reasonable value ($\kappa_{\text{HEAVY}} > 1.6$). If the kappa value was obtained through data trimming, we then run the forward model again with fixed κ_{HEAVY} and its associated uncertainty on the raw (untrimmed) data. In this way, we can obtain κ_{HEAVY} and T_{HEAVY} for the major component of the heavy ion distribution obtained by JADE-I. We could only do this for the heavy ions but not for the protons since their distribution seems to require a combination of multiple distribution functions. This issue is discussed in section 6.8.

The presence of sulfur isotope ^{34}S in Io's atmosphere has been revealed by measurements from Atacama Pathfinder Experiment (APEX) and its isotope ratio $^{34}\text{S}/^{32}\text{S}$ is found to range from 0.065 to 0.120 (Mouillet et al., 2013). In this study, we only treat $^{32}\text{S}^+$, $^{32}\text{S}^{2+}$, and $^{32}\text{S}^{3+}$ for sulfuric ion species and gave 10% additional errors to the heavy ion in situ data (i.e., $\sigma_{\text{TOTAL}} = \sqrt{\sigma_{\text{TOTAL}}^2 + (0.1 \times \text{Data})^2}$) to account for the potential contribution from these isotopes. This additional error is applied in the TOF region longer than the boundary obtained by plugging in $M=14$ and $Q=3$ (shown as magenta line at the bottom panel of Figure 5). This boundary is carefully chosen to include O^{3+} data in the E/Q -TOF spectra.

Figure 7 shows the forward model result for interval A presented in Table 1. The E/Q -TOF spectra for JADE-I data and forward model result are shown in the top two panels and the bottom panel shows comparisons at $E/Q=8.58$ keV/q in count-TOF space. In the bottom panel of Figure 7, the data are shown in black lines with gray error bars. The total forward model results at this energy step is shown in magenta dots and the results for individual ions are shown with different colored lines.

The uncertainties σ of the forward model results are calculated from MPFIT (Markwardt, 2009) by providing best fit parameters obtained from Powell minimization routine (Seiler & Seiler, 1989). Then a goodness of fit of the forward model results are represented by analyzing the variation of the chi-square value within 1σ of the best fit results. Figure 8 shows a chi-square space analysis within 1σ of the best fit parameters for interval A. Both the best fit parameters and their associated uncertainties (i.e., 1σ) are shown on the right. The center of each spectra is the best fit parameter from the forward model and the range of each x and y axis are determined by $\pm 1\sigma$ of the best fit parameters. For example, the spectra in the first row, second column of Figure 8 shows the chi-square space analysis for proton density $n(\text{H}^+)$ and flow speed V_{Flow} . The chi-square values are calculated by fixing the rest of the parameters but varying values of $n(\text{H}^+)$ and V_{Flow} within their 1σ . The minimum chi-square value is shown in blue and the maximum in red. As stated in equation (8), the chi-square values represent the difference between the data and the model. A successful chi-square minimization should have minimal difference between the model and the data (i.e., global minimum). Therefore, the goodness of fit can be validated by investigating if the center of each spectra has a minimum chi-square value. Chi-square space analysis for κ_{PROTON} is not shown in Figure 8 because they are fixed in this study so that their 1σ value cannot be obtained. However, a successful chi-square minimization does not always guarantee physically reasonable parameters (e.g., very small kappa and very high temperature) since the chi-square is just a difference between the model and the data.

Figure 8 also shows a correlation between each parameter. For example, an anti-correlation (i.e., one has to decrease the value if another value increases) between O^+ and S^{2+} is shown with negatively skewed ellipse. This is due to the fact that their count-TOF distributions appear in similar TOF range and E/Q ranges. But as shown in Figures 6 and 7, the instrument response functions for O^+ and S^{2+} have different shape and peak positions due to their distinct carbon foil effects (see section C1 for details) under the large electric potential difference (~ 8 kV) in JADE-I's TOF section. A perfect circle in the spectra (e.g., $n(\text{Na}^+)$ and $n(\text{O}^+)$) indicates that both parameters do not depend on each other and therefore have no correlation in the fit.

Figure 9 compares the count-TOF distributions from the in situ measurements with forward model results at 7.59, 8.58, 9.97, 11.27, 13.10, and 19.43 keV/q for intervals A and L. The forward model results are shown as colored lines, and the total of the forward model results are represented by magenta dots. These intervals

Figure 8. A chi-square space analysis results for interval A. Each panel shows a variation of chi-square values from the best fit parameters (center of each spectra) where their axis range is determined by $\pm 1\sigma$ from the center. This figure shows that the fitted parameters are at the minima within 1σ of the best fit parameters P . Proton kappa values are not shown in this figure since they are fixed in this study.

do not require any data trimming. The results for interval L has the highest flow speed where interval A had a lowest flow speed. A detailed discussion on ion properties for selected intervals are presented in sections 5 and 6.

4. Assumptions and Limitations

Before we present the ion properties obtained from the forward model, we first summarize the assumptions and limitation in our methodology. The instrument response functions (i.e., count-TOF distributions for eight individual ions) are based on a ray tracing simulation that incorporates the carbon foil effects (see Appendix A for details). O^+ and S^{2+} are separable because they have a distinct shape in their count-TOF distribution (Figure A5) due to differences in their carbon foil effects (mainly in the charge state modification process shown in Figure A1) and the large electric potential differences applied in JADE-I's TOF section (shown in Figure 1).

We were only able to validate simulation results for singly charged ions (including O^+) with laboratory measurements (see Figures A6 and A7) due to the limitation of the ion source at Southwest Research Institute. We did not obtain any laboratory measurements on Sulfur ions due to a possible contamination of the facility. Therefore, instrument response functions for sulfur ions could not be validated. However, we think the simulated spectra for sulfur ions are representative of the laboratory measurements based on following criteria: (1) Other singly charged ions show good agreement with the laboratory measurements (Figures A6 and A7) and (2) carbon foil effects do not depend on the initial charge states of incoming ions.

Figure 9. Forward model results for intervals A (left) and L (right) at several different energies in count-TOF space. These intervals did not require any data trimming for obtaining reasonable heavy kappa values.

For the distribution functions, we assume that all heavy ions share common temperature based on the fact that the Coulomb equilibration time is much smaller than the transport time from the Io plasma torus to the plasma sheet (Delamere et al., 2005). This assumption was also used in analysis of Voyager PLS (Bagenal et al., 2017; Dougherty et al., 2017) and Galileo PLS data Bagenal et al. (2016). Also, we had to fix the proton kappa value due to the complexity of their distributions.

The error estimates for the forward model results (i.e., fitted parameters) were propagated from Poisson statistics (including data compression errors presented in Wilson, 2017) of the measured ion counts. These errors are shown in Tables 2 and 3. We note that the flow speed errors are very small because the forward model is sensitive to the changes in the flow speed (i.e., the chi-square value changes rapidly for small change in flow speed). A similar issue is discussed in Appendix C1 of Nicolaou et al. (2014).

In this study, we have ignored the spacecraft velocity in our convected kappa distribution because (1) JADE-I's level 3 TOF data does not have any directional information and (2) they are relatively small (~ 8 km/s during intervals A–N) compared to the ion flow speed. Thus, the intrinsic error for the flow speed

Figure 10. Comparisons of in situ data (black) and the forward model results (colored lines) in E/Q -count space for all intervals presented in Table 1. The total forward model results are shown as magenta dots. The convected kappa distributions represent H^+ , O^+ , O^{2+} , O^{3+} , Na^+ , S^+ , S^{2+} , and S^{3+} (see top of the figure for the corresponding colors).

should include the spacecraft speed at some degree but it requires information on the angle between the spacecraft speed direction and the flow speed direction.

Also, we would like to note that kappa values cannot be less (function diverges) or equal (delta function) to 1.5 (Livadiotis & McComas, 2013) and although our best fits are greater than 1.5, the lower boundary (provided statistically by the $\kappa_{HEAVY} - \text{Error}_{\kappa_{HEAVY}}$) for the heavy kappa values in Tables 2 and 3 are below 1.5 for some intervals. Large errors of the heavy kappa values are due to insensitivity of the forward model on the JADE-I's E/Q -TOF spectra.

5. Results

This section presents the forward model results and bulk ion properties of the Jovian plasma sheet intervals described in Table 1. Figure 10 shows a comparison between the forward model results and in situ measurements from JADE-I in count- E/Q space. The in situ data and the forward model results are summed

Figure 11. Summary plot of plasma parameters obtained from the forward model for all intervals. From top to bottom, the panels show density, density ratio of O^+ and S^{2+} , temperature, pressure, and plasma β . Juno MAG data were used to estimate plasma β . Radial distance from the planet (R), magnetic latitude ($MLat$), and equatorial crossing distance ($MShell$) are shown in the bottom of the figure.

over all TOF ranges for all intervals shown in Figure 4 and Table 1. As shown in Figure 10, the convected kappa distribution that characterizes v_f , $n(H^+)$, $n(O^+)$, $n(O^{2+})$, $n(O^{3+})$, $n(Na^+)$, $n(S^+)$, $n(S^{2+})$, $n(S^{3+})$, T_{PROTON} , T_{HEAVY} , κ_{PROTON} , and κ_{HEAVY} successfully captures most of the distribution except in the <1 and >20 keV/ q ranges. We think these are background (<1 keV/ q) and high-energy populations (>20 keV/ q) which are not a significant contributor to the bulk plasma parameters observed by JADE-I. The peak position of O^+ and S^{2+} in E/Q space (shown in Figure 10) do not always match due to their difference in thermal speed η (i.e., $\eta(O^+) = \sqrt{2} \cdot \eta(S^{2+})$). The peaks of the distribution functions (i.e., in count- E/Q space) are not solely determined by the flow energy (E_f/Q), but there are a small contribution from the thermal speed. The misalignment of O^+ and S^{2+} peak positions becomes more notable when the ratio between thermal speed and flow speed becomes larger.

Figure 11 shows plasma parameters for all intervals with errors. The values of the plasma parameters and errors are shown in Tables 2 and 3. The ion number density ratios during these plasma sheet crossings are $n(H^+)/n_e = 1\text{--}11\%$, $n(O^+)/n_e = 5\text{--}14\%$, $n(O^{2+})/n_e = 3\text{--}6\%$, $n(O^{3+})/n_e = 0.3\text{--}0.8\%$, $n(Na^+)/n_e = 1\text{--}3\%$, $n(S^+)/n_e = 2\text{--}5\%$, $n(S^{2+})/n_e = 20\text{--}28\%$, and $n(S^{3+})/n_e = 5\text{--}8\%$. n_e is an estimate of the total electron number density derived from the total ion number density assuming charge neutrality $n_e = \sum n_i q_i$, where q_i is the number

Table 2*Forward Model Results for Intervals A Through G from During the Juno orbit 7 Plasma Sheet Crossing*

Parameters	Units	A	B	C	D	E	F	G
R^a	R_J	36.12	36.08	36.01	35.91	35.86	35.76	35.71
$MLat^b$	Deg	−0.43	−0.75	−1.17	−1.70	−1.86	−2.02	−2.01
ρ^c	R_J	34.32	34.28	34.23	34.13	34.09	34.00	33.96
v_{Corot}^d	km/s	431.5	431.0	430.3	429.1	428.6	427.5	427.0
v_f	km/s	307.9 ± 0.5	330.1 ± 0.7	370.3 ± 0.7	365.0 ± 0.7	374.6 ± 0.7	357.2 ± 0.7	387.1 ± 0.9
v_{SC}^e	km/s	8.18	8.19	8.20	8.22	8.22	8.24	8.25
v_f/v_{Corot}	%	71.35	76.58	86.05	85.06	87.40	83.56	90.66
$n(H^+) \times 10^4$	cm^{-3}	69 ± 4	105 ± 2	99 ± 2	134 ± 2	122 ± 2	129 ± 2	122 ± 2
$n(O^+) \times 10^4$	cm^{-3}	823 ± 62	399 ± 19	433 ± 31	295 ± 19	291 ± 19	291 ± 20	136 ± 18
$n(O^{2+}) \times 10^4$	cm^{-3}	419 ± 12	128 ± 4	192 ± 6	141 ± 4	124 ± 4	129 ± 4	115 ± 4
$n(O^{3+}) \times 10^4$	cm^{-3}	40 ± 6	11 ± 2	15 ± 3	13 ± 2	15 ± 2	14 ± 2	13 ± 2
$n(Na^+) \times 10^4$	cm^{-3}	145 ± 16	55 ± 6	127 ± 8	86 ± 5	100 ± 6	83 ± 6	71 ± 5
$n(S^+) \times 10^4$	cm^{-3}	216 ± 12	101 ± 5	171 ± 7	155 ± 5	129 ± 5	128 ± 5	103 ± 5
$n(S^{2+}) \times 10^4$	cm^{-3}	2324 ± 65	582 ± 20	1229 ± 32	922 ± 20	888 ± 20	892 ± 21	742 ± 19
$n(S^{3+}) \times 10^4$	cm^{-3}	696 ± 19	214 ± 6	246 ± 8	191 ± 5	192 ± 5	202 ± 5	145 ± 5
$n(O^+)/n(S^{2+})$		0.35 ± 0.03	0.68 ± 0.04	0.35 ± 0.03	0.32 ± 0.02	0.33 ± 0.02	0.33 ± 0.02	0.18 ± 0.02
$n(O^{n+})/n(S^{n+})^f$		0.40 ± 0.02	0.60 ± 0.03	0.39 ± 0.02	0.35 ± 0.02	0.36 ± 0.02	0.36 ± 0.02	0.27 ± 0.02
$n_i \times 10^4$	cm^{-3}	4731 ± 95	1595 ± 30	2512 ± 47	1937 ± 29	1860 ± 30	1868 ± 31	1447 ± 28
$n_e \times 10^4^g$	cm^{-3}	8946 ± 158	2756 ± 49	4454 ± 77	3408 ± 49	3286 ± 49	3321 ± 51	2620 ± 46
$n(H^+)/n_e$	%	0.77 ± 0.04	3.82 ± 0.09	2.22 ± 0.06	3.94 ± 0.07	3.70 ± 0.07	3.88 ± 0.08	4.65 ± 0.10
$n(O^+)/n_e$	%	9.20 ± 0.72	14.47 ± 0.75	9.73 ± 0.71	8.66 ± 0.57	8.86 ± 0.60	8.77 ± 0.61	5.20 ± 0.69
$n(O^{2+})/n_e$	%	4.68 ± 0.16	4.65 ± 0.18	4.31 ± 0.15	4.14 ± 0.13	3.76 ± 0.12	3.90 ± 0.13	4.40 ± 0.16
$n(O^{3+})/n_e$	%	0.45 ± 0.06	0.41 ± 0.09	0.34 ± 0.07	0.38 ± 0.06	0.45 ± 0.07	0.42 ± 0.07	0.48 ± 0.09
$n(Na^+)/n_e$	%	1.62 ± 0.18	1.98 ± 0.22	2.85 ± 0.20	2.53 ± 0.16	3.05 ± 0.17	2.49 ± 0.17	2.70 ± 0.20
$n(S^+)/n_e$	%	2.41 ± 0.14	3.65 ± 0.21	3.84 ± 0.17	4.56 ± 0.16	3.92 ± 0.16	3.85 ± 0.16	3.93 ± 0.19
$n(S^{2+})/n_e$	%	25.98 ± 0.86	21.13 ± 0.81	27.59 ± 0.87	27.05 ± 0.71	27.02 ± 0.74	26.86 ± 0.76	28.31 ± 0.88
$n(S^{3+})/n_e$	%	7.78 ± 0.25	7.75 ± 0.25	5.52 ± 0.19	5.59 ± 0.16	5.85 ± 0.17	6.08 ± 0.18	5.55 ± 0.20
T_{PROTON}	keV	9.55 ± 0.51	5.20 ± 0.10	4.80 ± 0.15	3.69 ± 0.06	6.24 ± 0.10	3.99 ± 0.07	4.05 ± 0.08
T_{HEAVY}	keV	1.21 ± 0.06	3.84 ± 0.09	3.95 ± 0.49	5.98 ± 0.11	4.75 ± 0.09	6.60 ± 1.09	12.94 ± 0.31
κ_{proton}^h	keV	3.50	3.50	3.50	3.50	3.50	3.50	3.50
κ_{heavy}	keV	2.11 ± 1.56	1.75 ± 1.55	1.75 ± 1.54	1.77 ± 1.56	1.81 ± 1.57	1.68 ± 1.54	1.63 ± 1.58
$\langle T_i \rangle^i$	keV	1.21 ± 0.05	3.84 ± 0.12	3.96 ± 0.33	5.97 ± 0.15	4.75 ± 0.12	6.59 ± 0.72	12.92 ± 0.41
$P_{THERMAL}$	nPa	0.101 ± 0.045	0.100 ± 0.008	0.160 ± 0.029	0.181 ± 0.014	0.144 ± 0.012	0.192 ± 0.024	0.283 ± 0.025
P_{MAG}	nPa	0.030	0.029	0.005	0.003	0.011	0.030	0.030
β		3.3 ± 1.5	3.4 ± 0.3	33.5 ± 6.0	61.1 ± 4.8	12.9 ± 1.1	6.4 ± 0.8	9.6 ± 0.8

^aRadial distance from the planet in spherical coordinate system. ^bMagnetic latitude from VIP4+CAN model. ^cAxial distance from Jupiter's rotational axis.^dRigid corotation speed ($12.57 \times \rho[R_J]$ km/s). ^eSpacecraft speed in a Jupiter centered system based on J2000 (see Wilson, 2017). ^fRatio of total oxygen ions to sulfur ions where $1 \leq n \leq 3$. ^gElectron density obtained from ion densities with quasi-neutrality assumption ($n_e = \sum n_i q_i$). ^hProton kappa indices are fixed in this study based on Collier and Hamilton (1995). ⁱAverage ion temperature $\langle T_i \rangle = \sum n_i m_i T_i / \rho$, where $\rho = \sum n_i m_i$ is the total mass density.

of charges for the i th ion. These density ratios do not add up to 100% because it is the mixing ratio within the total electron number density not the total ion number density (electrons will have a higher number density than ions since some ions have multiple charges such as S^{2+}). We use ion mixing ratios throughout this paper to follow the convention that has been used in previous Jupiter plasma papers (e.g., Bagenal et al., 2017).

The density of the ions are dominated by O^+ and S^{2+} ($M/Q \approx 16 \text{ amu}/q$) and their relative ratio varies among intervals as shown in the second panel from the top of Figure 11. The number density ratio between total

Table 3*Forward Model Results for Intervals H Through N During the Juno orbit 7 Plasma Sheet Crossing*

Parameters	Unit	H	I	J	K	L	M	N
R^a	R_J	35.61	35.56	35.43	35.37	35.31	35.24	35.16
$MLat^b$	Deg	−1.83	−1.65	−0.92	−0.52	−0.01	0.70	1.54
ρ^c	R_J	33.87	33.82	33.71	33.66	33.61	33.54	33.47
v_{Corot}^d	km/s	425.8	425.2	423.8	423.2	422.5	421.7	420.8
v_f	km/s	350.9 ± 1.4	350.8 ± 2.1	373.8 ± 1.2	388.6 ± 2.6	402.6 ± 1.6	365.4 ± 1.9	360.5 ± 1.5
v_{SC}^e	km/s	8.27	8.28	8.30	8.31	8.32	8.33	8.34
v_f/v_{Corot}	%	82.40	82.49	88.21	91.84	95.28	86.67	85.67
$n(H^+) \times 10^4$	cm^{-3}	110 ± 1	120 ± 1	134 ± 2	113 ± 2	135 ± 1	137 ± 1	126 ± 1
$n(O^+) \times 10^4$	cm^{-3}	127 ± 12	111 ± 11	203 ± 21	230 ± 24	233 ± 18	160 ± 15	110 ± 12
$n(O^{2+}) \times 10^4$	cm^{-3}	86 ± 3	65 ± 3	126 ± 4	124 ± 5	132 ± 3	112 ± 3	85 ± 3
$n(O^{3+}) \times 10^4$	cm^{-3}	10 ± 2	9 ± 2	14 ± 3	11 ± 3	14 ± 2	13 ± 2	10 ± 2
$n(Na^+) \times 10^4$	cm^{-3}	46 ± 4	24 ± 4	71 ± 6	57 ± 6	62 ± 5	52 ± 5	41 ± 4
$n(S^+) \times 10^4$	cm^{-3}	69 ± 4	32 ± 4	98 ± 5	73 ± 7	100 ± 5	88 ± 5	65 ± 4
$n(S^{2+}) \times 10^4$	cm^{-3}	402 ± 13	219 ± 12	675 ± 22	505 ± 25	566 ± 19	488 ± 16	410 ± 13
$n(S^{3+}) \times 10^4$	cm^{-3}	107 ± 4	70 ± 3	154 ± 5	128 ± 6	138 ± 4	123 ± 4	98 ± 3
$n(O^+)/n(S^{2+})$		0.32 ± 0.03	0.51 ± 0.06	0.30 ± 0.03	0.45 ± 0.05	0.41 ± 0.03	0.33 ± 0.03	0.27 ± 0.03
$n(O^{n+})/n(S^{n+})^f$		0.39 ± 0.02	0.58 ± 0.04	0.37 ± 0.02	0.52 ± 0.04	0.47 ± 0.03	0.41 ± 0.02	0.36 ± 0.02
$n_i \times 10^4$	cm^{-3}	957 ± 20	651 ± 18	$1,474 \pm 32$	$1,241 \pm 37$	$1,379 \pm 27$	$1,172 \pm 23$	945 ± 19
$n_e \times 10^4^g$	cm^{-3}	$1,680 \pm 33$	$1,093 \pm 29$	$2,610 \pm 52$	$2,150 \pm 61$	$2,381 \pm 44$	$2,043 \pm 38$	$1,656 \pm 32$
$n(H^+)/n_e$	%	6.56 ± 0.15	11.00 ± 0.31	5.14 ± 0.13	5.28 ± 0.18	5.66 ± 0.12	6.70 ± 0.14	7.59 ± 0.16
$n(O^+)/n_e$	%	7.56 ± 0.76	10.16 ± 1.07	7.77 ± 0.80	10.68 ± 1.14	9.78 ± 0.76	7.85 ± 0.74	6.65 ± 0.75
$n(O^{2+})/n_e$	%	5.11 ± 0.21	5.97 ± 0.29	4.83 ± 0.19	5.77 ± 0.29	5.53 ± 0.17	5.48 ± 0.19	5.16 ± 0.20
$n(O^{3+})/n_e$	%	0.61 ± 0.13	0.84 ± 0.18	0.53 ± 0.11	0.53 ± 0.15	0.58 ± 0.09	0.63 ± 0.10	0.62 ± 0.12
$n(Na^+)/n_e$	%	2.73 ± 0.25	2.22 ± 0.36	2.71 ± 0.23	2.65 ± 0.31	2.61 ± 0.20	2.54 ± 0.23	2.45 ± 0.23
$n(S^+)/n_e$	%	4.08 ± 0.24	2.91 ± 0.36	3.76 ± 0.22	3.39 ± 0.33	4.19 ± 0.23	4.29 ± 0.25	3.95 ± 0.24
$n(S^{2+})/n_e$	%	23.95 ± 0.91	20.03 ± 1.19	25.85 ± 0.98	23.49 ± 1.34	23.79 ± 0.90	23.87 ± 0.89	24.73 ± 0.91
$n(S^{3+})/n_e$	%	6.38 ± 0.26	6.40 ± 0.34	5.89 ± 0.24	5.97 ± 0.33	5.79 ± 0.20	6.00 ± 0.22	5.90 ± 0.23
T_{PROTON}	keV	4.26 ± 0.07	7.05 ± 0.09	3.10 ± 0.07	5.19 ± 0.12	4.40 ± 0.07	4.17 ± 0.06	4.69 ± 0.06
T_{HEAVY}	keV	4.34 ± 0.12	11.30 ± 0.44	9.36 ± 0.25	7.13 ± 0.24	25.53 ± 0.54	8.66 ± 0.22	9.17 ± 0.25
κ_{proton}^h	keV	3.50	3.50	3.50	3.50	3.50	3.50	3.50
κ_{heavy}	keV	2.29 ± 1.69	1.75 ± 1.70	1.73 ± 1.64	2.71 ± 3.17	1.68 ± 1.58	2.38 ± 2.58	1.86 ± 1.70
$\langle T_i \rangle^i$	keV	4.34 ± 0.15	11.27 ± 0.59	9.34 ± 0.33	7.13 ± 0.36	25.46 ± 5.83	8.64 ± 0.29	9.15 ± 0.32
$P_{THERMAL}$	nPa	0.066 ± 0.004	0.110 ± 0.005	0.208 ± 0.017	0.138 ± 0.014	0.519 ± 0.105	0.153 ± 0.008	0.130 ± 0.007
P_{MAG}	nPa	0.055	0.061	0.024	0.015	0.003	0.005	0.038
β		1.2 ± 0.1	1.8 ± 0.1	8.5 ± 0.7	9.2 ± 0.9	200.3 ± 40.7	29.3 ± 1.6	3.4 ± 0.2

^aRadial distance from the planet in spherical coordinate system. ^bMagnetic latitude from VIP4+CAN model. ^cAxial distance from Jupiter's rotational axis.^dRigid corotation speed ($12.57 \times \rho[R_J]$ km/s). ^eSpacecraft speed in a Jupiter centered system based on J2000 (see Wilson, 2017). ^fRatio of total oxygen ions to sulfur ions where $1 \leq n \leq 3$. ^gElectron density obtained from ion densities with quasi-neutrality assumption ($n_e = \sum n_i q_i$). ^hProton kappa indices are fixed in this study based on Collier and Hamilton (1995). ⁱAverage ion temperature $\langle T_i \rangle = \sum n_i m_i T_i / \rho$, where $\rho = \sum n_i m_i$ is the total mass density.

oxygen ions (O^{n+}) to total sulfur ions (S^{n+}) follows a similar trend as the O^+ and S^{2+} ratio but is slightly lower in most of the intervals.

The flow speed of the plasma is always less than the rigid corotation speed ($=12.57 \times \rho[R_J]$ km/s, where ρ is the axial distance from Jupiter's rotational axis). The proton temperature and the heavy ion temperature varies over the intervals with no specific trends. The thermal pressure is mainly driven by the O^+ and S^{2+} densities and is much higher than the magnetic pressure (derived from the Juno MAG data, Connerney et al., 2017). The plasma β (a ratio of the plasma pressure to the magnetic pressure) is typically greater than

1 and sometimes reaches ~ 200 which is due to high plasma pressure and weakening of the magnetic field strength at the magnetic equator.

6. Discussion

6.1. Number Density

The number density estimates of jovian magnetospheric ions using Voyager Plasma Science (PLS) (Bridge et al., 1977) have been presented in several studies (Bagenal & Sullivan, 1981; Bagenal et al., 2017; Bodisch et al., 2017; Dougherty et al., 2017; McNutt et al., 1981). They selected intervals where the peaks in the count- E/Q distributions were clearly resolvable (i.e., cold plasma). These were well described by convected Maxwell-Boltzmann distributions with eight different ion species (H^+ , O^+ , O^{2+} , Na^+ , S^+ , S^{2+} , S^{3+} , and SO_2^+). However, Voyager PLS only measured the E/Q of ions up to 5.95 keV/q , likely missing the high-energy tails of the energy distribution. On the other hand, Voyager 2's Low Energy Charged Particle Experiment (LECP) (Krimigis et al., 1977) measured energy distribution from $\sim 15 \text{ keV}$ to $\sim 150 \text{ MeV}$ for protons and heavier ions. Kane et al. (1992) and Kane et al. (1995) employed convected kappa distributions for two species (H^+ and S^{n+} with unknown charge states) to describe the ion distributions in the energy range from 30 keV to 5 MeV . Both PLS and LECP on Voyager only measured E/Q or E and could not measure M/Q and therefore had limitations on analyzing the plasma parameters.

The Galileo spacecraft was the first orbiter around Jupiter and extensively measured the plasma distributions near the centrifugal equator. The plasma instrument (PLS) on Galileo measured E/Q from 0.9 to 52 keV/q along with M/Q measurements (Frank et al., 1992). However, the mass resolution of the instrument was not sufficient to separate heavy ion species from the data (Frank & Paterson, 2000) and had a low signal-to-noise ratio on their TOF data (Bagenal et al., 2016). The analysis of plasma properties had to use fixed ion composition based on Cassini UVIS (Esposito et al., 2004) measurements of torus emission lines and a physical chemistry model from Delamere et al. (2005) (see Bagenal et al., 2016). The Galileo Energetic Particles Detector instrument measured ions from $\sim 20 \text{ keV}$ to $\sim 55 \text{ MeV}$. Kane et al. (1999) used three convected kappa distributions (H^+ , O^{n+} , and S^{n+} with unknown charge states). Clark et al. (2016) introduced a charge state determination technique on energetic ($\sim 0.3\text{--}5 \text{ MeV}$) oxygen and sulfur ions when there is an injection event. They found that the most probable charge states of oxygen and sulfur ions are single and double charges, respectively. However, the relative abundances between O^+ and S^{2+} are still not well understood considering that the ion number densities are mostly from thermal population (e.g., Bagenal & Sullivan, 1981; McNutt et al., 1981).

Although Ulysses SWICS (Gloeckler et al., 1992) and Cassini MIMI/CHEMS (Krimigis et al., 2004) could immediately separate O^+ and S^{2+} from their measurements, the statistics are insufficient (both spacecraft only had a single flyby for gravity assist where the closest approach to Jupiter for Ulysses spacecraft was at $6.31 R_J$ and Cassini spacecraft at $136 R_J$) to understand how their relative abundances change throughout the magnetosphere. JADE-I covers a much wider energy range (0.01 up to 46.2 keV/q) of the distribution than Voyager PLS and provides M/Q measurements with higher mass resolution ($2.5 \leq M/\Delta M \leq 11$) than Galileo PLS ($2 \leq M/\Delta M \leq 4.1$).

A direct comparison of ion number density ratio results from this study to Dougherty et al. (2017) and Bodisch et al. (2017) show that we observe slightly more S^{2+} than O^+ with similar mixing ratios for other minor ion species except SO_2^+ . Both studies (this study and Dougherty et al., 2017) have limitations on estimating O^+ and S^{2+} due to the limited information provided by the measurements. This study had to depend on simulation work (i.e., JIST) due to JADE-I's insufficient mass resolution to separate O^+ and S^{2+} and Dougherty et al. (2017) had to assume ion mixing ratios from Delamere et al. (2005) due to the absence M/Q information from Voyager PLS.

At $\sim 36 R_J$, we also identify a weak signal of O^{3+} but do not see any clear signals for SO_2^+ (64 amu/q) from JADE-I during the selected intervals. This is probably because these molecular ions quickly dissociate as they move away from the source point. Bodisch et al. (2017) reports they were only observed between 5.07 and $5.31 R_J$.

6.2. O^+ and S^{2+} ($M/Q \approx 16 \text{ amu/q}$)

Obtaining in situ relative abundances between O^+ and S^{2+} has been a challenging issue due to the similarities in their E/Q and M/Q . The plasma instruments flown to date at Jupiter have been unable to resolve them in the measurements except for the Ulysses SWICS and Cassini MIMI/CHEMS measurements. Their relative

abundances are important since they are the major constituents of the ions governing the magnetospheric dynamics at Jupiter.

In this study, we developed JIST, which is based on a detailed instrument response (that includes carbon foil effects) to determine the ion distributions of O^+ and S^{2+} . JIST shows that O^+ and S^{2+} have different shaped count-TOF distributions caused by their different interactions with the carbon foil (mainly the charge state modification) and strong nonlinear electric field in JADE-I's TOF section (see Appendix for details). This model was used to calculate the instrument response function that is an essential part of the forward model presented in this study.

The results shown in Figure 11 indicate that the relative abundance of O^+ and S^{2+} varies from 0.2 to 0.7 (mean value of 0.37 ± 0.12), accounting for 62–69% of the iogenic ions (i.e., O, S, and Na ions) and 50–66% of the Jovian magnetospheric ion number density. The number density ratios between total oxygen ions (O^{n+}) to total sulfur ions (S^{n+}) are 0.2–0.6 (mean value of 0.41 ± 0.09). This value is much lower than the 1.02 value derived by the Delamere et al. (2005) physical chemistry model. However, the physical chemistry model only can cover radial distances up to $\sim 10 R_J$ and the 1.02 value is obtained assuming freezing-in of the charge state. This assumption is used in previous studies (e.g., Bagenal et al., 2016, 2017; Bodisch et al., 2017; Dougherty et al., 2017) to obtain ion properties from Voyager PLS and Galileo PLS.

However, we have only analyzed one plasma sheet crossing in this study and at a different radial distance. An investigation with more plasma sheet crossing data will be conducted in the future. As the spacecraft's equatorial crossing distance will get closer to the planet in the later phase of the mission, we should be able to directly compare to Delamere et al. (2005).

6.3. Flow Speed

The flow speed obtained in this study ranges from 71–95% of the rigid corotation speed. These results are higher than previously reported values in this region (e.g., Dougherty et al., 2017; Kane et al., 1995; McNutt et al., 1981). The highest flow speed is observed in interval L which coincides with when Juno is at a magnetic latitude (MLat) of near zero and the magnitude of B_r is smallest (see Figure 11). Dougherty et al. (2017) reports that the significant breakdown of corotation occurs at $9\text{--}10 R_J$ and the flow speed reaches an asymptotic value of $\sim 225 \text{ km/s}$ beyond $\sim 20 R_J$. However, that analysis was heavily dependent on plasma populations having a cold enough temperature where the peak of each ion distributions was resolvable in the Voyager PLS data. Dougherty et al. (2017) pointed out that these cold ion populations may not be the dominant contributors to the ambient plasma flow. The flow speed was also derived by Krupp et al. (2001) using the Galileo EPD data covering $6\text{--}142 R_J$ and all local times. They used a power law for proton, oxygen, and sulfur ions to obtain spectral index. The flow speeds obtained from this study are comparable to Krupp et al. (2001) at similar local times.

Comparing flow speed to the Alfvén (magnetic field information from Juno MAG) speed, the flow speed is always higher than the Alfvén speed (see Figure 12). We also computed the sound speed with $\sqrt{\gamma \cdot (\sum n_i T_i + n_e T_e) / (\sum n_i m_i + n_e m_e)}$ under assumptions of $\gamma=5/3$ (monatomic gas), quasi-neutrality (i.e., $n_e = \sum n_i q_i$), and the electron temperature being identical to the average ion temperature ($T_e = \langle T_i \rangle = \sum n_i m_i T_i / \sum n_i m_i$). The average ion temperature varies from 1 to 10 keV. The sound and magnetosonic speed are higher than the flow speed for intervals G, I, J, L, M, and N and lower than flow speed for rest of the intervals (Figure 12). Considering that the electron temperature is typically lower than the ions in the jovian plasma sheet (Voyager PLS observed electron temperatures as low as 50 eV at $40 R_J$, see Scudder & Bridge, 1981), the sound speed can be much lower than the our estimates suggesting supersonic flow in the plasma sheet.

The results shown in this study indicate that the hot plasma populations have higher flow speed than the cold blobs observed by Voyager PLS. However, since this study only examines plasma parameters for one plasma sheet crossing by Juno (mostly hot plasma), it is difficult to determine whether the hot or cold plasma is more representative of the plasma sheet populations in the jovian magnetosphere.

6.4. Temperature

The temperature obtained in this study cannot be compared to Voyager PLS analysis results due to the difference in the selection of data (cold plasma for Voyager PLS and hot plasma in this study) and the difference in E/Q range for Voyager PLS and Juno JADE-I. Instead, we compare the results to the Voyager 2 LECP results

Figure 12. Comparisons between flow speed and other characteristic speeds (gray: rigid corotation, magenta: magnetosonic, blue: sound, and purple: Alfvén). Sound speed was calculated under assumption of quasi-neutrality and electron temperature being identical to the average ion temperature.

that used two-species (H^+ and S^{n+} with unknown charge states) convected kappa distributions (Kane et al., 1992, 1995).

The proton temperature ranges from $3 < T_{\text{PROTON}} [\text{keV}] < 10$, while Kane et al. (1992, 1995) found that the proton temperature is nearly constant with ~ 20 and 10 keV for the dayside and nightside of the magnetosphere, respectively. This difference may be due to the local time effects as the data presented in this study are sampled at $\sim 04:16$ local time (predawn sector), a region which both Voyager 1 and Voyager 2 did not encounter.

The heavy ion temperature in this study ranges from $1 < T_{\text{HEAVY}} [\text{keV}] < 25$ and is highly variable throughout the selected intervals. This result is slightly lower than the temperature ($\sim 30 \text{ keV}$ at $36 R_J$) shown in Figure 4 of Kane et al. (1995), and we think this is due to the difference in the number of ion species in the analysis.

6.5. Plasma β

The pressure of the plasma sheet is mainly dominated by the ion thermal pressure. The magnetic pressure is relatively low compared to the ion thermal pressure due to the decrease in the magnetic field strength (mainly the radial component of the magnetic field vector) at the plasma sheet crossing. Because of this, the resulting plasma β is very high, ranging from $1 < \beta < 200$. Thus, at these regions plasma forces dominate resulting in a nondipolar magnetic field.

6.6. Kappa (Spectral Index)

The kappa value represents the statistical correlation between particles in a stationary state system. The value of kappa can range from 1.5 to ∞ where the smallest value of kappa indicates a highly correlated system (or far-equilibrium) and the largest value indicates a noncorrelated system (near-equilibrium) (Livadiotis & McComas, 2011). In this study, we fixed κ_{PROTON} to 3.5 due to the presence of the high-energy populations that typically extends to the top of JADE-I's E/Q range. Different fixed kappa values result in different temperatures (e.g., larger temperatures for smaller κ_{PROTON} and smaller temperatures for larger κ_{PROTON} , see Figure 13).

Collier and Hamilton (1995) obtained a linear relationship between the proton temperature and kappa value ($\kappa_{\text{PROTON}} = 0.08 [keV^{-1}] \cdot T_{\text{PROTON}} [\text{keV}] + 2.86$) from Voyager LECP data during 33 plasma sheet crossings from 26 to $160 R_J$. Plugging the T_{PROTON} derived from this study to the relationship above yields $3.1 < \kappa_{\text{PROTON}} < 3.6$ showing that our fixed proton kappa value is in the range. The heavy kappa value in this study ranges

Figure 13. (left) Variations in (top) proton density, (middle) flow speed, and (bottom) proton temperature for different (fixed) proton kappa values. (right) Forward model results for intervals D and M with various kappa values and comparisons with the in situ data from JADE-I. High-energy populations are shown >20 keV/q.

$1.6 < \kappa_{HEAVY} < 2.7$. Kane et al. (1999) found heavy kappa value of 4.4 during a plasma sheet crossing near $40 R_j$ in the midnight sector, which is slightly higher than the heavy kappa values presented in this study.

6.7. Plasma Parameters for Different Proton Kappa Values

We now discuss how the fixed proton kappa value affects the other plasma parameters by comparing the proton density, flow speed, and proton temperature derived from the forward model results for different values of kappa. Keeping the proton kappa value as a free parameter sometimes resulted in unreasonably small proton kappa values ($\kappa < 1.51$) and large proton temperature ($T \gg 100$ keV). This is likely due to the presence of the high-energy populations in the JADE-I data.

Figure 13 shows the proton density, flow speed, proton temperature T_{PROTON} , and heavy ion temperature T_{HEAVY} for various kappa values for all intervals. The density and flow speed does not change much but the temperatures are significantly different for various kappa values. The temperature is not highly correlated with density, though a positive correlation appears for the initial events, leading to a polytropic index closer to an adiabatic process.

6.8. High-Energy Populations

The JADE-I observations revealed high-energy populations that extend to the top of the instrument's E/Q range. The forward model developed in this study did not model these high-energy populations since it uses a single convected kappa distribution per ion species. These high-energy populations are mainly composed of H^+ , O^+ , and/or S^{2+} (see middle/bottom panel of Figure 5). They are clearly observed when the flow speed is low during the plasma sheet crossing as shown in Figure 10. These ions are not well observed when the flow speeds are high (e.g., intervals K and L) since the flow energy reaches closer to the top of JADE-I's E/Q range. Interval L has the highest flow speed and the lowest magnetic latitude as predicted by the VIP4+CAN model (Connerney et al., 1981, 1998) where the magnitude of B_r is close to 0 (see Figure 13). The rest of the intervals show relatively low flow speed and the high-energy populations are clearly observed in JADE-I data. It is not clear where these populations are originating from and how they affect the core population. However, a rough calculation indicates that these high-energy populations appear to have 4 times the energy of the bulk flow energy (or 2 times the flow speed).

7. Summary and Conclusion

Obtaining the in situ relative abundance of O^+ and S^{2+} have been a challenging task in the Jovian magnetosphere since they have similar mass-per-charge ($M/Q \approx 16 \text{ amu}/q$). They also have similar energy-per-charge (E/Q) considering that all ions share a common flow speed. This paper presented a methodology to resolve the M/Q ambiguity between O^+ and S^{2+} measured by JADE-I using the carbon foil effects and omni-directional averaged convected kappa distributions with a common flow speed. We were able to separate O^+ and S^{2+} because they have very different charge state distributions after transiting the carbon foils. In addition, the large electric potential difference ($\sim 8 \text{ kV}$) in JADE-I's TOF section (Figure 1) creates a distinct count-TOF distribution for O^+ and S^{2+} (Figure A5). We assumed a common proton kappa value κ_{PROTON} of 3.5 based on Collier and Hamilton (1995) due to the existence of high-energy populations and introduced a methodology to derive heavy kappa value by empirically trimming the E/Q -TOF spectra (bottom panel of Figure 5). We also use a common temperature T_{HEAVY} for all heavy ions to reduce the number of free parameters (from 25 down to 13) in the forward model. With the assumptions and limitations discussed in section 4, we summarize our findings as follows:

1. The in situ relative abundances (i.e., the density ratio) for O^+ and S^{2+} varies from 0.2 to 0.7 (mean value of 0.37 ± 0.12). They contribute 62–69% of the iogenic ions and 50–66% of the Jovian magnetospheric ion number density. The density ratio of total oxygen ions to total sulfur ions (O^{n+}/S^{n+}) ranges from 0.2 to 0.6 (mean value of 0.41 ± 0.09) lower than the 1.02 value derived by the Delamere et al. (2005).
2. The density ratios near the equatorial plasma sheet crossing at $\sim 36 R_J$ in the predawn sector ($\sim 04:16$ local time) of the magnetosphere range are $n(O^+)/n_e = 5\text{--}14\%$, $n(S^{2+})/n_e = 20\text{--}28\%$, $n(H^+)/n_e = 1\text{--}11\%$, $n(S^{3+})/n_e = 5\text{--}8\%$, $n(O^{2+})/n_e = 3\text{--}6\%$, $n(S^+)/n_e = 2\text{--}5\%$, $n(Na^+)/n_e = 1\text{--}3\%$, and $n(O^{3+})/n_e = 0.3\text{--}0.8\%$.
3. Flow speed ranges from 71% to 95% of the rigid corotation speed and the speed is always super Alfvénic. This flow speed is higher than the Voyager PLS observations in the same region. This difference is likely due to the type of intervals selected as Voyager PLS analysis chose cold plasma and this study focuses hot plasma distributions.
4. Proton temperature ranges from 3 to 10 keV and the heavy ion temperature ranges from 1 to 25 keV. The proton temperatures are low compared to the Voyager 2 LECP analysis. This could be due to the lower energy limit of the LECP instrument that could not measure ions at thermal energy range.
5. Plasma β ranges from 1 to 200. The variation in the plasma β is mostly due to the change in magnetic field strength. The highest β is observed when the magnetic latitude (from VIP4+CAN) and the magnitude of B_t is smallest during selected intervals.

Appendix A: JADE-I Simulation Tool

A1. Model

The dynamics of transmitted ions are mainly governed by electrostatic fields in the ESA and TOF sections of JADE-I (see Figure 1). Each ion species has a unique count-TOF distribution as they have different mass and charge. They also have distinct carbon foil effects so that the unique TOF measurement can be obtained. When an ion travels through a thin carbon foil, it exits with a charge state that may be different from its initial charge state. This process is called charge state modification. Residual charge states (i.e., charge states after transit through the foil) of the transmitted ions can be either neutral, positive, or negative (the latter only for a limited number of atomic species) with single or multiple charges unless it is neutral. This effect varies for individual atomic species and depends on the incident energy of ions, foil thickness, and the chemical composition of the foil (e.g., Allegrini, Coulter, et al. 2016; Gonin et al., 1994; Kallenbach et al., 1993, 1995). Bürgi et al. (1990), Gonin et al. (1994), Kallenbach et al. (1995), and Gonin et al. (1995) provided a semiempirical model to calculate residual charge states after passage through a thin carbon foil ($1.1\text{--}10 \mu\text{g}/\text{cm}^2$) for various atomic species energy ranging from 0.5 to 5 keV/amu.

The trajectory of an ion varies after transit through the foil (i.e., angular scattering). This effect can be approximated by Meyer's theory (Meyer, 1971), given as

$$\psi_{1/2} E = k_F \quad (\text{A1})$$

where k_F (keV·Deg) is a unique value for a particular particle-foil combination (Funsten et al., 1993), E (keV) is the incident energy of the particle, and $\psi_{1/2}$ (Deg) is the angular scattering half width at half maximum of

the κ -distribution expressed as,

$$F(\theta) = F_0 \left(1 + \frac{\theta^2}{2\sigma^2 b} \right)^{-b}. \quad (\text{A2})$$

Moreover, $\psi_{1/2}$ can be defined as,

$$\psi_{1/2} = \sigma \sqrt{2b(2^{1/b} - 1)} \quad (\text{A3})$$

where σ is the half-width of the distribution at height $F_0(1 + \frac{1}{2b})^{-b}$ and b is the power index.

Ions also lose energy while passing through the foil. Typically, the residual energy (i.e., exit energy of the ion after the carbon foil) of the transmitted ions have dispersion in their distribution due to the energy straggling process (Kalz et al., 1978). Like other carbon foil effects, energy loss varies for individual ions and depends on the physical parameter of the foil (e.g., Allegrini et al., 2006, 2015).

JIST utilizes SIMION (Dahl, 2000), SRIM (Ziegler et al., 2010), charge state yield models (Bürge et al., 1990; Gonin et al., 1994; 1995; Kallenbach et al., 1995), and studies on angular scattering of incident ions (Ebert et al., 2014; Funsten et al., 1993) to compute the carbon foil effects and ion behavior in the JADE-I TOF section. SIMION calculates the ion and electron trajectories everywhere except in the carbon foil. SRIM calculates interaction of atoms with solids which can determine energy loss of transmitted ions. The JIST consists of three main parts:

1. *ESA simulation.* A population of ions with a distribution of energies, angles, masses, and charge states are released toward the entrance aperture of JADE-I. The ions that have the correct E/Q and incident direction pass between the ESA electrodes and are accelerated (at -10 kV) toward the carbon foil at the entrance of the TOF section. The energy, incident angle, position, charge state, and mass of those ions that reach the carbon foil are then passed to the carbon foil simulation.
2. *Carbon foil simulation.* The carbon foil effects on incident ions are computed based on the information obtained from the ESA simulation. The new information for the transmitted ions are residual charge states ($\{Q\}$), residual energy (E_{RES}), scattered angle (θ_s, ϕ_s), and transmitted position (x_T, y_T, z_T).
3. *TOF simulation.* Transmitted ions are released from the carbon foil with information obtained from the carbon foil simulation. The TOF simulation calculates the TOF of the transmitted ions that land on the microchannel plate (MCP) and outer anodes (Figure 1). Histograms for individual residual charge states are then generated to obtain the count-TOF distribution.

Table A1

Comparison of the JADE-I Energy Range for Each Target Ion, Total Energy of Ion at the Carbon Foil (Including -10 kV Post-Acceleration), and the Energy Coverage of the Charge State Model

Target ion	Measurable energy	Energy at carbon foil	Charge state model
H ⁺	0.01–46.2 keV	10.01 – 56.2 keV	★ 0.4–120 keV
He ⁺	0.01–46.2 keV	10.01 – 56.2 keV	◊ 2–20 keV
He ²⁺	0.02–92.4 keV	20.02–112.4 keV	◊ 2–20 keV
C ⁺	0.01–46.2 keV	10.01–56.2 keV	• 6–60 keV
N ⁺	0.01–46.2 keV	10.01–56.2 keV	◊ 7–70 keV
O ⁺	0.02–46.2 keV	10.01–56.2 keV	• 8–80 keV
O ²⁺	0.02–92.4 keV	20.02–112.4 keV	• 8–80 keV
O ³⁺	0.03–138.6 keV	30.03–168.6 keV	• 8–80 keV
Ne ⁺	0.01–46.2 keV	10.01–56.2 keV	◊ 10–101 keV
Na ⁺	0.01–46.2 keV	10.01–56.2 keV	† 11–115 keV
S ⁺	0.01–46.2 keV	10.01–56.2 keV	• 16–160 keV
S ²⁺	0.02–92.4 keV	20.02–112.4 keV	• 16–160 keV
S ³⁺	0.03–138.6 keV	30.03–168.6 keV	• 16–160 keV

Note. Total incident energy of the key ions in this study (i.e., O⁺ and S²⁺) are within the energy coverage of the model. The symbols in the fourth column indicates the model. ◊: Bürge et al. (1990), ★: Gonin et al. (1994), •: Kallenbach et al. (1995), and †: Gonin et al. (1995). JADE-I = Jovian Auroral Distributions Experiment Ion sensor.

Table A2

Fit Parameters of the Residual Energy Distribution Generated From SRIM for Total Incident Energy of 11 keV and Normal Incidence With Respect to the Foil

Species	<i>a</i>	<i>b</i>	<i>c</i>	<i>d</i>	<i>e</i>	<i>f</i>
H	−3.52	4.74×10^2	6.08×10^4	4.27×10^2	6.19×10^4	1.26×10^2
He	−1.38	2.41×10^2	1.39×10^5	8.34×10^2	1.41×10^5	1.80×10^2
C	2.36	1.12×10^2	5.81×10^4	1.72×10^3	6.21×10^4	4.46×10^2
N	2.7	1.02×10	5.77×10^4	1.93×10^3	6.13×10^4	7.17×10^2
O	−1.00	0.50	6.03×10^4	3.01×10	6.03×10^4	3.01×10
Ne	2.56	1.56×10^{13}	1.44×10^5	1.21×10^4	5.88×10^4	4.87×10^2
Na	2.81	1.55×10^{13}	1.48×10^5	1.25×10^4	5.95×10^4	3.83×10^2
S	1.00	1.00	1.24×10^5	3.01×10	1.24×10^5	3.01×10

Note. The effective foil thickness used here is 130 Å.

Each part of the JIST iterates for each energy step and for individual ions. The JIST target species are H⁺, He⁺, He²⁺, C⁺, N⁺, O⁺, O²⁺, O³⁺, Ne⁺, Na⁺, S⁺, S²⁺, and S³⁺. We extended the target ion list from O⁺ and S²⁺, which are the main focus of this study, to validate our model results with in situ and laboratory measurements.

For simplicity, we assumed the following in regard to carbon foil effects and instrument response.

1. JADE-I's carbon foils have flat surfaces (i.e., ignore microscopic variations).
2. Chemical composition, mass density, and thickness of the foils are constant across the whole instrument.
3. Carbon foils used in Ebert et al. (2014), Bürgi et al. (1990), Gonin et al. (1994), Kallenbach et al. (1995), and Gonin et al. (1995) are identical.
4. Angular scattering of an incident ion is equally likely in any direction in a 3-D cone defined by a 2-D scattered angle.
5. All carbon foil effects are independent of initial charge states.
6. The carbon foil effects are not interdependent (e.g., charge state modification is independent of angular scattering).
7. Ions hitting instrument structure other than the carbon foil or MCP are lost.
8. There is no delay or offset due to the electronics.
9. TOF contributions from SEs are ignored.

Further details on each part of the model are provided in following sections.

A2. ESA and TOF Simulation

SIMION is a software package that can calculate the trajectory of a charged particle under an electrostatic field and a magnetostatic field. JIST uses the JADE-I SIMION model (i.e., electrodes configuration) developed in McComas et al. (2017). Figure 1 shows a cross-sectional view of the JADE-I SIMION model on a user-defined *x*-*z* plane. Like any other top-hat ESA, JADE-I transmits ions within a certain *E*/*Q* range for each energy step. The relationship between incident *E*/*Q* (denoted as $E_{\text{IN}}/Q_{\text{IN}}$ hereafter) of the ions and V_{ESA} are given as

$$\frac{E_{\text{IN}}}{Q_{\text{IN}}} [eV/q] = k [eV/V] \cdot \frac{V_{\text{ESA}}}{Q_{\text{IN}}} [V/q], \quad (\text{A4})$$

where *k* is an analyzer constant (4.62 *e* for JADE-I).

JADE-I has an energy resolution $\Delta E_{\text{TOTAL}}/E_{\text{TOTAL}}$ (where E_{TOTAL} is the energy at the foil) of ~18–28% full width at half maximum (depending on energy) to allow ions with $E_{\text{IN}}/Q_{\text{IN}}$ ranging from approximately $0.7 E_{\text{IN}}/Q_{\text{IN}} \leq E_{\text{IN}}/Q_{\text{IN}} \leq 1.3 E_{\text{IN}}/Q_{\text{IN}}$. The ESA simulation takes into account the exact function of the energy resolution as a function of energy. For each energy step (64 for JADE-I), constant voltages are applied to each part of the instrument except the inner ESA (Figure 1). SIMION records ion information when they arrive at the user-defined boundary. For the ESA simulation, our boundary is defined at the carbon foils to limit data to only those ions that land on the foils.

Once carbon foil effects on the incident ions are calculated, proper angular conversion from the 2-D scattered ion angle (β) to the SIMION angle (θ_{RES} , ϕ_{RES}) is required to initiate the TOF simulation. Position information

$(x_{\text{RES}}, y_{\text{RES}}, z_{\text{RES}})$ is then simply transferred vertically downward (negative x -direction) to the bottom of the foil in the SIMION model. The carbon foil-modified incident ions are then mapped through the TOF section using SIMION. The hit location on the MCP and the travel time are recorded. The travel time of the secondary electrons are assumed to be zero for the time-of-flight (TOF) determination. The circular boundary (in the y - z plane) is identical to the outer anodes, and the vertical boundary (along the x axis) is the MCP location. Then we obtain the count-TOF distribution for individual residual charge states for given ion species per energy step. If the peak position of the count-TOF distribution does not match with the laboratory/in situ measurements, the foil thickness is adjusted to regenerate the residual energy distribution using SRIM.

A3. Carbon Foil Simulation

The second part of the model, carbon foil simulations, is further divided into three parts: charge state modification, angular scattering, and energy loss of the incident ions. The total energy (E_{TOTAL}) of an incident ion at the foil depends on its initial charge state (Q_{IN}), given as

$$E_{\text{TOTAL}}[\text{eV}] = Q_{\text{IN}}[e] \cdot (k[\text{eV}/V] \cdot \frac{V_{\text{ESA}}}{Q_{\text{IN}}}[V/e] + 10^4[V]), \quad (\text{A5})$$

where $10^4 V$ represents the post-acceleration in JADE-I. The third column of Table A2 shows the total energy range of the target ions at the foil based on equation (A5). This equation shows that S^{2+} will have twice the total energy of the O^+ if both are detected at the same energy step.

The residual charge states of incident ions are determined by the models presented in Bürgi et al. (1990), Gonin et al. (1994), Kallenbach et al. (1995), and Gonin et al. (1995). These models use carbon foil thickness, incident atomic species, and incident energy as an input and is capable of computing the residual charge states of most of the ions that are found in space plasma. Table A1 shows the measurable energy range of JADE-I for different target ions and the energy range of the charge state yield models. We neglected the initial charge state of the incident ions in the charge state yield models since they experience multiple electronic interactions. The model covers most of the JADE-I energy range, but there are species that require extrapolation in some of the instrument's energy bins (see Table A1). Based on the charge state yield models, there are significant differences in residual charge states between O^+ and S^{2+} in JADE-I's $E_{\text{IN}}/Q_{\text{IN}}$ range (see Figure A1). Similar calculations for the residual charge state were performed for other species that are listed in Table A2. The model takes a fixed ratio of residual charge states for each energy bin and randomly allocates the charges to the ions.

For the angular scattering calculation, this study uses $b=2$ (Lorentzian) in equation A(2), which is a good approximation for the JIST ions. The k_F values for H, He, C, N, O, Ne, and Ar ions are obtained from Ebert

Figure A1. Charge state yield of O^+ and S^{2+} in the $E_{\text{IN}}/Q_{\text{IN}}$ range from 0 to 50 keV/q (JADE-I has $E_{\text{IN}}/Q_{\text{IN}}$ range from 0.01 to 46.2 keV/q). The calculation is based on Kallenbach et al. (1995) and the -10 kV post-acceleration is considered. O^+ yields more negative ions (i.e., $\text{O}^{(-)}$) than positive ions (i.e., $\text{O}^{(n+)}$) and S^{2+} yields more positive ions (i.e., $\text{S}^{(n+)}$) than negative ions (i.e., $\text{S}^{(-)}$) at a given $E_{\text{IN}}/Q_{\text{IN}}$ where $1 \leq n \leq 3$. The count-TOF distribution for O^+ and S^{2+} are different mainly due to the distinct residual charge states between them (O^+ has more negative charge states than S^{2+} after the foil).

Figure A2. Estimated k_F value of Na and S using a simple linear fit to data points obtained from Ebert et al. (2014). Atomic number (Z) and k_F ($\text{keV} \cdot \text{Deg}$) are proportional to each other.

et al. (2014) and the values for Na and S are estimated using a linear fit (k_F versus atomic number Z). Figure A2 shows data points from Ebert et al. (2014) and the interpolated data point. Figure A3 shows an angular scattering distribution of O^+ and S^{2+} at incident energy of $1 \text{ keV}/q$ (outside the instrument). The scattered angle distributions are normalized for a given species per an incident angle to obtain a probabilistic distribution of scattered 2-D angles. JIST uses these distributions to generate cumulative probabilistic distributions.

SRIM can help determine the residual energy of the ions for a given chemical composition, density, and thickness of the carbon foil. However, it is challenging to find the right foil composition, density, and thickness in SRIM to reproduce the correct energy loss and straggling (Allegrini, Coulter, et al. 2016). JIST uses a chemical composition of 12% oxygen and 88% carbon, a mass density of 1.8 g/cm^3 , and a thickness of $90 \text{ \AA}$ (or $1.62 \mu\text{g/cm}^2$) as a starting point then change the foil thickness to refine the model based on the comparisons with the measurements. We calculate the energy loss of each target ion's energy range listed on the

third column of Table A2 with a step size of 0.5 keV . The ESA simulation results show that the incident angle with respect to the carbon foil ranges from 0° to 20° . We use a step size of 0.5° in our SRIM calculations. Figure A4 shows the energy loss of O^+ and S^{2+} at incident energy of $1 \text{ keV}/q$.

We then fit the histogram profile of each ion's incident energy and angle for a probabilistic approach to obtain the residual energy. We find that the best matching fit function for the residual energy profile of all of our target ions is

$$R(E_{\text{RES}}) = a + b \cdot \exp \left[-\frac{1}{2} \cdot \left(\frac{E_{\text{RES}} - c}{d} \right)^2 \right] \cdot \left[1 - \tanh \left(\frac{E_{\text{RES}} - e}{2f} \right) \right], \quad (\text{A6})$$

where $R(E_{\text{RES}})$ is the residual energies distribution that represents the probability distribution of residual energy of an incident ion for a particular atomic species with a given incident angle and energy. a , b , c , d , e , and f are the free parameters and E_{RES} is the residual energy of the incident ion. Table A2 shows fitted parameters a , b , c , d , e , and f for an incident energy of 1 keV with normal incidence (90°) with respect to the carbon foil for H, He, C, N, O, Ne, Na, and S. By normalizing total counts of $R(E_{\text{RES}})$, we obtain a probability

Figure A3. Simulation result of angular scattering of O^+ and S^{2+} based on Meyer's theory (Meyer, 1971) using data points from Ebert et al. (2014) and estimated k_F value of sulfur (see Figure A2). Half width at half maximum ($\psi_{1/2}$) for both species are shown in the middle of the figure.

Figure A4. Simulation results of O^+ and S^{2+} residual energy using a probabilistic approach based on SRIM results for $130 \text{ \AA}$. The residual energy distribution of S^{2+} is wider than O^+ . There is a slight difference in the peak position of the distribution between O^+ and S^{2+} .

distribution for residual energy. The model generates residual energy for individual ions per incident angle through a similar probabilistic approach that is used for scattered angle calculation.

While low-energy ($v \ll v_0$, where v_0 is a first orbital Bohr velocity) heavy ions with high charge states ($Q > 10$) could have charge state-dependent energy loss (Ogawa et al., 1992; Schenkel et al., 1997; Wilhelm et al., 2016; Wilhelm & Möller, 2016), which could limit the use of SRIM, we neglected the initial charge state dependency of the energy loss process. JADE-I has $\sim 10 \text{ kV}$ post-acceleration on incident ions before they arrive at the carbon foils and the expected ion species that JADE-I measures in the Jovian magnetosphere are known to have charge states that are $Q_{IN} \leq 3$ (Bagenal & Delamere, 2011).

Fama, Lantschner, et al. (2000), Fama, Eckardt, et al. (2000), and Famá et al. (2002) show that there is a dependence between energy loss and angular scattering. The model developed in this study assumes no dependence between angular scattering and energy loss for simplicity. The charge state modification process is not strongly connected to the other carbon foil effects (i.e., angular scattering and energy loss, Allegrini, Ebert & Funsten 2016). Thus, the residual charge states calculation for the tar-

get ions is independently performed in our model. Per each ion that reaches the carbon foils, the model simulates 100 different cases using the probabilistic approach to account for random nature (i.e., random energy loss and angular scattering) of the transmitted ions.

A4. Carbon Foil Thickness in the Model

Secondary electrons (SE) provide the start signal to the instrument and ions provide the stop signal. The TOF is the difference between the travel time of the SE and the ion. The model results for a simple SE TOF analysis (i.e., which includes the energy distribution after the foil but not the angular distribution of the SE) based on Allegrini, Coulter, et al. (2016) show that it takes about 2.2 ns (the peak of the SE's count-TOF distribution) to receive a start signal from SE for both the EM and FM. JADE-I's TOF bin has a minimum size of 1.45 ns . The TOF simulation of the model does not include SE effect for simplicity since the energy and angular distribution of the SE are not well characterized for the carbon foil used in JADE-I.

Each carbon foil simulation in the model (i.e., charge state yield, energy loss, and angular scattering) uses foil thickness as an input parameter. The charge state yield model uses foil thickness in $\mu\text{g}/\text{cm}^2$ and SRIM in $\text{\AA}$. For the angular scattering, we use a k_F value that is related to the foil thickness. k_F is unique for each specific projectile and is related to the foil thickness. Considering the similarity in the foil used in Ebert et al. (2014) and in JADE-I's EM and FM, we use a k_F value listed in Ebert et al. (2014) and fix this in the model.

Peak positions in the count-TOF distribution are mainly determined by the residual energy distribution of transmitted ions, making it necessary to first find the SRIM foil thickness. However, the SRIM thickness (i.e., the energy loss processes as a function of incident angle and energy of the ion) differs from the nominal foil thickness due to the grid size of the model and the SE effect. Thus, we call it an 'effective carbon foil thickness' to indicate that it is derived from our simulation. The model results show that the best matching effective foil thicknesses for JADE-I's EM and FM are $\sim 130 \text{ \AA}$ for the SRIM input parameter (or $\sim 2.34 \mu\text{g}/\text{cm}^2$ for $1.8 \text{ g}/\text{cm}^3$ foil density).

Once the effective carbon foil thickness is determined, we then obtain a foil thickness for the charge state-yield model by comparing peak amplitude from the model results to the laboratory measurements. We find that the foil thickness of $1.1 \mu\text{g}/\text{cm}^2$ for the charge state model best represents the simulated O^+ count-TOF distribution compared to the $1 \text{ keV}/q$ laboratory measurement (see Figure A6). Note that this is slightly thicker than the foil thickness shown in Ebert et al. (2014), from which we obtained the k_F values. However, considering that the foil can build an impurity layer that increases the foil thickness by $\geq 0.5 \mu\text{g}/\text{cm}^2$ (e.g., nominal foil thickness foil $0.5 \mu\text{g}/\text{cm}^2$ quoted by the manufacturer can have measured thickness of $1.84 \mu\text{g}/\text{cm}^2$, Allegrini et al., 2006; Funsten et al., 1993), $1.1 \mu\text{g}/\text{cm}^2$ is in the uncertainty range of the foil thickness in Ebert et al. (2014).

Appendix B: Empirical Models

B1. Empirical Peak Correction of the Instrument Response Functions

The mass tracer lines shown on E/Q -TOF spectra in Figures 5–7 are obtained from the following relationship between the measured energy-per-charge E/Q_{MEASURED} [eV/q] and the TOF bin t (note that this is different from TOF band τ),

$$E_T [\text{keV}] = \{ (E/Q_{\text{MEASURED}} [\text{eV}/q] + 10^4 [\text{V}]) \cdot Q [e] \} \cdot 10^{-3}, \quad (\text{B1})$$

$$t = \left(A_0 + \frac{A_1}{\sqrt{E_T [\text{keV}]}} \right) \cdot M [\text{amu}]^{P_0 + P_1 \cdot E_T [\text{keV}]} + (C_0 + C_1 \cdot E_T [\text{keV}] + C_2 \cdot E_T [\text{keV}]^2), \quad (\text{B2})$$

where A_0 , A_1 , C_0 , C_1 , C_2 , P_0 , and P_1 are constants obtained from in-flight calibrations using in situ magnetospheric data that contain multiply charged ion species. A_0 , A_1 , C_0 , C_1 , C_2 , P_0 , and P_1 are

$$\begin{aligned} A_0 &= -0.48, A_1 = 67.68, \\ C_0 &= -1.68, C_1 = -2.986 \times 10^{-3}, C_2 = 9.96 \times 10^{-5}, \\ P_0 &= 0.53, P_1 = -1.498 \times 10^{-4}. \end{aligned} \quad (\text{B3})$$

And the trace of secondary proton (i.e., $\text{H}^{(+)}$) peak position can be obtained by using $E_T [\text{keV}]$ with $Q=0.96$ in equation (B1), $M=2.6$ in equation (B2) with the following constants:

$$\begin{aligned} A_0 &= 36.4, A_1 = 49.4, \\ C_0 &= 17.8, C_1 = -0.287, C_2 = 1.59 \times 10^{-3}, \\ P_0 &= 0.541, P_1 = -0.21. \end{aligned} \quad (\text{B4})$$

These mass tracer functions are used to correct instrument response functions from JIST, which is shown in Figure 6. We note that these mass tracer lines are only valid for ions with charge state $1 \leq Q[e] \leq 3$.

B2. Proton Model

The light ions (H and He) in the simulation results are observed to be much narrower in the width of the count-TOF distribution compared to the JADE-I EM measurements. Since the protons are almost always observed in the Jovian magnetosphere, we develop separate model for protons using Gaussian triplets (each representing distribution of $\text{H}^{(+)}$, $\text{H}^{(0)}$, and $\text{H}^{(-)}$, respectively). The curly brackets represent residual charge states (i.e., charge states after the foil). Then the total count C at energy step E and TOF bin t can be written as

$$\begin{aligned} C_{H^+}(E, t) &= \zeta(E) \cdot \exp \left\{ \frac{[t - X(E)]^2}{2 \cdot U(E)} \right\} + \lambda(E) \cdot \exp \left\{ \frac{[t - Y(E)]^2}{2 \cdot V(E)} \right\} \\ &\quad + \xi(E) \cdot \exp \left\{ \frac{[t - Z(E)]^2}{2 \cdot W(E)} \right\}, \end{aligned} \quad (\text{B5})$$

where $\zeta(E)$, $\lambda(E)$, and $\xi(E)$ are the fractional counts determined by the charge state yield model (Bürge et al., 1990; Gonin et al., 1994, 1995; Kallenbach et al., 1995); $X(E)$, $Y(E)$, and $Z(E)$ ($=0.8 \cdot Y(E)$) are the centroids of the Gaussian derived from equation (B5); and $U(E)$, $V(E)$, and $W(E)$ are the width of each count-TOF distribution.

bution. The best matching width were obtained by comparing it with JADE-I EM laboratory measurements and they are related with the centroids as follows:

$$\begin{aligned} U(E) &= \log_{10}[X(E)/2], \\ V(E) &= \log_{10}[Y(E)], \\ W(E) &= \log_{10}[Z(E)/3]. \end{aligned} \quad (\text{B6})$$

B3. False Coincidences

False coincidences can occur when start and stop signals trigger within the TOF window. They are clearly related with incident flux (see Gilbert et al., 2014). We model these false coincidences based on JADE-E EM calibration data assuming that their functional form is power law distribution in count-TOF space, expressed as

$$FC(E) = \epsilon(E) \cdot t^{\nu}, \quad (\text{B7})$$

where ϵ is the energy-dependent scaling factor derived from ratio between maximum counts of measured count-TOF distribution at energy E to the count of false coincidence distribution (t^{ν}) at TOF bin t where the maximum counts are observed. The ν value for the protons is -2.08 and 0.7 for the non-protons. The modeled false coincidences are added to the JIST results in two separate ways. Proton false coincidences are added in the long TOF range (longer TOF bin than the H^+ tracer lines obtained from equations B2 and (B3)) and the non-protons in the short TOF range (shorter than TOF bin for than tracer lines for a given non-proton ions). The instrument response functions shown in Figure 6 contain these false coincidences.

Appendix C: Model Results

C1. O^+ and S^{2+} ($M/Q \approx 16 \text{ amu}/q$)

We now investigate if the count-TOF distributions of O^+ and S^{2+} can be distinguished from the JADE-I TOF measurement. Figure A5 shows the model results for O^+ and S^{2+} . The model result shows significant differences in the shape of the count-TOF distribution for O^+ and S^{2+} . Their non-neutral residual charge states are different as shown in Figure A1 and they essentially determine the shape of the count-TOF distribution in JADE-I's TOF section as positive ions are decelerated and negative ions/electrons are accelerated toward MCPs due to the $\sim 8 \text{ kV}$ electric potential differences as shown in Figure 1. The residual charge state of O^+ has a significantly larger portion of negative residual charge states (i.e., $\text{O}^{(-)}$) than S^{2+} . Because of this, the count-TOF distribution of O^+ has a short TOF bulge as shown in Figure A5. The distribution of S^{2+} is wider than O^+ since S^{2+} has dispersion in the residue energy distribution (Figure A4) and more positive residual charge states (i.e., $\text{S}^{(n+)}$, Figure A1) than O^+ . As the incident energy (or E_{TOTAL} at the foil) rises, the distributions become narrower, but the difference still exist for O^+ and S^{2+} . Thus, O^+ and S^{2+} can be separated from the TOF measurement of JADE-I.

C2. Comparison With Laboratory Measurement (JADE-I EM)

In this section, we compare the simulation results to the laboratory measurements of the JADE-I Engineering Model (EM) obtained at Southwest Research Institute's Ion Calibration Facility (see Appendix in McComas et al., 2009). The JADE-I EM is functionally identical to the JADE-I Flight Model (FM) and is used for ground support calibration purposes. Our test species were H^+ , He^+ , C^+ , N^+ , O^+ , and Ne^+ at 1, 2, 4, 5, 10, 20, and $30 \text{ keV}/q$. We then simulated the test species using the model with the identical configuration and scaled the simulation results to the peak position of the laboratory measurements for JADE-I's EM. Figure A6 compares the simulation results with the laboratory measurement of various species at $1 \text{ keV}/q$ incident energy (2, 4, 5, 10, 20, and $30 \text{ keV}/q$ comparisons are shown in Figure A7). As shown in Figures A6 and A7 the model is capable of separating the count-TOF distributions by residual charge states for each ion species which helps us to understand the TOF measurement of the instrument. The model shows good agreement in the main peak of the distribution. The count-TOF distributions for higher energies tend to have a single peak for all species. This is because the electric field in JADE-I's TOF section is not strong enough to separate residual charge distributions for ions with higher energies.

Appendix D: TOF Data Background Subtraction

Apart from the false coincidences in JADE-I TOF data, there are background counts that arise from the sources (e.g., penetrating particles and radiation) that are not affected by the electrooptics. Because of this,

Figure A5. Comparison of simulated E/Q -TOF (time-of-flight) spectra between (left column) O^+ and (right column) S^{2+} for $130 \text{ \AA}$. The TOF peaks were corrected using equations B2 and (B3). O^+ has higher counts in negative residual count-TOF distributions due to the difference in charge state yield.

the characteristics of these background counts do not have energy dependency in the measurements. Therefore, the backgrounds appear as flat energy spectra in counts (or power law in differential number flux). The JADE-I TOF data have 93 TOF bands over 64 energy steps covering from 0.01 to 46.2 keV/q . To properly subtract backgrounds from in situ TOF data, a standard form of count-TOF distribution for background must be identified beforehand. This can be obtained from the period when the strong radiation dominated during PJ 1. JADE-I has a separate anode for measuring these background counts (see Figure 12 in McComas et al. 2017). Since the TOF data is a coincidence measurement and the background anode is not a coincidence measurement, it should not be a linear function of the background anode counts. The relation between the TOF count rate (summed over TOF bands) and the background anode count rate can be represented by the

Figure A6. Comparison between JIST results and laboratory measurements of H^+ , He^+ , C^+ , N^+ , O^+ , and Ne^+ at 1 keV/q incident energy. Colored dots on orange line (i.e., JIST results for 130 Å) represent the count-TOF distributions due to the residual charge of the incident ions. Magenta line shows a proton model described in section B2.

following:

$$\begin{aligned} N_{\text{TOF}} &= \alpha B^\beta \cdot e^{-B\gamma}, \\ \alpha &= 4.5 \times 10^{-5}, \\ \beta &= 1.8, \\ \gamma &= 5.8 \times 10^{-6}, \end{aligned} \tag{D1}$$

where B is the counts per second in the background anode, averaged over the energy channel. The accumulation time for the plasma sheet intervals presented in this study is 30 s (one of the low rate science mode) and PJ 1 background example is in 2 s (highest time cadence of JADE-I). The level 3 TOF data are in count rates and they can be converted into counts by multiplying total accumulation time T_{acc} (use the values in

Figure A7. Comparison between simulation model results and laboratory measurements of H^+ , He^+ , C^+ , N^+ , O^+ , and Ne^+ at 2, 4, 5, 10, 20, and 30 keV/q of incident E_{IN}/Q_{IN} . Each column shows species at different energies with fixed time-of-flight ranges. All panels share a common y axis. Color legends are identical to Figure A6.

Table 1 for the plasma sheet intervals presented in this study) and the accumulation time per energy step for JADE-I, $\Delta t [s] = \frac{1}{32} - 2 \times 10^{-3}$.

For every accumulation, the TOF background counts are determined by equation (D1), then the standard form of background count-TOF distribution is scaled and subtracted from the TOF data with the following equation:

$$N[t_{UTC}, E, \tau] = DATA_{TOF}[t_{UTC}, E, \tau] - N_{TOF}[t_{UTC}] \cdot B_{TOF}[E, \tau], \quad (D2)$$

where $DATA_{TOF}$ is the level 3 JADE-I TOF data (in counts per second), t_{UTC} is time step of coordinated universal time (UTC), E is energy step, τ is each TOF band, and B_{TOF} is the standard form of background count-TOF distribution. The result of the background subtraction based on equation (D2) is shown in Figure D1 for the day which includes PJ 1. The method described here does a good job removing the counts at times with high background, but less well for low background times. The data in low TOF bands were removed where the

Figure D1. An example of JADE-I TOF background subtraction during PJ 1 encounter (2016-240T11:50:24 to 2016-240T13:39:40) is shown in E/Q -time spectra (top two panels) and E/Q -TOF spectra. The short TOF range was removed from the data in the bottom plot.

remnant background counts may exist. These TOF ranges are

$$\begin{array}{ll}
0 \leq \tau \leq 14 & \text{for } 0 \leq E \leq 25 \\
0 \leq \tau \leq 13 & \text{for } 26 \leq E \leq 36 \\
0 \leq \tau \leq 12 & \text{for } 37 \leq E \leq 41 \\
0 \leq \tau \leq 11 & \text{for } 42 \leq E \leq 45 \\
0 \leq \tau \leq 10 & \text{for } 46 \leq E \leq 47 \\
0 \leq \tau \leq 9 & \text{for } 48 \leq E \leq 50 \\
0 \leq \tau \leq 8 & \text{for } 51 \leq E \leq 52 \\
0 \leq \tau \leq 7 & \text{for } E = 53 \\
0 \leq \tau \leq 6 & \text{for } 54 \leq E \leq 55 \\
0 \leq \tau \leq 5 & \text{for } 56 \leq E \leq 57 \\
0 \leq \tau \leq 4 & \text{for } E = 58 \\
0 \leq \tau \leq 3 & \text{for } 59 \leq E \leq 60 \\
0 \leq \tau \leq 2 & \text{for } 61 \leq E \leq 63,
\end{array} \tag{D3}$$

where τ is each bin of 93 TOF bands and E is the energy step of JADE-I.

Appendix E: Updates on JADE-I's Geometric Factor and Detection Efficiency

The $G\epsilon$ (the geometric factor $\times$ the detection efficiency) for JADE-I quoted in Wilson et al. (2018) yields higher (approximately by a factor of 4) charge densities (i.e., $\sum n_i q_i$, where n is the number density and q is the number of charges of i th ion) than those derived from Juno Waves (Kurth et al., 2017) when compared to each other during the plasma sheet encounter. Waves can determine the electron density via local plasma frequency. We have revisited $G\epsilon$ with flight calibration data, and the work is presented in McComas et al. (2017). Below is the updated $G\epsilon$ (per pixel or anode) for the JADE-I instrument:

$$G\epsilon(E/Q [eV/q]) = 6.84 \times 10^{-5} + 2.52 \times 10^{-6} \cdot \log_{10}(E/Q [eV/q]) + 6.84 \times 10^{-5} \cdot E/Q [eV/q]^{-0.28}, \quad (E1)$$

where the unit of $G\epsilon$ is $cm^2 \cdot eV / eV \cdot sr$. For the TOF data sets, a factor of 12 should be multiplied to the $G\epsilon$ shown in (E1) since it is a combined product of 12 anodes. The $G\epsilon$ for the JADE-I TOF data is then,

$$G\epsilon(E/Q [eV/q]) = 8.21 \times 10^{-4} + 3.02 \times 10^{-5} \cdot \log_{10}(E/Q [eV/q]) + 8.21 \times 10^{-4} \cdot E/Q [eV/q]^{-0.28}. \quad (E2)$$

Acknowledgments

This work is supported by NASA Contract NNM06AA75C. The authors would like to thank Joshua S. Brody, David W. Cronk, Troy J. Diaz, and Benjamin Rodriguez for supporting JADE-I EM laboratory testing at Southwest Research Institute; Chip Beebe for a valuable advice about the JADE-I SIMION model. The flight data used in this study are the L3 JADE and MAG data in NASA's Planetary Data System (PDS), volumes JNO-J/SW-JAD-3-CALIBRATED-V1.0 and JNO-J-3-FGM-CAL-V1.0, respectively. The Juno spacecraft trajectory information was obtained from Laboratory for Atmospheric and Space Physics website (<http://lasp.colorado.edu/home/mop/missions/juno/trajectory-information/>). George Livadiotis contributed under NNX17AB74G NASA's grant for plasma-magnetic field coupling.

References

- Allegrini, F., Coulter, K., Ebert, R. W., Nicolaou, G., & Poenitzsch, V. Z. (2016). Investigation of the influence of surface composition on the charge state distribution of keV hydrogen exiting thin carbon foils for space plasma instrumentation. *Advances in Space Research*, 57(11), 2420–2426.
- Allegrini, F., Ebert, R., & Funsten, H. (2016). Carbon foils for space plasma instrumentation. *Journal of Geophysical Research: Space Physics*, 121, 3931–3950. <https://doi.org/10.1002/2016JA022570>
- Allegrini, F., Ebert, R. W., Fuselier, S. A., Nicolaou, G., Bedworth, P., Sinton, S., & Trattner, K. J. (2014). Charge state of 1 to 50 keV ions after passing through graphene and ultrathin carbon foils. *Optical Engineering*, 53(2), 24,101–24,101.
- Allegrini, F., Ebert, R. W., Nicolaou, G., & Grubbs, G. (2015). Semi-empirical relationships for the energy loss and straggling of 1–50 keV hydrogen ions passing through thin carbon foils. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 359, 115–119.
- Allegrini, F., McComas, D., Young, D., Berthelier, J.-J., Covinhes, J., Illiano, J.-M., et al. (2006). Energy loss of 1–50 keV H, He, C, N, O, Ne, and Arions transmitted through thin carbon foils. *Review of Scientific Instruments*, 77(4), 44501.
- Allegrini, F., Wimmer-Schweingruber, R. F., Wurz, P., & Bochsler, P. (2003). Determination of low-energy ion-induced electron yields from thin carbon foils. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 211(4), 487–494.
- Bagenal, F., Adriani, A., Allegrini, F., Bolton, S. J., Bonfond, B., Bunce, E. J., et al. (2017). Magnetospheric science objectives of the Juno mission. *Space Science Reviews*, 213, 219–287. <https://doi.org/10.1007/s11214-014-0036-8>
- Bagenal, F., & Delamere, P. A. (2011). Flow of mass and energy in the magnetospheres of Jupiter and Saturn. *Journal of Geophysical Research*, 116, 5209. <https://doi.org/10.1029/2010JA016294>
- Bagenal, F., Dougherty, L., Bodisch, K., Richardson, J., & Belcher, J. (2017). Survey of Voyager plasma science ions at Jupiter: 1. Analysis method. *Journal of Geophysical Research: Space Physics*, 122, 8241–8256. <https://doi.org/10.1002/2016JA023797>
- Bagenal, F., & Sullivan, J. D. (1981). Direct plasma measurements in the Io torus and inner magnetosphere of Jupiter. *Journal of Geophysical Research*, 86(A10), 8447–8466.
- Bagenal, F., Wilson, R. J., Siler, S., Paterson, W. R., & Kurth, W. S. (2016). Survey of Galileo plasma observations in Jupiter's plasma sheet. *Journal of Geophysical Research: Planets*, 121, 871–894. <https://doi.org/10.1002/2016JE005009>
- Bevington, P. R., & Robinson, D. K. (2003). *Data reduction and error analysis*. Boston: McGraw-Hill.
- Bodisch, K., Dougherty, L., & Bagenal, F. (2017). Survey of Voyager plasma science ions at Jupiter: 3. Protons and minor ions. *Journal of Geophysical Research: Space Physics*, 122, 8277–8294. <https://doi.org/10.1002/2017JA024148>
- Bolton, S. J., Adriani, A., Adumitroaie, V., Allison, M., Anderson, J., Atreya, S., et al. (2017). Jupiter's interior and deep atmosphere: The initial pole-to-pole passes with the Juno spacecraft. *Science*, 356(6340), 821–825.
- Bolton, S., Lunine, J., Stevenson, D., Connerney, J., Levin, S., Owen, T., et al. (2017). The Juno mission. *Space Science Reviews*, 213(1–4), 5–37.
- Bridge, H., Belcher, J., Butler, R., Lazarus, A., Mavretic, A., Sullivan, J., et al. (1977). The plasma experiment on the 1977 Voyager mission. *Space Science Reviews*, 21(3), 259–287.
- Bridge, H., Belcher, J., Lazarus, A., Sullivan, J., McNutt, R., Bagenal, F., et al. (1979). Plasma observations near Jupiter: Initial results from Voyager 1. *Science*, 204(4396), 987–991.
- Bürki, A., Oetliker, M., Bochsler, P., Geiss, J., & Coplan, M. A. (1990). Charge exchange of low-energy ions in thin carbon foils. *Journal of Applied Physics*, 68(6), 2547–2554.
- Clark, G., Mauk, B., Paranicas, C., Kollmann, P., & Smith, H. (2016). Charge states of energetic oxygen and sulfur ions in Jupiter's magnetosphere. *Journal of Geophysical Research: Space Physics*, 121, 2264–2273. <https://doi.org/10.1002/2015JA022257>
- Collier, M. R., & Hamilton, D. (1995). The relationship between kappa and temperature in energetic ion spectra at Jupiter. *Geophysical Research Letters*, 22(3), 303–306.
- Connerney, J., Acuna, M., & Ness, N. (1981). Modeling the Jovian current sheet and inner magnetosphere. *Journal of Geophysical Research*, 86(A10), 8370–8384.

- Connerney, J., Acuna, M., Ness, N., & Satoh, T. (1998). New models of Jupiter's magnetic field constrained by the Io flux tube footprint. *Journal of Geophysical Research*, 103(A6), 11,929–11,939.
- Connerney, J., Adriani, A., Allegrini, F., Bagenal, F., Bolton, S., Bonfond, B., et al. (2017). Jupiter's magnetosphere and aurorae observed by the Juno spacecraft during its first polar orbits. *Science*, 356(6340), 826–832.
- Connerney, J., Benn, M., Bjarno, J., Denver, T., Espley, J., Jorgensen, J., et al. (2017). The Juno magnetic field investigation. *Space Science Reviews*, 213(1-4), 39–138.
- Dahl, D. A. (2000). SIMION for the personal computer in reflection. *International Journal of Mass Spectrometry*, 200(1), 3–25.
- Delamere, P., Bagenal, F., & Steffl, A. (2005). Radial variations in the Io plasma torus during the Cassini era. *Journal of Geophysical Research*, 110, A12223. <https://doi.org/10.1029/2005JA011251>
- Dougherty, L., Bodisch, K., & Bagenal, F. (2017). Survey of Voyager plasma science ions at Jupiter: II. Heavy ions. *Journal of Geophysical Research: Space Physics*, 122, 8257–8276. <https://doi.org/10.1002/2017JA024053>
- Ebert, R. W., Allegrini, F., Fuselier, S. A., Nicolaou, G., Bedworth, P., Sinton, S., & Trattner, K. J. (2014). Angular scattering of 1–50 keV ions through graphene and thin carbon foils: Potential applications for space plasma instrumentation. *Review of Scientific Instruments*, 85(3), 33302.
- Esposito, L. W., Barth, C. A., Colwell, J. E., Lawrence, G. M., McClintock, W. E., Stewart, A. I. F., et al. (2004). The Cassini ultraviolet imaging spectrograph investigation. *Space Science Reviews*, 115(1-4), 299–361.
- Famá, M., Eckardt, J., Lantschner, G., & Arista, N. (2000). Isotopic effects in the angular dependence of the energy loss and angular distribution of protons and deuterons in Al foils at low energies. *Physical Review A*, 62(6), 62901.
- Famá, M., Lantschner, G., Eckardt, J., Arista, N., Gayone, J., Sanchez, E., & Lovey, F. (2002). Energy loss and angular dispersion of 2–200 keV protons in amorphous silicon. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 193(1), 91–96.
- Famá, M., Lantschner, G., Eckardt, J., Denton, C., & Arista, N. (2000). Energy angle distribution of low-energy hydrogen ions in thin aluminum and gold foils. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 164, 241–249.
- Frank, L. A., Ackerson, K. L., Lee, J. A., English, M. R., & Pickett, G. L. (1992). The plasma instrumentation for the Galileo Mission. *Space Science Reviews*, 60(1), 283–304. <https://doi.org/10.1007/BF00216858>
- Frank, L., & Paterson, W. (2000). Return to Io by the Galileo spacecraft: Plasma observations. *Journal of Geophysical Research*, 105(A11), 25,363–25,378.
- Funsten, H. O., McComas, D. J., & Barraclough, B. L. (1993). Ultrathin foils used for low-energy neutral atom imaging of the terrestrial magnetosphere. *Optical Engineering*, 32(12), 3090–3095.
- Gilbert, J., Gershman, D., Gloeckler, G., Lundgren, R., Zurbuchen, T., Orlando, T., et al. (2014). Invited article: Characterization of background sources in space-based time-of-flight mass spectrometers. *Review of Scientific Instruments*, 85(9), 91301.
- Gloeckler, G., Geiss, J., Balsiger, H., Bedini, P., Cain, J., Fischer, J., et al. (1992). The solar wind ion composition spectrometer. *Astronomy and Astrophysics Supplement Series*, 92, 267–289.
- Gonin, M., Kallenbach, R., & Bochsler, P. (1994). Charge exchange of hydrogen atoms in carbon foils at 0.4–120 keV. *Review of Scientific Instruments*, 65(3), 648–652.
- Gonin, M., Kallenbach, R., Bochsler, P., & Bürgi, A. (1995). Charge exchange of low energy particles passing through thin carbon foils: Dependence on foil thickness and charge state yields of Mg, Ca, Ti, Cr and Ni. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 101(4), 313–320.
- Hill, T. W., & Michel, F. C. (1976). Heavy ions from the galilean satellites and the centrifugal distortion of the Jovian magnetosphere. *Journal of Geophysical Research*, 81(25), 4561–4565. <https://doi.org/10.1029/JA081i025p04561>
- Kallenbach, R., Gonin, M., Bochsler, P., & Bürgi, A. (1995). Charge exchange of B, C, O, Al, Si, S, F and Cl passing through thin carbon foils at low energies: Formation of negative ions. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 103(2), 111–116.
- Kallenbach, R., Gonin, M., Bürgi, A., & Bochsler, P. (1993). Charge exchange of hydrogen ions in carbon foils. *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 83(1-2), 68–72.
- Kalz, D., Kreysch, G., & Müller-Jahreis, U. (1978). Energy loss straggling of low-energy protons in carbon. *Radiation Effects*, 36(1-2), 119–121.
- Kane, M., Mauk, B., Keath, E., & Krimigis, S. (1992). A convected κ distribution model for hot ions in the Jovian magnetodisc. *Geophysical Research Letters*, 19(14), 1435–1438.
- Kane, M., Mauk, B., Keath, E., & Krimigis, S. (1995). Hot ions in Jupiter's magnetodisc: A model for Voyager 2 low-energy charged particle measurements. *Journal of Geophysical Research*, 100(A10), 19,473–19,486.
- Kane, M., Williams, D., Mauk, B., McEntire, R., & Roelof, E. (1999). Galileo energetic particles detector measurements of hot ions in the neutral sheet region of Jupiter's magnetodisc. *Geophysical Research Letters*, 26(1), 5–8.
- Krimigis, S., Armstrong, T., Axford, W., Bostrom, C., Fan, C., Gloeckler, G., & Lanzerotti, L. (1977). The low energy charged particle (LECP) experiment on the Voyager spacecraft. *Space Science Reviews*, 21(3), 329–354.
- Krimigis, S., Mitchell, D., Hamilton, D., Livi, S., Dandouras, J., Jaskulek, S., et al. (2004). Magnetosphere imaging instrument (MIMI) on the Cassini mission to Saturn/Titan, *The cassini-huygens mission* (pp. 233–329). Switzerland: Springer.
- Krupp, N., Lagg, A., Livi, S., Wilken, B., Woch, J., Roelof, E., & Williams, D. (2001). Global flows of energetic ions in Jupiter's equatorial plane: First-order approximation. *Journal of Geophysical Research*, 106(A11), 26,017–26,032.
- Kurth, W., Hospodarsky, G., Kirchner, D., Mokrzycki, B., Averkamp, T., Robison, W., et al. (2017). The Juno waves investigation. *Space Science Reviews*, 213(1-4), 347–392.
- Livadiotis, G. (2017). *Kappa distributions: Theory and applications in plasmas*. Netherlands-UK: Elsevier.
- Livadiotis, G., & McComas, D. (2011). Invariant kappa distribution in space plasmas out of equilibrium. *The Astrophysical Journal*, 741(2), 88.
- Livadiotis, G., & McComas, D. (2013). Understanding kappa distributions: A toolbox for space science and astrophysics. *Space Science Reviews*, 175(1-4), 183–214.
- Markwardt, C. B. (2009). Non-linear least squares fitting in IDL with MPFIT. arXiv preprint arXiv:0902.2850.
- McComas, D. J., Alexander, N., Allegrini, F., Bagenal, F., Beebe, C., Clark, G., et al. (2017). The Jovian Auroral Distributions Experiment (JADE) on the Juno mission to Jupiter. *Space Science Reviews*, 123, 547–643. <https://doi.org/10.1007/s11214-013-9990-9>
- McComas, D., Allegrini, F., Bagenal, F., Casey, P., Delamere, P., Demkee, D., et al. (2008). The Solar Wind Around Pluto (SWAP) instrument aboard new horizons. *Space Science Reviews*, 140(1-4), 261–313.

- McComas, D., Allegrini, F., Baldonado, J., Blake, B., Brandt, P., Burch, J., et al. (2009). The two wide-angle imaging neutral-atom spectrometers (TWINS) NASA mission-of-opportunity. *Space Science Reviews*, 142(1-4), 157–231.
- McComas, D., Allegrini, F., Pollock, C., Funsten, H., Ritzau, S., & Gloeckler, G. (2004). Ultrathin (~10 nm) carbon foils in space instrumentation. *Review of Scientific Instruments*, 75(11), 4863–4870.
- McNutt, R., Belcher, J., & Bridge, H. (1981). Positive ion observations in the middle magnetosphere of Jupiter. *Journal of Geophysical Research*, 86(A10), 8319–8342.
- Meyer, L. (1971). Plural and multiple scattering of low-energy heavy particles in solids. *Physica Status Solidi*, 44(1), 253–268.
- Moulet, A., Lellouch, E., Moreno, R., Gurwell, M., Black, J. H., & Butler, B. (2013). Exploring Io's atmospheric composition with APEX: First measurement of $^{34}\text{SO}_2$ and tentative detection of KCl. *The Astrophysical Journal*, 776(1), 32.
- Nicolaou, G., McComas, D., Bagenal, F., & Elliott, H. (2014). Properties of plasma ions in the distant Jovian magnetosheath using Solar Wind Around Pluto data on new horizons. *Journal of Geophysical Research: Space Physics*, 119, 3463–3479. <https://doi.org/10.1002/2013JA019665>
- Ogawa, H., Katayama, I., Sugai, I., Haruyama, Y., Aoki, A., Tosaki, M., et al. (1992). Charge state dependent energy loss of light ions. *Nuclear Instruments and Methods in Physics Research Section B: Beam interactions with materials and atoms*, 69(1), 108–112.
- Olbert, S. (1968). Summary of experimental results from MIT detector on IMP-1, *Physics of the magnetosphere* (pp. 641–659). Dordrecht: Springer.
- Ritzau, S., & Baragiola, R. (1998). Electron emission from carbon foils induced by keV ions. *Physical Review B*, 58(5), 2529.
- Schenkel, T., Briere, M., Barnes, A., Hamza, A., Bethge, K., Schmidt-Böcking, H., & Schneider, D. (1997). Charge state dependent energy loss of slow heavy ions in solids. *Physical Review Letters*, 79(11), 2030.
- Scudder, J., & Bridge, H. (1981). A survey of the plasma electron environment of Jupiter: A view from Voyager. *Journal of Geophysical Research*, 86(A10), 8157–8179.
- Seiler, M. C., & Seiler, F. A. (1989). Numerical recipes in C: The art of scientific computing. *Risk Analysis*, 9(3), 415–416.
- Thomas, N., Bagenal, F., Hill, T., & Wilson, J. (2004). The Io neutral clouds and plasma torus. *Jupiter. The Planet, Satellites and Magnetosphere*, 1, 561–591.
- Vasyliunas, V. M. (1968). A survey of low-energy electrons in the evening sector of the magnetosphere with OGO 1 and OGO 3. *Journal of Geophysical Research*, 73(9), 2839–2884.
- Wilhelm, R. A., Gruber, E., Smejkal, V., Facsko, S., & Aumayr, F. (2016). Charge-state-dependent energy loss of slow ions. I. Experimental results on the transmission of highly charged ions. *Physical Review A*, 93(5), 52708.
- Wilhelm, R. A., & Möller, W. (2016). Charge-state-dependent energy loss of slow ions. II. Statistical atom model. *Physical Review A*, 93(5), 52709.
- Williams, D., McEntire, R., Jaskulek, S., & Wilken, B. (1992). The Galileo energetic particles detector. *Space Science Reviews*, 60(1-4), 385–412.
- Wilson, R. J. (2017). JADE standard product data record and archive volume software interface specification. PDS archive volume at <http://pds.nasa.gov/>
- Wilson, R., Bagenal, F., Valek, P. W., McComas, D., Allegrini, F., Ebert, R. W., et al. (2018). Solar wind properties during Juno's approach to Jupiter: Data analysis and resulting plasma properties utilizing a 1D forward model. *Journal of Geophysical Research: Space Physics*, 123, 2772–2786. <https://doi.org/10.1002/2017JA024860>
- Wüest, M. (1998). Time-of-flight ion composition measurement technique for space plasmas, *Measurement Techniques in Space Plasmas: Particles* (pp. 141–155). Washington, DC: American Geophysical Union.
- Ziegler, J. F., Ziegler, M. D., & Biersack, J. P. (2010). SRIM—The stopping and range of ions in matter (2010). *Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms*, 268(11), 1818–1823.