Accepted Manuscript

Comparable short-term morphodynamics of three estuarine-coastal systems in the southwest coastal region of England, UK

Temitope D. Timothy Oyedotun, Helene Burningham

PII: S2352-4855(18)30042-2
DOI: <https://doi.org/10.1016/j.rsma.2019.100749>
Article number: 100749
Reference: RSMA 100749

To appear in: *Regional Studies in Marine Science*

Received date : 13 February 2018
Revised date : 2 July 2019
Accepted date : 3 July 2019

Please cite this article as: T.D.T. Oyedotun and H. Burningham, Comparable short-term morphodynamics of three estuarine-coastal systems in the southwest coastal region of England, UK. *Regional Studies in Marine Science* (2019), <https://doi.org/10.1016/j.rsma.2019.100749>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Comparable short-term morphodynamics of three estuarine-coastal systems in the southwest coastal region of England, UK

Temitope D. Timothy Oyedotun^{a,b*} and Helene Burningham^b

^aDepartment of Geography, Faculty of Earth and Environmental Sciences, University of Guyana, P. O. Box 10 1110, Turkeyen Campus, Greater Georgetown, Guyana, South America.

^bCoastal and Estuarine Research Unit, UCL Department of Geography, Gower Street, London, WC1E 6BT.

*For correspondence: oyedotuntim@yahoo.com; temitope.oyedotun@uog.edu.gy

Abstract

Here, we describe and compare the changes in low tide channel position and the morphology of the three estuaries in southwest England from an analysis of contemporary (2008 – 2016) topographic surveys. The Hayle, Gannel, and Camel estuaries and their adjacent open-coast shorelines were subject of various studies in the past principally in terms of mining impacts on estuarine sediments and sedimentation, but no regional synthesis of the contemporary behaviour has yet been attempted. Light Detection and Ranging (LiDAR) data for the north coast of Cornwall are analysed for planform morphological changes of the systems. At the annual scale, intertidal bars and sandwaves migrate across the foreshore and into the inlet region, where flood-oriented and wave-forced movement is evident. This is evidenced at all the estuarine systems considered here. However, the steep upstream slope of the Hayle is almost devoid of bedforms, while the shallower upstream slope of the Gannel comprises similar scaled bedforms to the whole flood delta, and the megaripples of Camel flood delta are smaller on the backslope. Over the short-term covered by the LiDAR data considered here, there was no significant storm surges, and a relatively consistent wave climate. The occurrence of strong onshore wave conditions are possibly the drivers of contemporary behaviour, and with the inner estuary intertidal currents, for the movements of sediments within the estuarine environment, thereby causing the reorganisation of sediments and the shifting of channel position.

Keywords: contemporary behaviour; morphodynamics; estuaries; LiDAR; bedforms.

1. Introduction

The physical conditions, forms and morphology of the coastal and estuarine environments are far from being stable, especially as they respond in time and space to variations in sediment

movement and transportation (Pierce, 2004; Nicholls, et al., 2007; Shipman, 2010), wave, current or river processes (Lorang et al., 1993; Pierce, 2004; Wong et al., 2014), human activities (Bale et al., 2007; Jaffe et al., 2007; Syvitski et al., 2007; Lee and Ryu, 2008; Wang et al., 2015) and sea level change (Ranasinghe et al., 2013; Padmalal et al., 2014). Sediment-rich estuaries often readily exhibit considerable dynamicity especially in response to, and/or in association with, tidal channel migration (*e.g.* Levoy, et al., 2013; Montreuil et al., 2014). Estuarine systems are generally dynamic (Elias and Hansen, 2013) and are highly controlled by sediment supply to the system, the geological context and the hydrodynamic forces (FitzGerald, 1996; Nordstrom and Jackson, 1992, 2012). The estuarine shorelines, therefore, can progress through significant phases of erosion and/or deposition over time-scales of years to decades (Burningham, 2008) in response to the local hydrodynamic and successive forcing conditions (Savenije, 2006; Montreuil et al., 2014; Leuven et al., 2016).

To understand the complexities of these estuarine-coastal sedimentary interactions, the mechanisms at the intertidal environments have been widely studied by observations (*e.g.* Fan et al., 2006; Green, 2011; Zhu et al., 2014); through analytical solutions (*e.g.* Friedrichs and Aubrey, 1996; Friedrichs, 2011) or numerical modelling (*e.g.* Roberts et al., 2000; Pritchard et al., 2002; Mariotti and Fagherazzi, 2010; Hunt et al., 2015; Hu et al. 2015). These sampled studies are examples of attempts at illustrating the influence of driving forces (*e.g.* hydrodynamic conditions) on estuarine-coastal tidal bathymetry (tidal depth) and morphology. The connection of estuaries to the coasts are found in many places, throughout the world (Duong et al., 2016) and are likely to be in tens of thousands (Carter and Woodroffe, 1994). A wider review of the morphodynamics of estuaries-coastal interaction and response to water level fluctuations, sediment characteristics/movements, geological inheritance, physical processes, *etc.* has been presented and documented in literature (*e.g.* Pye and Blott, 2014; Nordstrom and Jackson, 2012; Duong et al., 2016; Leuven et al., 2016; Ranasinghe, 2016; Robins et al., 2016; Wei et al., 2016; Goudie, 2018). However, to accurately predict the estuarine and coastal morphological changes has proved to be difficult and challenging for coastal scientists especially within the context of climate change (Duong et al., 2017). In the absence of long-term historical and observational data, short-term observation of morphological process is key in the better understanding of the essential processes that drive estuarine-coastal behaviours (*e.g.* Murray, 2003). Little has, however, been focused on the short-term temporal and spatial variability of morphodynamics of estuarine-coastal system interaction and response within a region.

This work describes and compares the changes in low tide channel position and the morphology of the three estuarine systems in southwest England region of Great Britain from an analysis of contemporary (2008 – 2016) topographic (LiDAR) surveys. The aims of this study

are: to examine the comparable temporal and spatial variability in coastal change; and explore the geomorphic sensitivity as the visible manifestations of sediment movements in the estuary-coastal interactions. The contemporary morphodynamic behaviour of the systems considered here focuses on an eight-year (2008 – 2016) investigation of the recent morphological behaviour and short-term morphodynamics. The knowledge of this short-term morphological evolution is of great importance as it is essential to the long-term monitoring of sustainability of intertidal coastal-estuarine sediment movement, coastal and estuarine conservation, coastal defence plan and investigation of changes in the physical features and estuarine-coastal environment of this region.

2. Regional and local setting: Southwest England case study

The Hayle, Gannel and Camel estuaries and their adjacent open-coast shorelines (Fig. 1) have been the subject of various studies in the past primarily in terms of mining impacts on estuarine sediments/sedimentation (*e.g.* Pirrie et al., 1999, 2000; Pascoe, 2005; Brew and Gibberd, 2009; Uncles et al., 2015) and sediments characterisation (*e.g.* Oyedotun et al., 2012, 2013) but no regional synthesis of contemporary behaviour has yet been attempted. These three neighbouring systems provide an excellent opportunity for an investigation of the extent to which the contemporary estuarine behaviour exhibits any regional coherence.

2.1 The Estuaries

2.1.1 The Hayle Estuary is situated within St Ives Bay, which extends between Carbis Bay and Gwithian Towans headlands (Fig. 1c). The estuary is formed within the drowned valleys of the Rivers Hayle (western arm) and Angarrack (eastern arm). Geomorphologically, the Hayle is classified as a bar built (Defra, 2002) and spit enclosed (ABPmer et al., 2008) estuary. The Hayle and St Ives Bay comprise approximately 1.2 km² of largely intertidal sands, mud flats, and saltmarsh. The sedimentary cover overlies a rock floor, which lies at an average of 3.40 m below Ordnance Datum Newlyn (ODN). The tidal regime in St Ives Bay is macro-tidal (mean range at spring tides 5.3 m) and storm surges may add 1 m or more to predicted tidal levels (Pugh, 1987). The published historical (1915 - 2005) rate of sea level rise for Newlyn (southwest England) was 1.77 mm yr⁻¹ (Araújo & Pugh, 2008) while the current rate is *c.* 1.8 mm yr⁻¹ (PSMSL, 2016). The north Cornwall coast is exposed to a predominantly westerly wave climate with a 10% annual exceedance wave height of 2.5 – 3 m and a 1 in 50-year extreme offshore wave height of 20 m, with the possibility of wave heights, regularly exceeding 5 m during the winter months and common swells of 15 seconds or more (Royal Haskoning, 2011).

2.1.2 The Gannel Estuary is also situated in the southwest region of England, between Pentire Point East and Pentire Point West (Fig. 1b). The coastline is similarly macro-tidal (mean spring tide range 6.4 m) (Royal Haskoning, 2011). The Gannel is a ria system comprising sandy intertidal flats within a narrow bedrock valley that merges with the large sandy beach-dune system of Crantock at the seaward extent (Crantock Beach). Around 70% of the estuarine valley is intertidal (Davidson, et al. 1991), and saltmarsh occupies much of the upper intertidal zone. It has been suggested that the estuary (between the Devonian slate/sandstone headlands of Pentire Points East and West) functions as a self-contained sediment cell (Dyer, 2002; Hollick et al., 2006). However, there may be weak and intermittent alongshore transport and some limited exchange of sediments between the bay and open-coast, especially across the low intertidal, and during storm conditions (Royal Haskoning, 2011).

2.1.3 The Camel Estuary, (Fig. 1a) which is adjacent to Padstow Bay is a shallow, predominantly sandy ria formed as a product of post-glacial rise in sea level (Brew and Gibberd, 2009). It is a macro-tidal system with a mean spring tide range of 6.3 m at Padstow, decreasing to 2.8 m near the estuary head c. 12 km up the valley at Wadebridge. Principal sedimentary environments within the estuary include saltmarsh, mud flats, sand flats, subtidal channels, sand dunes and grazing marsh (Brew and Gibberd, 2009). The total intertidal area is around 6 km² with 92% of this being tidal flat (Buck 1993; Brew and Gibberd, 2009). The catchment comprises the River Camel, which drains the Devonian metasediments of the slate formations (Pirrie et al., 2000). The River Camel drains the western margin to the northeast, where granite bedrock is overlain with head deposits (sediments formed through a range of slope processes under periglacial conditions). Catchment geology is mainly Devonian slates and granite with some shales and sandstones.

Fig. 1 Location of the selected study sites on the Cornwall coast of southwest England; Inset: (a) - Camel Estuary, (b) - Gannel Estuary, (c) - Hayle Estuary. (T represents Transect positions extracted for morphological (features) observation, and ★ represents positions for Wave Data used)

3. Material and Methods

The techniques mostly used in mapping intertidal topography focus on the analysis of airborne stereo-photogrammetry, optical satellite imagery, airborne and satellite interferometry, Light Detection and Ranging (LiDAR) data (Mason et al., 2006; Gallay, 2013). The main purpose of using the LiDAR surveys of the study sites is because of its high resolution and accuracy on a larger scale (Lane et al., 2003) and its ability to provide high-density and precise topographic point measurements (Montreuil et al., 2014). Airborne LiDAR data has been widely used in geospatial analyses because of its accuracy and reliability (e.g. Young and Ashford, 2006; Brock

and Purkis, 2009; Young et al., 2009; Lim et al., 2011; Nunes et al., 2011; Young et al., 2011; Earlie et al., 2015). It has also been proven useful in determining and evaluating volumetric changes in diverse coastal landscapes all over the world (e.g. Sallenger et al., 2003; Zhang et al., 2005; Brooks and Spencer, 2010; Schmidt et al., 2011; Jaboyedoff et al., 2012; Allen et al., 2012; Earlie et al., 2015). The LiDAR data is analysed here to assess the recent morphological change within the estuarine intertidal zone.

3.1 Morphological data source and analyses

Airborne LiDAR surveys covering the sites were obtained from the Channel Coast Observatory (CCO, <http://www.channelcoast.org/>). A total of seven LiDAR topographic datasets for each of the estuaries were acquired and used in the analysis of the morphological changes (Table 1). The map date represents the date when the data was captured by the LiDAR instrument which is flown annually to capture the surface reflectance and the changes in the date of flight do not affect the results of the analyses presented here. The LiDAR datasets provide information with which we investigated the recent morphological behaviour of the systems. These LiDAR datasets, obtained at a high spatial resolution of 0.5 m², were analysed using topographic profiles and surface change analysis in GIS. The high resolution and the annual repetitive airborne LiDAR surveys mean that annual changes in sediment movement in the estuarine-coastal environment can be captured and assessed. The public availability of this form of data makes it an ideal type of data that can be used in assessing coastal change at different spatial scales (Earlie et al., 2015) and enables continuous assessment of estuarine-coastal morphodynamics at a short-term scale. This study utilises LiDAR data captured and available for seven (7) different years (from 2008 - 2016) to derive the rates of morphological movements and volumetric changes in the estuarine coastal of three (3) sites in southwest England. In this study, the zones of notable change of vertical erosion, deposition and no change were obtained by undertaking a difference calculation between consecutive surveys, and this was achieved in Quantum GIS (QGIS). A measure of ± 0.25 m was used to differentiate areas of minimal change from those exhibiting notable morphological change. Transects positions (indicated as T in Fig. 1) were also extracted from these spatial surfaces to explore the morphological features associated with the changes observed. This was undertaken in MATLAB. The 2009 Hayle LiDAR data could not be used due to data distribution in integer format, and not as float like others. The technical details and full information on the LiDAR system and datasets are detailed in Levoy et al., (2013) and Kashani et al., (2015), for example.

Table 1 LiDAR datasets used for Hayle, Gannel, and Camel systems

Source	Map Date
<u>Hayle</u> Channel Coast Observatory	09/09/2008
	10/09/2009
	10/04/2010
	19/03/2011
	06/04/2012
	17/04/2014
	11/03/2016
<u>Gannel</u> Channel Coast Observatory	27/02/2008
	16/04/2009
	09/02/2010
	09/02/2011
	05/04/2012
	05/03/2014
	11/03/2016
<u>Camel</u> Channel Coast Observatory	27/02/2008
	16/04/2009
	09/02/2010
	10/03/2011
	06/04/2012
	05/03/2014
	11/03/2016

To evaluate the importance of physical environmental forcings as agents of change within the coastal system, wave climate conditions, using the 20-year (1999 – 2009) hindcast hourly wave parameters supplied by Associated British Ports (ABP) Marine Environmental Research Ltd (ABPmer, henceforth), were analysed in MATLAB. Specifically, the wave parameters at West (Long. -5.67, Lat. 50.65) and Central West (Long. -5.33, Lat. 50.55), (indicated as 1 and 2 in the inset map of Fig. 1), which are within the study areas, were considered for contemporary wave condition analysis. Hindcast wave data were provided by ABP Marine Environmental Research Ltd, from its SEASTATES hindcast service (www.seastates.net). Key variables explored here are the frequency of wave direction and significant wave height, the frequency distribution of wave direction and wave period, time-series of significant wave height and wave approach. For the tidal conditions and sea levels, information from the Permanent Service for Mean Sea Level (<http://www.psmsl.org/>) for Newlyn, the closest gauging station to the study sites were considered. It should be noted that the wave climate around the study area (the Cornish coastline) is the most energetic of the UK coastal waters (Scott et al., 2011) as this region is exposed to high energy swells from the Atlantic Ocean and wind waves from the region's prevailing westerly winds (Earlie et al., 2015).

4. Results

4.1 Recent morphological change

4.1.1 Hayle Estuary

The change analysis of repeat LiDAR surveys for St Ives Bay and the Hayle (Fig. 2) show small lateral shifts in channel position, but the analysis shows that movement of low amplitude bars and high tide berms dominate the short-term beach dynamics. Much of the topographic change shown in LiDAR change analysis is balanced across the system as expected with onshore/offshore bar/berm movement. Between 2008 and 2011, the vast proportion of beaches (36.4% of the proportion) to the north of the Hayle inlet showed evidence of accretion, and in fact, the primary signature of change is accretional (positive), as the average net change was 0.013m. Erosion was almost entirely associated with the ebb tidal delta (Porth Kidney Sands) (~1.7%), except for some erosion in Carbis Bay (~1%) and some along the high-water shorelines of the inlet. It is worth noting that there is some asymmetry to the change in Carbis Bay where erosion dominates to the east and accretion dominates to the west. This could reflect small-scale rotation in the beach deposit in this small bay. The patterns of positive and negative change across the ebb tidal delta are best explained by the migration of swash bars across the broad intertidal flat: the location of the bars in the earlier time frame is shown as a significant erosional patch, and where they are present in the latter time frame is shown as a large accretional patch. Within the estuary, change is limited to two clear patches of accretion, one in the central basin, and one within the western arm. In both cases, between 0.5 and 1m of accretion has taken place across large banks within and alongside the channel. Although some of this change could be attributed to channel migration (there are certainly some linear erosional features alongside the channels, it seems likely that most of this sediment has been brought into the estuary through the inlet, and these features are comparable to flood-tidal deltas. Table 2 presents the frequency volume of classes of the magnitude of change in Fig. 2.

In Carbis Bay, however, the rotational change evidenced between 2008 and 2011 is reversed, where erosion now dominates the west, and accretion to the east. This seems to suggest that Carbis Bay switches between westerly and easterly skewed orientation. Due to problems with the 2009 data, it is not possible to establish whether this is an annual rotation, but consideration of the total change between 2008 and 2016 shows that the changes of 2011-2012, 2012-2014 and 2014-2016 have dominated over the slightly longer period. Conversely, it is the accretional signature in beach change from 2008-2011 that is maintained as a signature in the slightly longer term (2008-2016) (Fig. 2).

The change between 2011 and 2012 (with the average net change of 0.043 m³) is to some extent comparable to that shown for 2008-2011 (with the average net change of 0.013 m³). Most

change is focused on the ebb delta region, but in this time frame, very little change ($\sim\pm 0.25\text{m}$) is evident for beaches to the east and west of the inlet. There are patches of both small-scale accretion and erosion within the estuary, some of which looks to be the result of small shifts in channel position.

4.1.2 Gannel Estuary

The Crantock-Gannel system shows perhaps slightly smaller scales of change to those evidenced in St Ives Bay (Fig. 3). Here though, changes within the estuary part of the system are of comparable magnitude. Between 2008 and 2009, the most significant patterns of erosional and accretional occur on the beach, aligned in a cross-shore pattern. Along the southwest margin, a large area of erosion dominates, but further north on the beach, accretion dominates. The patterns suggest that sediment movement on this beach is in the form of large-scale migratory bedforms, but that these are neither shore-parallel nor shore-normal in structure or movement. Closer to the inlet, it is clear small lateral shifts in channel position have resulted in a succession of linear erosional and accretional signatures. This continues into the estuary, but here there is also evidence of larger, more diffuse areas of erosion and accretion that would be associated with bedform movement over the intertidal flats.

Fig. 2 Recent morphological change in Hayle Estuary

Between 2009 and 2010, somewhat interesting patterns of erosion and accretion continue to take place across the Crantock foreshore. Again, they are more generally concentrated to the southwest, but there is also further change alongside the channel and inlet. In this case, quite significant erosion (-2m to -1 m) is evident along most of the southern margin of the inlet, but perhaps in balance to this, a large deposit is shown around the seaward extent of the channel. Between 2010 and 2012, accretion (0.5m to 1m) is evident across most of the mid-foreshore, extending shore-parallel across most of the bay. To the seaward of this, some erosion (between -2 to -5m) is shown between 2012 and 2014, and this erosion is along the upper foreshore, closer to the inlet. Within the estuary, the rather erosional expression shown in the 2009-2010 change map is replaced by a distinctly accretional signature in 2010-2012 and erosional signature in 2012-2014 and 2014-2016 respectively.

When considering the sequence of events between 2008 and 2016, perhaps what is most striking about the series of changes shown here is the behaviour of depositional features on Crantock beach and the branching of stream channel between 2014 and 2016. Apart from this, quite a significant volume of material accumulated on the lower foreshore on the southwest side of the Bay between 2008 and 2009 that subsequently dispersed cross-shore (landward), and then moved alongshore (north-eastward). This suggests that sediment is delivered to the west part of the beach and is then redistributed north and east-ward over the following 3 or 4 years (2012 – 2016) across the rest of the foreshore. Table 2 presents the frequency volume of classes of the magnitude of change in Fig. 3.

Within the estuary, however, there appear to be different cycles of erosion and deposition that might relate specifically to channel meandering, but also seem to suggest quite significant delivery of sediment into the estuary from the inlet region. Certainly, over the 8-year period, the margins of the inlet show significant erosion and reshaping, whilst the estuary shows a complex mosaic of accretion and erosion, possibly associated with the movement of megaripples through the system.

Fig. 3 Recent morphological change in Gannel Estuary

4.1.2 Camel Estuary

Patterns of change in Padstow Bay and the Camel estuary are complicated, with multiple patches of erosion and accretion observed throughout the system, and throughout the time periods considered here (Fig. 4). Focusing initially on the estuary, between 2008 and 2009, the change was primarily characterised by accretion – large areas of intertidal flat experienced 0.25-2m vertical accretion. To some extent, quite a lot of this change was associated with channel margins and evidenced for migration is shown in the presence of spatially-matched, narrow patches of erosion and accretion either side of the main channels. From 2009 to 2011, the general patterns of change are spatially comparable, but the signature is reversed, and erosion dominated. Channel migration clearly continues, most notably along the stretch to the east of Padstow where the west bank is eroded and the east bank exhibits accretion. Changes between 2011 and 2012 appear very similar to those between 2008 and 2009, where again, erosion is largely aligned with channel margins, and large surfaces of the intertidal flat show an accretionary signature. Table 4 presents the frequency volume of classes of the magnitude of change in Fig. 4.

Channel migration driving erosion and accretion is not observed in the outer bays of Harbour Cove and Daymer Bay. These intertidal flats form the ebb-delta, and here deposition and erosion seem to follow the formation of large sedimentary deposits such as sandwaves and their migration across the flats. This is clearer in Daymer Bay, where the deposits are shore-parallel, but in Harbour Cove, features are more complex and variable in structure, though the dynamic zone is clearly the lower, rather than upper foreshore.

The net consequence of these changes (2008 to 2016) reveals quite substantial changes have occurred over the vast majority of the intertidal environment of the Camel, especially between 2012 and 2016. In comparison, the open coast beach in Padstow Bay and within the estuary dynamics exhibit very little change over a similar period (2008 to 2011). Although the lack of LiDAR data for the open coast site precludes a direct comparison, the small patches of accretion (0.25 m to 2m) evident are insignificant in comparison to the dynamics shown in the estuary, inlet, and ebb-delta regions.

Fig. 4 Recent morphological change in Camel Estuary

Table 2 Frequency analysis for classes of the magnitude of sediment movement (m³) in the Hayle system

Volume of sediment movement (change) (m ³)							
2008 – 2011		2011 – 2012		2012-2014		2014 - 2016	
Average net change (m ³)		0.013		-0.045		0.051	
Classes of Magnitude (m)		Absolute (m ³)	% of proportion	Absolute (m ³)	% of proportion	Absolute (m ³)	% of proportion
-5 - -2	3384	3705	0.11	4805	0.11	3106	0.07
-2 - -1	57361	39114	1.11	38234	0.90	37146	0.84
-0.5 - -0.25	283517	35349	10.04	292418	6.88	342298	7.74
-0.25 – 0.25	1919625	275723	78.4	2646324	62.23	2826643	63.91
0.25 – 0.5	898832	262276	7.5	883238	20.77	902823	20.41
0.5 – 1	229112	226041	6.4	283401	6.66	212215	4.80
1 – 2	106276	66396	1.9	103627	2.44	98578	2.23
2 – 5	24	29	0.001	40	0.001	68	0.0001
						210	0.01

Table 3 Frequency analysis for classes of the magnitude of sediment movement (m^3) in the Gannel system

Volume of sediment movement (change) (m ³)									
	2008 – 2009	2009 – 2010	2010 – 2012	2012 – 2014	2014-2016	2008 - 2012			
Average net change (m ³)		0.032	0.077	0.09	0.17	0.19			
Classes of Magnitude (m)	Absolute (m ³)	% of proportion	Absolute (m ³)	% of proportion	Absolute (m ³)	% of proportion			
-5 - -2	86	0.01	219	0.02	425	0.04			
-2 - -1	3943	0.54	3046	0.3	1238	0.13			
-0.5 - -0.25	49960	6.84	82112	8.15	73259	7.65			
-0.25 – 0.25	649718	89.02	82249	81	725140	75.71			
0.25 – 0.5		6.68	12777	12.7	123606	12.9			
0.5 – 1	48786	1.49	100180	13.5	12654	13.48			
1 – 2	10853	0.29	33304	4.5	30146	3.13			
2 – 5	2152	0.001	4484	0.61	4098	0.41			
	7		16	0.002	346	0.03			
			2	0.001	358	0.04			
				252	681091	17.05			
				1897	167188	6.16			
				1897	60397	0.97			
				252	9478	0.04			

*Instead of comparing directly here - the number of cells in classes of magnitude of change is presented as a % (proportion) rather than an absolute.

Table 4 Frequency analysis for classes of the magnitude of sediment movement (m³) in the Camel System

Volume of sediment movement (change) (m ³)												
2008 - 2009		2008 - 2011		2009 - 2011		2011 - 2012		2012-2016		2008 - 2016		
Average net change (m ³)		0.017		-0.094		0.083		-0.098		0.09		
Classes of Magnitude (m)	Absolute (m ³)	% of proportion	Absolute (m ³)	% of proportion	Absolute (m ³)	% of proportion	Absolute (m ³)	% of proportion	Absolute (m ³)	% of proportion	Absolute (m ³)	
-5 - -2	9036	0.14	112448	0.41	24593	0.4	15173	0.2	29989	0.50	25174	0.41
-2 - -1	24215	0.38	112448	1.66	106625	1.7	28175	0.5	15725	0.26	26147	0.43
-0.5 - -0.25	77626	1.22	574998	8.5	758198	12.3	269715	4.4	243261	4.09	248627	4.04
-0.25 - 0.25	5450930	85.5	5464654	83.7	5509911	87.1	5349288	86.4	4958796	83.32	5237067	85.16
0.25 - 0.5	541003	8.5	507484	7.3	276 93	4.5	455679	7.4	369591	6.21	342357	5.57
0.5 - 1	222355	3.5	303270	4.5	13 640	2.1	190315	3.1	235234	3.95	190014	3.09
1 - 2	47727	0.75	141760	2.1	31260	0.5	80505	1.3	98631	1.66	79689	1.30
2 - 5	276	0.01	356	0.01	311	0.01	332	0.001	459	0.01	434	0.01

*Instead of comparing directly here - the number of cells in classes of magnitude of change is represented as a % (proportion) rather than an absolute.

4.2 Ebb-inlet-flood delta morphodynamics

4.2.1 Hayle Estuary

Cross-sectional profiles are investigated using the LiDAR data (Moore et al. 2009) to examine the behaviour of specific cross-shore features. Cross-sections extracted from LiDAR data for St Ives Bay (2008 – 2016) are shown in Fig. 5. The western transects (T1, T2) show varied shifts – some accretion at T1 and T2, and more significant erosion at T2. The eastern beach transects (T4) show very small changes. Beach level fluctuates by up to 75 cm (as shown in the vertical accretion at T2 and erosion at T4), but again the story is not consistent across the bay. Transect T3 crosses the main ebb channel in the inlet. Here, significant vertical change has occurred over the 2-3-year period, but only on the eastern margin: the west bank has changed little in comparison. Referring to the spatial mapping of change, the deposits here (of >2 m) are associated with the migration of sandwaves/spits into the inlet from the upper foreshore of the beach.

Fig. 5 LiDAR cross-sections for transects 1 – 4 (T1 – T4) in Hayle Estuary (Transects positions are indicated in Fig. 1)

4.2.2 Gannel Estuary

At Crantock, the shoreface cross section (T1) reveal only small changes in the position of Mean High Water (MHW) and Mean Low Water (MLW), and small-scale shifts in the beach level during the eight-year (2008 – 2016, Fig. 6), except at T1 where there is a notable change between 2014 and 2016. The estuarine transects (T2 – T4) which cross the main ebb channel in the inlet and inner estuary reveal year to year localised erosion and deposition, which appears to be linked to sandbank movement and channel migration/shifting in the Gannel estuary. The inlet transects (T2) shows variable change over the 8-year period, whilst T4 (extending over the flood-delta just landward of the inlet) illustrates the role of mobile and migrating sandwaves (wavelength 8–10cm, height 10–30cm) in the re-organisation of sediment over the larger intertidal forms. These bedforms progressively move over the broad intertidal flood-delta platform into the estuary. As expected, the flood-delta is forced primarily by tidal currents, but the changes also suggest that this supply of sediment into the estuary might drive shifts in channel position.

Fig. 6 Transects positions of the LiDAR cross-sections in Gannel Estuary (Transects positions are indicated in Fig. 1). Please note different elevation (y-axis) and distance scales (x-axis)

4.2.3 Camel Estuary

Transects within the Camel estuary (Fig. 7) reveal smaller shifts in surface levels over the intertidal flats, with year to year localised erosion and deposition. Within the ebb-delta and inlet region, smaller scales of change are more observed (T3). Broad, but low features move across the lower foreshore in both T2 and T3: these do not show progressive change, and the envelope of variability is consistent across the foreshore. Within the inlet, however, there is distinct evidence of accretion on both margins of the inlet. The east bank displays progressive accumulation across the mid and upper foreshore. In places, the vertical accretion is *c.* 1m, and this has led to a narrowing of the mid-tide inlet by around 25m. Given the presence of intertidal bars across the foreshore to the north (T2), it is likely that this accretion has benefitted from sediment supply via these mobile foreshore bars.

Into the estuary, the wide transect at T4 shows the lateral extent of the channel network through the estuary. This transect seems to suggest that changes in elevation of the intertidal flat surface are less distinct than changes associated with channel movement. A small static channel exists close to the west bank, but there are several substantially larger channels within the mid-estuary where there is evidence of progressive migration (shown by the gradual retreat or advance of the channel margin) and channel switching (where channels seem to appear and disappear). Channel switching is particularly clear around 300 m into the transect, where a substantial channel was present in 2009 but completely gone by 2011. It is not clear whether shoals either side prograded and therefore enclosed and infilled this channel or whether sediment was delivered from up/down stream. But it is clear that channels in the mid-Camel estuary are significantly more active than in the Havel or Gannel. As the valley begins to narrow, the intertidal structure becomes less variable. At T3, the intertidal surface shows both minimal erosion and accretion over the 8 years, but channel position has remained relatively stable, and this is furthermore the case at T4.

Fig. 7 Transects positions of the LiDAR cross-sections in Camel Estuary (Transects positions are indicated in Fig. 1). Please note different elevation (y-axis) and distance scales (x-axis)

5. Discussion

5.1 Comparable contemporary morphological behaviour

At the annual scale, intertidal bars and sandwaves migrate across the foreshore and into the inlet region, where flood-oriented and wave-forced movement is evident. This is evidenced, to lesser and greater extent, at all the estuaries considered here. Flood dominance in the outer estuary encourages the net landward movement of sediment across inlet-associated beaches and supplies sediment to the macro-tidal estuaries. Findings here support the concept that an estuary has a tendency toward flood dominance until sediment-infilling becomes more advanced, thereby forcing a change in tidal hydrodynamics that leads to a switch to ebb dominance (Moore et al., 2009; Friedrichs, 2011; Hunt et al., 2015; Duong et al., 2016). The contemporary behaviour suggests shifts in the flood-ebb balance are occurring on an annual scale, particularly in the Camel estuary where large areas of accretion are followed a year or two later by large areas of erosion. Channel migration is an important driver of the erosion and accretion experienced in these estuaries, but these changes are focused on the channel-margins, a distinctly

different geomorphic signature to the large-scale accretion and lowering evidenced over the intertidal flats.

The flood delta zones of the three estuaries comprise similar bedforms – megaripples of 0.2-0.5m height and 10-20m wavelength that climb a shallow platform that rises gradually by 0.4-0.5° (Fig. 8). The megaripples reduce in size with distance across the platform, and beyond the size of the flood delta, it is only on the landward ramp that any significant difference between the three estuaries can be found. The steep upstream slope of the Hayle is almost devoid of bedforms, while the shallower upstream slope of the Gannel comprises similar scaled bedforms to the whole flood delta, and megaripples are smaller on the back slope of the Camel flood delta. The megaripples are quite varied and suggest superimposition but show no significant evidence of asymmetry – it is only the skewed and slightly concave profile of the flood delta shoal and the arcuate planform nature of the bedforms that imply any flood-dominance across these features.

Based on the results here, where both the ebb and flood deltas show evidence of flood-dominance and movement of sediment into the inlets and main estuary body via bedform migration, it would follow that the estuaries are infilling on a relatively short timescale. But channel migration within each system appears to be capable of eroding and releasing large volumes of sediment which are then perhaps transported via the ebb tide out of the estuaries to be deposited nearshore to the ebb deltas. It is, therefore, possible that the net effect over several years is balanced as sediment arriving at the ebb delta might simply be sourced from the release of sediment within the estuary, through channel dynamics.

The focus so far on the inlet, delta and channel dynamics, at the expense of discussion of the open coast are unsurprising given the difference in scales of change exhibited. Crantock beach is the perhaps the most dynamic ‘open coast’ site, but this is likely to be due to the lack of distinct ebb delta here. The foreshore almost operates in the way expected for ebb deltas, where swash bars form in the lower foreshore and migrate across the upper foreshore to the inlet margin (Hayes, 1975; Panagoulas et al., 2013). The open coast beach in Padstow Bay was the most stable environment within the entire system, although LiDAR coverage only allowed consideration of a single time interval. As with the other sites though, beach dynamics increased with proximity to the inlet, a behaviour that has been documented by others (FitzGerald, 1988; Hicks and Hume, 1996; Nordstrom and Jackson, 2012; Nunes et al., 2011; Pierce, 2004; Garnier et al., 2006; Young and Ashford, 2006; Short et al., 2013). The broader extent of beaches in St Ives Bay is possibly the best barometer of open coast vs. inlet and estuary dynamics, and here there was varying change depending on time frame. Widespread beach accretion occurred between 2008 and 2011, but this was followed by a negligible change between 2011 and 2012, comparable to the behaviour of beaches in outer Padstow Bay.

Beach rotation displayed in Carbis Bay points to a change in forcing between 2008-2011 and 2011-2012. One could speculate that morphological changes between 2011 and 2012, which comprised erosion to west Carbis Bay and accretion in the east, might be the result of an increased forcing from the northeast, which would be out of the norm given the westerly-dominated climate in this region.

Over the short-term covered by the LiDAR data considered here, there were no significant storm surges, and a relatively consistent wave climate. Although sea level change throughout the 2000s was variable (PSMSL, 2016), a sustained rise was achieved during the dominant accretion episodes (2008-2011 in St Ives Bay, and 2008-2009 on the Camel estuary) so it seems unlikely that this was responsible for the changes observed. The relative stability reported in the beaches may be because of the absence of storminess, but also possibly the limitation of the data to fully illustrate short-term changes (*i.e.* consistent annual intervals would certainly help to improve clarity).

Fig. 8 Cross-sections of the flood tidal deltas in the Hayle, Gannel, and Camel estuaries. Scales of topographic profiles are equivalent (distance and elevation are in metres).

5.2 Contemporary morphodynamics and forcing factors

Spatio-temporal shifts in coastal system morphology reflect processes operating in the coastal-estuarine environment at a different range of magnitudes and frequencies. It has been shown that the coastal and estuarine environment can recover relatively rapidly following a single storm event (Montreuil and Bullard, 2012), but a continuous sequence of storm events is able to cause major and sustained alterations because of insufficient recovery time (Fan et al., 2006; Nunes et al., 2011; Montreuil and Bullard, 2012).

The recent LIDAR data (2008-2016) examined allowed the exploration of temporal variability of short-term morphodynamics. The intertidal environment showed expressions of erosion and accretion that in places were commensurate with the scales of change observed over the contemporary time scale. The beaches exhibited only small changes in the position of MHW and MLW, despite in some cases there being evidence of significant intertidal deposits that moved across the foreshore. In the estuaries, year to year localised erosion and deposition took place, which appears to be linked to sandbank movement and channel migration/shifting. Topographic cross-sections covering the flood-tidal delta, just landward of the inlet, illustrate the role of mobile and migrating sandwaves in the re-organisation of sediment over the larger intertidal forms. These progressively move into the inner estuary, and so seem to be forced by tidal currents, and this force shifts in channel position.

On a regional scale, the character of sea level is much more variable (Teasdale et al., 2011) because of a more local factor such as tectonic and glacio-isostatic movements (Teasdale, et al. 2011). The trend of the consistent rise in sea level in the Celtic Sea continues in the 21st century, with some fluctuations, annual mean sea level rise subsided between 2002 and 2005 before increasing again in 2010 (PSMSL, 2016). The annual rate for the entire southwest region remains *c.* 1.77 mm yr⁻¹ (PSMSL, 2016). With the recent observation and the expected future increase in sea level, wave climate, tidal processes and the morphodynamic evolution of the north Cornwall coastlines are expected to be affected.

The largest proportion of waves, in this region, typically occurs from the westerly position with 56.2% (Table 5). Waves from the north are far more infrequent, with 17.6% and given the reduced fetch; waves from the southeast are very rare (Fig. 9). The coastal systems considered in this study all occupy a north-westerly aspect. Basic metrics of the wave climate split based on direction (westerly vs. northerly). The northerly waves are smaller (in terms of height and period) than westerly waves but have a slightly wider spread indicating that westerlies are more likely swell-dominated, and northerlies comprise an increased wind-wave component.

One of the most important coastal factors which influences change on coastlines is the state of the tide. The high waves are potentially most likely to cause a notable hazard on the coastline

at high tide. The spring tidal range for this region is around 5-6 m (UKHO, 2003) and the coast can be described as macro-tidal. Tidal currents are described as generally weak ($< \sim 0.75 \text{ ms}^{-1}$) except in local areas around the headlands (Dyer, 2002). The lowest monthly water levels are around 0 to 0.5m, with minimum surges of -0.1 to -0.5m. The maximum monthly water levels range from 5.5 to 6.5 m (highest astronomical tide level is 6.13 m above chart datum) and monthly mean maximum surges are up to 1 m (from the Permanent Service for Mean Sea Level, <http://www.psmsl.org/>). The envelope of variability over a 12-year period is relatively consistent, where year to year fluctuations occur on similar scales. The highest monthly mean water level recorded was 6.42 m in October 2004 and the lowest was -0.01 in February 1996, but these levels occur relatively frequently, so not considerably out of the normal range (PSMSL, 2016).

Fig. 9 Wave roses for hindcast data, showing A) frequency distribution of wave direction and significant wave height and B) frequency distribution of wave direction and wave period (West Point and Central West Point locations are indicated as 1 and 2 in Fig. 1).

Fig. 10 Time series of significant wave height at the West and Central West points.

Table 5 Wave climate summary for westerly (225-315°N) and northerly (315-45°N) waves.

	Percentile	Westerly waves		Northerly waves	
		West point	Central West point	West point	Central West point
Proportion (%)		56.2	60.9	17.6	19.2
Vector mean direction	202.58				
Significant wave height (m)	50th	1.67	1.56	0.96	0.89
	90th	3.99	3.69	2.27	2.11
	99th	6.52	6.04	4.39	4.12
Wave period (s)	50th	5.62	5.53	3.90	3.77
	90th	8.12	7.84	5.58	5.43
	99th	10.05	9.77	7.41	7.22
Wave spread (°)	50th	31.00	29.73	34.91	33.73
	90th	41.18	39.57	52.82	50.90
	99th	63.17	60.95	72.22	70.70

Continued sea level rise in the region (PSMSL, 2016), the occurrence of strong onshore wave conditions (Figs. 9, 10) are possibly the drivers of flood-ebb delta dynamics and the small-scale coastline retreat observed in the contemporary behaviour. The forcing factors for the period 2008-2016 showed little evidence of storm events and surges that were out of the ordinary. Therefore, the relative constant tidal conditions, wave parameters, and sea level during the eight-year period are thought to cause small changes in the position of MHW and ebb channel. The

inner estuary intertidal currents, however, are responsible for the movements of sediments within the estuarine environment, thereby causing the re-organisation of sediments and the shifting of channel position. The changes at the annual scale are mainly on bedform (sandwave) movement within, and landward of the inlet, in a flood-oriented direction in the estuaries which facilitates shifts in sandbanks and forces changes in channel position.

The geological framework, sedimentary characteristics and antecedent geomorphology are known to determine the sensitivity of estuaries to changes in forcing factors (Roberts, 2000; Blott et al., 2006; Green, 2011; Lim et al., 2011; Hunt et al., 2015). Though the inner channels of the estuaries are, to some extent, protected, particularly by narrow rock-controlled inlets, there are, however, alternating zones of stable and unstable channels present in the short-term period considered. The inlets are observed to have positional stability as a result of the geological framework, but evidence of some morphological instability is noted in terms of channel throat widening and/or deepening to varying degrees as the tidal regime and processes probably cause fluctuations in sediment movement and supply. What was clear in the contemporary analysis here, was the tendency for morphological change around the inlet margins because of upper foreshore spit and bar development and extension into the inlets. Furthermore, the ebb tidal deltas were shown to be dynamic intertidal platforms where year to year changes in the distribution of large-scale bedforms (such as swash bars) far exceeded the changes observed along the high or low water shorelines. The net product of significant foreshore activity was far less than the gross magnitudes of change.

6. Conclusions

Changes over the contemporary time scale are focused on bedform (sandwave, megaripples, small spits) movement into, within, and landward of the inlet. Shifts in these bedforms are driven by waves in the outer estuary/ebb delta region and inlet upper foreshore, and by tides in the channels and flood delta region. Within the estuaries, tidal force drives the channel and sandwave (sediment) movement while adjacent to the inlet, the waves drive shoreline movement. There is difficulty in finding the explicit link between the morphodynamic behaviour observed in all the systems and the processes in coastal climate or the relative sea level. It is most probable that the systems are responding to more than one aspect of forcing at a time. The result presented here has investigated, for the first time, the contemporary morphological behaviour of the macrotidal estuarine systems in southwest England. The use of LiDAR datasets in this study has also highlighted the importance of LiDAR datasets as an important source of information for coastal morphological and sediments dynamics.

Acknowledgements

We will like to thank and acknowledge the following organisations for their contributions to this research work: Channel Coast Observatory (<http://www.channelcoast.org/>) for LiDAR data used; ABP Marine Environmental Research (ABPmer) Ltd, for hindcast wave data (Hindcast wave data were provided by ABP Marine Environmental Research Ltd, through its SEASTATES hindcast service (www.seastates.net); and Permanent Service for Mean Sea Level (<http://www.psmsl.org/>) for Mean Sea Level for Newlyn. The authors appreciate the anonymous reviewers for their valuable comments and suggestions on this paper.

References

- ABP MER, University of Plymouth, University College London, Discovery Software, HR Wallingford, & Delft Hydraulics 2008, *Development and Demonstration of Systems-Based Estuary Simulators. R & D Technical Report FD 2117/TR*, Joint DEFRA/EA Flood and Coastal Erosion Risk Management R & D Programme, FD2117/TR.
- Allen, T. R., Oertel, G. F., Gares, P. A. 2012. Mapping coastal morphodynamics with geospatial techniques, Cape Henry, Virginia, USA. *Geomorphology* 137, 138–149. doi:10.1016/j.geomorph.2010.10.040
- Araújo, I.B., and Pugh, D.T., 2008. Sea level rise at Newlyn 1915-2005: Analysis of trends for future flooding risks, *Journal of Coastal Research*. 24 (4C), 203-212
- Bale, A.J., Uncles, R.J., Villena Lincoln, A., Widdows, J., 2007. An assessment of the potential impact of dredging activity on the Tamar estuary over the last century: bathymetric and hydrodynamic changes. *Hydrobiologia* 588, 83–95.
- Bitner-Gregersen, E. M., Evans, K. C., and Johnson, M. C. 2014. Some uncertainties associated with wind and wave description and their importance for engineering applications. *Ocean Engineering* 8, 11-25
<http://dx.doi.org/10.1016/j.oceaneng.2014.05.002>
- Blott, S.J., Pye, K., van der Wal, D., & Neal, A. 2006. Long-term morphological exchange and its causes in the Mersey Estuary, NW England. *Geomorphology* 81, 185-206
- Brew, D.S. & Gibberd, B.B. 2009. Geomorphological change and its impact on habitats in the Camel Estuary, Cornwall, UK. *Geoscience in South-West England* 12, 95-100
- Brock, J. C., Purkiss, S. J. 2009. The emerging role of LiDAR remote sensing in coastal research and resource management. *J Coast Res Spec Issue* 53,1–5
- Buck, A.L. 1993. *An Inventory of UK Estuaries. Volume 2 South-west Britain*. Peterborough, Joint Nature Conservation Committee.
- Burningham, H. 2008. Contrasting geomorphic response to structural control: The Loughros

- Estuaries, Northwest Ireland. *Geomorphology* 97, 300-320
- Cabanes, C.A., Cazenave, A., & Le Provost 2001. Sea level rise during the past 40 years determined from satellite and in-situ observations. *Science* 294, 840-842
- Carter, R.W.G., Woodroffe, C.D., 1994. *Coastal Evolution: Late Quaternary shoreline Morphodynamics*. Cambridge University Press (517 pp.).
- Church, J.A. & White, A. 2006. A 20th Century acceleration in global sea level rise. *Geophysical Research Letters* 33, L01602
- Davidson, N. C., Laffoley, D. D., Way, L. S., Key, R., Drake, C. M., Pienkowski, M. W., Mitchell, R., & Duff, K. L. 1991, *Nature conservation and estuaries in Great Britain*, Nature Conservancy Council, Peterborough, UK.
- Defra 2002, *Futurecoast*, Defra.
- Defra 2006, *Flood and Coastal Defence Appraisal Guidance FCDAG3 Economic Appraisal*, Supplementary Note to Operating Authorities - Climate Change Impacts, October 2006.
- Deng, J., Woodroffe, C., Rogers, K., Harff, J. 2017. Morphogenetic modelling of coastal and estuarine evolution. *Earth-Science Reviews* 171, 254–271
- Duong, T. M., Ranasinghe, R., Walstra, D., Roelvink, D. 2016. Assessing climate change impacts on the stability of small tidal inlet systems: Why and how? *Earth-Science Reviews* 154, 369–380
- Dyer, K. R. 2002, "FutureCoast Estuary Assessment," In *Futurecoast CD-ROM prepared for Defra.*, Halcrow, ed.
- Earlie, S. C., Masserlink, G., Russell, P. E., Shail, K. R. 2015. Application of airborne LiDAR to investigate rates of recession in rocky coast environments. *J Coast Conserv.* 19, 831–845 <http://dx.doi.org/10.1007/s11852-014-0340-1>
- Elias, P.L., Hansen, J.E., 2013. Understanding processes controlling sediment transports at the mouth of highly energetic inlet system (San Francisco Bay, CA). *Mar. Geol.* 345, 207–220. <http://dx.doi.org/10.1016/j.margeo.2012.07.003>.
- Fan, D., Guo, Y., Wang, P. and Shi, J. Z. 2006. Cross-shore variations in morphodynamic processes of an open-coast mudflat in the Changjiang Delta, China: With an emphasis on storm impacts, *Cont. Shelf Res.* 26(4): 517–538.
- FitzGerald, D. M. 1968, "Shoreline erosional-depositional processes associated with tidal inlets," In *Hydrodynamics and Sediment Dynamics of Tidal Inlets.*, D. G. Aubrey & L. Weishar, eds., New York: Springer-Verlag, Inc., pp. 186-225.
- Friedrichs, C. T. 2011. Tidal flat morphodynamics: A synthesis, in *Treatise on Estuarine and Coastal Science*, edited by E. Wolanski and D. McLusky, chap. 3.06, pp. 137–170,

- Academic Press, Waltham, Mass.
- Friedrichs, C. T., and D. G. Aubrey 1996. Uniform bottom shear stress and equilibrium hypsometry of intertidal flats, In: Pattiaratchi, C (ed) *Coastal and Estuarine Studies* 50, 405–429, AGU, Washington, D. C.
- Goudie, A. 2018. Rias: Global distribution and causes. *Earth-Science Reviews* 177, 425–435
- Gallay, M. 2013. Direct Acquisition of Data: Airborne laser scanning. In Clarke, L. E. and Nield, J. (Ed) *Geomorphological Techniques (Online Edition)*. British Society for Geomorphology, London, UK. ISSN 2047 – 0371. Chapter 2, Section 1.4.
- Garnier, R., Calvete, D., Falqués, A., Caballeria, M., 2006. Generation and nonlinear evolution of shore-oblique/transverse sand bars. *J. Fluid Mech* 567, 321–360. <http://dx.doi.org/10.1017/S0022112006002126>.
- Green, M. O. 2011., Very small waves and associated sediment resuspension on an estuarine intertidal flat, *Estuarine Coastal Shelf Sci.* 93(4), 549–559.
- Hayes, M.O., 1975. Morphology of sand accumulation in estuaries: an introduction to the symposium. In: Cronin, L.E. (Ed.), *Estuarine Research*. Academic Press, New York, pp. 3–22.
- Hicks, D.M. and Hume, T.M., 1996. Morphology and size of ebb tidal deltas at natural inlets on open-sea and pocket-beach coasts, North Island, New Zealand. *J. Coast. Res.* 12 (1), 47–63.
- Holgate, S.J. & Woodworth, P.L. 2004. Evidence for enhanced coastal sea level rise during the 1990s. *Geophysical Research Letters* 31, (L07305)
- Hollick, L.M., Shail, R.K. and Levbridge, B.E. 2006. Devonian rift-related sedimentation and Variscan tectonics - new data on the Looe and Gramscatho basins from the resurvey of the Newquay District. *Geoscience in south-west England* 11, 191-198.
- Hu, Z., Wang, Z. B., Zhang, T. J., Stive, M. J. F. and Bouma, T. J. 2015. Predicting long-term and short-term tidal flat morphodynamics using a dynamic equilibrium theory, *J. Geophys. Res. Earth Surf.* 120: 1803–1823, doi:10.1002/2015JF003486.
- Hunt, S., Bryan, R.R., Mullarney, J.C., 2015. The influence of wind and waves on the existence of stable intertidal morphology in meso-tidal estuaries. *Geomorphology* 228, 158–174. <http://dx.doi.org/10.1016/j.geomorph.2014.09.001>.
- Jaboyedol, M., Coppikof, T., Abellan, A., Derron, M. H., Loye, A., Metzger, R., Pedrazzini, A. 2012. Use of LiDAR in landslide investigations: a review. *Nat Hazards* 61, 5–28
- Jaffe, B.E. Smith, B.E., Smith, R.E., Foxgrover, A.C., 2007. Anthropogenic influence on sedimentation and intertidal mudflat change in San Pablo Bay, California: 1856 – 1983. *Estuar. Coast. Shelf Sci.* 73, 175–187.

- Kashani, A. G., Olsen, M. J., Parrish, C. E., and Wilson, N. 2015. A Review of LiDAR Radiometric Processing: From Ad Hoc Intensity Correction to Rigorous Radiometric Calibration. *Sensors* 15, 28099-28128; doi:10.3390/s151128099
- Lane, S.N., Westaway, R.M., Hicks, D.M., 2003. Estimation of erosion and deposition volumes in a large, gravel-bed, braided river using synoptic remote sensing. *Earth Surf. Process. Landf.* 28, 249–271. <https://doi.org/10.1002/esp.483>
- Lee, H. J., Ryu, S. O. 2008. Changes in topography and surface sediments by the Saemangeum dyke in an estuarine complex, west coast of Korea. *Cont. Shelf Res.* 28 (9): 1177–1189.
- Levoy, F., Anthony, E.J., Monfort, O., Robin, N., Bretel, P., 2013. Formation and migration of transverse bars along a tidal sandy coast deduced from multi-temporal LiDAR datasets. *Mar. Geol.* 342, 39–52. <http://dx.doi.org/10.1016/j.margeo.2013.06.007>.
- Leuliette, E.W., Nerem, R.S., & Mitchum, G. 2004. Calibration of TOPEX/Poseidon and Jason altimeter data to construct a continuous record of mean sea level change. *Marine Geodesy* 27, 79-94
- Leuven, J. R. F. W., Kleinhans, M. G., Weisscher, S. A. M., van der Vegt, M. 2016. Tidal sand bar dimensions and shapes in estuaries. *Earth-Science Reviews* 161, 204–223
- Lim, M., Rosser, N. J, Petley, D. N., Keen, M. 2011. Quantifying the controls and influence of tide and wave impacts on coastal rock cliff erosion. *J Coast Res* 27, 46–56
- Lorang, M.S., Stanford, J. A., Hauer, F. R., Jourdain, J. H. 1993. Dissipative and reflective beaches in a large lake and the physical effects of lake level regulation. *Ocean and Coastal Management* 19, 263–287
- Mariotti, G., Fagherazzi, S. 2010. A numerical model for the coupled long-term evolution of salt marshes and tidal flats. *J. Geophys. Res.* 115, F01004. <http://dx.doi.org/10.1029/2009JF001326>.
- Mason, D.C., Scott, T.R., & Wang, H.-J. 2006. Extraction of tidal channel networks from airborne scanning laser altimetry. *ISPRS Journal of Photogrammetry and Remote Sensing* 61(2): 67–83
- Montreuil, A-L., and Bullard, J. E. 2012. A 150-year record of coastline dynamics within a sediment cell: Eastern England. *Geomorphology* 179, 168 – 185.
- Montreuil, A-L., Levoy, F., Bretel, P., Anthony, E. J. 2014. Morphological diversity and complex sediment recirculation on the ebb delta of a microtidal inlet (Normandy, France): A multiple LiDAR dataset approach. *Geomorphology* 219, 114-125 <http://dx.doi.org/10.1016/j.geomorph.2014.05.008>
- Moore, R. D., Wolf, J., Souza, A. J. and Flint, S. S. 2009. Morphological evolution of the

- Dee Estuary, Eastern Irish Sea, UK: A tidal asymmetry approach. *Geomorphology* 103, 588 – 596.
- Nicholls, R.J., Wong, P.P., Burkett, V., Codignotto, J., Hay, J., McLean, R., Page-Edwards, S., Woodroffe, C.D., 2007. Coastal systems and low-lying areas. In: Parry, M.L., Canziani, O.F., Palutikof, J.P., Van Der Linden, P.J., Hanson, C. (Eds.), *Climate Change 2007: Impacts, Adaptation and Vulnerability. Contribution of Working Group II to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK, pp. 315–356.
- Nordstrom, K.F., Jackson, N.L. 1992. Two dimensional change on sand beaches in estuaries. *Zeitschrift für Geomorphologie* 36: 465–478.
- Nordstrom, K.F., Jackson, N.L. 2012. Physical processes and landforms on beaches in short fetch environments in estuaries, small lakes and reservoirs: A review. *Earth-Science Reviews* 111, 232–247
- Nunes, M., Ferreira, O., Loureiro, C. and Baily, B. 2011. Beach and cliff retreat induced by storm groups at Forte Novo, Algarve (Portugal). *Journal of Coastal Research* (SI 64), 795–799
- Oyedotun, T. D. T., Burningham, H., French, J. R., 2012. Characterisation of estuary and adjacent beach sediments in the Camel Estuary, south-west England. *Geosci. South-West England* 13, 70–76
- Oyedotun, T. D. T., Burningham, H., French, J. R., 2013. Sediment sorting and mixing in the Camel Estuary, UK. In: Conley, D. C., Masserlink, G., Russell, P. E. and O'Hare, T. J. (Eds), *Proceedings 12th International Coastal Symposium (Plymouth, England)*, *Journal of Coastal Research* Special Issue 65, 1563–1568.
- Padmalal, D., Kumaran, K.P.N., Nair, K.M., Limaye, R.B., Mohan, S.V., 2014. Consequences of sea level and climate changes on the morphodynamics of a tropical coastal lagoon during Holocene: an evolutionary model. *Quat. Int.* 333, 156-172.
<http://dx.doi.org/10.1016/j.quaint.2013.12.018>.
- Pascoe, H., 2007. *The book of Hayle: portrait of a Cornish coastal town*. Halsgrove, Devon
- Pierce, L.R., 2004. Late waves, coarse clastic beach variability and management implications, Loch Lomond, Scotland, UK. *Journal of Coastal Research* 20, 562–585
- Pirrie, D., Beer, A. J., Camm, G. S. 1999. Early diagenetic sulphide minerals in the Hayle Estuary, Cornwall. *Geosci. South-West England* 9, 325–332
- Pirrie, D., Power, M.R., Payne, G.S., & Wheeler, P.D. 2000. Impact of mining on sedimentation: The Camel and Gannel estuaries, Cornwall. *Geoscience in South-West England*, 10, 021-028

- Pritchard, D., Hogg, A. J., Roberts, W. 2002. Morphological modelling of intertidal mudflats: The role of cross-shore tidal currents, *Cont. Shelf Res.*, 22(11–13), 1887–1895.
- PSMSL, 2016. Newlyn tide gauge data.
[<http://www.psmsl.org/data/obtaining/stations/202.php>]
- Pugh, D. T. 1987. *Tides, surges and mean sea level: a handbook for engineers and scientists* Chichester, Wiley.
- Pye, K., Blott, S. J. 2014. The geomorphology of UK estuaries: the role of geological controls, antecedent conditions and human activities. *Estuar Coast Shelf Sci.* 159, 196–214
- Royal Haskoning 2011, *Cornwall and Isles of Scilly SMP2*, Royal Haskoning
- Ranasinghe, R., Duong, T.M., Uhlenbrook, S., Roelvink, D., Stive, M., 2013. Climate change impact assessment for inlet-interrupted coastlines. *Nat. Clim. Chang.* 3, 83–87.
<http://dx.doi.org/10.1038/NCLIMATE1664>.
- Roberts, W., Le Hir, P., Whitehouse, R. J. S. 2000. Investigation using simple mathematical models of the effect of tidal currents and waves on the profile shape of intertidal mudflats, *Cont. Shelf Res.* 20(10–11), 1079–1091.
- Robins, P.E., Skov, M.W., Lewis, M.J., Giménez, L., Davies, A.G., Malham, S.K., Neill, S.P., McDonald, J.E., Whitton, T.A., Jackson, C.E., Jago, C.F. 2016. Impact of climate change on UK estuaries: a review of past trends and potential projections. *Estuarine, Coastal and Shelf Science* 169: 119–135. doi:10.1016/j.ecss.2015.12.016.
- Sallenger, A. H. Jr, Krabill, W. B., Swart, R. N., Brock, J., List, J., Hansen, M., Holman, R. A., Manizade, S., Sontag, J., Meredith, A., Morgan, K., Yunkel, J. K., Frederick, E. B., Stockdon, H. 2003. Evaluation of Airborne Topographic LiDAR for quantifying beach changes. *J Coast Res* 19, 125–137.
- Savenije, H.H., 2006. *Salinity and Tides in Alluvial Estuaries*. Elsevier.
- Schmidt, K. A., Hadley, F. C., Wijekoon, N. 2011. Vertical accuracy and use of topographic LiDAR data in coastal marshes. *J Coast Res* 27, 116–132
- Shipman, H., 2010. The geomorphic setting of Puget Sound: implications for shoreline erosion and the impacts of erosion control structures. In: Shipman, H., Dethier, M. N., Gelfenbaum, G., Irish, K.L., Dinicola, R.S. (Eds.), *Puget Sound Shorelines and the Impacts of Armoring: Proceedings of a State of the Science Workshop, May 2009*: U.S. Geological Survey Scientific Investigations Report 2010–5254, 19–33.
- Short, A.D., Johnson, D.W.T., 2013. Beach morphodynamics. In: Shroder, J. (Editor in Chief), Sherman, D.J. (Ed.), *Treatise on Geomorphology*. Academic Press, San Diego, CA, 10, Coastal Geomorphology, pp. 106–129.
- Syvistki, J.P.M., Milliman, J.D., 2007. *Geology, geography, and humans battle for dominance*

- over the delivery of fluvial sediment to the coastal ocean. *J. Geol.* 115, 1–19.
- Teasdale, P.A., Collins, P.E.F., Firth, C.R., & Cundy, B.A. 2011. Recent estuarine sedimentation rates from shallow intertidal environments in western Scotland: implications for future sea level trends and coastal wetland development. *Quaternary Science Reviews*, 30, 109-129
- UK Hydrographic Office (UKHO). 2003. Admiralty Tide Tables: Volume 1, United Kingdom and Ireland (including European channel ports). United Kingdom Hydrographic Office, London.
- Uncles, R. J., Stephen, J. A., Harris, C. 2015. Estuaries of south west England: salinity, suspended particulate matter, loss-on-ignition and morphology. *Prog Oceanogr* 137, 385–408.
- Wei, W., Dai, Z., Mei, X., Liu, J. P., Gao, S., & Li, S. 2017. Shoal morphodynamics of the Changjiang (Yangtze) estuary: Influences from river damming, estuarine hydraulic engineering and reclamation projects. *Marine Geology* 386, 32–43. <https://doi.org/10.1016/j.margeo.2017.02.013>
- Woodworth, P.L., White, N.J., Jevrejeva, S., Holgate, S., Church, J.A., & Gehrels, W.R. 2009. Evidence for the accelerations of sea level on multi-decade and century timescales. *International Journal of Climatology* 29 (6), 777-789
- Wang, Z.B., Van Maren, D.S., Ding, P.X., Yang, S.L., Van Prooijen, B.C., De Vet, P.L.M., Winterwerp, J.C., Vriend, H.J., Stive, M.J.F., He, Q. (2015) Human impacts on morphodynamic thresholds in estuarine systems. *Continental Shelf Research* 111, 174 – 183. <http://dx.doi.org/10.1016/j.csr.2015.08.009>
- Wong, P.P., Losada, I.J., Gratiuso, J.P., Hinkel, J., Khattabi, A., McInnes, K.L., Saito, Y., Sallenger, A., 2014. Coastal systems and low-lying areas. In: Field, C.B., Barros, V.R., Dokken, D.J., Mach, K.J., Mastrandrea, M.D., Bilir, T.E., Chatterjee, M., Ebi, K.L., Estrada, Y.O., Genova, R.C., Girma, B., Kissel, E.S., Levy, A.N., MacCracken, S., Mastrandrea, P.R., White, L.L. (Eds.), *Climate Change 2014: Impacts, Adaptation, and Vulnerability. Part A: Global and Sectoral Aspects. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, NY, USA, 61–109.
- Young, A. P., Ashford, S. A. 2006. Application of airborne LiDAR for sea cliff volumetric change and beach-sediment budget contributions. *J Coast Res* 22, 307–318.
- Young, A. P., Flick, R. E., Gutierrez, R., Guza, R. T. 2009. Comparison of short-term sea cliff retreat measurement methods in Del Mar, California. *Geomorphology* 112, 318–323.

- Young, A. P., Guza, R. T., O'Reilly, W. C., Flick, R. E., Gutierrez, R. 2011. Short term retreat statistics of a slowly eroding coastal cliff. *Nat Hazards Earth Syst Sci* 11, 205–217.
- Zhang, K., Whitman, D., Leatherman, S., Robertson, W. 2005. Quantification of beach changes caused by hurricane Floyd along Florida's Atlantic coast using airborne laser surveys. *J Coast Res* 21, 123–134.
- Zhu, Q., Yang, S., Ma, Y. 2014. Intra-tidal sedimentary processes associated with combined wave–current action on an exposed, erosional mudflat, southern Yangtze River Delta, China, *Mar. Geol.* 347, 95–106, doi:10.1016/j.margeo.2013.11.005.