Towards first-principles-level polarization energies in force fields. A Gaussian basis for the atom-condensed Kohn-Sham method (ACKS2)

Patrick Gütlein,[†] Lucas Lang,[†] Karsten Reuter,[†] Jochen Blumberger,^{‡,¶} and
Harald Oberhofer^{*,†}

[†]*Chair for Theoretical Chemistry and Catalysis Research Center, Technische Universität
München, Lichtenbergstrasse 4, D-85747 Garching, Germany*

[‡]*Department of Physics and Astronomy, University College London, London WC1E 6BT,
UK.*

[¶]*Institute for Advanced Study, Technische Universität München, Lichtenbergstrasse 2 a,
D-85748 Garching, Germany.*

E-mail: harald.oberhofer@tum.de

Abstract

The last twenty years of force field development have shown that even well parametrized classical models need to at least approximate the dielectric response of molecular systems—based e.g. on atomic polarizabilities—in order to correctly render their structural and dynamic properties. Yet, despite great advances most approaches tend to be based on *ad hoc* assumptions and often insufficiently capture the dielectric response of the system to external perturbations, such as e.g. charge carriers in semiconducting materials. A possible remedy was recently introduced with the atom-condensed Kohn-Sham density-functional theory approximated to second order (ACKS2), which is fully

derived from first principles. Unfortunately, specifically its reliance on first-principles derived parameters so far precluded the widespread adoption of ACKS2. Opening up ACKS2 for general use, we here present a reformulation of the method in terms of Gaussian basis functions, which allows us to determine many of the ACKS2 parameters analytically. Two sets of parameters depending on exchange-correlation interactions are still calculated numerically, but we show that they could be straightforwardly parametrized owing to the smoothness of the new basis. Our approach exhibits three crucial benefits for future applications in force fields, *i*) efficiency, *ii*) accuracy, and *iii*) transferability. We numerically validate our Gaussian augmented ACKS2 model for a set of small hydrocarbons which shows a very good agreement with density-functional theory reference calculations. To further demonstrate the method’s accuracy and transferability for realistic systems, we calculate polarization responses and energies of anthracene and tetracene, two major building blocks in organic semiconductors.

1 Introduction

Owing to their novel materials properties and the vast chemical space they span, molecular organic semiconductors (OS) have attracted the interest of both science and industry for many years.^{1–5} In parallel to applied OS research, there is a vibrant community aiming to understand the intrinsic electronic properties and fundamental mechanisms involved in charge and energy transfer processes.^{6–12} To the theorist, an extensive toolbox of methods is available to model these compounds, often involving molecular dynamics simulations as an integral component to sample structures and responses to external perturbations.^{9,10,13}

Since the dawn of molecular simulation, classical force fields (FF) have been used to efficiently sample the phase space of systems too large to be tackled from first principles.^{14–17} Unfortunately, a correct dielectric response to external perturbation poses a severe challenge for an effective FF treatment where electrostatic interactions are usually treated in terms of

fixed charges or multipoles.^{18–22} A number of approaches have been put forward to tackle this problem, typically based on an atom-centered expansion of classical multipoles.^{23–25} An early and successful example of this is the electronegativity equalization method (EEM),^{26,27} where a population of fluctuating atomic charges is applied to model the formation of molecular bonds as well as rearrangements of the charge density upon perturbation (cf. Fig. 1).

Figure 1: Illustration of the perturbation of the charge density of a tetracene molecule (black scaffold) upon responding to a point charge of $+1 e$ on the right (black sphere). Charge accumulation vs. ground state is depicted in blue ($1 \times 10^{-3} e a_0^{-3}$) and reduction ($-1 \times 10^{-3} e a_0^{-3}$) in red density difference iso-surfaces. Calculated with FHI-aims and the settings described in the text.

This simple representation is further refined in subsequently developed methods, like the chemical potential equalization method^{28,29} or charge equilibration model.³⁰ Later models, like the atom-atom charge transfer theory³¹ and the split-charge equilibration scheme,³² tackle the more fundamental shortcomings of EEM, related to the overestimation of long range charge transfer in dielectrics and vacuum. Yet, so far only *ad hoc* corrections employing inter-atomic charge constraints based on topological considerations alleviated these limitations, while still manacled to daunting calibration procedures due to the nature of the split-charge variable.^{33,34}

Aspiring to eliminate the fundamental issues of previous methods, the atom-condensed Kohn-Sham density-functional theory approximated to second order (ACKS2)^{34,35} has been developed on the foundations of EEM. It is a linear electronic response theory derived entirely from first principles and hence provides an exact *ab initio* many-body polarization correction to semi-empirical interaction potentials. **Thereby, ACKS2 rests on an atom-projected description of changes to charge density and Kohn-Sham potential due to external perturbations such as applied electric fields or other molecules. It thus also bears some conceptional**

similarity to distributed polarizability approaches pioneered e.g. by Misquitta and Stone,^{36,37} where perturbative responses of molecules are separated into individual atomic contributions. However, in terms of practical applications, the ACKS2 method has so far only been used to complement more traditional EEM models,³⁸ due to the fact that full ACKS2 response calculations still rely on system dependent parameters derived from density functional theory (DFT) calculations. For the same reasons, ACKS2 has to date never been applied to the calculation of electronic polarization energies, to the best of our knowledge.

In this work we thus present an efficient implementation of the ACKS2 method allowing, for the first time, the calculation of electronic polarization energies on top of the polarization of the electronic density. In order to demonstrate its efficacy we then apply our implementation to calculate the charge responses and energies of small and medium sized organic molecules to several external fields. Following the original parametrization recipe,³⁵ we choose a Gaussian representation of the electronic linear response, i.e. changes to the electron density and Kohn-Sham potential, in which many of the ACKS2 energy terms can be calculated analytically. This results in a formally exact description (up to second order) of the electronic polarization response without the need for time consuming self-consistent iteration of the electron density for each external perturbation, as e.g. in DFT. Yet, a few parameters depending on exchange and correlation terms of the KS DFT remain. These need to be parametrized from a single, unperturbed DFT calculation, which is facilitated by the smoothness of our new basis. We demonstrate the accuracy of our approach by calculating the electronic polarization energy of a set of hydrocarbons and π -conjugated organic molecules forming crystalline semiconductors. Compared to DFT reference calculations, we obtain very small errors, while the computational cost currently exceeds standard force field calculations—i.e. with fixed bonding topologies and either fixed- or variable charge representations—due to the few parameters still dependent on electronic structure calculations. These will be discussed in detail below. Hence, this work lays the foundation for MD simulations with rigorous and efficient ACKS2-based polarization responses.

1.1 Theoretical Background of the ACKS2 Method

In order to determine the electronic response to an external perturbation regular Kohn-Sham (KS) DFT offers two distinct approaches. First, one could perform self-consistent DFT calculations of both perturbed and un-perturbed system state, to take their difference to compute the changes to the electronic density, the KS potential, and thus the energy. Second, for small perturbations, one could employ a linear response scheme, using density functional perturbation theory (DFPT),^{39,40} to determine these properties in a single calculation.

The ACKS2 method follows the latter approach in that it is a simplified linear electronic response model derived from regular KS DFT. At its heart lies a linear equation, derived using the Euler-Lagrange formalism, which describes the response of the electron density to the external perturbation v_{ext} , while a second coupled linear equation accounts for changes of the auxiliary Kohn-Sham potential induced by the density changes. A third and final equation adds a constraint on the total number of electrons for any density rearrangements. Note that this formulation only considers induced changes of the electron density and KS potential and hence cannot describe properties of the unperturbed system state. In order to provide an efficient electronic response scheme for force field simulations, two crucial approximations are introduced to the full KS linear response formalism to arrive at the ACKS2 method. First, in analogy to linear DFPT,³⁹ the functional derivatives of the energy terms are linearized. This is a particularly important simplification of the KS DFT equations, as—in the limit of a linear electronic response—it removes the necessity of a self-consistent iteration scheme. Second, representations of density and potential response are chosen to be atom-condensed, i.e. all basis sets and parameters are mapped to atom centers similar to previous equilibration schemes like EEM, in order to provide transferability between different geometries.

Considering polarization models, the significant innovation of the ACKS2 approach over e.g. the EEM method is now to not only account for the density change $\Delta\rho$ but also to explicitly treat all induced changes to the auxiliary KS potential Δv_{KS} , both in mean-field as well as xc components. To make them computationally accessible, both are expanded in

a series of atom-centered basis functions,

$$\Delta\rho(\mathbf{r}) = \sum_i^N c_i g_i(\mathbf{r}), \quad (1a)$$

$$\Delta v_{\text{KS}}(\mathbf{r}) = \sum_j^M d_j h_j(\mathbf{r}), \quad (1b)$$

where g_i (h_j) denote the N (M) basis functions while c_i (d_j) denote the expansion coefficients of the induced density (potential) changes $\Delta\rho$ (Δv_{KS}).

Given the atom-centered basis set expansions of eqs. 1, the parametrized form of the KS-DFT linear response, i.e. the working equations of ACKS2, read:

$$\begin{bmatrix} [\eta_{i,i}]_{N,N} & [O_{i,j}]_{N,M} & [D_i]_N \\ [O_{j,i}]_{M,N} & [\chi_{j,j}]_{M,M} & 0 \\ [D_i]_N & 0 & 0 \end{bmatrix} \begin{bmatrix} [c_i]_N \\ [d_j]_M \\ \Delta\tilde{\mu} \end{bmatrix} = \begin{bmatrix} [V_i]_N \\ 0 \\ 0 \end{bmatrix}. \quad (2)$$

The matrix on the left hand side of eq. (2) contains the entire KS electronic structure-dependent linear response information of the system and assigns parameters to each basis function. Contrary to earlier schemes,^{28–30} the ACKS2 hardness sub-matrix $\eta_{N,N}$ accounts on a DFT level for both, changes of the Coulomb energy—which was accounted for in earlier schemes—and of the exchange-correlation energies—which was either estimated empirically or neglected. Both energy contributions described by $\eta_{N,N}$ are thereby due to rearrangements of the electron density. The non-interacting linear response kernel parameters $\chi_{M,M}$, on the other hand, incorporate changes of the KS energy induced by changes of the KS potential.

$$\eta_{kl} = \int \int g_k(\mathbf{r}) \left(\frac{1}{|\mathbf{r} - \mathbf{r}'|} + \frac{\partial^2 E^{\text{xc}}[\rho]}{\partial \rho(\mathbf{r}) \partial \rho(\mathbf{r}')} \right) g_l(\mathbf{r}') d\mathbf{r} d\mathbf{r}', \quad (3a)$$

$$\chi_{kl} = \int \int h_k(\mathbf{r}) \left(\frac{\partial^2 E^{\text{KS}}[v_{\text{KS}}]}{\partial v_{\text{KS}}(\mathbf{r}) \partial v_{\text{KS}}(\mathbf{r}')} \right) h_l(\mathbf{r}') d\mathbf{r} d\mathbf{r}'. \quad (3b)$$

Energy changes due to concurrent perturbations of the charge density and KS potential are

mediated through the basis set overlap integrals, $O_{kl} = \int g_k(\mathbf{r})h_l(\mathbf{r})d\mathbf{r}$. Additionally, the last row of the left hand side working matrix (eq. (2)) contains the integrals of the density basis functions, $D_k = \int g_k(\mathbf{r})d\mathbf{r}$, to establish a constraint on the total charge of the system. In analogy to KS DFT the change in chemical potential $\Delta\tilde{\mu}$ thereby mediates the charge constraint in the form of a Lagrange multiplier. The right hand side of eq. (2), introduces external perturbations acting on the charge density, $V_k = \int g_k(\mathbf{r})\Delta v_{\text{ext}}(\mathbf{r})d\mathbf{r}$. Finally, the ACKS2 based linear electronic response, represented by the expansion coefficients c_i and d_j , is calculated by solving eq. (2).

In our reformulation of the ACKS2 approach, a subset of parameters (O , D , and the Coulomb integral in η) can be calculated analytically, i.e. independently of the KS-DFT based electronic structure, while the remainders (χ and the exchange and correlation contributions to η) are evaluated exactly from a reference KS-DFT calculation. Hence, the ACKS2 method theoretically reproduces the exact electronic response of the underlying DFT reference up to second order in the limit of linear response and a complete basis set representation. While the limitation to second order effects at first may seem overly restrictive, recent work e.g. by Giese and York,^{41,42} has shown that carefully conducted second order expansions can yield polarization responses and even geometries nearly indistinguishable from the full KS DFT case. In section 2 we show ACKS2 to go even one step further, yielding excellent agreement not only for the induced molecular dipole moments but also the polarization energy.

1.1.1 ACKS2 Energy Expression

To be useful in actual applications, ACKS2 needs to be able to reproduce a molecule’s electronic response to an external potential. Here we focus on the induced dipole moment and the energetic cost of that response known as the polarization energy. While the induced dipole moment follows straightforwardly from the density response (eq. (1a)), the expression

for the energy response expanded in a Taylor-series up to second order is more involved:³⁵

$$\begin{aligned}\Delta E &= E_{v_0+\Delta v_{\text{ext}}}[\rho_0 + \Delta\rho, v_{\text{KS},0} + \Delta v_{\text{KS}}] - E_{v_0}[\rho_0, v_{\text{KS},0}] \\ &\approx \int \rho_0(\mathbf{r})v_{\text{ext}}(\mathbf{r})d\mathbf{r} + \frac{1}{2} \sum_{i,i'}^{N,N'} c_i \eta_{i,i'} c_{i'} + \frac{1}{2} \sum_{j,j'}^{M,M'} d_j \chi_{j,j'} d_{j'} - \sum_{i,j}^{N,M} c_i O_{i,j} d_j + \sum_i^N c_i V_i,\end{aligned}\tag{4a}$$

$$\Delta E_{\text{ACKS2}}^{\text{pol.}} = \frac{1}{2} \sum_{i,i'}^{N,N'} c_i \eta_{i,i'} c_{i'} + \frac{1}{2} \sum_{j,j'}^{M,M'} d_j \chi_{j,j'} d_{j'} - \sum_{i,j}^{N,M} c_i O_{i,j} d_j + \sum_i^N c_i V_i,\tag{4b}$$

$$\Delta E_{\text{ACKS2}}^{\text{intra}} = \frac{1}{2} \sum_{i,i'}^{N,N'} c_i \eta_{i,i'} c_{i'} + \frac{1}{2} \sum_{j,j'}^{M,M'} d_j \chi_{j,j'} d_{j'} - \sum_{i,j}^{N,M} c_i O_{i,j} d_j,\tag{4c}$$

$$\Delta E_{\text{ACKS2}}^{\text{inter}} = \sum_i^N c_i V_i.\tag{4d}$$

The first term on the right hand side of eq. (4a) represents the interaction energy of the unperturbed reference charge density (of both electrons and nuclei), ρ_0 , with the external perturbation v_{ext} . The ACKS2 methodology does not account for this term, as it operates in the space of the *induced* charge density $\Delta\rho$ and KS-potential Δv_{KS} only. Yet, it could easily and efficiently be calculated e.g. from a static force field’s multipole representation of the unperturbed charge density.^{19,21,22} The energy of induced charge and KS potential polarization $\Delta E_{\text{ACKS2}}^{\text{pol.}}$ provide a static response correction accounting for induction effects. The term $\Delta E_{\text{ACKS2}}^{\text{intra}}$ describes the intramolecular polarization energy necessary to rearrange the charge density of the reference state, while $\Delta E_{\text{ACKS2}}^{\text{inter}}$ is the interaction energy of the induced charge density with the external perturbation. Evaluation of the ACKS2 polarization energy, and hence the computational expense to be expected in its application are determined by the vector-matrix-vector multiplications and a single vector-vector scalar product, c.f. eq. (4b). The induced dipole moment is then a simple scalar product of the density expansion coefficients and density basis function times dipole operator integrals, which can

simply be tabulated for a specific Gaussian type density basis set,

$$\Delta\mu_{\text{ACKS2}} = \sum_i c_i \int g_i(\mathbf{r}) \mathbf{r} d\mathbf{r}. \quad (5)$$

1.2 A Gaussian Basis for ACKS2

Inspired by the point charges in classical FFs, earlier studies expanded the response of the electron density and KS potential in Fukui and Hirshfeld functions.^{34,35} While these implementations show a relatively high accuracy, they produce a large computational overhead as the basis functions directly depend on the electronic structure of the molecule. In contrast, our implementation employs a set of primitive Cartesian Gaussian functions (GF) centered at the atomic positions. Next to a greatly increased computational efficiency, this choice of representation has the added advantages of a tunable accuracy and—even more importantly—basis set transferability. Thereby, the accuracy of the ACKS2 representation can straightforwardly be increased by adding higher angular momentum basis functions. As will be made clear below, the (atom-centered) basis functions are solely element specific, which ensures the transferability among different molecular structures.

By derivation, the ACKS2 parameters $\chi_{M,M}$, $\eta_{M,M}$, D_N , $O_{M,N}$ and V_N in eq. (2) are KS-DFT expectation values and need to be obtained from DFT calculations. In contrast to earlier incarnations of ACKS2, our Gaussian representation allows many of the parameters to be calculated analytically. Nevertheless, the evaluation of $\chi_{M,M}$ as well as exchange-correlation contributions to $\eta_{M,M}$ currently necessitates a KS-DFT electronic structure, whereas coulomb-interaction portions of $\eta_{M,M}$ as well as all terms in D_N , $O_{M,N}$, and V_N are obtained directly by recursive GF integration schemes. Hence, computational cost of the analytically-derived parameters depends on the specific software implementation of the recursion relations introduced in the supporting information. The software thereby used is an in-house proof-of-concept Python package to be made available upon reasonable request. With a view on the use of ACKS2 for force fields, preliminary studies indicate that the

ACKS2 parameters that require explicit DFT calculation depend very smoothly on nuclear geometry, see below. Hence, DFT calculations on only one or a few molecular geometries should be sufficient for parametrization of an ACKS2-based force field and drastically improve the computational cost, **which in this case would then solely be due to the necessary inversion of the ACKS2 response matrices.**

2 Results

2.1 Fitting the New Basis Set

We illustrate our Gaussian basis set implementation of ACKS2 for carbon and hydrogen. We choose a minimal basis of Cartesian s- and p-type Gaussian functions for both the electron density and Kohn-Sham potential responses to balance good accuracy with high computational efficiency of the ACKS2 method as well as a manageable procedure for basis set optimization. The s-type functions describe atom-atom charge transfer, whereas the p-type functions account for intra-atomic dipole polarization. The latter are particularly important for planar or linear molecular fragments often encountered, e.g. in organic semiconductor materials and biochemical systems. Together, both types of functions can be expected to cover the basic electronic response to external electrostatic perturbations, both in-plane and perpendicular to the molecular plane.

An atomic species-dependent sp-GF basis set is generated independently for both carbon and hydrogen by optimizing the width σ of the Gaussian functions:

$$f_{a,b,c}(\mathbf{r}, \sigma, \mathbf{R}_A) = (x - X_A)^a (y - Y_A)^b (z - Z_A)^c \times \exp\left(-\frac{1}{2\sigma^2}(\mathbf{r} - \mathbf{R}_A)^2\right), \quad (6)$$

where $\mathbf{R}_A = \{X_A, Y_A, Z_A\}$ denotes the position of atom A and $\mathbf{r} = \{x, y, z\}$ is the position vector. To find the optimal σ for each elementary s- and p-function, atomic dimers of H and C are subjected individually to a set of external electrostatic perturbations. The

basis sets are then optimized for the ACKS2 response to match the full KS-DFT reference by variation of the width (or radial decay) σ of the Gaussian functions separately for the representations of electron density and KS-potential response, respectively. The space of possible perturbations is sampled by a radial and angular distribution of point charges (PC) of +1e placed between 1.3 times and up to 3.5 times the van-der-Waals radii, complemented by homogeneous electric field (HEF) potentials of similar strengths, yielding a total of 110 and 130 different perturbation potentials for carbon and hydrogen, respectively. A more detailed description of our fitting set can be found in the supporting information.

The quality of the basis set is measured by the relative mean error (RME) of the ACKS2 response from the respective full KS-DFT reference,

$$\text{RME} = \frac{1}{N_{\text{data}}} \sum_i^{N_{\text{data}}} \left| \frac{x_i^{\text{ACKS2}} - x_i^{\text{DFT}}}{x_i^{\text{DFT}}} \right|, \quad (7)$$

where x represents the property of interest, e.g. induced dipole moment, intramolecular rearrangement energy $\Delta E_{\text{ACKS2}}^{\text{intra}}$ (eq. (4c)) and intermolecular interaction energy $\Delta E_{\text{ACKS2}}^{\text{inter}}$ (eq. (4d)), respectively. Screening of the four-dimensional parameter space (σ_s and σ_p for the electron density and KS potential response functions, respectively) exposed a great number of local minima. Therefore, we applied a global particle swarm optimizer⁴³ to select the global minimum among several local minima with similar fitness scores, as shown in table 1. For the present basis set representation, the radial decay matches the order of magnitude of atomic valence shells. This is not surprising, given that the basis needs to represent changes to the density and the potential, respectively. However, this being an atom-condensed approximation of these quantities, likely dependent on the angular momentum order of the expansion, one should not expect a physical relevance of our σ values beyond the correct order of magnitude.

The results obtained with the best basis set are shown in fig. 2. For H₂ the relative mean error of induced dipole moments and polarization energy are 4.4% and 4.1% with respect

to KS-DFT, while they are 2.5% and 6.5% for C_2 , respectively. The optimized sp-GF basis shows excellent agreement with DFT references for induced dipole moments (A, D) and for the polarization energy (B, E) over three orders of magnitude, from 0.01 to 1.0 eV. In general, ACKS2 predicts the response to HEF perturbations (blue) better than to PC perturbations (red), in particular for the carbon dimer. A more detailed view of the differences in panels (C, F) show deviations for HEF perturbations to follow a rather narrow distribution up to 5% , while distributions for PC perturbations exhibit a flat-bottomed tail up to 20% error.

An exemplary illustration of the density changes due to external perturbations is shown in fig. 3. Here, a one-dimensional cut of the electronic response to a single HEF or PC along the bond of a hydrogen dimer reveals the simplicity of the presented approach based on a sp-GF minimal basis set. Evidently, the fine details of the density response around the nuclei are not and cannot be resolved as the basis functions are not flexible enough. Nevertheless, the very good agreement of integrated properties—i.e. energies and induced dipoles—with respect to DFT demonstrate that only a sufficiently accurate averaged ACKS2 description of charge and potential response is required. **Our effective basis thus follows the spirit of other effective methods such as e.g. pseudo-potentials, which are designed to reproduce orbital energies without reproducing the nodal features of the orbitals close to the nucleus. A more in-depth study of the convergence of the density response near the cores with basis set size is beyond the scope of this work.**

Table 1: Optimized radial parameters, i.e. Gaussian widths σ [a_0] for an sp-type ACKS2 basis set for hydrogen and carbon. Note the different parameters for the density and the potential basis, respectively.

	$\sigma_{\Delta\rho}^{(s)}$	$\sigma_{\Delta\rho}^{(p)}$	$\sigma_{\Delta v_{KS}}^{(s)}$	$\sigma_{\Delta v_{KS}}^{(p)}$
H	0.608	0.863	0.490	1.000
C	0.446	1.174	0.637	5.652

Figure 2: Performance of the fitted sp-GF minimal basis for the hydrogen (A, B, C) and carbon (D, E, F) training sets. Panel (A) and (D) illustrate the induced dipole moment, panel (B) and (E) depict the polarization energy of the ACKS2 method, c.f. eq. (4a), compared to the DFT reference. The relative error is summarized in panels (C) and (F).

2.2 Small Molecule Test-Set

To test the transferability of the Gaussian basis from H₂ and C₂ to more general CH-containing molecules, we curated a set of ten small hydrocarbons. They were selected for their different chemical features, i.e. four linear alkanes, three compounds containing single, double and triple bonds C₂H_{2,4,6} and four unsaturated cyclic hydrocarbons, cf. table 2. Similar to the training procedure, each molecule is separately exposed to a set of external perturbations sampling the three-dimensional space by a radial and angular distribution of HEF and PC potentials, yielding 30 data points per molecule and 300 data points in total for the test set.

Figure 3: One-dimensional cuts of the induced electron density rearrangements in H_2 . The external perturbations are aligned with the molecular bond axis. The response to a PC with center-of-mass distance of $2.0 \text{ \AA}$ and single elemental charge and a homogeneous electric field matching the PC potential, are illustrated.

Figure 4: Performance of the ACKS2 method in predicting induced dipole moment (A) and polarization energy (B). The left column depicts the absolute induced dipole moment and polarization energy, while the right column displays the relative errors of the linear one-dim. alkanes (C), $\text{C}_2\text{H}_{\{2,4,6\}}$ compounds (D) and cyclic molecules subsets (E).

Again, our ACKS2 implementation is capable to predict the DFT reference electronic response properties over an order of magnitude with good accuracy, as illustrated in fig. 4. The induced dipole moment, panel (A), and polarization energy, panel (B), match the DFT reference data very well. The slight underestimation of both induced dipole and polarization

energy can be attributed to a slight overestimation of the intramolecular polarization energy contributions compared to the intermolecular interactions. We believe this to be a side effect of the minimal basis set representation. To our knowledge, this is for the first time that the polarization energy $\Delta E_{\text{ACKS2}}^{\text{pol}}$ has been determined solely within the ACKS2 framework. Our results further show that atom-centered basis sets, though optimized for simple diatomics, are transferable to other molecules with different chemical features (bonds between heteroatoms, double, triple bond etc). Unfortunately, simple stacking of more basis functions does not necessarily improve the basis set representation, as the ACKS2 method does not follow the variational principle of KS-DFT due to the perturbative treatment of the KS response matrix χ .^{34,35} Again, deviations are generally smaller for HEF perturbations (blue) compared to external PCs (red) as illustrated by the distributions of relative errors given fig. 4 C, D, and E.

The subset of linear alkanes exhibits narrow distributions in the 7-10% range, with a slight tail towards larger errors for PC perturbations. Hence, accuracy slightly falls off compared to the training set data averaging at relative errors of about 4.5%. Nevertheless, such small errors provide evidence for the transferability of the simple GF basis set representation. A more detailed view, given in table 2, indicates small increases in error with increasing chain size except ethane, as it resembles the geometric structure of the carbon training dimer C_2 , the closest in the present test set.

The $\text{C}_2\text{H}_{\{2,4,6\}}$ compounds mimic the effect of different chemical bonding types and local electronic structure. They display broader distributions of errors, fig. 4, with shallow tails up to the 30% range, but are again centered at the 10% mark. A closer look reveals an increased error with larger bond orders, table 2, due to enhanced non-linear response effects in π -conjugated organic systems, in particular triple bonded carbon.

Finally, the set of cyclic unsaturated hydrocarbons, relevant for application in OS material modeling, reproduces the DFT reference reasonably well. The distribution of errors, cf. fig. 4, is again centered around the 10% mark. It closely resembles the deviations of the

linear alkane chains, but exhibits a slightly enhanced tail up to the 30% mark, similar to the $\text{C}_2\text{H}_{\{2,4,6\}}$ molecules.

2.3 Acenes

Figure 5: Numerical validation for anthracene and tetracene polarization properties. In panel (A), the induced dipole moment of the ACKS2 method is depicted vs. the DFT reference, panel (B) shows the polarization energy defined in eq. (4a). RME gives the relative mean error for each property.

While the test-set comprised of simple molecules already demonstrates the efficacy of ACKS2 for different chemical species and bonding situations, we now turn to two larger molecules that form semiconducting organic crystals, anthracene and tetracene.⁴⁴ The electronic response properties of these acenes are not trivial and exhibit complex effects as illustrated in fig. 1. For a number of HEF and PC perturbations the ACKS2 response again matches the DFT reference properties very well as depicted in fig. 5. Hence, the present work provides a crucial first step towards an ACKS2-based electronic response function for large, application-relevant molecules.

2.4 Polarizability and Raman intensity

We now turn to the calculation of polarizability and the nuclear derivative of polarizability determining the intensity of Raman spectra, where other effective polarization models often struggle to yield accurate results.^{45,46}

To this end, we compute the isotropic polarizability—i.e. the average of the diagonal of the polarizability tensor, $\alpha^{\text{iso}} = \frac{1}{3}(\alpha_{xx} + \alpha_{yy} + \alpha_{zz})$ —and its dependence on the molecular structure, by perturbing the equilibrium geometry of benzene along the symmetric ring-breathing mode.

Here, the polarizability is calculated numerically by a finite electric field differentiation of the total molecular dipole moment with a field strength of $0.01 \text{ V \AA}^{-1}$. Figure 6 depicts the isotropic polarizability for small displacements along the benzene breathing mode. Separately, fig. 6 highlights changes of α^{iso} with respect to the polarizability in the vibrational equilibrium geometry— $\Delta\alpha^{\text{iso}} = \alpha_{\text{vib}}^{\text{iso}} - \alpha_{\text{equ}}^{\text{iso}}$. The overall trend of total isotropic molecular polarizabilities of benzene, relative mean error of 12.6%, matches the accuracy of previously introduced molecular test sets of small organic molecules and acenes, see figs. 4 and 5. However, a closer look reveals a slight, albeit systematic growth of the relative error of the isotropic molecular polarizabilities outside the equilibrium configuration. This is most likely a consequence of the small basis set, fitted to yield a reasonable approximation of the overall electronic response properties, but potentially too insensitive to capture quantitatively the small changes in polarization response with nuclear displacements.

Finally, we point out that each of the distorted geometries still necessitated a full DFT calculation to extract the ACKS2 response matrices as discussed in detail in the next section.

2.5 Non-analytic parameters of ACKS2

Envisioning the ACKS2 method as polarization correction to force fields, the current bottleneck remains the evaluation of the Kohn-Sham DFT electronic structure dependent pa-

Figure 6: Illustration of the isotropic polarizability and its relative changes for a benzene symmetric breathing mode.

rameters, namely the exchange-correlation contributions to the hardness— $\eta_{i,j}^{\text{xc}}$ —as well as the non-interacting linear response kernel— $\chi_{i,j}$. Our preliminary studies of those ACKS2 parameters indicate a smooth dependency on nuclear geometry due to our smooth Gaussian representation of the response properties. We exemplify this on benzene, c.f. fig. 7a, where we symmetrically alter the equilibrium distance of all C-H bonds ($\Delta d_{\text{C,H}}^{\text{bond}} = 1.09 \text{ \AA}$) over a range of $-0.075 \text{ \AA}$ to $0.075 \text{ \AA}$ by equally displacing the C atoms inwards (outwards) and the H atoms outwards (inwards) for a bond expansion (contraction). This then also changes the C-C distances in the ring, where the respective change of the pairwise distance between two adjacent C atoms (ortho), two second-nearest C atoms (meta) and two third-nearest C atoms (para) on the ring follows simple geometric dependencies on the C-H bond length ($\Delta d_{\text{C,C}}^{\text{ortho}} = 1.40 \text{ \AA}$, $\Delta d_{\text{C,C}}^{\text{meta}} = 2.42 \text{ \AA}$, $\Delta d_{\text{C,C}}^{\text{para}} = 2.79 \text{ \AA}$). As illustrated in fig. 7b, the corresponding response matrix elements of the exchange-correlation contributions as well as the non-interacting linear response kernel vary smoothly near the equilibrium interatomic positions. For the sake of simplicity we here focus on a pure s-basis for both density and potential and plot carbon-carbon as well as bonded carbon-hydrogen interaction elements. Hydrogen-hydrogen terms as well as hydrogen-carbon terms of atoms, that are not directly bonded,

are not shown as we find them negligibly small. The exchange-correlation contributions decay to zero rapidly with distance, as shown in the top graph of fig. 7b. Hence, same-center contributions, depicted in fig. 7c, and off-center matrix elements of atoms partaking in a bond are the most important features of $\eta_{i,j}^{\text{xc}}$ and could be tabulated or interpolated smoothly from a DFT reference. The elements of the non-interacting linear response kernel $\chi_{i,j}$ generally decay less rapidly and involve considerable non-local contributions due to the long-range effects of the Kohn-Sham potential changes. Nevertheless, they vary smoothly with interatomic distances, which should allow for interpolation from a few DFT reference calculations. Considering basis functions of higher angular momentum like p-type GF, the ACKS2 model requires descriptors that are able to account for an angular resolution of the latter response parameters as well compared to a simple interatomic distance metric of an s-type representation, as illustrated here. One candidate for such a metric could e.g. be a simple GF basis overlap, which naturally encodes distances and relative orientations of basis functions. We found that both $\eta_{i,j}^{\text{xc}}$ and $\chi_{i,j}$ still vary smoothly with their respective overlap matrix elements (see supporting information). Hence, for a given molecule or set of molecules, a careful parametrization of an ACKS2 model from a set of a few DFT reference calculations should straightforwardly be possible, and greatly facilitate the method’s application e.g. as a polarization correction in molecular dynamics calculations.

2.6 Comparison to Other Polarizeable Methods

Setting our results into perspective, even state of the art approximations to DFT such as the semi-empirical modified neglect of differential overlap (MNDO) tend to struggle to accurately reproduce polarization responses with relative mean errors (RME) for the induced dipoles of 25%,⁴⁷ compared to hybrid-level DFT. Similar failures can be found for straightforward applications of density functional tight binding DFTB2.⁴² Both of these can be substantially improved though, by coupling the respective ground state electronic structures

to carefully parametrized implementations of the chemical potential equalization (CPE). A single post-processing step of CPE, for example, yields an RME of 12% and 3% for induced dipole moments and molecular polarizabilities, respectively for MNDO,⁴⁷ and 5% for molecular polarizabilities in DFTB2.⁴² Using a careful parametrization, similar results—RME of 12% for induced dipole moments and RME of 4% for molecular isotropic polarizabilities—can also be found for a direct application of CPE²⁹ to a set of small organic molecules, very similar to the set used here (cf. section 2.2). On the more empirical side, the popular Drude oscillator model for example has been shown to yield polarizabilities down to 3%,⁴⁸ if carefully parametrized, albeit at the potential cost of a loss of transferability. Furthermore, relying on fixed charges, such an approach could potentially struggle with polarizations along conjugated bonds such as in the acenes discussed in section 2.3 above.

Additionally, we point out that these studies did not include any correction to the polarization energy terms, which are conceptionally more involved (see section 1.1.1) to accurately estimate and typically would require even more sophisticated parameter training. In ACKS2, on the other hand, density response and polarization energy can be computed from the same set of matrices. Finally, given that ACKS2 is parametrized directly from DFT, currently without any empirical fitting, the accuracy of the model rests on a single parameter, the quality of the basis set.

3 Discussion

In conclusion, we presented a highly efficient Gaussian basis set implementation of the ACKS2 method, that is transferable from diatomics to general molecules and that shows good accuracy when compared to reference DFT calculations. A new minimal GF basis set has been developed for carbon and hydrogen species and consequent numerical validation illustrated the power of the ACKS2 method despite the rather simplistic training procedure.

Employing the new Gaussian basis set for ACKS2 responses, we found it to yield admirable accuracies compared to DFT reference calculations. The electronic response prop-

erties of small molecules such as alkanes, unsaturated cyclic hydrocarbons, as well as single, double, and triple bonded compounds demonstrated the accuracy of ACKS2 for a large number of bonding perturbation scenarios. As observables, we thereby calculated induced dipole moments and presented the first evaluation of the polarization energy within the ACKS2 formalism, which matches DFT energies to a remarkable degree. Although one-dimensional cuts through the ACKS2 density and potential response show great differences to their DFT counterparts, the observables nevertheless agree well due to error cancellations when integrated over all space. Finally, we demonstrated the good performance of ACKS2 on application-relevant molecules in organic semiconductor research, tetracene and anthracene.

Future work will focus on the improvement of basis sets, extension to further compounds, and in particular the development of strategies for electronic structure free incorporation of ACKS2 polarization energy in force field based molecular dynamics simulation. Due to the Gaussian basis introduced in this work, most of the ACKS2 parameters can be determined analytically, yet a few of them still rely on a full electronic structure evaluation. In the current state, the exchange-correlation contributions to the hardness η and the KS non-interacting linear response kernel χ are evaluated from the KS orbitals of the reference system. For an application of the ACKS2 model in a force field this computational bottleneck needs to be removed. We envision the elimination of the computationally expensive DFT parametrization process in a second step by geometric scaling or interpolation schemes, facilitated by the smoothness of the ACKS2 parameters with respect to nuclear displacements, to finally obtain linear electronic responses that only depend on nuclear geometry, and hence are ideally suited for force fields. In such a framework, conventional inter-atomic potentials would account for the intramolecular interactions, permanent electrostatic interactions and dispersion corrections, while the ACKS2 model provides static electronic polarization.

4 Methods

Throughout this study, DFT reference calculations as well as the KS electronic structure dependent ACKS2 parameter evaluations were carried out with the FHI-aims full potential all electron DFT simulation package.⁴⁹ We applied the PBE generalized gradient approximated density functional. Integrations were conducted using “tight” integration grids with wave functions expanded in a *tier 3* numeric atomic orbital basis, to ensure the numerical convergence of our results.

While PBE certainly is not the perfect functional for both, the C₂ dimer used in the basis set parametrization, as well as the polarizability of acenes, the exact choice of functional is not relevant. As shown earlier,⁴⁹ even an imperfect description of the electronic structure of elementary dimers seems sufficient to yield accurate and transferable basis sets (cf. section 2.1). On the other hand, in our accuracy tests, ACKS2 shows itself able to very well reproduce the DFT reference. We have no reason to suspect this not to hold for other, modern DFT functionals such as the promising SCAN meta-GGA,⁵⁰ that can be expected to yield better polarization properties of molecules.

5 Contributions

L.L. initially implemented the Gaussian-based ACKS2 method. P.G. refined and extended this implementation. Furthermore, P.G. implemented the ACKS2 energy terms and performed all training and testing calculations. The initial idea to the project was conceived by H.O. and K.R. H.O. and J.B. derived the ACKS2 energy expressions, designed and supervised the project. All authors contributed to the writing of the manuscript.

6 Competing Interests

The authors declare no competing interests.

7 Data Availability

Training data for the Gaussian parametrization as well as raw data of the test-sets are available from the corresponding author upon reasonable request.

8 Code Availability

All software used in this work is present as an in-house proof-of-concept Python package to be made available upon reasonable request.

Acknowledgement

The authors gratefully gratefully acknowledge support from the Solar Technologies Go Hybrid Initiative of the State of Bavaria. P.G. further acknowledges the support of the Technische Universität München – Institute for Advanced Study, funded by the German Excellence Initiative (and the European Union Seventh Framework Programme under Grant Agreement №291763). H.O. acknowledges support from the German Research Foundation (DFG), priority program 1928 COORNETS, Grant OB425/3-1. J.B. acknowledges TUM-IAS for the award of a generous Hans-Fischer Fellowship.

Supporting Information Available

Theoretical foundations of the ACKS2 approach and technical details of the evaluation of the new Gaussian basis can be found in the supplementary material. It also contains a description of the training and test-sets used in this work, as well as the description of how to extract the reference data from DFT calculations. This information is available free of charge via the Internet at <http://pubs.acs.org>

References

- (1) Forrest, S. R. The Path to Ubiquitous and Low-Cost Organic Electronic Appliances on Plastic. *Nature* **2004**, *428*, 911–918.
- (2) Brédas, J.-L.; Norton, J. E.; Cornil, J.; Coropceanu, V. Molecular understanding of organic solar cells: the challenges. *Acc. Chem. Res.* **2009**, *42*, 1691–1699.
- (3) Wang, L.; Nan, G.; Yang, X.; Peng, Q.; Li, Q.; Shuai, Z. Computational methods for design of organic materials with high charge mobility. *Chem. Soc. Rev.* **2010**, *39*, 423–434.
- (4) Minemawari, H.; Yamada, T.; Matsui, H.; Tsutsumi, J.; Haas, S.; Chiba, R.; Kumai, R.; Hasegawa, T. Inkjet Printing of Single-Crystal Films. *Nature* **2011**, *475*, 364–367.
- (5) Stavrinidou, E.; Gabrielsson, R.; Gomez, E.; Crispin, X.; Nilsson, O.; Simon, D. T.; Berggren, M. Electronic Plants. *Sci. Adv.* **2015**, *1*, e1501136.
- (6) Coropceanu, V.; Cornil, J.; da Silva Filho, D. A.; Olivier, Y.; Silbey, R.; Brédas, J.-L. Charge Transport in Organic Semiconductors. *Chem. Rev.* **2007**, *107*, 926–952, PMID: 17378615.
- (7) Jailaubekov, A. E.; Willard, A. P.; Tritsch, J. R.; Chan, W.-L.; Sai, N.; Gearba, R.; Kaake, L. G.; Williams, K. J.; Leung, K.; Rossky, P. J.; Zhu, X.-Y. Hot charge-transfer excitons set the time limit for charge separation at donor/acceptor interfaces in organic photovoltaics. *Nat. Mater.* **2012**, *12*, 66 EP–.
- (8) Oberhofer, H.; Reuter, K.; Blumberger, J. Charge Transport in Molecular Materials: An Assessment of Computational Methods. *Chem. Rev.* **2017**, *117*, 10319–10357, PMID: 28644623.
- (9) Nelson, J.; Kwiatkowski, J. J.; Kirkpatrick, J.; Frost, J. M. Modeling Charge Trans-

- port in Organic Photovoltaic Materials. *Acc. Chem. Res.* **2009**, *42*, 1768–1778, PMID: 19848409.
- (10) Troisi, A. Charge transport in high mobility molecular semiconductors: classical models and new theories. *Chem. Soc. Rev.* **2011**, *40*, 2347–2358.
 - (11) Giannini, S.; Carof, A.; Blumberger, J. Crossover from Hopping to Band-Like Charge Transport in an Organic Semiconductor Model: Atomistic Nonadiabatic Molecular Dynamics Simulation. *J. Phys. Chem. Lett.* **2018**, *9*, 3116–3123, PMID: 29787275.
 - (12) Spencer, J.; Gajdos, F.; Blumberger, J. FOB-SH: Fragment orbital-based surface hopping for charge carrier transport in organic and biological molecules and materials. *J. Chem. Phys.* **2016**, *145*, 064102.
 - (13) Rühle, V.; Lukyanov, A.; May, F.; Schrader, M.; Vehoff, T.; Kirkpatrick, J.; Baumeier, B.; Andrienko, D. Microscopic Simulations of Charge Transport in Disordered Organic Semiconductors. *J. Chem. Theor. Comput.* **2011**, *7*, 3335–3345, PMID: 22076120.
 - (14) Rappe, A. K.; Casewit, C. J.; Colwell, K. S.; Goddard, W. A.; Skiff, W. M. UFF, a full periodic table force field for molecular mechanics and molecular dynamics simulations. *J. Am. Chem. Soc.* **1992**, *114*, 10024–10035.
 - (15) Ponder, J. W.; Wu, C.; Ren, P.; Pande, V. S.; Chodera, J. D.; Schnieders, M. J.; Haque, I.; Mobley, D. L.; Lambrecht, D. S.; DiStasio, R. A.; Head-Gordon, M.; Clark, G. N. I.; Johnson, M. E.; Head-Gordon, T. Current Status of the AMOEBA Polarizable Force Field. *J. Phys. Chem. B* **2010**, *114*, 2549–2564, PMID: 20136072.
 - (16) Senftle, T. P.; Hong, S.; Islam, M. M.; Kylasa, S. B.; Zheng, Y.; Shin, Y. K.; Junkermeier, C.; Engel-Herbert, R.; Janik, M. J.; Aktulga, H. M.; Verstraelen, T.; Grama, A.; van Duin, A. C. T. The ReaxFF reactive force-field: development, applications and future directions. *Npj Comput. Mater.* **2016**, *2*, 15011 EP–.

- (17) Xu, P.; Guidez, E. B.; Bertoni, C.; Gordon, M. S. Perspective: Ab initio force field methods derived from quantum mechanics. *J. Chem. Phys.* **2018**, *148*, 090901.
- (18) Rahman, A.; Stillinger, F. H. Molecular Dynamics Study of Liquid Water. *J. Chem. Phys.* **1971**, *55*, 3336–3359.
- (19) Brooks, B. R.; Brucoleri, R. E.; Olafson, B. D.; States, D. J.; Swaminathan, S.; Karplus, M. CHARMM: A program for macromolecular energy, minimization, and dynamics calculations. *J. Comput. Chem.* **1983**, *4*, 187–217.
- (20) Haughney, M.; Ferrario, M.; McDonald, I. R. Molecular-dynamics simulation of liquid methanol. *J. Phys. Chem.* **1987**, *91*, 4934–4940.
- (21) Cornell, W. D.; Cieplak, P.; Bayly, C. I.; Gould, I. R.; Merz, K. M.; Ferguson, D. M.; Spellmeyer, D. C.; Fox, T.; Caldwell, J. W.; Kollman, P. A. A Second Generation Force Field for the Simulation of Proteins, Nucleic Acids, and Organic Molecules. *J. Am. Chem. Soc.* **1995**, *117*, 5179–5197.
- (22) Scott, W. R. P.; Hünenberger, P. H.; Tironi, I. G.; Mark, A. E.; Billeter, S. R.; Fenn, J.; Torda, A. E.; Huber, T.; Krüger, P.; van Gunsteren, W. F. The GROMOS Biomolecular Simulation Program Package. *J. Phys. Chem. A* **1999**, *103*, 3596–3607.
- (23) Wang, H.; Yang, W. Determining polarizable force fields with electrostatic potentials from quantum mechanical linear response theory. *J. Chem. Phys.* **2016**, *144*, 224107.
- (24) Nyman, J.; Pundyke, O. S.; Day, G. M. Accurate force fields and methods for modelling organic molecular crystals at finite temperatures. *Phys. Chem. Chem. Phys.* **2016**, *18*, 15828–15837.
- (25) Cieplak, P.; Dupradeau, F.-Y.; Duan, Y.; Wang, J. Polarization effects in molecular mechanical force fields. *J. Phys. Condens. Matter* **2009**, *21*, 333102.

- (26) Mortier, W. J.; Van Genechten, K.; Gasteiger, J. Electronegativity equalization: application and parametrization. *J. Am. Chem. Soc.* **1985**, *107*, 829–835.
- (27) Mortier, W. J.; Ghosh, S. K.; Shankar, S. Electronegativity-equalization method for the calculation of atomic charges in molecules. *J. Am. Chem. Soc.* **1986**, *108*, 4315–4320.
- (28) York, D. M.; Yang, W. A chemical potential equalization method for molecular simulations. *J. Chem. Phys.* **1996**, *104*, 159–172.
- (29) Chelli, R.; Procacci, P. A transferable polarizable electrostatic force field for molecular mechanics based on the chemical potential equalization principle. *J. Chem. Phys.* **2002**, *117*, 9175–9189.
- (30) Rappe, A. K.; Goddard, W. A. Charge equilibration for molecular dynamics simulations. *J. Phys. Chem.* **1991**, *95*, 3358–3363.
- (31) Chelli, R.; Procacci, P.; Righini, R.; Califano, S. Electrical response in chemical potential equalization schemes. *J. Chem. Phys.* **1999**, *111*, 8569–8575.
- (32) Nistor, R. A.; Polihronov, J. G.; Müser, M. H.; Mosey, N. J. A generalization of the charge equilibration method for nonmetallic materials. *J. Chem. Phys.* **2006**, *125*, 094108.
- (33) Lee Warren, G.; Davis, J. E.; Patel, S. Origin and control of superlinear polarizability scaling in chemical potential equalization methods. *J. Chem. Phys.* **2008**, *128*, 144110.
- (34) Verstraelen, T.; Ayers, P. W.; Speybroeck, V. V.; Waroquier, M. ACKS2: Atom-condensed Kohn-Sham DFT approximated to second order. *J. Chem. Phys.* **2013**, *138*, 074108.
- (35) Verstraelen, T.; Vandenbrande, S.; Ayers, P. W. Direct computation of parameters for accurate polarizable force fields. *J. Chem. Phys.* **2014**, *141*, 194114.

- (36) Misquitta, A. J.; Stone, A. J. Distributed polarizabilities obtained using a constrained density-fitting algorithm. *J. Chem. Phys.* **2006**, *124*, 024111.
- (37) Misquitta, A. J.; Stone, A. J. Accurate induction energies for small organic molecules: 1. Theory. *J. Chem. Theor. Comput.* **2008**, *4*, 7–18.
- (38) Islam, M. M.; Kolesov, G.; Verstraelen, T.; Kaxiras, E.; van Duin, A. C. T. eReaxFF: A Pseudoclassical Treatment of Explicit Electrons within Reactive Force Field Simulations. *J. Chem. Theor. Comput.* **2016**, *12*, 3463–3472, PMID: 27399177.
- (39) Baroni, S.; Giannozzi, P.; Testa, A. Green’s-function approach to linear response in solids. *Phys. Rev. Lett.* **1987**, *58*, 1861.
- (40) Gonze, X. Adiabatic density-functional perturbation theory. *Phys. Rev. A* **1995**, *52*, 1096.
- (41) Giese, T. J.; York, D. M. Density-functional expansion methods: Evaluation of LDA, GGA, and meta-GGA functionals and different integral approximations. *J. Chem. Phys.* **2010**, *133*, 244107.
- (42) Giese, T. J.; York, D. M. Density-functional expansion methods: grand challenges. *Theor. Chem. Acc.* **2012**, *131*, 1145.
- (43) Garrett, A. inspyred: Bio-inspired Algorithms in Python. 2017; <http://aarongarrett.github.io/inspyred/>.
- (44) Anthony, J. E. The Larger Acenes: Versatile Organic Semiconductors. *Angew. Chem. Int. Ed.* **2008**, *47*, 452–483.
- (45) Kaminski, S.; Giese, T. J.; Gaus, M.; York, D. M.; Elstner, M. Extended Polarization in Third-Order SCC-DFTB from Chemical-Potential Equalization. *J. Phys. Chem. A* **2012**, *116*, 9131–9141.

- (46) Kaminski, S.; Gaus, M.; Elstner, M. Improved Electronic Properties from Third-Order SCC-DFTB with Cost Efficient Post-SCF Extensions. *J. Phys. Chem. A* **2012**, *116*, 11927–11937.
- (47) Giese, T. J.; York, D. M. Improvement of semiempirical response properties with charge-dependent response density. *J. Chem. Phys.* **2005**, *123*, 164108.
- (48) Baker, C. M.; Anisimov, V. M.; MacKerell Jr, A. D. Development of CHARMM polarizable force field for nucleic acid bases based on the classical Drude oscillator model. *J. Phys. Chem. B* **2010**, *115*, 580–596.
- (49) Blum, V.; Gehrke, R.; Hanke, F.; Havu, P.; Havu, V.; Ren, X.; Reuter, K.; Scheffler, M. Ab initio molecular simulations with numeric atom-centered orbitals. *Comp. Phys. Commun.* **2009**, *180*, 2175–2196.
- (50) Sun, J.; Ruzsinszky, A.; Perdew, J. P. Strongly Constrained and Appropriately Normed Semilocal Density Functional. *Phys. Rev. Lett.* **2015**, *115*, 036402.

Table 2: Linear response prediction for HEF and PC perturbations of hydrocarbon test set. The relative errors of the ACKS2 polarization energy ($\text{RME}(\Delta E^{\text{pol.}})$ [%]) and induced dipole moment ($\text{RME}(\Delta\mu)$ [%]) compared to the DFT reference are listed.

alkanes	$\text{RME}(\Delta E^{\text{pol.}})$	$\text{RME}(\Delta\mu)$
methane	10.2	10.5
ethane	8.5	9.0
propane	10.9	9.9
n-butane	11.9	10.7
$\text{C}_2\text{H}_{\{2,4,6\}}$		
ethane	8.5	9.0
ethylene	13.1	11.3
acetylene	21.4	21.8
cyclic		
cyclopropanene	12.4	11.2
cyclobutadiene	16.4	14.6
cyclopentadiene	14.8	13.2
benzene	18.4	16.4

Figure 7: Illustration of pairwise distance dependency of KS-DFT derived ACKS2 parameters for atomic s-type Gaussian basis functions for breathing mode of benzene.

(a) Benzene molecular motion and relative position of carbon atoms. The pairwise interatomic distances are applied as metric for the ACKS2 parameters following the color coding scheme.

(b) Off-center KS-DFT parameters for symmetric breathing mode.

(c) Same-center KS-DFT parameters for symmetric breathing mode.

Graphical TOC Entry