

Table 2: Adjusted analysis of renal impairment predictors, full cohort

Variable	Crude Odds Ratio [95%CI] of eGFR ≤ 60 mL/min/1.73m²	Adjusted* Odds Ratio [95%CI] of eGFR ≤ 60 mL/min/1.73m²
Age (by 10 years older)	4.6 [3.5-6.0]	5.4 [3.9-7.5]
Sex (female vs. male)	2.9 [1.6-5.2]	5.0 [2.6-9.8]
Ancestry**	1.6 [0.7-3.8]	1.4 [0.5-3.7]
HIV status (yes vs. no)	2.4 [1.4-4.1]	3.4 [1.8-6.4]
BMI (kg/m ²) (by 1 increase in BMI)	1.0 [1.0-1.1]	1.0 [1.0-1.1]
Pack years (by additional 10)	1.0 [1.0-1.2]	0.9 [0.8-1.0]
Diabetes (yes vs. no)	4.8 [2.4-9.7]	2.9 [1.2-6.7]
Hypertension (yes vs. no)	2.7 [1.5-4.8]	0.9 [0.5-1.9]
Educational level (REF: No		
– Short (<3 years)	0.9 [0.2-3.1]	0.6 [0.1-2.9]
– Vocational	0.6 [0.2-1.8]	1.0 [0.3-3.4]
– Middle length	0.8 [0.3-2.2]	0.8 [0.2-2.6]
– University degree	0.6 [0.2-1.8]	1.3 [0.4-4.8]

* Adjusted for age, sex, HIV, BMI, pack years, diabetes and hypertension

** Other Scandinavian vs Danish