Is there a relationship between saccadic eye movements and cognitive function in multiple sclerosis?

J.A. Nij Bijvank^{1,2}, A. Eijlers³, L.J. Balk², H.S. Tan¹, B.M.J Uitdehaag², A. Petzold^{1,2,4}, L.J. van Rijn¹

¹Department of ophthalmology, VU University Medical Center, Amsterdam, The Netherlands ²Neurology, Amsterdam neuroscience, VUmc MS Center Amsterdam, The Netherlands ³Anatomy & Neurosciences, Amsterdam neuroscience, VU University Medical Center, Amsterdam, The Netherlands ⁴Moorfields Eye Hospital, London, United Kingdom



> Consider precise quantitative saccadic testing as a potential surrogate outcome for cognitive function in longitudinal studies and treatment trials

- Rao SM, Leo GJ, Bernardin L, Unverzagt F. Cognitive dysfunction in multiple sclerosis. I. Frequency, patterns and prediction. Neurol. 1991; 41:685-691 6. Amato MP, Zipoli V, Portaccio E. Multiple sclerosis-related cognitive changes: a review of cross-sectional and longitudinal studies. J. Neurol. Sci. 2006; 245:41-46

Disclosures (none related to this work)

JANB, AE, HST, LJR: no disclosures - LJB: The VUmc MS Centre received research support from TEVA - BMJU: has received consultancy fees from Biogen Idec, Genzyme, Merck Serono, Novartis, Roche and Teva - AP: member of the steering committee for the OCTiMS study (Novartis), no consulting fees. Performs OCT QC for the Passos study (Novartis), receives consulting fees.





E-mail: j.nijbijvank@vumc.nl

Amsterdam

Neuroscience