

**IRF2BPL mutations cause autosomal dominant dystonia
with anarthria, slow saccades and seizures**

Christos Ganos¹, Simone Zittel², Ute Hidding²,

Claudia Funke³, Saskia Biskup³ and Kailash P. Bhatia⁴

1Department of Neurology, Charité University Medicine Berlin, Berlin, Germany

2Department of Neurology, University Medical Center Hamburg-Eppendorf, Hamburg, Germany.

3Praxis für Humangenetik Tübingen, Paul-Ehrlich-Str. 23, 72076 Tuebingen

4Department of Clinical and Movement Neurosciences, Queen Square Institute of Neurology,
University College London, London, United Kingdom

Corresponding Author

Dr. med. Christos Ganos

Movement Disorders and Body Control Lab

Movement Disorders and Neuromodulation Unit

Department of Neurology, Charité University Medicine Berlin

Charitéplatz 1, 10117 Berlin, Germany

eMail: christos.ganos@charite.de

Word count

563

Running title

Complex dystonia with anarthria/aphonia due to IRF2BPL mutations

Key words

Dystonia, anarthria, aphonia, IRF2BPL, seizures

Conflict of Interest

The authors disclose no conflicts of interest regarding this manuscript.

Funding sources

This research project was supported by a grant from the VolkswagenStiftung (Freigeist) held by CG.

In the ever-expanding spectrum of genetic dystonia syndromes, the presence of associated clinical signs can provide useful clues to guide diagnostic reasoning and inform treatment approaches. We have previously delineated the range of known etiologies associated with dystonia and anarthria/aphonia, and provided an algorithmic approach to reach diagnosis¹. Since then, mutations in the *KMT2B* gene have been added to this list and indeed, two of the patients we previously reported (cases 15 and 20)¹ were subsequently found to harbor pathogenic mutations in this gene. Here, we wish to highlight a novel genetic etiology, namely mutations in the *IRF2BPL* gene, recently reported to cause a broad phenotypic range of neurological syndromes²⁻⁴, with a particular focus on the syndromic association of dystonia with anarthria/aphonia.

In 2013 we described an autosomal dominant dystonic syndrome in a female patient (index) and her son (figure 1A), with a characteristic phenotypic presentation consisting of leg-onset generalized dystonia, severe anarthria/aphonia, slow saccades and epilepsy, including photic myoclonus (video; with informed written consent from all individuals for online publication and dissemination of the videos; additional clinical information is to be found in original description)⁵. Thorough clinical, paraclinical and genetic investigations at the time, including exome sequencing had been unrevealing⁵. Brain MRI of the index patient revealed mild supratentorial and cerebellar brain atrophy (figure 1B) and neurophysiology showed axonal and demyelinating neuropathy of the peroneal and sural nerves. Due to newly performed trio-based exome analyses of the affected mother, the affected son and the unaffected brother of the index patient we filtered all common variants which were unique only in the affected family members. With this approach we identified a heterozygous nonsense variant (c.355C>T; p.Gln119*) within the gene *IRF2BPL*, which was confirmed by Sanger sequencing (figure 1C).

IRF2BPL mutations were recently reported to cause a range of neurological syndromes, from progressive childhood-onset motor regression with severe speech abnormalities, including anarthria/aphonia, and photosensitive seizures, over to developmental epileptic encephalopathy²⁻⁴. Dystonia was reported in 9 cases and oculomotor abnormalities, including (horizontal) saccadic speed slowing, were also documented²⁻⁴. Onset of dystonia was variable. Indeed, in the 4 out of 7 cases

reported by Macrogliese et al. dystonic symptoms developed between childhood and adolescence ². Remarkably, in one further patient symptoms of neurological degeneration were first noted at 17 years of age⁴.

The cases we present here, fit well within the phenotypic presentation of *IRF2BPL* mutation syndrome, but also provide important new aspects. First, all previous reported patients, where genetic evaluation of parents was available, harbored *de novo* mutations. In contrast, our cases show that *IRF2BPL* mutations could also manifest later in life and be, thereby, transmitted in an autosomal dominant manner. Importantly, they also reveal marked variability in the onset of symptoms (23 years for index vs early childhood for her son). Second, we believe that mutations at the *IRF2BPL* gene should be – to date – considered pathognomonic for the remarkably rare syndromic association of dystonia with anarthria/aphonia, slow (horizontal) saccades and seizures. Moreover, our patients had certain features, such as keratoconus (male) and gingival enlargement (index), which may prove to be further useful diagnostic clues. Third, we note the lack of treatment response to pallidal deep brain stimulation in the male patient.

To conclude, *IRF2BPL* mutations should be added to the long list of conditions that cause dystonia with anarthria/aphonia, particularly in cases where slow horizontal saccades and epileptic seizures, including photic myoclonus, are present.

Financial Disclosure

CG holds research grants from the VolkswagenStiftung (Freigeist Fellowship) and the German Parkinson Society and was also supported by the Deutsche Forschungsgemeinschaft (DFG; GA2031/1-1 and GA2031/1-2).

SZ holds a research grant from Werner Otto Stiftung and has received research support from Merz.

UH reports no disclosures

CF reports no disclosures

SB reports no disclosures.

KPB holds research grants from NIHR RfPB, MRC Wellcome Strategic grant (WT089698) and PD UK (G-1009), and has received honoraria/financial support to speak/attend meetings from GSK, Boehringer-Ingelheim, Ipsen, Merz, Sun Pharma, Allergan, Teva Lundbeck and Orion pharmaceutical companies. K.B. receives royalties from Oxford University press and a stipend for MDCP editorship.

Reference:

1. Ganos C, Crowe B, Stamelou M, et al. The clinical syndrome of dystonia with anarthria/aphonia. *Parkinsonism Relat Disord* 2016;24:20-27.
2. Marcogliese PC, Shashi V, Spillmann RC, et al. IRF2BPL Is Associated with Neurological Phenotypes. *Am J Hum Genet* 2018;103:245-260.
3. Skorvanek M, Dusek P, Rydzanicz M, et al. Neurodevelopmental disorder associated with IRF2BPL gene mutation: Expanding the phenotype? *Parkinsonism Relat Disord* 2019.
4. Tran Mau-Them F, Guibaud L, Duplomb L, et al. De novo truncating variants in the intronless IRF2BPL are responsible for developmental epileptic encephalopathy. *Genet Med* 2019;21:1008-1014.
5. Ganos C, Biskup S, Kruger S, et al. Dystonia with aphonia, slow horizontal saccades, epilepsy and photic myoclonus: a novel syndrome? *Parkinsonism Relat Disord* 2014;20:328-331.

Figure 1. **A.** Pedigree of the reported family (previously published and used with permission). **B.** Midsagittal plane of brain MRI of the index patient (II.2) at the age of 56 years demonstrating cerebellar atrophy. **C.** NGS (upper lane: family members; lower lane: wildtype references) and Sanger data of the region of interest of the *IRF2BPL* gene

Video: Segment A demonstrates index patient at the age of 56 with generalized dystonia, including, jaw-opening dystonia, profound saccadic slowing, particularly in the horizontal plane, anarthria and photic myoclonus. No evidence of cerebellar ataxia in finger-to-nose testing. Gingiva hyperplasia, as well as distal muscle atrophy of both legs without evidence of spasticity are noted. Segment B shows the same patient at the age of 61 years with now spontaneous myoclonic jerks during examination and severe atrophy of distal leg muscles, with accompanying color and temperature changes of the skin. Segment C shows son of index patient at the age of 27 years with generalized dystonia, including jaw-opening dystonia, aphonia, and markedly slowed horizontal saccades. No evidence of cerebellar ataxia was noted (Segments A and C previously published¹).