Geophysical Research Letters

RESEARCH LETTER

10.1029/2019GL086650

Key Points:

- The Dongqiao Formation was deposited on top of ophiolite in subaerial to shallow marine environments during Oxfordian to Kimmeridgian time
- The Dongqiao Formation includes detritus from the underlying Dongqiao ophiolite and Lhasa-affinity continental crust
- The Dongqiao ophiolite was obducted onto Lhasa-affinity continental crust by 163 Ma

Supporting Information:

- Supporting Information S1
- Data Set S1

Correspondence to:

X. Hu,
huxm@nju.edu.cn

Citation:

Ma, A., Hu, X., Kapp, P., BouDagher-Fadel, M., & Lai, W. (2020). Pre-Oxfordian (>163 Ma) ophiolite obduction in Central Tibet. *Geophysical Research Letters*, 47, e2019GL086650. <https://doi.org/10.1029/2019GL086650>

Received 16 DEC 2019

Accepted 11 APR 2020

Accepted article online 17 APR 2020

Pre-Oxfordian (>163 Ma) Ophiolite Obduction in Central Tibet

Anlin Ma¹, Xiumian Hu¹ , Paul Kapp² , Marcelle BouDagher-Fadel³, and Wen Lai¹

¹State Key Laboratory of Mineral Deposit Research, School of Earth Sciences and Engineering, Nanjing University, Nanjing, China, ²Department of Geosciences, University of Arizona, Tucson, Arizona, USA, ³Department of Earth Sciences, University College London, London, UK

Abstract The timing of Bangong-Nujiang suture ophiolite obduction between the Lhasa and Qiangtang terranes in central Tibet is important for understanding the closure history of the Meso-Tethys but remains poorly constrained. We investigated subaerial to shallow marine strata of the Dongqiao Formation that sit unconformably on Bangong-Nujiang suture ophiolites that crystallized in a supra-subduction zone setting at 189–181 Ma. Based on foraminiferal and coral studies, the depositional age of the Dongqiao Formation is constrained to be Oxfordian and Kimmeridgian (Late Jurassic). Provenance analyses including detrital modes, geochemistry of detrital chromian spinels, and U-Pb age populations of detrital zircons suggest the Dongqiao Formation was sourced from uplifted Bangong-Nujiang suture ophiolites and sedimentary and metamorphic rocks of Lhasa terrane affinity to the south. We conclude that Bangong-Nujiang suture ophiolites were obducted soon after crystallization (prior to Oxfordian time; >163 Ma) onto the Lhasa terrane or a microcontinent of Lhasa terrane affinity.

Plain Language Summary Ophiolite obduction often occurs when a passive continental margin enters an oceanic subduction zone and thus may mark when an arc-continent or continent-continent collision begins. In central Tibet, the Dongqiao ophiolite represents a relict of Meso-Tethys oceanic lithosphere. Initial research during the 1980s provided rough constraints on the timing, polarity, and mechanism of Dongqiao ophiolite obduction. However, there has been little advancement in our knowledge of the obduction history since. Here we report results of sedimentologic, stratigraphic, and provenance studies on the Dongqiao Formation overlying the Dongqiao ophiolite. Our foraminiferal and coral biostratigraphic data show that subaerial to shallow marine strata of the Dongqiao Formation were deposited between 163 and 152 million years ago. We demonstrate that the clastic rocks in the Dongqiao Formation received detritus from both the underlying Dongqiao ophiolite and Lhasa terrane affinity continental crust. We propose that a continental margin of Lhasa terrane affinity entered a north-dipping oceanic trench, and the Dongqiao ophiolite was obducted southwards on to it, no later than 163 million years ago.

1. Introduction

It is important in plate tectonics to understand why and how dense submarine oceanic lithosphere rocks were thrust (obducted) on top of less dense continental crust to form ophiolites and cause orogeny (Dewey, 1976; Hacker et al., 1996). It has been proposed that ophiolite obduction may be triggered by various tectonic events including backarc basin closure, accretionary prism underthrusting, ridge-trench collision, and arc-continent or continent-continent collision, among others (Agard et al., 2011; Alavi, 1994; Coleman, 1981; Dewey & Casey, 2011; Wakabayashi & Dilek, 2003).

In central Tibet, Late Mesozoic convergence between the Lhasa and Qiangtang terranes resulted in the consumption of Bangong-Nujiang (Meso-Tethys) oceanic lithosphere, a transition from marine to nonmarine deposition, and crustal thickening (Kapp et al., 2007; Ma et al., 2018; Murphy et al., 1997; Raterman et al., 2014). However, the timing of initial Lhasa-Qiangtang collision is disputed, with estimates ranging from Middle Jurassic to Late Cretaceous time (see a review by Li, Yin, et al., 2019). Other fundamental issues such as when Bangong-Nujiang suture ophiolites were obducted, and whether ophiolite obduction marks the initiation of intercontinental Lhasa-Qiangtang collision also remain strongly debated (e.g., Girardeau et al., 1984; Kapp et al., 2003; Li, Guilmette, et al., 2019).

Figure 1. (a) Digital elevation model showing tectonic division of the Tibetan Plateau, ophiolite distribution in the Bangong-Nujiang suture zone, and the Xainzha ophiolite within the northern Lhasa terrane. (b) Geological map of the Dongqiao area modified from the 1:250,000 scale geological maps of the Zigetangco, Bangoin, and Amdo counties (ITGS, 2002, 2003, 2005). Black dashed lines outline possible suture zones bounding a micro-continent. References for ophiolite ages are provided in supplementary Table S1. YSZ, Yarlung-Tsangpo suture zone; BNSZ, Bangong-Nujiang suture zone; JSZ, Jinsha suture zone; QT, Qiangtang; L, Lhasa.

The subaerial to shallow marine Dongqiao Formation (same as the Zigetang Formation of Girardeau et al., 1984) in the Dongqiao area sits depositionally on ophiolite (Figures 1a and 1b) and thus can place an upper age limit for the timing of ophiolite obduction and shed light on the obduction mechanism. Previous studies only roughly constrained the depositional age of the Dongqiao Formation to be at some time during the Late Jurassic to Early Cretaceous based on first-order biostratigraphic studies (Girardeau et al., 1984; Kidd et al., 1988; Marcoux et al., 1987; Wang & Dong, 1984).

In this study, we used foraminiferal and coral biostratigraphy to constrain the depositional age of the Dongqiao Formation. We also conducted detailed sedimentologic and provenance studies to better constrain the paleogeographic setting of the Dongqiao Formation. We discuss the timing and possible tectonic significance of Dongqiao ophiolite obduction in light of our results.

2. Geological Background

The Tibetan Plateau formed as the result of accretion of the Qiangtang, Lhasa, and Indian continental terranes to the southern continental margin of Asia, successively from north to south during Triassic to Cenozoic time (e.g., Dewey et al., 1988; Yin & Harrison, 2000). These terranes are bounded by the Jinsha, Bangong-Nujiang, and Yarlung-Tsangpo sutures which mark the closure of Paleo-, Meso- and Neo-Tethys oceans, respectively (Figure 1a). Relevant to this study are the Qiangtang terrane, Amdo basement, Bangong-Nujiang suture zone, and Lhasa terrane (Figure 1a). The Qiangtang terrane is covered in many places by Triassic to Jurassic shallow-marine to littoral limestones and siliciclastic rocks (Figure 1b; Ma et al., 2017, 2018). The Amdo basement is structurally bounded within the Bangong-Nujiang suture zone and consists of 915–840 Ma and 530–470 Ma orthogneisses and metasedimentary rocks of Qiangtang affinity intruded by 185–170 Ma calc-alkaline granitoids (Figure 1b); it is inferred to have collided with the Qiangtang terrane at 170–165 Ma (Guynn et al., 2006; Kapp & DeCelles, 2019).

Ophiolites in the Bangong-Nujiang suture zone yield mainly Triassic to Jurassic crystallization ages based on zircon U-Pb geochronology (see Figure 1 for location and supplementary Table S1 for compiled data).

Gabbro in the Dongqiao and Jiangco ophiolites crystallized between 189 and 181 Ma (Table S1). Farther south (~70 km), the Baila ophiolite has yielded crystallization ages of 172–164 Ma (Tang et al., 2018), 244 Ma, and 149–148 Ma (Zhong et al., 2017). At the base of the Dongqiao ophiolite is a ~8-m-thick amphibolite- to greenschist-facies metamorphic sole that yielded two $^{40}\text{Ar}/^{39}\text{Ar}$ hornblende dates of ~180 and ~175 Ma (Zhou et al., 1997). The Bangong-Nujiang suture ophiolites in the Dongqiao area are unconformably overlain by the Dongqiao Formation, which includes conglomerate in its lower part and sandstone and limestone up-section (Girardeau et al., 1984). Corals, algae, foraminifers and bivalves in the Dongqiao Formation broadly suggest a Late Jurassic–Early Cretaceous age (Girardeau et al., 1984; Marcoux et al., 1987; Wang & Dong, 1984). Late Jurassic rocks exposed between Amdo and Dongqiao include volcanic rocks, clastic rocks, and limestones that were deposited mostly in shallow marine environments (Figure 1b; ITGS, 2002, 2005).

Triassic to Jurassic deep-marine turbidite-bearing sandstones and shales within the suture zone and possibly on the northern Lhasa terrane, mapped as Jurassic flysch by Girardeau et al. (1984), include the Mugangangri, Quehala, Xihu, and Jienu groups (Figure 1b; ITGS, 2002). The Mugangangri and Jienu groups have a provenance of Qiangtang terrane affinity (Li et al., 2020). Paleozoic rocks between Dongqiao and Baila are assigned the same lithostratigraphic units as those in the Lhasa terrane and are mainly composed of limestone with subordinate marble, dolomite, and schist (ITGS, 2002). The structural geology between Dongqiao and Baila is very complex, but previous mapping shows that ophiolitic rocks and Paleozoic strata structurally overlie (i.e., are thrust over) the Mesozoic turbiditic strata (redrafted cross-sections of Girardeau et al., 1984 are provided in Figure S1). Intermediate and felsic volcanic rocks and granitoids of 166–160 Ma are also locally exposed between Dongqiao and Baila (Li et al., 2020).

3. Stratigraphy and Sedimentology

The Dongqiao Formation in the Zigetangco section is >117 m thick (Figure 2a). It dips gently to the south and overlies the Dongqiao ophiolite serpentinite (Figure 2h). The section can be divided into two parts. The lower part consists of >28 m of red conglomerate and sandstone, followed by 8 m of opaque Fe-oxide layers interbedded with gray micaceous muddy siltstones and sandstones (Figure 2h). The conglomerate is clast-supported, massive or horizontally stratified, and poorly sorted, with the largest clasts being of boulder size (Figure 2i). Gravels have been strongly silicified such that primary clast compositions are difficult to identify, though chromite clasts were previously reported (Girardeau et al., 1984; Leeder et al., 1988). The red sandstone is massive, fine to coarse grained, and composed mainly of detrital spinels (Figures S2a and S2b). In the upper part, gray sandstone, limestone, and mixed carbonate and siliciclastic rocks are interbedded for ~43 m. The gray sandstone at the base of the upper part contains well-rounded quartzite pebbles. Bioclasts are abundant throughout the upper part, including echinoderm, coral, foraminifera, bivalve, brachiopod, and algae. Oncoid rudstone layers and shell beds were also documented. Parallel bedding (Figure 2j) and wood remains are present in the sandstone. Two beds of coral framestone were observed (Figures 2k and 2l). Apart from the Zigetangco section, sandstone was only found in the Pari site, of gray color, while limestone is dominant at most other sites.

Overall, the Zigetangco section shows a transition upward from sub-aerial alluvial to shallow marine facies. The red conglomerate, sandstone, and Fe-oxide rocks in the lower part are interpreted to have been deposited in a proximal alluvial fan along a steep slope, with iron-oxide and silcrete duricrusts related to ferrosiallitic laterite generation as a result of strong chemical weathering of the ophiolite in a humid tropical climate (Girardeau et al., 1984; Leeder et al., 1988). This provides additional evidence that the Dongqiao Formation was deposited unconformably above the ophiolite. The gray sandstone and limestone with abundant marine fossils in the upper part indicate a shallow marine environment close to the shoreline where rivers delivered ophiolite fragments, siliciclastic detritus, and wood remains. The coral framestone represents in-situ patch reefs (Girardeau et al., 1984). Widespread limestone distributes to the north of the Zigetangco section, which implies a proximal to distal (compared to Lhasa-affinity source) paleogeography from south to north. The nature of the contacts between the Dongqiao limestone and Mesozoic turbiditic strata and Qiangtang Mesozoic strata are uncertain.

Figure 2. Measured Zigetangco section (a), foraminifera (b–f) and coral (g) plate, and field photos (h–l) in the Zigetangco section. (b) *Everticyclammina virguliana* (Koechlin), 15DQ01, site b in Figure 1b; (c) *Mesoendothyra croatica* Gušić, 15DQ01; (d) *Pseudocyclammina lituus* (Yokoyama), 15QM02, site c in Figure 1b; (e) (i) *Daxia* sp.; (ii) *Textularia* sp., 15DQ01; (f) *Redmondoides lugeoni* (Septfontaine), 15DQ01; (g) *Cladocoropsis mirabilis* Felix, 17AD25, site c in Figure 1b. Scale bars: 0.3 mm in (b)–(f); 1 cm in (g); (h) the Zigetangco section, showing transitions from ophiolite to red conglomerate and sandstone, to light-colored sandstone and limestone on the top; (i) red conglomerate composed mainly of silicified boulder and pebble clasts, lower part of the measured section; (j) parallel lamination in sandstone at ~94 m of the measured section; (k) reefal limestone; (l) close-up of the reefal limestone, showing coral frame structure, at ~105 m of the measured section.

4. Samples and Methods

We selected three sandstone samples for detrital zircon U–Pb geochronology, three for detrital spinel electron microprobe analysis, five for petrographic analysis, and eight limestone samples for coral and foraminifera identification in the Dongqiao Formation in our measured Zigetangco section (Figure 2a). We also collected one sandstone sample from the Dongqiao Formation along strike to the west in the Pari area (Figure 1b) for provenance analysis and eighteen limestone samples in the Dongqiao Formation and other Upper Jurassic strata in the region for biostratigraphy. Two quartzose sandstone samples (Figures S2c and S2d) of possible Paleozoic age were also collected for provenance analysis (Figure 1b). Details of the analytical methods and data are provided in the Supporting Information.

5. Stratigraphic and Provenance Results

5.1. Biostratigraphy

Foraminifera in the Zigetangco section include *Pseudocyclammina* sp. and *Everticyclammina virguliana* (Figure 2b). Foraminifera in other sites in the Dongqiao Formation and the Upper Jurassic strata in this region include *Siphovalvulina* sp., *Mesoendothyra croatica*, *Redmondoides lugeoni*, *Everticyclammina virguliana*, *Everticyclammina* sp., *Textularia* sp., *Valvulina* sp., *Pseudocyclammina* sp., *Pseudocyclammina sphaeroidalis*, *Daxia* sp., and *Nautiloculina* sp. (Figures 2c–2f and Table S2). Calcareous algae in the Zigetangco section include *Thaumatoporella* sp., *Salpingaporella annulata*, *Trinocladus perplexus*, and *Cayeuxia piae* Frolo (Table S2 and Figure S3a), with *Cayeuxia piae* also found in the Upper Jurassic. Corals in the Dongqiao Formation and the Upper Jurassic include *Dermosmilia laxata* (Étallon) (Figure S3f), *Cladocoropsis mirabilis* Felix (Figures 2g and S3b–S3be and S3bg) and *Thecosmilia shunghuensis* Liao (Table S2 and Figure S3e).

5.2. Provenance Data

Red sandstone in the lower part of the Dongqiao Formation in the Zigetangco section is dominated by detrital spinels and opaque minerals, while gray sandstone in the upper part is litho-quartzose (Figure 3a) with average QFL = 78:3:19 (Figure 3e). Monocrystalline quartz dominates over polycrystalline quartz (maximum of 18% of all quartz). Lithic fragments are dominated by quartz-muscovite schist (Figure 3b), limestone, and chert (Figure 3c), with subordinate felsic volcanic fragments. Muscovite and chromian spinel (Figure 3d) are common detrital minerals, and few ferromagnesian minerals were serpentinized and chloritized.

We determined a total of 438 U-Pb detrital zircon ages in four sandstone samples in the Dongqiao Formation. Cathodoluminescence images show that most of the zircons are texturally homogeneous, with a few showing oscillatory zoning and core-mantle-rim structure (Figure S4), indicating a protracted zircon growth history. The four samples show similar age spectra (Figure S5) and thus were plotted together (Figure 3f). Most of the zircons have U/Th values lower than 10, suggestive of an igneous origin (Figure 3f; e.g., Rubatto, 2002). Two dominant peaks are at 510–490 Ma and 1,300–1,100 Ma, with two broad subordinate peaks at 1,100–510 Ma and 2,000–1,500 Ma (Figure 3f). No zircons are younger than 460 Ma.

The 60 detrital spinels analyzed from three Dongqiao Formation samples can be divided into two groups based on the geochemical composition (Figures 3g and 3h). Cr# ($\text{Cr} \times 100 / (\text{Cr} + \text{Al})$ atomic ratio, 59–39) is relatively lower and TiO_2 (0.33–0.01%) is relatively higher in Group 1 ($n = 41$) compared to Group 2, in which Cr# and TiO_2 are 77–61 and 0.14–0%, respectively.

6. Discussion

6.1. Depositional Age

In both the Dongqiao Formation and nearby Upper Jurassic limestone, foraminifera including *Siphovalvulina* sp., *Mesoendothyra croatica*, *Redmondoides lugeoni*, *Everticyclammina virguliana*, *Everticyclammina* sp., *Textularia* sp., *Valvulina* sp., *Pseudocyclammina* sp., *Pseudocyclammina sphaeroidalis*, *Daxia* sp., and *Nautiloculina* sp. indicate an Oxfordian–Kimmeridgian age (Boudaughier-Fadel, 2018). Corals including *Dermosmilia laxata* (Étallon), *Cladocoropsis mirabilis* Felix, and *Thecosmilia shunghuensis* Liao indicate an Oxfordian–Kimmeridgian age (Dong & Wang, 1983; Liao et al., 2012). The fossils are intact, inconsistent with long-distance transport and reworking from older strata. Collectively, the identified fossils indicate an Oxfordian–Kimmeridgian depositional age for the Dongqiao Formation.

Previously, three foraminifera, *Kurnubia* sp. (or off. *Kurnubia*), *Pseudocyclammina* sp., and *Nautiloculina* sp., were used to argue for a broad Late Jurassic–Early Cretaceous age (Girardeau et al., 1984). However, *Kurnubia* sp., if present, rules out an Early Cretaceous age because it ranges from Bajocian to early Tithonian (Boudaughier-Fadel, 2018). Our study also narrows *Pseudocyclammina* sp. to *Pseudocyclammina sphaeroidalis*, which is of Kimmeridgian age (Boudaughier-Fadel, 2018). Finally, *Nautiloculina* species ranges from Jurassic (late Bajocian) to Early Cretaceous (Aptian) (Boudaughier-Fadel, 2018), but its coexistence with other Jurassic fossils points to a Jurassic age.

Figure 3. Provenance data of sandstones in the Dongqiao Formation. (a) Litho-quartzose sandstones, 15DQ11; (b) muscovite-quartz schist fragments, 17 AD48; (c) chert and limestone fragments, 17 AD44; (d) detrital chromian spinels, 17 AD44. Q, quartz; Qp, polycrystalline quartz; Pl, plagioclase; Lc, limestone fragments; (e) petrographic ternary plot. Cr-spinel (S) is incorporated into the lithic fragment (L) so that the Cr-spinel dominated red sandstone in the lower part of the section can also be shown in this ternary plot. F, feldspar; (f) kernel density estimation plots of U-Pb ages of detrital zircon from the Dongqiao Formation and Paleozoic (?), compared with those previously determined for the Qiangtang terrane (Gehrels et al., 2011; Pullen et al., 2011; Wang et al., 2016), Amdo basement (Guynn et al., 2012), Muganggri and Jienu groups (Li et al., 2020), and Lhasa terrane (Gehrels et al., 2011; Leier et al., 2007; Li et al., 2014; Zhu et al., 2011); (g) Cr# versus Mg# diagram of the detrital chromian spinels from the Dongqiao Formation. Field of abyssal peridotite is modified from Dick and Bullen (1984) and forearc peridotite from Ishii (1992); (h) TiO₂ versus Al₂O₃ tectonic discrimination diagram of the detrital chromian spinels, with fields modified from Kamenetsky et al. (2001). The geochemical data of chromian spinels from the Dongqiao ophiolite (Deng, 1988; Liu et al., 2016) are also plotted for comparison. MORB, mid-ocean ridge basalt; SSZ, supra-subduction zone; OIB, ocean island basalt; LIP, large igneous province.

6.2. Provenance Interpretation

Chert fragments and detrital chromian spinels in the Dongqiao Formation indicate a source from uplifted ophiolite and ocean plate stratigraphy. The high percentage of quartz, metamorphic, and sedimentary fragments implies an additional provenance from continental crust. Limestone fragments may have been derived from within the basin or have been recycled from older lithified limestone assemblages exposed within or adjacent to the Bangong-Nujiang suture zone. Muscovite and spinel are attributed to schist and ultramafic rocks, respectively, in the source region.

The continental crust and ophiolite provenance can be further determined using U-Pb ages of detrital zircons and major element abundances of detrital chromian spinels, respectively. U-Pb detrital zircon age

Figure 4. Cartoon showing the tectonic, paleogeographic, and depositional setting of the Dongqiao Formation. The Dongqiao ophiolite was obducted southward onto the Lhasa terrane, or a microcontinent of Lhasa-affinity. Local sources shed detritus northward into the Dongqiao Formation, while limestone was deposited more distally to the north. These stratigraphic relationships are consistent with those suggested by Girardeau et al. (1984) and Kidd et al. (1988).

distributions in the Dongqiao Formation are similar to those within pre-Jurassic sandstones of the Lhasa terrane and Paleozoic (?) quartzose sandstones in the Dongqiao area (Figure 3f). Given the sandstone detrital composition, detrital zircon U-Pb ages, and presence of alluvial fan conglomerates, the Dongqiao Formation is interpreted to have been sourced locally from metasedimentary rocks of Lhasa terrane affinity and the Dongqiao ophiolite. Jurassic calc-alkaline rocks are abundant in the southern Qiangtang and northern Lhasa terranes, Baila area, and Amdo basement but are absent in the Dongqiao area (e.g., Zhu et al., 2016; Li et al., 2020; Figure 1b). This may explain why syn-depositional zircons were not identified in the locally sourced Dongqiao Formation. The Amdo basement and Qiangtang terrane are unlikely sources because they exhibit a major U-Pb zircon age population at ~1,000–500 Ma, which is not a dominant age peak in the Dongqiao Formation (Figure 3f).

In binary plots of TiO_2 vs. Cr_2O_3 and $\text{Cr}\#$ vs. $\text{Mg}\#$ (after Kamenetsky et al., 2001 and Dick & Bullen, 1984, respectively), the Group 1 chromian spinels are similar to those from cumulate (including troctolite, olivine pyroxenite, and wehrlite) and Group 2 chromian spinels are similar to those from harzburgite and dunite from the underlying Dongqiao ophiolite (Deng, 1988; Liu et al., 2016; Figures 3g and 3h). This provides additional evidence that the Dongqiao ophiolite contributed detritus to the Dongqiao Formation. Both groups of chromian spinels plot in the forearc peridotite field on a $\text{Cr}\#$ vs $\text{Mg}\#$ diagram and the SSZ peridotite field on a TiO_2 vs. Cr_2O_3 diagram, which indicate that the Dongqiao ophiolite source crystallized in a forearc spreading setting as previously suggested (Liu et al., 2016; Zhou et al., 1997).

6.3. Implication for the History of Ophiolite Obduction

The Dongqiao ophiolite crystallized at 189–181 Ma (Table S1) and includes a metamorphic sole with ~180–175 Ma hornblende $^{40}\text{Ar}/^{39}\text{Ar}$ dates (Zhou et al., 1997). These dating results can be explained by rapid cooling shortly after Dongqiao ophiolite generation in a forearc spreading setting above a north-dipping intra-oceanic subduction zone—similar to other large-slab ophiolites such as the Bay of Islands and Semail ophiolites (Dewey & Casey, 2011).

The Dongqiao Formation has a Lhasa terrane provenance and unconformably overlies the Dongqiao ophiolite, which in turn is locally thrust on top of Lhasa-affinity Paleozoic strata (Figure 4; Girardeau et al., 1984; Kidd et al., 1988). Based on these geological relationships and provenance of the Dongqiao Formation, we conclude that the Dongqiao ophiolite was obducted onto the Lhasa terrane, or a micro-continent of Lhasa-provenance-affinity prior to the deposition of the Dongqiao Formation at ~163–152 Ma according to our biostratigraphic data. The hypothetical micro-continent may be bounded by the Dongqiao ophiolite

in the north and the Baila ophiolite belt to the south (e.g., Zeng et al., 2016; Figure 1b). Additional evidence for existence of a micro-continent is the presence of isotopically-evolved I-type 166–160 Ma granitoids within it, which were interpreted to have been generated by partial melting of Proterozoic-Archean crust (Li et al., 2020). The history of Dongqiao ophiolite obduction and subsequent Dongqiao Formation deposition may be comparable to the Late Cretaceous geological evolution of Oman, where the Semail ophiolite crystallized during the earliest Late Cretaceous (e.g., Rioux et al., 2016), shortly prior to its obduction onto the Arabian passive continental margin and subsequent burial by Late Campanian–early Maastrichtian locally sourced stream-dominated and marine-influenced fan deposits (Abbasi et al., 2014; Alsharhan & Nasir, 1996).

Our tectonic interpretation suggests that a Lhasa terrane affinity continental margin entered and ultimately led to the demise of a north-dipping oceanic subduction zone within the Meso-Tethys Ocean during Middle Jurassic time (between ~180–175 Ma cooling of the metamorphic sole and ~163–152 Ma deposition of the Dongqiao Formation). It remains to be definitively determined, however, whether Dongqiao ophiolite obduction marks only the demise of an intra-oceanic subduction zone within the Meso-Tethys Ocean or also the onset of Lhasa-Qiangtang collision. In a broader context, termination of the Meso-Tethys north-dipping subduction zone may have induced plate kinematic reorganizations and generation of ophiolites above newly initiated oceanic subduction zones to the south. This could potentially explain generation of Middle to Late Jurassic ophiolites near Baila and Xainza in the Lhasa terrane (Fan et al., 2017; Tang et al., 2018; Zhong et al., 2017) and within the Yarlung-Tsangpo suture farther to the south (~177–150 Ma; e.g., Pedersen et al., 2001; Hébert et al., 2012).

7. Conclusions

(1) The Dongqiao Formation was deposited on ophiolite and thus post-dates ophiolite obduction. (2) Biostratigraphy tightly constrains deposition of the Dongqiao Formation, and the minimum age of ophiolite obduction to be 163–152 Ma. (3) The Dongqiao Formation includes sandstones deposited in subaerial to shallow marine environments and which were derived from local sources to the south including the ophiolite and Paleozoic (?) strata of Lhasa terrane affinity. We conclude that the Dongqiao ophiolite was obducted southward onto the Lhasa terrane or a microcontinent of Lhasa terrane affinity after ophiolite crystallization (189–181 Ma) and before the Oxfordian (>163 Ma). The obduction marks the demise of a north-dipping subduction zone in the Bangong-Nujiang Ocean, which in turn may have induced plate kinematic reorganizations along the southern Asian margin to the south during Middle to Late Jurassic time.

Acknowledgments

The data supporting the conclusions can be found in the Supporting Information and at the website (<https://osf.io/5nv7d/>). We thank Weiwei Xue, Zhong Han, Bo Zhou, and Hanpu Fu for assistance in the field, and Weihua Liao for coral identification. This manuscript benefited from comments and suggestions by Christopher Spencer, Ryan Leary, and an anonymous reviewer, and editorial handling by Steve Jacobsen. This work was financially supported by the Distinguished Young Scholar (41525007) and Tethys Major Project which are funded by the National Natural Science Foundation of China (91755209).

References

- Abbasi, I. A., Hersi, O. S., & Al-Harthi, A. (2014). Late Cretaceous conglomerates of the Qahlah Formation, North Oman. *Geological Society, London, Special Publications*, 392(1), 325–341. <https://doi.org/10.1144/SP392.17>
- Agard, P., Omrani, J., Jolivet, L., Whitechurch, H., Vrielynck, B., Spakman, W., et al. (2011). Zagros orogeny: A subduction-dominated process. *Geological Magazine*, 148(5–6), 692–725. <https://doi.org/10.1017/S001675681100046X>
- Alavi, M. (1994). Tectonics of the Zagros orogenic belt of Iran: New data and interpretations. *Tectonophysics*, 229(3–4), 211–238. [https://doi.org/10.1016/0040-1951\(94\)90030-2](https://doi.org/10.1016/0040-1951(94)90030-2)
- Alsharhan, A., & Nasir, S. (1996). Sedimentological and geochemical interpretation of a transgressive sequence: The Late Cretaceous Qahlah Formation in the western Oman Mountains, United Arab Emirates. *Sedimentary Geology*, 101(3–4), 227–242. [https://doi.org/10.1016/0037-0738\(95\)00067-4](https://doi.org/10.1016/0037-0738(95)00067-4)
- Boudaughier-Fadel, M. K. (2018). *Evolution and geological significance of larger benthic foraminifera*. London: UCL Press. <https://doi.org/10.14324/111.9781911576938>
- Coleman, R. G. (1981). Tectonic setting for ophiolite obduction in Oman. *Journal of Geophysical Research*, 86(B4), 2497–2508. <https://doi.org/10.1029/JB086iB04p02497>
- Deng, W. (1988). Chemical compositions of spinel group from metamorphic peridotites and cumulates in north Tibetan ophiolite. *Scientia Geologica Sinica*, 2, 121–127.
- Dewey, J. (1976). Ophiolite obduction. *Tectonophysics*, 31(1–2), 93–120. [https://doi.org/10.1016/0040-1951\(76\)90169-4](https://doi.org/10.1016/0040-1951(76)90169-4)
- Dewey, J. F., & Casey, J. F. (2011). The origin of obducted large-slab ophiolite complexes. In D. Brown, & P. D. Ryan (Eds.), *Arc-continent collision*. *Frontiers in earth sciences* (pp. 431–443). Berlin: Springer. https://doi.org/10.1007/978-3-540-88558-0_15
- Dewey, J., Shackleton, R., Chang, C., & Sun, Y. (1988). The tectonic evolution of the Tibetan Plateau. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 327(1594), 379–413. <https://doi.org/10.1098/rsta.1988.0135>
- Dick, H. J., & Bullen, T. (1984). Chromian spinel as a petrogenetic indicator in abyssal and alpine-type peridotites and spatially associated lavas. *Contributions to Mineralogy and Petrology*, 86(1), 54–76. <https://doi.org/10.1007/BF00373711>
- Dong, D., & Wang, M. (1983). New materials of the upper Jurassic Stromatoporoidea in the Anduo County of North Xizang. *Acta Palaeontologica Sinica*, 22(4), 413–427.
- Fan, S., Ding, L., Murphy, M. A., Yao, W., & Yin, A. (2017). Late Paleozoic and Mesozoic evolution of the Lhasa Terrane in the Xainza area of southern Tibet. *Tectonophysics*, 721, 415–434. <https://doi.org/10.1016/j.tecto.2017.10.022>

- Gehrels, G., Kapp, P., DeCelles, P., Pullen, A., Blakey, R., Weislogel, A., et al. (2011). Detrital zircon geochronology of pre-tertiary strata in the Tibetan-Himalayan orogen. *Tectonics*, 30, TC5016. <https://doi.org/10.1029/2011tc002868>
- Girardeau, J., Marcoux, J., Allegre, C. J., Bassoulet, J. P., Tang, Y. K., Xiao, X. C., et al. (1984). Tectonic environment and geodynamic significance of the Neo-Cimmerian Dongqiao Ophiolite, Bangong-Nujiang suture zone, Tibet. *Nature*, 307(5946), 27–31. <https://doi.org/10.1038/307027a0>
- Guynn, J., Kapp, P., Gehrels, G. E., & Ding, L. (2012). U–Pb geochronology of basement rocks in Central Tibet and paleogeographic implications. *Journal of Asian Earth Sciences*, 43(1), 23–50. <https://doi.org/10.1016/j.jseas.2011.09.033>
- Guynn, J., Kapp, P., Pullen, A., Heizler, M., Gehrels, G., & Ding, L. (2006). Tibetan basement rocks near Amdo reveal “missing” Mesozoic tectonism along the Bangong suture, central Tibet. *Geology*, 34(6), 505–508. <https://doi.org/10.1130/G22453.1>
- Hacker, B., Mosenfelder, J., & Gnoss, E. (1996). Rapid emplacement of the Oman ophiolite: Thermal and geochronologic constraints. *Tectonics*, 15(6), 1230–1247. <https://doi.org/10.1029/96TC01973>
- Hébert, R., Bezard, R., Guilmette, C., Dostal, J., Wang, C., & Liu, Z. (2012). The Indus–Yarlung Zangbo ophiolites from Nanga Parbat to Namche Barwa syntaxes, southern Tibet: First synthesis of petrology, geochemistry, and geochronology with incidences on geodynamic reconstructions of Neo-Tethys. *Gondwana Research*, 22(2), 377–397. <https://doi.org/10.1016/j.gr.2011.10.013>
- Institute of Tibetan Geological Survey. (2002). Geological Report of the 1:250, 000 Regional Geological Survey in Baingoin Area (in Chinese).
- Institute of Tibetan Geological Survey. (2003). Geological Report of the 1:250, 000 Regional Geological Survey in Zigetangco Area (in Chinese).
- Institute of Tibetan Geological Survey. (2005). Geological Report of the 1:250, 000 Regional Geological Survey in Amdo Area (in Chinese).
- Ishii, T. (1992). Petrological studies of peridotites from diapiric serpentinite seamounts in the Izu-Ogasawara-Mariana forearc, Leg 125. Paper presented at the proceedings of the ocean drilling program, scientific results.
- Kamenetsky, V. S., Crawford, A. J., & Meffre, S. (2001). Factors controlling chemistry of magmatic spinel: An empirical study of associated olivine, Cr-spinel and melt inclusions from primitive rocks. *Journal of Petrology*, 42(4), 655–671. <https://doi.org/10.1093/petrology/42.4.655>
- Kapp, P., & DeCelles, P. G. (2019). Mesozoic–Cenozoic geological evolution of the Himalayan–Tibetan orogen and working tectonic hypotheses. *American Journal of Science*, 319(3), 159–254. <https://doi.org/10.2475/03.2019.01>
- Kapp, P., DeCelles, P. G., Gehrels, G. E., Heizler, M., & Ding, L. (2007). Geological records of the Lhasa–Qiangtang and Indo–Asian collisions in the Nima area of central Tibet. *Geological Society of America Bulletin*, 119(7–8), 917–932. <https://doi.org/10.1130/b26033.1>
- Kapp, P., Murphy, M. A., Yin, A., Harrison, T. M., Ding, L., & Guo, J. H. (2003). Mesozoic and Cenozoic tectonic evolution of the Shiquanhe area of western Tibet. *Tectonics*, 22(4), 1029. <https://doi.org/10.1029/2001tc001332>
- Kidd, W., Pan, Y., Chang, C., Coward, M. P., Dewey, J. F., Gansser, A., et al. (1988). Geological mapping of the 1985 Chinese–British Tibetan (Xizang–Qinghai) plateau Geotraverse route. *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 327(1594), 287–305. <https://doi.org/10.1098/rsta.1988.0130>
- Leeder, M. R., Smith, A. B., & Yin, J. X. (1988). Sedimentology, paleoecology and palaeoenvironmental evolution of the 1985 Lhasa to Golmud Geotraverse. *Philosophical Transactions Of the Royal Society a-Mathematical Physical And Engineering Sciences*, 327(1594), 107–143. <https://doi.org/10.1098/rsta.1988.0123>
- Leier, A. L., Kapp, P., Gehrels, G. E., & DeCelles, P. G. (2007). Detrital zircon geochronology of Carboniferous–Cretaceous strata in the Lhasa terrane, southern Tibet. *Basin Research*, 19(3), 361–378. <https://doi.org/10.1111/j.1365-2117.2007.00330.x>
- Li, G. W., Sandiford, M., Liu, X. H., Xu, Z. Q., Wei, L. J., & Li, H. Q. (2014). Provenance of Late Triassic sediments in central Lhasa terrane, Tibet and its implication. *Gondwana Research*, 25(4), 1680–1689. <https://doi.org/10.1016/j.gr.2013.06.019>
- Li, S., Guilmette, C., Yin, C., Ding, L., Zhang, J., Wang, H., & Baral, U. (2019). Timing and mechanism of Bangong–Nujiang ophiolite emplacement in the Gerze area of Central Tibet. *Gondwana Research*, 71, 179–193. <https://doi.org/10.1016/j.gr.2019.01.019>
- Li, S., Yin, C., Guilmette, C., Ding, L., & Zhang, J. (2019). Birth and demise of the Bangong–Nujiang Tethyan Ocean: A review from the Gerze area of Central Tibet. *Earth-Science Reviews*, 102907. <https://doi.org/10.1016/j.earscirev.2019.102907>
- Li, S. M., Wang, Q., Zhu, D. C., Cawood, P. A., Stern, R. J., Weinberg, R., et al. (2020). Reconciling orogenic drivers for the evolution of the Bangong–Nujiang Tethys during Middle–Late Jurassic. *Tectonics*. <https://doi.org/10.1029/2019TC005951>
- Liao, W., Ji, Z., & Wu, G. (2012). Late Jurassic Scleractinian corals from Gerze, northwestern Xizang (Tibet). *Acta Palaeontologica Sinica*, 51(3), 290–307.
- Liu, T., Zhai, Q.-G., Wang, J., Bao, P.-S., Qiangba, Z., Tang, S.-H., & Tang, Y. (2016). Tectonic significance of the Dongqiao ophiolite in the north-central Tibetan plateau: Evidence from zircon dating, petrological, geochemical and Sr–Nd–Hf isotopic characterization. *Journal of Asian Earth Sciences*, 116, 139–154. <https://doi.org/10.1002/j.jseas.2015.11.014>
- Ma, A., Hu, X., Garzanti, E., Han, Z., & Lai, W. (2017). Sedimentary and tectonic evolution of the southern Qiangtang basin: Implications for the Lhasa–Qiangtang collision timing. *Journal of Geophysical Research: Solid Earth*, 122, 4790–4813. <https://doi.org/10.1002/2017jb014211>
- Ma, A., Hu, X., Kapp, P., Han, Z., Lai, W., & BouDagher-Fadel, M. (2018). The disappearance of a Late Jurassic remnant sea in the southern Qiangtang Block (Shamuluo Formation, Njangco area): Implications for the tectonic uplift of central Tibet. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 506, 30–47. <https://doi.org/10.1016/j.palaeo.2018.06.005>
- Marcoux, J., Girardeau, J., Fourcade, E., Bassoulet, J.-P., Philip, J., Jaffrezo, M., et al. (1987). Geology and biostratigraphy of the Jurassic and Lower Cretaceous series to the north of the Lhasa Block (Tibet, China). *Geodinamica Acta*, 1(4–5), 313–325. <https://doi.org/10.1080/09853111.1987.11105148>
- Murphy, M. A., Yin, A., Harrison, T. M., Dürr, S. B., Chen, Z., Ryerson, F. J., et al. (1997). Did the Indo–Asian collision alone create the Tibetan plateau? *Geology*, 25(8), 719–722. [https://doi.org/10.1130/0091-7613\(1998\)026<0958:DTIACA>2.3.CO;2](https://doi.org/10.1130/0091-7613(1998)026<0958:DTIACA>2.3.CO;2)
- Pedersen, R., Searle, M., & Corfield, R. (2001). U–Pb zircon ages from the Spontang ophiolite, Ladakh Himalaya. *Journal of the Geological Society*, 158(3), 513–520. <https://doi.org/10.1144/jgs.158.3.513>
- Pullen, A., Kapp, P., Gehrels, G. E., Ding, L., & Zhang, Q. H. (2011). Metamorphic rocks in central Tibet: Lateral variations and implications for crustal structure. *Geological Society of America Bulletin*, 123(3–4), 585–600. <https://doi.org/10.1130/b30154.1>
- Rateman, N. S., Robinson, A. C., & Cowgill, E. S. (2014). Structure and detrital zircon geochronology of the Domar fold-thrust belt: Evidence of pre-Cenozoic crustal thickening of the western Tibetan Plateau. *Geological Society of America Special Papers*, 507, 89–104. [https://doi.org/10.1130/2014.2507\(05\)](https://doi.org/10.1130/2014.2507(05))
- Rioux, M., Garber, J., Bauer, A., Bowring, S., Searle, M., Kelemen, P., & Hacker, B. (2016). Synchronous formation of the metamorphic sole and igneous crust of the Semail ophiolite: New constraints on the tectonic evolution during ophiolite formation from high-precision U–Pb zircon geochronology. *Earth and Planetary Science Letters*, 451(2), 185–195. <https://doi.org/10.1016/j.palaeo.2018.06.005>

- Rubatto, D. (2002). Zircon trace element geochemistry: Partitioning with garnet and the link between U–Pb ages and metamorphism. *Chemical Geology*, 184(1–2), 123–138. [https://doi.org/10.1016/S0009-2541\(01\)00355-2](https://doi.org/10.1016/S0009-2541(01)00355-2)
- Tang, Y., Zhai, Q.-G., Hu, P.-Y., Wang, J., Xiao, X.-C., Wang, H.-T., et al. (2018). Rodingite from the Beila ophiolite in the Bangong–Nujiang suture zone, northern Tibet: New insights into the formation of ophiolite-related rodingite. *Lithos*, 316, 33–47. <https://doi.org/10.1016/j.lithos.2018.07.006>
- Wakabayashi, J., & Dilek, Y. (2003). What constitutes ‘emplacement’ of an ophiolite?: Mechanisms and relationship to subduction initiation and formation of metamorphic soles. *Geological Society, London, Special Publications*, 218(1), 427–447. <https://doi.org/10.1144/GSL.SP.2003.218.01.22>
- Wang, J. G., Wu, F. Y., Garzanti, E., Hu, X. M., Ji, W. Q., Liu, Z. C., & Liu, X. C. (2016). Upper Triassic turbidites of the northern Tethyan Himalaya (Langjixue Group): The terminal of a sediment-routing system sourced in the Gondwanide Orogen. *Gondwana Research*, 34, 84–98. <https://doi.org/10.1016/j.jgr.2016.03.005>
- Wang, M., & Dong, D. (1984). Stromatoporoids from the Dongqiao Formation (Upper Jurassic–Lower Cretaceous) in northern Xizang (Tibet) [in Chinese with English abstract]. *Acta Palaeontologica Sinica*, 12(3), 343–352.
- Yin, A., & Harrison, T. M. (2000). Geologic evolution of the Himalayan–Tibetan orogen. *Annual Review of Earth and Planetary Sciences*, 28(1), 211–280. <https://doi.org/10.1146/annurev.earth.28.1.211>
- Zeng, Y. C., Chen, J. L., Xu, J. F., Wang, B. D., & Huang, F. (2016). Sediment melting during subduction initiation: Geochronological and geochemical evidence from the Darutso high-Mg andesites within ophiolite melange, central Tibet. *Geochemistry, Geophysics, Geosystems*, 17(12), 4859–4877. <https://doi.org/10.1002/2016GC006456>
- Zhong, Y., Liu, W.-L., Xia, B., Liu, J.-N., Guan, Y., Yin, Z.-X., & Huang, Q.-T. (2017). Geochemistry and geochronology of the Mesozoic Lanong ophiolitic melange, northern Tibet: Implications for petrogenesis and tectonic evolution. *Lithos*, 292, 111–131. <https://doi.org/10.1016/j.lithos.2017.09.003>
- Zhou, M. F., Malpas, J., Robinson, P. T., & Reynolds, P. H. (1997). The dynamothermal aureole of the Dongqiao ophiolite (northern Tibet). *Canadian Journal of Earth Sciences*, 34(1), 59–65. <https://doi.org/10.1139/e17-005>
- Zhu, D. C., Li, S.-M., Cawood, P. A., Wang, Q., Zhao, Z.-D., Liu, S.-A., & Wang, L.-Q. (2016). Assembly of the Lhasa and Qiangtang terranes in central Tibet by divergent double subduction. *Lithos*, 245, 7–17. <https://doi.org/10.1139/e17-005>
- Zhu, D. C., Zhao, Z. D., Niu, Y., Dilek, Y., & Mo, X. X. (2011). Lhasa terrane in southern Tibet came from Australia. *Geology*, 39(8), 727–730. <https://doi.org/10.1130/g31895.1>

References From the Supporting Information

- Andersen, T. (2002). Correction of common lead in U–Pb analyses that do not report 204Pb. *Chemical Geology*, 192(1–2), 59–79. [https://doi.org/10.1016/S0009-2541\(02\)00195-X](https://doi.org/10.1016/S0009-2541(02)00195-X)
- Barnes, S. J., & Roeder, P. L. (2001). The range of spinel compositions in terrestrial mafic and ultramafic rocks. *Journal of Petrology*, 42(12), 2279–2302. <https://doi.org/10.1093/petrology/42.12.2279>
- Black, L., & Gulson, B. (1978). The age of the mud tank carbonatite, strangways range, northern territory. *Journal of Australian Geology and Geophysics*, 3, 227–232.
- Boudagher-Fadel, M. K. (2018a). Revised diagnostic first and last occurrences of Mesozoic and Cenozoic planktonic foraminifera. UCL Office of the Vice-Provost Research, Professional Papers Series(2), 1–5.
- Boudagher-Fadel, M. K. (2018b). *Evolution and geological significance of larger benthic foraminifera, second edition*. London: UCL Press. <https://doi.org/10.14324/111.9781911576938>
- Gradstein, F. M., Ogg, J. G., Schmitz, M., & Ogg, G. (2012). *The geologic time scale 2012*. New York: Elsevier. <https://doi.org/10.4095/215638>
- Horstwood, M. S., Köhler, J., Gehrels, G., Jackson, S. E., McLean, N. M., Paton, C., et al. (2016). Community-derived standards for LA-ICP-MS U–(Th)–Pb geochronology—uncertainty propagation, age interpretation and data reporting. *Geostandards and Geoanalytical Research*, 40(3), 311–332. <https://doi.org/10.1111/j.1751-908X.2016.00379.x>
- Huang, Q. S., Shi, R. D., Liu, D. L., Zhang, X. R., Fan, S. Q., & Ding, L. (2013). Os isotopic evidence for a carbonaceous chondritic mantle source for the Nagqu ophiolite from Tibet and its implications. *Chinese Science Bulletin*, 58(1), 92–98. <https://doi.org/10.1007/s11434-012-5384-8>
- Huang, Q. T., Li, J. F., Xia, B., Yin, Z. X., Zheng, H., Shi, X. L., & Hu, X. C. (2015). Petrology, geochemistry, chronology and geological significance of Jiang Tso ophiolite in middle segment of Bangonghu–Nujiang suture zone, Tibet. *Journal of China University of Geosciences*, 40(1), 34–48. <https://doi.org/10.3799/dqkx.2015.003>
- Jackson, S. E., Pearson, N. J., Griffin, W. L., & Belousova, E. A. (2004). The application of laser ablation-inductively coupled plasma-mass spectrometry to in situ U–Pb zircon geochronology. *Chemical Geology*, 211(1–2), 47–69. <https://doi.org/10.1016/j.chemgeo.2004.06.017>
- Li, J. F., Xia, B., Xia, L. Z., Xu, L. F., Liu, W. L., Cai, Z. R., & Yang, Z. Q. (2013). Geochronology of the Dong Tso Ophiolite and the tectonic environment. *Acta Geologica Sinica (English Edition)*, 87(6), 1604–1616. <https://doi.org/10.1111/1755-6724.12162>
- Liu, T., Zhai, Q.-G., Wang, J., Bao, P.-S., Qiangba, Z., Tang, S.-H., & Tang, Y. (2016). Tectonic significance of the Dongqiao ophiolite in the north-central Tibetan plateau: Evidence from zircon dating, petrological, geochemical and Sr–Nd–Hf isotopic characterization. *Journal of Asian Earth Sciences*, 116, 139–154. <https://doi.org/10.1002/j.jseas.2015.11.014>
- Qu, X., Xin, H., Zhao, Y., & Wang, R. (2010). Opening time of Bangong Lake middle Tethys oceanic basin of the Tibet plateau: Constraints from petro-geochemistry and zircon U–Pb LAICPMS dating of mafic ophiolites. *Earth Science Frontiers*, 17(3), 53–63.
- Shi, R. (2007). SHRIMP dating of the Bangong Lake SSZ-type ophiolite: Constraints on the closure time of ocean in the Bangong Lake–Nujiang River, northwestern Tibet. *Chinese Science Bulletin*, 52(7), 936–941. <https://doi.org/10.1007/s11434-007-0134-z>
- Spencer, C. J., Kirkland, C. L., & Taylor, R. J. (2016). Strategies towards statistically robust interpretations of in situ U–Pb zircon geochronology. *Geoscience Frontiers*, 7(4), 581–589. <https://doi.org/10.1016/j.gsf.2015.11.006>
- Sun, L., Bai, Z., Xun, D., Li, H., & Sun, B. (2011). Geological characteristics and zircon U–Pb SHRIMP dating of the plagiogranite in Amduo ophiolites, Tibet. *Geological Survey and Research*, 34(1), 10–15.
- Tang, Y., Zhai, Q.-G., Hu, P.-Y., Wang, J., Xiao, X.-C., Wang, H.-T., et al. (2018). Rodingite from the Beila ophiolite in the Bangong–Nujiang suture zone, northern Tibet: New insights into the formation of ophiolite-related rodingite. *Lithos*, 316, 33–47. <https://doi.org/10.1016/j.lithos.2018.07.006>
- Tang, Y., Zhai, Q., Hu, P., Xiao, X., & Wang, H. (2018). Petrology, geochemistry and geochronology of the Zhongcang ophiolite, northern Tibet: Implications for the evolution of the Bangong–Nujiang Ocean. *Geoscience Frontiers*, 9(5), 1369–1381. <https://doi.org/10.1016/j.gsf.2018.05.007>

- Vermeesch, P., Resentini, A., & Garzanti, E. (2016). An R package for statistical provenance analysis. *Sedimentary Geology*, 336, 14–25. <https://doi.org/10.1016/j.sedgeo.2016.01.009>
- Wang, B.-D., Wang, L.-Q., Chung, S.-L., Chen, J.-L., Yin, F.-G., Liu, H., et al. (2016). Evolution of the Bangong–Nujiang Tethyan Ocean: Insights from the geochronology and geochemistry of mafic rocks within ophiolites. *Lithos*, 245, 18–33. <https://doi.org/10.1016/j.lithos.2015.07.016>
- Wu, Y., Cheng, S. Y., Qin, M. K., Guo, D. F., Guo, G. L., Zhang, C., & Yang, J. S. (2018). Zircon U-Pb ages of Dongcuo Ophiolite in Western Bangonghu–Nujiang suture zone and their geological significance. *Earth Science*, 43(4).
- Xia, B., Xu, L., Wei, Z., Zhang, Y., Wang, R., Li, J., & Wang, Y. (2008). SHRIMP zircon dating of gabbro from the Dongqiao ophiolite in Tibet and its geological implications. *Acta Geologica Sinica*, 82(4), 528–531.
- Xu, M., Li, C., Xu, W., Xie, C., Hu, P., & Wang, M. (2014). Petrology, geochemistry and geochronology of gabbros from the Zhongcang ophiolitic mélange, Central Tibet: Implications for an intra-oceanic subduction zone within the neo-Tethys Ocean. *Journal of Earth Science*, 25(2), 224–240.
- Zhang, Y. X., Zhang, K. J., Li, B., Wang, Y., Wei, Q. G., & Tang, X. C. (2007). Zircon SHRIMP U-Pb geochronology and petrogenesis of the plagiogranites from the Lagkor Lake ophiolite, Gerze, Tibet, China. *Chinese Science Bulletin*, 52(5), 651–659. <https://doi.org/10.1007/s11434-007-0084-5>
- Zhong, Y., Liu, W.-L., Xia, B., Liu, J.-N., Guan, Y., Yin, Z.-X., & Huang, Q.-T. (2017). Geochemistry and geochronology of the Mesozoic Lanong ophiolitic mélange, northern Tibet: Implications for petrogenesis and tectonic evolution. *Lithos*, 292, 111–131. <https://doi.org/10.1016/j.lithos.2017.09.003>