INVASIVE CONJUNCTIVAL MELANOMA MIMICKING OCULAR SURFACE SQUAMOUS NEOPLASIA: A CASE SERIES

¹Beatrice Gallo, ^{2,3}Caroline Thaung, ¹Gordon Hay, ¹Amit K. Arora, ^{1,2,3}Victoria M.L. Cohen, ¹Bertil Damato, ^{1, 2,3}Mandeep S. Sagoo

AFFILIATIONS

¹Ocular Oncology Service, Moorfields Eye Hospital, City Road, London, EC1V 2PD, United Kingdom.

²University College London, Institute of Ophthalmology, Bath Street, London, EC1V 9EL, United Kingdom.

³NIHR Biomedical Research Centre for Ophthalmology at Moorfields Eye Hospital and University College London Institute of Ophthalmology, London, United Kingdom.

CORRESPONDING AUTHOR

Beatrice Gallo

Ocular Oncology Service, Moorfields Eye Hospital, City Road, London, EC1V 2PD, United Kingdom.

e-mail address: beatricegallo.bg@gmail.com

phone number: +447401710381

SYNOPSIS

Conjunctival melanoma in our patients showed features suggesting ocular surface squamous neoplasia. Given the heterogeneous and sometimes misleading clinical presentation, the histopathological analysis remains the diagnostic gold standard. Inappropriate management can adversely affect the prognosis.

ABSTRACT

Background: Conjunctival melanoma is the second most common conjunctival malignant tumor after squamous cell carcinoma, usually arising from primary acquired melanosis and less commonly from a conjunctival naevus or de novo. We report four cases of conjunctival melanoma masquerading as ocular surface squamous neoplasia.

Methods: Four patients (2 females and 2 males; mean age 60.7 years; range 41-72 years) were referred for suspicious conjunctival lesions. In all cases the lesions had a perilimbal location, were non-pigmented (cases 1 and 3) or mildly pigmented (cases 2 and 4), had a fleshy (cases 1, 2 and 4) or papillomatous (case 3) appearance and involved the corneal surface. In each case, our main clinical differential diagnosis included conjunctival intraepithelial neoplasia and squamous cell carcinoma. All four patients underwent an excisional biopsy with double freeze-thaw cryotherapy and alcohol keratoepitheliectomy.

Results: In all 4 cases the histopathological diagnosis was of invasive conjunctival melanoma with extension to the deep surgical margins. Adjuvant therapy consisting of strontium-90 β radiotherapy (all 4 patients) and topical Mitomycin C (patient 2) was administered.

Conclusion: Conjunctival melanoma can clinically resemble ocular surface squamous neoplasia. Clinical impressions therefore need to be confirmed histopathologically.

Keywords: conjunctiva, tumor, melanoma, ocular surface squamous neoplasia.

INTRODUCTION

Conjunctival melanoma (CM) is rare (1-2% of all ocular melanomas) but carries a 32% risk of metastatic spread to regional lymph nodes and other organs at 15 years (1). After squamous cell carcinoma (SCC), CM is the most common conjunctival malignant tumor. The incidence is 0.2-0.8/million population per year (1) with an upward trend over the last three decades (2, 3). The mean age at presentation is 60 years (4). Sunlight and ultraviolet light exposure is an important risk factor (5).

CM usually develops from primary acquired melanosis (PAM) with atypia (75%), and less frequently from a pre-existing conjunctival naevus (20%) or *de novo* (5%) (5). According to an outcome multivariable analysis based on CM origin and location, *de novo* melanoma and nodular tumour are the main risk factors for melanoma-related metastasis and death (6). Treatment of CM includes surgical resection of the conjunctival component with the 'no touch' technique, superficial keratectomy of the corneal component and cryotherapy to the adjacent margins. For tumours invading the sclera a lamellar resection is necessary. Exenteration is reserved for cases of orbital invasion or otherwise unresectable recurrent disease. Adjuvant treatments are used in cases of incomplete surgical excision and include topical chemotherapy with Mitomycin C (0.02% or 0.04%) and in case of deep margin involvement adjunctive radiotherapy delivered with a ruthenium or iodine plaque, strontium applicator or proton beam radiotherapy (5).

Typical clinical features of conjunctival melanoma include a pigmented thickened conjunctival mass with feeder vessels. However, not all melanomas conform to the classical picture. Nodular melanoma is often associated with flat conjunctival pigmentation, consisting either of primary acquired melanosis or secondary pagetoid spread from the nodular tumour (7). Less frequently, CM is non-pigmented. Inadvertent incisional biopsy or incomplete excision are associated with local recurrences

and increased mortality (6). Herein, we describe four cases of invasive CM masquerading as Ocular Surface Squamous Neoplasia (OSSN), with the caution that this atypical mode of presentation should be recognized to avoid mismanagement.

SUBJECTS AND METHODS

We included in our series four patients referred to the Ocular Oncology Service of Moorfields Eye Hospital with suspected conjunctival squamous neoplasia between 2011 and 2019. This study was registered with the Research and Development Department of Moorfields Eye Hospital (CaRS_10) and adhered to the tenets of the Declaration of Helsinki.

Case 1

A 66-year-old Caucasian female presented with a three-year history of a conjunctival lesion in her left eye, noting a slowly enlarging red mark with a foreign body sensation. Although the lesion was initially ascribed to actinic damage, a papilloma was also considered because of a past history of a left lower lid papilloma 4 years prior. There was no clear history of prolonged exposure to UV radiation. Her general health was unremarkable. Family history was positive for leukemia (sister).

On examination, the visual acuity was 6/6 OD and 6/5 OS. The intraocular pressures were normal. The right eye was healthy. Examination of the left eye revealed a pale, gelatinous, temporal limbal conjunctival lesion, with basal dimensions measuring 5 mm transversely and 5.5 mm longitudinally, with the tumour encroaching the cornea for 2 mm. The conjunctival tumor was freely mobile with no attachment to the underlying sclera. There were dilated feeder vessels and prominent intrinsic tumor vascularity (Fig 1a).

Case 2

A 41-year-old Caucasian female with a history of prolonged cumulative sun exposure was referred with a three-year history of a conjunctival lesion in the right eye, originally diagnosed clinically as pterygium, which recently became more pigmented and uncomfortable. Her past ocular history included a recent episode of orbital cellulitis. Past medical history included nephrectomy for pyelonephritis and appendectomy. There was no family history of cancer.

On examination, the visual acuity was 6/6 bilaterally. Intraocular pressures were normal. The left eye was healthy. The right eye showed a temporal, perilimbal tumor with basal dimensions of 5 mm transversely and 3 mm longitudinally. The lesion was flat and mobile, without adhesion to the underlying Tenon's capsule and sclera, with conspicuous intrinsic vasculature and dilated feeder vessels. The tumor was pale but pigment dusting was present in the corneal epithelium from the 8 to 10 o'clock meridians (Fig 1b). The rest of the eye examination was unremarkable.

Case 3

A 64-year-old Caucasian male had a past ocular history of left ocular trauma with a wire injury 48 years before which resulted in a red conjunctival mark. Over the preceding three months there was an increase in size of the mark with mild discomfort. Medical history included prostate cancer treated by radiotherapy.

On examination, visual acuities were 6/18 OD and 6/9 OS. The intraocular pressures and right eye were within normal limits. In the left eye there was a large temporal conjunctival tumor abutting the limbus, measuring 14 mm transversely and 8 mm longitudinally. It was bi-lobed, pink, elevated, with prominent intrinsic and extrinsic vasculature and a feeder vessel between the two lobes of the lesion. The conjunctival component was mobile, with no areas of scleral invasion seen, while the corneal component was fixed and presented as an opaque vascularised infiltration from the 1 to 7 o'clock

meridians, extending into cornea for 3 mm superiorly and 5 mm inferiorly (Fig 1c). A papilloma was observed on the upper lid.

Case 4

A 72-year-old Caucasian male presented with a two-year history of a conjunctival lesion in his left eye, enlarged over the last year. The lesion was initially treated as a non-resolving episcleritis until recent review when a suspicion of ocular surface squamous neoplasia arose and then was referred to our service. There was no history of prolonged UV radiation exposure. His medical history included diabetes mellitus, hypertension and myocardial infarction. Family history was positive for uterine cancer (mother).

On examination, the visual acuity was 6/6 bilaterally. The intraocular pressures were normal. The right eye examination was unremarkable. Examination of the left eye revealed a gelatinous, infero-temporal perilimbal conjunctival lesion, with basal dimensions measuring 8 mm transversely and 5 mm longitudinally, non-pigmented except in the portion encroaching the cornea for 1 mm which contained some pigmentation. The conjunctival tumor was freely mobile and without invasion of the deeper layers. Three large feeder vessels and a prominent intrinsic tumor vascularity were observed (Fig 1d).

The differential diagnosis in these 4 cases included conjunctival intraepithelial neoplasia and squamous cell carcinoma, conjunctival papilloma (cases 1 and 3), pyogenic granuloma (case 1) and chronic inflammatory sun-related conjunctival damage (case 2). All patients underwent a conjunctival excisional biopsy with corneal alcohol epitheliectomy and double freeze-thaw cryotherapy applied to the conjunctival edges and base of the excised area.

RESULTS

Histopathology examination revealed invasive conjunctival malignant melanoma and melanosis with severe atypia, both incompletely excised (pT1a, pNx, pMx; TNM8) in all 4 cases. In patients 1, 2 and 4 the tumor cells were predominantly of epithelioid type, while in patient 3 they were mainly of spindle type (Fig 2 a-h) . Immunohistochemistry was positive for MelanA and HMB45 (Supplementary material).

All patients underwent adjuvant treatment with strontium-90 β plaque radiotherapy and patient 2 also received 3 cycles of topical Mitomycin C 0.04% four times a day. All 4 patients are in remission without local or systemic relapse (median follow-up time 16.5 months, range 10-103 months).

DISCUSSION

Conjunctival melanoma is a rare ocular surface cancer, which can be fatal. Pattern recognition for early referral to a specialist ocular oncology centre is essential for appropriate management. Complete excision in a dry surgical field, with minimal handling “no touch” excision of lesional tissue and cryotherapy is performed to limit recurrence and metastasis (8). Clinical features on presentation are heterogeneous. In 70% of cases CM presents as a pigmented thickened conjunctival tumour with enlarged feeder vessels, while in the remaining cases there is a minimally elevated and non-pigmented or mildly pigmented lesion, thus making clinical diagnosis challenging.

Conjunctival SCC is an epithelial malignant tumor in the spectrum of ocular surface squamous neoplasia (OSSN) and its main differential diagnosis includes conjunctival intraepithelial neoplasia, papilloma and granuloma. It usually presents as a gelatinous non-pigmented vascularized mass, located more frequently in the interpalpebral area and with possible infiltration of the adjacent corneal epithelium. OSSN can masquerade as CM particularly in dark-skinned individuals where the tumour takes up melanin to become a pigmented epibulbar mass (9, 10). In this report we highlight the

opposite scenario, where tumours with the clinical appearance of suspected OSSN were found to be CM on histopathological diagnosis.

The cases reported herein had clinical features highly suspicious for OSSN such as the absence of pigmentation in two patients (cases 1 and 3), mild pigmentation in two patients (cases 2 and 4), papillomatous architecture (case 3), cork-screw intrinsic vascularity (case 1), and no deep fixity of the conjunctival component (all cases). All were located at the limbal or perilimbal location, a common feature in OSSN as well as CM.

Features other than colour and tumour location that help differentiate CM from OSSN include the lack of keratinization and leukoplakia in CM. However, there is a clinical overlap between these two conjunctival tumors, hence histopathology remains the gold standard for the diagnosis, as highlighted in the 8th American Joint Committee Cancer (AJCC) Staging Manual (11). Other less invasive diagnostic methods including impression cytology, vital dyes, in vivo confocal microscopy, anterior segment-OCT are also used in some centres (12, 13). However, currently, histopathology is superior in reporting tissue architecture at the macroscopic and microscopic level. Vital dyes like toluidine blue 0.05% and methylene blue 1% are useful as screening but not diagnostic tools for ocular surface squamous neoplasia, as they have a high sensitivity (92% for toluidine blue 0.05%, 97% for Methylene blue 1%) but a low specificity (from 31% to 60% for toluidine blue 0.05%; 50% for Methylene blue 1%)(13, 14). Rose Bengal and lissamine green have also been used. AS-OCT is a non-invasive imaging technique used in several fields of ophthalmology (15, 16) . For conjunctival tumours, it is useful for directing diagnosis and management, as it provides high quality images of the internal structure and extension (12). If AS-OCT had been performed in our cases, it would have helped direct the clinical diagnosis, though it does not yet replace histopathology and management would still have been tumour excision. As a diagnostic or incisional biopsy may lead to tumour

seeding, our approach is complete excision with a 3 mm margin and cryotherapy to prevent recurrence in such cases.

These four cases of CM with atypical presentation shared some clinical features, such as perilimbal location, the lack of pigmentation and light iris colour. PAM, which raises the index of suspicion for CM, was absent in the clinical examination of the ocular surface, including double eversion of the upper eyelid. However, histopathological analysis of the excised tumours in all showed the presence of PAM with atypical melanocytes, highlighting again the importance of wide excision and histopathological correlation.

Recent studies showed that low-pigmented CM is seen more frequently in eyes with light iris color and is associated with a higher rate of recurrences, distant metastases and death. Thus, it has been hypothesized that a low tumor pigmentation is associated with worse outcomes (17), with a possible link between eumelanin/pheomelanin ratio and the grade of severity of CM. It is possible, however, that worse outcomes occur in pale tumours because they mimic other entities such as OSSN and definitive management is delayed.

There is a trend towards treatment of conjunctival squamous neoplasms without histopathological confirmation of the diagnosis (18,19). Our series is an example of conjunctival melanomas showing clinical features suggestive of squamous conjunctival neoplasia, serving as a warning against topical chemotherapy for suspected conjunctival carcinoma without histopathological confirmation of the diagnosis.

Ophthalmologists need to be aware that surgical clearance and histopathological analysis remains the gold standard for the diagnosis of CM especially in such cases where CM masquerades as OSSN, to avoid erroneous management that can adversely affect the prognosis.

CONTRIBUTORS

Study concept and design: BG and MSS. Acquisition, analysis or interpretation of data: all authors.

Drafting of the manuscript and critical revision for important intellectual content: all authors. Final approval of the version to be published: all authors.

FUNDING

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

ACKNOWLEDGEMENT

The research was supported by the National Institute for Health Research (NIHR) Biomedical Research Centre based at Moorfields Eye Hospital NHS Foundation Trust and UCL Institute of Ophthalmology. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health.

COMPETING INTERESTS

None declared.

REFERENCES

1. Seregard S. Conjunctival melanoma. *Surv Ophthalmol* 1998;42(4):321-350.
2. Yu G P, Hu D N, McCormick S, Finger P T. Conjunctival melanoma: is it increasing in the United States? *Am J Ophthalmol* 2003;135:800-6.
3. Tuomaala S, Kivela T. Conjunctival melanoma: is it increasing in the United States (letter)? *Am J Ophthalmol* 2003;136:1189-1190; author reply.

4. Tuomala S, Eskelin S, Tarkkanen A, Kivela T. Population-based assessment of clinical characteristics predicting outcome of conjunctival melanoma in whites. *Invest Ophthalmol Vis Sci* 2002;43:3399-3408.
5. Vasalaki M, Fabian I D, Reddy M A, Cohen V M L, Sagoo M S. Ocular oncology: advances in retinoblastoma, uveal melanoma and conjunctival melanoma. *British Medical Bulletin* 2017;1-13. doi 10.1093/bmb/ldw053.
6. Shields C L, Markowitz J S, Belinsky I, Schwartzstein H, George N S, Lally S E *et al.* Conjunctival Melanoma: outcomes based on tumor origin in 382 consecutive cases. *Ophthalmology* 2011; 118:389-95. doi 10.1016/j.opthta.2010.06.021.
7. Shields C L, Alset A E, Boal N S, Casey M G, Knapp A N, Sugarman J A *et al.* Conjunctival tumors in 5002 cases. Comparative analysis of benign versus malignant counterparts. The 2016 James D. Allen Lecture. *Am J Ophthalmol* 2017; 173:106-133. doi: 10.1016/j.ajo.2016.09.034.
8. Shields J A, Shields C L, De Potter P. Surgical management of conjunctival tumors. The 1994 Lynn B. McMahan lecture. *Arch Ophthalmol* 1997;115:808-815.
9. Kenawy N, Lake S L, Coupland S E, Damato B E. Conjunctival melanoma and melanocytic intraepithelial neoplasia. *Eye* 2013; 27:142-152. doi10.1038/eye.2012.254.
10. Gichuhi S, Macharia E, Kabiru J, Zindamoyen AM, Rono H, Ollando E *et al.* Clinical Presentation of Ocular Surface Squamous Neoplasia in Kenya. *JAMA Ophthalmol* 2015;133(11):1305-1313. doi: 10.1001/jamaophthLMOL.2015.3335.
11. Amin M B, Edge S, Greene F. *AJCC Cancer Staging Manual 8th edition*. Springer 2017.
12. Nanji A A, Mercado C, Galor A, Dubovy S, Karp C L. Updates in ocular surface tumor diagnostics. *Int Ophthalmol Clin* 2017;57 (3):47-62.
13. Gichuhi S, Macharia E, Kabiru J, Zindamoyen A M, Rono H, Ollando E *et al.* Toluidine Blue 0.05% vital staining for the diagnosis of ocular surface squamous neoplasia in Kenya. *JAMA Ophthalmol* 2015;133:1314-21. doi: 10.1001/jamaophthalmol.2015.3345.

14. Steffen J, Rice J, Lecuona K, Carrara H. Identification of ocular surface squamous neoplasia by in vivo staining with methylene blue. *Br J Ophthalmology* 2014; 98:13-15. doi: 10.1136/bjophthalmol-2013-303956.
15. Vajzovic L M, Karp C L, Haft P, Shousha M A, Dubovy S R, Hurmeric V et al. Ultra high-resolution anterior segment optical coherence tomography in the evaluation of anterior corneal dystrophies and degenerations. *Ophthalmol* 2011; 118(7):1291–1296. doi: 10.1016/j.optha.2010.12.015.
16. Ramos J L, Li Y, Huang D. Clinical and research applications of anterior segment optical coherence tomography - A review. *Clin Exp Ophthalmol* 2009; 37(1):81–89. doi: 10.1111/j.1442-9071.2008.01823.x.
17. Brouer N J, Marinkovic M, Luyten G P M, Shields C L, Jager M J. Br. Lack of tumour pigmentation in conjunctival melanoma is associated with light iris colour and worse prognosis. *Br J Ophthalmology* 2018;0:1-6. doi:10.1136/bjophthalmol-2018-312018.
18. Chaugule S S, Park J, Finger P T. Topical chemotherapy for giant ocular surface squamous neoplasia: is surgery necessary? *Indian J Ophthalmol* 2018;66(1):55-60. doi: 10.4103/ijo.IJO_590_17.
19. Polski A, Saber M S, Kim J W, Berry J L. Extending far and wide: the role of biopsy and staging in the management of ocular surface squamous neoplasia. *Clin Exp Ophthalmol* 2018;1-8. doi: 10.1111/ceo.13382.

FIGURE 1a-d

FIGURE 2 a-h

SUPPLEMENTARY MATERIAL

FIGURE 1 LEGENDS

Figure 1 a: Case 1 - colour anterior segment photograph of the left eye of a 66-year-old female, showing a temporal bulbar conjunctival lesion crossing the limbus for 1 mm. The lesion measured 5 mm x 5.5 mm, was fleshy, had 4 feeder vessels and cork-screw intrinsic vessels. Note the absence of pigmentation.

Figure 1 b: Case 2 - colour anterior segment photograph of a 41-year-old female right eye showing a temporal bulbar conjunctival lesion encroaching the limbus for 2 mm. The lesion measured 3 mm x 5 mm, had 3 dilated feeder vessels; the conjunctival portion was non-pigmented while the corneal portion had a mildly pigmented foamy infiltration.

Figure 1 c: Case 3 - colour anterior segment photograph of the left eye of a 64-year-old male, with a pink-yellow temporal and inferior perilimbal lesion encroaching the limbus for 2 mm superiorly and 5 mm inferiorly, measuring 14 mm x 8 mm. Note the bi-lobed aspect and the corneal epithelial infiltration by fine vessels resembling a conjunctival papilloma.

Figure 1 d: Case 4 – colour anterior segment photograph of the left eye of a 72-year-old man, with an infero-temporal perilimbal lesion, non-pigmented except at the limbus for 1 mm where some faint pigmentation was observed, measuring 8 mm x 5 mm.

FIGURE 2 LEGENDS

Figure 2 a-b: Case 1 - Photomicrograph of conjunctiva showing loose nests of atypical epithelioid melanocytes within the stroma as well as the surface epithelium. There is minimal melanin.

Figure 2 c-d: Case 2 - Photomicrograph of conjunctiva showing nests of atypical cells within the epithelium and stroma. The latter is associated with a chronic inflammatory infiltrate. There is little melanin.

Figure 2 e-f: Case 3 - Photomicrograph of conjunctiva showing large nests of spindle cells within the epithelium and stroma, accompanied by a modest chronic inflammatory infiltrate.

Figure 2 g-h: Case 4 - Photomicrograph of conjunctiva showing flattened detached epithelium overlying stroma infiltrated by sheets of atypical epithelioid melanocytes and a modest chronic inflammatory infiltrate.

SUPPLEMENTARY MATERIAL LEGENDS

a: Case 4 – Immunohistochemistry for MelanA highlights the extent of tumour infiltration within the stroma.

b: Case 4 – HMB45 immunohistochemistry is positive in tumour cells within both the superficial and deep stroma.