

Risk of COVID-19 among front-line health-care workers and the general community: a prospective cohort study



Long H Nguyen*, David A Drew*, Mark S Graham*, Amit D Joshi, Chuan-Guo Guo, Wenjie Ma, Raaj S Mehta, Erica T Warner, Daniel R Sikavi, Chun-Han Lo, Sohee Kwon, Mingyang Song, Lorelei A Mucci, Meir J Stampfer, Walter C Willett, A Heather Eliassen, Jaime E Hart, Jorge E Chavarro, Janet W Rich-Edwards, Richard Davies, Joan Capdevila, Karla A Lee, Mary Ni Lochlainn, Thomas Varsavsky, Carole H Sudre, M Jorge Cardoso, Jonathan Wolf, Tim D Spector, Sebastien Ourselin†, Claire J Steves†, Andrew T Chan†, on behalf of the Coronavirus Pandemic Epidemiology Consortium‡

Summary

Background Data for front-line health-care workers and risk of COVID-19 are limited. We sought to assess risk of COVID-19 among front-line health-care workers compared with the general community and the effect of personal protective equipment (PPE) on risk.

Methods We did a prospective, observational cohort study in the UK and the USA of the general community, including front-line health-care workers, using self-reported data from the COVID Symptom Study smartphone application (app) from March 24 (UK) and March 29 (USA) to April 23, 2020. Participants were voluntary users of the app and at first use provided information on demographic factors (including age, sex, race or ethnic background, height and weight, and occupation) and medical history, and subsequently reported any COVID-19 symptoms. We used Cox proportional hazards modelling to estimate multivariate-adjusted hazard ratios (HRs) of our primary outcome, which was a positive COVID-19 test. The COVID Symptom Study app is registered with ClinicalTrials.gov, NCT04331509.

Findings Among 2 035 395 community individuals and 99 795 front-line health-care workers, we recorded 5545 incident reports of a positive COVID-19 test over 34 435 272 person-days. Compared with the general community, front-line health-care workers were at increased risk for reporting a positive COVID-19 test (adjusted HR 11·61, 95% CI 10·93–12·33). To account for differences in testing frequency between front-line health-care workers and the general community and possible selection bias, an inverse probability-weighted model was used to adjust for the likelihood of receiving a COVID-19 test (adjusted HR 3·40, 95% CI 3·37–3·43). Secondary and post-hoc analyses suggested adequacy of PPE, clinical setting, and ethnic background were also important factors.

Interpretation In the UK and the USA, risk of reporting a positive test for COVID-19 was increased among front-line health-care workers. Health-care systems should ensure adequate availability of PPE and develop additional strategies to protect health-care workers from COVID-19, particularly those from Black, Asian, and minority ethnic backgrounds. Additional follow-up of these observational findings is needed.

Funding Zoe Global, Wellcome Trust, Engineering and Physical Sciences Research Council, National Institutes of Health Research, UK Research and Innovation, Alzheimer's Society, National Institutes of Health, National Institute for Occupational Safety and Health, and Massachusetts Consortium on Pathogen Readiness.

Copyright © 2020 The Author(s). Published by Elsevier Ltd. This is an Open Access article under the CC BY-NC-ND 4.0 license.

Introduction

Since its emergence, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), which causes COVID-19, has become a global health threat.¹ As of July 22, 2020, more than 15 million cases of COVID-19 have been documented worldwide, with nearly 618 000 deaths.² In the UK and the USA, Black, Asian, and minority ethnic communities have been disproportionately affected.^{3,4} With ongoing community transmission from asymptomatic individuals, disease burden is expected to rise. As a result, there will be an ongoing need for front-line health-care workers in patient-facing roles. Because this work requires close personal exposure to patients with SARS-CoV-2, front-line health-care workers are at high

risk of infection, contributing to further spread.⁵ Initial estimates suggest that front-line health-care workers could account for 10–20% of all diagnoses,^{6,7} with some early evidence that people from Black, Asian, and minority ethnic backgrounds are at higher risk.³

Based on experience with other respiratory viruses, consistent use of personal protective equipment (PPE) is important to reduce nosocomial transmission.⁸ Guidelines from the UK and the USA recommend mask use for health-care workers caring for people with COVID-19.^{9,10} However, global shortages of masks, respirators, face shields, and gowns, caused by surging demand and supply chain disruptions, have led to efforts to conserve PPE through extended use or reuse, and

Lancet Public Health 2020; 5: e475–83

Published Online
July 31, 2020
[https://doi.org/10.1016/S2468-2667\(20\)30164-X](https://doi.org/10.1016/S2468-2667(20)30164-X)

See [Comment](#) page e461

*Contributed equally as first authors

†Contributed equally

‡Members listed in the appendix

Division of Gastroenterology (L H Nguyen MD, D A Drew PhD, A D Joshi PhD, C-G Guo MS, W Ma ScD, R S Mehta MD, C-H Lo MD, S Kwon MD, M Song ScD, Prof A T Chan MD), **Clinical and Translational Epidemiology Unit** (E T Warner ScD, L H Nguyen, A D Drew, A D Joshi, C-G Guo, W Ma, R S Mehta, C-H Lo, S Kwon, M Song, Prof A T Chan), **Center on Genomics, Vulnerable Populations, and Health Disparities** (E T Warner), and **Department of Medicine** (D R Sikavi MD), **Massachusetts General Hospital and Harvard Medical School, Boston, MA, USA; Department of Biostatistics** (L H Nguyen, W Ma, R S Mehta), **Department of Epidemiology** (Prof L A Mucci ScD, Prof M J Stampfer MD, Prof W C Willett MD, A H Eliassen ScD, J E Chavarro MD, J W Rich-Edwards ScD, C-H Lo, M Song), **Department of Nutrition** (M Song, Prof W C Willett, J E Chavarro), **Department of Environmental Health** (J E Hart ScD), and **Department of Immunology and Infectious Disease** (Prof A T Chan), **Harvard T H Chan School of Public Health, Boston, MA, USA; School of Biomedical Engineering and Imaging Sciences** (M S Graham PhD, T Varsavsky MSc, C H Sudre PhD, M J Cardoso PhD,

Prof S Ourselin PhD), and Department of Twin Research and Genetic Epidemiology (K A Lee MBBCh, M N Lochlainn MBBCh, Prof T D Spector MD, C J Steves PhD), King's College London, London, UK; Department of Medicine, Li Ka Shing Faculty of Medicine, University of Hong Kong, Hong Kong Special Administrative Region, China (C-G Guo); Channing Division of Network Medicine (Prof M J Stampfer, J E Hart, J E Chavarro, A H Eliassen), and Division of Women's Health (J W Rich-Edwards), Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA, USA; Zoe Global, London, UK (R Davies MA, J Capdevila PhD, J Wolf MA); Broad Institute of Massachusetts Institute of Technology and Harvard, Cambridge, MA, USA (Prof A T Chan); and Massachusetts Consortium on Pathogen Readiness, Cambridge, MA, USA (Prof A T Chan)

Correspondence to: Prof Andrew T Chan, Massachusetts General Hospital and Harvard Medical School, Boston, MA 02114, USA achan@mgh.harvard.edu

See Online for appendix

Research in context

Evidence before this study

We searched PubMed for articles published between Jan 1 and June 30, 2020, with the terms "covid-19", "healthcare workers", and "personal protective equipment". We did not restrict our search by language or type of publication. The prolonged course of the COVID-19 pandemic, coupled with sustained challenges supplying adequate personal protective equipment (PPE) for front-line health-care workers, have strained global health-care systems in an unprecedented fashion. Despite growing awareness of this problem, there are few studies to inform policy makers on the risk of COVID-19 among health-care workers and the effect of PPE on disease burden. Previous reports of infections in health-care workers are based on cross-sectional data with limited information on individual-level risk factors. Our PubMed search yielded no population-scale investigations.

Added value of this study

We did a prospective observational study of 2 135 190 individuals, comprised of front-line health-care workers and the general community who were voluntary

disinfection protocols have been developed, for which scientific consensus on best practice is scarce.¹¹⁻¹³

Although addressing the needs of front-line health-care workers during the COVID-19 pandemic is a high priority,^{6,14} data to inform such efforts are scarce, and particularly so among Black, Asian, and minority ethnic communities. Thus, we did a prospective population-based study using data from a smartphone-based application (app) to investigate the risk of testing positive for COVID-19, the risk of developing symptoms associated with SARS-CoV-2 infection, or both, among individuals in the UK and the USA.

Methods

Study design and participants

COVID Symptom Study (previously known as COVID Symptom Tracker) is a free smartphone app developed by Zoe Global (London, UK) in collaboration with Massachusetts General Hospital (Boston, MA, USA) and King's College London (London, UK). It offers participants a guided interface to report baseline demographic information and comorbidities, daily information on symptoms, and COVID-19 testing. Participants are encouraged to log daily, even when asymptomatic, for longitudinal collection of incident symptoms and COVID-19 testing results. The app was launched in the UK on March 24, 2020, and the USA on March 29, 2020.

We did a prospective, observational cohort study using the COVID Symptom Study app. Participants were recruited through social media outreach and invitations from the investigators of long-running cohort studies to their volunteers (appendix p 3). At enrolment, participants consented to use of information for research and

users of the COVID Symptom Study smartphone application (app). From self-reported data obtained via this app, we found that front-line health-care workers had at least a threefold increased risk of COVID-19. Compared with front-line health-care workers who reported adequate availability of PPE, those with inadequate PPE had an increase in risk. However, adequate availability of PPE did not seem to completely reduce risk among health-care workers caring for patients with COVID-19. We also found that Black, Asian, and minority ethnic health-care workers might be disproportionately affected.

Implications of all the available evidence

Front-line health-care workers, particularly those who are from Black, Asian, and minority ethnic backgrounds, could be at substantially greater risk of COVID-19. Health-care systems should ensure adequate availability of PPE and develop additional strategies to protect health-care workers from COVID-19.

agreed to applicable privacy policies and terms of use. Our study was approved by the Partners Human Research Committee (protocol 2020P000909) and King's College London ethics committee (REMAS ID 18210, LRS-19/20-18210).

Procedures

Information obtained through the COVID Symptom Study app has been described in detail.¹⁵ Briefly, on first use, participants were asked to provide demographic factors and were questioned separately about a series of COVID-19 risk factors (appendix pp 4-8). At enrolment and with daily reminders, participants were asked if they felt physically normal, and if they reported not feeling well they were asked about their symptoms (appendix p 11). Participants were also asked if they had been tested for COVID-19 and the results (none, negative, pending, or positive). Our primary outcome was a report of a positive COVID-19 test. Follow-up started when participants first reported on the COVID Symptom Study app and continued until a report of a positive COVID-19 test or the time of last data entry, whichever occurred first.

Participants were also asked if they worked in health care and, if yes, whether they had direct patient contact. For our primary analysis, we defined front-line health-care workers as participants who reported direct patient contact. Prespecified secondary analyses among front-line health-care workers investigated PPE availability and contact with patients with COVID-19, as well as the primary site of clinical practice. A post-hoc analysis among front-line health-care workers assessed the effect of race and ethnicity. Beginning March 29, 2020, in an updated version of the app used by 84% of participating health-care workers, we included mandatory questions

about availability of PPE for participants who identified as a front-line health-care worker. Among these individuals, we asked whether they cared for patients with suspected or documented SARS-CoV-2 infection and the frequency with which they used PPE (always, sometimes, or never). We asked if they had enough PPE when needed, if they had to reuse PPE, or if they did not have enough because of shortages. We classified PPE as adequate if they never required PPE or if they reported always having the PPE they needed. We classified PPE as inadequate if they reported they did not have enough PPE or if it was not available. We also asked health-care workers to report the site of their patient care (inpatient, nursing home, outpatient, home health, ambulatory clinic, or other).

Statistical analysis

We used standard calculations to determine the minimum detectable hazard ratio (HR) for our primary categorical exposure (health-care worker status) and risk of reporting a positive COVID-19 test with 80% power.¹⁶

We used Cox proportional hazards modelling stratified by age, date, and country to estimate multivariable-adjusted HRs and 95% CIs. Covariates were selected a priori based on putative risk factors, including sex (male or female), race or ethnic origin (non-Hispanic white, Hispanic or Latinx, Black, Asian, or more than one or other), history of diabetes (yes or no), heart disease (yes or no), lung disease (yes or no), kidney disease (yes or no), current smoking status (yes or no), and body-mass index (17.0–19.9 kg/m², 20.0–24.9 kg/m², 25.0–29.9 kg/m², and ≥30.0 kg/m²). Data imputation replaced no more than 5% of missing values for a given metadatum, with numerical values replaced with the median and categorical variables imputed using the mode.

Because our primary outcome (positive COVID-19 test) required a participant to receive a test, we did several prespecified secondary analyses. First, we tested a symptom-based classifier predictive of positive COVID-19 testing.¹⁷ Briefly, using logistic regression and symptoms preceding testing, we have previously described that loss of smell or taste, fatigue, persistent cough, and loss of appetite predicted COVID-19 positivity with high specificity (appendix p 2). Second, to account for country-specific predictors of obtaining testing, we did separate inverse probability weighting in the UK and the USA as a function of demographic and clinical factors, followed by inverse probability-weighted Cox proportional hazards modelling stratified by age and date with additional adjustment for the covariates used in previous models (appendix p 2). To assess factors associated with PPE adequacy, work in especially high-risk clinical settings, or greater exposure to patients with COVID-19 (including a post-hoc analysis of race and ethnicity), we used multivariable logistic regression models to estimate adjusted odds ratios (ORs) and 95% CIs. Two-sided

p values less than 0.05 were considered statistically significant. All analyses were done using R version 3.6.1.

The COVID Symptom Study app is registered with ClinicalTrials.gov, NCT04331509.

Role of the funding source

Zoe Global developed the app for data collection. The funders had no role in study design, data analysis, data interpretation, or writing of the report. LHN, DAD, MSG, SO, CJS, and ATC had access to raw data. The corresponding author had full access to all data in the study and had final responsibility for the decision to submit for publication.

Results

Between March 24 and April 23, 2020, we enrolled 2810103 consecutive users of the COVID Symptom Study app to our study. 2627695 participants in the UK and 182408 in the USA provided baseline information about feeling physically normal or having symptoms (appendix p 14). 134885 (4.8%) participants reported being a front-line health-care worker. The prevalence of COVID-19 was 2747 cases per 100000 front-line health-care workers compared with 242 cases per 100000 people in the general community (figure). The highest infection rates were reported in the US states New York, New Jersey, and Louisiana and in areas around London and the Midlands in the UK.

After excluding 670298 participants with less than 24 h of follow-up and 4615 individuals who reported a positive COVID-19 test at baseline, we included 2135190 participants in our prospective inception cohort, of whom 99795 (4.7%) identified as front-line health-care workers (appendix p 14). Based on this cohort, we had 80% power to detect a minimum HR of 1.16 for risk of reporting a positive COVID-19 test between health-care workers and the general community. In this cohort, we recorded 24.4 million entries, or 11.5 logs per participant, with median follow-up of 18.9 days (IQR 5.1–26.1). Median age was 44 years (IQR 32–57). Compared with the general community, front-line health-care workers were more frequently female, had a slightly higher prevalence of body-mass index 30.0 kg/m² or higher, were slightly more likely to smoke (particularly in the UK), and were more likely to use non-steroidal anti-inflammatory drugs (table 1; appendix pp 9–10). At baseline, 20.2% of front-line health-care workers reported at least one symptom associated with SARS-CoV-2 infection compared with 14.4% of the general population; fatigue, loss of smell or taste, and hoarse voice were especially frequent (appendix p 11). When comparing health-care workers who were asked about PPE use with those who were not asked, no difference was noted in baseline factors including age (median 45 years vs 40 years), female sex (81% vs 82%), or body-mass index (median 25.9 kg/m² vs 25.7 kg/m²).

We recorded 5545 incident reports of positive COVID-19 testing over 34435272 person-days. In the UK, 1.1% of

For more on R see
<https://r-project.org>




Figure: Risk of testing positive for COVID-19 among front-line health-care workers (A) Between March 24 and April 23, 2020, considerable disparities were noted in prevalence of a positive COVID-19 test among front-line health-care workers compared with the general community, in both the UK and the USA. (B) Prevalence of a positive COVID-19 test reported by front-line health-care workers in the UK and the USA. Regions in grey did not have sufficient data for analysis. app=COVID-19 Symptom Study smartphone application.

	Front-line health-care workers (n=99 795)	General community (n=2 035 395)
Country		
UK	85.4%	93.9%
USA	14.6%	6.1%
Age, years	42 (33–53)	44 (33–56)
<25	4.5%	4.7%
25–34	24.7%	19.2%
35–44	25.1%	21.5%
45–54	23.6%	19.5%
55–64	17.5%	16.2%
≥65	3.9%	13.1%
Missing data for age	1.1%	5.7%
Sex		
Male	17.0%	37.0%
Female	83.0%	63.0%
Race or ethnic origin*		
Non-Hispanic white	88.2%	92.5%
Hispanic or Latinx	1.1%	0.5%
Black	1.2%	0.6%
Asian	4.4%	2.2%
More than one or other	2.4%	2.9%
Missing data for race or ethnic origin, or prefer not to say	2.7%	1.3%

(Table 1 continues in next column)

	Front-line health-care workers (n=99 795)	General community (n=2 035 395)
(Continued from previous column)		
Body-mass index (kg/m ²)	25.8 (22.8–30.2)	25.3 (22.5–29.1)
17.0–19.9	5.8%	8.3%
20.0–24.9	38.1%	39.2%
25.0–29.9	30.1%	31.5%
≥30.0	25.9%	21.1%
Missing data for body-mass index	0.5%	0.5%
Comorbidities		
Diabetes	2.5%	3.1%
Heart disease	1.6%	2.4%
Lung disease	13.1%	12.2%
Kidney disease	0.6%	0.7%
Cancer	0.5%	1.3%
Missing data for cancer	0.3%	0.3%
Pregnant (% of females)	0.9%	1.0%
Medication use		
Non-steroidal anti-inflammatory drugs	8.2%	6.1%
Immunosuppressants	2.5%	3.2%
Chemotherapy or immunotherapy	0.1%	0.3%
Angiotensin-converting enzyme inhibitor	5.0%	4.9%
Missing data for angiotensin-converting enzyme inhibitor	10.1%	4.3%
Current smoking		
Yes	10.2%	8.5%
Missing data for smoking status	0.2%	0.1%

Data are % or median (IQR). % are calculated based on the total number of participants with available data. Polytomous variables might not add up to 100% because of rounding. Questions about history of cancer, angiotensin-converting enzyme inhibitor use, and smoking status have been asked since launch in the USA and March 29, 2020, in the UK; questions about race and ethnic origin were asked since April 17, 2020, in both the UK and the USA. Percentages within each category are based on the total population responding when the question was first asked. *Non-Hispanic white defined as UK White, US White, and no designation of other race or ethnic origin. Hispanic or Latinx designated as Hispanic and Latino. Black defined as UK Black, Black British, US Black, and African American. White defined as UK White and US White. Asian defined as UK Asian, Asian British, UK Chinese, Chinese British, US Asian, and US Native Hawaiian or other Pacific Islander. More than one or other defined as UK mixed race White and Black or Black British UK, mixed race other, UK Middle Eastern or Middle Eastern British, US American Indian or Alaska Native, other, and denoted more than one race.

Table 1: Baseline characteristics of front-line health-care workers compared with the general community

health-care workers reported being tested compared with 0.2% of the general community (health-care workers to community testing ratio 5.5), whereas 4.1% of US health-care workers were tested versus 1.1% of the general community (testing ratio 3.7). Compared with the general community, front-line health-care workers had a twelve-fold increase in risk of a positive test after multivariable

	Event/person-days	Incidence (30-day)	Age-adjusted hazard ratio (95% CI)	Multivariate-adjusted hazard ratio (95% CI)	Inverse probability-weighted hazard ratio (95% CI)
Overall (primary analysis)					
General community	3623/32 980 571	0.33%	1 (ref)	1 (ref)	1 (ref)
Front-line health-care worker	1922/1 454 701	3.96%	11.68 (10.99–12.40)	11.61 (10.93–12.33)	3.40 (3.37–3.43)
According to race or ethnic origin (post-hoc analysis)					
Non-Hispanic white, general community	1498/23 941 092	0.19%	1 (ref)	1 (ref)	1 (ref)
Black, Asian, and minority ethnic, general community	227/1 362 956	0.50%	2.49 (2.16–2.86)	2.51 (2.18–2.89)	1.74 (1.71–1.77)
Non-Hispanic white, front-line health-care worker	726/935 860	2.33%	12.47 (11.33–13.72)	12.58 (11.42–13.86)	3.52 (3.48–3.56)
Black, Asian, and minority ethnic, front-line health-care worker	98/72 556	4.05%	21.68 (17.61–26.68)	21.88 (17.78–26.94)	4.88 (4.76–5.01)

All models were stratified by 5-year age group, calendar date at study entry, and country. Multivariate risk factor models were adjusted for sex (male or female), history of diabetes (yes or no), heart disease (yes or no), lung disease (yes or no), kidney disease (yes or no), current smoking (yes or no), and body-mass index (17.0–19.9 kg/m², 20.0–24.9 kg/m², 25.0–29.9 kg/m², and ≥30 kg/m²). Black, Asian, and minority ethnic was defined among individuals who had race or ethnicity information and did not identify as non-Hispanic white.

Table 2: Risk of reporting a positive test for COVID-19 among front-line health-care workers compared with the general community

adjustment (adjusted HR 11.61, 95% CI 10.93–12.33; table 2; appendix p 15). The association seemed stronger in the UK (adjusted HR 12.52, 95% CI 11.77–13.31) compared with the USA (2.80, 2.09–3.75; $p_{\text{difference}} < 0.0001$; appendix p 12).

We considered the possibility that noted differences could be related to testing eligibility. A multivariable-adjusted Cox proportional hazards model with inverse probability weighting for predictors of testing also showed a higher risk of infection among front-line health-care workers (adjusted HR 3.40, 95% CI 3.37–3.43; table 2), which was higher in the UK (3.43, 3.18–3.69) than in the USA (1.97, 1.36–2.85; $p_{\text{difference}} < 0.0001$; appendix p 12). In a prespecified secondary analysis, a validated model was used based on a combination of symptoms predictive of COVID-19 infection.¹⁶ Compared with the general community, health-care workers initially free of symptoms had an increased risk of predicted COVID-19 (adjusted HR 2.05, 95% CI 1.99–2.10), which was higher in the UK (2.09, 2.02–2.15) than in the USA (1.31, 1.14–1.51; $p_{\text{difference}} < 0.0001$).

In a post-hoc analysis, compared with individuals in the general community from a non-Hispanic white ethnic background, the risk for a positive COVID-19 test was increased for individuals from Black, Asian, and minority ethnic backgrounds in the general community (adjusted HR 2.51, 95% CI 2.18–2.89), for Black, Asian, and minority ethnic health-care workers (21.88, 17.78–26.94), and for non-Hispanic white health-care workers (12.58, 11.42–13.86; table 2). In post-hoc analyses, the association of race and health-care worker status with risk of COVID-19 was assessed. Black, Asian, and minority ethnic health-care workers had an increased risk of COVID-19 (adjusted HR 1.81, 95% CI 1.45–2.24) compared with non-Hispanic white health-care workers.

	Adequate PPE	Reused PPE	Inadequate PPE
Overall			
Event/person-days	592/332 901	146/80 728	157/60 916
Unadjusted hazard ratio (95% CI)	1 (ref)	1.46 (1.21–1.76)	1.32 (1.10–1.57)
Multivariate-adjusted hazard ratio (95% CI)	1 (ref)	1.46 (1.21–1.76)	1.31 (1.10–1.56)
No exposure to patients with COVID-19			
Event/person-days	186/227 654	19/37 599	48/35 159
Unadjusted hazard ratio (95% CI)	1 (ref)	0.96 (0.60–1.55)	1.53 (1.11–2.11)
Multivariate-adjusted hazard ratio (95% CI)	1 (ref)	0.95 (0.59–1.54)	1.52 (1.10–2.09)
Exposure to patients with suspected COVID-19			
Event/person-days	126/54 676	36/19 378	26/14 083
Unadjusted hazard ratio (95% CI)	2.40 (1.91–3.02)	3.23 (2.24–4.66)	1.87 (1.24–2.83)
Multivariate-adjusted hazard ratio (95% CI)	2.39 (1.90–3.00)	3.20 (2.22–4.61)	1.83 (1.21–2.78)
Exposure to patients with documented COVID-19			
Event/person-days	280/50 571	91/23 751	83/11 675
Unadjusted hazard ratio (95% CI)	4.93 (4.07–5.97)	5.12 (3.94–6.64)	5.95 (4.57–7.76)
Multivariate-adjusted hazard ratio (95% CI)	4.83 (3.99–5.85)	5.06 (3.90–6.57)	5.91 (4.53–7.71)

All models were stratified by 5-year age group, calendar date at study entry, and country. Multivariate risk factor models were adjusted for sex (male or female), history of diabetes (yes or no), heart disease (yes or no), lung disease (yes or no), kidney disease (yes or no), current smoking (yes or no), and body-mass index (17.0–19.9 kg/m², 20.0–24.9 kg/m², 25.0–29.9 kg/m², and ≥30.0 kg/m²). PPE=personal protective equipment.

Table 3: Risk of reporting a positive test for COVID-19, according to availability of PPE and exposure to patients with COVID-19 among front-line health-care workers (prespecified secondary analysis)

Risk estimates were similar among male (adjusted HR 14.02, 95% CI 12.38–15.82) compared with female (11.27, 10.53–12.14) front-line health-care workers.

Among front-line health-care workers, availability and use of PPE, COVID-19 patient exposures, and subsequent risk for testing positive were assessed in prespecified analyses. Compared with health-care workers who reported adequate PPE, front-line health-care workers

reporting PPE reuse had an increased risk of a positive COVID-19 test (adjusted HR 1.46, 95% CI 1.21–1.76), with inadequate PPE associated with a comparable increase in risk after multivariable adjustment (1.31, 1.10–1.56; table 3).

In a prespecified secondary analysis, front-line health-care workers with inadequate PPE caring for patients with documented COVID-19 had an increased risk for COVID-19 after multivariable adjustment (adjusted HR 5.91, 95% CI 4.53–7.71) compared with those with adequate PPE not caring for patients with suspected or documented COVID-19 (table 3). Risk for front-line health-care workers reusing PPE and exposed to patients with documented COVID-19 was also increased (adjusted HR 5.06, 95% CI 3.90–6.57). Notably, even among

front-line health-care workers reporting adequate PPE, the risk for COVID-19 was increased for those caring for patients with suspected COVID-19 (adjusted HR 2.39, 95% CI 1.90–3.00) and for those caring for patients with documented COVID-19 (4.83, 3.99–5.85), compared with health-care workers who did not care for either group (table 3).

In a post-hoc analysis, differences were noted in PPE adequacy according to race and ethnicity, with non-white health-care workers more frequently reporting reuse of or inadequate access to PPE, even after adjusting for exposure to patients with COVID-19 (adjusted OR 1.49, 95% CI 1.36–1.63; table 4).

In a prespecified secondary analysis, risk of COVID-19 by practice location was assessed. Compared with risk for the general community, risk for front-line health-care workers was increased in all health-care settings, but was highest for those working in inpatient settings (adjusted HR 24.30, 95% CI 21.83–27.06) and nursing homes (16.24, 13.39–19.70; table 5). Notably, health-care workers in nursing homes most frequently (16.9%) reported inadequate PPE, whereas inpatient providers reported reuse of PPE most often (23.7%; table 5). In a post-hoc analysis, compared with non-Hispanic white health-care workers, Black, Asian, and minority ethnic health-care workers were more likely to work in higher risk clinical settings, including inpatient hospital or nursing homes (adjusted OR 1.13, 95% CI 1.03–1.23) and to care for patients with suspected or documented COVID-19 (1.20, 1.09–1.30). These noted differences were most pronounced among Black health-care workers (appendix p 13).

Discussion

Among 2135190 people in the UK and USA using the COVID-19 Symptom Study app between March 24 and April 23, 2020, we noted that front-line health-care workers had at least a threefold increased risk of reporting a positive COVID-19 test and predicted COVID-19 infection, compared with the general community, even after

	Health-care workers reporting reuse of or inadequate PPE	Multivariate-adjusted odds ratio (95% CI)
Overall		
Non-Hispanic white, front-line health-care worker	27.7%	1 (ref)
Black, Asian, and minority ethnic, front-line health-care worker	36.7%	1.49 (1.36–1.63)
According to racial or ethnic subgroup*		
Non-Hispanic white`	27.7%	1 (ref)
Hispanic or Latinx	49.6%	2.64 (2.03–3.45)
Black	33.5%	1.30 (1.02–1.65)
Asian	35.6%	1.42 (1.24–1.63)
More than one race or other race	34.7%	1.33 (1.12–1.57)

Multivariate risk factor models were adjusted for 5-year age group, sex, and exposure to patients with COVID-19 (none, suspected, and documented). Black, Asian, and minority ethnic was defined among individuals who had race or ethnicity information and did not identify as non-Hispanic white. PPE=personal protective equipment. *Non-Hispanic white defined as UK White, US White, and no designation of other race or ethnic origin. Hispanic or Latinx designated as Hispanic and Latino. Black defined as UK Black, Black British, US Black, and African American. White defined as UK White and US White. Asian defined as UK Asian, Asian British, UK Chinese, Chinese British, US Asian, and US Native Hawaiian or other Pacific Islander. More than one or other defined as UK mixed race White and Black or Black British UK, mixed race other, UK Middle Eastern or Middle Eastern British, US American Indian or Alaska Native, other, and denoted more than one race.

Table 4: Risk of reporting PPE inadequacy or reuse among front-line health-care workers, according to race or ethnic origin (post-hoc analysis)

	Event/ person-days	Incidence (30-day)	Age-adjusted hazard ratio (95% CI)	Multivariate-adjusted hazard ratio (95% CI)	Health-care workers reporting reuse of PPE	Health-care workers reporting inadequate PPE
General community	3623/32 980 571	0.33%	1 (ref)	1 (ref)
Front-line health-care worker						
Inpatient	564/184 293	9.18%	23.58 (21.20–26.25)	24.30 (21.83–27.06)	23.7%	11.9%
Nursing homes	118/52 901	6.69%	16.48 (13.60–19.97)	16.24 (13.39–19.70)	15.4%	16.9%
Outpatient hospital clinics	51/45 217	3.38%	10.75 (8.10–14.27)	11.21 (8.44–14.89)	16.3%	12.2%
Home health sites	36/38 642	2.79%	7.79 (5.58–10.87)	7.86 (5.63–10.98)	14.7%	15.9%
Ambulatory clinics	44/66 408	1.99%	6.64 (4.90–9.01)	6.94 (5.12–9.41)	19.3%	11.8%
Other	73/64 310	3.41%	9.42 (7.42–11.96)	9.52 (7.49–12.08)	12.0%	13.8%

Model was stratified by 5-year age group, calendar date at study entry, and country and adjusted for sex (male or female), history of diabetes (yes or no), heart disease (yes or no), lung disease (yes or no), kidney disease (yes or no), current smoking (yes or no), and body-mass index (17.0–19.9 kg/m², 20.0–24.9 kg/m², 25.0–29.9 kg/m², and ≥30.0 kg/m²). Ambulatory clinics include free-standing (non-hospital) primary care or specialty clinics and school-based clinics. PPE=personal protective equipment.

Table 5: Front-line health-care workers and risk of testing positive for COVID-19, by site of care delivery (prespecified secondary analysis)

accounting for other risk factors. Post-hoc analyses showed that Black, Asian, and minority ethnic health-care workers are at especially high risk of SARS-CoV-2 infection, with at least a fivefold increased risk of COVID-19 compared with the non-Hispanic white general community. Among front-line health-care workers, reuse of PPE or inadequate PPE were each associated with a subsequent increased risk of COVID-19. Although health-care workers caring for patients with COVID-19 who reported inadequate PPE had the highest risk of SARS-CoV-2 infection, increased susceptibility to infection was evident even among those reporting adequate PPE. Front-line health-care workers who worked in inpatient settings (where providers most frequently reported PPE reuse) and nursing homes (where providers most frequently reported inadequate PPE) had the greatest risk. Non-white health-care workers were disproportionately affected by scant PPE adequacy and more likely to work in clinical settings with greater exposure to patients with COVID-19.

Our findings could help provide greater context for previous cross-sectional reports from public health authorities suggesting 10–20% of SARS-CoV-2 infections occur among health-care workers.^{6,7} Our results offer individual-level data additionally accounting for workplace risk factors that complement these limited reports by providing a more precise assessment of the magnitude of increased risk among health-care workers during the initial phases of the COVID-19 pandemic. Taken in the context of the requirement for testing to establish a COVID-19 diagnosis, our range of results based on either reporting a positive test for COVID-19 or symptoms predictive of COVID-19 offer several complementary estimates for risk among front-line health-care workers.

We also provide evidence that sufficient availability of PPE, quality of PPE, or both reduce the risk of COVID-19, but reuse of PPE or inadequate PPE might confer comparably increased risk, which is compatible with findings from one of the first studies to specifically investigate PPE reuse.¹⁸ The greater risk associated with PPE reuse could be related to either self-contamination during repeated application and removal of PPE or breakdown of materials from extended wear. Of note, during the period of this study, disinfection protocols before PPE reuse were not widely available.^{11–13} Thus, results should be not extended to reflect risk of PPE reuse after such disinfection, which has now been implemented in various settings. An assessment of the PPE supply chain and equitable access to PPE should be a part of the deliberate and informed decision making about resource allocation.

However, even with adequate PPE, health-care workers who cared for patients with COVID-19 remained at increased risk, highlighting the importance of not only ensuring PPE quality and availability but also other aspects of appropriate use, including correct application and removal of PPE and clinical situation (practice location). Moreover, these data underscore the possibility

for health-care workers to perpetuate infections or contribute to community spread, particularly when asymptomatic or mildly symptomatic, and justify calls to increase testing to reduce hospital-based transmission.⁵

Notably, we recorded a significant difference in risk for health-care workers in the UK compared with the USA. This discrepancy could be attributable to country-specific or region-specific variation in population density, socioeconomic deprivation, overall availability or quality of PPE, and type of health-care settings, and these findings require further investigation. Our results might also reflect differences in access to testing among health-care workers compared with the general community in the UK compared with the USA. However, in secondary analyses using inverse probability-weighted Cox modelling adjusting for the probability of receiving a test, we also found that health-care workers in the UK were at higher risk of reporting a positive test. Furthermore, health-care workers were at greater risk of developing symptoms predictive of eventual COVID-19, which does not reflect access to testing. Thus, the higher risk noted in the UK could reflect a higher infection rate because of differences in the quality and appropriate use of PPE across practice settings¹⁹ or country-specific differences in PPE recommendations for health-care workers or the general public (eg, cloth face coverings).^{20,21} Ideally, we would assess risk within a population that has undergone uniform screening. However, the current shortage of PCR-based testing kits does not make such an approach feasible but could justify targeted screening of front-line health-care workers.^{5,22} Future studies using serological testing to ascertain SARS-CoV-2 infection will require assessments of test performance and the ability to distinguish recent or active infection from past exposure.

Our results are supported by historical data during similar infectious disease outbreaks. Ebola virus has a basic reproduction number (ie, the number of new cases generated from one individual) comparable with that for SARS-CoV-2. During the Ebola virus disease crisis, health-care workers comprised 3·9% of all cases, 21–32-times greater than the general public.²³ During the severe acute respiratory syndrome epidemic, health-care workers comprised 20–40% of cases,^{24–26} and inadequate PPE was associated with increased risk.²⁶ The experience with influenza A virus subtype H1N1 reaffirmed the importance of PPE,²⁷ with much higher infection rates among health-care workers in dedicated containment units.²⁸

Our study has several strengths. First, we used a smartphone app to rapidly obtain prospective data from a large multinational cohort in real time, offering actionable risk estimates to inform the public health response to an ongoing pandemic.²⁹ By recruiting participants through existing cohorts,³⁰ our results provide proof of concept of the feasibility of leveraging existing infrastructure and engaged participants to address a key knowledge gap. Second, we obtained information from participants who

did not have a positive COVID-19 test, which offered an opportunity to prospectively assess risk factors with minimal recall bias. Third, our study design recorded initial onset of symptoms, which minimises biases related to capturing only severe cases through hospitalisation records or death reports. Finally, we gathered information on a wide range of known or suspected risk factors for COVID-19 generally not available in existing registries or population-scale surveillance efforts.

We acknowledge several limitations. First, details for some exposures were shortened to ensure our survey was brief. For example, we did not ask about specific occupations, experience level, type of PPE used (eg, masks, respirators, or powered air-purifying respirators), receipt of PPE training (eg, mask fit-testing or application and removal of PPE), and frequency of exposure to patients with SARS-CoV-2 infection or aerosolising procedures. We are pursuing additional questionnaires to more deeply investigate these topics in a subset of participants. Second, our findings are based on self-report. However, alternative exposure measures, such as PPE supply, or assessment of additional outcomes would have been difficult to obtain within the context of a fast-moving pandemic. In future studies, linkage to other sources (eg, electronic health records) might be possible. Third, our cohort is not a random sampling of the population. Although this limitation is inherent to any study requiring voluntary provision of information, we acknowledge that data collection through smartphone adoption has comparatively lower penetrance among some socioeconomic groups and adults older than age 65 years, despite smartphone use by 81% of US adults.³¹ This limitation could have resulted in selection bias, although our primary conclusions were robust to several sensitivity and secondary analyses. In future studies, we plan greater targeted outreach of under-represented populations. Our primary outcome was based on the report of a positive COVID-19 test. During the study period, this outcome would generally reflect a positive PCR-based swab, which should be moderately specific, compared with antibody testing, which was not widely available. However, any misclassification of positive testing should be non-differential according to occupation.

In conclusion, we reported increased risk for SARS-CoV-2 infection among front-line health-care workers compared with the general community, using either self-reported data on COVID-19 testing positivity or a symptom-based predictor of positive infection status. This risk was especially high among Black, Asian, and minority ethnic health-care workers and individuals in direct contact with patients with COVID-19 who reported inadequate PPE availability or were required to reuse PPE. Ensuring the adequate allocation of PPE is important to alleviate structural inequities in COVID-19 risk. However, because infection risk was increased even with adequate PPE, our results suggest the need to ensure proper use of PPE and adherence to other infection control measures. Further intervention studies

investigating modifiable risk factors for health-care worker-related SARS-CoV-2 infection, ideally accounting for differential exposure according to race and ethnic background and care location, are urgently needed to support our observational findings.

Contributors

LHN, DAD, MSG, JW, SO, CJS, TDS, and ATC contributed to study concept and design. LHN, DAD, MSG, JW, SO, RD, JC, CJS, TDS, and ATC contributed to acquisition of data. LHN, DAD, MSG, ADJ, C-GG, WM, RSM, DRS, C-HL, SK, and MS contributed to data analysis. LHN, DAD, and ATC contributed to initial drafting of the manuscript. All authors contributed to interpretation of data and critical revision of the report. SO, CJS, TDS, and ATC contributed to study supervision.

Declaration of interests

JW, RD, and JC are employees of Zoe Global. TDS is a consultant to Zoe Global. DAD and ATC previously served as investigators on a clinical trial of diet and lifestyle using a separate smartphone application that was supported by Zoe Global. All other authors declare no competing interests.

Data sharing

Data collected in the COVID-19 Symptom Study smartphone application are being shared with other health researchers through the UK National Health Service-funded Health Data Research UK (HDRUK) and Secure Anonymised Information Linkage consortium, housed in the UK Secure Research Platform (Swansea, UK). Anonymised data are available to be shared with HDRUK researchers according to their protocols in the public interest. US investigators are encouraged to coordinate data requests through the Coronavirus Pandemic Epidemiology Consortium. Data updates can be found online.

Acknowledgments

LHN is supported by the American Gastroenterological Association Research Scholars Award. DAD is supported by the National Institute of Diabetes and Digestive and Kidney Diseases (K01DK120742). ATC is the Stuart and Suzanne Steele Massachusetts General Hospital (MGH) Research Scholar and a Stand Up to Cancer scientist. National Institutes of Health grants related to this project include UM1 CA186107 (to AHE and MJS), U01 CA176726 (to AHE and WCW), U01 CA167552 (to WCW and LAM), U01 HL145386 (to JEC), R24 ES028521 (to JEC), P30ES000002 (to JEH), and grant contract 200-2017-M-94186 (to JEC). The Massachusetts Consortium on Pathogen Readiness and Mark and Lisa Schwartz supported investigators from MGH. Investigators from King's College London were supported by the Wellcome Trust and Engineering and Physical Sciences Research Council (WT212904/Z/18/Z, WT203148/Z/16/Z, T213038/Z/18/Z), the National Institute for Health Research Guy's and St Thomas' NHS Foundation Trust and King's College London Biomedical Research Centre, the UK Medical Research Council and British Heart Foundation (MR/M016560/1), UK Research and Innovation London Medical Imaging and Artificial Intelligence Centre for Value Based Healthcare, and the Alzheimer's Society (AS-JF-17-011). We thank all individuals who downloaded the COVID Symptom Study smartphone application (app), including participants of cohort studies within the Coronavirus Pandemic Epidemiology (COPE) Consortium (appendix p 3). We thank investigators of cohort studies enrolled in the COPE Consortium for their assistance in disseminating the COVID Symptom Study smartphone app to their study participants, and the MGH Clinical and Translational Epidemiology Unit Clinical Research Coordination team. We thank Zoe Global for providing in-kind support for all aspects of building, running, and supporting the COVID-19 Symptom Study smartphone app and service to users worldwide. Finally, we thank Stand Up to Cancer for their assistance in media and social media outreach.

Editorial note: the Lancet Group takes a neutral position with respect to territorial claims in published maps and institutional affiliations.

References

- Anderson RM, Heesterbeek H, Klinkenberg D, Hollingsworth TD. How will country-based mitigation measures influence the course of the COVID-19 epidemic? *Lancet* 2020; **395**: 931–34.

For the HDRUK Innovation Gateway see <https://healthdatagateway.org>

For the Coronavirus Pandemic Epidemiology Consortium see <https://www.monganinstitute.org/cope-consortium>

For data updates see <https://covid.joinzoe.com>

- 2 Johns Hopkins University of Medicine. Coronavirus resource center. July 22, 2020. <https://coronavirus.jhu.edu/map.html> (accessed July 22, 2020).
- 3 Kirby T. Evidence mounts on the disproportionate effect of COVID-19 on ethnic minorities. *Lancet Respir Med* 2020; **8**: 547–48.
- 4 Garg S, Kim L, Whitaker M, et al. Hospitalization rates and characteristics of patients hospitalized with laboratory-confirmed coronavirus disease 2019: COVID-NET, 14 States, March 1–30, 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**: 458–64.
- 5 Black JRM, Bailey C, Przewrocka J, Dijkstra KK, Swanton C. COVID-19: the case for health-care worker screening to prevent hospital transmission. *Lancet* 2020; **395**: 1418–20.
- 6 CDC COVID-19 Response Team. Characteristics of health care personnel with COVID-19: United States, February 12–April 9, 2020. *MMWR Morb Mortal Wkly Rep* 2020; **69**: 477–81.
- 7 Lazzarini M, Putoto G. COVID-19 in Italy: momentous decisions and many uncertainties. *Lancet Glob Health* 2020; **8**: e641–42.
- 8 Verbeek JH, Rajamaki B, Ijaz S, et al. Personal protective equipment for preventing highly infectious diseases due to exposure to contaminated body fluids in healthcare staff. *Cochrane Database Syst Rev* 2020; **4**: CD011621.
- 9 US Centers for Disease Control and Prevention. Interim infection prevention and control recommendations for healthcare personnel during the coronavirus disease 2019 (COVID-19) pandemic. July 9, 2020. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/infection-control-recommendations.html> (accessed July 15, 2020).
- 10 Public Health England. COVID-19 personal protective equipment (PPE): 5—summary of PPE recommendations for health and social care workers. June 18, 2020. <https://www.gov.uk/government/publications/wuhan-novel-coronavirus-infection-prevention-and-control/covid-19-personal-protective-equipment-ppe#summary-of-ppe-recommendations-for-health-and-social-care-workers> (accessed July 15, 2020).
- 11 Fischer R, Morris DH, van Doremalen N, et al. Assessment of N95 respirator decontamination and re-use for SARS-CoV-2. *medRxiv* 2020; published online April 24. <https://doi.org/10.1101/2020.04.11.20062018> (preprint).
- 12 Schwartz A, Stiegel M, Greeson N, et al. Decontamination and reuse of N95 respirators with hydrogen peroxide vapor to address worldwide personal protective equipment shortages during the SARS-CoV-2 (COVID-19) pandemic. *Appl Biosaf* 2020; **25**: 67–70.
- 13 Livingston E, Desai A, Berkwitz M. Sourcing personal protective equipment during the COVID-19 pandemic. *JAMA* 2020; **323**: 1912–14.
- 14 The Lancet. COVID-19: protecting health-care workers. *Lancet* 2020; **395**: 922.
- 15 Drew DA, Nguyen LH, Steves CJ, et al. Rapid implementation of mobile technology for real-time epidemiology of COVID-19. *Science* 2020; **368**: 1362–67.
- 16 Rosner B. Fundamentals of biostatistics, 4th edn. Boston, MA, Brooks/Cole, 1995.
- 17 Menni C, Valdes AM, Freidin MB, et al. Real-time tracking of self-reported symptoms to predict potential COVID-19. *Nat Med* 2020; **26**: 1037–40.
- 18 Chou R, Dana T, Buckley DI, Selph S, Fu R, Totten AM. Epidemiology of and risk factors for coronavirus infection in health care workers: a living rapid review. *Ann Intern Med* 2020; **173**: 120–36.
- 19 Kenber B, Lay K, Fisher L, Smith HL. Fifth of frontline doctors complain of unusable coronavirus PPE. *The Times (London)*, May 8, 2020.
- 20 Thomas JP, Srinivasan A, Wickramarachchi CS, Dheshi PK, Hung YM, Kamath AV. Evaluating the national PPE guidance for NHS healthcare workers during the COVID-19 pandemic. *Clin Med* 2020; **20**: 242–47.
- 21 Cheng KK, Lam TH, Leung CC. Wearing face masks in the community during the COVID-19 pandemic: altruism and solidarity. *Lancet* 2020; published online April 16. [https://doi.org/10.1016/S0140-6736\(20\)30918-1](https://doi.org/10.1016/S0140-6736(20)30918-1).
- 22 Treibel TA, Manisty C, Burton M, et al. COVID-19: PCR screening of asymptomatic health-care workers at London hospital. *Lancet* 2020; **395**: 1608–10.
- 23 Houlihan CF, McGowan CR, Dicks S, et al. Ebola exposure, illness experience, and Ebola antibody prevalence in international responders to the West African Ebola epidemic 2014–2016: a cross-sectional study. *PLoS Med* 2017; **14**: e1002300.
- 24 Koh D, Lim MK, Chia SE. SARS: health care work can be hazardous to health. *Occup Med* 2003; **53**: 241–43.
- 25 Seto WH, Tsang D, Yung RW, et al. Effectiveness of precautions against droplets and contact in prevention of nosocomial transmission of severe acute respiratory syndrome (SARS). *Lancet* 2003; **361**: 1519–20.
- 26 Lau JT, Fung KS, Wong TW, et al. SARS transmission among hospital workers in Hong Kong. *Emerg Infect Dis* 2004; **10**: 280–86.
- 27 Marshall C, Kelso A, McBryde E, et al. Pandemic (H1N1) 2009 risk for frontline health care workers. *Emerg Infect Dis* 2011; **17**: 1000–06.
- 28 Chen MI, Lee VJ, Barr I, et al. Risk factors for pandemic (H1N1) 2009 virus seroconversion among hospital staff, Singapore. *Emerg Infect Dis* 2010; **16**: 1554–61.
- 29 Chan AT, Brownstein JS. Putting the public back in public health: surveying symptoms of Covid-19. *N Engl J Med* 2020; published online June 5. <https://doi.org/10.1056/NEJMp2016259>.
- 30 Chan AT, Drew DA, Nguyen LH, et al. The Coronavirus Pandemic Epidemiology (COPE) Consortium: a call to action. *Cancer Epidemiol Biomarkers Prev* 2020; **29**: 1283–89.
- 31 Pew Research Center for Internet & Technology. Mobile fact sheet. June 12, 2020. <https://www.pewresearch.org/internet/fact-sheet/mobile/> (accessed July 15, 2020).