1 PCD Detect: Enhancing ciliary features though image averaging and classification

2

3 Amelia Shoemark^{1,2}, Andreia L. Pinto¹, Mitali P. Patel⁴, Farheen Daudvohra¹, Claire Hogg^{1,3}, Hannah
4 M. Mitchison^{4,5}, Thomas Burgoyne^{1,6} #

5

6 ¹ Paediatric Respiratory Medicine, Primary Ciliary Dyskinesia Centre, Royal Brompton & Harefield
7 NHS Trust, London, SW3 6NP, UK

8 ² School of Medicine, University of Dundee, Dundee, DD1 9SY, UK

9 ³ Department of Paediatrics, Imperial College London, London, SW3 6LY, UK

10 ⁴ Genetics and Genomic Medicine, University College London, UCL Great Ormond Street Institute of
11 Child Health and NIHR Great Ormond Street Hospital Biomedical Research Centre, London, WC1N
12 1EH, UK

13 ⁵ NIHR Great Ormond Street Hospital Biomedical Research Centre, London, WC1N 1EH, UK

14 ⁶ UCL institute of Ophthalmology, London, EC1V 9EL, UK

15 # All correspondence to t.burgoyne@rbht.nhs.uk, Paediatric Respiratory Medicine, Primary Ciliary
16 Dyskinesia Centre, Royal Brompton & Harefield NHS Trust, London, SW3 6NP, UK Tel: 44-(0)20-
17 7352-8121, ORCID: 0000-0002-8428-720X

18

19 Author Contributions: T.B.: wrote the programs; A.S. and T.B.: designed and performed experiments
20 as well as analysed and interpreted data; A.L.P. and F.D.: analysed and interpreted data; A.S.,
21 H.M.M.: provided genetic results; A.S., T.B., H.M.M and C.H.: wrote the manuscript.

22

23 Running Head: PCD Detect: Cilia ultrastructural analysis software

24 Keywords: primary ciliary dyskinesia, cilia ultrastructure, image averaging, image classification,
25 transmission electron microscopy
26

27 Abstract

28 Primary ciliary dyskinesia (PCD) is an inherited disorder of the motile cilia. Early accurate diagnosis is
29 important to help prevent lung damage in childhood and to preserve lung function. Confirmation of a
30 diagnosis traditionally relied on assessment of ciliary ultrastructure by transmission electron
31 microscopy (TEM), however >50 known PCD genes has made the identification of bi-allelic mutations
32 a viable alternative to confirm diagnosis. TEM and genotyping lack sensitivity and research to improve
33 accuracy of both is required. TEM can be challenging when a subtle or partial ciliary defect is present
34 or affected cilia structures are difficult to identify due to poor contrast. Here we demonstrate software
35 to enhance TEM ciliary images and reduce background by averaging ciliary features. This includes an
36 option to classify features into groups based on their appearance, to generation multiple averages
37 when a nonhomogeneous abnormality is present. We validated this software on images taken from
38 subjects with well characterised PCD caused by variants in the outer dynein arm (ODA) heavy chain
39 gene *DNAH5*. Examining more difficult to diagnose cases, we detected (i) regionally restricted
40 absence of the ODAs away from the ciliary base, in a subject carrying mutations in *DNAH9*; (ii) loss of
41 the typically poorly contrasted inner dynein arms; (iii) sporadic absence of part of the central pair
42 complex in subjects carrying mutations in *HYDIN*, including one case with an unverified genetic
43 diagnosis. We show this easy to use software can assist in detailing relationships between genotype
44 and ultrastructural phenotype, improving diagnosis of PCD.

45 Introduction

46 Motile cilia are hair-like projections on the apical surface of human epithelial cells that have a classical
47 9+2 microtubule arrangement of their core axoneme structure (Fig. 1). Respiratory motile cilia beat in
48 a highly coordinated manner essential for the movement and clearance of mucus, bacteria and debris
49 from the airways. Primary ciliary dyskinesia (PCD) is an inherited disorder of motile cilia that results in
50 failure of mucus clearance, chronic lung disease, rhinosinusitis and hearing impairment, with
51 additional organ laterality defects, subfertility and occasional hydrocephalus. The estimated
52 prevalence is ~1 per 10,000 births, but it is more prevalent in populations where consanguinity is
53 common (7). An early diagnosis of PCD is vital to implement appropriate treatment with the aim to
54 preserve lung function and prevent lung damage. Diagnosis is often complex, due to disease
55 heterogeneity and incomplete understanding of the molecular and genetic causes of PCD. Diagnosis
56 currently involves combinations of tests, including measurement of nasal nitric oxide (nNO) as well as
57 acquiring cells by nasal brushings for examination of cilia motility functions using high speed video
58 microscopy (HSVM), immunofluorescence microscopy (IF), transmission electron microscopy (TEM)
59 and genotyping (18). Confirmation of a diagnosis traditionally relied on assessment of ciliary
60 ultrastructure by TEM, however advancements in genetic testing has made the identification of bi-
61 allelic mutations in a known PCD gene a viable alternative to confirm diagnosis. Mutations in more
62 than 50 different genes are known to cause PCD, defects which account for 65-75% cases.
63 Therefore, more than a quarter of sufferers have no genetic cause yet established despite screening,
64 which can be because they carry no detectable mutations in the known genes, or carry mutations in
65 known genes of unknown significance or have no diagnosis with only single mono-allelic (rather than
66 biallelic) mutations so far identified (26). In these patients, confirmation with a functional test is
67 required.

68 TEM diagnosis involves examining a large number (>50) of cilia to verify the frequency of particular
69 ultrastructural defects (28). The most commonly reported PCD defect affecting patients that can be
70 identified using this approach is the absence of outer dynein arms (ODAs) (Fig. 1). Absent ODAs are
71 readily detected due to loss of a well contrasted (high signal to noise) structure in the TEM. The most
72 common cause of this defect is a mutation in the *DNAH5* gene that encodes a large ODA heavy chain
73 that forms a substantial part of the ODA (20). Unfortunately, there are a range of other ciliary
74 structures affected in PCD that are more challenging to visualise by TEM. These include inner dynein
75 arms (IDAs) that typically have poor contrast and are often masked by the close positioning or
76 overlapping of the radial spokes and nexin links. Also challenging to image are the central pair
77 complex (CP) projections such as the CP C2b component HYDIN, which are small and difficult to
78 detect due to the resolution limits of diagnostic electron microscopes (21). Furthermore, there are
79 specific genetic defects causing PCD that are known to result in partial loss or sporadic loss of a
80 ciliary structure. This includes mutations in the *DNAH9* and *DNAH11* genes which encode heavy
81 chain subunits of the ODA, that are both localised only to a sub-region of the axoneme, the distal or
82 proximal region, respectively. PCD-causing variants in *DNAH9* and *DNAH11* consequently lead to a
83 loss of ODAs restricted to the distal (cilia tip) or proximal (cilia base) regions, as these proteins serve

complementary roles in either region (8, 12). Due to the difficulty of detecting the presence or absence of some ciliary structures, many patients end up having multiple nasal brushings, leading to delayed diagnosis. Approximately 30% of patients are considered to have a normal ciliary ultrastructure after TEM analysis (14). Subtle defects, e.g. *HYDIN* which have been missed, are likely to account for a number of these cases.

Recently, electron tomography and 3D ciliary averaging have been used to successfully identify subtle cilia defects in samples previously considered to have normal ultrastructure. However this is a time consuming, expensive technique requiring specialist tools and knowledge (4, 8, 12, 17, 27). To address this clinical need to improve the detection of subtle or partial structural defects in cilia of patients that are likely to have PCD, we have developed PCD detect, a package of programs (toolkit) that allows the selection and averaging of ciliary features (microtubular doublets (MTDs) and CPs) from TEM images. This assists in removing background while enhancing signal, allowing ciliary structures that are critical for motility to be better structurally resolved and evaluated. Of importance for some PCD cases where there is partial or sporadic loss of a ciliary structure, and/or genetic variants identified that are of uncertain clinical significance, the image classification tool has been developed to provide multiple (class) averaged outputs to better study and improve detection of these defects.

101

102 **Materials and methods**

103 **Ethical approval**

The study was approved by the ethics review board of the Institute of Child Health/Great Ormond Street Hospital, London (UK) (08/H0713/82). All subjects gave informed signed consent for genetic testing and use of surplus diagnostic samples for development of new diagnostic tools.

107

108 **Preparation of nasal brushings for electron microscopy analysis**

Respiratory epithelial cells were obtained from subjects' nasal turbinate using 3-mm bronchial cytology brushes and placing the biopsy into media 199 (Thermo Fisher Scientific). Each biopsy was fixed and embedded following the protocol outlined in reference (24). In brief, an equal volume of 2.5% glutaraldehyde in 0.05M sodium cacodylate buffer pH 7.4 was added to the solution containing the biopsies in 199 media and left overnight or longer (up to 2 weeks) at 4°C. The biopsies were incubated in 1% osmium tetroxide in 0.05M sodium cacodylate buffer pH 7.4 for 1 hour before embedding in 2% agar aqueous solution. The samples were then dehydrated and infused with araldite using a Lynx Microscopy Tissue Processor. This involved incubating in increasing concentrations of methanol (70%, 90% and 100%) followed by propylene oxide, mixtures of propylene oxide:araldite (1:1 and 1:3 respectively) and araldite. Finally, the samples were embedded in araldite and incubated at 65°C for 72 hrs. 90nm sections from archived araldite-embedded nasal brushings

were cut onto electron microscopy grids and stained using 2% methanolic uranyl acetate for 7 minutes and Reynolds lead citrate solution for 5 minutes. The sections were viewed on a JEOL 1400+ TEM and when needed the stage was tilted to provide ciliary cross-sections without obliqueness. Images were acquired at 40,000x magnification using an AMT 16X CCD camera.

Image criteria

For optimal analysis of cilia using the PCD detect toolkit, it is important to use clear images (strong signal to background), without stain artefacts, taken at a single magnification, also to include non-oblique cross sections taken at the focus where it is possible to see the most ultrastructural detail. Demonstration images are included along with the toolbox to demonstrate the kind of images that are required. Blinded averaged images generated from control and patient cilia were examined to determine sensitivity and specificity. It is recommended to pick MTD features from at least 5 cilia cross sections (equivalent to 45 or more MTD features) from different cells for a sensitivity $\geq 75\%$ and specificity of 100%. More than 10 cilia cross sections from different cells should be used in PCD detect when examining the CP (equivalent to more than 10 CP features), for a sensitivity $>63\%$ and specificity $>83\%$.

Software development

The toolkit includes programs to extract cilia from TEM images (Cilia Crop), perform averaging of cilia structures (Cilia Averaging), group and generate class averages (Cilia Classification) and to overlay results for analysis (Cilia Analysis). All programs were written in Matlab (Mathworks).

Cilia Crop: This is a tool to select multiple regions of interest from a larger image. (An example is shown in Fig. S1 A).

Cilia Averaging: This tool extracts manually chosen features (some of these are shown in Fig. S2) and performs a rotational and translational search (see Fig. S1 B). By calculating the cross-correlation, the best alignment of features is determined. The outputs from this tool include two averaged images comprising (i) all features averaged together (ii) only features best matching the reference based on the cross-correlation cut-off value. The latter average outputs are used in this study, as these should contain less background by not including poor quality features (high background, artefacts, tilted or distorted). All averaged images generated include a corresponding colour contour map where dark blue indicates an electron dense feature and yellow indicates the absence of a feature. When using a reference feature (rather than reference-free), it is important to select a well orientated (non-oblique) feature (MTD or CP) that has good contrast and no stain artefacts.

Cilia Classification: The classification tool (Fig. S3 A) was developed to allow two methods of grouping features based on their similarity (image classification) as determined by (i) principle

component analysis (PCA) and (ii) hierarchical classification. When using PCA, histograms of the first six coefficients are generated and the plots that indicate more than one feature class (multiple peaks) are selected. Using these, k-means clustering then separates the features into two or more classes, which is followed by averaging them together to generate the separate class averages. The hierarchical classification approach was designed to analyse the feature alignment and similarity, using cross-correlation results generated previously through the averaging tool. Using this data, the first class is generated by averaging features aligned to the reference feature (has the most matching other features based on cross-correlation). Subsequent classes are generated by averaging to the reference features that are most dissimilar to the reference features used to generate the first class. The hierarchical classification approach allows a feature to be included in more than one class if there is enough similarity (based on cross-correlation) to more than one of the reference features used to generate the classes. This is to allow any features that cannot be easily classed to contribute towards multiple class averages. It is important to note that class averages can be generated when there is sporadic loss of structure(s), but class averages can also be given by other types of variation between groups of features, such as tilt (obliqueness). In this study, when using PCA (Fig. S4), the number of classes was set to 2 and the PCA and k-means calculated from an averaged version of each feature (each feature acts as reference and all other features are averaged to it). By picking the plots that indicate multiple classes (more than one peak), k-means are calculated to group features before generating class averages.

Cilia Analysis: This tool allows the visualisation and comparison of averaged features (Fig. S3B). By opening two features, an overlay on both is provided. The features can be aligned automatically or manually, to allow similarities and differences in the structures to be assessed (Fig. S3 B2-4).

Toolkit validation

Image acquisition

The toolkit was initially tested using artificially generated images with added noise (see Fig. S5) before using ciliary images taken from patients with PCD. Details of the PCD subjects in this study are provided in Table 1.

Input into PCD Detect

For the validation, the four programs were run sequentially. In cilia crop, cropping size is selected by boxing an entire cross-sectional cilium, with cilia cropped out into a sub-directory by pressing the Select ROI (region of interest) button and selecting the centre of each cilium within an image. The folder containing the cropped images is then opened in the Cilia Averaging tool (Fig. S1 B). Each image is checked manually to make sure all cilia are orientated in the same direction by flipping images either horizontally or vertically. A box is drawn around the feature of interest to provide its size before picking the features to be averaged together. For the validation, the feature of interest was

either the MTD (*DNAH5*, *DNAH9*, *CCDC103* or *CCDC40* cases) or the CP (*HYDIN* cases). Reference free averaging was used, as well as a search range of 360° with increments of 1° and a translational search in X and Y of 10 pixels with an increment size of 1 pixel.

After the Cilia Averaging tool performs the rotational and translational search aligning the features, an average feature is generated. The data is opened and further processed within the Cilia Classification tool (Fig. S3 A), using both the PCA and hierarchy image classification schemes to group features based on their structure and generate class averages. When using the hierarchy classification, a cross-correlation cut-off of 0.6 (any features that have a cross-correlation when compared to the reference below this value are not included in the class average) and class cut off of 10% (if a reference is found to have less than 10% of the total number of features matching it, no class average is generated) were set. Finally, the analysis tool is used to overlay features between classes as well as between PCD cases and control images. A workflow of PCD Detect is shown in Fig 2.

Results

Averaged MTDs show complete or sporadic absence of the ODAs in subjects with a pathogenic variant in *DNAH5*

The utility of PCD Detect was first evaluated in the analysis of nasal epithelial cilia structures from PCD patients carrying causal *DNAH5* mutations, already established to have clear-cut defects in conventional TEM analysis, involving loss of the ciliary ODAs. TEM images of transverse orientation cilia were acquired from diagnostic nasal brushing samples taken from control subjects as well as the *DNAH5* PCD subjects (Fig. 3 A, B and Table 1). Using the image averaging tool, the MTDs were selected and reference-free alignment and averaging was performed. The averaged MTD generated from a control and three *DNAH5* subjects are shown in Fig. 3 C, D. Averages from further controls are given in Fig S6. In all three *DNAH5* cases (*DNAH5* [1-3]), there is a clearly visible absence of the ODA, verifying the ability of PCD Detect to identify PCD. Interestingly in two of these cases, *DNAH5* [1] and *DNAH5* [2], a small visible structure (white filled arrowheads in Fig. 3 D) can be seen, indicating that part of the ODA remains, whilst ODA are completely absent in the *DNAH5* [3] case (denoted by the white outline arrowheads in Fig. 3 D). Of these cases, *DNAH5* [1] carries two stop-gain mutations (homozygosity confirmed by parental segregation analysis), while in *DNAH5* [2] and the previously reported case *DNAH5* [3] (11) the genetic diagnosis is not complete since only one *DNAH5* variant has so far been identified in each, although it is a highly implicated rare stop-gain allele in *DNAH5* [3]; *DNAH5* [2] carries a single uncharacterised missense mutation (Table 1).

A further subject *DNAH5* [4], with an uncharacterised missense mutation (p.Thr3410Met) plus a previously reported *DNAH5* stop-gain mutation p.Arg1761* (10), was examined using the classification tool to group MTDs and form class averages. Many of the ciliary cross sections look normal by TEM before using the enhancement tools (Fig. 4 B). Class averages generated using the hierarchy classification scheme are shown in Fig. 4 C, D and plots from the hierarchy classification as

well as the PCA classification results are shown in Fig. S4. When using this tool, subject *DNAH5* [4] with the combined nonsense and missense mutation (Fig. 4 C), had two visibly different MTD classes, with and without the ODA. This reflects a sporadic absence of the ODAs that was not found in the other three *DNAH5* cases, as all classes generated had absent ODA (results for *DNAH5* [1] shown in Fig. 4 D, E and other cases not shown).

Multiple MTD classes generated from a subject with pathogenic variants in *DNAH9* due to loss of the ODA at the distal region of the cilia

To further test the classification tool, we examined ciliary images from a subject that carried a homozygous proven-pathogenic variant in *DNAH9* (p.Asp4123Asn) (12) (Table 1). *DNAH9* protein is a component only of the ODA in the distal cilium, not the proximal region (determining these regions is shown in Fig. 5 A) and it is known that *DNAH9* mutations cause a defect where the ODA are only absent in the distal region of the ciliary axoneme (12, 13). Images were acquired in both the proximal (Fig. 5 B) and distal regions (Fig. 5 C) of the ciliary axoneme. By averaging the MTDs from these images and using hierarchy classification, two distinct classes were generated when compared to control cilia (Fig. 5 D, E). When separating the ciliary images into two groups depending on the region, the classification results in single distinct averaged structures for the distal and proximal regions, indicating a clear structural difference (Fig. 5 F, G). Images at the proximal region of the cilia resulted in an averaged structure where the ODA is present, whereas the distal ciliary images resulted in an averaged structure showing absence of the ODA.

The absence of IDAs can be detected in cases with pathogenic variants in *CCDC40* and *CCDC103*

As the absence of IDAs can be difficult to detect by TEM, we determined whether averaging MTDs was able to enhance this structure. We examined cilia from subjects with known pathogenic variants in *CCDC40* (subject homozygous for the splice site mutation c.2712-1G>T) (2) or *CCDC103* (subject homozygous for the well-studied missense mutation p.His154Pro) (30) and compared them against a control (Table 1, Fig. 6 A-C). Mutations in both these genes confer a loss of IDAs in the cilia of affected patients. When acquiring images, as the *CCDC40* variant also results (as expected (2)) in a proportion of cilia with disarrangement MTDs, only normal 9+2 arranged cilia were selected for IDA imaging. When comparing ciliary cross-section images from both *CCDC40* and *CCDC103* variants against a control, it was clear in conventional TEM that IDA loss, in addition to their being of poor contrast, can be masked by the nearby or overlapping presence of the electron dense radial spoke and nexin structures as well as remnants of some IDA isoforms (Fig. 6 A-C). By averaging the MTDs together using PCD Detect, it became apparent that the IDAs were absent in the *CCDC40* and less dense in the *CCDC103* variant cases when compared to a control (Fig. 6 D-F).

264

265 **Confirmation of *HYDIN* mutations by visualisation of the central pair component C2b**

266 To examine the absence of the C2b component that arises due to pathogenic variants in *HYDIN*, cilia
267 from three affected subjects (cases *HYDIN* [1-3]) carrying pathogenic variants and a subject with
268 variants of unknown significance in *HYDIN* were compared to control cilia (Fig. 7 A, B). *HYDIN* [1] and
269 [2] carry compound heterozygous variants of pathogenic consequence while *HYDIN* [3] carries only a
270 single variant but one that was previously published (6). Subject *HYDIN* [4] is also less well proven for
271 genetic diagnosis, carrying two missense *HYDIN* variants that have not been previously reported or
272 functionally tested for their pathogenicity (Table 1). The CP complex from the ciliary images were
273 averaged together in each case and the hierarchy classification system used to generate class
274 averages (Fig. 7 C-E). In the *HYDIN* mutation subjects, two main class averages were generated from
275 the classification tool, indicating a partial defect in the C2b component, where this component is
276 present in some cilia and absent in others (white filled arrowheads in Fig. 7 D, E). As the subjects with
277 predicted null-effect causal variants (*HYDIN* [1-3]) generated the same result as the missense
278 mutation-carrying subject (*HYDIN* [4]), we could confirm that all these *HYDIN* [4] variants are highly
279 likely to be pathogenic in effect. To highlight the absence of the C2b component in the *HYDIN* [4]
280 subject, we overlaid the averaged CP complex on top of the control using the 'cilia analysis' tool
281 (white outlined arrowhead in Fig. 7 E). While examining *HYDIN* [4], further cases were assessed that
282 either had previous diagnostic tests (semi-circling patches of cilia visualised by HSVM) indicating a
283 potential *HYDIN* defect (PHD) or had variants of unknown significance (VUS) in *HYDIN* (Table S1).
284 These individuals were all found to have a C2b component present in the averaged images generated
285 (Fig. S7), suggesting that *HYDIN* variants are unlikely to be their disease cause.

286

287 **Discussion**

288 To examine important structures in respiratory cilia that can be affected in PCD and other conditions
289 affecting the airway motile cilia (19), we have developed and implemented PCD Detect, a user-
290 friendly package of tools for image averaging and classification. Due to the uniquely conserved
291 structure and abundance of motile cilia in patient respiratory epithelial samples, these are ideal
292 structures for this type of image analysis. By averaging some of the structures that are repeated
293 within cilia cross-sections, background noise can be reduced and the signal and image resolution
294 enhanced. As some PCD defects are well known to be difficult to detect from TEM images due to their
295 small size as well as lack of contrast or resolution, the averaging technique this package employs can
296 provide an additional approach to assist in the diagnosis of PCD in problematic cases. It has become
297 increasingly recognised that diagnosis of PCD by TEM can be further complicated by the partial or
298 sporadic absence of a ciliary structure (8, 12, 21, 29, 34). The classification tool can assist in these
299 cases by grouping cilia axonemal structures (classes) based on their similarities.

The first samples we used to test PCD Detect were from subjects with pathogenic variants in *DNAH5*, chosen because these typically present with a clear absence of the ODA. When averaging the first three subjects' MTDs, the ODAs were evidently absent, but in some cases a remnant of the ODA was observed. The ODA is attached to the MTD via a docking complex (15, 31). Depending on the type of variant in *DNAH5*, PCD-causing mutations may result in the loss of a portion or the entire docking complex, which may explain the remaining structure that is present in some cases (32).

When examining the MTDs in a subject (*DNAH5* [4]) carrying a missense variant in *DNAH5* that is uncharacterised for its likely pathogenic effect, using a combination of image averaging and classification, two classes of structure were found, with and without the ODA present. A partial loss of the ODA has been previously shown for a *DNAH5* splice site mutation (16). Certain gene variants may lead to reduced synthesis of *DNAH5* or of *DNAH5* protein that is transported and integrated into the cilium but lacks stability, leading to loss of a portion of the ODAs. Our averaging and classification tool allowed identification of this complex defect. The success of the average classification was further demonstrated in a subject with pathogenic variants in *DNAH9* which are known to lead to a loss of ODA only in the distal region of the cilium, whereas the proximal cilium displayed a normal ultrastructure due to the compensating presence of *DNAH11* (8, 12). When using all images acquired from a *DNAH9* PCD subject for the averaging and classification scheme, two class averages were generated, with and without the ODA. By splitting the image input for PCD Detect into two groups, either proximal (determined by the presence of epithelial cell apical microvilli) or distal, only single averages of distinct structure were generated when using the classification tool. This shows that PCD Detect can successfully group and average similar structures from a non-homogeneous dataset. The separation of images into the distal and proximal was done manually, but we aim to develop a system that can detect microvilli and segregate images automatically, which can be added to PCD Detect in the future.

The IDAs are typically difficult to detect in conventional ciliary TEM, linked to the fact that within the 96nm axonemal repeat there are four overlapping homologous double headed ODAs, whereas there are six single-headed and one double-headed IDA of varying morphology (17). This results in a lower contrast of the IDA than the ODA. Methods to enhance the contrast and reduce background can be highly beneficial in TEM analysis when needing to determine if the IDAs are present or absent. The cilia from subjects carrying pathogenic variants in *CCDC40* and *CCDC103* were both examined, as mutations in both these genes give rise to loss of the IDAs, with the distinction that *CCDC40* mutations are also linked to microtubular disorganisation and *CCDC103* mutations linked to ODA loss (3, 23). The averaged MTDs generated successfully showed a clear absence of the IDA for both mutation types.

To examine CP complex defects in PCD, the cilia of three subjects' that carry pathogenic variants in *HYDIN* were examined as well as cilia from a number of other subjects with less well confirmed genetic results suggestive of *HYDIN* defects or no genetic tests yet performed. Using the averaging and classification scheme, all the three well genetically defined cases were found to have two class averages, as well as one individual (*HYDIN* [4]) carrying biallelic VUS, all of these cases showing the

339 presence and absence of the C2b projection known to be associated with pathogenic variants in
340 *HYDIN*. PCD Detect was also able to confirm the presence of the C2b structure in several other
341 analysed cases that either had suspected *HYDIN* defects based on previous clinical tests, or who
342 carried other *HYDIN* VUS, thus excluding *HYDIN* defects as their disease cause. This is the first time
343 two distinct CP structures have been shown to exist in the positively diagnosed *HYDIN* subjects. The
344 finding of the same two class-averaged structures in the less genetically well-unverified *HYDIN* [4]
345 confirmed the likely causal nature of their *HYDIN* variants, a conclusion that was supported by the
346 subject's history and other diagnostic results.

347 PCD Detect is a free toolkit that has advantages over other TEM image enhancing techniques such
348 as electron tomography as it requires less specialised knowledge, uses information from a larger
349 number of cilia and does not require prior information about which structure is affected. It does require
350 multiple TEM images of well orientated cilia, knowledge of how to interpret the final averaged images
351 and it has a lower resolution and lack of 3D data that is achievable through electron tomography. This
352 system package has built upon averaging and image enhancement methods previously developed to
353 assist in diagnosing PCD (1, 5, 9, 22, 25). The simple, automated angular and translational searches,
354 cross correlation and class averaging features present advantages over previously designed
355 averaging programs. Previous methods to enhance ciliary structures such as the IDAs have involved
356 using the cilium's axonemal symmetry to average MTD (9). This involves manually rotating an image
357 of a cilium through steps of 40° and averaging each time. This is therefore limited to an average of
358 only 9 MTD per cilium. In PCD Detect, by allowing the toolkit to extract the features, there is no limit to
359 the number of structures that can be averaged together. The averaging scheme in PCD Detect has a
360 user-friendly interface along with a robust feature comparison and alignment system using cross-
361 correlation, allowing the exclusion of poor-quality features or artefacts. Furthermore, the addition of a
362 classification tool allows analysis of more complex, non-consistent PCD defects associated with non-
363 homogenous structural defects.

364 The averaging and classification scheme available within PCD Detect is particularly well suited for
365 difficult to diagnose cases that have atypical, poor characterised or difficult to detect ciliary structural
366 defects. Many of these patients have previously had to have multiple nasal brushings to make a
367 diagnosis (Table 1), as their initial TEM diagnostic results were either normal or inconclusive and only
368 after multiple inspections was a defect detected. PCD Detect has the potential to reduce the need for
369 additional nasal brushings, speeding up the diagnostic turnover times as well as reducing the use of
370 diagnostic resources.

371 In summary, PCD Detect is a toolkit to enhance diagnostic images of PCD patient cilia, through an
372 averaging and classification scheme. It has not been designed for routine use, but instead to assess
373 difficult cases that have had a suggestive HSVM result, TEM result or a variant of unknown
374 significance by genetics when following the conventional diagnostic pathway. PCD Detect should be
375 used alongside other test results as part of a multi-test diagnosis not as a standalone tool.
376 Furthermore, it is important that high quality TEM images are used to provide reliable averaged
377 images. We were able to show successful enhancing of structural detail in samples that have clear-

cut as well as more subtle ciliary structural defects and defects affecting only a portion of the axoneme or a minor structure such as the C2b projection of the CP. PCD Detect can in this way form an associated technique that uplifts current conventional PCD genetic diagnosis. This is something that can be helpful where genetic diagnosis is more difficult, due to genetic variants of uncertain significance being identified. It has notable application for the growing number of affected individuals in whom functional tests such as conventional TEM are equivocal and they carry genetic variants that have been identified in known genes but are of less certain causality, for example if they are not previously reported or where single amino acids are affected (33), or in the specific case of *HYDIN* where variants are hard to call because of the *HYDIN2* copy gene (21). This study shows PCD Detect has the potential to benefit global PCD diagnostic centres. Furthermore, the principles of the toolkit could also be applied to image data beyond ciliary structure analysis, including homogenous biological structures affected in diseases where diagnostics involves TEM or for investigative structural biological studies.

Acknowledgements

We are very grateful to the families who participated in this study and thank the UK PCD Family Support Group for their continued support, as well as Dr James Geraets (University of Helsinki) for his image analysis advice. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. The authors participate in the COST Action BEAT-PCD: Better Evidence to Advance Therapeutic options for PCD network (BM1407). H.M.M. is supported by Great Ormond Street Children's Charity (V1299, V2217) and the NIHR Great Ormond Street Hospital Biomedical Research Centre (GOSH BRC).

References

1. **Afzelius BA, Dallai R, Lanzavecchia S, Bellon PL.** Flagellar structure in normal human spermatozoa and in spermatozoa that lack dynein arms. *Tissue and Cell* 27: 241–247, 1995. doi: 10.1016/S0040-8166(95)80044-1.
2. **Antony D, Becker-Heck A, Zariwala MA, Schmidts M, Onoufriadis A, Forouhan M, Wilson R, Taylor-Cox T, Dewar A, Jackson C, Goggin P, Loges NT, Olbrich H, Jaspers M, Jorissen M, Leigh MW, Wolf WE, Daniels MLA, Noone PG, Ferkol TW, Sagel SD, Rosenfeld M, Rutman A, Dixit A, O’Callaghan C, Lucas JS, Hogg C, Scambler PJ, Emes RD, Uk10k, Chung EMK, Shoemark A, Knowles MR, Omran H, Mitchison HM.** Mutations in CCDC39 and CCDC40 are the Major Cause of Primary Ciliary Dyskinesia with Axonemal Disorganization and Absent Inner Dynein Arms. *Human Mutation* 34: 462–472, 2013. doi: 10.1002/humu.22261.
3. **Becker-Heck A, Zohn IE, Okabe N, Pollock A, Lenhart KB, Sullivan-Brown J, McSheene J, Loges NT, Olbrich H, Haeffner K, Fliegauf M, Horvath J, Reinhardt R, Nielsen KG, Marthin JK, Baktai G, Anderson KV, Geisler R, Niswander L, Omran H, Burdine RD.** The coiled-coil domain containing protein CCDC40 is essential for motile cilia function and left-right axis formation. *Nature Genetics* 43: 79–84, 2011. doi: 10.1038/ng.727.
4. **Burgoyne T, Lewis A, Dewar A, Luther P, Hogg C, Shoemark A, Dixon M.** Characterizing the ultrastructure of primary ciliary dyskinesia transposition defect using electron tomography. *Cytoskeleton (Hoboken)* 71: 294–301, 2014. doi: 10.1002/cm.21171.
5. **Carson, JL, Hu, S-CS, Collier AM.** Computer-Assisted Analysis of Radial Symmetry in Human Airway Epithelial Cilia: Assessment of Congenital Ciliary Defects in Primary Ciliary Dyskinesia. *Ultrastructural Pathology* 24: 169–174, 2000. doi: 10.1080/01913120050132903.
6. **Davidson AE, Schwarz N, Zelinger L, Stern-Schneider G, Shoemark A, Spitzbarth B, Gross M, Laxer U, Sosna J, Sergouniotis PI, Waseem NH, Wilson R, Kahn RA, Plagnol V, Wolfrum U, Banin E, Hardcastle AJ, Cheetham ME, Sharon D, Webster AR.** Mutations in ARL2BP, Encoding ADP-Ribosylation-Factor-Like 2 Binding Protein, Cause Autosomal-Recessive Retinitis Pigmentosa. *Am J Hum Genet* 93: 321–329, 2013. doi: 10.1016/j.ajhg.2013.06.003.
7. **Dehlink E, Hogg C, Carr SB, Bush A.** Clinical phenotype and current diagnostic criteria for primary ciliary dyskinesia. *Expert Review of Respiratory Medicine* 10: 1163–1175, 2016. doi: 10.1080/17476348.2016.1242414.
8. **Dougherty GW, Loges NT, Klinkenbusch JA, Olbrich H, Pennekamp P, Menchen T, Raidt J, Wallmeier J, Werner C, Westermann C, Ruckert C, Mirra V, Hjeij R, Memari Y, Durbin R, Kolb-Kokocinski A, Praveen K, Kashef MA, Kashef S, Eghtedari F, Häffner K, Valmari P, Baktai G, Aviram M, Bentur L, Amirav I, Davis EE, Katsanis N, Brueckner M, Shaposhnykov A, Pigino G, Dworniczak B, Omran H.** DNAH11 Localization in the Proximal Region of Respiratory Cilia Defines Distinct Outer Dynein Arm Complexes. *Am J Respir Cell Mol Biol* 55: 213–224, 2016. doi: 10.1165/rcmb.2015-0353OC.
9. **Escudier E, Couprie M, Duriez B, Roudot-Thoraval F, Millepied M-C, Prulière-Escabasse V, Labatte L, Coste A.** Computer-assisted Analysis Helps Detect Inner Dynein Arm Abnormalities. *Am J Respir Crit Care Med* 166: 1257–1262, 2002. doi: 10.1164/rccm.2111070.
10. **Failly M, Bartoloni L, Letourneau A, Munoz A, Falconnet E, Rossier C, Santi MM de, Santamaria F, Sacco O, DeLozier-Blanchet CD, Lazor R, Blouin J-L.** Mutations in DNAH5 account for only 15% of a non-preselected cohort of patients with primary ciliary dyskinesia. *Journal of Medical Genetics* 46: 281–286, 2009. doi: 10.1136/jmg.2008.061176.
11. **Fassad MR, Patel MP, Shoemark A, Cullup T, Hayward J, Dixon M, Rogers AV, Ollosson S, Jackson C, Goggin P, Hirst RA, Rutman A, Thompson J, Jenkins L, Aurora P, Moya E,**

- 451 Chetcuti P, O'Callaghan C, Morris-Rosendahl DJ, Watson CM, Wilson R, Carr S, Walker
452 W, Pitno A, Lopes S, Morsy H, Shoman W, Pereira L, Constant C, Loebinger MR, Chung
453 EMK, Kenia P, Rumman N, Fasseeh N, Lucas JS, Hogg C, Mitchison HM. Clinical utility of
454 NGS diagnosis and disease stratification in a multiethnic primary ciliary dyskinesia cohort. .
- 455 12. Fassad MR, Shoemark A, Legendre M, Hirst RA, Koll F, le Borgne P, Louis B, Daudvohra
456 F, Patel MP, Thomas L, Dixon M, Burgoyne T, Hayes J, Nicholson AG, Cullup T, Jenkins
457 L, Carr SB, Aurora P, Lemullos M, Aubusson-Fleury A, Papon J-F, O'Callaghan C,
458 Amselem S, Hogg C, Escudier E, Tassin A-M, Mitchison HM. Mutations in Outer Dynein Arm
459 Heavy Chain DNAH9 Cause Motile Cilia Defects and Situs Inversus. *The American Journal of*
460 *Human Genetics* 103: 984–994, 2018. doi: 10.1016/j.ajhg.2018.10.016.
- 461 13. Fliegauf M, Olbrich H, Horvath J, Wildhaber JH, Zariwala MA, Kennedy M, Knowles MR,
462 Omran H. Mislocalization of DNAH5 and DNAH9 in Respiratory Cells from Patients with Primary
463 Ciliary Dyskinesia. *Am J Respir Crit Care Med* 171: 1343–1349, 2005. doi:
464 10.1164/rccm.200411-1583OC.
- 465 14. Goutaki M, Maurer E, Halbeisen FS, Amirav I, Barbato A, Behan L, Boon M, Casaulta C,
466 Clement A, Crowley S, Haarman E, Hogg C, Karadag B, Koerner-Rettberg C, Leigh MW,
467 Loebinger MR, Mazurek H, Morgan L, Nielsen KG, Omran H, Schwert N, Scigliano S,
468 Werner C, Yiallourous P, Zivkovic Z, Lucas JS, Kuehni CE. The international primary ciliary
469 dyskinesia cohort (iPCD Cohort): methods and first results. *Eur Respir J* 49, 2017. doi:
470 10.1183/13993003.01181-2016.
- 471 15. Hjeij R, Onoufriadis A, Watson CM, Slagle CE, Klena NT, Dougherty GW, Kurkowiak M,
472 Loges NT, Diggle CP, Morante NFC, Gabriel GC, Lemke KL, Li Y, Pennekamp P, Menchen
473 T, Konert F, Marthin JK, Mans DA, Letteboer SJF, Werner C, Burgoyne T, Westermann C,
474 Rutman A, Carr IM, O'Callaghan C, Moya E, Chung EMK, Sheridan E, Nielsen KG,
475 Roepman R, Bartscherer K, Burdine RD, Lo CW, Omran H, Mitchison HM. CCDC151
476 Mutations Cause Primary Ciliary Dyskinesia by Disruption of the Outer Dynein Arm Docking
477 Complex Formation. *The American Journal of Human Genetics* 95: 257–274, 2014. doi:
478 10.1016/j.ajhg.2014.08.005.
- 479 16. Kispert A, Petry M, Olbrich H, Volz A, Ketelsen U-P, Horvath J, Melkaoui R, Omran H,
480 Zariwala M, Noone PG, Knowles M. Genotype-phenotype correlations in PCD patients
481 carrying DNAH5 mutations. *Thorax* 58: 552–554, 2003. doi: 10.1136/thorax.58.6.552-b.
- 482 17. Lin J, Yin W, Smith MC, Song K, Leigh MW, Zariwala MA, Knowles MR, Ostrowski LE,
483 Nicastro D. Cryo-electron tomography reveals ciliary defects underlying human RSPH1 primary
484 ciliary dyskinesia. *Nat Commun* 5: 5727, 2014. doi: 10.1038/ncomms6727.
- 485 18. Lucas JS, Barbato A, Collins SA, Goutaki M, Behan L, Caudri D, Dell S, Eber E, Escudier
486 E, Hirst RA, Hogg C, Jorissen M, Latzin P, Legendre M, Leigh MW, Midulla F, Nielsen KG,
487 Omran H, Papon J-F, Pohunek P, Redfern B, Rigau D, Rindlisbacher B, Santamaria F,
488 Shoemark A, Snijders D, Tonia T, Titieni A, Walker WT, Werner C, Bush A, Kuehni CE.
489 European Respiratory Society guidelines for the diagnosis of primary ciliary dyskinesia.
490 *European Respiratory Journal* 49: 1601090, 2017. doi: 10.1183/13993003.01090-2016.
- 491 19. Mitchison HM, Shoemark A. Motile cilia defects in diseases other than primary ciliary
492 dyskinesia: The contemporary diagnostic and research role for transmission electron
493 microscopy. *Ultrastruct Pathol* 41: 415–427, 2017. doi: 10.1080/01913123.2017.1370050.
- 494 20. Olbrich H, Häffner K, Kispert A, Völkel A, Volz A, Sasmaz G, Reinhardt R, Hennig S,
495 Lehrach H, Konietzko N, Zariwala M, Noone PG, Knowles M, Mitchison HM, Meeks M,
496 Chung EMK, Hildebrandt F, Sudbrak R, Omran H. Mutations in *DNAH5* cause primary ciliary
497 dyskinesia and randomization of left–right asymmetry. *Nature Genetics* 30: ng817, 2002. doi:
498 10.1038/ng817.
- 499 21. Olbrich H, Schmidts M, Werner C, Onoufriadis A, Loges NT, Raidt J, Banki NF, Shoemark
500 A, Burgoyne T, Al Turki S, Hurles ME, UK10K Consortium, Köhler G, Schroeder J,

- 501 **Nürnberg G, Nürnberg P, Chung EMK, Reinhardt R, Marthin JK, Nielsen KG, Mitchison**
 502 **HM, Omran H.** Recessive HYDIN mutations cause primary ciliary dyskinesia without
 503 randomization of left-right body asymmetry. *Am J Hum Genet* 91: 672–684, 2012. doi:
 504 10.1016/j.ajhg.2012.08.016.
- 505 22. **O'Toole ET, Giddings TH, Porter ME, Ostrowski LE.** Computer-assisted image analysis of
 506 human cilia and Chlamydomonas flagella reveals both similarities and differences in axoneme
 507 structure. *Cytoskeleton (Hoboken)* 69: 577–590, 2012. doi: 10.1002/cm.21035.
- 508 23. **Panizzi JR, Becker-Heck A, Castleman VH, Al-Mutairi DA, Liu Y, Loges NT, Pathak N,**
 509 **Austin-Tse C, Sheridan E, Schmidts M, Olbrich H, Werner C, Häffner K, Hellman N,**
 510 **Chodhari R, Gupta A, Kramer-Zucker A, Olale F, Burdine RD, Schier AF, O'Callaghan C,**
 511 **Chung EMK, Reinhardt R, Mitchison HM, King SM, Omran H, Drummond IA.** *CCDC103*
 512 mutations cause primary ciliary dyskinesia by disrupting assembly of ciliary dynein arms. *Nature*
 513 *Genetics* 44: 714–719, 2012. doi: 10.1038/ng.2277.
- 514 24. **Rutland J, Griffin WM, Cole PJ.** Human Ciliary Beat Frequency in Epithelium from
 515 Intrathoracic and Extrathoracic Airways. *Am Rev Respir Dis* 125: 100–105, 1982. doi:
 516 10.1164/arrd.1982.125.1.100.
- 517 25. **Schroeder JA.** Application of laboratory and digital techniques for visual enhancement during
 518 the ultrastructural assessment of cilia. *Ultrastructural Pathology* 41: 399–407, 2017. doi:
 519 10.1080/01913123.2017.1363335.
- 520 26. **Shapiro AJ, Davis SD, Polineni D, Manion M, Rosenfeld M, Dell SD, Chilvers MA, Ferkol**
 521 **TW, Zariwala MA, Sagel SD, Josephson M, Morgan L, Yilmaz O, Olivier KN, Milla C,**
 522 **Pittman JE, Daniels MLA, Jones MH, Janahi IA, Ware SM, Daniel SJ, Cooper ML, Nogee**
 523 **LM, Anton B, Eastvold T, Ehrne L, Guadagno E, Knowles MR, Leigh MW, Lavergne V.**
 524 Diagnosis of Primary Ciliary Dyskinesia. An Official American Thoracic Society Clinical Practice
 525 Guideline. *Am J Respir Crit Care Med* 197: e24–e39, 2018. doi: 10.1164/rccm.201805-0819ST.
- 526 27. **Shoemark A, Burgoyne T, Kwan R, Dixon M, Patel MP, Rogers AV, Onoufriadis A, Scully**
 527 **J, Daudvohra F, Cullup T, Loebinger MR, Wilson R, Chung EMK, Bush A, Mitchison HM,**
 528 **Hogg C.** Primary ciliary dyskinesia with normal ultrastructure: three-dimensional tomography
 529 detects absence of DNAH11. *Eur Respir J* 51, 2018. doi: 10.1183/13993003.01809-2017.
- 530 28. **Shoemark A, Dixon M, Corrin B, Dewar A.** Twenty-year review of quantitative transmission
 531 electron microscopy for the diagnosis of primary ciliary dyskinesia. *J Clin Pathol* 65: 267–271,
 532 2012. doi: 10.1136/jclinpath-2011-200415.
- 533 29. **Shoemark A, Frost E, Dixon M, Ollosson S, Kilpin K, Patel M, Scully J, Rogers AV,**
 534 **Mitchison HM, Bush A, Hogg C.** Accuracy of Immunofluorescence in the Diagnosis of Primary
 535 Ciliary Dyskinesia. *Am J Respir Crit Care Med* 196: 94–101, 2017. doi: 10.1164/rccm.201607-
 536 13510C.
- 537 30. **Shoemark A, Moya E, Hirst RA, Patel MP, Robson EA, Hayward J, Scully J, Fassad MR,**
 538 **Lamb W, Schmidts M, Dixon M, Patel-King RS, Rogers AV, Rutman A, Jackson CL,**
 539 **Goggin P, Rubbo B, Ollosson S, Carr S, Walker W, Adler B, Loebinger MR, Wilson R,**
 540 **Bush A, Williams H, Boustred C, Jenkins L, Sheridan E, Chung EMK, Watson CM, Cullup**
 541 **T, Lucas JS, Kenia P, O'Callaghan C, King SM, Hogg C, Mitchison HM.** High prevalence of
 542 *CCDC103* p.His154Pro mutation causing primary ciliary dyskinesia disrupts protein
 543 oligomerisation and is associated with normal diagnostic investigations. *Thorax* 73: 157–166,
 544 2018. doi: 10.1136/thoraxjnl-2017-209999.
- 545 31. **Takada S, Wilkerson CG, Wakabayashi K, Kamiya R, Witman GB.** The Outer Dynein Arm-
 546 Docking Complex: Composition and Characterization of a Subunit (Oda1) Necessary for Outer
 547 Arm Assembly. *MBoC* 13: 1015–1029, 2002. doi: 10.1091/mbc.01-04-0201.
- 548 32. **Wallmeier J, Shiratori H, Dougherty GW, Edelbusch C, Hjeij R, Loges NT, Menchen T,**
 549 **Olbrich H, Pennekamp P, Raidt J, Werner C, Minegishi K, Shinohara K, Asai Y, Takaoka K,**

- 550 **Lee C, Griesse M, Memari Y, Durbin R, Kolb-Kokocinski A, Sauer S, Wallingford JB,**
551 **Hamada H, Omran H.** TTC25 Deficiency Results in Defects of the Outer Dynein Arm Docking
552 Machinery and Primary Ciliary Dyskinesia with Left-Right Body Asymmetry Randomization. *The*
553 *American Journal of Human Genetics* 99: 460–469, 2016. doi: 10.1016/j.ajhg.2016.06.014.
- 554 33. **Wheway G, Mitchison HM.** Opportunities and Challenges for Molecular Understanding of
555 Ciliopathies—The 100,000 Genomes Project. *Front Genet* 10, 2019. doi:
556 10.3389/fgene.2019.00127.
- 557 34. **Zariwala MA, Omran H, Ferkol TW.** The Emerging Genetics of Primary Ciliary Dyskinesia.
558 *Proc Am Thorac Soc* 8: 430–433, 2011. doi: 10.1513/pats.201103-023SD.
- 559

561 Figure legends

562 **Fig 1.** Cilia ultrastructure. (A) Transmission electron microscopy image of a cross-section of healthy
563 9+2 motile cilium. (B) Diagram highlighting the different components of a cilium. (C) Diagram showing
564 the detailed structure of the central pair complex including the projections from the C1 and C2
565 microtubule. Scale bar (A) 100 nm.

566 **Fig 2.** PCD Detect workflow showing the function of the different programs in the toolkit and the order
567 in which to use them. Information is given in regard to the type of ciliary images required for PCD
568 Detect and more information about the *DNAH5* [1], *DNAH5* [4], *HYDIN* [2] and *HYDIN* [4] are provided
569 in Fig 3,4 and 7 and table 1.

570 **Fig 3.** Image averaging shows a clear absence of the ODA in subjects with pathogenic variants in
571 *DNAH5*. (A) TEM image of a cilium from a control subject and (B) cilia from three unrelated subjects
572 with pathogenic variants in *DNAH5*. (C) Average control MTD and (D) shows averaged MTD with
573 absent ODAs from the subjects carrying pathogenic variants in *DNAH5*. The white filled arrowheads
574 indicate MTD that have a projection emanating from the MTD, the white outlined arrowheads indicate
575 a complete absence of the ODA. Scale bars (A – B) 100nm and (C – D) 10nm

576 **Fig 4.** A subject with an atypical pathogenic variant in *DNAH5* has a sporadic absence of the ODA
577 leading to the generation of two averaged classes. (A) TEM image of a cilia from a control subject and
578 (B) a subject with a pathogenic variant of unknown significance (VUS) in *DNAH5* (compound
579 heterozygous for a missense mutation p.Thr3410Met, plus a stop-gain mutation p.Arg1761*). (C)
580 Two averaged classes generated for this subject shows the presence and absence (white filled
581 arrowheads) of the ODA. (D) In a subject with pathogenic variants in *DNAH5* (homozygous for the
582 stop-gain mutation p.Arg4429*) the class averages generated both show a loss of the ODA (white
583 filled arrowheads), indicating complete absence of the ODAs. (E) The difference which is the
584 subtraction of the two classes is shown in yellow (white outlined arrowhead) and is overlaid and
585 replaces part of the class 1 MTD image (red). Scale bars (A-B) 100nm (C-E) 10nm.

586 **Fig 5.** Analysis of MTDs from a subject with pathogenic variants in *DNAH9* gives two class averages
587 when using all cross-sectional images, but only single classes when splitting the images into proximal
588 or distal ciliary regions. (A) *DNAH11* instead of *DNAH9* is localised to the base of healthy cilia close
589 to the cell surface where microvilli are observed. Defects in *DNAH9* do not affect the ODA structure
590 within this region, making cilia cross-sections surrounded by microvilli ideal internal controls when
591 comparing to ODAs from the tip region. (B) TEM images at the proximal ciliary regions determined by
592 the presence of microvilli (black arrows) and (C) the distal ciliary region. (D) Average MTD from a
593 healthy control compared to (E – G) a subject with pathogenic variants in *DNAH9*. (E) All cross-
594 sectional ciliary images generate two classes showing the presence and absence of the ODA (white
595 filled arrowhead). Separating the ciliary images into two groups at the proximal and distal region of the
596 cilium generates single classes (F) showing the presence and (G) absence of the ODA (white filled
597 arrowhead). Scale bars (B-C) 100nm and (D-G) 10nm.

Fig 6. Feature averaging helps in the detection of the absence of the IDA in subjects that have pathogenic variants in *CCDC40* and *CCDC103*. TEM image of a cilium from (A) a healthy control subject and subjects with *CCDC40* (B) and *CCDC103* (C) mutations. (D-F) Averaged MTD from the different subjects. The pathogenic variants in (E) *CCDC40* and (F) *CCDC103* demonstrate a clear absence of the IDA when compared to (D) the healthy control MTD. Scale bars (A-C) 100nm and (D-F) 10nm

Fig 7. Subjects carrying pathogenic variants in *HYDIN* have two distinct CP complex structures. TEM images of cilia from (A) a control healthy subject, (B) three subjects (*HYDIN* [1-3]) with variants in *HYDIN* and one subject (*HYDIN* [4]) that has variants of unknown significance in *HYDIN*. (C-E) Average CP structures from the subjects examined. (D-E) Subjects with pathogenic *HYDIN* variants gave two class averages, showing both a complete CP complex and with an absence of the C2b component (white filled arrowheads) when compared to (C) the healthy control subject CP. (E) An overlay of genetically undiagnosed *HYDIN* [4] (blue) and the healthy control subject CP (red), highlights the missing C2b component (white outlined arrowhead) in *HYDIN* [4] confirming the likely pathogenicity of their *HYDIN* variants. Scale bars (A – B) 100nm and (C – E) 20nm.

615

616 **Supplementary information**

617 The PCD Detect programs have been built for both 32- and 64-bit windows machines and are include
618 in the supplementary zip files. The toolkit includes a guide (PCD Detect Guide.pdf) within the zip files
619 and there are also compressed test images. These files are as following; PCD Detect 32-bit part 1.zip,
620 PCD Detect 32-bit part 2.zip, PCD Detect 64-bit.zip, Test images for PCD Detect.zip. Supplementary
621 figures, table and programs are available to download from Figshare;
622 <https://figshare.com/s/bc7920e94ad14fba2ee8>.

623 <https://doi.org/10.5522/04/12327839>

Fig S1. PCD Detect Crop and Average screenshots. (A) Cilia Crop that has been designed to extract ciliary cross-sections from transmission electron microscopy images. (B) Cilia Average program used to pick cilia features (in this case MTDs) by performing a rotational and translational search before averaging them together.

Fig S2. Montages of some of the features (MTDs and CPs) of subjects carrying different gene mutations (Table 1) extracted by PCD Average before performing a rotation and translation search. Scale bars (A-I) 25nm.

Fig S3. Screenshots of PCD Detect Classification and Analysis programs. (A) Cilia Classification has been designed for subject samples that have a non-homogeneous defect in order to group images based on structural similarity, before generating a number of averaged outputs. (B) Analysis program designed to overlay images and highlight ciliary structural differences. The hollow arrowheads indicate the presence of the C1a and C2a CP complexes (see Fig. 1). Whereas, the white arrow shows the position of the C2b CP component that is absent in the HYDIN case (in blue) compared to the control (in red). (B1) The program interface showing an overlay of two average images of the CP, control in red and a HYDIN defect in blue. (B1-B4) Shows the different images that can be generated by the program when comparing two averaged images. (B2) A subtraction image showing the difference between the red and blue images. (B3) The subtraction image overlaid (in yellow) onto the red image (image 1). (B4) Image showing overlapping detail from the two images.

Fig S4. The outputs from hierarchy and PCA classification of MTDs. (A-B) Classification outputs for a subject that has an atypical pathogenic variant in *DNAH5* and (C-D) pathogenic variants in *DNAH5*. (A, C) Graphs generated for hierarchy classification show the feature that has been picked as the reference and the corresponding features used to generate the first class average. (B, D) Graphs of the first six PCA coefficients calculated from MTD features, as well as cluster analysis using k-means to group the features into two classes. The white filled arrowheads indicate absence of the ODA. Scale bar (B, D) 10nm.

Fig S5. Test of selected images with added noise that have been put through PCD Detect.

Fig S6. Examples of averaged structures generated from healthy controls. Scale bars (A) 100nm (B-C) 10nm.

Fig S7. Subjects predicted to have a potential HYDIN defect (PHD) were found to have present C2b CP component. (A) Cilia from a control healthy subject and (B) three subjects predicted to have a PHD. (C-D) Average CP structures from the subjects examined. When generating class averages

661 using the same settings only one class is given for Control [1] and PHD [1], whereas multiple classes
662 are given for PHD [2] and PHD [3]. This is likely to be due to a larger number of features input into
663 PCD detect for PHD [2-3] leading to the detection of minor variations of the features including image
664 tilt. Even so, there is no visible defect of the CP when comparing to the control. The white filled
665 arrowheads indicate the position of the C2b CP component. Scale bars (A – B) 100nm and (C – E)
666 20nm.

667

668 **PCD Detect 32-bit part 1.zip.** Zip file containing the first half of the 32-bit PCD Detect programs.

669

670 **PCD Detect 32-bit part 2.zip.** Zip file containing the second half of the 32-bit PCD Detect programs.

671

672 **PCD Detect 64-bit.zip.** Zip file containing the 64-bit PCD Detect programs.

673

674 **Test images for PCD Detect.zip.** Zip file containing test images for PCD Detect.

675

676 **PCD Detect Guide.pdf** Step-by-step instructions on installing and using the PCD Detect programs.

Figure 1

PCD Detect Workflow

Considerations before inputting images into PCD Detect

- a) Stained sample that has good image contrast
- b) Little or no background and stain artefacts
- c) Well orientated ciliary cross sections
- d) Image taken at a single magnification
- e) Images focussed so that as ultrastructural detail can be seen as possible

Recommendation to check for distal vs proximal defects

- a) Manually separate images at the base of cilia (visible microvilli) and the rest of ciliary structure (no nearby microvilli); see Fig 4

Optional consideration

- a) Separate images based on the cell the cilia originate from (was not used in this study)

Figure 2

Figure 3

Figure 5

Figure 6

Figure 7

Subject	Patients symptoms and clinical diagnostic criteria	Average nNO nL min ⁻¹ (Normal nasal nitric oxide (nNO) >77 nL min ⁻¹)	Average light microscopy CBF (Hz)	No of nasal brushing to make PCD diagnosis	Genotype	Number of ciliary images input into PCD detect to generate MTD and/or CP averages	Number of Features selected in PCD detect	Features used by PCD detect to generate final average
Control [1]		-	-	-	-	MTD – 13 CP - 29	MTD – 112 CP – 29	MTD - 13 CP -15
Control [2]		-	-	-	-	MTD – 9 CP - 24	MTD - 76 CP -24	MTD - 22 CP – 15
Control [3]		-	-	-	-	MTD – 11 CP - 25	MTD - 99 CP - 25	MTD - 32 CP – 11
<i>DNAH5</i> [1]	Situs inversus, neonatal respiratory distress at term and persistent rhinitis from birth	122	0 (static)	2	DNAH5 NM_001369.2: c.13285C>T (p.Arg4429*) + c.13285C>T (p.Arg4429*)	MTD – 13	MTD - 117	Class 1 -14 Class 2 -13
<i>DNAH5</i> [2]	Persistent rhinitis and prolonged otorrhoea post grommet insertion	19	0 (static)	2	Unsolved - Heterozygous DNAH5 NM_001369.2: c.7477G>C (p.Glu2493Arg)	MTD – 13	MTD - 109	MTD - 18
<i>DNAH5</i> [3]	Situs inversus, neonatal respiratory distress at term and early onset rhinitis	Too young	0 (static)	1	Unsolved - Heterozygous DNAH5 NM_001369.2: c.6261T>G (p.Tyr2087*)	MTD – 11	MTD - 91	MTD - 19
<i>DNAH5</i> [4]	Rhinitis, recurrent ear infections and otorrhoea	775 (± 430 SEM)	6.98 (± 0.76 SEM)	2	DNAH5 NM_001369.2: c.10229C>T (p.Thr3410Met) + c.5281C>T (p.Arg1761*)	MTD – 15	MTD - 135	Class 1 - 21 Class 2 -19
<i>DNAH9</i>	Congenital heart disease, heterotaxy, rhinitis and recurrent ear infections	306 (± 48 SEM)	9.10 (± 0.36 SEM)	4	DNAH9 NM_001372: c.12367G>A (p.Asp4123Asn) + c.12367G>A (p.Asp4123Asn)	All - 16 Base -14 Tip - 7	All - 142 Base -125 Tip - 63	All Class 1 - 35 All Class 2 - 25 Base - 49 Tip - 32
<i>CCDC40</i>	-	-	-	2	CCDC40 NM_017950.3: c.2712-1G>T + c.2712-1G>T	MTD - 14	MTD - 119	MTD - 48
<i>CCDC103</i>	Situs inversus, neonatal respiratory distress, rhinitis and otitis media	250 (± 50 SEM)	8.56 (± 0.58 SEM)	4	CCDC103 NM_001258395.1: c.461A>C (His154Pro) + c.461A>C (His154Pro)	MTD - 14	MTD - 118	Class 1 - 46 Class 2 - 34
<i>HYDIN</i> [1]	Normal situs, chronic wet cough, rhinitis and Otorrhoea.	30	8.94	3	HYDIN NM_001270974.1: c.8487del (p.Pro2830Hisfs*23) + c.8489C>A (p.Pro2830Gln)	CP – 21	CP - 21	Class 1 – 6 Class 2 – 3

Subject	Patients symptoms and clinical diagnostic criteria	Average nNO nL min ⁻¹ (Normal nasal nitric oxide (nNO) >77 nL min ⁻¹)	Average light microscopy CBF (Hz)	No of nasal brushing to make PCD diagnosis	Genotype	Number of ciliary images input into PCD detect to generate MTD and/or CP averages	Number of Features selected in PCD detect	Features used by PCD detect to generate final average
HYDIN [2]	Neonatal respiratory distress, early onset rhinitis, otitis media and otorrhoea	31	5.98	3	HYDIN NM_001270974.1: c.13709del (p.Pro4570Leufs*2) + c.13709del (p.Pro4570Leufs*2)	CP - 35	CP - 25	Class 1 - 19 Class 2 - 4
HYDIN [3]	Neonatal respiratory distress, rhinosinusitis, poor hearing, immotile sperm and retinitis pigmentosa	5	9.86	2	Unsolved - Heterozygous HYDIN NM_001270974.1: c.3786-1G>T	CP - 15	CP - 15	Class 1 - 11 Class 2 - 6
HYDIN [4]	wet cough and intermittent rhinitis	442 (± 239 SEM)	9.95 (± 0.42 SEM)	3	HYDIN NM_001270974.1: c.1491G>C (p.Met497Ile) + c.2702G>T (p.Gly901Val)	CP - 27	CP - 27	Class 1 - 8 Class 2 - 6

Table 1. Diagnostic, genetic and image process information for the subjects involved in this study. Normal nasal nitric oxide (nNO) >77 nL min⁻¹.