

Ramsey goodness of paths

Alexey Pokrovskiy*

Benny Sudakov[†]

Abstract

Given a pair of graphs G and H , the Ramsey number $R(G, H)$ is the smallest N such that every red-blue coloring of the edges of the complete graph K_N contains a red copy of G or a blue copy of H . If graph G is connected, it is well known and easy to show that $R(G, H) \geq (|G| - 1)(\chi(H) - 1) + \sigma(H)$, where $\chi(H)$ is the chromatic number of H and σ the size of the smallest color class in a $\chi(H)$ -coloring of H . A graph G is called H -good if $R(G, H) = (|G| - 1)(\chi(H) - 1) + \sigma(H)$. The notion of Ramsey goodness was introduced by Burr and Erdős in 1983 and has been extensively studied since then. In this short note we prove that n -vertex path P_n is H -good for all $n \geq 4|H|$. This proves in a strong form a conjecture of Allen, Brightwell, and Skokan.

1 Introduction

Given a pair of graphs G and H , the Ramsey number $R(G, H)$ is the smallest N such that every red-blue coloring of the edges of the complete graph K_N contains a red copy of G or a blue copy of H . It is a corollary of the celebrated theorem of Ramsey that these numbers are always finite. Let $\chi(H)$ be the chromatic number of H , i.e. the smallest number of colors needed to color the vertices of H so that no pair of adjacent vertices have the same colour, and $\sigma(H)$ be the the size of the smallest color class in a $\chi(H)$ -colouring of H . It was observed by Burr [4] that for connected G with $|G| \geq \sigma(H)$ Ramsey numbers always satisfy the following easy lower bound

$$R(G, H) \geq (|G| - 1)(\chi(H) - 1) + \sigma(H). \quad (1)$$

To prove (1), consider a 2-edge-coloring of the complete graph on $N = (|G| - 1)(\chi(H) - 1) + \sigma(H) - 1$ vertices consisting of $\chi(H) - 1$ disjoint red cliques of size $|G| - 1$ as well as one disjoint red clique of size $\sigma(H) - 1$. This coloring has no red G because all red connected components have size $\leq |G| - 1$, and there is no blue H since the partition of this H induced by red cliques would give a coloring of H by $\chi(H)$ colors with one color class smaller than $\sigma(H)$, contradicting the definition of $\sigma(H)$.

For some graphs the bound in (1) is quite far from the truth. For example Erdős [8] showed that $R(K_n, K_n) \geq \Omega(2^{n/2})$ which is much larger than the quadratic bound we get from (1). However there are many known pairs of graphs for which $R(G, H) = (|G| - 1)(\chi(H) - 1) + \sigma(H)$. In this case we say that G is H -good. The notion of Ramsey goodness was introduced by Burr and Erdős [5] in 1983 and was extensively studied since then, see, e.g., [1, 6, 7, 11, 13, 14] and their references.

In this short note we study the question of when the n -vertex path P_n is H -good, for some fixed graph H . This problem goes back to the work of Erdős, Faudree, Rousseau, and Schelp [9], who in 1985 proved that there is a function f such that P_n is H -good for all $n \geq f(|H|)$. The function $f(|H|)$ is not explicit in [9], but $f(H) = O(|H|^4)$ can be proved using their method. Häggkvist [12] (for $k = 2$) and later Pokrovskiy [16] obtained a general upper bound on the Ramsey number of path

*Department of Mathematics, ETH, 8092 Zurich, Switzerland. dr.alexey.pokrovskiy@gmail.com. Research supported in part by SNSF grant 200021-149111.

[†]Department of Mathematics, ETH, 8092 Zurich, Switzerland. benjamin.sudakov@math.ethz.ch. Research supported in part by SNSF grant 200021-149111.

versus complete k -partite graphs, showing that $R(P_n, K_{m_1, \dots, m_k}) \leq (k-1)(n-1) + km - k + 1$. Here and later, K_{m_1, \dots, m_k} denotes a complete k -partite graph with parts of order m_1, \dots, m_k together with all the edges connecting vertices in different parts. Although this bound is not strong enough to prove goodness, it holds for all values of the parameters. More recently, Pei and Li [15] showed that if $n \geq 8|H| + 3\sigma(H)^2 + c\chi^8(H)$, then P_n is H -good. For general H (e.g., when $H = K_{m,m}$) this result requires n to be quadratic in $|H|$. Allen, Brightwell, and Skokan [1] conjectured that P_n is H -good already when n is linear in $|H|$.

Conjecture 1.1 ([1]). *Let H be a fixed graph with chromatic number k and let $n \geq k|H|$. Then $R(P_n, H) = (n-1)(k-1) + \sigma(H)$.*

Let $R(C_{\geq n}, H)$ be the smallest N such that any 2-edge-coloring of K_N contains either a red cycle of length at least n or a blue H . Notice that we always have $R(P_n, H) \leq R(C_{\geq n}, H)$. Motivated by the above conjecture, in this note we prove the following theorem.

Theorem 1.2. *Given integers $m_1 \leq m_2 \leq \dots \leq m_k$ and $n \geq 3m_k + 5m_{k-1}$, we have*

$$R(C_{\geq n}, K_{m_1, \dots, m_k}) = (k-1)(n-1) + m_1.$$

Notice that the vertices of a k -chromatic graph H can be partitioned into k independent sets of sizes m_1, \dots, m_k with $\sigma(H) = m_1 \leq m_2 \leq \dots \leq m_k$. This is equivalent to H being a subgraph of K_{m_1, \dots, m_k} . Since $4|H| \geq 4m_k + 4m_{k-1} \geq 3m_k + 5m_{k-1}$, Theorem 1.2 implies the following.

Corollary 1.3. *Let H be a fixed graph with chromatic number k and let $n \geq 4|H|$. Then $R(P_n, H) = (n-1)(k-1) + \sigma(H)$.*

For $k \geq 4$, this corollary proves Conjecture 1.1 in a very strong form, showing that the condition $n \geq \chi(H)|H|$ is unnecessary, and $n \geq 4|H|$ suffices. For $k \leq 3$, our result is slightly weaker than the conjecture, but is a large improvement on the best previously known [15] quadratic dependence of n on $|H|$. Moreover, for certain graphs H , Theorem 1.2 shows that P_n is H -good even when n is smaller than $4|H|$. For example if H is balanced (i.e. if $|H| = \sigma(H)\chi(H)$), then this theorem implies that P_n is H -good as long as $n \geq 8|H|/\chi(H)$.

2 Proof of the main theorem

The proof of Theorem 1.2 uses a combination of the Pósa rotation-extension technique and induction on k . Let $P = p_1p_2 \dots p_t$ be a path in a graph G which also contains an edge $p_t p_i$. We say that a path $Q = p_1p_2 \dots p_i p_t p_{t-1} \dots p_{i+1}$ is a rotation of P . We say that a path Q is *derived* from P if there is a sequence of paths $P_0 = P, P_1, \dots, P_s = Q$ with P_j being a rotation of P_{j-1} for each j . We say that a vertex x is an ending vertex for P if it is the final vertex of some path derived from P . We say that a set X of ending vertices of P is *connected* if for every $x \in X$ there is a path P_x ending with x and a sequence of paths $P = P_0, P_1, \dots, P_s = P_x$ with P_j being a rotation of P_{j-1} for each j and P_j ending with some $x' \in X$. That is, for a P_x ending in $x \in X$, the set X also contains ending vertices of a sequence of intermediate paths whose rotations produce P_x . Notice that for a path P , the set of all ending vertices for P is connected. For a set of vertices S in a graph G , we use $N_G(S)$ to mean the set of vertices $v \in V(G) \setminus S$ for which there is a vertex $s \in S$ such that vs is an edge of G i.e $N_G(S)$ is the neighbourhood of the set S outside S . When there is no ambiguity of what the underlying graph is, we abbreviate this to $N(S)$.

We will need a variant of the celebrated lemma of Pósa from [18].

Lemma 2.1. *Let P be a path of maximum length in a graph G starting at some vertex p_0 , X a connected set of ending vertices for P , and S the set of all ending vertices for P . Then we have*

(i) $|N(X)| \leq 2|S|$.

(ii) *There is a cycle in G containing $N(X) \cup X$.*

Proof. Fix an orientation of P . Let S^- be the set of left neighbours on P of vertices in S and let S^+ be the set of right neighbours on P of vertices in S . Notice that by maximality of $|P|$, we have $N(X) \subseteq V(P)$ (indeed otherwise if we have $y \in N(X) \setminus V(P)$, then we can obtain a longer path starting from p_0 by choosing a rotation of P ending with a neighbour of y , and extending it with y .) First we will show that $N(X) \cup X \subseteq S^- \cup S^+ \cup X$.

Let x be a vertex in X and $P = P_0, P_1, \dots, P_s$ be a sequence of paths such that P_s ends with x , P_i ends with some $v \in X$, and P_i is a rotation of P_{i-1} (such a sequence exists since X is a connected set of endpoints.) Notice that if $yz \in E(P_s)$, then either $yz \in E(P)$ or $\{y, z\} \subseteq X \cup X^- \cup X^+$ (this is proved by induction using the fact that from the definition of “rotation”, we have $yz \in P_i \implies yz \in P_{i-1}$ or $\{y, z\} \subseteq X \cup X^- \cup X^+$.)

Let $p_0, \dots, p_k = x$ be the vertex sequence of P_s . For $p_i \in N(x)$, we know $p_1 p_2 \dots p_i p_k p_{k-1} \dots p_{i+1}$ is a rotation of P and hence $p_{i+1} \in S$. Since $p_i p_{i+1} \in E(P_s)$, we have that either $p_i p_{i+1} \in E(P)$ which implies that $p_i \in S^- \cup S^+ \subseteq X \cup S^- \cup S^+$, or we have $\{p_i, p_{i+1}\} \subseteq X \cup X^- \cup X^+ \subseteq X \cup S^- \cup S^+$. This implies that $N(X) \subseteq \bigcup_{x \in X} N(x) \subseteq X \cup S^- \cup S^+$ as required. Now part (i) of the lemma comes from $|N(X) \cup X| \leq |S^-| + |S^+| + |X| \leq 2|S| + |X|$.

Although part (ii) of the lemma already appeared in [3], we include its short proof for the sake of completeness. Let p_0, \dots, p_k be the vertex sequence of P , and let p_i be the first element of $(N(X) \cup X) \cap \{p_0, \dots, p_k\}$. Since we are considering paths starting from p_0 , we must have $p_0 \notin X$. If $p_0 \in N(X)$, then we have a cycle on $V(P)$ formed by taking a rotation of P ending with $p \in N(p_0)$ and adding the edge $p_0 p$. Therefore, we can assume that $p_i \neq p_0$. Notice that since $p_{i-1} \in N(p_i)$ we must have $p_i \notin X$ (otherwise p_{i-1} would be an element of $(N(X) \cup X) \cap \{p_0, \dots, p_k\}$ preceding p_i) which implies that $p_i \in N(X)$. Let $x \in X$ be a neighbour of p_i , and let Q be a rotation of P ending with x . Let $p_0 q_1 \dots q_k$ be the vertex sequence of Q . Recall that for all i either $q_t q_{t+1} \in E(P)$ or $\{q_t, q_{t+1}\} \subseteq X^- \cup X^+ \cup X$. Since Q starts with p_0 , and $X \cap \{p_0, \dots, p_i\} = \emptyset$, each of the edges $p_0 p_1, \dots, p_{i-1} p_i$ must be edges of Q , and so Q starts with the sequence $p_0 p_1 \dots p_i$. Therefore letting C be the cycle formed from $Q \setminus \{p_0, \dots, p_{i-1}\}$ by adding the edge $x p_i$ gives a cycle containing $X \cup N(X)$. \square

We now prove the main result of this note.

Proof of Theorem 1.2. The lower bound of the theorem follows from (1), so it remains to show that $R(C_{\geq n}, K_{m_1, \dots, m_k}) \leq (k-1)(n-1) + m_1$. The proof is by induction on k . In case $k = 1$ we need to find either a red copy of $C_{\geq n}$ or a blue 1-partite graph K_m^1 on m vertices. Since such a graph contains no edges its copy exist in any 2-edge-coloring of K_m . This implies that $R(H, K_m^1) \leq m$ for any graph H .

Now let $k \geq 2$ and suppose that the theorem holds for all $k' < k$. Let Γ be a 2-edge-colored complete graph on $(k-1)(n-1) + m_1$ vertices. Suppose that Γ contains no blue K_{m_1, \dots, m_k} and let G be the subgraph spanned by the red edges of Γ .

Claim 2.2. *For any set B with $|B| \geq m_1$, we have $|N_G(B) \cup B| \geq n - m_2 + m_1$.*

Proof. Suppose that $|N_G(B) \cup B| \leq n - m_2 + m_1 - 1$. Let $\Gamma' = \Gamma \setminus (N_G(B) \cup B)$. We have $|\Gamma'| = |\Gamma| - |N_G(B) \cup B| \geq (k-2)(n-1) + m_2$. By induction Γ' either contains a red $C_{\geq n}$ or a blue K_{m_2, \dots, m_k} . In the second case, since $|B| \geq m_1$, and all the edges between B and Γ' are blue, B can be joined to K_{m_2, \dots, m_k} in order to produce a blue K_{m_1, \dots, m_k} . \square

Let A be a maximum size subset of G satisfying $|A| < 2m_1$ and $|N_G(A)| \leq 2|A|$. This implies that $|N_G(A) \cup A| \leq 3|A| < 6m_1 < n - m_2 + m_1$, and so by Claim 2.2 we have $|A| < m_1$. Let G' be the induced subgraph of G on $V(G) \setminus A$. Note that every subset X with $|X| \leq m_1$ in G' satisfies that $|N_{G'}(X)| > 2|X|$ —indeed otherwise we would have

$$|N_G(X \cup A)| \leq |N_G(X) \setminus A| + |N_G(A)| = |N_{G'}(X)| + |N_G(A)| \leq 2|X| + 2|A| = 2|A \cup X|.$$

This would contradict the maximality of A .

Let P be a maximum length path in G' , and S the set of ending vertices for P . By Lemma 2.1 we have $|N_{G'}(S)| \leq 2|S|$ which implies that $|S| \geq m_1$. Let X be a connected set of exactly m_1 ending vertices for P . This set can be obtained by repeatedly applying rotations, adding one new ending vertex at a time until we have m_1 ending vertices. This guarantees that the resulting set is connected, since we always keep all previous ending vertices. From Claim 2.2 we obtain

$$|N_{G'}(X) \cup X| \geq n - m_2 + m_1 - |A| \geq 2m_2 + 5m_1.$$

Since $|X| = m_1$ and $|N_{G'}(X) \cup X| = |N_{G'}(X)| + |X|$, the above is equivalent to $|N_{G'}(X)| \geq 2m_2 + 4m_1$. Combining this with Lemma 2.1 gives

$$|S| \geq \frac{1}{2}|N_{G'}(X)| \geq m_2 + 2m_1.$$

Since A has at most $2|A| \leq 2m_1$ neighbours in G , we can choose $S' \subseteq S$ with $|S'| = m_2$ such that there are no edges between S' and A in G . Note that then $N_{G'}(S') = N_G(S')$.

By Lemma 2.1, there is a cycle C in G' containing $S' \cup N_{G'}(S') = S' \cup N_G(S')$. If $|C| \geq n$, then the complete graph Γ contains a red cycle of length at least n . Otherwise, if $|C| \leq n - 1$, then let $\Gamma' = \Gamma \setminus (S' \cup N_G(S'))$. We have $|\Gamma'| \geq |\Gamma| - |C| \geq (k - 2)(n - 1) + m_1$. Therefore, by induction, Γ' contains a blue K_{m_1, m_3, \dots, m_k} . All the edges between S' and Γ' are blue, and so S' can be joined to K_{m_1, m_3, \dots, m_k} in order to produce a blue K_{m_1, \dots, m_k} . \square

3 Concluding remarks

An interesting open question is to determine all n for which P_n is H -good. It is possible to show that P_n is not K_{m_1, m_2} -good when $n \leq 2m_2 - 2$. To see this, let $n = 2m_2 - 2$, $N = n + m_1 - 1$ and consider the edge coloring of the complete graph K_N , consisting of two blue cliques of orders $m_2 + m_1 - 1$ and $m_2 - 2$ respectively with all the edges between them red. It is easy to see that this coloring contains no red path P_n and no blue K_{m_1, m_2} . This implies that constant “4” in Corollary 1.3 cannot be made less than 2.

Our main result gives either a red cycle of length at least n or a blue copy of H . One could ask whether under the same conditions, we can guarantee a cycle of length *exactly* n . Indeed, this was asked by Allen, Brightwell, and Skokan [1], who conjectured that $R(C_n, H) = (n - 1)(\chi(H) - 1) + \sigma(H)$ when $n \geq \chi(H)|H|$. In a forthcoming paper [17], we will prove a strengthening of this conjecture for large $\chi(H)$ and $\sigma(H)$. Our result shows that C_n is H -good when $n \geq C|H|$ for some constant C and $\sigma(H) \geq \chi(H)^{11}$.

We also like to mention a related problem of Erdős, Faudree, Rousseau, and Schelp [10], who conjectured that $R(C_n, K_m) = (n - 1)(m - 1) + 1$ for $n \geq m$. The best known bound for this conjecture is due to Nikiforov [13], who proved it when $n \geq 4m + 1$.

Since paths are special case of trees, it is natural to consider the Ramsey goodness problem for trees as well. For example an old result of Chvátal [6] says that any tree T is good for every complete

graph. Motivated by question of Erdős, Faudree, Rousseau and Schelp, in the recent paper [2] we study the Ramsey goodness of bounded degree trees with respect to general graphs.

Acknowledgment. Part of this work was done when the second author visited Freie University Berlin. He would like to thank Humboldt Foundation for a generous support during this visit and Freie University for its hospitality and stimulating research environment. Both authors would like to thank J. Skokan, M. Stein and D. Hefetz for helpful discussions, as well as D. Korandi for finding a mistake in an earlier draft of this note.

References

- [1] P. Allen, G. Brightwell and J. Skokan. Ramsey-goodness and otherwise. *Combinatorica* **33** (2013), 125–160.
- [2] I. Balla, A. Pokrovskiy and B. Sudakov, Ramsey goodness of bounded degree trees, preprint.
- [3] S. Brandt, H. Broersma, R. Diestel, M. Kriesell. Global connectivity and expansion: long cycles and factors in f -connected graphs. *Combinatorica* **26** (2006), 17–36.
- [4] S. Burr. Ramsey numbers involving graphs with long suspended paths. *J. London Math. Soc.* **24** (1981), 405–413.
- [5] S. Burr and P. Erdős. Generalizations of a Ramsey-theoretic result of Chvátal. *J. Graph Theory* **7** (1983), 39–51.
- [6] V. Chvátal. Tree-complete graph Ramsey number. *J. Graph Theory* **1** (1977), 93.
- [7] D. Conlon, J. Fox, C. Lee and B. Sudakov. Ramsey numbers of cubes versus cliques. *Combinatorica*, to appear.
- [8] P. Erdős. Some remarks on the theory of graphs. *Bull. Am. Math. Soc.* **53** (1947), 292–294.
- [9] P. Erdős, R. Faudree, C. Rousseau and R. Schelp. Multipartite graph-sparse graph Ramsey numbers. *Combinatorica* **5** (1985), 311–318.
- [10] P. Erdős, R. Faudree, C. Rousseau and R. Schelp. On cycle-complete graph Ramsey numbers. *J. Graph Theory* **2** (1978), 53–64.
- [11] G. Fiz Pontiveros, S. Griffiths, R. Morris, D. Saxton and J. Skokan. The Ramsey number of the clique and the hypercube. *J. Lond. Math. Soc.* **89** (2014), 680–702.
- [12] R. Häggkvist. On the path-complete bipartite Ramsey number. *Discrete Math.* **75** (1989), 243–245.
- [13] V. Nikiforov. The cycle-complete graph Ramsey numbers. *Combin. Probab. Comput.* **14** (2005), 349–370.
- [14] V. Nikiforov and C. Rousseau. Ramsey goodness and beyond. *Combinatorica* **29** (2009), 227–262.
- [15] C. Pei and Y. Li. Ramsey numbers involving a long path. *Discrete Math.* **339** (2016), 564–570.
- [16] A. Pokrovskiy. Calculating Ramsey numbers by partitioning coloured graphs, preprint.
- [17] A. Pokrovskiy and B. Sudakov. Ramsey goodness of cycles, preprint.
- [18] L. Pósa. Hamiltonian circuits in random graphs. *Discrete Math.* **14** (1976), 359–364.