

Authors:

Petzold, Axel; Moorfields Eye Hospital NHS Foundation Trust and The National Hospital for Neurology and Neurosurgery, Queen Square, Institute of Neurology, University College London; Amsterdam UMC - Locatie VUMC, Neurology and Ophthalmology;

Biousse, Valerie; Emory University, Ophthalmology;

Bursztyn, Lulu; Western University;

Costello, Fiona; University of Calgary, Clinical Neurosciences

Crum, Alison; Utah System of Higher Education

Digre, Kathleen; John Moran Eye Center - University of Utah, Department of Neurology- Ophthalmology and Visual Science

Fraser, Clare; Save Sight Institute, The University of Sydney

Fraser, J. Alex; Department of Clinical Neurological Sciences and Department of Ophthalmology, Western University, London, Ontario, Canada

Katz, Bradley; Utah System of Higher Education

Jurkute, Neringa; UCL, Institute of Ophthalmology

Newman, Nancy; Emory University, School of Medicine, Department of Neuro-Ophthalmology

Lautrup-Battistini, Jette; Copenhagen University Hospital Glostrup

Lawlor, Mitchell; The University of Sydney, Save Sight Institute

Liskova, Petra; Charles University, Ophthalmology

Lorenz, Birgit; Justus Liebig University Giessen, Ophthalmology

Malmqvist, Lasse; Rigshospitalet Glostrup, Ophthalmology

Pergallo, Jason; Emory University, Ophthalmology

Sibony, Patrick; Stony Brook University

Subramanian, Prem; Sue Anschutz-Rodgers UC Health Eye Center, University of Colorado School of Medicine

Rejdak, Robert; Lublin, Ophthalmology

Touitou, Valerie; Groupe Hospitalier La Pitié Salpêtrière-Charles Foix,
DHU Vision et Handicaps

Warner, Judith; University of Utah Health Sciences Center, Department
of Ophthalmology and Visual Sciences

Wegener, Marianne; Rigshospitalet Glostrup, Ophthalmology

Wong, Sui; Moorfields Eye Hospital City Road Campus

Yu-Wai-Man, Patrick; Cambridge University; Moorfields Eye Hospital NHS
Foundation Trust, Medical Retina

Hamann, Steffen; Rigshospitalet Glostrup, Ophthalmology

ODDS consortium and ERN-EYE

Abstract

Peripapillary hyperreflective ovoid mass-like structures (PHOMS) are a new retinal optical coherence tomography (OCT) finding. The *Optic Disc Drusen Studies Consortium* had made recommendations to distinguish PHOMS from true optic disc drusen (ODD) in 2018. While publications on PHOMS have increased since then, the accuracy of the definition of PHOMS and reliability of detection is unknown. In this multi-rater study we demonstrate that the 2018 definition of PHOMS resulted in a poor multi-rater kappa of 0.356. We performed a Delphi consensus process to develop a consistent and refined definition of PHOMS with clear principles around the nature of PHOMS and how they differ from normal anatomy. Fifty explanatory teaching slides, provided as supplementary material, allowed our expert group of raters to achieve a good level of agreement (kappa 0.701, 50 OCT scans, 21 raters). We recommend adopting the here refined definition for PHOMS.

Key words: OCT, ODD, PHOMS

To the Editor: in 2018 a consortium of ophthalmologists and neurologists reported a novel optical coherence tomography (OCT) finding, peripapillary hyperreflective ovoid mass-like structures (PHOMS) [1]. The early definition of PHOMS was made in the context that they often occur alongside optic disc drusen (ODD), but there are several features to demonstrate that PHOMS are distinct from ODD and should not be confused with ODD [2; 3]. This definition did not define PHOMS in other contexts or distinguish PHOMS from normal anatomical variants. The definition of PHOMS has traditionally been one of exclusion [1; 3]. In the present multirater validation exercise, we determined the accuracy of our definition. A total of 100 OCT optic disc scans were reviewed for presence of PHOMS by 15 independent raters. There were three possible answers: (I) yes, (ii) no, (iii) possible. The level of agreement was determined by Fleiss multi-rater kappa in R. The overall multi-rater kappa was poor at 0.356.

We reviewed the disagreements and conducted a Delphi approach amongst the authors to create an internally consistent definition of PHOMS. At the end of this process we defined PHOMS by:

- i. *location*: strictly peripapillary and sitting on top of Bruch's membrane in the OCT B-scan. Frequently; a gap can be observed in the B-scan scans of PHOMS aligned through the centre of the optic disc;
- ii. *effect on adjacent retinal layers*: there is typically an upward deflection (like a ski slope, see supplementary material) of at least two of the other retinal layers;
- iii. *signal* appearance which is similar to the reflectivity to the retinal nerve fiber and ganglion cell layers because there is evidence that they may represent axonal stasis [4], and because the normal egress of axons at the optic disc can resemble PHOMS. The latter is can be observed in tilted discs, discs with an elevated border

and myopia [5]. This downward slope of essentially normal axons can easily be mistaken for PHOMS.

After this process a repeat two-rater assessment (AP, SH) was performed using this refined definition of PHOMS on 106 OCT optic nerve B-scans and revealed an excellent kappa of 0.811. Six scans remained ambiguous even after discussion among raters (for example slides #33 and #34 in the supplementary material). We then created fifty teaching slides to explain the new consensus definition (supplementary material) and after reviewing these slides our 21 expert raters were then asked to review 50 more OCT B-scans.

A good level of agreement was achieved with a kappa of 0.701.

In conclusion, a good inter-rater agreement can be achieved for PHOMS even on a single B-scan on optic nerve OCT. Future studies should test joint access to the entire volume scans on a shared OCT viewer platform which will have to meet contemporary Information Governance and Data Protection requirements. Importantly, the image characteristics of PHOMS have been defined more succinctly and teaching slides are provided (see supplementary material). As PHOMS are increasingly being recognised in a range of pathologies, we recommend adopting our proposed definition.

References

[1] Malmqvist, L.; Bursztyn, L.; Costello, F.; Digre, K.; Fraser, J. A.; Fraser, C.; Katz, B.; Lawlor, M.; Petzold, A.; Sibony, P.; Warner, J.; Wegener, M.; Wong, S. and Hamann, S. (2018). *The Optic Disc Drusen Studies Consortium Recommendations for Diagnosis of Optic Disc Drusen Using Optical Coherence Tomography*, *Journal of Neuro-Ophthalmology* 38 : 299-307.

- [2] Lee, K. M.; Woo, S. J. and Hwang, J.-M. (2018). *Peripapillary Hyperreflective Ovoid Mass-Like Structures: Is It Optic Disc Drusen or Not?*, Journal of Neuro-Ophthalmology 38 : 567-568.
- [3] Malmqvist, L.; Bursztyn, L.; Costello, F.; Digre, K.; Fraser, J. A.; Fraser, C.; Katz, B.; Lawlor, M.; Petzold, A.; Sibony, P.; Warner, J.; Wegener, M.; Wong, S. and Hamann, S. (2018). *Peripapillary Hyperreflective Ovoid Mass-Like Structures*, Journal of Neuro-Ophthalmology 38 : 568-570.
- [4] Wang, X.; Lou, N.; Eberhardt, A.; Yang, Y.; Kusk, P.; Xu, Q.; Förster, B.; Peng, S.; Shi, M.; Ladrón-de-Guevara, A. and others (2020). *An ocular glymphatic clearance system removes β -amyloid from the rodent eye*, Science Translational Medicine 12.
- [5] Sibony, P.; Bioussé, V.; Bursztyn, L.; Costello, F.; Digre, K.; Fraser, J.; Fraser, C.; Huna-Baron, R.; Katz, B.; Lawlor, M.; Malmqvist, L.; Newman, N.; Peragallo, J.; Petzold, A.; Subramanian, P.; Warner, J.; Wegener, M.; Wong, S. and Hamann, S. (). *Comment on: Morphologic Features of Buried Optic Disc Drusen on En Face Optical Coherence Tomography and Optical Coherence Tomography Angiography*, American Journal of Ophthalmology .