

[Open Peer Review on Qeios](#)

The association of smoking status with SARS-CoV-2 infection, hospitalisation and mortality from COVID-19: A living rapid evidence review with Bayesian meta-analyses (version 11)

David Simons¹, Lion Shahab², Jamie Brown², Olga Perski²

¹ Royal Veterinary College, RVC

² University College London, University of London

Funding: The author(s) received no specific funding for this work.

Potential competing interests: The author(s) declared that no potential competing interests exist.

Abstract

Aims: To estimate the association of smoking status with rates of i) infection, ii) hospitalisation, iii) disease severity, and iv) mortality from SARS-CoV-2/COVID-19 disease.

Design: Living rapid review of observational and experimental studies with random-effects hierarchical Bayesian meta-analyses. Published articles and pre-prints were identified via MEDLINE and medRxiv.

Setting: Community or hospital. No restrictions on location.

Participants: Adults who received a SARS-CoV-2 test or a COVID-19 diagnosis.

Measurements: Outcomes were SARS-CoV-2 infection, hospitalisation, disease severity and mortality stratified by smoking status. Study quality was assessed (i.e. 'good', 'fair' and 'poor').

Findings: v11 (searches up to 2021-02-16) included 405 studies with 62 'good' and 'fair' quality studies included in unadjusted meta-analyses. 121 studies (29.9%) reported current, former and never smoking status with the remainder using broader categories. Recorded smoking prevalence among people with COVID-19 was generally lower than national prevalence. Current compared with never smokers were at reduced risk of SARS-CoV-2 infection (RR = 0.71, 95% Credible Interval (CrI) = 0.61-0.82, $\tau = 0.34$). Data for former smokers were inconclusive (RR = 1.03, 95% CrI = 0.95-1.11, $\tau = 0.17$) but favoured there being no important association (4% probability of RR ≥ 1.1). Former compared with never smokers were at increased risk of hospitalisation (RR = 1.19, CrI = 1.1-1.29, $\tau = 0.13$), greater disease severity (RR = 1.8, CrI = 1.27-2.55, $\tau = 0.46$) and mortality (RR = 1.56, CrI = 1.23-2, $\tau = 0.43$). Data for current smokers on hospitalisation, disease severity and mortality were inconclusive (RR = 1.1, 95% CrI = 0.99-1.21, $\tau = 0.15$; RR 1.26,

95% CrI = 0.92-1.73, $\tau = 0.32$; RR = 1.12, 95% CrI = 0.84-1.47, $\tau = 0.42$, respectively) but favoured there being no important associations with hospitalisation and mortality (49% and 56% probability of RR ≥ 1.1 , respectively) and a small but important association with disease severity (83% probability of RR ≥ 1.1).

Conclusions: Compared with never smokers, current smokers appear to be at reduced risk of SARS-CoV-2 infection while former smokers appear to be at increased risk of hospitalisation, greater disease severity and mortality from COVID-19. However, it is uncertain whether these associations are causal.

v7 of this living review article has been published in *Addiction*

Introduction

COVID-19 is a respiratory disease caused by the SARS-CoV-2 virus. Large age and gender differences in case severity and mortality have been observed in the ongoing COVID-19 pandemic (Guan, Ni, et al., 2020); however, these differences are currently unexplained. SARS-CoV-2 enters epithelial cells through the angiotensin-converting enzyme 2 (ACE-2) receptor (Hoffmann et al., 2020). Some evidence suggests that gene expression and subsequent receptor levels are elevated in the airway and oral epithelium of current smokers (Brake et al., 2020; Cai, 2020), which could put smokers at higher risk of contracting SARS-CoV-2. Other studies, however, suggest that nicotine downregulates the ACE-2 receptor (Oakes et al., 2018). These uncertainties notwithstanding, both former and current smoking is known to increase the risk of respiratory viral (Abadom et al., 2016; Denholm et al., 2010) and bacterial (Almirall et al., 1999; Feldman and Anderson, 2013) infections and is associated with worse outcomes once infected. Cigarette smoke reduces the respiratory immune defence through peri-bronchiolar inflammation and fibrosis, impaired mucociliary clearance and disruption of the respiratory epithelium (Dye and Adler, 1994). There is also reason to believe that behavioural factors (e.g. regular hand-to-mouth movements) involved in smoking may increase SARS-CoV-2 infection and transmission in current smokers. However, early data from the COVID-19 pandemic have not provided clear evidence for a negative impact of current or former smoking on SARS-CoV-2 infection or COVID-19 disease outcomes, such as hospitalisation or mortality (Vardavas and Nikitara, 2020). It has also been hypothesised that nicotine might protect against a hyper-inflammatory response to SARS-CoV-2 infection, which may lead to adverse outcomes in patients with COVID-19 disease (Farsalinos, Niaura, et al., 2020).

There are several reviews that fall within the scope of smoking and COVID-19 (Alqahtani et al., 2020; Berlin et al., 2020; Emami et al., 2020; Farsalinos, Barbouni, et al., 2020; Grundy et al., 2020; Patanavanich and Glantz, 2020; Vardavas and Nikitara, 2020). We aimed to produce a rapid synthesis of available evidence pertaining to the rates of infection, hospitalisation, disease severity and mortality from SARS-CoV-2/COVID-19 stratified by smoking status. Given the increasing availability of data on this topic, this is a living review with regular updates. As evidence accumulates, the review will be expanded to include studies reporting COVID-19 outcomes by alternative nicotine use (e.g., nicotine replacement therapy or e-cigarettes).

Methods

Study design

This is a living evidence review which is updated as new evidence becomes available (Elliott et al., 2014). We adopted recommended best practice for rapid evidence reviews, which involved limiting the search to main databases and having one reviewer extract the data and another verify (Tricco et al., 2015). This study was not pre-registered but evolved from a report written for a UK medical society (Simons, Brown, et al., 2020). The most recent version of this living review is available [here](#). Version 7 of this living review has been published in a peer-reviewed journal (Simons, Shahab, et al., 2020). A completed Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) checklist is included in Supplementary file 1.

Eligibility criteria

Studies were included if they:

1. Were primary research studies using experimental (e.g. randomised controlled trial), quasi-experimental (e.g. pre- and post-test) or observational (e.g. case-control, retrospective cohort, prospective cohort) study designs;
2. Included adults aged 16+ years;
3. Recorded as outcome i) results of a SARS-CoV-2 diagnostic test (including antibody assays), ii) clinical diagnosis of COVID-19, iii) hospitalisation with COVID-19, iv) severity of COVID-19 disease in those hospitalised or v) mortality from COVID-19; Reported any of the outcomes of interest by self-reported or biochemically verified smoking status (e.g. current smoker, former smoker, never smoker) or current vaping and nicotine replacement therapy (NRT) use;
4. Were available in English;
5. Were published in a peer-reviewed journal, as a pre-print or a public health report by reputable bodies (e.g. governments, scientific societies).

Search strategy

The following terms were searched for in Ovid MEDLINE (2019-search date) as free text or Medical Subject Headings:

1. Tobacco Smoking/ or Smoking Cessation/ or Water Pipe Smoking/ or Smoking/ or Smoking Pipes/ or Cigar Smoking/ or Smoking Prevention/ or Cigarette Smoking/ or smoking.mp. or Pipe Smoking/ or Smoking, Non-Tobacco Products/ or Smoking Water Pipes/
2. Nicotine/ or nicotine.mp. or Electronic Nicotine Delivery Systems/ or Nicotine Chewing Gum/
3. vaping.mp. or Vaping/
4. 1 or 2 or 3

5. Coronavirus/ or Severe Acute Respiratory Syndrome/ or Coronavirus Infections/ or covid.mp.
6. 4 and 5

The following terms were searched for in titles, abstracts and full texts in medRxiv (no time limitations):

1. covid (this term captures both covid and SARS-CoV-2) AND smoking
2. covid AND nicotine
3. covid AND vaping

Additional articles/reports of interest were identified through mailing lists, Twitter, the International Severe Acute Respiratory and Emerging Infection Consortium (ISARIC) and the US Centers for Disease Control and Prevention (CDC). Where updated versions of pre-prints or public health reports were available, old versions were superseded.

Selection of studies

One reviewer screened titles, abstracts and full texts against the inclusion criteria.

Data extraction

Data were extracted by one reviewer and verified (i.e. independently checked against pre-prints and published reports) by another on i) author (year); ii) date published; iii) country; iv) study design; v) study setting; vi) sample size; vii) sex; viii) age; ix) smoking status (e.g. current, former, never, not stated, missing) and whether it was biochemically verified; x) use of alternative nicotine products; xi) SARS-CoV-2 testing; xii) SARS-CoV-2 infection; xiii) diagnosis of COVID-19; xiv) hospitalisation with COVID-19; xv) disease severity in those hospitalised with COVID-19; xvi) mortality; xvii) adjustment of smoking specific risk estimates for relevant covariates (e.g. age, sex); and xviii) whether a representative or random sampling method was used.

Quality appraisal

The quality of included studies was assessed to determine suitability for inclusion in meta-analyses. Studies were judged as 'good' quality if they: i) had <20% missing data on smoking status and used a reliable self-report measure that distinguished between current, former and never smoking status; AND ii) used biochemical verification of smoking status and reported results from adjusted analyses; OR reported data from a representative/random sample. Studies were rated as 'fair' if they fulfilled only criterion i) and were otherwise rated as 'poor'. The quality appraisal was conducted by one reviewer and verified by a second.

Evidence synthesis

A narrative synthesis was conducted. Data from 'good' and 'fair' quality studies were pooled in R v.3.6.3 (R Team, 2013). In a living review where new data are regularly added to the analyses, it may be more appropriate to use a Bayesian (as opposed to frequentist) approach where prior knowledge is used in combination with new data to estimate a posterior risk distribution. A Bayesian approach mitigates the issue of performing multiple statistical tests, which can inflate family-wise error. A series of random-effects hierarchical Bayesian meta-analyses were performed with the brms (Bürkner, 2018) package to estimate the relative risk for each comparison with accompanying 95% credible intervals (CrIs). We first defined prior distributions for the true pooled effect size (μ) and the between-study heterogeneity (τ), with μ specified as a normal distribution with a mean equal to the derived point estimate from each comparison of interest in the immediately preceding version of this living review, and τ specified as a half-Cauchy distribution with a mean of 0 and standard deviation of 1. The half-Cauchy distribution was selected to reflect prior knowledge that high levels of between-study heterogeneity are more likely than lower levels. Markov Chain Monte Carlo methods (20,000 burn-ins followed by 80,000 iterations) were then used to generate a risk distribution for each study, in addition to a pooled effect for the posterior risk distribution. We report forest plots with the pooled effect for the posterior risk distribution displayed as the median relative risk with an accompanying 95% CrIs. We used the empirical cumulative distribution function (ECDF) to estimate the probability of there being a 10% reduction or 10% increase in relative risk (RR) (i.e. $RR \geq 1.1$ or $RR \leq 0.9$). Due to a lack of indication as to what constitutes a clinically or epidemiologically meaningful effect (e.g. with regards to onward disease transmission or requirements for intensive care beds), we deemed a 10% change in risk as small but important. Where data were inconclusive (as indicated by CrIs crossing $RR = 1.0$), to disambiguate whether data favoured no effect or there being a small but important association, we estimated whether there was $\geq 75\%$ probability of $RR \geq 1.1$ or $RR \leq 0.9$.

Two sensitivity analyses were performed. First, a minimally informative prior for μ was specified as a normal distribution with a mean of 0 and standard deviation of 1 and τ as described above. Second, an informative prior as described above for μ was used with τ specified as a half-Cauchy distribution with a mean of 0.3 and standard deviation of 1 to reflect greater between-study heterogeneity.

To aid in the visualisation of smoking prevalence in the included studies, the weighted mean prevalence of current and former smoking was calculated for countries with ≥ 3 studies and plotted for comparison with national prevalence estimates. It should be noted that prevalence estimates in the included studies were not adjusted for age, sex, socioeconomic position, or geographic region within countries.

Results

In the current review version (v11) with searches up to 2021-02-16, a total of 1133 records were identified, with 405 studies included in a narrative synthesis and 62 studies included in meta-analyses (see Figure 1).




Figure 1. PRISMA flow diagram of included studies.

Characteristics of included studies are presented in Table 1. Studies were conducted across 41 countries. 109 studies were conducted in the USA, 70 in China, 44 in the UK, 28 in Spain, 20 in France, 18 in Mexico, 16 in Italy, 13 in Multiple, 8 in Turkey, 7 in Brazil and Iran, 5 in Israel and Switzerland, 4 in Finland and India, with 3 each from Australia, Austria, Japan, Saudi Arabia, and South Korea and a single study from 10 further countries. The majority of studies used observational designs (see Supplementary table S1). 256 (63.2%) were conducted in hospital settings, 106 studies (26.2%) included individuals from community and hospital settings, 40 studies (9.9%) were conducted exclusively in the community, with one study each conducted in a homeless shelter and a quarantine centre, and one study that did not state the study setting. Studies had a median of 502 (interquartile range = 146-2038) participants. The majority of studies (87.8%) used reverse transcriptase polymerase chain reaction (RT-PCR) for confirmation of SARS-CoV-2 infection, 12.2% used an antibody test to confirm prior infection and 6.8% of studies relied on a combination of RT-PCR or antibody assays.

Smoking status

Categorisation of smoking status was heterogeneous (see Table 1). 236 studies collected data on smoking status through routine electronic health records (EHRs), 129 studies used a bespoke case report form for COVID-19 and 40 studies did not state the source for information on smoking status. None of the studies verified smoking status biochemically. Notably,

only 121 (29.9%) studies reported current, former and never smoking status (see Supplementary table S2a), with a further 26 studies reporting only ever and never smoking status (see Supplementary table S2b). The remaining 252 studies reported current, current/former or current and former smoking status but did not explicitly state whether remaining participants were never smokers or if data were missing on smoking status (see Supplementary table S2c). 133 studies explicitly reported the proportion with missing data on smoking status, which ranged from 0% to 97.6%.

Use of alternative nicotine products

Eight studies recorded the use of alternative nicotine products in current and/or former smokers but did not report COVID-19 outcomes stratified by alternative nicotine use (Crooks et al., 2020; Ebinger et al., 2020; Gallichotte et al., 2020; Girardeau et al., 2020; Islam et al., 2020; Kantele et al., 2020; Miyara et al., 2020; Rimland et al., 2020). One additional study grouped together current smokers and vapers (Schubl et al., 2020).

Quality appraisal

Three studies were performed in random or representative population samples and were rated as 'good' quality, and 94 studies were rated as 'fair' quality, of which 62 studies reported results stratified by smoking status for the outcomes of interest and could be included in meta-analyses. The remaining 308 studies were rated as 'poor' quality (see Table 1).

Table 1. All studies included in narrative review and meta-analysis

ref	Lead author	Date published	Country	Sample size	Study setting	Median (IQR)	Female %	Current smoker %	Former smokers %	Current/former smokers %	Never smokers %
Huang et al. (2020)	Huang, Wang	2020-01-24	China	41	Hospital	49 (41-58)	27.0	7.3	-	-	-
Li et al. (2020)	Li	2020-02-12	China	17	Hospital	45 (33-57)	47.1	17.6	-	-	-
Zhang et al. (2020)	Zhang, Dong	2020-02-19	China	140	Hospital	57 [^] (25-87)	46.3	1.4	5.00	-	-
Yang et al. (2020)	Yang, Yu	2020-02-24	China	52	Hospital	60 (47-73)	37.0	3.8	-	-	-
Guan et al. (2020)	Guan, Ni	2020-02-28	China	1,099	Hospital	47 (35-58)	41.9	12.5	1.91	-	84.
Liu et al. (2020)	Liu, Tao	2020-02-28	China	78	Hospital	38 (33-57)	50.0	-	-	6.41	-
Qi et al. (2020)	Qi	2020-03-03	China	267	Hospital	48 (35-65)	45.2	19.9	-	-	-
Huang et al. (2020)	Huang, Yang	2020-03-05	China	36	Hospital	69 (60-78)	30.6	-	-	11.11	-
Xu et al. (2020)	Xu	2020-03-08	China	53	Hospital	NA	47.2	11.3	-	-	-
Zhou et al. (2020)	Zhou, Yu	2020-03-11	China	191	Hospital	56 (46-67)	38.0	5.8	-	-	-

Author(s)	First Name	Date	Country	Sample Size (n)	Setting	Age (mean)	IC ₅₀ (days)	IC ₉₀ (days)	IC ₉₅ (days)	IC ₉₉ (days)	IC _{99.5} (days)
Liu et al. (2020)	Liu, Ming	2020-03-12	China	41	Hospital	39 (30-48)	58.5	9.8	-	-	-
Mo et al. (n.d.)	Mo	2020-03-16	China	155	Hospital	54 (53-66)	44.5	3.9	-	-	-
Shi et al. (2020)	Shi, Yu	2020-03-18	China	487	Hospital	46 (27-65)	46.8	-	-	8.21	-
Zhang et al. (2020)	Zhang, Cai	2020-03-20	China	645	Hospital	NA	49.1	6.4	-	-	-
Dong et al. (2020)	Dong, Cao	2020-03-20	China	9	Hospital	44 (30-46)	66.7	11.1	-	-	-
Wan et al. (2020)	Wan	2020-03-21	China	135	Hospital	47 (36-55)	46.7	6.7	-	-	-
Jin et al. (2020)	Jin	2020-03-24	China	651	Hospital	46 (32-60)	49.2	6.3	-	-	-
Wang et al. (2020)	Wang, Pan	2020-03-24	China	125	Hospital	41 (26-66)	43.2	-	-	12.80	-
Lian et al. (2020)	Lian	2020-03-25	China	788	Hospital	NA	38.5	6.9	-	-	-
Hu et al. (2020)	Hu	2020-03-25	China	323	Hospital	61 [^] (23-91)	48.6	-	-	11.76	-
Guan et al. (2020)	Guan, Liang	2020-03-26	China	1,590	Hospital	49 (33-64)	42.7	-	-	6.98	93.
Chen et al. (2020)	Chen	2020-03-26	China	548	Hospital	62 (44-70)	37.6	4.4	2.55	-	-
Guo et al. (2020)	Guo	2020-03-27	China	187	Hospital	59 (45-73)	51.3	9.6	-	-	-
CDCMMWR (2020)	Chow (US CDC)	2020-03-31	USA	7,162	Community and Hospital	NA	-	1.3	2.30	-	-
Kim et al. (2020)	Kim	2020-04-01	South Korea	28	Hospital	43 (30-56)	46.4	17.9	-	-	-
Feng et al. (2020)	Feng	2020-04-10	China	476	Hospital	53 (40-64)	43.1	9.2	-	-	-
Rentsch et al. (2020)	Rentsch	2020-04-14	USA	3,528	Community and Hospital	66 (60-70)	4.6	27.2	30.60	-	36.
Goyal et al. (2020)	Goyal	2020-04-17	USA	393	Hospital	62.2 (49-74)	39.3	5.1	-	-	-
Zheng et al. (2020)	Zheng, Gao	2020-04-19	China	66	Hospital	47 [^] (NA)	25.8	12.1	-	-	-
Gold et al. (2020)	Gold (US CDC)	2020-04-20	USA	305	Hospital	NA	50.5	5.2	-	-	-
Argenziano et al. (2020)	Argenziano	2020-04-22	USA	1,000	Hospital	63 (50-75)	40.4	4.9	17.90	-	77.
Richardson et al. (2020)	Richardson	2020-04-22	USA	5,700	Hospital	63 (52-75)	39.7	-	-	9.79	52.
Fontanet et al. (2020)	Fontanet	2020-04-23	France	661	Community and Hospital	37 (16-47)	62.0	10.4	-	-	-
Shi et al. (2020)	Shi, Ren	2020-04-23	China	134	Hospital	46 (34-58)	51.5	-	-	10.45	-

Hadjadj et al. (2020)	Hadjadj	2020-04-23	France	50	Hospital	55 (50-63)	22.0	2.0	18.00	-	80.
Liao et al. (2020)	Liao, Feng	2020-04-24	China	1,848	Hospital	55 (48-61)	54.7	-	-	0.43	-
Gil-Agudo et al. (2020)	Gil-Agudo	2020-04-24	Spain	7	Hospital	68 (34-75)	28.6	-	-	42.86	57.
Yao et al. (2020)	Yao	2020-04-24	China	108	Hospital	52 (37-58)	60.2	3.7	-	-	-
Zuo et al. (2020)	Zuo, Yalavarthi	2020-04-24	USA	50	Hospital	61 (46-76)	34.0	-	-	36.00	-
Solis and Carreño (2020)	Solis	2020-04-25	Mexico	650	Hospital	46 (NA)	42.1	9.4	-	-	-
Yu et al. (2020)	Yu, Cai	2020-04-27	China	95	Hospital	NA	44.2	8.4	-	-	-
Ziehr et al. (2020)	Ziehr	2020-04-29	USA	66	Hospital	58 (23-87)	35.0	-	-	33.33	63.
Zheng et al. (2020)	Zheng, Xiong	2020-04-30	China	73	Hospital	43 [^] (NA)	45.2	-	-	10.96	89.
Kalan et al. (2020)	Kalan	2020-05-01	Iran	193	Hospital	52.6 [^] (37-67)	36.3	7.3	-	-	85.
Kolin et al. (2020)	Kolin	2020-05-05	UK	1,474	Community and Hospital	58 (49-67)	46.6	14.5	40.16	-	44.
Borobia et al. (2020)	Borobia	2020-05-06	Spain	2,226	Hospital	61 (46-78)	52.0	7.1	-	-	-
Giacomelli et al. (2020)	Giacomelli	2020-05-06	Italy	233	Hospital	61 (50-72)	31.9	-	-	30.04	70.
Shah et al. (2020)	Shah	2020-05-06	USA	316	Hospital	63 (43-72)	48.1	16.5	17.72	-	42.
Williamson et al. (2020)	The OpenSAFELY Collaborative	2020-05-07	UK	17,425,445	Community and Hospital	NA	50.1	17.0	32.93	-	45.
Allenbach et al. (2020)	Allenbach	2020-05-08	France	152	Hospital	77 (60-83)	31.1	-	-	6.58	-
Robilotti et al. (2020)	Robilotti	2020-05-08	USA	423	Hospital	NA	50.0	2.1	37.59	-	58.
Lubetzky et al. (2020)	Lubetzky	2020-05-08	USA	54	Hospital	57 (29-83)	62.0	-	-	22.22	-
Yin et al. (2020)	Yin, Yang	2020-05-10	China	106	Hospital	73 (61-85)	39.6	-	-	16.98	-
Rica et al. (2020)	de la Rica	2020-05-11	Spain	48	Hospital	66 [^] (33-88)	33.0	-	-	20.83	-
Cho et al. (2020)	Cho	2020-05-11	UK	1,331	Community and Hospital	NA	49.2	19.0	26.97	-	54.
Yanover et al. (2020)	Yanover	2020-05-13	Israel	4,353	Community and Hospital	35 (22-54)	44.5	11.8	2.96	-	85.
Hamer et al. (2020)	Hamer	2020-05-13	UK	387,109	Hospital	56.2 (48-64)	55.1	9.7	34.84	-	55.
Targher et al.	Targher	2020-05-	China	339	Hospital	48.4 [^]	52.8	8.3	-	-	-

(2020)	Author	Date	Country	Case Count	Setting	Age Range (NA)	ICU	ICU	ICU	ICU	ICU
Carrillo-Vega et al. (2020)	Carrillo-Vega	2020-05-14	Mexico	10,544	Community and Hospital	46.5^ (30-62)	42.3	8.9	-	-	-
Regina et al. (2020)	Regina	2020-05-14	Switzerland	200	Hospital	70 (55-81)	40.0	4.5	-	-	-
Almazeedi et al. (2020)	Almazeedi	2020-05-15	Kuwait	1,096	Hospital	41 (25-57)	19.0	4.0	-	-	-
Lusignan et al. (2020)	de Lusignan	2020-05-15	UK	3,802	Community	58 (34-73)	57.6	10.9	46.11	-	29.
Palaiodimos et al. (2020)	Palaiodimos	2020-05-15	USA	200	Hospital	64 (50-73.5)	51.0	-	-	32.50	67.
Mejia-Vilet et al. (2020)	Mejia-Vilet	2020-05-16	Mexico	329	Hospital	49 (41-60)	36.0	-	-	6.99	-
Chen et al. (2020)	Chen, Jiang	2020-05-16	China	135	Hospital	NA	42.2	-	-	9.63	-
Li et al. (2020)	Li, Chen	2020-05-16	China	1,008	Hospital	55 (44-65)	43.6	5.7	-	-	-
Valenti et al. (2020)	Valenti	2020-05-18	Italy	789	Community	40.7^ (NA)	35.0	25.9	-	-	-
Feuth et al. (2020)	Feuth	2020-05-18	Finland	28	Hospital	56 (47-72)	46.0	10.7	28.57	-	60.
Ge et al. (2020)	Ge	2020-05-18	China	51	Hospital	70 (58-79)	27.5	13.7	-	-	-
Parrotta et al. (2020)	Parrotta	2020-05-18	USA	76	Community and Hospital	44.9 (13-71)	61.8	2.6	26.32	-	68.
Shekhar et al. (2020)	Shekhar	2020-05-18	USA	50	Hospital	55.5 (20-85)	54.0	48.0	-	-	-
Rimland et al. (2020)	Rimland	2020-05-19	USA	11	Hospital	59 (48-65)	18.2	9.1	-	-	-
Basse et al. (2020)	Basse	2020-05-19	France	141	Hospital	62 (52-72)	72.0	17.7	-	-	-
Freites et al. (2020)	Freites	2020-05-19	Spain	123	Hospital	59.88^ (44-74)	69.9	3.3	-	-	-
Alshami et al. (2020)	Alshami	2020-05-19	Saudi Arabia	128	Quarantine Centre	39.6^ (24-55)	53.9	15.6	2.34	-	-
Shi et al. (2020)	Shi, Zhao	2020-05-20	China	101	Hospital	71 (59-80)	40.6	-	-	4.95	-
Al-Hindawi et al. (2020)	Al-Hindawi	2020-05-20	UK	31	Hospital	61 (NA)	12.9	3.2	70.97	-	25.
Wu et al. (2020)	Wu	2020-05-21	Italy	174	Hospital	61.2^ (50-71)	30.5	-	-	33.33	-
Kim et al. (2020)	Kim, Garg	2020-05-22	USA	2,491	Hospital	62 (50-75)	46.8	6.0	25.77	-	-
Docherty et al. (2020)	Docherty	2020-05-22	Multiple	20,133	Hospital	72.9 (58-82)	40.0	4.2	21.68	-	44.
Petrilli et al. (2020)	Petrilli	2020-05-22	USA	5,279	Community and Hospital	54 (38-66)	51.5	5.5	17.09	-	61.
Klang et al. (n.d.)	Klang	2020-05-22	USA	3,406	Hospital	NA	61.8	-	-	23.28	-

Author(s)	Study Name	Date	Country	Sample Size	Setting	Age Range	Prevalence (%)	OR	95% CI	OR	95% CI
Vaquero et al. (2020)	Vaquero-Roncero	2020-05-24	Spain	146	Hospital	66 [^] (59-72)	32.2	-	-	6.85	-
Ip et al. (2020)	Ip	2020-05-25	USA	2,512	Hospital	64 (52-76)	37.6	3.1	17.83	-	64.
Heili-Frades (2020)	Heili-Frades	2020-05-25	Spain	4,712	Hospital	62 (47-77)	50.5	4.9	17.40	-	-
Berumen et al. (2020)	Berumen	2020-05-26	Mexico	102,875	Hospital	NA	49.1	-	-	9.64	-
Garibaldi et al. (2020)	Garibaldi	2020-05-26	USA	832	Hospital	63 (49-75)	47.0	5.5	22.60	-	-
Soto-Mota et al. (2020)	Soto-Mota	2020-05-27	Mexico	400	Hospital	NA	30.0	-	-	12.00	-
Li et al. (2020)	Li, Long	2020-05-28	China	145	Not Stated	49 [^] (13-80)	61.0	-	-	5.52	-
Louis et al. (2020)	Louis	2020-05-28	USA	22	Hospital	66.5 [^] (55-77)	36.4	-	-	45.45	-
Kuderer et al. (2020)	Kuderer	2020-05-28	Multiple	928	Community and Hospital	66 (57-76)	50.0	4.6	35.13	-	50.
Gianfrancesco et al. (2020)	Gianfrancesco	2020-05-29	Multiple	600	Community and Hospital	56 (45-67)	71.0	-	-	21.50	64.
Chaudhry et al. (2020)	Chaudhry	2020-05-29	USA	40	Community and Hospital	52 (45.5-61)	60.0	-	-	15.00	-
Niedzwiedz et al. (2020)	Niedzwiedz	2020-05-29	UK	392,116	Community and Hospital	NA	54.9	9.8	34.81	-	55.
Valle et al. (2020)	del Valle	2020-05-30	USA	1,484	Hospital	62 (52-72)	40.6	5.5	23.32	-	-
Bello-Chavolla et al. (2020)	Bello-Chavolla	2020-05-31	Mexico	177,133	Community and Hospital	42.6 (26-59)	48.9	-	-	9.28	-
Batty et al. (2020)	Batty	2020-06-01	UK	908	Hospital	57.27 [^] (48-66)	44.3	11.2	-	-	-
Israel et al. (2020)	Israel	2020-06-01	Israel	24,906	Community and Hospital	40 (27-59)	48.7	16.8	12.66	-	70.
Hao et al. (2020)	Hao	2020-06-01	China	788	Hospital	46 (35-56)	48.4	6.9	-	-	-
Lassale et al. (2020)	Lassale	2020-06-01	UK	900	Hospital	57.2 [^] (48-66)	44.4	11.4	41.89	-	46.
Eugen-Olsen et al. (2020)	Eugen-Olsen	2020-06-02	Denmark	407	Hospital	64 (47-77)	57.7	20.6	36.86	-	39.
Martinez-Portilla et al. (2020)	Martinez-Portilla	2020-06-02	Mexico	224	Community and Hospital	29 (26-33)	100.0	-	-	3.12	-
Raisi-Estabragh et al. (2020)	Raisi-Estabragh	2020-06-02	UK	4,510	Hospital	NA	48.8	-	-	51.80	-
Luo et al. (2020)	Luo	2020-06-02	China	625	Hospital	46 (NA)	47.7	3.0	-	-	-
Paulun et al.		2020-06-02				46 (22-70)					

Boulware et al. (2020)	Boulware	2020-06-03	Multiple	821	Community	40 (33-50)	51.6	3.3	-	-	-
Ikitimur et al. (2020)	Ikitimur	2020-06-03	Turkey	81	Hospital	55^ (38-72)	44.0	-	-	28.40	-
Sierpiński et al. (2020)	Sierpinski	2020-06-03	Poland	1,942	Community	50 (NA)	60.0	6.3	-	-	-
Wang et al. (2020)	Wang, Oekelen	2020-06-05	USA	58	Community and Hospital	67 (NA)	48.0	-	-	36.21	-
Perrone et al. (2020)	Perrone	2020-06-05	Italy	1,189	Hospital	NA	21.2	-	-	21.87	-
Sharma et al. (2020)	Sharma	2020-06-05	India	501	Hospital	35.1^ (18-51)	36.0	-	-	4.19	-
Magagnoli et al. (2020)	Magagnoli	2020-06-05	USA	807	Hospital	70 (60-75)	4.3	-	-	15.86	-
Ramlall et al. (2020)	Ramlall	2020-06-06	USA	11,116	Community and Hospital	52 (34.7-69.5)	55.2	-	-	26.80	73.
Giannouchos et al. (2020)	Giannouchos	2020-06-07	Mexico	236,439	Community and Hospital	42.5^ (25-59)	49.1	9.1	-	-	-
Romão et al. (2020)	Romao	2020-06-08	Portugal	34	Community	41^ (26-66)	67.7	-	-	26.47	-
Cen et al. (2020)	Cen	2020-06-08	China	1,007	Hospital	61 (49-68)	51.0	-	-	8.74	-
Houlihan et al. (2020)	Houlihan	2020-06-09	UK	200	Community	34 (29-44)	61.0	11.0	16.50	-	66.
Lan et al. (2020)	Lan	2020-06-09	USA	104	Community	49^ (34-63)	47.1	-	-	24.04	-
Russell et al. (2020)	Russell	2020-06-09	UK	156	Community and Hospital	65.18^ (50-79)	42.3	7.1	25.00	-	37.
Veras et al. (2020)	Veras	2020-06-09	Brazil	32	Hospital	58.9^ (40-77)	47.0	-	-	25.00	-
Rossi et al. (2020)	Rossi	2020-06-09	France	246	Hospital	68^ (53-83)	39.0	-	-	25.20	-
Martin-Jimenez et al. (2020)	Martin-Jiminez	2020-06-09	Spain	339	Hospital	81.6 (72-87)	39.5	-	-	30.68	-
Rajter et al. (2020)	Rajter	2020-06-10	USA	280	Hospital	59.6^ (41-77)	45.5	5.7	10.71	-	74.
Zhou et al. (2020)	Zhou, He	2020-06-10	China	238	Hospital	55.5 (35-67)	57.0	2.9	-	-	-
Woolford et al. (2020)	Woolford	2020-06-11	UK	4,510	Community and Hospital	70.5 (NA)	51.2	13.0	38.12	-	48.
Hultcrantz et al. (2020)	Hultcrantz	2020-06-11	USA	127	Community and Hospital	68 (41-91)	46.0	-	-	26.77	72.
Hernández-Garduño (2020)	Hernandez, Garduno	2020-06-11	Mexico	32,583	Community and Hospital	45 (34-56)	48.7	-	-	11.02	-
Sterlin et al. (2020)	Sterlin	2020-06-11	France	135	Hospital	61 (50-72)	41.0	3.7	38.52	-	57.

Maraschini et al. (2020)	Maraschini	2020-06-12	Italy	146	Hospital	32.5 [^] (27-38)	100.0	-	9.59	-	80.
Wang et al. (2020)	Wang, Zhong	2020-06-12	USA	7,592	Community and Hospital	NA	45.1	3.6	17.08	-	51.
McQueenie et al. (2020)	McQueenie	2020-06-12	UK	428,199	Community and Hospital	NA	54.9	-	-	44.36	55.
Miyara et al. (2020)	Miyara	2020-06-12	France	479	Community and Hospital	NA	44.7	6.7	31.60	-	59.
Apea et al. (2021)	Apea	2020-06-12	UK	1,737	Hospital	63.4 [^] (NA)	30.4	-	-	9.96	-
Garassino et al. (2020)	Garassino	2020-06-12	Multiple	200	Community and Hospital	68 (61.8-75)	30.0	24.0	55.50	-	18.
Zeng et al. (2020)	Zeng	2020-06-16	China	1,031	Hospital	60.3 [^] (46-74)	47.8	-	-	10.18	-
Suleyman et al. (2020)	Suleyman	2020-06-16	USA	463	Hospital	57.5 [^] (40-74)	55.9	-	-	34.56	-
Chen et al. (2020)	Chen, Yu	2020-06-16	China	1,859	Hospital	59 (45-68)	50.0	2.4	3.55	-	94.
Kibler et al. (2020)	Kibler	2020-06-16	France	702	Community and Hospital	82 [^] (75-88)	56.0	3.7	-	-	-
Olivares et al. (2020)	Olivares	2020-06-16	Chile	21	Hospital	61 [^] (26-85)	76.2	-	-	9.52	-
Elezkurtaj et al. (2020)	Elezkurtaj	2020-06-17	Germany	26	Hospital	70 (61.8-78.3)	34.6	-	-	19.23	-
Zuo et al. (2020)	Zuo, Estes	2020-06-17	China	172	Hospital	61 [^] (25-95)	44.0	-	-	26.16	-
Killerby (2020)	Killerby	2020-06-17	USA	531	Community and Hospital	51.6 (38-62)	57.1	-	-	17.14	71.
Gu et al. (2020)	Gu	2020-06-18	USA	5,698	Community and Hospital	47 [^] (26-67)	62.0	7.0	24.68	-	50.
Wei et al. (2020)	Wei	2020-06-18	USA	147	Hospital	52 [^] (34-70)	41.0	14.3	-	-	-
Crovetto et al. (2020)	Crovetto	2020-06-19	Spain	874	Community and Hospital	33.7 [^] (28-38)	100.0	1.1	-	-	-
Govind et al. (2020)	Govind	2020-06-20	UK	6,309	Community and Hospital	46.5 [^] (31-61)	38.3	66.3	26.77	-	5.5
Sisó-Almirall et al. (2020)	Siso-Almirall	2020-06-20	Spain	322	Community and Hospital	56.7 [^] (38-74)	50.0	-	-	25.16	-
Salton et al. (2020)	Salton	2020-06-20	Italy	173	Hospital	64.4 [^] (NA)	34.9	-	-	29.48	-
Duan et al. (2020)	Duan	2020-06-22	China	616	Hospital	64 (53-70)	57.5	3.7	-	-	-

Lenka et al. (2020)	Lenka	2020-06-22	USA	32	Hospital	62.2 [^] (51-73)	37.5	-	-	50.00	-
Fisman et al. (2020)	Fisman	2020-06-23	Canada	21,922	Community and Hospital	NA	57.0	-	-	2.35	-
Madariaga et al. (2020)	Madariaga	2020-06-23	USA	103	Community and Hospital	41.8 [^] (27-55)	48.5	-	-	25.24	74.
Jin et al. (2020)	Jin, Gu	2020-06-25	China	6	Hospital	60.5 [^] (51-75)	33.3	33.3	-	-	-
Mendy et al. (2020)	Mendy	2020-06-27	USA	689	Community and Hospital	49.5 (35.2-67.5)	47.0	-	-	24.67	-
Sigel et al. (2020)	Sigel	2020-06-28	USA	493	Hospital	60 (55-67)	24.1	-	-	28.60	-
Souza et al. (2020)	de Souza	2020-06-28	Brazil	8,443	Hospital	NA	53.0	-	-	1.68	-
Nguyen et al. (2020)	Nguyen	2020-06-29	USA	689	Community and Hospital	55 (40-68)	57.0	-	-	24.82	-
Melo et al. (2020)	de Melo	2020-06-29	Brazil	181	Hospital	55.3 [^] (34-76)	60.8	9.9	12.15	-	38.
Auvinen et al. (2020)	Auvinen	2020-06-29	Finland	61	Hospital	53 (41-67)	36.0	18.0	27.87	-	54.
Magleby et al. (2020)	Magleby	2020-06-30	USA	678	Hospital	68 (50-81)	38.9	-	-	28.61	-
Hewitt et al. (2020)	Hewitt	2020-06-30	Multiple	1,564	Hospital	74 (61-83)	42.3	7.7	38.55	-	52.
Mohamud et al. (2020)	Mohamud	2020-07-02	USA	6	Hospital	65.8 [^] (55-78)	16.7	-	-	16.67	-
Trubiano et al. (2020)	Trubiano	2020-07-02	Australia	2,935	Community and Hospital	39 (29-53)	63.5	-	-	8.82	-
Patel et al. (2020)	Patel	2020-07-03	USA	129	Hospital	60.8 [^] (47-74)	45.0	37.2	-	-	-
Merzon et al. (2020)	Merzon	2020-07-03	Israel	7,807	Community and Hospital	46.2 [^] (NA)	58.6	-	-	16.18	-
Bello-Chavolla et al. (2020)	Bello-Chavolla, Antonio-Villa	2020-07-04	Mexico	60,121	Community and Hospital	45.5 [^] (29-61)	47.0	-	-	10.48	-
Zacharioudakis et al. (2020)	Zacharioudakis	2020-07-04	USA	314	Hospital	64 (54-72)	34.7	-	-	22.78	-
Antonio-Villa et al. (2020)	Antonio-Villa	2020-07-04	Mexico	34,263	Community and Hospital	40 [^] (29-50)	62.9	9.7	-	-	-
Kimmig et al. (2020)	Kimmig	2020-07-06	USA	111	Hospital	63 [^] (48-78)	44.1	7.2	36.04	-	56.
Senkal (2020)	Senkal	2020-07-07	Turkey	611	Hospital	57 [^] (18-98)	40.6	11.3	-	-	-
Xie et al. (2020)	Xie	2020-07-07	China	619	Hospital	NA	52.0	-	-	8.24	-

Elmunzer et al. (2020)	Elmunzer	2020-07-09	Multiple	1,992	Hospital	60 [^] (43-76)	43.0	6.3	28.56	-	59.
Alizadehsani et al. (2020)	Alizadehsani	2020-07-09	Iran	319	Hospital	45.48 [^] (26-63)	55.5	-	-	0.31	-
Maucourant et al. (2020)	Maucourant	2020-07-10	Sweden	27	Hospital	57 (18-78)	22.2	11.1	25.93	-	40.
Fan et al. (2020)	Fan	2020-07-11	UK	1,425	Community and Hospital	NA	46.7	12.2	40.07	-	46.
Shi et al. (2020)	Shi, Resurreccion	2020-07-11	UK	1,521	Community and Hospital	61.5 [^] (57-66.8)	45.9	-	-	54.90	-
REACT Study Investigators et al. (2020)	Riley	2020-07-11	UK	120,620	Community and Hospital	NA	54.0	2.2	-	-	-
Hippisley-Cox et al. (2020)	Hippisley-Cox	2020-07-13	UK	8,275,949	Community and Hospital	48.5 [^] (30-66)	50.3	17.2	21.44	-	57.
Zhang et al. (2020)	Zhang, Cao	2020-07-14	China	289	Hospital	57 (22-88)	46.6	3.5	6.23	-	-
Eiros et al. (2020)	Eiros	2020-07-14	Spain	139	Community and Hospital	52 (41-57)	72.0	4.3	50.36	-	-
Marcos et al. (2020)	Marcos	2020-07-14	Spain	918	Hospital	72.8 [^] (58-87)	42.2	6.1	-	15.25	-
Hoertel et al. (2020)	Hoertel, Sanchez, Rico	2020-07-14	France	7,345	Hospital	NA	49.3	8.5	-	-	-
Shi et al. (2020)	Shi, Zuo	2020-07-15	USA	172	Hospital	61.48 [^] (25-96)	44.0	-	-	26.16	-
Hussein et al. (2020)	Hussein	2020-07-15	USA	502	Hospital	60.9 [^] (45-76)	52.0	9.0	22.11	-	-
Bian et al. (2020)	Bian	2020-07-15	China	28	Hospital	56 [^] (42-67)	42.9	7.1	-	-	-
Zhan et al. (2020)	Zhan	2020-07-16	China	75	Hospital	57 (25-75)	48.0	-	-	12.00	-
Omrani et al. (2020)	Omrani	2020-07-16	Qatar	1,409	Community and Hospital	39 (30-50)	17.2	-	-	9.23	-
Gupta et al. (2020)	Gupta	2020-07-16	USA	496	Hospital	70 (60-78)	46.0	-	-	7.26	-
Soares et al. (2020)	Soares	2020-07-16	Brazil	10,713	Community and Hospital	NA	55.0	2.0	-	-	-
Abolghasemi et al. (2020)	Abolghasemi	2020-07-17	Iran	24	Hospital	49 [^] (29-64)	37.5	-	-	4.17	-
Merkely et al. (2020)	Merkely	2020-07-17	Hungary	10,474	Community	48.7 [^] (30-66)	53.6	28.0	20.46	-	51.
Fox et al. (2020)	Fox	2020-07-17	UK	55	Community and Hospital	63 (23-88)	31.0	1.8	10.91	-	56.
Pandolfi et al. (2020)	Pandolfi	2020-07-17	Italy	33	Hospital	62 (52-65)	21.1	3.0	24.24	-	72.
Girardeau et al.		2020-07-				30 (29-					

Girardeau et al. (2020)	Girardeau	2020-07-17	France	10	Community	50.0 (33)	50.0	40.0	10.00	-	-
Kurashima et al. (2020)	Kurashima	2020-07-17	Japan	53	Hospital	62.9 [^] (49-76)	35.8	-	-	50.94	-
McGrail and Edwards (2020)	Edwards	2020-07-19	USA	209	Hospital	62.5 (NA)	38.8	-	-	18.66	-
Martinez-Resendez et al. (2020)	Martinez-Resendez	2020-07-20	Mexico	8	Hospital	57 (48-69)	25.0	-	-	12.50	-
Hoertel et al. (2020)	Hoertel	2020-07-20	France	12,612	Hospital	58.7 [^] (39-77)	49.6	-	-	9.28	-
Wang et al. (2020)	Wang, Shu	2020-07-20	China	59	Hospital	67.4 [^] (56-78)	35.6	-	-	15.25	-
Bernaola et al. (2020)	Bernaola	2020-07-21	Spain	1,645	Hospital	NA	38.5	2.5	10.88	-	86.
Schneeweiss et al. (2020)	Schneeweiss	2020-07-22	USA	24,313	Community and Hospital	67 [^] (53-80)	53.0	-	-	2.88	-
Concha-Mejia and Rincon-Sanchez (2020)	Mejia	2020-07-24	Colombia	72	Community and Hospital	46 (28-64)	47.0	8.3	11.11	-	-
Izquierdo et al. (2020)	Izquierdo	2020-07-24	Spain	71,192	Community and Hospital	42 [^] (18-66)	59.0	10.0	-	-	-
Santos et al. (2020)	Santos	2020-07-25	USA	23	Community and Hospital	NA	-	-	-	8.70	-
Reiter et al. (2020)	Reiter	2020-07-26	Austria	235	Community	44.2 [^] (32-55)	70.0	22.6	22.55	-	54.
Motta et al. (2020)	Motta	2020-07-26	USA	374	Hospital	64.7 [^] (46-82)	41.4	-	-	33.16	66.
Altamimi et al. (2020)	Altamimi	2020-07-27	Qatar	68	Hospital	49 [^] (40-58)	2.0	16.2	-	-	-
Thompson et al. (2020)	Thompson	2020-07-27	UK	470	Hospital	71 (57-82)	46.0	14.0	27.23	-	58.
Zobairy et al. (2020)	Zobairy	2020-07-28	Iran	203	Community and Hospital	49.2 [^] (32-65)	44.8	5.9	-	-	-
Zhou et al. (2020)	Zhou, Sun	2020-07-29	China	144	Hospital	47 (38-56)	46.5	9.0	-	-	-
Kumar et al. (2020)	Kumar	2020-07-29	India	91	Hospital	47 [^] (41-52)	21.0	44.0	-	-	-
Qu et al. (2020)	Qu	2020-07-29	China	246	Hospital	53.6 [^] (38-68)	53.3	42.3	-	-	-
Higuchi et al. (2020)	Higuchi	2020-07-30	Japan	57	Hospital	52 (35-70)	43.9	12.3	29.82	-	57.
Zhao et al. (2020)	Zhao, Chen	2020-07-30	USA	641	Hospital	60 (NA)	40.1	21.7	-	-	-
Fond et al. (2020)	Fond	2020-07-30	France	1,092	Hospital	62.5 (51-76)	45.7	11.4	-	-	-
Jun et al. (2020)	Jun	2020-08-01	USA	3,086	Hospital	66 (56-77)	40.9	3.7	21.32	-	52.
Morshed et al.		2020-08-				37 (31-					

Morshed et al. (2020)	Morshed	2020-08-02	Bangladesh	103	Community	57.1^ (53)	28.2	31.1	-	-	-
Iversen et al. (2020)	Iversen	2020-08-03	Denmark	28,792	Community and Hospital	44.4^ (31-57)	78.9	16.0	6.52	-	76.
Ebinger et al. (2020)	Ebinger	2020-08-04	USA	6,062	Community	41.5^ (29-53)	67.8	1.7	-	-	-
Hadi et al. (2020)	Hadi	2020-08-05	USA	370	Community and Hospital	48.2^ (34-62)	29.5	-	-	15.14	84.
Tao et al. (2020)	Tao	2020-08-06	China	70	Community	33.2^ (12-53)	48.6	-	-	15.71	-
Klang et al. (2020)	Klang, Soffer	2020-08-09	USA	1,320	Hospital	NA	41.5	-	-	24.70	-
Zhou et al. (2021)	Zhou, Ma	2020-08-10	China	429	Hospital	58.3^ (42-74)	50.6	-	-	8.39	-
Altibi et al. (2020)	Altibi	2020-08-11	USA	706	Hospital	66.7^ (51-81)	43.0	4.0	37.25	-	58.
Izzi-Engbeaya et al. (2020)	Izzi-Engbeaya	2020-08-11	UK	889	Hospital	65.8^ (48-83)	40.0	-	-	21.26	33.
Rizzo et al. (2020)	Rizzo	2020-08-11	USA	76,819	Hospital	54 (38-67)	55.2	6.7	20.78	-	50.
Jehi et al. (2020)	Jehi	2020-08-11	USA	4,536	Community and Hospital	NA	-	7.3	28.48	-	49.
Holman et al. (2020)	Holman	2020-08-13	UK	10,989	Community and Hospital	NA	38.8	5.5	42.62	-	49.
Ouyang et al. (2020)	Ouyang	2020-08-14	China	217	Hospital	46.5^ (30-62)	53.5	16.6	-	-	-
Valenzuela et al. (2020)	Valenzuela	2020-08-14	Chile	29	Hospital	56.9^ (43-70)	6.9	17.2	-	-	-
Monteiro et al. (2020)	Monteiro	2020-08-14	USA	112	Hospital	61 (45-74)	34.0	6.2	17.86	-	68.
Philipose et al. (2020)	Philipose	2020-08-14	UK	466	Hospital	67 (6-97)	41.8	6.0	73.18	-	16.
Weerahandi et al. (2020)	Weerahandi	2020-08-14	USA	394	Community	63 (55-70)	37.0	5.3	25.89	-	55.
Parra-Bracamonte et al. (2020)	Parra-Bracamonte	2020-08-14	Mexico	331,298	Community and Hospital	44 (33-56)	46.2	-	-	7.39	-
Peters et al. (2020)	Peters	2020-08-15	Netherlands	1,893	Hospital	66.8^ (52-81)	39.4	4.9	-	-	-
Islam et al. (2020)	Islam	2020-08-18	Bangladesh	1,016	Community and Hospital	37 (28-49)	35.9	18.2	-	-	-
Chand et al. (2020)	Chand	2020-08-19	USA	300	Hospital	58.2^ (45-70)	39.3	22.3	-	-	-
Aksu et al. (2020)	Aksu	2020-08-19	Turkey	123	Community and Hospital	49.7^ (36-63)	33.3	11.4	-	-	-
Alkurt et al. (2020)	Alkurt	2020-08-20	Turkey	932	Community and	34.8^ (25-44)	64.4	24.5	-	-	-

Author(s)	Study Name	Date	Country	Sample Size	Setting	ICU Mortality (%)	Case Fatality Rate (%)	Case Fatality Rate (95% CI)	Case Fatality Rate (95% CI)	Case Fatality Rate (95% CI)	Case Fatality Rate (95% CI)
Ward et al. (2020)	Ward	2020-08-21	UK	99,908	Community	NA	56.1	10.6	-	-	-
Salerno et al. (2020)	Salerno	2020-08-22	USA	15,920	Hospital	49 (30-65)	57.0	-	-	36.78	55.
Rashid et al. (2020)	Rashid	2020-08-22	UK	517	Hospital	72.8 [^] (59-86)	31.9	9.9	29.01	-	29.
Pan et al. (2020)	Pan	2020-08-22	USA	12,084	Community and Hospital	45.5 [^] (27-63)	54.3	-	-	17.51	-
Fillmore et al. (2020)	Fillmore	2020-08-24	USA	22,914	Community and Hospital	NA	-	37.5	40.65	-	15.
Zhou et al. (2020)	Zhou, Qin	2020-08-25	China	51	Hospital	57.37 [^] (42-72)	29.4	-	-	78.43	21.
Ibrahim et al. (2020)	Ibrahim	2020-08-27	USA	38	Hospital	63 [^] (51-75)	47.0	10.5	-	-	-
Oliveira et al. (2020)	Oliveira	2020-08-31	USA	131	Hospital	61 (49.5-71.5)	64.9	-	-	17.56	26.
Yoo et al. (2020)	Yoo	2020-08-31	USA	4,840	Hospital	66.4 (54.9-77.8)	43.5	4.4	21.38	-	53.
Zhan et al. (2020)	Zhan, Liu	2020-08-31	China	405	Hospital	56 [^] (17-95)	54.1	-	-	11.36	88.
Mohamed-Hussein et al. (2020)	Hussein, Galal	2020-09-01	Egypt	444	Community	33.1 [^] (21-45)	56.8	13.1	9.01	-	77.
Villar-Garcia et al. (2020)	Vilar-Garcia	2020-09-01	Spain	7,699,568	Community and Hospital	43 (24-59)	50.9	17.1	-	-	-
Ibarra-Nava et al. (2020)	Ibarra-Nava	2020-09-01	Mexico	416,546	Community and Hospital	NA	46.9	7.4	-	-	-
Rubio-Rivas et al. (2020)	Rubio-Rivas	2020-09-01	Spain	186	Hospital	64.3 [^] (51-77)	30.6	4.3	20.43	-	75.
Mamtani et al. (2020)	Mamtani	2020-09-02	USA	403	Hospital	55 [^] (41-68)	32.3	9.7	12.66	-	68.
Ren et al. (2020)	Ren	2020-09-02	China	432	Hospital	NA	57.9	10.0	-	-	90.
Mutambudzi et al. (2020)	Mutambudzi	2020-09-03	UK	120,075	Community and Hospital	NA	54.2	11.7	26.39	-	61.
Yan et al. (2020)	Yan	2020-09-07	China	578	Hospital	49.2 [^] (35-63)	49.3	9.2	-	-	-
Mancilla-Galindo et al. (2020)	Mancilla-Galindo	2020-09-08	Mexico	183,779	Community and Hospital	45 [^] (28-61)	46.0	7.6	-	-	-
Ullah et al. (2020)	Ullah	2020-09-08	UK	212	Community and Hospital	66.7 (54.2-80.5)	44.8	11.3	48.11	-	37.
Hamadah et al. (2020)	Hamadah	2020-09-10	Kuwait	1,123	Hospital	40 (1-93)	18.7	3.9	-	-	-
					Community						

Dashti et al. (2020)	Dashti	2020-09-13	USA	12,347	Community and Hospital	47 (32-62)	53.3	4.6	15.87	-	57.
Sami et al. (2020)	Sami	2020-09-14	Iran	490	Community and Hospital	56.6 [^] (41-71)	39.0	14.1	-	-	-
Pongpirul et al. (2020)	Pongpirul	2020-09-16	Thailand	193	Community and Hospital	37 (29-53)	41.5	-	-	15.03	66.
Burrell et al. (2020)	Burrell	2020-09-16	Australia	204	Hospital	63.5 (53-72)	31.4	-	-	13.24	-
Nicholson et al. (2020)	Nicholson	2020-09-17	USA	1,042	Hospital	64 (53-75)	43.2	8.3	22.17	-	37.
Ariza et al. (2020)	Ariza	2020-09-18	Colombia	351	Community and Hospital	30.5 (NA)	54.0	6.8	-	-	-
Carrat et al. (2020)	Carrat	2020-09-18	France	14,628	Community	NA	60.3	12.0	40.83	-	45.
Favara et al. (2020)	Favara	2020-09-20	UK	434	Community	40 (19-66)	82.0	8.5	-	-	-
Invernizzi et al. (2020)	Invernizzi	2020-09-20	Italy	54	Hospital	49.9 [^] (34-65)	29.7	-	-	24.07	-
Zhu et al. (2020)	Zhu	2020-09-21	China	432	Community and Hospital	49 (35-60)	47.9	14.4	-	-	-
O'Reilly et al. (2020)	O'Reilly	2020-09-21	Australia	1,334	Hospital	NA	-	-	-	28.49	-
Meini et al. (n.d.)	Meini	2020-09-23	Italy	461	Hospital	NA	51.2	10.4	25.81	-	63.
Silva Neto et al. (2020)	da Silva Neto	2020-09-23	Brazil	91	Community and Hospital	49 [^] (29-68)	49.4	-	-	19.78	-
Ioannou et al. (2020)	Ioannou	2020-09-23	USA	88,747	Community and Hospital	NA	9.0	20.6	37.55	-	29.
Torres-Macho et al. (2020)	Torres-Macho	2020-09-23	Spain	1,968	Hospital	NA	44.0	-	-	23.37	-
Li et al. (2020)	Li, Cai	2020-09-28	China	98	Hospital	68.5 (63-75)	58.2	-	-	11.22	-
Wang et al. (2020)	Wang	2020-09-29	USA	1,078	Hospital	NA	38.2	3.7	24.86	-	49.
Lopez-Medrano et al. (2020)	Lopez-Medrano	2020-09-30	Spain	261	Hospital	NA	43.7	-	-	37.16	-
Collard et al. (2020)	Collard	2020-10-01	Netherlands	1,604	Hospital	65.7 [^] (50-80)	39.5	4.9	-	-	-
Makaronidis et al. (2020)	Makaronidis	2020-10-01	UK	567	Community	39.4 [^] (27-51)	69.1	9.3	-	-	-
Yadaw et al. (2020)	Yadaw	2020-10-01	USA	5,051	Community and Hospital	NA	-	3.6	15.92	-	51.
Talavera et al. (2020)	Talavera	2020-10-01	Spain	576	Hospital	67.2 [^] (52-81)	43.4	-	-	20.49	-
		2020-10-			Community						

Jakob et al. (2020)	Jakob	01	Multiple	2,155	and Hospital	NA	40.3	6.6	7.29	-	34.
Incerti et al. (2020)	Incerti	2020-10-02	USA	13,658	Hospital	62 (49-75)	48.1	6.3	22.64	-	45.
Luo et al. (2020)	Luo, Rizvi	2020-10-03	USA	102	Hospital	68 (61-75)	52.0	-	-	26.47	-
Alharthy et al. (2020)	Alharthy	2020-10-03	Saudi Arabia	352	Hospital	50.6 [^] (37-63)	12.8	49.4	-	-	-
Robinson et al. (2021)	Robinson	2020-10-05	USA	3,248	Hospital	51 [^] (34-68)	72.0	4.0	17.64	-	61.
Adrish et al. (2020)	Adrish	2020-10-05	USA	1,173	Hospital	NA	38.6	14.0	14.66	-	71.
Erber et al. (2020)	Erber	2020-10-06	Germany	4,554	Community	38.5 [^] (NA)	70.4	-	-	18.02	-
Chaudhary et al. (2020)	Chaudhary	2020-10-06	Nepal	220	Hospital	31.5 (25-37)	17.7	11.4	7.73	-	80.
Raines et al. (2021)	Raines	2020-10-07	USA	453	Community and Hospital	60.8 [^] (46-74)	10.7	-	-	52.98	41.
Roederer et al. (2021)	Roederer	2020-10-09	France	818	Community	NA	20.4	36.9	8.80	-	53.
Zinellu et al. (2021)	Zinellu	2020-10-11	Italy	105	Hospital	72 (59.5-80)	33.3	30.5	10.48	-	59.
Ramachandran et al. (2020)	Ramachandran	2020-10-12	USA	188	Hospital	NA	-	18.6	-	-	-
Lamure et al. (2020)	Lamure	2020-10-12	France	89	Hospital	67 (19-92)	34.0	5.6	32.58	-	48.
Ghinai et al. (2020)	Ghinai	2020-10-12	USA	1,435	Homeless Shelters	NA	27.6	36.6	17.49	-	33.
Best et al. (2020)	Best	2020-10-12	USA	3,471	Hospital	63.5 [^] (47-79)	51.2	-	-	28.64	-
Savarraj et al. (2020)	Savarraj	2020-10-18	USA	48	Hospital	50 [^] (33-67)	48.0	10.4	-	-	-
Israel et al. (2020)	Israel, Schaffer	2020-10-18	Israel	26,959	Hospital	NA	50.6	6.8	15.22	-	77.
El-Solh et al. (2020)	El-Solh	2020-10-20	USA	7,816	Hospital	69 (60-74)	5.5	-	-	45.25	-
Perico et al. (2020)	Perico	2020-10-22	Italy	423	Community	44.3 [^] (34-54)	36.4	21.7	17.97	-	60.
Zhou et al. (2020)	Zhou, Song	2020-10-22	China	124	Hospital	67 (30-86)	48.0	19.1	-	-	-
Chudasama et al. (2020)	Chudasama	2020-10-23	UK	1,706	Community and Hospital	68 (48-85)	42.5	13.8	40.97	-	45.
Salama et al. (2021)	Salama	2020-10-23	Multiple	377	Hospital	55.9 [^] (41-70)	40.8	5.8	16.98	-	77.
Wang et al. (2020)	Wang, Zheutlin	2020-10-26	USA	3,273	Hospital	65 (53-77)	42.7	3.5	20.68	-	53.
Zhou et al. (2020)	Zhou, He, Yang	2020-10-27	China	1,087	Hospital	NA	51.7	-	-	85.00	15.
Hoertel et al.	Hoertel, Sanchez	2020-10-									

Fricker et al. (2020)	Fricker, Catherine, Vernet	2020-10-27	France	12,210	Hospital	NA	50.1	9.0	-	-	-
Arleo et al. (2020)	Arleo	2020-10-27	USA	70	Community and Hospital	56.6 [^] (48-65)	80.0	1.4	28.57	-	70.
Bermejo-Martin et al. (2020)	Bermejo-Martin	2020-10-27	Spain	250	Community and Hospital	NA	64.0	6.0	-	-	-
Joubert et al. (2020)	Joubert	2020-10-29	France	74	Community and Hospital	NA	-	5.4	-	-	-
Kortela et al. (2020)	Kortela	2020-11-01	Finland	3,008	Community and Hospital	51 (36-69)	59.6	11.0	14.03	-	29.
Gianfrancesco et al. (2020)	Gianfrancesco, Leykina	2020-11-03	USA	1,324	Community and Hospital	NA	75.9	-	-	26.66	68.
Gallichotte et al. (2020)	Gallichotte	2020-11-05	USA	239	Community	41 [^] (16-76)	-	20.1	16.74	-	57.
Lin et al. (2020)	Lin	2020-11-05	USA	2,821	Hospital	62.7 [^] (NA)	45.0	2.9	12.51	-	3.3
Kim et al. (2020)	Kim, Han	2020-11-09	South Korea	4,787	Hospital	55 (38-68)	60.1	5.5	2.84	-	91.
Galal et al. (2020)	Galal	2020-11-12	Egypt	430	Community	37.4 [^] (24-50)	63.7	6.0	7.67	-	86.
Sourij et al. (2020)	Sourij	2020-11-16	Austria	238	Hospital	71.1 [^] (58-83)	36.1	1.7	15.97	-	82.
Clavario et al. (2020)	Clavario	2020-11-16	Italy	110	Community	61.7 (53.5-69.2)	40.9	40.9	-	-	-
Saeed et al. (2020)	Saeed	2020-11-16	United Arab Emirates	173	Hospital	NA	34.1	6.4	-	-	-
Mansour et al. (2020)	Mansour	2020-11-16	Iran	353	Hospital	61.7 [^] (45-78)	42.5	7.1	-	-	-
Cadegiani et al. (2020)	Cadegiani	2020-11-18	Brazil	130	Community	42 [^] (NA)	0.0	7.7	-	-	-
Ilic et al. (2021)	Ilic	2020-11-19	Serbia	107	Community and Hospital	39.1 [^] (27-50)	-	29.9	-	-	-
Benaim et al. (2020)	Benaim	2020-11-19	Israel	693	Hospital	59.8 (NA)	47.9	-	-	5.05	-
Ho et al. (2020)	Ho	2020-11-19	UK	235,928	Community and Hospital	NA	-	-	-	45.38	-
Singh et al. (2020)	Singh	2020-11-20	UK	930	Hospital	71.4 [^] (54-87)	44.8	-	-	19.03	-
Márquez-Salinas et al. (2020)	Marquez-Salinas	2020-11-24	Mexico	1,068	Hospital	53 (44-63)	36.8	-	-	14.98	-
Díez-Manglano et al. (2020)	Diez-Manglanas	2020-11-24	Spain	4,393	Hospital	53 [^] (NA)	40.8	6.6	-	-	-
Bellan et al. (2020)	Bellan	2020-11-26	Italy	1,697	Hospital	71 (58-80)	41.0	3.2	-	-	-
Woolcott and		2020-11-			Community	42 (34-					

Castilla-Bancayán (2020)	Woolcott	2020-11-26	Mexico	1,636,050	and Hospital	54)	51.9	-	-	8.76	-
Yao et al. (2021)	Yao, Hasegawa	2020-11-26	Japan	101	Hospital	60 [^] (17-97)	39.6	-	-	28.71	-
Chen et al. (2020)	Chen, Varathraja	2020-11-29	USA	10,123	Community and Hospital	40 (28-54)	53.6	4.4	9.71	-	45.
Serling-Boyd et al. (2020)	Serling-Boyd	2020-11-30	USA	831	Hospital	NA	76.0	3.0	23.23	-	50.
Simons et al. (2020)	Simons	2020-11-30	UK	446	Hospital	64.9 (52.4-76.2)	35.9	9.4	38.57	-	52.
Dupraz et al. (2020)	Dupraz	2020-11-30	Switzerland	219	Community	NA	54.8	11.0	-	-	-
Barasa et al. (2020)	Barasa	2020-11-30	USA	394	Hospital	NA	47.7	14.7	36.04	-	41.
Ren et al. (2020)	Ren, Guo	2020-11-30	China	481	Hospital	NA	45.7	7.7	0.62	-	91.
Martinez-Lacalzada et al. (2020)	Martinez-Lacalzada	2020-12-01	Spain	10,433	Hospital	65.8 [^] (49-82)	42.8	5.2	23.77	-	66.
Li et al. (2020)	Li, Long, Zhang	2020-12-03	China	954	Hospital	NA	-	-	-	5.87	94.
Martini et al. (2020)	Martini	2020-12-04	Italy	146	Hospital	NA	49.0	-	-	46.58	53.
O'Gallagher et al. (2020)	O'Gallagher	2020-12-04	UK	1,721	Hospital	71 (56-83)	43.4	6.6	18.48	-	74.
Alguwaihes et al. (2020)	Alguwaihes	2020-12-05	Saudi Arabia	439	Hospital	55 (19-101)	31.7	2.1	-	-	-
Zuo et al. (2020)	Zuo, Warnock	2020-12-05	USA	118	Hospital	61 [^] (44-78)	46.0	-	-	23.73	-
Zhang et al. (2020)	Zhang, Li	2020-12-06	UK	1,746	Community and Hospital	68.8 [^] (59-78)	47.1	10.1	35.09	-	44.
Dai et al. (2020)	Dai	2020-12-09	China	1,574	Hospital	57.3 [^] (41-73)	48.2	-	-	9.21	90.
Vila-Córcoles et al. (2020)	Vila-Corcoles	2020-12-10	Spain	79,083	Community	NA	52.4	-	-	16.12	-
Bisso et al. (2020)	Bisso	2020-12-11	Argentina	168	Hospital	67 (58-75)	34.0	10.7	-	-	-
Rentsch et al. (2020)	Rentsch, Beckman	2020-12-11	USA	4,297	Hospital	68 (58-75)	6.6	36.8	39.31	-	1.9
Thiabaud et al. (2020)	Thiabaud	2020-12-11	Switzerland	3,582	Hospital	68 (54-79)	40.5	-	-	6.62	-
Vila-Corcoles et al. (2020)	Vila-Corcoles, Satue-Gracia	2020-12-11	Spain	282	Community and Hospital	65.9 [^] (53-78)	50.3	8.9	-	-	-
Lévy et al. (2020)	Levy	2020-12-12	France	61	Hospital	60 (50-69)	20.0	8.2	-	-	-
Kantele et al. (2020)	Kantele	2020-12-13	Finland	1,095	Community and Hospital	38 (31-48)	82.7	16.3	25.84	-	56.

Iftime et al. (2020)	Iftimie	2020-12-14	Spain	468	Hospital	NA	44.9	-	-	7.91	-
Ho et al. (2020)	Ho, Narasimhan	2020-12-15	USA	9,991	Community and Hospital	58 [^] (39-76)	45.9	4.1	18.08	-	77.
Caliskan and Saylan (2020)	Caliskan	2020-12-16	Turkey	565	Hospital	48 (38-58)	-	20.9	13.98	-	65.
Muñoz et al. (2020)	Munoz	2020-12-16	Spain	314	Community	45 (40-53)	52.5	6.7	-	-	-
Crooks et al. (2020)	Crooks	2020-12-16	UK	2,964	Hospital	NA	52.6	11.7	-	-	-
Núñez-Gil et al. (2020)	Nunez-Gil	2020-12-17	Multiple	2,798	Hospital	67 (53-78)	40.0	6.2	-	-	-
Gori et al. (2020)	Gori	2020-12-17	Italy	1,352	Hospital	68 (58-77)	28.4	3.6	18.42	-	66.
Rowlands et al. (2020)	Rowlands	2020-12-18	UK	580	Community	63.8 [^] (56-70)	52.2	10.2	38.10	-	51.
Richard et al. (2020)	Richard	2020-12-18	Switzerland	8,344	Community	46.9 [^] (NA)	53.5	15.2	17.11	-	66.
Schubl et al. (2020)	Schuble*	2020-12-19	USA	1,557	Community	NA	68.9	2.4	-	-	-
Ugur Chousein et al. (2020)	Chousein	2020-12-21	Turkey	114	Hospital	51.1 [^] (36-66)	32.5	16.7	20.18	-	63.
Modrák et al. (2020)	Modrak	2020-12-22	Czechia	213	Hospital	69 (58-80)	51.0	-	-	13.15	-
Kara Polat et al. (2020)	Polat	2020-12-22	Turkey	1,322	Community	NA	47.6	35.6	15.58	-	48.
Kjetland et al. (2020)	Kjetland	2020-12-24	Norway	7,839	Community and Hospital	45.3 [^] (33-56)	77.0	-	-	41.00	-
Lewnard et al. (2021)	Lewnard	2021-01-02	USA	1,115	Community	NA	52.6	4.4	14.17	-	81.
Nezhadmoghadam and Tamez-Peña (2021)	Nezhadmoghadam	2021-01-04	Mexico	33,325	Hospital	NA	-	10.2	-	-	-
Covid-19 in pregnancy et al. (2021)	Vousden	2021-01-05	UK	1,148	Community and Hospital	NA	100.0	8.6	-	-	-
Boyd and Martin-Loeches (2021)	Boyd	2021-01-07	Ireland	38	Hospital	NA	26.3	-	-	5.26	-
Paleiron et al. (2021)	Paleiron	2021-01-09	France	1,688	Community	28 (23-35)	13.0	48.3	22.87	-	28.
Ader et al. (2021)	Ader	2021-01-09	Multiple	583	Hospital	63 (54-71)	28.3	3.1	-	-	-
Giannini et al. (2021)	Giannini	2021-01-10	Italy	91	Hospital	74 [^] (61-87)	45.0	23.1	-	-	-
Shade et al. (2021)	Shade	2021-01-10	USA	3,779	Community and Hospital	NA	47.2	3.3	15.74	-	-
Park et al. (2021)	Park	2021-01-11	South Korea	2,269	Hospital	55.5 [^] (35-75)	64.1	4.1	-	-	-
Ferrari et al. (2021)	Ferrari	2021-01-	Brazil	188	Community	61 (19-	65.0	-	-	20.71	70.

Author(s)	Study Name	Date	Country	Sample Size	Setting	Prevalence (%)	95% CI Lower	95% CI Upper	OR	95% CI Lower	95% CI Upper
Zhang et al. (2021)	Zhang, Yang	2021-01-12	UK	1,485	Community and Hospital	68.2 [^] (59-77)	47.2	48.5	39.19	-	11.0
Ebrahimian et al. (2021)	Ebrahimian	2021-01-13	USA	226	Hospital	NA	-	-	-	10.62	-
Vahidy et al. (2021)	Vahidy	2021-01-13	USA	96,473	Hospital	51.2 [^] (32-69)	59.6	-	-	26.18	73.0
Mendes et al. (2021)	Mendes	2021-01-14	Switzerland	265	Hospital	85.9 [^] (79-92)	57.0	5.7	-	-	-
Saurabh et al. (2021)	Saurabh	2021-01-14	India	911	Community	43.1 [^] (23-62)	35.2	8.1	0.66	-	91.0
Ayoubkhani et al. (2021)	Ayoubkhani	2021-01-15	UK	47,780	Hospital	NA	45.1	8.4	40.94	-	42.0
Thakur et al. (2021)	Thakar	2021-01-18	India	250	Hospital	NA	42.4	-	-	49.20	-
Zhong et al. (2021)	Zhong	2021-01-18	China	91	Hospital	47.3 [^] (30-64)	49.5	-	-	18.68	-
Cummins et al. (2021)	Cummins	2021-01-20	UK	1,781	Community and Hospital	NA	44.8	10.2	-	-	-
Sun et al. (2021)	Sun	2021-01-21	USA	323	Community and Hospital	NA	57.6	-	-	39.32	-
Lowe et al. (2021)	Lowe	2021-01-25	USA	7,102	Hospital	50.3 [^] (NA)	61.2	2.4	12.81	-	84.0
Caglar and Kacer (2021)	Caglar	2021-01-25	Turkey	120	Hospital	57 (47-67)	51.7	-	-	37.50	-
Quan et al. (2021)	Quan	2021-01-27	USA	2,038	Hospital	64 [^] (47-80)	50.4	-	-	39.74	-
Strangfeld et al. (2021)	Strangfeld	2021-01-27	Multiple	3,729	Community and Hospital	57 [^] (41-72)	68.0	-	-	20.81	64.0
Tardif et al. (2021)	Tardif	2021-01-27	Canada	4,488	Community and Hospital	NA	53.9	9.6	33.60	-	56.0
Jafari et al. (2021)	Jafari	2021-01-28	USA	1,979	Hospital	66 (57-74)	32.0	12.1	-	-	-
Prats-Urbe et al. (2021)	Prats-Urbe	2021-01-30	Spain	696	Hospital	63 [^] (52-73)	30.9	-	-	16.09	74.0
Nuño et al. (2021)	Nuno	2021-02-01	USA	4,730	Hospital	61 (46-73)	43.6	10.1	-	-	-
Abajo et al. (2021)	de Abajo	2021-02-03	Spain	625	Hospital	NA	39.4	4.6	29.28	-	39.0
Mora et al. (2021)	Mora	2021-02-03	USA	1,058	Community	39.7 [^] (27-52)	52.5	4.5	14.56	-	80.0
Molenaar et al. (2021)	Molenaar	2021-02-03	USA	696	Community and Hospital	33.1 [^] (NA)	100.0	4.5	-	-	-
Leister et al. (2021)	Leister	2021-02-03	Austria	3,301	Community and Hospital	43.6 [^] (33-54)	68.0	23.3	-	-	-

Didikoglu et al. (2021)	Didikoglu	2021-02-04	UK	384,816	Community and Hospital	68.3 [^] (60-76)	54.9	9.8	35.19	-	55.
Estiri et al. (2021)	Estiri	2021-02-04	USA	16,709	Hospital	NA	57.2	-	-	8.62	-
Tavakol et al. (2021)	Tavakol	2021-02-04	Iran	206	Community and Hospital	40.9 [^] (29-52)	57.8	-	-	12.62	-
Cai et al. (2021)	Cai, Yang	2021-02-05	China	455	Hospital	NA	52.6	7.5	-	-	-
Lohia et al. (2021)	Lohia	2021-02-05	USA	1,871	Hospital	64.1 [^] (48-80)	48.4	-	-	37.63	-
Vila-Corcoles et al. (2021)	Vila-Corcoles, Satue-Gracia, Vila-Rovira	2021-02-05	Spain	79,083	Community	65.8 [^] (54-77)	52.4	16.1	-	-	-
International Severe Acute Respiratory and emerging Infections Consortium et al. (2020)	ISARIC	2021-02-11	Multiple	240,149	Hospital	60 (NA)	49.0	-	-	3.63	21.
Gégout petit et al. (2021)	Gegout Petit	2021-02-12	France	2,006	Community	NA	55.0	16.8	-	-	-
Zhang et al. (2021)	Zhang, Wang	2021-02-13	China	172	Hospital	47.9 [^] (29-66)	46.5	7.0	1.16	91.86	-

* grouped together current smokers and vapers

Smoking prevalence by country

Unadjusted smoking prevalence compared with overall estimates for national adult smoking prevalence split by country and study setting is presented in Figure 2a and 2b. Lower than expected current smoking prevalence was generally observed, especially in studies with hospitalised samples. Former smoking prevalence was more similar to expected prevalence when reported; however, study-based prevalence was typically higher than national estimates. National smoking prevalence estimates used for comparison are presented in Supplementary table 3.




Figure 2a. Weighted mean prevalence of current smoking in included studies, split by country (solid lines). The circles represent individual studies, with colour corresponding to study setting (i.e. community, community and hospital, hospital) and size corresponding to relative study sample size. For comparison, national current smoking prevalence is shown by the dashed red lines. Countries with three or more eligible studies are shown




Figure 2b. Prevalence of former smoking in included studies, split by country (solid lines). The circles represent individual studies, with colour corresponding to study setting (i.e. community, community and hospital, hospital) and size corresponding to relative study sample size. For comparison, national former smoking prevalence is shown by dashed red lines. Countries with three or more eligible studies are shown.

SARS-CoV-2 testing by smoking status

Four studies provided data on access to SARS-CoV-2 diagnostic testing for those meeting local testing criteria by smoking status. In a cohort study of US military veterans aged 54-75 (Rentsch et al., 2020), current smokers were more likely to receive a test: 42.3% (1,603/3,789) of the sample were current smokers compared with 23.8% of all veterans aged 50+

years using any tobacco product between 2010-2015 (Odani, 2018). In the UK Biobank cohort (Niedzwiedz et al., 2020), a multivariable analysis showed former (RR = 1.29, 95% CI = 1.14-1.45, $p < .001$) and current (RR = 1.44, 95% CI = 1.20-1.71, $p < .001$) compared with never smokers to be more likely to receive a test. In an Australian rapid assessment screening clinic for COVID-19 (Trubiano et al., 2020), 9.4% (397/4,226) of the self-referred sample (subsequently assessed by a healthcare professional to decide on testing) were current smokers. Of these self-referrals, healthcare professionals decided that current compared with former or never smokers were less likely to require a test (RR = 0.93, 95% CI = 0.86-1.0, $p = 0.045$). In a further study using the UK Biobank cohort (Didikoglu et al., 2021), current (RR = 1.23, 95% CI = 1.19-1.26, $p < 0.001$) and former smokers (RR = 1.20, 95% CI = 1.18-1.23, $p < 0.001$) were more likely to receive a test compared with never smokers.

SARS-CoV-2 infection by smoking status

76 studies provided data on SARS-CoV-2 infection for people meeting local testing criteria by smoking status (see Table 2). Meta-analyses were performed for 3 'good' and 27 'fair' quality studies (see Figure 3 and 4). Current smokers were at reduced risk of testing positive for SARS-CoV-2 compared with never smokers (RR = 0.71, 95% Credible Interval (CrI) = 0.61-0.82, $\tau = 0.34$). The three good quality studies each reported point estimates less than 1, although the CrI for one of the three studies crossed 1. The probability of current smokers being at reduced risk of infection compared with never smokers (RR ≤ 0.9) was >99%. Former compared with never smokers were at increased risk of testing positive, but data were inconclusive (RR = 1.03, 95% CrI = 0.95-1.11, $\tau = 0.17$) and favoured there being no important association. The probability of former smokers being at increased risk of infection (RR ≥ 1.1) compared with never smokers was 4%. Results were materially unchanged in sensitivity analyses.

Table 2 SARS-CoV-2 infection by smoking status

Author	Total population tested	SARS-CoV-2 negative						SARS-CoV-2 positive			
		N (%)	Current smoker (%)	Former smoker (%)	Current/former smoker (%)	Never smoker (%)	Not stated (%)	N (%)	Current smoker (%)	Former smoker (%)	Current/former smoker (%)
Rentsch	3,528	2974 (84.30%)	1444 (48.55%)	704 (23.67%)	-	826 (27.77%)	-	554 (15.70%)	159 (28.70%)	179 (32.31%)	-
Fontanet	661	490 (74.13%)	64 (13.06%)	-	-	426 (86.94%)	-	171 (25.87%)	5 (2.92%)	-	-
Cho	1,331	793 (59.58%)	142 (17.91%)	214 (26.99%)	-	437 (55.11%)	-	538 (40.42%)	111 (20.63%)	145 (26.95%)	-
Shah	243	212 (87.24%)	52 (24.53%)	47 (22.17%)	-	113 (53.30%)	-	29 (11.93%)	0 (0.00%)	9 (31.03%)	-
Kolin	1,474	805 (54.61%)	141 (17.52%)	307 (38.14%)	-	354 (43.98%)	3 (0.37%)	669 (45.39%)	72 (10.76%)	285 (42.60%)	-
de Lusignan	3,291	2740 (83.26%)	366 (13.36%)	1450 (52.92%)	-	924 (33.72%)	-	551 (16.74%)	47 (8.53%)	303 (54.99%)	-
Valenti	789	689 (87.33%)	197 (28.59%)	-	-	-	492 (71.41%)	40 (5.07%)	7 (17.50%)	-	-

Parrotta	76	39 (51.32%)	1 (2.56%)	10 (25.64%)	-	27 (69.23%)	1 (2.56%)	37 (48.68%)	1 (2.70%)	10 (27.03%)	-
Berumen	102,875	71353 (69.36%)	-	-	7173 (10.05%)	64180 (89.95%)	-	31522 (30.64%)	-	-	2748 (8.72%)
Israel	24,906	20755 (83.33%)	3783 (18.23%)	2671 (12.87%)	-	14301 (68.90%)	-	4151 (16.67%)	406 (9.78%)	483 (11.64%)	-
del Valle	1,108	143 (12.91%)	27 (18.88%)	53 (37.06%)	-	-	63 (44.06%)	965 (87.09%)	55 (5.70%)	293 (30.36%)	-
Romao	34	20 (58.82%)	-	-	5 (25.00%)	-	15 (75.00%)	14 (41.18%)	-	-	4 (28.57%)
Ramlall	11,116	4723 (42.49%)	-	-	-	-	-	6393 (57.51%)	-	-	1643.001 (25.70%)
Sharma	501	267 (53.29%)	-	-	1 (0.37%)	-	266 (99.63%)	234 (46.71%)	-	-	20 (8.55%)
Eugen-Olsen	407	290 (71.25%)	76 (26.21%)	104 (35.86%)	-	102 (35.17%)	-	117 (28.75%)	8 (6.84%)	46 (39.32%)	-
Raisi-Estabragh	4,510	3184 (70.60%)	-	-	1653 (51.92%)	-	1531 (48.08%)	1326 (29.40%)	-	-	683 (51.51%)
Houlihan	177	97 (54.80%)	14 (14.43%)	14 (14.43%)	-	69 (71.13%)	-	80 (45.20%)	7 (8.75%)	19 (23.75%)	-
McQueenie	428,199	424355 (99.10%)	-	-	189299 (44.61%)	235056 (55.39%)	-	1311 (0.31%)	-	-	669 (51.03%)
Woolford	4,474	3161 (70.65%)	441 (13.95%)	1194 (37.77%)	-	1526 (48.28%)	-	1313 (29.35%)	145 (11.04%)	525 (39.98%)	-
Lan	104	83 (79.81%)	-	-	24 (28.92%)	-	59 (71.08%)	21 (20.19%)	-	-	1 (4.76%)
Hernandez, Garduno	32,583	20279 (62.24%)	-	-	2399 (11.83%)	17861 (88.08%)	-	12304 (37.76%)	-	-	1191 (9.68%)
Govind	6,215	6207 (99.87%)	4104 (66.12%)	1669 (26.89%)	-	342 (5.51%)	-	102 (1.64%)	78 (76.47%)	20 (19.61%)	-
Gu	4,699	3815 (81.19%)	360 (9.44%)	1142 (29.93%)	-	2313 (60.63%)	-	884 (18.81%)	40 (4.52%)	264 (29.86%)	-
Kibler	702	680 (96.87%)	25 (3.68%)	-	-	-	655 (96.32%)	22 (3.13%)	1 (4.55%)	-	-
Auvinen	61	33 (54.10%)	10 (30.30%)	8 (24.24%)	-	15 (45.45%)	-	28 (45.90%)	1 (3.57%)	9 (32.14%)	-
Antonio-Villa	34,263	23338 (68.11%)	2293 (9.83%)	-	-	-	21045 (90.17%)	10925 (31.89%)	1023 (9.36%)	-	-
Merzon	7,807	7025 (89.98%)	-	-	1136 (16.17%)	-	5889 (83.83%)	782 (10.02%)	-	-	127 (16.24%)
Trubiano	2,935	2827 (96.32%)	-	-	256 (9.06%)	-	2586 (91.48%)	108 (3.68%)	-	-	3 (2.78%)
Shi, Resurreccion	1,521	1265 (83.17%)	-	-	681 (53.83%)	-	584 (46.17%)	256 (16.83%)	-	-	154 (60.16%)
Riley	120,620	120461 (99.87%)	2594 (2.15%)	-	-	19914 (16.53%)	97953 (81.32%)	159 (0.13%)	3 (1.89%)	-	-
Alizadehsani	319	196 (61.44%)	-	-	-	-	196 (100.00%)	123 (38.56%)	-	-	1 (0.81%)
Merkely	10,474	10336 (98.68%)	2904 (28.10%)	2107 (20.39%)	-	5310 (51.37%)	15 (0.15%)	70 (0.67%)	16 (22.86%)	15 (21.43%)	-
Edwards	209	118	-	-	31 (26.27%)	-	87	91	-	-	8 (8.79%)

		(56.46%)				(73.73%)	(43.54%)				
Reiter	235	175 (74.47%)	-	-	93 (53.14%)	82 (46.86%)	-	60 (25.53%)	-	-	13 (21.67%)
Izquierdo	71,192	NA (NA%)	-	-	-	-	-	1006 (1.41%)	111 (11.03%)	-	-
Ward	99,908	94416 (94.50%)	10202 (10.81%)	-	-	-	84214 (89.19%)	5492 (5.50%)	433 (7.88%)	-	-
Ebinger	6,062	5850 (96.50%)	99 (1.69%)	-	-	-	5668 (96.89%)	212 (3.50%)	3 (1.42%)	-	-
Salerno	15,920	14753 (92.67%)	-	-	5517 (37.40%)	8278 (56.11%)	958 (6.49%)	1167 (7.33%)	-	-	339 (29.05%)
Iversen	28,792	27629 (95.96%)	4430 (16.03%)	1799 (6.51%)	-	21217 (76.79%)	246 (0.89%)	1163 (4.04%)	177 (15.22%)	78 (6.71%)	-
Hippisley-Cox	8,275,949	NA (NA%)	-	-	-	-	-	19486 (0.24%)	1354 (6.95%)	5715 (29.33%)	-
Fillmore	22,914	21120 (92.17%)	8137 (38.53%)	8416 (39.85%)	-	3227 (15.28%)	1340 (6.34%)	1794 (7.83%)	452 (25.20%)	899 (50.11%)	-
Alkurt	119	NA (NA%)	-	-	-	-	-	119 (100.00%)	14 (11.76%)	-	-
Petrilli	10,620	5341 (50.29%)	3454 (64.67%)	816 (15.28%)	-	541 (10.13%)	530 (9.92%)	5279 (49.71%)	3268 (61.91%)	902 (17.09%)	-
Bello-Chavolla	150,200	98567 (65.62%)	-	-	9624 (9.76%)	-	88943 (90.24%)	51633 (34.38%)	-	-	4366 (8.46%)
Ariza	351	322 (91.74%)	21 (6.52%)	-	-	-	301 (93.48%)	29 (8.26%)	3 (10.34%)	-	-
Carrat	14,393	13426 (93.28%)	1652 (12.30%)	5620 (41.86%)	-	6154 (45.84%)	-	967 (6.72%)	98 (10.13%)	353 (36.50%)	-
Meini	461	243 (52.71%)	39 (16.05%)	66 (27.16%)	-	138 (56.79%)	-	218 (47.29%)	9 (4.13%)	53 (24.31%)	-
Favara	434	354 (81.57%)	28 (7.91%)	-	-	-	326 (92.09%)	80 (18.43%)	9 (11.25%)	-	-
Erber	4,554	4446 (97.63%)	-	-	806 (18.13%)	-	3640 (81.87%)	108 (2.37%)	-	-	11 (10.19%)
Roederer	815	390 (47.85%)	175 (44.87%)	32 (8.21%)	-	183 (46.92%)	-	425 (52.15%)	127 (29.88%)	40 (9.41%)	-
Makaronidis	567	127 (22.40%)	16 (12.60%)	-	-	-	111 (87.40%)	440 (77.60%)	37 (8.41%)	-	-
Ioannou	88,747	78616 (88.58%)	17138 (21.80%)	29245 (37.20%)	-	22327 (28.40%)	9906 (12.60%)	10131 (11.42%)	1135 (11.20%)	4073 (40.20%)	-
Perico	423	260 (61.47%)	69 (26.54%)	35 (13.46%)	-	156 (60.00%)	-	163 (38.53%)	23 (14.11%)	41 (25.15%)	-
Vila-Corcoles	2,324	1944 (83.65%)	-	-	-	-	-	380 (16.35%)	-	-	27 (7.11%)
O'Reilly	1,334	1284 (96.25%)	-	-	376 (29.28%)	-	908 (70.72%)	50 (3.75%)	-	-	4 (8.00%)
Ghinai	1,435	1004 (69.97%)	412 (41.04%)	155 (15.44%)	-	341 (33.96%)	96 (9.56%)	431 (30.03%)	113 (26.22%)	96 (22.27%)	-
Kortela	2,993	2419 (80.82%)	300 (12.40%)	340 (14.06%)	-	636 (26.29%)	1143 (47.25%)	574 (19.18%)	26 (4.53%)	80 (13.94%)	-
Gallichotte	239	190 (79.50%)	40 (21.05%)	28 (14.74%)	-	110 (57.89%)	-	49 (20.50%)	8 (16.33%)	12 (24.49%)	-

Saeed	173	69 (39.88%)	2 (2.90%)	-	-	-	67 (97.10%)	104 (60.12%)	9 (8.65%)	-	-
Woolcott	1,636,050	878840 (53.72%)	-	-	85816 (9.76%)	-	793024 (90.24%)	757210 (46.28%)	-	-	57451 (7.59%)
Barasa	394	277 (70.30%)	49 (17.69%)	105 (37.91%)	-	110 (39.71%)	13 (4.69%)	117 (29.70%)	9 (7.69%)	37 (31.62%)	-
Paleiron	1,688	409 (24.23%)	236 (57.70%)	77 (18.83%)	-	96 (23.47%)	-	1279 (75.77%)	579 (45.27%)	309 (24.16%)	-
Didikoglu	43,428	35695 (82.19%)	3919 (10.98%)	13841 (38.78%)	-	17939 (50.26%)	-	7733 (17.81%)	867 (11.21%)	2966 (38.36%)	-
Kantele	1,095	1059 (96.71%)	176 (16.62%)	272 (25.68%)	-	611 (57.70%)	-	36 (3.29%)	3 (8.33%)	11 (30.56%)	-
Polat	1,322	NA (NA%)	-	-	-	-	-	23 (1.74%)	4 (17.39%)	2 (8.70%)	-
Richard	8,344	6798 (81.47%)	1065 (15.67%)	1171 (17.23%)	-	4456 (65.55%)	106 (1.56%)	531 (6.36%)	47 (8.85%)	83 (15.63%)	-
Schubl	1,557	1392 (89.40%)	33 (2.37%)	-	-	-	1359 (97.63%)	165 (10.60%)	4 (2.42%)	-	-
Nezhadmoghadam	33,325	19958 (59.89%)	2151 (10.78%)	-	-	-	17807 (89.22%)	13367 (40.11%)	1251 (9.36%)	-	-
Mora	1,058	857 (81.00%)	40 (4.67%)	118 (13.77%)	-	698 (81.45%)	1 (0.12%)	201 (19.00%)	8 (3.98%)	36 (17.91%)	-
Molenaar	696	591 (84.91%)	29 (4.91%)	-	-	-	562 (95.09%)	105 (15.09%)	2 (1.90%)	-	-
Vila-Corcoles, Satue-Gracia, Vila- Rovira	4,113	3577 (86.97%)	591 (16.52%)	-	-	-	2986 (83.48%)	536 (13.03%)	41 (7.65%)	-	-
Gegout Petit	2,006	1964 (97.91%)	334 (17.01%)	-	-	1545 (78.67%)	85 (4.33%)	42 (2.09%)	4 (9.52%)	-	-
Leister	3,301	3269 (99.03%)	764 (23.37%)	-	-	-	2505 (76.63%)	32 (0.97%)	5 (15.62%)	-	-




Figure 3. Forest plot for risk of testing positive for SARS-CoV-2 in current vs. never smokers. * Indicates 'good' quality studies. The prior from the previous review version (v10) was RR = 0.69.




Figure 4. Forest plot for risk of testing positive for SARS-CoV-2 in former vs. never smokers. * Indicates 'good' quality studies. The prior from the previous review version (v10) was RR = 1.02.

Hospitalisation for COVID-19 by smoking status

48 studies examined hospitalisation for COVID-19 disease, stratified by smoking status (see Table 3). Meta-analyses were performed for 16 'fair' quality studies (see Figure 5 and 6). Current (RR = 1.1, 95% CrI = 0.99-1.21, $\tau = 0.15$) and former (RR = 1.19, CrI = 1.1-1.29, $\tau = 0.13$) compared with never smokers were at increased risk of hospitalisation with COVID-19. However, data for current smokers were inconclusive and favoured there being no important association. The probability of current and former smokers being at increased risk of hospitalisation (RR ≥ 1.1) compared with never smokers was 49% and 98%, respectively. Results were materially unchanged in two sensitivity analyses.

Table 3 COVID-19 hospitalisation by smoking status

Author	Population with outcome	Community							Hospitalised			
		N (%)	Current smoker (%)	Former smoker (%)	Current/former smoker (%)	Never smoker (%)	Never/unknown smoker (%)	Not stated (%)	N (%)	Current smoker (%)	Former smoker (%)	Current smoker (%)
Rentsch	554	269 (48%)	69 (25.65%)	90 (33.46%)	-	110 (40.89%)	-	-	285 (51%)	90 (31.58%)	89 (31.23%)	-
Chow (US CDC)	6,637	5143 (77%)	61 (1.19%)	80 (1.56%)	-	-	-	5002 (97.26%)	1494 (22%)	27 (1.81%)	78 (5.22%)	-
Argenziano	1,000	151 (15%)	14 (9.27%)	18 (11.92%)	-	119 (78.81%)	-	-	849 (84%)	35 (4.12%)	161 (18.96%)	-
Lubetzky	54	15 (27%)	-	-	4 (26.67%)	-	-	11 (73.33%)	39 (72%)	-	-	8 (2)
Carillo-Vega	9,946	3922 (39%)	408 (10.40%)	-	-	-	-	3514 (89.60%)	6024 (60%)	486 (8.07%)	-	-
Yanover	4,353	4180 (96%)	484 (11.58%)	118 (2.82%)	-	3578 (85.60%)	-	-	173 (3%)	30 (17.34%)	11 (6.36%)	-
Hamer	387,109	386349 (99%)	37333 (9.66%)	134542 (34.82%)	-	214474 (55.51%)	-	-	760 (0%)	93 (12.24%)	313 (41.18%)	-
Heili-Frades	4,712	1973 (41%)	121 (6.13%)	222 (11.25%)	-	-	1630 (82.62%)	1630 (82.62%)	2739 (58%)	112 (4.09%)	598 (21.83%)	-
Freites	123	69 (56%)	1 (1.45%)	-	-	-	-	68 (98.55%)	54 (43%)	3 (5.56%)	-	-
Berumen	102,875	18832 (18%)	-	-	1546 (8.21%)	-	17286 (91.79%)	-	12690 (12%)	-	-	120
Gianfrancesco	600	323 (53%)	-	-	61 (18.89%)	-	-	262 (81.11%)	277 (46%)	-	-	68 (
Chaudhry	40	19 (47%)	-	-	0 (0.00%)	-	-	19 (100.00%)	21 (52%)	-	-	6 (2
Giannouchos	89,756	58485 (65%)	4679 (8.00%)	-	-	-	53806 (92.00%)	-	31271 (34%)	2721 (8.70%)	-	-
Wang, Oekelen	57	22 (38%)	-	-	6 (27.27%)	-	-	16 (72.73%)	36 (63%)	-	-	15 (
Miyara	470	132 (28%)	14 (10.61%)	41 (31.06%)	-	77 (58.33%)	-	-	338 (71%)	18 (5.33%)	111 (32.84%)	-
Suleyman	463	108 (23%)	-	-	23 (21.30%)	-	-	85 (78.70%)	355 (76%)	-	-	137
Garassino	196	48 (24%)	10 (20.83%)	27 (56.25%)	-	11 (22.92%)	-	-	152 (77%)	38 (25.00%)	84 (55.26%)	-
Siso-Almirall	260	119 (45%)	-	-	31 (26.05%)	-	-	88 (73.95%)	141 (54%)	-	-	50 (
Gu	884	511 (57%)	30 (5.87%)	126 (24.66%)	-	355 (69.47%)	-	-	373 (42%)	10 (2.68%)	138 (37.00%)	-
Killerby	531	311 (58%)	-	-	37 (11.90%)	222 (71.38%)	-	52 (16.72%)	220 (41%)	-	-	54 (
Nguyen	689	333 (48%)	-	-	57 (17.12%)	-	-	276 (82.88%)	356 (51%)	-	-	114
Mendv	689	473	-	-	84 (17.76%)	-	-	389	216	-	-	86 (

Author	Total	Category 1	Category 2	Category 3	Category 4	Category 5	Category 6	Category 7	Category 8	Category 9	Category 10	Category 11	Category 12
Soares	10,713	9561 (89%)	132 (1.38%)	-	-	-	-	9429 (98.62%)	-	1152 (10%)	77 (6.68%)	-	-
Zobairy	203	65 (32%)	1 (1.54%)	-	-	-	-	64 (98.46%)	-	138 (67%)	11 (7.97%)	-	-
Izquierdo	1,006	743 (73%)	52 (7.00%)	-	-	-	-	691 (93.00%)	-	263 (26%)	16 (6.08%)	-	-
Rizzo	76,819	60039 (78%)	3931 (6.55%)	11379 (18.95%)	-	-	30042 (50.04%)	-	14687 (24.46%)	16780 (21%)	1254 (7.47%)	4585 (27.32%)	-
Pan	12,084	8548 (70%)	-	-	1263 (14.78%)	-	-	-	7285 (85.22%)	3536 (29%)	-	-	874
Petrilli	5,279	2538 (48%)	147 (5.79%)	337 (13.28%)	-	-	1678 (66.12%)	-	376 (14.81%)	2741 (51%)	141 (5.14%)	565 (20.61%)	-
Vilar-Garcia	328,892	291254 (88%)	64792 (22.25%)	-	-	-	-	-	226462 (77.75%)	37638 (11%)	9526 (25.31%)	-	-
Ibarra-Nava	416,546	302693 (72%)	26773 (8.84%)	-	-	-	-	-	275920 (91.16%)	113853 (27%)	8875 (7.80%)	-	-
Dashti	12,347	8946 (72%)	353 (3.95%)	1099 (12.28%)	-	-	5133 (57.38%)	-	2361 (26.39%)	3401 (27%)	210 (6.17%)	860 (25.29%)	-
da Silva Neto	91	44 (48%)	-	-	4 (9.09%)	-	-	40 (90.91%)	-	47 (51%)	-	-	14 (15%)
Israel, Schaffer	26,676	13706 (51%)	944 (6.89%)	2166 (15.80%)	-	-	10596 (77.31%)	-	-	12970 (48%)	880 (6.78%)	1936 (14.93%)	-
Ioannou	10,131	6624 (65%)	716 (10.81%)	2484 (37.50%)	-	-	2542 (38.38%)	-	882 (13.32%)	3507 (34%)	419 (11.95%)	1593 (45.42%)	-
Zhang, Li	1,596	576 (36%)	63 (10.94%)	190 (32.99%)	-	-	318 (55.21%)	-	5 (0.87%)	1020 (63%)	116 (11.37%)	429 (42.06%)	-
Parra-Bracamonte	331,298	235840 (71%)	-	-	16676 (7.07%)	-	-	-	219164 (92.93%)	95458 (28%)	-	-	751
Jehi	4,536	3578 (78%)	247 (6.90%)	943 (26.36%)	-	-	1795 (50.17%)	-	593 (16.57%)	958 (21%)	82 (8.56%)	349 (36.43%)	-
Arleo	70	36 (51%)	1 (2.78%)	10 (27.78%)	-	-	25 (69.44%)	-	-	34 (48%)	0 (0.00%)	10 (29.41%)	-
Kortela	604	246 (40%)	12 (4.88%)	14 (5.69%)	-	-	55 (22.36%)	-	165 (67.07%)	328 (54%)	14 (4.27%)	66 (20.12%)	-
Didikoglu	7,733	5239 (67%)	554 (10.57%)	1845 (35.22%)	-	-	2841 (54.23%)	-	-	2494 (32%)	313 (12.55%)	1121 (44.95%)	-
Ho, Narasimhan	9,991	5082 (50%)	189 (3.72%)	744 (14.64%)	-	-	4148 (81.62%)	-	-	4909 (49%)	217 (4.42%)	1062 (21.63%)	-
Saurabh	911	69 (7%)	6 (8.70%)	0 (0.00%)	-	-	63 (91.30%)	-	-	842 (92%)	68 (8.08%)	6 (0.71%)	-
Munoz	314	294 (93%)	20 (6.80%)	-	-	-	-	-	274 (93.20%)	20 (6%)	1 (5.00%)	-	-
Cummins	1,781	586 (32%)	77 (13.14%)	-	-	-	-	-	509 (86.86%)	1195 (67%)	104 (8.70%)	-	-




Figure 5. Forest plot for risk of hospitalisation in current vs. never smokers. The prior from the previous review version (v10) was RR = 1.08.




Figure 6. Forest plot for risk of hospitalisation in former vs. never smokers. The prior from the previous version (v10) was RR = 1.18.

Disease severity by smoking status

85 studies reported disease severity in hospitalised patients stratified by smoking status (see Table 4). Severe (as opposed to non-severe) disease was broadly defined as requiring intensive treatment unit (ITU) admission, requiring oxygen as a hospital inpatient or in-hospital death. Meta-analyses were performed for 11 'fair' quality studies (see Figure 7 and 8). Current (RR = 1.26, 95% CrI = 0.92-1.73, $\tau = 0.32$) and former (RR = 1.8, 95% CrI = 1.27-2.55, $\tau = 0.46$) compared with never smokers were at increased risk of greater disease severity. However, while data for current smokers only were inconclusive, they favoured there being a small but important association. The probability of current and former smokers having increased risk of greater disease severity (RR ≥ 1.1) compared with never smokers was 80% and >99%,

respectively. Results were materially unchanged in two sensitivity analyses.

Table 4 COVID-19 disease severity by smoking status

Author	Population with severity	Non severe disease							Severe disease			
		N (%)	Current smoker (%)	Former smoker (%)	Current/former smoker (%)	Never smoker (%)	Never/unknown smoker (%)	Not stated (%)	N (%)	Current smoker (%)	Former smoker (%)	C
Guan, Ni	1,085	913 (84%)	108 (11.83%)	12 (1.31%)	-	793 (86.86%)	-	-	172 (15%)	29 (16.86%)	9 (5.23%)	-
Zhang, Dong	9	3 (33%)	0 (0.00%)	3 (100.00%)	-	0 (0.00%)	-	-	6 (66%)	2 (33.33%)	4 (66.67%)	-
Wan	9	8 (88%)	8 (100.00%)	0 (0.00%)	-	0 (0.00%)	-	-	1 (11%)	1 (100.00%)	0 (0.00%)	-
Huang, Wang	3	3 (100%)	3 (100.00%)	0 (0.00%)	-	0 (0.00%)	-	-	0 (0%)	0 (NaN%)	0 (NaN%)	-
Rentsch	285	168 (58%)	47 (27.98%)	53 (31.55%)	-	68 (40.48%)	-	-	117 (41%)	43 (36.75%)	36 (30.77%)	-
Hu	323	151 (46%)	-	-	12 (7.95%)	-	139 (92.05%)	-	172 (53%)	-	-	26
Wang, Pan	125	100 (80%)	-	-	9 (9.00%)	-	91 (91.00%)	-	25 (20%)	-	-	7
Kim	27	21 (77%)	3 (14.29%)	-	-	-	18 (85.71%)	-	6 (22%)	2 (33.33%)	0 (0.00%)	-
Shi, Yu	474	425 (89%)	-	-	34 (8.00%)	-	391 (92.00%)	-	49 (10%)	-	-	6
Liao, Feng	148	92 (62%)	-	-	5 (5.43%)	-	-	87 (94.57%)	56 (37%)	3 (5.36%)	-	-
Shi, Ren	134	88 (65%)	-	-	8 (9.09%)	-	-	80 (90.91%)	46 (34%)	-	-	6
Hadjadj	50	15 (30%)	1 (6.67%)	2 (13.33%)	-	12 (80.00%)	-	-	35 (70%)	0 (0.00%)	7 (20.00%)	-
Zheng, Xiong	73	43 (58%)	-	-	6 (13.95%)	37 (86.05%)	-	-	30 (41%)	-	-	2
de la Rica	48	26 (54%)	-	-	6 (23.08%)	-	-	20 (76.92%)	20 (41%)	-	-	4
Yin, Yang	106	47 (44%)	-	-	6 (12.77%)	-	-	41 (87.23%)	59 (55%)	-	-	12
Allenbach	147	100 (68%)	-	-	9 (9.00%)	-	-	91 (91.00%)	47 (31%)	-	-	0
Goyal	393	263 (66%)	14 (5.32%)	-	-	-	-	249 (94.68%)	130 (33%)	6 (4.62%)	-	-
Feng	454	333 (73%)	27 (8.11%)	-	-	-	-	306 (91.89%)	121 (26%)	17 (14.05%)	-	-
Yao	108	83 (76%)	1 (1.20%)	-	-	-	-	82 (98.80%)	25 (23%)	3 (12.00%)	-	-
Regina	200	163 (81%)	9 (5.52%)	-	-	-	-	154 (94.48%)	37 (18%)	0 (0.00%)	-	-
Feuth	28	21 (75%)	1 (4.76%)	7 (33.33%)	-	13 (61.90%)	-	-	7 (25%)	2 (28.57%)	1 (14.29%)	-

Mejia-Vilet	329	214 (65%)	-	-	13 (6.07%)	-	-	201 (93.93%)	115 (34%)	-	-	10
Chen, Jiang	135	54 (40%)	-	-	4 (7.41%)	-	-	50 (92.59%)	81 (60%)	-	-	9
Vaquero-Roncero	146	75 (51%)	-	-	4 (5.33%)	-	-	71 (94.67%)	71 (48%)	-	-	6
Kim, Garg	2,490	1692 (67%)	112 (6.62%)	395 (23.35%)	-	-	1185 (70.04%)	-	798 (32%)	38 (4.76%)	247 (30.95%)	-
Wu	174	92 (52%)	-	-	47 (51.09%)	-	45 (48.91%)	-	82 (47%)	11 (13.41%)	-	-
Chaudhry	40	34 (85%)	-	-	5 (14.71%)	-	-	29 (85.29%)	6 (15%)	-	-	1
Garibaldi	832	532 (63%)	25 (4.70%)	107 (20.11%)	-	-	-	400 (75.19%)	300 (36%)	21 (7.00%)	81 (27.00%)	-
Kuderer	928	686 (73%)	35 (5.10%)	210 (30.61%)	-	370 (53.94%)	-	29 (4.23%)	242 (26%)	8 (3.31%)	116 (47.93%)	-
Romao	14	14 (100%)	-	-	4 (28.57%)	-	-	10 (71.43%)	0 (0%)	-	-	-
Giannouchos	89,756	78050 (86%)	6322 (8.10%)	-	-	-	71728 (91.90%)	-	11706 (13%)	1089 (9.30%)	-	-
Cen	1,007	720 (71%)	-	-	70 (9.72%)	-	-	650 (90.28%)	287 (28%)	-	-	18
Maraschini	132	89 (67%)	-	11 (12.36%)	-	78 (87.64%)	-	-	43 (32%)	-	3 (6.98%)	-
Russell	156	128 (82%)	9 (7.03%)	31 (24.22%)	-	51 (39.84%)	-	37 (28.91%)	28 (17%)	2 (7.14%)	8 (28.57%)	-
Siso-Almirall	260	212 (81%)	-	-	60 (28.30%)	-	-	152 (71.70%)	48 (18%)	-	-	21
Gu	884	511 (57%)	30 (5.87%)	126 (24.66%)	-	355 (69.47%)	-	-	134 (15%)	3 (2.24%)	61 (45.52%)	-
Mendy	689	598 (86%)	-	-	133 (22.24%)	-	-	465 (77.76%)	91 (13%)	-	-	37
Jin, Gu	6	2 (33%)	-	-	0 (0.00%)	-	-	2 (100.00%)	4 (66%)	-	-	2
Senkal	611	446 (73%)	48 (10.76%)	-	-	-	-	398 (89.24%)	165 (27%)	21 (12.73%)	-	-
Patel	129	89 (68%)	26 (29.21%)	-	-	-	58 (65.17%)	5 (5.62%)	40 (31%)	22 (55.00%)	-	-
Maucourant	27	10 (37%)	1 (10.00%)	2 (20.00%)	-	2 (20.00%)	-	5 (50.00%)	17 (62%)	2 (11.76%)	5 (29.41%)	-
Xie	619	469 (75%)	-	-	32 (6.82%)	-	-	437 (93.18%)	150 (24%)	-	-	19
Fox	55	30 (54%)	1 (3.33%)	4 (13.33%)	-	17 (56.67%)	-	8 (26.67%)	25 (45%)	0 (0.00%)	2 (8.00%)	-
Zhang, Cao	240	162 (67%)	2 (1.23%)	6 (3.70%)	-	-	-	154 (95.06%)	78 (32%)	4 (5.13%)	4 (5.13%)	-
Kurashima	53	10 (18%)	-	-	3 (30.00%)	-	-	7 (70.00%)	43 (81%)	-	-	24
Zhan	75	NA (NA%)	-	-	-	-	-	-	75 (100%)	-	-	9

Omrani	858	806 (93%)	-	-	121 (15.01%)	-	-	685 (84.99%)	52 (6%)	-	-	9
Marcos	918	555 (60%)	38 (6.85%)	-	69 (12.43%)	-	-	448 (80.72%)	363 (39%)	18 (4.96%)	-	71
Hoertel, Sanchez, Rico	7,345	6014 (81%)	433 (7.20%)	-	-	-	-	5581 (92.80%)	1331 (18%)	190 (14.27%)	-	-
Qi	267	217 (81%)	22 (10.14%)	-	-	-	195 (89.86%)	-	50 (18%)	31 (62.00%)	-	-
Monteiro	112	84 (75%)	3 (3.57%)	14 (16.67%)	-	63 (75.00%)	-	4 (4.76%)	28 (25%)	4 (14.29%)	6 (21.43%)	-
Morshed	103	87 (84%)	28 (32.18%)	-	-	-	59 (67.82%)	-	16 (15%)	4 (25.00%)	-	-
Zhou, Sun	144	108 (75%)	11 (10.19%)	-	-	-	-	97 (89.81%)	36 (25%)	2 (5.56%)	-	-
Hippisley-Cox	-	NA (NA%)	-	-	-	-	-	-	1286 (NA%)	56 (4.35%)	427 (33.20%)	-
Zhao, Chen	641	398 (62%)	87 (21.86%)	-	-	-	-	311 (78.14%)	195 (30%)	52 (26.67%)	-	-
Qu	246	226 (91%)	90 (39.82%)	-	-	-	-	136 (60.18%)	20 (8%)	14 (70.00%)	-	-
Petrilli	2,729	1739 (63%)	97 (5.58%)	325 (18.69%)	-	1067 (61.36%)	-	250 (14.38%)	990 (36%)	44 (4.44%)	236 (23.84%)	-
Ren	432	314 (72%)	26 (8.28%)	-	-	288 (91.72%)	-	-	118 (27%)	17 (14.41%)	-	-
Yan	578	450 (77%)	31 (6.89%)	-	-	-	-	419 (93.11%)	128 (22%)	20 (15.62%)	-	-
Nicholson	1,042	550 (52%)	37 (6.73%)	106 (19.27%)	-	211 (38.36%)	-	196 (35.64%)	401 (38%)	41 (10.22%)	92 (22.94%)	-
Zhu	432	285 (65%)	46 (16.14%)	-	-	-	-	239 (83.86%)	147 (34%)	16 (10.88%)	-	-
Kalan	193	122 (63%)	9 (7.38%)	-	-	102 (83.61%)	-	11 (9.02%)	71 (36%)	5 (7.04%)	-	-
Burrell	204	85 (41%)	-	-	7 (8.24%)	-	75 (88.24%)	3 (3.53%)	119 (58%)	-	-	20
Chudasama	1,706	NA (NA%)	-	-	-	-	-	-	1706 (100%)	235 (13.77%)	699 (40.97%)	-
Lamure	89	NA (NA%)	-	-	-	-	-	-	25 (28%)	1 (4.00%)	5 (20.00%)	-
Zhou, He, Yang	1,087	990 (91%)	-	-	849 (85.76%)	141 (14.24%)	-	-	97 (8%)	-	-	75
Zhou, Qin	51	NA (NA%)	-	-	-	-	-	-	51 (100%)	-	-	40
Zhan, Liu	405	257 (63%)	-	-	21 (8.17%)	236 (91.83%)	-	-	148 (36%)	-	-	25
Li, Long, Zhang	954	838 (87%)	-	-	34 (4.06%)	804 (95.94%)	-	-	116 (12%)	-	-	22
Jakob	2,155	1400 (64%)	-	-	92 (6.57%)	99 (7.07%)	-	669 (47.79%)	755 (35%)	51 (6.75%)	58 (7.68%)	-
Aksu	123	34 (27%)	3 (8.82%)	-	-	-	31 (91.18%)	-	89 (72%)	11 (12.36%)	-	-

Adrish	1,173	162 (13%)	-	-	36 (22.22%)	-	126 (77.78%)	-	1011 (86%)	-	-	30
Hoertel, Sanchez, Vernet	12,210	11018 (90%)	921 (8.36%)	-	-	-	-	10097 (91.64%)	1192 (9%)	181 (15.18%)	-	-
Vila- Corcoles, Satue-Gracia	282	218 (77%)	-	-	21 (9.63%)	-	-	197 (90.37%)	64 (22%)	-	-	4
Boyd	38	0 (0%)	-	-	-	-	-	-	38 (100%)	-	-	2
Caliskan	565	474 (83%)	96 (20.25%)	45 (9.49%)	-	333 (70.25%)	-	-	91 (16%)	22 (24.18%)	34 (37.36%)	-
Ebrahimian	226	134 (59%)	-	-	12 (8.96%)	-	122 (91.04%)	-	92 (40%)	-	-	12
Ho, Narasimhan	4,909	3859 (78%)	169 (4.38%)	833 (21.59%)	-	-	2857 (74.03%)	-	1050 (21%)	48 (4.57%)	229 (21.81%)	-
Quan	2,038	1452 (71%)	-	549 (37.81%)	-	-	-	903 (62.19%)	586 (28%)	-	-	26
Saurabh	911	783 (85%)	65 (8.30%)	3 (0.38%)	-	715 (91.32%)	-	-	128 (14%)	9 (7.03%)	3 (2.34%)	-
Chousein	114	94 (82%)	17 (18.09%)	17 (18.09%)	-	60 (63.83%)	-	-	20 (17%)	2 (10.00%)	6 (30.00%)	-
Tavakol	206	182 (88%)	-	-	24 (13.19%)	-	158 (86.81%)	-	24 (11%)	-	-	2
Yao, Hasegawa	101	78 (77%)	-	-	17 (21.79%)	-	71 (91.03%)	-	23 (22%)	-	-	12
Cummins	1,195	1043 (87%)	96 (9.20%)	-	-	-	-	947 (90.80%)	152 (12%)	9 (5.92%)	-	-
Nuno	4,730	3536 (74%)	333 (9.42%)	-	-	-	-	3203 (90.58%)	1194 (25%)	147 (12.31%)	-	-




Figure 7. Forest plot for the risk of severe disease in current vs. never smokers. The prior from the previous review version (v10) was RR = 1.26.




Figure 8. Forest plot for the risk of severe disease in former vs. never smokers. The prior from the previous review version (v10) was RR = 1.52.

Mortality by smoking status

89 studies reported mortality from COVID-19 by smoking status (see Table 5), with 19 'fair' quality studies included in meta-analyses (see Figure 9 and 10). Current (RR = 1.12, 95% CrI = 0.84-1.47, $\tau = 0.42$) and former (RR = 1.56, 95% CrI = 1.23-2, $\tau = 0.43$) compared with never smokers were at increased risk of in-hospital mortality from COVID-19. However, data for current smokers were inconclusive and favoured there being no important association. The probability of current and former smokers being at greater risk of in-hospital mortality (RR ≥ 1.1) compared with never smokers was 60% and >99%, respectively. Results were materially unchanged in two sensitivity analyses.

Table 5 COVID-19 attributed mortality by smoking status

Author	Population with mortality	Recovered							Died		
		N (%)	Current smoker (%)	Former smoker (%)	Current/former smoker (%)	Never smoker (%)	Never/unknown smoker (%)	Not stated (%)	N (%)	Current smoker (%)	Former smoker (%)
Chen	274	161 (58%)	5 (3.11%)	5 (3.11%)	-	-	-	151 (93.79%)	113 (41%)	7 (6.19%)	2 (1.77%)
Zhou, Yu	191	137 (71%)	6 (4.38%)	-	-	-	-	131 (95.62%)	54 (28%)	5 (9.26%)	-
Yang, Yu	52	20 (38%)	2 (10.00%)	-	-	-	18 (90.00%)	-	32 (61%)	-	-
Borobia	2,226	1766 (79%)	113 (6.40%)	-	-	-	-	1653 (93.60%)	460 (20%)	44 (9.57%)	-
Giacomelli	233	185 (79%)	-	-	53 (28.65%)	132 (71.35%)	-	-	48 (20%)	-	-
Yao	108	96 (88%)	1 (1.04%)	-	-	-	-	95 (98.96%)	12 (11%)	3 (25.00%)	-
Carillo-Vega	9,946	8983 (90%)	795 (8.85%)	-	-	-	-	8188 (91.15%)	963 (9%)	99 (10.28%)	-
Ge	51	39 (76%)	6 (15.38%)	-	-	-	-	33 (84.62%)	12 (23%)	1 (8.33%)	-
Chen, Jiang	135	NA (NA%)	-	-	-	-	-	-	31 (22%)	-	-
Heili-Frades	4,712	4086 (86%)	210 (5.14%)	659 (16.13%)	-	-	3217 (78.73%)	-	626 (13%)	23 (3.67%)	161 (25.72%)
Kim, Garg	2,490	2070 (83%)	128 (6.18%)	481 (23.24%)	-	-	1461 (70.58%)	-	420 (16%)	22 (5.24%)	161 (38.33%)
Al-Hindawi	31	15 (48%)	0 (0.00%)	10 (66.67%)	-	5 (33.33%)	-	-	16 (51%)	1 (6.25%)	12 (75.00%)
Louis	22	16 (72%)	-	-	7 (43.75%)	-	-	9 (56.25%)	6 (27%)	-	-
Soto-Mota	400	200 (50%)	-	-	23 (11.50%)	-	-	177 (88.50%)	200 (50%)	-	-
Garibaldi	747	634 (84%)	36 (5.68%)	129 (20.35%)	-	-	-	469 (73.97%)	113 (15%)	6 (5.31%)	36 (31.86%)
Docherty	13,364	8199 (61%)	370 (4.51%)	1832 (22.34%)	-	4179 (50.97%)	-	1818 (22.17%)	5165 (38%)	214 (4.14%)	1350 (26.14%)
Kuderer	928	807 (86%)	38 (4.71%)	262 (32.47%)	-	425 (52.66%)	-	31 (3.84%)	121 (13%)	5 (4.13%)	64 (52.89%)
Ramlall	11,116	10498 (94%)	-	-	2771 (26.40%)	7727 (73.60%)	-	-	618 (5%)	-	-
Wang, Oekelen	57	43 (75%)	-	-	14 (32.56%)	-	-	29 (67.44%)	14 (24%)	-	-
Martinez-Portilla	224	217 (96%)	-	-	7 (3.23%)	-	-	210 (96.77%)	7 (3%)	-	-
Cen	1,007	964 (95%)	-	-	87 (9.02%)	-	-	877 (90.98%)	43 (4%)	-	-
Klang	3,406	2270 (66%)	-	-	492 (21.67%)	-	-	1778 (78.33%)	1136 (33%)	-	-

Wang, Zhong	5,510	4874 (88%)	247 (5.07%)	1083 (22.22%)	-	3544 (72.71%)	-	-	636 (11%)	28 (4.40%)	214 (33.65%)
Miyara	338	211 (62%)	13 (6.16%)	58 (27.49%)	-	141 (66.82%)	-	-	46 (13%)	1 (2.17%)	23 (50.00%)
Rajter	255	209 (81%)	-	-	28 (13.40%)	181 (86.60%)	-	-	53 (20%)	-	-
Zeng	1,031	866 (84%)	-	-	69 (7.97%)	-	-	797 (92.03%)	165 (16%)	-	-
Chen, Yu	1,859	1651 (88%)	32 (1.94%)	54 (3.27%)	-	1565 (94.79%)	-	-	208 (11%)	13 (6.25%)	12 (5.77%)
Garassino	190	124 (65%)	-	-	92 (74.19%)	32 (25.81%)	-	-	66 (34%)	-	61 (92.42%)
Gu	884	864 (97%)	40 (4.63%)	250 (28.94%)	-	219 (25.35%)	-	-	20 (2%)	0 (0.00%)	14 (70.00%)
Zhou, He	-	NA (NA%)	-	-	-	-	-	-	NA (NA%)	-	-
Sigel	88	70 (79%)	-	-	37 (52.86%)	-	-	33 (47.14%)	18 (20%)	-	-
Nguyen	356	308 (86%)	-	-	91 (29.55%)	-	-	217 (70.45%)	45 (12%)	-	-
de Souza	8,443	7826 (92%)	-	-	95 (1.21%)	-	7571 (96.74%)	160 (2.04%)	617 (7%)	-	-
Mendy	532	663 (124%)	-	-	160 (24.13%)	-	-	503 (75.87%)	26 (4%)	-	-
Shi, Resurreccion	256	210 (82%)	-	-	128 (60.95%)	-	-	82 (39.05%)	46 (17%)	-	-
Xie	619	591 (95%)	-	-	43 (7.28%)	-	-	548 (92.72%)	28 (4%)	-	-
Fox	54	35 (64%)	1 (2.86%)	4 (11.43%)	-	18 (51.43%)	-	12 (34.29%)	19 (35%)	0 (0.00%)	2 (10.53%)
Zhang, Cao	289	240 (83%)	10 (4.17%)	6 (2.50%)	-	-	-	224 (93.33%)	49 (16%)	4 (8.16%)	8 (16.33%)
Gupta	496	255 (51%)	-	-	15 (5.88%)	-	80 (31.37%)	160 (62.75%)	241 (48%)	-	-
Soares	1,152	696 (60%)	38 (5.46%)	-	-	-	658 (94.54%)	-	456 (39%)	39 (8.55%)	-
Thompson	470	301 (64%)	39 (12.96%)	79 (26.25%)	-	183 (60.80%)	-	-	169 (35%)	27 (15.98%)	49 (28.99%)
Bernaola	1,645	1382 (84%)	35 (2.53%)	146 (10.56%)	-	1201 (86.90%)	-	-	263 (15%)	6 (2.28%)	33 (12.55%)
Islam	654	631 (96%)	103 (16.32%)	-	-	-	-	507 (80.35%)	23 (3%)	3 (13.04%)	-
Philipose	466	267 (57%)	19 (7.12%)	204 (76.40%)	-	44 (16.48%)	-	-	199 (42%)	9 (4.52%)	137 (68.84%)
Fillmore	1,794	1566 (87%)	408 (26.05%)	758 (48.40%)	-	279 (17.82%)	-	98 (6.26%)	228 (12%)	44 (19.30%)	141 (61.84%)
Pan	3,536	3302 (93%)	-	-	862 (26.11%)	-	-	2440 (73.89%)	234 (6%)	-	-
Zhao, Chen	474	398 (83%)	87 (21.86%)	-	-	-	-	311 (78.14%)	82 (17%)	36 (43.90%)	-
Holman	10,989	NA	-	-	-	-	-	-	10989	609	4684

Author	Total	Category 1	Category 2	Category 3	Category 4	Category 5	Category 6	Category 7	Category 8	Category 9	Category 10
Chand	300	143 (47%)	23 (16.08%)	-	-	-	-	120 (83.92%)	157 (52%)	44 (28.03%)	-
Oliveira	131	105 (80%)	-	-	16 (15.24%)	-	83 (79.05%)	6 (5.71%)	26 (19%)	-	-
Vilar-Garcia	328,892	316605 (96%)	71215 (22.49%)	-	-	-	-	245390 (77.51%)	12287 (3%)	3103 (25.25%)	-
Ibarra-Nava	416,546	370038 (88%)	27001 (7.30%)	-	-	-	-	343037 (92.70%)	46508 (11%)	3817 (8.21%)	-
Rubio-Rivas	186	147 (79%)	7 (4.76%)	32 (21.77%)	-	-	108 (73.47%)	-	39 (20%)	1 (2.56%)	6 (15.38%)
Ren	432	289 (66%)	25 (8.65%)	-	-	-	264 (91.35%)	-	143 (33%)	18 (12.59%)	-
Ullah	212	158 (74%)	22 (13.92%)	67 (42.41%)	-	-	63 (39.87%)	-	6 (3.80%)	54 (25%)	2 (3.70%)
Dashti	3,401	2892 (85%)	190 (6.57%)	689 (23.82%)	-	-	1756 (60.72%)	-	257 (8.89%)	509 (14%)	20 (3.93%)
Nicholson	1,040	829 (79%)	70 (8.44%)	163 (19.66%)	-	-	320 (38.60%)	-	276 (33.29%)	211 (20%)	16 (7.58%)
Kalan	193	188 (97%)	14 (7.45%)	-	-	-	162 (86.17%)	-	12 (6.38%)	5 (2%)	0 (0.00%)
Incerti	13,658	11495 (84%)	785 (6.83%)	2450 (21.31%)	-	-	5450 (47.41%)	2810 (24.45%)	-	2163 (15%)	81 (3.74%)
Ioannou	10,131	9033 (89%)	1054 (11.67%)	3549 (39.29%)	-	-	3339 (36.96%)	-	1091 (12.08%)	1098 (10%)	81 (7.38%)
Lamure	89	59 (66%)	4 (6.78%)	16 (27.12%)	-	-	31 (52.54%)	-	8 (13.56%)	30 (33%)	1 (3.33%)
Yadaw	5,051	4635 (91%)	162 (3.50%)	709 (15.30%)	-	-	2394 (51.65%)	-	1370 (29.56%)	416 (8%)	17 (4.09%)
Zinellu	105	77 (73%)	24 (31.17%)	8 (10.39%)	-	-	45 (58.44%)	-	-	28 (26%)	8 (28.57%)
Zhang, Li	399	NA (NA%)	-	-	-	-	-	-	399 (100%)	60 (15.04%)	186 (46.62%)
Wang, Shu	59	18 (30%)	-	-	0 (0.00%)	-	-	18 (100.00%)	41 (69%)	-	-
Wang, Zheutlin	2,448	1706 (69%)	57 (3.34%)	315 (18.46%)	-	-	954 (55.92%)	-	380 (22.27%)	742 (30%)	25 (3.37%)
Torres-Macho	1,968	1643 (83%)	-	-	335 (20.39%)	-	-	1308 (79.61%)	325 (16%)	-	-
Raines	440	408 (92%)	-	-	222 (54.41%)	186 (45.59%)	-	-	32 (7%)	-	-
Parra-Bracamonte	331,298	292988 (88%)	-	-	21269 (7.26%)	-	-	271719 (92.74%)	38310 (11%)	-	-
Li, Long, Zhang	954	876 (91%)	-	-	48 (5.48%)	828 (94.52%)	-	-	78 (8%)	-	-
Bellan	407	285 (70%)	30 (10.53%)	-	-	-	191 (67.02%)	64 (22.46%)	122 (29%)	24 (19.67%)	-
Alharthy	352	239 (67%)	109 (45.61%)	-	-	-	130 (54.39%)	-	113 (32%)	65 (57.52%)	-
Adrish	1,173	783 (66%)	-	-	205 (26.18%)	578 (73.82%)	-	-	390 (33%)	-	-

Hoertel, Sanchez, Vernet	12,210	12210 (100%)	898 (7.35%)	-	-	-	-	9904 (81.11%)	1408 (11%)	204 (14.49%)	-
Sourij	238	180 (75%)	4 (2.22%)	26 (14.44%)	-	-	150 (83.33%)	-	58 (24%)	0 (0.00%)	12 (20.69%)
Simons	446	318 (71%)	30 (9.43%)	109 (34.28%)	-	-	179 (56.29%)	-	128 (28%)	12 (9.38%)	63 (49.22%)
Bisso	168	122 (72%)	13 (10.66%)	-	-	-	-	109 (89.34%)	46 (27%)	5 (10.87%)	-
Caliskan	565	490 (86%)	97 (19.80%)	48 (9.80%)	-	-	345 (70.41%)	-	75 (13%)	21 (28.00%)	31 (41.33%)
Ebrahimian	226	128 (56%)	-	-	9 (7.03%)	-	-	119 (92.97%)	98 (43%)	-	-
Estiri	16,709	15879 (95%)	-	-	1304 (8.21%)	-	-	14575 (91.79%)	830 (4%)	-	-
Ferrari	198	165 (83%)	-	-	27 (16.36%)	-	138 (83.64%)	-	33 (16%)	-	-
Ho, Narasimhan	4,909	3553 (72%)	163 (4.59%)	743 (20.91%)	-	-	2647 (74.50%)	-	1356 (27%)	54 (3.98%)	319 (23.53%)
Mansour	111	83 (74%)	5 (6.02%)	-	-	-	-	78 (93.98%)	28 (25%)	6 (21.43%)	-
Park	2,269	2105 (92%)	92 (4.37%)	-	-	-	-	2013 (95.63%)	164 (7%)	2 (1.22%)	-
Quan	2,038	1587 (77%)	-	-	589 (37.11%)	-	-	998 (62.89%)	442 (21%)	-	-
Saurabh	911	870 (95%)	70 (8.05%)	4 (0.46%)	-	-	796 (91.49%)	-	41 (4%)	4 (9.76%)	2 (4.88%)
Strangfeld	3,729	3339 (89%)	-	-	664 (19.89%)	-	2190 (65.59%)	-	485 (14.53%)	390 (10%)	-
Nezhadmoghadam	13,367	7757 (58%)	686 (8.84%)	-	-	-	-	7071 (91.16%)	5610 (41%)	565 (10.07%)	-
Nuno	4,730	4401 (93%)	427 (9.70%)	-	-	-	-	3974 (90.30%)	329 (6%)	53 (16.11%)	-




Figure 9. Forest plot for the risk of mortality in current vs. never smokers. The prior from the previous review version (v10) was RR = 1.05.




Figure 10. Forest plot for the risk of mortality in former smokers vs. never smokers. The prior from the previous version (v10) was RR = 1.40.

Discussion

This living rapid review found uncertainty in the majority of 405 studies arising from the recording of smoking status. Notwithstanding this uncertainty, compared with overall adult national prevalence estimates, recorded current smoking rates in most studies were lower than expected. In a subset of good and fair quality studies ($n = 30$), current but not former smokers had a reduced risk of testing positive for SARS-CoV-2 but current smokers appeared somewhat more likely to present for testing and/or receive a test. Data for current smokers on the risk of hospitalisation, disease severity and mortality were inconclusive, and favoured there being no important associations with hospitalisation and mortality and a small but important increase in the risk of severe disease. Former smokers were at increased risk of hospitalisation,

disease severity and mortality compared with never smokers.

Issues complicating interpretation

Interpretation of results from studies conducted during the first phase of the SARS-CoV-2 pandemic is complicated by several factors (see Figure 11):

1. *Exposure to SARS-CoV-2*

- i) Exposure to the SARS-CoV-2 virus is heterogeneous with different subgroups at heightened risk of infection at different stages of the pandemic, at least partly due to differential contact matrices by age, sex and socioeconomic position (CMMID COVID-19 working group et al., 2020), which are associated with smoking status.
- ii) The probability of viral exposure depends largely on local prevalence, which varies over time. This likely introduces bias in studies assessing the rate of infection by smoking status conducted in the early phase of the pandemic.

2. *Infection with SARS-CoV-2*

- i) Infection following viral exposure depends on individual differences in, for example, genetic susceptibility or immunocompetence, which are poorly understood at present and may be confounded with smoking. For example, the household secondary attack rate for COVID-19 is estimated at 17% (Fung et al., 2020).
- ii) Heated and humidified air may act to disrupt the ability of the virus to persist in the airway mucosa of smokers. There is some evidence that transient localised hyperthermia can inhibit replication of rhinoviruses, a non-enveloped virus that causes the common cold (Conti et al., 1999). However, as SARS-CoV-2 is an enveloped virus (Schoeman and Fielding, 2019), it is unclear whether a similar protective effect against viral replication or invasion by heated and humidified air may occur.

3. *Symptomatic COVID-19*

- i) An estimated 20% (95% CI = 17-25%) of COVID-19 cases are asymptomatic (Buitrago-Garcia et al., 2020), with some evidence suggesting younger people are more likely to be asymptomatic (Kronbichler et al., 2020). Testing is hence likely limited in some subgroups, with the potential for these groups to include an overrepresentation of current smokers.
- ii) On the other hand, current and former smokers may be more likely to meet local criteria for community testing due to increased prevalence of symptoms consistent with SARS-CoV-2 infection, such as cough, increased sputum production or altered sense of smell or taste (Hopkinson et al., 2020). Evidence from a small number of studies indicates that current smokers may be more likely to present for testing, hence increasing the denominator in comparisons with never smokers and potentially inflating the rate of negative tests in current smokers. Infection positivity rates estimated among random samples are more informative. We identified one population study conducted in Hungary reporting on seroprevalence and smoking status (Merkely et al., 2020); however, the response rate was only 58.8% and the current smoking rate was 10 percentage points below national prevalence estimates, which raises some doubt about representativeness of the final sample. Similarly, two further representative population surveys (Carrat et al., 2020; Richard et al., 2020) reported a current smoking rate of more than 10 percentage points below national prevalence (12% vs. 25% and 15% vs. 27% daily smoking prevalence, respectively) (Anderl, 2019; FSO,

2020).

4. *Testing positive for SARS-CoV-2*

i) Smokers with COVID-19 may be less likely to receive a SARS-CoV-2 test or present to hospital due to lack of access to healthcare and may be more likely to die in the community from sudden complications (i.e. self-selection bias) and thus not be recorded (Brown, 2020).

ii) Diagnostic criteria for SARS-CoV-2 infection and COVID-19 have changed during the course of the pandemic (Organisation, n.d.). It was not possible to extract details on the specific RT-PCR or antibody-based techniques or platforms used across the included studies due to reporting gaps. Different platforms have varying sensitivity and specificity to detect SARS-CoV-2 infection. In addition, testing for acute infection requires swabbing of the mucosal epithelium, which may be disrupted in current smokers, potentially altering the sensitivity of assays (Lusignan et al., 2020).

5. *Hospitalisation with COVID-19*

i) Reasons for hospitalisation vary by country and time in the pandemic. For example, early cases may have been hospitalised for isolation and quarantine reasons and not due to medical necessity. It is plausible this may have skewed early data towards less severe cases. In addition, the observed association between former smoking and greater disease severity may be explained by collider bias (Griffith et al., 2020), where conditioning on a collider (e.g. testing or hospitalisation) by design or analysis may introduce a spurious association between current or former smoking (a potential cause of testing or hospitalisation) and SARS-CoV-2 infection/adverse outcomes from COVID-19 (potentially exacerbated by smoking) (Murray, 2020).

ii) The majority of included studies relied on EHRs as the source of information on smoking status. Research shows large discrepancies between EHRs and actual behaviour (Polubriaginof et al., 2018). Known failings of EHRs include implausible longitudinal changes, such as former smokers being recorded as never smokers at subsequent hospital visits (Polubriaginof et al., 2018). Misreporting on the part of the patient (perhaps due to perceived stigma) has also been observed, with biochemical measures showing higher rates of smoking compared with self-report in hospitalised patients in the US (Benowitz et al., 2009). It is hence possible that under-reporting of current and former smoking status in hospitals occurred across the included studies.

iii) The majority of included studies were conducted in hospital settings. It is plausible that a non-trivial proportion of patients were infected with SARS-CoV-2 while being an inpatient for a different medical reason. If so, this may have biased the hospitalised populations towards older and more frail groups, who are less likely to be smokers (Mangera et al., 2017).

iv) Individuals with severe COVID-19 symptoms may have stopped smoking immediately before admission to hospital and may therefore not have been recorded as current smokers (i.e. reverse causality).

6. *COVID-19 disease severity and death*

i) Given lack of knowledge of the disease progression and long-term outcomes of COVID-19, it is unclear whether studies conducted thus far in the pandemic have monitored patients for a sufficient time period to report complete survival outcomes or whether they are subject to early censoring. Adding to this, COVID-19 related mortality has been differentially defined across countries and epidemic phases. For example, in some UK reporting, death within 28 days

of a COVID-19 diagnosis is required for attributing the cause of death to the virus. However, according to the UK Office for National Statistics, COVID-19 deaths are recorded only if this was stated on the death certificate.

ii) If there is a protective effect of nicotine on COVID-19 disease outcomes, abrupt nicotine withdrawal upon hospitalisation may lead to worse disease outcomes including death (Farsalinos, Niaura, et al., 2020).

iii) During periods of heightened demand of limited healthcare resources, current and former smokers with extensive comorbidities may have reduced priority for intensive care admission, thus leading to higher in-hospital mortality.

iv) COVID-19 outcomes are currently limited to in-hospital death or survival to discharge. This binary outcome does not capture potential long-term morbidity attributed to COVID-19, such as stroke, amputation or acute cardiac events, which may be moderated by smoking status.




Figure 11. A schematic of some of the interpretation issues for the association of smoking status and COVID-19 infection, hospitalisation, disease severity and mortality. Numbers refer to the issues listed in-text in the above section. Issues presented on the right-hand side relate directly to smoking status.

Limitations

This living rapid evidence review was limited by having a single reviewer extracting data with a second independently verifying the data extracted to minimise errors, restricting the search to one electronic database and one pre-print server and by not including at least three large population surveys due to their reliance on self-reported suspected or confirmed SARS-CoV-2 infection (which means they do not meet our eligibility criteria) (Bowyer et al., 2020; Hopkinson et al., 2020; Jackson et al., 2020). We also did not include a large, UK-based, representative seroprevalence study (Ward et al., 2020) in our meta-analyses as the odds of testing positive in former smokers was not reported. However, the odds of infection for current smokers (OR = 0.64, 95% CI = 0.58-0.71) was in concordance with the pooled estimate in our meta-analysis. Population surveys – particularly with linked data on confirmed infection or antibodies – will be included in future review versions to help mitigate some of the limitations of healthcare based observational studies. The comparisons of current and former smoking prevalence in the included studies with national prevalence estimates did not adjust observed prevalence for the demographic profile of those tested/admitted to hospital. Other reviews focused on this comparison have applied adjustments for sex and age, and continue to find lower than expected prevalence – notwithstanding the issues complicating interpretation described above (Farsalinos, Barbouni, et al., 2020).

Implications for research, policy and practice

Further scientific research is needed to resolve the mixed findings summarised in our review. First, clinical trials of the posited therapeutic effect of nicotine could have important implications both for smokers and for improved understanding of how the SARS-CoV-2 virus causes disease in humans. Such trials should focus on medicinal nicotine (as smoked tobacco is a dirty delivery mechanism that could mask beneficial effects) and potentially differentiate between different modes of delivery (i.e. inhaled vs. ingested) since this can affect pharmacokinetics (Shahab et al., 2013) and potential therapeutic effects. A second research priority would be a large, representative (randomly sampled) population survey with a validated assessment of smoking status which distinguishes between recent and long-term ex-smokers – ideally biochemically verified – and assesses seroprevalence and links to health records.

In the meantime, public-facing messages about the possible protective effect of smoking or nicotine are premature. In our view, until there is further research, the quality of the evidence does not justify the huge risk associated with a message likely to reach millions of people that a lethal activity, such as smoking, may protect against COVID-19. It continues to be appropriate to recommend smoking cessation and emphasise the role of alternative nicotine products to support smokers to stop as part of public health efforts during COVID-19. At the very least, smoking cessation reduces acute risks from cardiovascular disease and could reduce demands on the healthcare system (Stead et al., 2013). GPs and other healthcare providers can play a crucial role – brief, high-quality and free online training is available at [National Centre for Smoking Cessation and Training](#).

Conclusion

Across 405 studies, recorded current but not past smoking prevalence was generally lower than national prevalence

estimates. Current smokers were at reduced risk of testing positive for SARS-CoV-2 and former smokers were at increased risk of hospitalisation, disease severity and mortality compared with never smokers.

Acknowledgements

An original short review for the Royal College of Physicians was converted to an extended living review after a request by Martin Dockrell, Tobacco Control Lead, Public Health England. All scientific decisions were made by the authors independently of funders and external organisations. The authors would like to thank Rosemary Koper for her assistance in running the electronic searches and data extraction up until v7, and all authors who responded to requests for additional data.

Declaration of conflicts of interest

DS and OP have no conflicts of interest to declare. LS has received a research grant and honoraria for a talk and travel expenses from manufacturers of smoking cessation medications (Pfizer and Johnson & Johnson). JB has received unrestricted research funding to study smoking cessation from companies who manufacture smoking cessation medications. All authors declare no financial links with tobacco companies or e-cigarette manufacturers or their representatives.

Funding statement

DS is supported by a PhD studentship from the UK Biotechnology and Biological Sciences Research Council [BB/M009513/1]. OP receives salary support from Cancer Research UK (C1417/A22962). JB, LS, & OP are members of SPECTRUM, a UK Prevention Research Partnership Consortium (MR/S037519/1). UKPRP is an initiative funded by the UK Research and Innovation Councils, the Department of Health and Social Care (England) and the UK devolved administrations, and leading health research charities.

Review versions

The most up-to-date review version is available [here](#)

Previous review versions

- [Version 1](#)
- [Version 2](#)
- [Version 3](#)
- [Version 4](#)

- [Version 5](#)
- [Version 6](#)
- [Version 7](#)
- [Version 8](#)
- [Version 9](#)
- [Version 10](#)

Data availability

All data contributing to the current and future review versions are available [here](#)

All code required to reproduce the current and future analyses are available [here](#)

References

Abadom TR, Smith AD, Tempia S, et al. (2016) Risk factors associated with hospitalisation for influenza-associated severe acute respiratory illness in South Africa: A case-population study. *Vaccine* 34(46): 5649–5655. DOI:

[10.1016/j.vaccine.2016.09.011](https://doi.org/10.1016/j.vaccine.2016.09.011).

Abajo FJ de, Rodríguez-Miguel A, Rodríguez-Martín S, et al. (2021) IN-HOSPITAL CONTINUATION WITH ANGIOTENSIN RECEPTOR BLOCKERS IS ASSOCIATED WITH A LOWER MORTALITY RATE THAN CONTINUATION WITH ANGIOTENSIN CONVERTING ENZYME INHIBITORS IN COVID-19 PATIENTS: A RETROSPECTIVE COHORT STUDY. DOI: [10.1101/2021.02.01.21250853](https://doi.org/10.1101/2021.02.01.21250853).

Abolghasemi S, Mardani M, Sali S, et al. (2020) COVID-19 and kidney transplant recipients. *Transplant Infectious Disease* 22(6). DOI: [10.1111/tid.13413](https://doi.org/10.1111/tid.13413).

Ader F, Peiffer-Smadja N, Poissy J, et al. (2021) Antiviral drugs in hospitalized patients with COVID-19 - the DisCoVeRy trial. DOI: [10.1101/2021.01.08.20248149](https://doi.org/10.1101/2021.01.08.20248149).

Adrish M, Chilimuri S, Mantri N, et al. (2020) Association of smoking status with outcomes in hospitalised patients with COVID-19. *BMJ Open Respiratory Research* 7(1): e000716. DOI: [10.1136/bmjresp-2020-000716](https://doi.org/10.1136/bmjresp-2020-000716).

Aksu K, Naziroğlu T and Özkan P (2020) Factors determining COVID-19 pneumonia severity in a country with routine BCG vaccination. *Clinical & Experimental Immunology* 202(2): 220–225. DOI: [10.1111/cei.13507](https://doi.org/10.1111/cei.13507).

Alguwaihes AM, Al-Sofiani ME, Megdad M, et al. (2020) Diabetes and Covid-19 among hospitalized patients in Saudi Arabia: A single-centre retrospective study. *Cardiovascular Diabetology* 19(1): 205. DOI: [10.1186/s12933-020-01184-4](https://doi.org/10.1186/s12933-020-01184-4).

Alharthy A, Aletreby W, Faqihi F, et al. (2020) Clinical Characteristics and Predictors of 28-Day Mortality in 352 Critically Ill Patients with COVID-19: A Retrospective Study: *Journal of Epidemiology and Global Health*. DOI: [10.2991/jegh.k.200928.001](https://doi.org/10.2991/jegh.k.200928.001).

Al-Hindawi A, Sokhi J, Cuddihy J, et al. (2020) COVID-19 in London, a Case Series Demonstrating Late Improvement in Survivors. *medRxiv*: 2020.05.16.20103853. DOI: [10.1101/2020.05.16.20103853](https://doi.org/10.1101/2020.05.16.20103853).

Alizadehsani R, Sani ZA, Behjati M, et al. (2020) Risk Factors Prediction, Clinical Outcomes, and Mortality of COVID-19

Patients. *medRxiv*: 2020.07.07.20148569. DOI: [10.1101/2020.07.07.20148569](https://doi.org/10.1101/2020.07.07.20148569).

Alkurt G, Murt A, Aydin Z, et al. (2020) Seroprevalence of Coronavirus Disease 2019 (COVID-19) Among Health Care Workers from Three Pandemic Hospitals of Turkey. *medRxiv*: 2020.08.19.20178095. DOI: [10.1101/2020.08.19.20178095](https://doi.org/10.1101/2020.08.19.20178095).

Allenbach Y, Saadoun D, Maalouf G, et al. (2020) Development of a multivariate prediction model of intensive care unit transfer or death: A French prospective cohort study of hospitalized COVID-19 patients. *PLOS ONE* Moreira J (ed.) 15(10): e0240711. DOI: [10.1371/journal.pone.0240711](https://doi.org/10.1371/journal.pone.0240711).

Almazeedi S, Al-Youha S, Jamal MH, et al. (2020) Characteristics, risk factors and outcomes among the first consecutive 1096 patients diagnosed with COVID-19 in Kuwait. *EClinicalMedicine* 24: 100448. DOI: [10.1016/j.eclinm.2020.100448](https://doi.org/10.1016/j.eclinm.2020.100448).

Almirall J, González CA, Balanzó X, et al. (1999) Proportion of community-acquired pneumonia cases attributable to tobacco smoking. *Chest* 116: 375–9. DOI: [10.1378/chest.116.2.375](https://doi.org/10.1378/chest.116.2.375).

Alqahtani JS, Oyelade T, Aldhahir AM, et al. (2020) Prevalence, Severity and Mortality associated with COPD and Smoking in patients with COVID-19: A Rapid Systematic Review and Meta-Analysis. *medRxiv*: 2020.03.25.20043745. DOI: [10.1101/2020.03.25.20043745](https://doi.org/10.1101/2020.03.25.20043745).

Alshami AA, Alattas RA, Anan HF, et al. (2020) Silent Disease and Loss of Taste and Smell are Common Manifestations of SARS-COV-2 Infection in a Quarantine Facility: First report from Saudi Arabia. *medRxiv*: 2020.05.13.20100222. DOI: [10.1101/2020.05.13.20100222](https://doi.org/10.1101/2020.05.13.20100222).

Altamimi H, Alahmad Y, Khazal F, et al. (2020) The Outcome of COVID-19 Patients with Acute Myocardial Infarction. *medRxiv*: 2020.07.21.20156349. DOI: [10.1101/2020.07.21.20156349](https://doi.org/10.1101/2020.07.21.20156349).

Altibi AM, Bhargava P, Liaqat H, et al. (2020) Comparative Clinical Outcomes and Mortality in Prisoner and Non-Prisoner Populations Hospitalized with COVID-19: A Cohort from Michigan. *medRxiv*: 2020.08.08.20170787. DOI: [10.1101/2020.08.08.20170787](https://doi.org/10.1101/2020.08.08.20170787).

Andler R (2019) BAISSÉ DE LA PRÉVALENCE DU TABAGISME QUOTIDIEN PARMIS LES ADULTES : RÉSULTATS DU BAROMÈTRE DE SANTÉ PUBLIQUE FRANCE 2018 / REDUCTION OF DAILY SMOKING RATE AMONG ADULTS: RESULTS FROM THE 2018 SANTÉ PUBLIQUE FRANCE HEALTH BAROMETER.: 7.

Antonio-Villa NE, Bello-Chavolla OY, Vargas-Vazquez A, et al. (2020) Health-care workers with COVID-19 living in Mexico City: Clinical characterization and related outcomes. *medRxiv*: 2020.07.02.20145169. DOI: [10.1101/2020.07.02.20145169](https://doi.org/10.1101/2020.07.02.20145169).

Apea VJ, Wan YI, Dhairyawan R, et al. (2021) Ethnicity and outcomes in patients hospitalised with COVID-19 infection in East London: An observational cohort study. *BMJ open* 11. DOI: [10.1136/bmjopen-2020-042140](https://doi.org/10.1136/bmjopen-2020-042140).

Argenziano MG, Bruce SL, Slater CL, et al. (2020) Characterization and clinical course of 1000 patients with coronavirus disease 2019 in New York: Retrospective case series. *BMJ*: m1996. DOI: [10.1136/bmj.m1996](https://doi.org/10.1136/bmj.m1996).

Ariza BE, Torres YX, Salgado D, et al. (2020) Seroprevalence and seroconversion rates to SARS-CoV-2 in interns, residents, and medical doctors in a University Hospital in Bogota, Colombia. *medRxiv*: 2020.09.15.20195313. DOI: [10.1101/2020.09.15.20195313](https://doi.org/10.1101/2020.09.15.20195313).

Arleo T, Tong D, Shabto J, et al. (2020) *Clinical Course and Outcomes of coronavirus disease 2019 (COVID-19) in Rheumatic Disease Patients on Immunosuppression: A case Cohort Study at a Single Center with a Significantly Diverse Population*. preprint, October. *Epidemiology*. DOI: [10.1101/2020.10.26.20219154](https://doi.org/10.1101/2020.10.26.20219154).

- Auvinen R, Nohynek H, Syrjänen R, et al. (2020) Comparison of the clinical characteristics and outcomes of hospitalized adult COVID-19 and influenza patients: A prospective observational study. *medRxiv*: 2020.06.29.20140632. DOI: [10.1101/2020.06.29.20140632](https://doi.org/10.1101/2020.06.29.20140632).
- Ayoubkhani D, Khunti K, Nafilyan V, et al. (2021) Epidemiology of post-COVID syndrome following hospitalisation with coronavirus: A retrospective cohort study. DOI: [10.1101/2021.01.15.21249885](https://doi.org/10.1101/2021.01.15.21249885).
- Barasa S, Kiage-Mokaya J, Cruz-Madrid K, et al. (2020) *Smoking increases the risk of COVID-19 positivity, while Never-smoking reduces the risk*. preprint, December. *Respiratory Medicine*. DOI: [10.1101/2020.11.30.20241380](https://doi.org/10.1101/2020.11.30.20241380).
- Basse C, Diakite S, Servois V, et al. (2020) Characteristics and outcome of SARS-CoV-2 infection in cancer patients. *medRxiv*: 2020.05.14.20101576. DOI: [10.1101/2020.05.14.20101576](https://doi.org/10.1101/2020.05.14.20101576).
- Batty GD, Deary I, Luciano M, et al. (2020) Psychosocial factors and hospitalisations for COVID-19: Prospective cohort study of the general population. *medRxiv*: 2020.05.29.20100735. DOI: [10.1101/2020.05.29.20100735](https://doi.org/10.1101/2020.05.29.20100735).
- Bellan M, Patti G, Hayden E, et al. (2020) Fatality rate and predictors of mortality in an Italian cohort of hospitalized COVID-19 patients. *Scientific Reports* 10(1): 20731. DOI: [10.1038/s41598-020-77698-4](https://doi.org/10.1038/s41598-020-77698-4).
- Bello-Chavolla OY, Bahena-López JP, Antonio-Villa NE, et al. (2020) Predicting Mortality Due to SARS-CoV-2: A Mechanistic Score Relating Obesity and Diabetes to COVID-19 Outcomes in Mexico. *The Journal of Clinical Endocrinology & Metabolism* 105(8). DOI: [10.1210/clinem/dgaa346](https://doi.org/10.1210/clinem/dgaa346).
- Bello-Chavolla OY, Antonio-Villa NE, Vargas-Vázquez A, et al. (2020) Profiling pre-symptomatic and asymptomatic cases with confirmed SARS-CoV-2 infection in Mexico City. *medRxiv*: 2020.07.02.20145516. DOI: [10.1101/2020.07.02.20145516](https://doi.org/10.1101/2020.07.02.20145516).
- Benaim AR, Sobel JA, Almog R, et al. (2020) *At the dawn of winter: Comparing COVID-19 and influenza presentation and trajectory*. preprint, November. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.11.19.20235077](https://doi.org/10.1101/2020.11.19.20235077).
- Benowitz NL, Schultz KE, Haller CA, et al. (2009) Prevalence of smoking assessed biochemically in an urban public hospital: A rationale for routine cotinine screening. *American Journal of Epidemiology* 170(7): 885–891. DOI: [10.1093/aje/kwp215](https://doi.org/10.1093/aje/kwp215).
- Berlin I, Thomas D, Le Faou A-L, et al. (2020) COVID-19 and Smoking. *Nicotine & Tobacco Research*. DOI: [10.1093/NTR/NTAA059](https://doi.org/10.1093/NTR/NTAA059).
- Bermejo-Martin JF, González-Rivera M, Almansa R, et al. (2020) Viral RNA load in plasma is associated with critical illness and a dysregulated host response in COVID-19. *Critical Care* 24(1): 691. DOI: [10.1186/s13054-020-03398-0](https://doi.org/10.1186/s13054-020-03398-0).
- Bernaola N, Mena R, Bernaola A, et al. (2020) Observational Study of the Efficiency of Treatments in Patients Hospitalized with Covid-19 in Madrid. *medRxiv*: 2020.07.17.20155960. DOI: [10.1101/2020.07.17.20155960](https://doi.org/10.1101/2020.07.17.20155960).
- Berumen J, Schmulson M, Alegre J, et al. (2020) Risk of infection and hospitalization by Covid-19 in Mexico: A case-control study. *medRxiv*: 2020.05.24.20104414. DOI: [10.1101/2020.05.24.20104414](https://doi.org/10.1101/2020.05.24.20104414).
- Best JH, Mohan SV, Kong AM, et al. (2020) Baseline Demographics and Clinical Characteristics Among 3471 US Patients Hospitalized with COVID-19 and Pulmonary Involvement: A Retrospective Study. *Advances in Therapy* 37(12): 4981–4995. DOI: [10.1007/s12325-020-01510-y](https://doi.org/10.1007/s12325-020-01510-y).
- Bian H, Zheng Z-H, Wei D, et al. (2020) Meplazumab treats COVID-19 pneumonia: An open-labelled, concurrent controlled add-on clinical trial. *medRxiv*: 2020.03.21.20040691. DOI: [10.1101/2020.03.21.20040691](https://doi.org/10.1101/2020.03.21.20040691).

- Bisso IC, Huespe I, Lockhart C, et al. (2020) *Clinical characteristics of critically ill patients with COVID-19*. preprint, December. Intensive Care; Critical Care Medicine. DOI: [10.1101/2020.12.09.20246413](https://doi.org/10.1101/2020.12.09.20246413).
- Borobia AM, Carcas AJ, Arnalich F, et al. (2020) A Cohort of Patients with COVID-19 in a Major Teaching Hospital in Europe. *Journal of Clinical Medicine* 9(6): 1733. DOI: [10.3390/jcm9061733](https://doi.org/10.3390/jcm9061733).
- Boulware DR, Pullen MF, Bangdiwala AS, et al. (2020) A Randomized Trial of Hydroxychloroquine as Postexposure Prophylaxis for Covid-19. *New England Journal of Medicine*. DOI: [10.1056/NEJMoa2016638](https://doi.org/10.1056/NEJMoa2016638).
- Bowyer RCE, Varsavsky T, Sudre CH, et al. (2020) Geo-social gradients in predicted COVID-19 prevalence and severity in Great Britain: Results from 2,266,235 users of the COVID-19 *Symptoms Tracker app*. DOI: [10.1101/2020.04.23.20076521](https://doi.org/10.1101/2020.04.23.20076521).
- Boyd S and Martin-Loeches I (2021) The incidence of venous thromboembolism in critically ill patients with COVID-19 compared with critically ill non-COVID patients. *Ir J Med Sci* NA. DOI: [10.1007/s11845-020-02503-0](https://doi.org/10.1007/s11845-020-02503-0).
- Brake SJ, Barnsley K, Lu W, et al. (2020) Smoking Upregulates Angiotensin-Converting Enzyme-2 Receptor: A Potential Adhesion Site for Novel Coronavirus SARS-CoV-2 (Covid-19). *J Clin Med* 9, Page 841: 841. DOI: [10.3390/jcm9030841](https://doi.org/10.3390/jcm9030841).
- Brown J (2020) The Smoking Toolkit Study, 2020. Available at: https://docs.google.com/presentation/d/e/2PACX-1vQmMeD0wPM7iEawo39m2QZnVBOo2e8YIQPjm2R0OgYBvKhBbYqnarSn5vxyk3-56w/embed?start=false&loop=false&delayms=3000&usp=embed_facebook.
- Buitrago-Garcia D, Egli-Gany D, Counotte MJ, et al. (2020) Occurrence and transmission potential of asymptomatic and presymptomatic SARS-CoV-2 infections: A living systematic review and meta-analysis. *PLOS Medicine* 17(9): e1003346. DOI: [10.1371/journal.pmed.1003346](https://doi.org/10.1371/journal.pmed.1003346).
- Burrell AJ, Pellegrini B, Salimi F, et al. (2020) Outcomes of COVID-19 patients admitted to Australian intensive care units during the early phase of the pandemic. *The Medical Journal of Australia*: 1. DOI: [10.5694/mja2.50883](https://doi.org/10.5694/mja2.50883).
- Bürkner P-C (2018) Advanced Bayesian Multilevel Modeling with the R Package brms. *The R Journal* 10(1): 395. DOI: [10.32614/RJ-2018-017](https://doi.org/10.32614/RJ-2018-017).
- Cadegiani FA, McCoy J, Wambier CG, et al. (2020) *5-Alpha-Reductase Inhibitors Reduce Remission Time of COVID-19: Results From a Randomized Double Blind Placebo Controlled Interventional Trial in 130 SARS-CoV-2 Positive Men*. preprint, November. Infectious Diseases (except HIV/AIDS). DOI: [10.1101/2020.11.16.20232512](https://doi.org/10.1101/2020.11.16.20232512).
- Caglar A and Kacer I (2021) Anxiety levels in patients admitted to the emergency department with myocardial infarction or COVID-19 pneumonia. *PSYCHOL HEALTH MED* NA: 1–9. DOI: [10.1080/13548506.2021.1876893](https://doi.org/10.1080/13548506.2021.1876893).
- Cai G (2020) Bulk and Single-Cell Transcriptomics Identify Tobacco-Use Disparity in Lung Gene Expression of ACE2, the Receptor of 2019-nCov. DOI: [10.20944/PREPRINTS202002.0051.V3](https://doi.org/10.20944/PREPRINTS202002.0051.V3).
- Cai H, Yang L, Lu Y, et al. (2021) High body mass index is a significant risk factor for the progression and prognosis of imported COVID-19: A multicenter, retrospective cohort study. *BMC Infect Dis* 21. DOI: [10.1186/s12879-021-05818-0](https://doi.org/10.1186/s12879-021-05818-0).
- Caliskan T and Saylan B (2020) Smoking and comorbidities are associated with COVID-19 severity and mortality in 565 patients treated in Turkey: A retrospective observational study. *Rev Assoc Med Bras* 66: 1679–1684. DOI: [10.1590/1806-9282.66.12.1679](https://doi.org/10.1590/1806-9282.66.12.1679).
- Carrat F, Lamballerie X de, Rahib D, et al. (2020) Seroprevalence of SARS-CoV-2 among adults in three regions of France following the lockdown and associated risk factors: A multicohort study. *medRxiv*: 2020.09.16.20195693. DOI:

[10.1101/2020.09.16.20195693](https://doi.org/10.1101/2020.09.16.20195693).

Carrillo-Vega MF, Salinas-Escudero G, Garcia-Peña C, et al. (2020) Early estimation of the risk factors for hospitalisation and mortality by COVID-19 in Mexico. *medRxiv*: 2020.05.11.20098145. DOI: [10.1101/2020.05.11.20098145](https://doi.org/10.1101/2020.05.11.20098145).

CDCMMWR (2020) Preliminary Estimates of the Prevalence of Selected Underlying Health Conditions Among Patients with Coronavirus Disease 2019 — United States, February 12–March 28, 2020. *MMWR. Morbidity and Mortality Weekly Report* 69. DOI: [10.15585/mmwr.mm6913e2](https://doi.org/10.15585/mmwr.mm6913e2).

Gen Y, Chen X, Shen Y, et al. (2020) Risk factors for disease progression in patients with mild to moderate coronavirus disease 2019—a multi-centre observational study. *Clinical Microbiology and Infection*: S1198743X20303414. DOI: [10.1016/j.cmi.2020.05.041](https://doi.org/10.1016/j.cmi.2020.05.041).

Chand S, Kapoor S, Orsi D, et al. (2020) COVID-19-Associated Critical Illness—Report of the First 300 Patients Admitted to Intensive Care Units at a New York City Medical Center: *Journal of Intensive Care Medicine*. DOI: [10.1177/0885066620946692](https://doi.org/10.1177/0885066620946692).

Chaudhary A, Singh UN, Paudel P, et al. (2020) Characteristics and outcomes of hospitalized adults with COVID-19 in Nepal: A multicenter, prospective cohort study. *medRxiv*: 2020.10.03.20206128. DOI: [10.1101/2020.10.03.20206128](https://doi.org/10.1101/2020.10.03.20206128).

Chaudhry F, Bulka H, Rathnam AS, et al. (2020) COVID-19 in Multiple Sclerosis Patients and Risk Factors for Severe Infection. *medRxiv*: 2020.05.27.20114827. DOI: [10.1101/2020.05.27.20114827](https://doi.org/10.1101/2020.05.27.20114827).

Chen C, Jiang J, Xu X, et al. (2020) Dynamic liver function indexes monitoring and clinical characteristics in three types of COVID-19 patients. *medRxiv*: 2020.05.13.20099614. DOI: [10.1101/2020.05.13.20099614](https://doi.org/10.1101/2020.05.13.20099614).

Chen H, Varatharajah Y, Ramirez SS de, et al. (2020) *A Retrospective Longitudinal Study of COVID-19 as Seen by a Large Urban Hospital in Chicago*. preprint, November. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.11.29.20240606](https://doi.org/10.1101/2020.11.29.20240606).

Chen L, Yu J, He W, et al. (2020) Risk factors for death in 1859 subjects with COVID-19. *Leukemia*: 1–11. DOI: [10.1038/s41375-020-0911-0](https://doi.org/10.1038/s41375-020-0911-0).

Chen T, Wu D, Chen H, et al. (2020) Clinical characteristics of 113 deceased patients with coronavirus disease 2019: Retrospective study. *BMJ* 368. DOI: [10.1136/bmj.m1091](https://doi.org/10.1136/bmj.m1091).

Cho ER, Slutsky AS and Jha P (2020) Smoking and the risk of COVID-19 infection in the UK Biobank Prospective Study. *medRxiv*: 2020.05.05.20092445. DOI: [10.1101/2020.05.05.20092445](https://doi.org/10.1101/2020.05.05.20092445).

Chudasama YV, Zaccardi F, Gillies CL, et al. (2020) *Patterns of Multimorbidity and Risk of Severe SARS-CoV-2 Infection: An observational study in the U.K.* preprint, October. *Epidemiology*. DOI: [10.1101/2020.10.21.20216721](https://doi.org/10.1101/2020.10.21.20216721).

Clavario P, De Marzo V, Lotti R, et al. (2020) *Assessment of functional capacity with cardiopulmonary exercise testing in non-severe COVID-19 patients at three months follow-up*. preprint, November. *Cardiovascular Medicine*. DOI: [10.1101/2020.11.15.20231985](https://doi.org/10.1101/2020.11.15.20231985).

CMMID COVID-19 working group, Jarvis CI, Van Zandvoort K, et al. (2020) Quantifying the impact of physical distance measures on the transmission of COVID-19 in the UK. *BMC Medicine* 18(1): 124. DOI: [10.1186/s12916-020-01597-8](https://doi.org/10.1186/s12916-020-01597-8).

Collard D, Nurmohamed NS, Kaiser Y, et al. (2020) *Cardiovascular risk factors are independently associated with COVID-19 mortality: A prospective cohort study*. preprint, October. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.10.01.20205229](https://doi.org/10.1101/2020.10.01.20205229).

- Concha-Mejia A and Rincon-Sanchez RA (2020) CCOFEE-GI Study: Colombian COVID19 First Experience in Gastroenterology. Characterization of digestive manifestations in patients diagnosed with COVID-19 at a highly complex institution in Bogota D.C., Colombia. *medRxiv*: 2020.07.24.20161604. DOI: [10.1101/2020.07.24.20161604](https://doi.org/10.1101/2020.07.24.20161604).
- Conti C, Marco A de, Mastromarino P, et al. (1999) Antiviral Effect of Hyperthermic Treatment in Rhinovirus Infection. *Antimicrobial Agents and Chemotherapy* 43(4): 822–829. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC89212/> (accessed 1 October 2020).
- Covid-19 in pregnancy, Vousden N, Bunch K, et al. (2021) The incidence, characteristics and outcomes of pregnant women hospitalized with symptomatic and asymptomatic SARS-CoV-2 infection in the UK from March to September 2020: A national cohort study using the UK Obstetric Surveillance System (UKOSS). DOI: [10.1101/2021.01.04.21249195](https://doi.org/10.1101/2021.01.04.21249195).
- Crooks CJ, West J, Fogarty A, et al. (2020) Predicting the need for escalation of care or death from repeated daily clinical observations and laboratory results in patients with SARS-CoV-2 during 2020: A retrospective population-based cohort study from the United Kingdom. DOI: [10.1101/2020.12.14.20248181](https://doi.org/10.1101/2020.12.14.20248181).
- Crovetto F, Crispi F, Llurba E, et al. (2020) SEROPREVALENCE AND CLINICAL SPECTRUM OF SARS-CoV-2 INFECTION IN THE FIRST VERSUS THIRD TRIMESTER OF PREGNANCY. *medRxiv*: 2020.06.17.20134098. DOI: [10.1101/2020.06.17.20134098](https://doi.org/10.1101/2020.06.17.20134098).
- Cummins L, Ebyarimpa I, Cheetham N, et al. (2021) Factors associated with COVID-19 related hospitalisation, critical care admission and mortality using linked primary and secondary care data. DOI: [10.1101/2021.01.19.20241844](https://doi.org/10.1101/2021.01.19.20241844).
- Dai M, Tao L, Chen Z, et al. (2020) Influence of Cigarettes and Alcohol on the Severity and Death of COVID-19: A Multicenter Retrospective Study in Wuhan, China. *Front Physiol* 11. DOI: [10.3389/fphys.2020.588553](https://doi.org/10.3389/fphys.2020.588553).
- Dashti H, Roche E, Bates D, et al. (2020) SARS2 simplified scores to estimate risk of hospitalization and death among patients with COVID-19. DOI: [10.1101/2020.09.11.20190520](https://doi.org/10.1101/2020.09.11.20190520).
- Denholm JT, Gordon CL, Johnson PD, et al. (2010) Hospitalised adult patients with pandemic (H1N1) 2009 influenza in Melbourne, Australia. *Medical Journal of Australia* 192(2): 84–86. DOI: [10.5694/j.1326-5377.2010.tb03424.x](https://doi.org/10.5694/j.1326-5377.2010.tb03424.x).
- Didikoglu A, Maharani A, Pendleton N, et al. (2021) Early life factors and COVID-19 infection in England: A prospective analysis of UK Biobank participants. *Early Hum Dev* 155. DOI: [10.1016/j.earlhumdev.2021.105326](https://doi.org/10.1016/j.earlhumdev.2021.105326).
- Díez-Manglano J, Marquínez MNS, García AÁ, et al. (2020) *Healthcare workers hospitalized due to COVID-19 have no higher risk of death than general population. Data from the Spanish SEMI-COVID-19 Registry*. preprint, November. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.11.23.20236810](https://doi.org/10.1101/2020.11.23.20236810).
- Docherty AB, Harrison EM, Green CA, et al. (2020) Features of 20 133 UK patients in hospital with covid-19 using the ISARIC WHO Clinical Characterisation Protocol: Prospective observational cohort study. *BMJ* 369. DOI: [10.1136/bmj.m1985](https://doi.org/10.1136/bmj.m1985).
- Dong X, Cao Y-y, Lu X-x, et al. (2020) Eleven faces of coronavirus disease 2019. *Allergy* 75(7): 1699–1709. DOI: [10.1111/all.14289](https://doi.org/10.1111/all.14289).
- Duan L, Zhang S, Guo M, et al. (2020) Epidemiological and clinical characteristics in patients with SARS-CoV-2 antibody negative probable COVID-19 in Wuhan. *medRxiv*: 2020.06.18.20134619. DOI: [10.1101/2020.06.18.20134619](https://doi.org/10.1101/2020.06.18.20134619).
- Dupraz J, Butty A, Duperrex O, et al. (2020) *Prevalence of SARS-CoV-2 in household members and other close contacts of COVID-19 cases: A serologic study in canton of Vaud, Switzerland*. preprint, November. *Epidemiology*. DOI:

[10.1101/2020.11.27.20239244](https://doi.org/10.1101/2020.11.27.20239244).

Dye JA and Adler KB (1994) Occasional review Effects of cigarette smoke on epithelial cells of the respiratory tract. *Thorax* 49: 825–834. DOI: [10.1136/thx.49.8.825](https://doi.org/10.1136/thx.49.8.825).

Ebinger J, Botwin GJ, Albert CM, et al. (2020) SARS-CoV-2 Seroprevalence Across a Diverse Cohort of Healthcare Workers. *medRxiv*: 2020.07.31.20163055. DOI: [10.1101/2020.07.31.20163055](https://doi.org/10.1101/2020.07.31.20163055).

Ebrahimian S, Homayounieh F, Rockenbach MABC, et al. (2021) Artificial intelligence matches subjective severity assessment of pneumonia for prediction of patient outcome and need for mechanical ventilation: A cohort study. *Sci. rep* 11. DOI: [10.1038/s41598-020-79470-0](https://doi.org/10.1038/s41598-020-79470-0).

Eiros R, Barreiro-Perez M, Martin-Garcia A, et al. (2020) Pericarditis and myocarditis long after SARS-CoV-2 infection: A cross-sectional descriptive study in health-care workers. *medRxiv*: 2020.07.12.20151316. DOI: [10.1101/2020.07.12.20151316](https://doi.org/10.1101/2020.07.12.20151316).

Elezkurtaj S, Greuel S, Ihlow J, et al. (2020) Causes of Death and Comorbidities in Patients with COVID-19. *medRxiv*: 2020.06.15.20131540. DOI: [10.1101/2020.06.15.20131540](https://doi.org/10.1101/2020.06.15.20131540).

Elliott JH, Turner T, Clavisi O, et al. (2014) Living Systematic Reviews: An Emerging Opportunity to Narrow the Evidence-Practice Gap. *PLoS Medicine* 11(2). DOI: [10.1371/journal.pmed.1001603](https://doi.org/10.1371/journal.pmed.1001603).

Elmunzer BJ, Spitzer RL, Foster LD, et al. (2020) Digestive Manifestations in Patients Hospitalized with COVID-19. *medRxiv*: 2020.07.07.20143024. DOI: [10.1101/2020.07.07.20143024](https://doi.org/10.1101/2020.07.07.20143024).

El-Solh AA, Meduri UG, Lawson Y, et al. (2020) CLINICAL COURSE AND OUTCOME OF COVID-19 ACUTE RESPIRATORY DISTRESS SYNDROME: DATA FROM A NATIONAL REPOSITORY. *medRxiv*: 2020.10.16.20214130. DOI: [10.1101/2020.10.16.20214130](https://doi.org/10.1101/2020.10.16.20214130).

Emami A, Javanmardi F, Pirbonyeh N, et al. (2020) Prevalence of Underlying Diseases in Hospitalized Patients with COVID-19: A Systematic Review and Meta-Analysis. *Archives of academic emergency medicine* 8(1): e35.

Erber J, Kappler V, Haller B, et al. (2020) *Strategies for infection control and prevalence of anti-SARS-CoV-2 IgG in 4,554 employees of a university hospital in Munich, Germany*. preprint, October. *Epidemiology*. DOI: [10.1101/2020.10.04.20206136](https://doi.org/10.1101/2020.10.04.20206136).

Estiri H, Strasser ZH, Klann JG, et al. (2021) Predicting COVID-19 mortality with electronic medical records. *npj digit. med* 4. DOI: [10.1038/s41746-021-00383-x](https://doi.org/10.1038/s41746-021-00383-x).

Eugen-Olsen J, Altintas I, Tingleff J, et al. (2020) Low levels of the prognostic biomarker suPAR are predictive of mild outcome in patients with symptoms of COVID-19 - a prospective cohort study. *medRxiv*: 2020.05.27.20114678. DOI: [10.1101/2020.05.27.20114678](https://doi.org/10.1101/2020.05.27.20114678).

Fan X, Yin C, Wang J, et al. (2020) Pre-diagnostic circulating concentrations of insulin-like growth factor-1 and risk of COVID-19 mortality: Results from UK Biobank. *medRxiv*: 2020.07.09.20149369. DOI: [10.1101/2020.07.09.20149369](https://doi.org/10.1101/2020.07.09.20149369).

Farsalinos K, Niaura R, Le Houezec J, et al. (2020) Editorial: Nicotine and SARS-CoV-2: COVID-19 may be a disease of the nicotinic cholinergic system. *Toxicology Reports*. DOI: [10.1016/j.toxrep.2020.04.012](https://doi.org/10.1016/j.toxrep.2020.04.012).

Farsalinos K, Barbouni A and Niaura R (2020) Systematic review of the prevalence of current smoking among hospitalized COVID-19 patients in China: Could nicotine be a therapeutic option? *Internal and Emergency Medicine*. DOI: [10.1007/s11739-020-02355-7](https://doi.org/10.1007/s11739-020-02355-7).

- Favara DM, McAdam K, Cooke A, et al. (2020) SARS-CoV-2 antigen and antibody prevalence among UK staff working with cancer patients during the COVID-19 pandemic. DOI: [10.1101/2020.09.18.20197590](https://doi.org/10.1101/2020.09.18.20197590).
- Feldman C and Anderson R (2013) Cigarette smoking and mechanisms of susceptibility to infections of the respiratory tract and other organ systems. *J. Infect* 67: 169–84. DOI: [10.1016/j.jinf.2013.05.004](https://doi.org/10.1016/j.jinf.2013.05.004).
- Feng Y, Ling Y, Bai T, et al. (2020) COVID-19 with Different Severities: A Multicenter Study of Clinical Features. *American Journal of Respiratory and Critical Care Medicine* 201(11): 1380–1388. DOI: [10.1164/rccm.202002-0445OC](https://doi.org/10.1164/rccm.202002-0445OC).
- Ferrari BL, Ferreira CG, Menezes M, et al. (2021) Determinants of COVID-19 Mortality in Patients With Cancer From a Community Oncology Practice in Brazil. *JCO Glob Oncol* 7: 46–55. DOI: [10.1200/GO.20.00444](https://doi.org/10.1200/GO.20.00444).
- Feuth T, Saaresranta T, Karlsson A, et al. (2020) Is sleep apnoea a risk factor for Covid-19? Findings from a retrospective cohort study. *medRxiv*: 2020.05.14.20098319. DOI: [10.1101/2020.05.14.20098319](https://doi.org/10.1101/2020.05.14.20098319).
- Fillmore NR, La J, Szalat RE, et al. (2020) Prevalence and outcome of Covid-19 infection in cancer patients: A national VA study. *medRxiv*: 2020.08.21.20177923. DOI: [10.1101/2020.08.21.20177923](https://doi.org/10.1101/2020.08.21.20177923).
- Fisman D, Greer AL and Tuite A (2020) Derivation and Validation of Clinical Prediction Rule for COVID-19 Mortality in Ontario, Canada. *medRxiv*: 2020.06.21.20136929. DOI: [10.1101/2020.06.21.20136929](https://doi.org/10.1101/2020.06.21.20136929).
- Fond G, Pauly V, Orleans V, et al. (2020) Increased in-hospital mortality from COVID-19 in patients with schizophrenia. *L'Encéphale*: S0013700620301883. DOI: [10.1016/j.encep.2020.07.003](https://doi.org/10.1016/j.encep.2020.07.003).
- Fontanet A, Tondeur L, Madec Y, et al. (2020) Cluster of COVID-19 in northern France: A retrospective closed cohort study. *medRxiv*: 2020.04.18.20071134. DOI: [10.1101/2020.04.18.20071134](https://doi.org/10.1101/2020.04.18.20071134).
- Fox TA, Troy-Barnes E, Kirkwood AA, et al. (n.d.) Clinical outcomes and risk factors for severe COVID-19 infection in patients with haematological disorders receiving chemo- or immunotherapy. *British Journal of Haematology* n/a(n/a). DOI: [10.1111/bjh.17027](https://doi.org/10.1111/bjh.17027).
- Freites D, Leon L, Mucientes A, et al. (2020) Risk factors for hospital admission related to COVID-19 in inflammatory rheumatic diseases. *medRxiv*: 2020.05.14.20101584. DOI: [10.1101/2020.05.14.20101584](https://doi.org/10.1101/2020.05.14.20101584).
- Fung HF, Martinez L, Alarid-Escudero F, et al. (2020) The household secondary attack rate of SARS-CoV-2: A rapid review. *Clinical Infectious Diseases*. DOI: [10.1093/cid/ciaa1558](https://doi.org/10.1093/cid/ciaa1558).
- Galal I, Hussein AAM, Amin MT, et al. (2020) Determinants of Persistent Post COVID-19 symptoms: Value of a Novel COVID-19 symptoms score. DOI: [10.1101/2020.11.11.20230052](https://doi.org/10.1101/2020.11.11.20230052).
- Gallichotte EN, Quicke KM, Sexton NR, et al. (2020) Longitudinal Surveillance for SARS-CoV-2 Among Staff in Six Colorado Long-Term Care Facilities: Epidemiologic, Virologic and Sequence Analysis. *medRxiv*: 2020.06.08.20125989. DOI: [10.1101/2020.06.08.20125989](https://doi.org/10.1101/2020.06.08.20125989).
- Garassino MC, Whisenant JG, Huang L-C, et al. (2020) COVID-19 in patients with thoracic malignancies (TERAVOLT): First results of an international, registry-based, cohort study. *The Lancet Oncology* 21(7): 914–922. DOI: [10.1016/S1470-2045\(20\)30314-4](https://doi.org/10.1016/S1470-2045(20)30314-4).
- Garibaldi BT, Fiksel J, Muschelli J, et al. (2020) Patient trajectories and risk factors for severe outcomes among persons hospitalized for COVID-19 in the Maryland/DC region. *medRxiv*: 2020.05.24.20111864. DOI: [10.1101/2020.05.24.20111864](https://doi.org/10.1101/2020.05.24.20111864).
- Ge H, Zhu M, Du J, et al. (2020) Cardiac Structural and Functional Characteristics in Patients with Coronavirus Disease

- 2019: A Serial Echocardiographic Study. *medRxiv*: 2020.05.12.20095885. DOI: [10.1101/2020.05.12.20095885](https://doi.org/10.1101/2020.05.12.20095885).
- Gégout petit A, Jeulin H, Legrand K, et al. (2021) Seroprevalence of SARS-CoV-2, symptom profiles and seroneutralization during the first COVID-19 wave in a suburban area, France. DOI: [10.1101/2021.02.10.21250862](https://doi.org/10.1101/2021.02.10.21250862).
- Ghinai I, Davis ES, Mayer S, et al. (2020) Risk Factors for Severe Acute Respiratory Syndrome Coronavirus 2 Infection in Homeless Shelters in Chicago, Illinois—March–May, 2020. *Open Forum Infectious Diseases* 7(11): ofaa477. DOI: [10.1093/ofid/ofaa477](https://doi.org/10.1093/ofid/ofaa477).
- Giacomelli A, Ridolfo AL, Milazzo L, et al. (2020) 30-day mortality in patients hospitalized with COVID-19 during the first wave of the Italian epidemic: A prospective cohort study. *Pharmacological Research* 158: 104931. DOI: [10.1016/j.phrs.2020.104931](https://doi.org/10.1016/j.phrs.2020.104931).
- Gianfrancesco M, Hyrich KL, Al-Adely S, et al. (2020) Characteristics associated with hospitalisation for COVID-19 in people with rheumatic disease: Data from the COVID-19 Global Rheumatology Alliance physician-reported registry. *Annals of the Rheumatic Diseases* 79(7): 859–866. DOI: [10.1136/annrheumdis-2020-217871](https://doi.org/10.1136/annrheumdis-2020-217871).
- Gianfrancesco MA, Leykina LA, Izadi Z, et al. (2020) Race/ethnicity association with COVID-19 outcomes in rheumatic disease: Data from the COVID-19 Global Rheumatology Alliance Physician Registry. *Arthritis & Rheumatology*: art.41567. DOI: [10.1002/art.41567](https://doi.org/10.1002/art.41567).
- Giannini S, Passeri G, Tripepi G, et al. (2021) Effectiveness of In-Hospital Cholecalciferol Use on Clinical Outcomes in Comorbid COVID-19 Patients: A Hypothesis-Generating Study. *Nutrients* 13. DOI: [10.3390/nu13010219](https://doi.org/10.3390/nu13010219).
- Giannouchos T, Sussman R, Mier JM, et al. (2020) Characteristics and risk factors for COVID-19 diagnosis and adverse outcomes in Mexico: An analysis of 89,756 laboratory-confirmed COVID-19 cases. *medRxiv*: 2020.06.04.20122481. DOI: [10.1101/2020.06.04.20122481](https://doi.org/10.1101/2020.06.04.20122481).
- Gil-Agudo A, Rodriguez-Cola M, Jimenez-Velasco I, et al. (2020) Clinical features of coronavirus disease 2019 (COVID-19) in a cohort of patients with disability due to spinal cord injury. *medRxiv*: 2020.04.20.20072918. DOI: [10.1101/2020.04.20.20072918](https://doi.org/10.1101/2020.04.20.20072918).
- Girardeau Y, Gallois Y, De Bonnecaze G, et al. (2020) Confirmed central olfactory system lesions on brain MRI in COVID-19 patients with anosmia: A case-series. DOI: [10.1101/2020.07.08.20148692](https://doi.org/10.1101/2020.07.08.20148692).
- Gold JAW, Wong KK, Szablewski CM, et al. (2020) Characteristics and Clinical Outcomes of Adult Patients Hospitalized with COVID-19 — Georgia, March 2020. *MMWR. Morbidity and Mortality Weekly Report* 69(18): 545–550. DOI: [10.15585/mmwr.mm6918e1](https://doi.org/10.15585/mmwr.mm6918e1).
- Gori M, Berzuini C, D’Elia E, et al. (2020) Exposure to Renin-Angiotensin System Inhibitors Is Associated with Reduced Mortality of Older Hypertensive Covid-19 Patients. DOI: [10.1101/2020.12.15.20247999](https://doi.org/10.1101/2020.12.15.20247999).
- Govind R, Freitas DF de, Pritchard MR, et al. (2020) Clozapine treatment and risk of COVID-19. *medRxiv*: 2020.06.17.20133595. DOI: [10.1101/2020.06.17.20133595](https://doi.org/10.1101/2020.06.17.20133595).
- Goyal P, Choi JJ, Pinheiro LC, et al. (2020) Clinical Characteristics of Covid-19 in New York City. *New England Journal of Medicine* 382(24): 2372–2374. DOI: [10.1056/NEJMc2010419](https://doi.org/10.1056/NEJMc2010419).
- Griffith G, Morris TT, Tudball M, et al. (2020) Collider bias undermines our understanding of COVID-19 disease risk and severity. *medRxiv*: 2020.05.04.20090506. DOI: [10.1101/2020.05.04.20090506](https://doi.org/10.1101/2020.05.04.20090506).
- Grundy* EJ, Suddek* T, Filippidis FT, et al. (2020) Smoking, SARS-CoV-2 and COVID-19: A review of reviews

- considering implications for public health policy and practice. *Tobacco Induced Diseases* 18(July). DOI: [10.18332/tid/124788](https://doi.org/10.18332/tid/124788).
- Gu T, Mack JA, Salvatore M, et al. (2020) COVID-19 outcomes, risk factors and associations by race: A comprehensive analysis using electronic health records data in Michigan Medicine. *medRxiv*: 2020.06.16.20133140. DOI: [10.1101/2020.06.16.20133140](https://doi.org/10.1101/2020.06.16.20133140).
- Guan W-j, Ni Z-y, Hu Y, et al. (2020) Clinical Characteristics of Coronavirus Disease 2019 in China. *New England Journal of Medicine*. DOI: [10.1056/NEJMoa2002032](https://doi.org/10.1056/NEJMoa2002032).
- Guan W-j, Liang W-h, Zhao Y, et al. (2020) Comorbidity and its impact on 1590 patients with COVID-19 in China: A nationwide analysis. *European Respiratory Journal* 55(5). DOI: [10.1183/13993003.00547-2020](https://doi.org/10.1183/13993003.00547-2020).
- Guo T, Fan Y, Chen M, et al. (2020) Cardiovascular Implications of Fatal Outcomes of Patients With Coronavirus Disease 2019 (COVID-19). *JAMA Cardiology* 5(7): 811–818. DOI: [10.1001/jamacardio.2020.1017](https://doi.org/10.1001/jamacardio.2020.1017).
- Gupta R, Agrawal R, Bukhari Z, et al. (2020) Higher Comorbidities and Early Death is Characteristic of Hospitalized African-American Patients with COVID-19. *medRxiv*: 2020.07.15.20154906. DOI: [10.1101/2020.07.15.20154906](https://doi.org/10.1101/2020.07.15.20154906).
- Hadi YB, Naqvi SFZ, Kupec JT, et al. (2020) Characteristics and outcomes of COVID-19 in patients with HIV: A multicentre research network study. *AIDS* 34(13): F3–F8. DOI: [10.1097/QAD.0000000000002666](https://doi.org/10.1097/QAD.0000000000002666).
- Hadjadj J, Yatim N, Barnabei L, et al. (2020) Impaired type I interferon activity and exacerbated inflammatory responses in severe Covid-19 patients. *medRxiv*: 2020.04.19.20068015. DOI: [10.1101/2020.04.19.20068015](https://doi.org/10.1101/2020.04.19.20068015).
- Hamadah H, Alahmad B, Behbehani M, et al. (2020) COVID-19 clinical outcomes and nationality: Results from a Nationwide registry in Kuwait. *BMC Public Health* 20(1): 1384. DOI: [10.1186/s12889-020-09490-y](https://doi.org/10.1186/s12889-020-09490-y).
- Hamer M, Kivimäki M, Gale CR, et al. (2020) Lifestyle risk factors, inflammatory mechanisms, and COVID-19 hospitalization: A community-based cohort study of 387,109 adults in UK. *Brain, Behavior, and Immunity* 87: 184–187. DOI: [10.1016/j.bbi.2020.05.059](https://doi.org/10.1016/j.bbi.2020.05.059).
- Hao S-R, Zhang S-Y, Lian J-S, et al. (2020) Liver Enzyme Elevation in Coronavirus Disease 2019: A Multicenter, Retrospective, Cross-Sectional Study. *The American Journal of Gastroenterology*. DOI: [10.14309/ajg.0000000000000717](https://doi.org/10.14309/ajg.0000000000000717).
- Heili-Frades S (n.d.) COVID-19 Outcomes in 4712 consecutively confirmed SARS-CoV2 cases in the city of Madrid. *medRxiv*. DOI: [10.1101/2020.05.22.20109850](https://doi.org/10.1101/2020.05.22.20109850).
- Hernández-Garduño E (2020) Obesity is the comorbidity more strongly associated for Covid-19 in Mexico. A case-control study. *Obesity Research & Clinical Practice*. DOI: [10.1016/j.orcp.2020.06.001](https://doi.org/10.1016/j.orcp.2020.06.001).
- Hewitt J, Carter B, Vilches-Moraga A, et al. (2020) The effect of frailty on survival in patients with COVID-19 (COPE): A multicentre, European, observational cohort study. *The Lancet Public Health* 5(8): e444–e451. DOI: [10.1016/S2468-2667\(20\)30146-8](https://doi.org/10.1016/S2468-2667(20)30146-8).
- Higuchi T, Nishida T, Iwahashi H, et al. (2020) Early Clinical Factors Predicting the Development of Critical Disease in Japanese Patients with COVID-19: A Single-Center Retrospective, Observational Study. *medRxiv*: 2020.07.29.20159442. DOI: [10.1101/2020.07.29.20159442](https://doi.org/10.1101/2020.07.29.20159442).
- Hippisley-Cox J, Young D, Coupland C, et al. (2020) Risk of severe COVID-19 disease with ACE inhibitors and angiotensin receptor blockers: Cohort study including 8.3 million people. *Heart*: heartjnl-2020-317393. DOI: [10.1136/heartjnl-2020-317393](https://doi.org/10.1136/heartjnl-2020-317393).

- Ho FK, Celis-Morales CA, Gray SR, et al. (2020) Modifiable and non-modifiable risk factors for COVID-19, and comparison to risk factors for influenza and pneumonia: Results from a UK Biobank prospective cohort study. *BMJ open* 10. DOI: [10.1136/bmjopen-2020-040402](https://doi.org/10.1136/bmjopen-2020-040402).
- Ho KS, Narasimhan B, Sheehan J, et al. (2020) Controversy over Smoking in COVID-19 - A Real World Experience in New York City. *J Med Virol* NA. DOI: [10.1002/jmv.26738](https://doi.org/10.1002/jmv.26738).
- Hoertel N, Rico MS, Vernet R, et al. (2020) Association between SSRI Antidepressant Use and Reduced Risk of Intubation or Death in Hospitalized Patients with Coronavirus Disease 2019: A Multicenter Retrospective Observational Study. *medRxiv*: 2020.07.09.20143339. DOI: [10.1101/2020.07.09.20143339](https://doi.org/10.1101/2020.07.09.20143339).
- Hoertel N, Sánchez M, Vernet R, et al. (2020) *Dexamethasone use and Mortality in Hospitalized Patients with Coronavirus Disease 2019: A Multicenter Retrospective Observational Study*. preprint, October. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.10.23.20218172](https://doi.org/10.1101/2020.10.23.20218172).
- Hoertel N, Rico MS, Vernet R, et al. (2020) Observational Study of Haloperidol in Hospitalized Patients with Covid-19. *medRxiv*: 2020.07.15.20150490. DOI: [10.1101/2020.07.15.20150490](https://doi.org/10.1101/2020.07.15.20150490).
- Hoffmann M, Kleine-Weber H, Schroeder S, et al. (2020) SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell* 181(2): 271–280.e8. DOI: [10.1016/j.cell.2020.02.052](https://doi.org/10.1016/j.cell.2020.02.052).
- Holman N, Knighton P, Kar P, et al. (2020) Risk factors for COVID-19-related mortality in people with type 1 and type 2 diabetes in England: A population-based cohort study. *The Lancet Diabetes & Endocrinology* 0(0). DOI: [10.1016/S2213-8587\(20\)30271-0](https://doi.org/10.1016/S2213-8587(20)30271-0).
- Hopkinson NS, Rossi N, Moustafa JE-S, et al. (2020) Current tobacco smoking and risk from COVID-19: Results from a population symptom app in over 2.4 million people. *medRxiv*: 2020.05.18.20105288. DOI: [10.1101/2020.05.18.20105288](https://doi.org/10.1101/2020.05.18.20105288).
- Houlihan CF, Vora N, Byrne T, et al. (2020) Pandemic peak SARS-CoV-2 infection and seroconversion rates in London frontline health-care workers. *The Lancet*: S0140673620314847. DOI: [10.1016/S0140-6736\(20\)31484-7](https://doi.org/10.1016/S0140-6736(20)31484-7).
- Hu L, Chen S, Fu Y, et al. (2020) Risk Factors Associated with Clinical Outcomes in 323 COVID-19 Patients in Wuhan, China. *medRxiv*: 2020.03.25.20037721. DOI: [10.1101/2020.03.25.20037721](https://doi.org/10.1101/2020.03.25.20037721).
- Huang C, Wang Y, Li X, et al. (2020) Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *The Lancet* 395(10223): 497–506. DOI: [10.1016/S0140-6736\(20\)30183-5](https://doi.org/10.1016/S0140-6736(20)30183-5).
- Huang Y, Yang R, Xu Y, et al. (2020) Clinical characteristics of 36 non-survivors with COVID-19 in Wuhan, China. *medRxiv*: 2020.02.27.20029009. DOI: [10.1101/2020.02.27.20029009](https://doi.org/10.1101/2020.02.27.20029009).
- Hultcrantz M, Richter J, Rosenbaum C, et al. (2020) COVID-19 infections and outcomes in patients with multiple myeloma in New York City: A cohort study from five academic centers. *medRxiv*: 2020.06.09.20126516. DOI: [10.1101/2020.06.09.20126516](https://doi.org/10.1101/2020.06.09.20126516).
- Hussein MH, Toraih EA, Attia AS, et al. (2020) Asthma in COVID-19: An extra chain fitting around the neck? *medRxiv*: 2020.07.13.20153130. DOI: [10.1101/2020.07.13.20153130](https://doi.org/10.1101/2020.07.13.20153130).
- Ibarra-Nava I, Flores-Rodriguez KG, Ruiz-Herrera V, et al. (2020) Ethnic disparities in COVID-19 mortality in Mexico: A cross-sectional study based on national data. *medRxiv*: 2020.08.26.20182543. DOI: [10.1101/2020.08.26.20182543](https://doi.org/10.1101/2020.08.26.20182543).
- Ibrahim D, Dulipsingh L, Zapatka L, et al. (2020) Factors Associated with Good Patient Outcomes Following Convalescent Plasma in COVID-19: A Prospective Phase II Clinical Trial. *medRxiv*: 2020.08.27.20183293. DOI:

[10.1101/2020.08.27.20183293](https://doi.org/10.1101/2020.08.27.20183293).

lftime S, López-Azcona AF, Vallverdu I, et al. (2020) *First and second waves of coronavirus disease-19: A comparative study in hospitalized patients in Reus, Spain*. preprint, December. Infectious Diseases (except HIV/AIDS). DOI:

[10.1101/2020.12.10.20246959](https://doi.org/10.1101/2020.12.10.20246959).

lkitimur H, Uysal BB, Cengiz M, et al. (n.d.) "Determining Host Factors Contributing to Disease Severity in a Family Cluster of 29 Hospitalized SARS-CoV-2 Patients: Could Genetic Factors Be Relevant in the Clinical Course of COVID-19?".

Journal of Medical Virology n/a(n/a). DOI: [10.1002/jmv.26106](https://doi.org/10.1002/jmv.26106).

llic I, Zdravkovic M, Timcic S, et al. (2021) Pneumonia in healthcare workers during a COVID-19 outbreak at a cardiovascular hospitals. *International Journal of Infectious Diseases* 103: 188–193. DOI: [10.1016/j.ijid.2020.11.156](https://doi.org/10.1016/j.ijid.2020.11.156).

Incerti D, Rizzo S, Li X, et al. (2020) *Risk factors for mortality among hospitalized patients with COVID-19*. preprint, September. Infectious Diseases (except HIV/AIDS). DOI: [10.1101/2020.09.22.20196204](https://doi.org/10.1101/2020.09.22.20196204).

International Severe Acute Respiratory and emerging Infections Consortium, Escher M, Hall M, et al. (2020) ISARIC Clinical Data Report 10 February 2021. DOI: [10.1101/2020.07.17.20155218](https://doi.org/10.1101/2020.07.17.20155218).

Invernizzi A, Torre A, Parrulli S, et al. (2020) Retinal findings in patients with COVID-19: Results from the SERPICO-19 study. *EClinicalMedicine* 27: 100550. DOI: [10.1016/j.eclinm.2020.100550](https://doi.org/10.1016/j.eclinm.2020.100550).

Ioannou GN, Locke E, Green P, et al. (2020) Risk Factors for Hospitalization, Mechanical Ventilation, or Death Among 10131 US Veterans With SARS-CoV-2 Infection. *JAMA Network Open* 3(9): e2022310. DOI:

[10.1001/jamanetworkopen.2020.22310](https://doi.org/10.1001/jamanetworkopen.2020.22310).

Ip A, Berry DA, Hansen E, et al. (2020) Hydroxychloroquine and Tocilizumab Therapy in COVID-19 Patients - An Observational Study. *medRxiv*: 2020.05.21.20109207. DOI: [10.1101/2020.05.21.20109207](https://doi.org/10.1101/2020.05.21.20109207).

Islam MZ, Riaz BK, Islam AS, et al. (2020) Risk factors associated with morbidity and mortality outcomes of COVID-19 patients on the 14th and 28th day of the disease course: A retrospective cohort study in Bangladesh. *medRxiv*: 2020.08.17.20176586. DOI: [10.1101/2020.08.17.20176586](https://doi.org/10.1101/2020.08.17.20176586).

Israel A, Schäffer AA, Cicurel A, et al. (2020) Large population study identifies drugs associated with reduced COVID-19 severity. *medRxiv*: 2020.10.13.20211953. DOI: [10.1101/2020.10.13.20211953](https://doi.org/10.1101/2020.10.13.20211953).

Israel A, Feldhamer I, Lahad A, et al. (2020) Smoking and the risk of COVID-19 in a large observational population study. *medRxiv*: 2020.06.01.20118877. DOI: [10.1101/2020.06.01.20118877](https://doi.org/10.1101/2020.06.01.20118877).

Iversen K, Bundgaard H, Hasselbalch RB, et al. (2020) Risk of COVID-19 in health-care workers in Denmark: An observational cohort study. *The Lancet Infectious Diseases* 0(0). DOI: [10.1016/S1473-3099\(20\)30589-2](https://doi.org/10.1016/S1473-3099(20)30589-2).

Izquierdo JL, Almonacid C, Gonzalez Y, et al. (2020) The impact of COVID-19 on patients with asthma. *medRxiv*: 2020.07.24.20161596. DOI: [10.1101/2020.07.24.20161596](https://doi.org/10.1101/2020.07.24.20161596).

Izzi-Engbeaya C, Distaso W, Amin A, et al. (2020) Severe COVID-19 and Diabetes: A Retrospective Cohort Study from Three London Teaching Hospitals. *medRxiv*: 2020.08.07.20160275. DOI: [10.1101/2020.08.07.20160275](https://doi.org/10.1101/2020.08.07.20160275).

Jackson SE, Brown J, Shahab L, et al. (2020) COVID-19, smoking and inequalities: A study of 53 002 adults in the UK. *Tobacco Control*: tobaccocontrol-2020-055933. DOI: [10.1136/tobaccocontrol-2020-055933](https://doi.org/10.1136/tobaccocontrol-2020-055933).

Jafari D, Gandomi A, Makhnevich A, et al. (2021) Trajectories of Hypoxemia & Respiratory System Mechanics of COVID-19 ARDS in the NorthCARDS dataset. DOI: [10.1101/2021.01.26.21250492](https://doi.org/10.1101/2021.01.26.21250492).

- Jakob CEM, Borgmann S, Duygu F, et al. (2020) First results of the ‘Lean European Open Survey on SARS-CoV-2-Infected Patients (LEOSS)’. *Infection*. DOI: [10.1007/s15010-020-01499-0](https://doi.org/10.1007/s15010-020-01499-0).
- Jehi L, Ji X, Milinovich A, et al. (2020) Development and validation of a model for individualized prediction of hospitalization risk in 4,536 patients with COVID-19. *PLOS ONE* Orueta JF (ed.) 15(8): e0237419. DOI: [10.1371/journal.pone.0237419](https://doi.org/10.1371/journal.pone.0237419).
- Jin C, Gu J, Yuan Y, et al. (2020) Treatment of Six COVID-19 Patients with Convalescent Plasma. *medRxiv*: 2020.05.21.20109512. DOI: [10.1101/2020.05.21.20109512](https://doi.org/10.1101/2020.05.21.20109512).
- Jin X, Lian J-S, Hu J-H, et al. (2020) Epidemiological, clinical and virological characteristics of 74 cases of coronavirus-infected disease 2019 (COVID-19) with gastrointestinal symptoms. *Gut* 69(6): 1002–1009. DOI: [10.1136/gutjnl-2020-320926](https://doi.org/10.1136/gutjnl-2020-320926).
- Joubert A, Andry F, Bertolotti A, et al. (2020) *Distinguishing non severe cases of dengue from COVID-19 in the context of co-epidemics: A cohort study in a SARS-CoV-2 testing center on Reunion island*. preprint, October. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.10.20.20214718](https://doi.org/10.1101/2020.10.20.20214718).
- Jun T, Nirenberg S, Kovatch P, et al. (2020) Sex-specificity of mortality risk factors among hospitalized COVID-19 patients in New York City: Prospective cohort study. *medRxiv*: 2020.07.29.20164640. DOI: [10.1101/2020.07.29.20164640](https://doi.org/10.1101/2020.07.29.20164640).
- Kalan ME, Ghobadi H, Taleb ZB, et al. (2020) Descriptive characteristics of hospitalized adult smokers and never-smokers with COVID-19. *Tobacco Induced Diseases* 18(May). DOI: [10.18332/tid/122759](https://doi.org/10.18332/tid/122759).
- Kantele A, Laaveri T, Kareinen L, et al. (2020) SARS-CoV-2 infections among healthcare workers at Helsinki University Hospital, Finland, spring 2020: Serosurvey, symptoms and risk factors. *Travel Med Infect Dis* 39. DOI: [10.1016/j.tmaid.2020.101949](https://doi.org/10.1016/j.tmaid.2020.101949).
- Kara Polat A, Oguz Topal I, Karadag AS, et al. (2020) The impact of COVID-19 in patients with psoriasis: A multicenter study in Istanbul. *Dermatol Ther* NA. DOI: [10.1111/dth.14691](https://doi.org/10.1111/dth.14691).
- Kibler M, Carmona A, Marchandot B, et al. (2020) Risk and severity of COVID-19 and ABO blood group in transcatheter aortic valve patients. *medRxiv*: 2020.06.13.20130211. DOI: [10.1101/2020.06.13.20130211](https://doi.org/10.1101/2020.06.13.20130211).
- Killerby ME (2020) Characteristics Associated with Hospitalization Among Patients with COVID-19 — Metropolitan Atlanta, Georgia, March–April 2020. *MMWR. Morbidity and Mortality Weekly Report* 69. DOI: [10.15585/mmwr.mm6925e1](https://doi.org/10.15585/mmwr.mm6925e1).
- Kim ES, Chin BS, Kang CK, et al. (2020) Clinical Course and Outcomes of Patients with Severe Acute Respiratory Syndrome Coronavirus 2 Infection: A Preliminary Report of the First 28 Patients from the Korean Cohort Study on COVID-19. *Journal of Korean Medical Science* 35(13). DOI: [10.3346/jkms.2020.35.e142](https://doi.org/10.3346/jkms.2020.35.e142).
- Kim H-J, Han D, Kim J-H, et al. (2020) An Easy-to-Use Machine Learning Model to Predict the Prognosis of Patients With COVID-19: Retrospective Cohort Study. *Journal of Medical Internet Research* 22(11): e24225. DOI: [10.2196/24225](https://doi.org/10.2196/24225).
- Kim L, Garg S, O’Halloran A, et al. (2020) Interim Analysis of Risk Factors for Severe Outcomes among a Cohort of Hospitalized Adults Identified through the U.S. Coronavirus Disease 2019 (COVID-19)-Associated Hospitalization Surveillance Network (COVID-NET). *medRxiv*: 2020.05.18.20103390. DOI: [10.1101/2020.05.18.20103390](https://doi.org/10.1101/2020.05.18.20103390).
- Kimmig LM, Wu D, Gold M, et al. (2020) IL-6 Inhibition in Critically Ill COVID-19 Patients Is Associated With Increased Secondary Infections. *Frontiers in Medicine* 7: 583897. DOI: [10.3389/fmed.2020.583897](https://doi.org/10.3389/fmed.2020.583897).
- Kjetland EF, Kalleberg KT, Søråas CL, et al. (2020) Risk factors for community transmission of SARS-CoV-2. A cross-

sectional study in 116,678 people. DOI: [10.1101/2020.12.23.20248514](https://doi.org/10.1101/2020.12.23.20248514).

Klang E, Soffer S, Nadkarni G, et al. (2020) Sex Differences in Age and Comorbidities for COVID-19 Mortality in Urban New York City. *SN Comprehensive Clinical Medicine* 2(9): 1319–1322. DOI: [10.1007/s42399-020-00430-w](https://doi.org/10.1007/s42399-020-00430-w).

Klang E, Kassim G, Soffer S, et al. (n.d.) Morbid Obesity as an Independent Risk Factor for COVID-19 Mortality in Hospitalized Patients Younger than 50. *Obesity n/a(n/a)*. DOI: [10.1002/oby.22913](https://doi.org/10.1002/oby.22913).

Kolin DA, Kulm S and Elemento O (2020) Clinical and Genetic Characteristics of Covid-19 Patients from UK Biobank. *medRxiv*: 2020.05.05.20075507. DOI: [10.1101/2020.05.05.20075507](https://doi.org/10.1101/2020.05.05.20075507).

Kortela E, Kirjavainen V, Ahava MJ, et al. (2020) *Real-life clinical sensitivity of SARS-CoV-2 RT-PCR test in symptomatic patients*. preprint, November. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.11.01.20223107](https://doi.org/10.1101/2020.11.01.20223107).

Kronbichler A, Kresse D, Yoon S, et al. (2020) Asymptomatic patients as a source of COVID-19 infections: A systematic review and meta-analysis. *International Journal of Infectious Diseases* 98: 180–186. DOI: [10.1016/j.ijid.2020.06.052](https://doi.org/10.1016/j.ijid.2020.06.052).

Kuderer NM, Choueiri TK, Shah DP, et al. (2020) Clinical impact of COVID-19 on patients with cancer (CCC19): A cohort study. *The Lancet* 395(10241): 1907–1918. DOI: [10.1016/S0140-6736\(20\)31187-9](https://doi.org/10.1016/S0140-6736(20)31187-9).

Kumar A, Prasad G, Srivastav S, et al. (2020) A Retrospective Study on Efficacy and Safety of Guduchi Ghan Vati for Covid-19 Asymptomatic Patients. *medRxiv*: 2020.07.23.20160424. DOI: [10.1101/2020.07.23.20160424](https://doi.org/10.1101/2020.07.23.20160424).

Kurashima K, Kagiyama N, Ishiguro T, et al. (2020) IgG antibody seroconversion and the clinical progression of COVID-19 pneumonia: A retrospective, cohort study. DOI: [10.1101/2020.07.16.20154088](https://doi.org/10.1101/2020.07.16.20154088).

Lamure S, Duléry R, Blasi RD, et al. (2020) Determinants of outcome in Covid-19 hospitalized patients with lymphoma: A retrospective multicentric cohort study. *EClinicalMedicine* 27. DOI: [10.1016/j.eclinm.2020.100549](https://doi.org/10.1016/j.eclinm.2020.100549).

Lan F-Y, Suharlim C, Kales SN, et al. (2020) Association between SARS-CoV-2 infection, exposure risk and mental health among a cohort of essential retail workers in the United States. *medRxiv*: 2020.06.08.20125120. DOI: [10.1101/2020.06.08.20125120](https://doi.org/10.1101/2020.06.08.20125120).

Lassale C, Gaye B, Hamer M, et al. (2020) Ethnic disparities in hospitalisation for COVID-19 in England: The role of socioeconomic factors, mental health, and inflammatory and pro-inflammatory factors in a community-based cohort study. *Brain, Behavior, and Immunity* 88: 44–49. DOI: [10.1016/j.bbi.2020.05.074](https://doi.org/10.1016/j.bbi.2020.05.074).

Leister I, Ponocny-Seliger E, Kollaritsch H, et al. (2021) Antibody seroprevalence and rate of asymptomatic infections with SARS-CoV-2 in Austrian hospital personnel. DOI: [10.1101/2021.02.01.21250898](https://doi.org/10.1101/2021.02.01.21250898).

Lenka J, Chhabria MS, Sharma N, et al. (2020) Clinical characteristics and outcomes of critically ill patients with COVID-19 in a tertiary community hospital in upstate New York. *medRxiv*: 2020.06.18.20135046. DOI: [10.1101/2020.06.18.20135046](https://doi.org/10.1101/2020.06.18.20135046).

Lewnard JA, Mora AM, Nkwocha O, et al. (2021) Prevalence and clinical profile of SARS-CoV-2 infection among farmworkers in Monterey County, California: June–November, 2020. DOI: [10.1101/2020.12.27.20248894](https://doi.org/10.1101/2020.12.27.20248894).

Lévy Y, Wiedemann A, Hejblum BP, et al. (2020) *CD177, a specific marker of neutrophil activation, is a hallmark of COVID-19 severity and death*. preprint, December. *Allergy; Immunology*. DOI: [10.1101/2020.12.12.20246934](https://doi.org/10.1101/2020.12.12.20246934).

Li F, Cai Y, Gao C, et al. (2020) *Clinical Course And Risk Factors For In-hospital Death In Critical COVID-19 In Wuhan, China*. preprint, September. *Public; Global Health*. DOI: [10.1101/2020.09.26.20189522](https://doi.org/10.1101/2020.09.26.20189522).

Li J, Chen Y, Chen S, et al. (2020) Derivation and validation of a prognostic model for predicting in-hospital mortality in

patients admitted with COVID-19 in Wuhan, China: The PLANS (Platelet Lymphocyte Age Neutrophil Sex) model.

medRxiv: 2020.05.13.20100370. DOI: [10.1101/2020.05.13.20100370](https://doi.org/10.1101/2020.05.13.20100370).

Li J, Li S, Cai Y, et al. (2020) Epidemiological and Clinical Characteristics of 17 Hospitalized Patients with 2019 Novel Coronavirus Infections Outside Wuhan, China. *medRxiv*: 2020.02.11.20022053. DOI: [10.1101/2020.02.11.20022053](https://doi.org/10.1101/2020.02.11.20022053).

Li J, Long X, Zhang Q, et al. (2020) Tobacco smoking confers risk for severe COVID-19 unexplainable by pulmonary imaging. *Journal of Internal Medicine*: joim.13190. DOI: [10.1111/joim.13190](https://doi.org/10.1111/joim.13190).

Li J, Long X, Zhu C, et al. (n.d.) Olfactory Dysfunction in Recovered Coronavirus Disease 2019 (COVID-19) Patients. *Movement Disorders* n/a(n/a). DOI: [10.1002/mds.28172](https://doi.org/10.1002/mds.28172).

Lian J, Jin X, Hao S, et al. (n.d.) Analysis of Epidemiological and Clinical Features in Older Patients With Coronavirus Disease 2019 (COVID-19) Outside Wuhan. *Clinical Infectious Diseases*. DOI: [10.1093/cid/ciaa242](https://doi.org/10.1093/cid/ciaa242).

Liao Y, Feng Y, Wang B, et al. (2020) Clinical Characteristics and Risk factors for developed COVID-19 patients transferring to designated hospital from Jiangnan Fangcang shelter Hospital: A retrospective, observational study. *medRxiv*: 2020.04.21.20074724. DOI: [10.1101/2020.04.21.20074724](https://doi.org/10.1101/2020.04.21.20074724).

Lin KJ, Schneeweiss S, Tesfaye H, et al. (2020) Pharmacotherapy for Hospitalized Patients with COVID-19: Treatment Patterns by Disease Severity. *Drugs* 80: 1961–1972. DOI: [10.1007/s40265-020-01424-7](https://doi.org/10.1007/s40265-020-01424-7).

Liu R, Ming X, Xu O, et al. (2020) Association of Cardiovascular Manifestations with In-hospital Outcomes in Patients with COVID-19: A Hospital Staff Data. *medRxiv*: 2020.02.29.20029348. DOI: [10.1101/2020.02.29.20029348](https://doi.org/10.1101/2020.02.29.20029348).

Liu W, Tao Z-W, Wang L, et al. (2020) Analysis of factors associated with disease outcomes in hospitalized patients with 2019 novel coronavirus disease. *Chinese Medical Journal* 133(9): 1032–1038. DOI: [10.1097/CM9.0000000000000775](https://doi.org/10.1097/CM9.0000000000000775).

Lohia P, Sreeram K, Nguyen P, et al. (2021) Preexisting respiratory diseases and clinical outcomes in COVID-19: A multihospital cohort study on predominantly African American population. *Respir Res* 22. DOI: [10.1186/s12931-021-01647-6](https://doi.org/10.1186/s12931-021-01647-6).

Lopez-Medrano F, Perez-Jacoiste Asin MA, Fernandez-Ruiz M, et al. (2020) *Combination therapy with tocilizumab and corticosteroids for aged patients with severe COVID-19 pneumonia: A single-center retrospective study*. preprint, September. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.09.26.20202283](https://doi.org/10.1101/2020.09.26.20202283).

Louis S, Dhawan A, Newey C, et al. (2020) Continuous Electroencephalography (cEEG) Characteristics and Acute Symptomatic Seizures in COVID-19 Patients. *medRxiv*: 2020.05.26.20114033. DOI: [10.1101/2020.05.26.20114033](https://doi.org/10.1101/2020.05.26.20114033).

Lowe KE, Zein J, Hatipoglu U, et al. (2021) Association of Smoking and Cumulative Pack-Year Exposure With COVID-19 Outcomes in the Cleveland Clinic COVID-19 Registry. *JAMA Intern Med* NA. DOI: [10.1001/jamainternmed.2020.8360](https://doi.org/10.1001/jamainternmed.2020.8360).

Lubetzky M, Aull M, Craig-Shapiro R, et al. (2020) Kidney Allograft Recipients Diagnosed with Coronavirus Disease-2019: A Single Center Report. *medRxiv*: 2020.04.30.20086462. DOI: [10.1101/2020.04.30.20086462](https://doi.org/10.1101/2020.04.30.20086462).

Luo H, Liu S, Wang Y, et al. (2020) Age differences in clinical features and outcomes in patients with COVID-19, Jiangsu, China: A retrospective, multi-center cohort study. *medRxiv*: 2020.06.01.20086025. DOI: [10.1101/2020.06.01.20086025](https://doi.org/10.1101/2020.06.01.20086025).

Luo J, Rizvi H, Preeshagul IR, et al. (2020) COVID-19 in patients with lung cancer. *Annals of Oncology* 31(10): 1386–1396. DOI: [10.1016/j.annonc.2020.06.007](https://doi.org/10.1016/j.annonc.2020.06.007).

Lusignan S de, Dorward J, Correa A, et al. (2020) Risk factors for SARS-CoV-2 among patients in the Oxford Royal College of General Practitioners Research and Surveillance Centre primary care network: A cross-sectional study. *The*

Lancet Infectious Diseases: S1473309920303716. DOI: [10.1016/S1473-3099\(20\)30371-6](https://doi.org/10.1016/S1473-3099(20)30371-6).

Madariaga MLL, Guthmiller J, Schrantz S, et al. (2020) Clinical predictors of donor antibody titer and correlation with recipient antibody response in a COVID-19 convalescent plasma clinical trial. *medRxiv*: 2020.06.21.20132944. DOI: [10.1101/2020.06.21.20132944](https://doi.org/10.1101/2020.06.21.20132944).

Magagnoli J, Narendran S, Pereira F, et al. (2020) Outcomes of Hydroxychloroquine Usage in United States Veterans Hospitalized with COVID-19. *Med*: S2666634020300064. DOI: [10.1016/j.medj.2020.06.001](https://doi.org/10.1016/j.medj.2020.06.001).

Magleby R, Westblade LF, Trzebucki A, et al. (n.d.) Impact of SARS-CoV-2 Viral Load on Risk of Intubation and Mortality Among Hospitalized Patients with Coronavirus Disease 2019. *Clinical Infectious Diseases*. DOI: [10.1093/cid/ciaa851](https://doi.org/10.1093/cid/ciaa851).

Makaronidis J, Mok J, Balogun N, et al. (2020) Seroprevalence of SARS-CoV-2 antibodies in people with an acute loss in their sense of smell and/or taste in a community-based population in London, UK: An observational cohort study. *PLOS Medicine* 17(10): e1003358. DOI: [10.1371/journal.pmed.1003358](https://doi.org/10.1371/journal.pmed.1003358).

Mamtani M, Athavale AM, Abraham M, et al. (2020) ASSOCIATION OF HYPERGLYCEMIA WITH HOSPITAL MORTALITY IN COVID-19 PATIENTS WITHOUT DIABETES: A COHORT STUDY. *medRxiv*: 2020.08.31.20185157. DOI: [10.1101/2020.08.31.20185157](https://doi.org/10.1101/2020.08.31.20185157).

Mancilla-Galindo J, Vera-Zertuche JM, Navarro-Cruz AR, et al. (2020) Development and Validation of the Patient History COVID-19 (PH-Covid19) Scoring System: A Multivariable Prediction Model of Death in Mexican Patients with COVID-19. *medRxiv*: 2020.09.05.20189142. DOI: [10.1101/2020.09.05.20189142](https://doi.org/10.1101/2020.09.05.20189142).

Mangera Z, Lewis A, Hutchinson J, et al. (2017) Smoking prevalence in UK hospital admissions from a national observational study. *European Respiratory Journal* 50(suppl 61). DOI: [10.1183/1393003.congress-2017.PA1268](https://doi.org/10.1183/1393003.congress-2017.PA1268).

Mansour A, Sajjadi-Jazi SM, Kasaeian A, et al. (2020) Clinical characteristics and outcomes of diabetics hospitalized for COVID-19 infection: A single-centered, retrospective, observational study. *EXCLI J* 19: 1533–1543. DOI: [10.17179/excli2020-2988](https://doi.org/10.17179/excli2020-2988).

Maraschini A, Corsi E, Salvatore MA, et al. (2020) Coronavirus and birth in Italy: Results of a national population-based cohort study. *medRxiv*: 2020.06.11.20128652. DOI: [10.1101/2020.06.11.20128652](https://doi.org/10.1101/2020.06.11.20128652).

Marcos M, Belhassen-Garcia M, Puente AS, et al. (2020) Development of a severity of disease score and classification model by machine learning for hospitalized COVID-19 patients. *medRxiv*: 2020.07.13.20150177. DOI: [10.1101/2020.07.13.20150177](https://doi.org/10.1101/2020.07.13.20150177).

Martinez-Lacalzada M, Viteri-Noël LA, Manzano L, et al. (2020) Predicting critical illness on initial diagnosis of COVID-19: Development and validation of the PRIORITY model for outpatient applicability. DOI: [10.1101/2020.11.27.20237966](https://doi.org/10.1101/2020.11.27.20237966).

Martinez-Portilla RJ, Sotiriadis A, Torres-Torres J, et al. (2020) Risk factors for mortality in pregnant women with SARS-CoV-2 infection. *medRxiv*: 2020.05.31.20107276. DOI: [10.1101/2020.05.31.20107276](https://doi.org/10.1101/2020.05.31.20107276).

Martinez-Resendez MF, Castilleja-Leal F, Torres-Quintanilla A, et al. (2020) Initial experience in Mexico with convalescent plasma in COVID-19 patients with severe respiratory failure, a retrospective case series. *medRxiv*: 2020.07.14.20144469. DOI: [10.1101/2020.07.14.20144469](https://doi.org/10.1101/2020.07.14.20144469).

Martini F, D'Alessio A, Bracchi F, et al. (2020) On Cancer, COVID-19, and CT Scans: A Monocentric Retrospective Study. *Journal of Clinical Medicine* 9(12): 3935. DOI: [10.3390/jcm9123935](https://doi.org/10.3390/jcm9123935).

Martin-Jimenez P, Munoz-Garcia MI, Seoane D, et al. (2020) Cognitive impairment is a common comorbidity in COVID-19

- deceased patients. A hospital-based retrospective cohort study. *medRxiv*: 2020.06.08.20125872. DOI: [10.1101/2020.06.08.20125872](https://doi.org/10.1101/2020.06.08.20125872).
- Maucourant C, Filipovic I, Ponzetta A, et al. (2020) Natural killer cell activation related to clinical outcome of COVID-19. *medRxiv*: 2020.07.07.20148478. DOI: [10.1101/2020.07.07.20148478](https://doi.org/10.1101/2020.07.07.20148478).
- Márquez-Salinas A, Fermín-Martínez CA, Antonio-Villa NE, et al. (2020) *Adaptive metabolic and inflammatory responses identified using accelerated aging metrics are linked to adverse outcomes in severe SARS-CoV-2 infection*. preprint, November. *Geriatric Medicine*. DOI: [10.1101/2020.11.03.20225375](https://doi.org/10.1101/2020.11.03.20225375).
- McGrail DE and Edwards D (2020) COVID-19 Case Series at UnityPoint Health St. Lukes Hospital in Cedar Rapids, IA. *medRxiv*: 2020.07.17.20156521. DOI: [10.1101/2020.07.17.20156521](https://doi.org/10.1101/2020.07.17.20156521).
- McQueenie R, Foster H, Jani BD, et al. (2020) Multimorbidity, Polypharmacy, and COVID-19 infection within the UK Biobank cohort. *medRxiv*: 2020.06.10.20127563. DOI: [10.1101/2020.06.10.20127563](https://doi.org/10.1101/2020.06.10.20127563).
- Meini S, Fortini A, Andreini R, et al. (n.d.) The Paradox of the Low Prevalence of Current Smokers Among Covid-19 Patients Hospitalized in Non-Intensive Care Wards: Results From an Italian Multicenter Case-Control Study. *Nicotine & Tobacco Research*. DOI: [10.1093/ntr/ntaa188](https://doi.org/10.1093/ntr/ntaa188).
- Mejia-Vilet JM, Cordova-Sanchez BM, Fernandez-Camargo D, et al. (2020) A Risk Score to Predict Admission to Intensive Care Unit in Patients With COVID-19: The ABC-GOALS Score. *medRxiv*: 2020.05.12.20099416. DOI: [10.1101/2020.05.12.20099416](https://doi.org/10.1101/2020.05.12.20099416).
- Melo AC de, Thuler LCS, Silva JL da, et al. (2020) Cancer inpatient with COVID-19: A report from the Brazilian National Cancer Institute. *medRxiv*: 2020.06.27.20141499. DOI: [10.1101/2020.06.27.20141499](https://doi.org/10.1101/2020.06.27.20141499).
- Mendes A, Herrmann FR, Genton L, et al. (2021) Incidence, characteristics and clinical relevance of acute stroke in old patients hospitalized with COVID-19. *BMC geriatr* 21. DOI: [10.1186/s12877-021-02006-2](https://doi.org/10.1186/s12877-021-02006-2).
- Mendy A, Apewokin S, Wells AA, et al. (2020) Factors Associated with Hospitalization and Disease Severity in a Racially and Ethnically Diverse Population of COVID-19 Patients. *medRxiv*: 2020.06.25.20137323. DOI: [10.1101/2020.06.25.20137323](https://doi.org/10.1101/2020.06.25.20137323).
- Merkely B, Szabó AJ, Kosztin A, et al. (2020) Novel coronavirus epidemic in the Hungarian population, a cross-sectional nationwide survey to support the exit policy in Hungary. *GeroScience*. DOI: [10.1007/s11357-020-00226-9](https://doi.org/10.1007/s11357-020-00226-9).
- Merzon E, Tworowski D, Gorohovski A, et al. (2020) Low plasma 25(OH) vitamin D3 level is associated with increased risk of COVID-19 infection: An Israeli population-based study. *medRxiv*: 2020.07.01.20144329. DOI: [10.1101/2020.07.01.20144329](https://doi.org/10.1101/2020.07.01.20144329).
- Miyara M, Tubach F, Martinez V, et al. (2020) Low rate of daily smokers in patients with symptomatic COVID-19. *medRxiv*: 2020.06.10.20127514. DOI: [10.1101/2020.06.10.20127514](https://doi.org/10.1101/2020.06.10.20127514).
- Mo P, Xing Y, Xiao Y, et al. (n.d.) Clinical characteristics of refractory COVID-19 pneumonia in Wuhan, China. *Clinical Infectious Diseases*. DOI: [10.1093/cid/ciaa270](https://doi.org/10.1093/cid/ciaa270).
- Modrák M, Bürkner P-C, Sieger T, et al. (2020) Detailed disease progression of 213 patients hospitalized with Covid-19 in the Czech Republic: An exploratory analysis. DOI: [10.1101/2020.12.03.20239863](https://doi.org/10.1101/2020.12.03.20239863).
- Mohamed-Hussein A, Galal I, Saad M, et al. (2020) Post-COVID-19 Functional Status: Relation to age, smoking, hospitalization and comorbidities. *medRxiv*: 2020.08.26.20182618. DOI: [10.1101/2020.08.26.20182618](https://doi.org/10.1101/2020.08.26.20182618).

- Mohamud AY, Griffith B, Rehman M, et al. (2020) Intraluminal Carotid Artery Thrombus in COVID-19: Another Danger of Cytokine Storm? *American Journal of Neuroradiology*. DOI: [10.3174/ajnr.A6674](https://doi.org/10.3174/ajnr.A6674).
- Molenaar NM, Rommel A-S, Witte L de, et al. (2021) Seroprevalence of SARS-CoV-2 during pregnancy and associated outcomes: Results from an ongoing prospective cohort study, New York City. DOI: [10.1101/2021.02.01.21250943](https://doi.org/10.1101/2021.02.01.21250943).
- Monteiro ACC, Suri R, Emeruwa IO, et al. (2020) Obesity and Smoking as Risk Factors for Invasive Mechanical Ventilation in COVID-19: A Retrospective, Observational Cohort Study. *medRxiv*: 2020.08.12.20173849. DOI: [10.1101/2020.08.12.20173849](https://doi.org/10.1101/2020.08.12.20173849).
- Mora AM, Lewnard JA, Kogut K, et al. (2021) Risk factors for SARS-CoV-2 infection among farmworkers in Monterey County, California. DOI: [10.1101/2021.02.01.21250963](https://doi.org/10.1101/2021.02.01.21250963).
- Morshed MS, Mosabbir AA, Chowdhury P, et al. (2020) Clinical manifestations of patients with Coronavirus Disease 2019 (COVID- 19) attending at hospitals in Bangladesh. *medRxiv*: 2020.07.30.20165100. DOI: [10.1101/2020.07.30.20165100](https://doi.org/10.1101/2020.07.30.20165100).
- Motta JK, Ogunnaike RO, Shah R, et al. (2020) Clinical Outcomes With the Use of Prophylactic Versus Therapeutic Anticoagulation in COVID-19. *medRxiv*: 2020.07.20.20147769. DOI: [10.1101/2020.07.20.20147769](https://doi.org/10.1101/2020.07.20.20147769).
- Muñoz FJT, García-Guijarro E, García-Domingo P, et al. (2020) A safe protocol to identify low risk patients with COVID-19 pneumonia for outpatient management. *medRxiv*: 2020.12.15.20229286. DOI: [10.1101/2020.12.15.20229286](https://doi.org/10.1101/2020.12.15.20229286).
- Murray E (2020) *Causation in Smoking and COVID-19*. Available at: <https://twitter.com/EpiEllie/status/1258607277357006849?s=20>.
- Mutambudzi M, Niedzwiedz CL, Macdonald EB, et al. (2020) Occupation and risk of severe COVID-19: Prospective cohort study of 120,075 UK Biobank participants. *medRxiv*: 2020.05.22.20109892. DOI: [10.1101/2020.05.22.20109892](https://doi.org/10.1101/2020.05.22.20109892).
- Nezhadmoghadam F and Tamez-Peña J (2021) Unsupervised Discovery of Risk Profiles on Negative and Positive COVID-19 Hospitalized Patients. DOI: [10.1101/2020.12.30.20248908](https://doi.org/10.1101/2020.12.30.20248908).
- Nguyen AB, Upadhyay GA, Chung B, et al. (2020) Outcomes and Cardiovascular Comorbidities in a Predominantly African-American Population with COVID-19. *medRxiv*: 2020.06.28.20141929. DOI: [10.1101/2020.06.28.20141929](https://doi.org/10.1101/2020.06.28.20141929).
- Nicholson CJ, Wooster L, Sigurslid HH, et al. (2020) Estimating Risk of Mechanical Ventilation and Mortality Among Adult COVID-19 patients Admitted to Mass General Brigham: The VICE and DICE Scores. *medRxiv*: 2020.09.14.20194670. DOI: [10.1101/2020.09.14.20194670](https://doi.org/10.1101/2020.09.14.20194670).
- Niedzwiedz CL, O'Donnell CA, Jani BD, et al. (2020) Ethnic and socioeconomic differences in SARS-CoV-2 infection: Prospective cohort study using UK Biobank. *BMC Medicine* 18(1): 160. DOI: [10.1186/s12916-020-01640-8](https://doi.org/10.1186/s12916-020-01640-8).
- Nuño M, García Y, Rajasekar G, et al. (2021) COVID-19 Hospitalizations in Five California Hospitals. DOI: [10.1101/2021.01.29.21250788](https://doi.org/10.1101/2021.01.29.21250788).
- Núñez-Gil IJJ, Fernández-Ortiz A, Eid CM, et al. (2020) Underlying heart diseases and acute COVID-19 outcomes. *Cardiology Journal* 0(0). DOI: [10.5603/cj.a2020.0183](https://doi.org/10.5603/cj.a2020.0183).
- Oakes JM, Fuchs RM, Gardner JD, et al. (2018) Nicotine and the renin-angiotensin system. *Am. J. Physiol. - Regul. Integr. Comp. Physiol* 315: 895–906. DOI: [10.1152/ajpregu.00099.2018](https://doi.org/10.1152/ajpregu.00099.2018).
- Odani S (2018) Tobacco Product Use Among Military Veterans — United States, 2010–2015. *MMWR. Morbidity and Mortality Weekly Report* 67. DOI: [10.15585/mmwr.mm6701a2](https://doi.org/10.15585/mmwr.mm6701a2).
- O’Gallagher K, Shek A, Bean DM, et al. (2020) *Pre-existing cardiovascular disease rather than cardiovascular risk factors*

- drives mortality in COVID-19*. preprint, December. *Cardiovascular Medicine*. DOI: [10.1101/2020.12.02.20242933](https://doi.org/10.1101/2020.12.02.20242933).
- Olivares F, Munoz D, Fica A, et al. (2020) Covid-19 in Chile. The experience of a Regional reference Center. Preliminary report. *medRxiv*: 2020.06.14.20130898. DOI: [10.1101/2020.06.14.20130898](https://doi.org/10.1101/2020.06.14.20130898).
- Oliveira E, Parikh A, Lopez-Ruiz A, et al. (2020) ICU Outcomes and Survival in Patients with Severe COVID-19 in the Largest Health Care System in Central Florida. *medRxiv*: 2020.08.25.20181909. DOI: [10.1101/2020.08.25.20181909](https://doi.org/10.1101/2020.08.25.20181909).
- Omrani AS, Almaslamani MA, Daghfal J, et al. (2020) The First Consecutive 5000 Patients with Coronavirus Disease 2019 from Qatar; a Nation-wide Cohort Study. *medRxiv*: 2020.07.15.20154690. DOI: [10.1101/2020.07.15.20154690](https://doi.org/10.1101/2020.07.15.20154690).
- O'Reilly GM, Mitchell RD, Mitra B, et al. (2020) Epidemiology and clinical features of emergency department patients with suspected and confirmed COVID-19: A multisite report from the COVID-19 Emergency Department Quality Improvement Project for July 2020 (COVED-3). *Emergency Medicine Australasia*: 1742–6723.13651. DOI: [10.1111/1742-6723.13651](https://doi.org/10.1111/1742-6723.13651).
- Organisation WH (n.d.) *Laboratory testing for 2019 novel coronavirus (2019-nCoV) in suspected human cases*. Available at: <https://www.who.int/publications-detail-redirect/10665-331501>.
- Ouyang J, Shan X, Wang X, et al. (2020) Clinical characteristics of COVID-19 and the model for predicting the occurrence of critically ill patients: A retrospective cohort study. *medRxiv*: 2020.08.13.20173799. DOI: [10.1101/2020.08.13.20173799](https://doi.org/10.1101/2020.08.13.20173799).
- Palaiodimos L, Kokkinidis DG, Li W, et al. (2020) Severe obesity, increasing age and male sex are independently associated with worse in-hospital outcomes, and higher in-hospital mortality, in a cohort of patients with COVID-19 in the Bronx, New York. *Metabolism* 108: 154262. DOI: [10.1016/j.metabol.2020.154262](https://doi.org/10.1016/j.metabol.2020.154262).
- Paleiron N, Mayet A, Marbac V, et al. (2021) Impact of Tobacco Smoking on the risk of COVID-19. A large scale retrospective cohort study. *Nicotine Tob Res NA*. DOI: [10.1093/ntr/ntab004](https://doi.org/10.1093/ntr/ntab004).
- Pan A, Khan O, Meeks J, et al. (2020) Disparities in COVID-19 Hospitalizations and Mortality among Black and Hispanic Patients: Cross-Sectional Analysis from the Greater Houston Metropolitan Area. *medRxiv*: 2020.08.19.20177956. DOI: [10.1101/2020.08.19.20177956](https://doi.org/10.1101/2020.08.19.20177956).
- Pandolfi L, Fossali T, Frangipane V, et al. (2020) Broncho-alveolar inflammation in COVID-19 patients: A correlation with clinical outcome. *medRxiv*: 2020.07.17.20155978. DOI: [10.1101/2020.07.17.20155978](https://doi.org/10.1101/2020.07.17.20155978).
- Park BE, Lee JH, Park HK, et al. (2021) Impact of Cardiovascular Risk Factors and Cardiovascular Diseases on Outcomes in Patients Hospitalized with COVID-19 in Daegu Metropolitan City. *J Korean Med Sci* 36. DOI: [10.3346/jkms.2021.36.e15](https://doi.org/10.3346/jkms.2021.36.e15).
- Parra-Bracamonte GM, Lopez-Villalobos N and Parra-Bracamonte FE (2020) Clinical characteristics and risk factors for mortality of patients with COVID-19 in a large data set from Mexico. *Annals of Epidemiology* 52: 93–98.e2. DOI: [10.1016/j.annepidem.2020.08.005](https://doi.org/10.1016/j.annepidem.2020.08.005).
- Parrotta E, Kister I, Charvet L, et al. (2020) COVID-19 outcomes in MS: Observational study of early experience from NYU Multiple Sclerosis Comprehensive Care Center. *Neurology - Neuroimmunology Neuroinflammation* 7(5): e835. DOI: [10.1212/NXI.0000000000000835](https://doi.org/10.1212/NXI.0000000000000835).
- Patanavanich R and Glantz SA (2020) Smoking is Associated with COVID-19 Progression: A Meta-Analysis. *medRxiv*. DOI: [10.14171/j.2095-5944.sg.2014.02.004](https://doi.org/10.14171/j.2095-5944.sg.2014.02.004).
- Patel M, Chowdhury J, Mills N, et al. (2020) ROX Index Predicts Intubation in Patients with COVID-19 Pneumonia and

Moderate to Severe Hypoxemic Respiratory Failure Receiving High Flow Nasal Therapy. *medRxiv*: 2020.06.30.20143867. DOI: [10.1101/2020.06.30.20143867](https://doi.org/10.1101/2020.06.30.20143867).

Perico L, Tomasoni S, Peracchi T, et al. (2020) COVID-19 and lombardy: TESTing the impact of the first wave of the pandemic. *EBioMedicine* 61. DOI: [10.1016/j.ebiom.2020.103069](https://doi.org/10.1016/j.ebiom.2020.103069).

Perrone F, Piccirillo MC, Ascierio PA, et al. (2020) Tocilizumab for patients with COVID-19 pneumonia. The TOCOVID-19 prospective phase 2 trial. *medRxiv*: 2020.06.01.20119149. DOI: [10.1101/2020.06.01.20119149](https://doi.org/10.1101/2020.06.01.20119149).

Peters EJ, Collard D, Assen S van, et al. (2020) Outcomes of Persons With COVID-19 in Hospitals With and Without Standard Treatment With (Hydroxy)chloroquine. *medRxiv*: 2020.08.14.20173369. DOI: [10.1101/2020.08.14.20173369](https://doi.org/10.1101/2020.08.14.20173369).

Petrilli CM, Jones SA, Yang J, et al. (2020) Factors associated with hospital admission and critical illness among 5279 people with coronavirus disease 2019 in New York City: Prospective cohort study. *BMJ* 369. DOI: [10.1136/bmj.m1966](https://doi.org/10.1136/bmj.m1966).

Philipose Z, Smati N, Wong CSJ, et al. (2020) Obesity, old age and frailty are the true risk factors for COVID-19 mortality and not chronic disease or ethnicity in Croydon. *medRxiv*: 2020.08.12.20156257. DOI: [10.1101/2020.08.12.20156257](https://doi.org/10.1101/2020.08.12.20156257).

Polubriaginof F, Salmasian H, Albert DA, et al. (2018) Challenges with Collecting Smoking Status in Electronic Health Records. *AMIA Annual Symposium Proceedings* 2017: 1392–1400. Available at: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5977725/> (accessed 25 September 2020).

Pongpirul WA, Wiboonchutikul S, Charoenpong L, et al. (2020) Clinical course and potential predicting factors of pneumonia of adult patients with coronavirus disease 2019 (COVID-19): A retrospective observational analysis of 193 confirmed cases in Thailand. *medRxiv*: 2020.06.24.20139642. DOI: [10.1101/2020.06.24.20139642](https://doi.org/10.1101/2020.06.24.20139642).

Prats-Uribe A, Tobed M, Villacampa JM, et al. (2021) Timing of elective tracheotomy and duration of mechanical ventilation amongst patients admitted to intensive care with severe COVID-19: A multicentre prospective cohort study. DOI: [10.1101/2021.01.22.21249651](https://doi.org/10.1101/2021.01.22.21249651).

Qi D, Yan X, Tang X, et al. (2020) Epidemiological and clinical features of 2019-nCoV acute respiratory disease cases in Chongqing municipality, China: A retrospective, descriptive, multiple-center study. *medRxiv*: 2020.03.01.20029397. DOI: [10.1101/2020.03.01.20029397](https://doi.org/10.1101/2020.03.01.20029397).

Qu J, Chang LK, Tang X, et al. (2020) Clinical characteristics of COVID-19 and its comparison with influenza pneumonia. *Acta Clinica Belgica* 0(0): 1–9. DOI: [10.1080/17843286.2020.1798668](https://doi.org/10.1080/17843286.2020.1798668).

Quan D, Luna Wong L, Shallal A, et al. (2021) Impact of Race and Socioeconomic Status on Outcomes in Patients Hospitalized with COVID-19. *J Gen Intern Med* NA. DOI: [10.1007/s11606-020-06527-1](https://doi.org/10.1007/s11606-020-06527-1).

Raines AM, Tock JL, McGrew SJ, et al. (2021) Correlates of death among SARS-CoV-2 positive veterans: The contribution of lifetime tobacco use. *Addictive Behaviors* 113: 106692. DOI: [10.1016/j.addbeh.2020.106692](https://doi.org/10.1016/j.addbeh.2020.106692).

Raisi-Estabragh Z, McCracken C, Bethell MS, et al. (n.d.) Greater risk of severe COVID-19 in Black, Asian and Minority Ethnic populations is not explained by cardiometabolic, socioeconomic or behavioural factors, or by 25(OH)-vitamin D status: Study of 1326 cases from the UK Biobank. *Journal of Public Health*. DOI: [10.1093/pubmed/fdaa095](https://doi.org/10.1093/pubmed/fdaa095).

Rajter JC, Sherman M, Fatteh N, et al. (2020) ICON (Ivermectin in COvid Nineteen) study: Use of Ivermectin is Associated with Lower Mortality in Hospitalized Patients with COVID19. *medRxiv*: 2020.06.06.20124461. DOI: [10.1101/2020.06.06.20124461](https://doi.org/10.1101/2020.06.06.20124461).

Ramachandran P, Kathirvelu B, Chakraborti A, et al. (2020) COVID-19 in Cancer Patients From New York City: A

Comparative Single Center Retrospective Analysis. *Cancer Control* 27(1): 107327482096045. DOI: [10.1177/1073274820960457](https://doi.org/10.1177/1073274820960457).

Ramlall V, Thangaraj P, Meydan C, et al. (2020) Identification of Immune complement function as a determinant of adverse SARS-CoV-2 infection outcome. *medRxiv*: 2020.05.05.20092452. DOI: [10.1101/2020.05.05.20092452](https://doi.org/10.1101/2020.05.05.20092452).

Rashid M, Wu J, Timmis A, et al. (2020) Clinical Characteristics and Outcomes of COVID-19 Positive Acute Coronary Syndrome Patients; a multisource Electronic Healthcare Records Study from England. *medRxiv*: 2020.08.20.20175091. DOI: [10.1101/2020.08.20.20175091](https://doi.org/10.1101/2020.08.20.20175091).

REACT Study Investigators, Riley S, Ainslie KEC, et al. (2020) Community prevalence of SARS-CoV-2 virus in England during May 2020: REACT study. DOI: [10.1101/2020.07.10.20150524](https://doi.org/10.1101/2020.07.10.20150524).

Regina J, Papadimitriou-Olivgeris M, Burger R, et al. (2020) Epidemiology, risk factors and clinical course of SARS-CoV-2 infected patients in a Swiss university hospital: An observational retrospective study. *medRxiv*: 2020.05.11.20097741. DOI: [10.1101/2020.05.11.20097741](https://doi.org/10.1101/2020.05.11.20097741).

Reiter T, Pajenda S, Wagner L, et al. (2020) Covid-19 serology in nephrology health care workers. *medRxiv*: 2020.07.21.20136218. DOI: [10.1101/2020.07.21.20136218](https://doi.org/10.1101/2020.07.21.20136218).

Ren HG, Guo X, Tu L, et al. (2020) *Clinical Characteristics and Risk Factors for Myocardial Injury and Arrhythmia in COVID-19 patients*. preprint, December. Cardiovascular Medicine. DOI: [10.1101/2020.11.30.20190926](https://doi.org/10.1101/2020.11.30.20190926).

Ren HG, Guo X, Blighe K, et al. (2020) Risk Factors for ICU Admission, Mechanical Ventilation and Mortality in Hospitalized Patients with COVID-19 in Hubei, China. *medRxiv*: 2020.08.31.20184952. DOI: [10.1101/2020.08.31.20184952](https://doi.org/10.1101/2020.08.31.20184952).

Rentsch CT, Kidwai-Khan F, Tate JP, et al. (2020) Covid-19 Testing, Hospital Admission, and Intensive Care Among 2,026,227 United States Veterans Aged 54-75 Years. *medRxiv*: 2020.04.09.20059964. DOI: [10.1101/2020.04.09.20059964](https://doi.org/10.1101/2020.04.09.20059964).

Rentsch CT, Beckman JA, Tomlinson L, et al. (2020) *Early initiation of prophylactic anticoagulation for prevention of COVID-19 mortality: A nationwide cohort study of hospitalized patients in the United States*. preprint, December. Infectious Diseases (except HIV/AIDS). DOI: [10.1101/2020.12.09.20246579](https://doi.org/10.1101/2020.12.09.20246579).

Rica R de la, Borges M, Aranda M, et al. (2020) Low albumin levels are associated with poorer outcomes in a case series of COVID-19 patients in Spain: A retrospective cohort study. *medRxiv*: 2020.05.07.20094987. DOI: [10.1101/2020.05.07.20094987](https://doi.org/10.1101/2020.05.07.20094987).

Richard A, Wisniak A, Perez-Saez J, et al. (2020) Seroprevalence of anti-SARS-CoV-2 IgG antibodies, risk factors for infection and associated symptoms in Geneva, Switzerland: A population-based study. *medRxiv*: 2020.12.16.20248180. DOI: [10.1101/2020.12.16.20248180](https://doi.org/10.1101/2020.12.16.20248180).

Richardson S, Hirsch JS, Narasimhan M, et al. (2020) Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* 323(20): 2052–2059. DOI: [10.1001/jama.2020.6775](https://doi.org/10.1001/jama.2020.6775).

Rimland CA, Morgan CE, Bell GJ, et al. (2020) Clinical characteristics and early outcomes in patients with COVID-19 treated with tocilizumab at a United States academic center. *medRxiv*: 2020.05.13.20100404. DOI: [10.1101/2020.05.13.20100404](https://doi.org/10.1101/2020.05.13.20100404).

- Rizzo S, Chawla D, Zalocusky K, et al. (2020) Descriptive epidemiology of 16,780 hospitalized COVID-19 patients in the United States. *medRxiv*: 2020.07.17.20156265. DOI: [10.1101/2020.07.17.20156265](https://doi.org/10.1101/2020.07.17.20156265).
- Robilotti EV, Babady NE, Mead PA, et al. (2020) Determinants of Severity in Cancer Patients with COVID-19 Illness. *medRxiv*: 2020.05.04.20086322. DOI: [10.1101/2020.05.04.20086322](https://doi.org/10.1101/2020.05.04.20086322).
- Robinson L, Wang L, Fu X, et al. (2021) COVID-19 severity in asthma patients: A multi-center matched cohort study. *Journal of Allergy and Clinical Immunology* 147(2): AB241. DOI: [10.1016/j.jaci.2020.12.026](https://doi.org/10.1016/j.jaci.2020.12.026).
- Roederer T, Mollo B, Vincent C, et al. (2021) Seroprevalence and risk factors of exposure to COVID-19 in homeless people in Paris, France: A cross-sectional study. *The Lancet Public Health*: S2468266721000013. DOI: [10.1016/S2468-2667\(21\)00001-3](https://doi.org/10.1016/S2468-2667(21)00001-3).
- Romão VC, Oliveira-Ramos F, Cruz-Machado AR, et al. (2020) A COVID-19 outbreak in a rheumatology department upon the early days of the pandemic. *medRxiv*: 2020.06.05.20107011. DOI: [10.1101/2020.06.05.20107011](https://doi.org/10.1101/2020.06.05.20107011).
- Rossi B, Nguyen LS, Zimmermann P, et al. (2020) Effect of tocilizumab in hospitalized patients with severe pneumonia COVID-19: A cohort study. *medRxiv*: 2020.06.06.20122341. DOI: [10.1101/2020.06.06.20122341](https://doi.org/10.1101/2020.06.06.20122341).
- Rowlands AV, Gillies C, Chudasama Y, et al. (2020) Association of working shifts, inside and outside of healthcare, with risk of severe COVID-19: An observational study. DOI: [10.1101/2020.12.16.20248243](https://doi.org/10.1101/2020.12.16.20248243).
- Rubio-Rivas M, Ronda M, Padullés A, et al. (2020) Beneficial Effect of Corticosteroids in Preventing Mortality in Patients Receiving Tocilizumab to Treat Severe COVID-19 Illness. *medRxiv*: 2020.08.31.20182428. DOI: [10.1101/2020.08.31.20182428](https://doi.org/10.1101/2020.08.31.20182428).
- Russell B, Moss C, Papa S, et al. (2020) Factors Affecting COVID-19 Outcomes in Cancer Patients: A First Report From Guy's Cancer Center in London. *Frontiers in Oncology* 10: 1279. DOI: [10.3389/fonc.2020.01279](https://doi.org/10.3389/fonc.2020.01279).
- Saeed GA, Al Helali AA, Almazrouei S, et al. (2020) Chest CT features of COVID-19 in the region of Abu Dhabi, UAE- A single institute study. preprint, November. Radiology; Imaging. DOI: [10.1101/2020.11.14.20229096](https://doi.org/10.1101/2020.11.14.20229096).
- Salama C, Han J, Yau L, et al. (2021) Tocilizumab in Patients Hospitalized with Covid-19 Pneumonia. *New England Journal of Medicine* 384(1): 20–30. DOI: [10.1056/NEJMoa2030340](https://doi.org/10.1056/NEJMoa2030340).
- Salerno S, Zhao Z, Sankar SP, et al. (2020) Understanding the patterns of repeated testing for COVID-19: Association with patient characteristics and outcomes. *medRxiv*: 2020.07.26.20162453. DOI: [10.1101/2020.07.26.20162453](https://doi.org/10.1101/2020.07.26.20162453).
- Salton F, Confalonieri P, Santus P, et al. (2020) Prolonged low-dose methylprednisolone in patients with severe COVID-19 pneumonia. *medRxiv*: 2020.06.17.20134031. DOI: [10.1101/2020.06.17.20134031](https://doi.org/10.1101/2020.06.17.20134031).
- Sami R, Soltaninejad F, Amra B, et al. (2020) A one-year hospital-based prospective COVID-19 open-cohort in the Eastern Mediterranean region: The Khorshid COVID Cohort (KCC) study. *medRxiv*: 2020.05.11.20096727. DOI: [10.1101/2020.05.11.20096727](https://doi.org/10.1101/2020.05.11.20096727).
- Santos C, Rhee Y, Hollinger E, et al. (2020) Comparative Incidence and Outcomes of COVID-19 in Kidney or Kidney-Pancreas Transplant Recipients Versus Kidney or Kidney-Pancreas Waitlisted Patients: A Pilot Study. *medRxiv*: 2020.07.20.20157990. DOI: [10.1101/2020.07.20.20157990](https://doi.org/10.1101/2020.07.20.20157990).
- Saurabh S, Verma MK, Gautam V, et al. (2021) Tobacco, alcohol use and other risk factors for developing symptomatic COVID-19 vs asymptomatic SARS-CoV-2 infection: A case-control study from western Rajasthan, India. *Trans R Soc Trop Med Hyg* NA. DOI: [10.1093/trstmh/traa172](https://doi.org/10.1093/trstmh/traa172).

- Savarraj JP, Burkett AB, Hinds SN, et al. (n.d.) Three-month outcomes in hospitalized COVID-19 patients.: 7. DOI: [10.1101/2020.10.16.20211029](https://doi.org/10.1101/2020.10.16.20211029).
- Schneeweiss MC, Leonard S, Weckstein A, et al. (2020) Renin-Angiotensin-Aldosterone-System inhibitor use in patients with COVID-19 infection and prevention of serious events: A cohort study in commercially insured patients in the US. *medRxiv*: 2020.07.22.20159855. DOI: [10.1101/2020.07.22.20159855](https://doi.org/10.1101/2020.07.22.20159855).
- Schoeman D and Fielding BC (2019) Coronavirus envelope protein: Current knowledge. *Virology Journal* 16(1): 69. DOI: [10.1186/s12985-019-1182-0](https://doi.org/10.1186/s12985-019-1182-0).
- Schubl SD, Figueroa C, Palma AM, et al. (2020) Risk Factors for SARS-CoV-2 Seropositivity in a Health Care Worker Population. DOI: [10.1101/2020.12.17.20248430](https://doi.org/10.1101/2020.12.17.20248430).
- Senkal N (2020) Chronic ACE Inhibitor use is Associated with Decreased Odds of Severe Disease in Patients with COVID-19. *The Anatolian Journal of Cardiology*. DOI: [10.14744/AnatoJCardiol.2020.57431](https://doi.org/10.14744/AnatoJCardiol.2020.57431).
- Serling-Boyd N, D’Silva KM, Hsu TY, et al. (2020) Coronavirus disease 2019 outcomes among patients with rheumatic diseases 6 months into the pandemic. *Annals of the Rheumatic Diseases*: annrheumdis–2020–219279. DOI: [10.1136/annrheumdis-2020-219279](https://doi.org/10.1136/annrheumdis-2020-219279).
- Shade JK, Doshi AN, Sung E, et al. (2021) COVID-HEART: Development and Validation of a Multi-Variable Model for Real-Time Prediction of Cardiovascular Complications in Hospitalized Patients with COVID-19. DOI: [10.1101/2021.01.03.21249182](https://doi.org/10.1101/2021.01.03.21249182).
- Shah SJ, Barish PN, Prasad PA, et al. (2020) Clinical features, diagnostics, and outcomes of patients presenting with acute respiratory illness: A comparison of patients with and without COVID-19. *medRxiv*: 2020.05.02.20082461. DOI: [10.1101/2020.05.02.20082461](https://doi.org/10.1101/2020.05.02.20082461).
- Shahab L, Brose LS and West R (2013) Novel delivery systems for nicotine replacement therapy as an aid to smoking cessation and for harm reduction: Rationale, and evidence for advantages over existing systems. *CNS Drugs* 27(12): 1007–1019. DOI: [10.1007/s40263-013-0116-4](https://doi.org/10.1007/s40263-013-0116-4).
- Sharma AK, Ahmed A, Baig VN, et al. (2020) Characteristics and Outcomes of Hospitalized Young Adults with Mild to Moderate Covid-19 at a University Hospital in India. *medRxiv*: 2020.06.02.20106310. DOI: [10.1101/2020.06.02.20106310](https://doi.org/10.1101/2020.06.02.20106310).
- Shekhar R, Sheikh AB, Upadhyay S, et al. (2020) Early experience with COVID-19 patients at academic hospital in Southwestern United States. *Infectious Diseases* 52(8): 596–599. DOI: [10.1080/23744235.2020.1774645](https://doi.org/10.1080/23744235.2020.1774645).
- Shi H, Zuo Y, Yalavarthi S, et al. (2020) Neutrophil calprotectin identifies severe pulmonary disease in COVID-19. *medRxiv*: 2020.05.06.20093070. DOI: [10.1101/2020.05.06.20093070](https://doi.org/10.1101/2020.05.06.20093070).
- Shi P, Ren G, Yang J, et al. (2020) Clinical characteristics of imported and second-generation COVID-19 cases outside Wuhan, China: A multicenter retrospective study. *medRxiv*: 2020.04.19.20071472. DOI: [10.1101/2020.04.19.20071472](https://doi.org/10.1101/2020.04.19.20071472).
- Shi Q, Zhao K, Yu J, et al. (2020) Clinical characteristics of 101 COVID-19 nonsurvivors in Wuhan, China: A retrospective study. *medRxiv*: 2020.03.04.20031039. DOI: [10.1101/2020.03.04.20031039](https://doi.org/10.1101/2020.03.04.20031039).
- Shi Y, Yu X, Zhao H, et al. (2020) Host susceptibility to severe COVID-19 and establishment of a host risk score: Findings of 487 cases outside Wuhan. *Critical Care* 24(1): 108. DOI: [10.1186/s13054-020-2833-7](https://doi.org/10.1186/s13054-020-2833-7).
- Shi Z, Resurreccion WK, Wang C-H, et al. (2020) Association of Cancer with Risk and Mortality of COVID-19: Results from the UK Biobank. *medRxiv*: 2020.07.10.20151076. DOI: [10.1101/2020.07.10.20151076](https://doi.org/10.1101/2020.07.10.20151076).

- Sierpiński R, Pinkas J, Jankowski M, et al. (2020) Sex differences in the frequency of gastrointestinal symptoms and olfactory or taste disorders among 1,942 non-hospitalized patients with COVID-19. *Polish Archives of Internal Medicine*. DOI: [10.20452/pamw.15414](https://doi.org/10.20452/pamw.15414).
- Sigel K, Swartz T, Golden E, et al. (n.d.) Covid-19 and People with HIV Infection: Outcomes for Hospitalized Patients in New York City. *Clinical Infectious Diseases*. DOI: [10.1093/cid/ciaa880](https://doi.org/10.1093/cid/ciaa880).
- Silva Neto PV da, Carvalho JCS de, Pimentel VE, et al. (2020) *Prognostic value of sTREM-1 in COVID-19 patients: A biomarker for disease severity and mortality*. preprint, September. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.09.22.20199703](https://doi.org/10.1101/2020.09.22.20199703).
- Simons D, Brown J, Shahab L, et al. (2020) Smoking and COVID-19: Rapid evidence review for the Royal College of Physicians, London (UK). *Qeios*. DOI: [10.32388/VGJCUN](https://doi.org/10.32388/VGJCUN).
- Simons D, Perski O, Shahab L, et al. (2020) The association of smoking status with hospitalisation for COVID-19 compared with other respiratory viruses a year previous: A case-control study at a single UK National Health Service trust. *medRxiv*: 2020.11.26.20238469. DOI: [10.1101/2020.11.26.20238469](https://doi.org/10.1101/2020.11.26.20238469).
- Simons D, Shahab L, Brown J, et al. (2020) The association of smoking status with SARS-CoV-2 infection, hospitalisation and mortality from COVID-19: A living rapid evidence review with Bayesian meta-analyses (version 7). *Addiction* n/a(n/a). DOI: [10.1111/add.15276](https://doi.org/10.1111/add.15276).
- Singh BM, Bateman J, Viswanath A, et al. (2020) *The risk of COVID hospital admission and COVID mortality during the first COVID 19 wave with a special emphasis on Ethnic Minorities: An observational study of a single, deprived, multi ethnic UK health economy*. preprint, November. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.11.20.20224691](https://doi.org/10.1101/2020.11.20.20224691).
- Sisó-Almirall A, Kostov B, Mas-Heredia M, et al. (2020) PROGNOSTIC FACTORS IN SPANISH COVID-19 PATIENTS: A CASE SERIES FROM BARCELONA. *medRxiv*: 2020.06.18.20134510. DOI: [10.1101/2020.06.18.20134510](https://doi.org/10.1101/2020.06.18.20134510).
- Soares R de CM, Mattos LR and Raposo LM (2020) Risk Factors for Hospitalization and Mortality due to COVID-19 in Espírito Santo State, Brazil.: *tpmd200483*. DOI: [10.4269/ajtmh.20-0483](https://doi.org/10.4269/ajtmh.20-0483).
- Solís P and Carreño H (2020) COVID-19 Fatality and Comorbidity Risk Factors among Diagnosed Patients in Mexico. DOI: [10.1101/2020.04.21.20074591](https://doi.org/10.1101/2020.04.21.20074591).
- Soto-Mota A, Garza BAM, Rodriguez EM, et al. (2020) THE LOW-HARM SCORE FOR PREDICTING MORTALITY IN PATIENTS DIAGNOSED WITH COVID-19: A MULTICENTRIC VALIDATION STUDY. *medRxiv*: 2020.05.26.20111120. DOI: [10.1101/2020.05.26.20111120](https://doi.org/10.1101/2020.05.26.20111120).
- Sourij H, Aziz F, Bräuer A, et al. (2020) COVID-19 fatality prediction in people with diabetes and prediabetes using a simple score upon hospital admission. *Diabetes, Obesity and Metabolism*: dom.14256. DOI: [10.1111/dom.14256](https://doi.org/10.1111/dom.14256).
- Souza FSH, Hojo-Souza NS, Santos EB, et al. (2020) Predicting the disease outcome in COVID-19 positive patients through Machine Learning: A retrospective cohort study with Brazilian data. *medRxiv*: 2020.06.26.20140764. DOI: [10.1101/2020.06.26.20140764](https://doi.org/10.1101/2020.06.26.20140764).
- Swiss Federal Statistics Office (FSO) (2020) La part des gros fumeurs s'est réduite de moitié en 25 ans - Enquête suisse sur la santé 2017: Consommation de tabac Communiqué de presse. *Office fédéral de la statistique*. Available at: [/content/bfs/fr/home/aktuell/neue-veroeffentlichungen.assetdetail.11907023.html](https://content/bfs/fr/home/aktuell/neue-veroeffentlichungen.assetdetail.11907023.html) (accessed 23 February 2021).

- Stead LF, Buitrago D, Preciado N, et al. (2013) Physician advice for smoking cessation. *Cochrane Database of Systematic Reviews* 2017(12). DOI: [10.1002/14651858.CD000165.pub4](https://doi.org/10.1002/14651858.CD000165.pub4).
- Sterlin D, Mathian A, Miyara M, et al. (2020) IgA dominates the early neutralizing antibody response to SARS-CoV-2. *medRxiv*: 2020.06.10.20126532. DOI: [10.1101/2020.06.10.20126532](https://doi.org/10.1101/2020.06.10.20126532).
- Strangfeld A, Schäfer M, Gianfrancesco MA, et al. (2021) Factors associated with COVID-19-related death in people with rheumatic diseases: Results from the COVID-19 Global Rheumatology Alliance physician-reported registry. *Annals of the Rheumatic Diseases*: annrheumdis–2020–219498. DOI: [10.1136/annrheumdis-2020-219498](https://doi.org/10.1136/annrheumdis-2020-219498).
- Suleyman G, Fadel RA, Malette KM, et al. (2020) Clinical Characteristics and Morbidity Associated With Coronavirus Disease 2019 in a Series of Patients in Metropolitan Detroit. *JAMA Network Open* 3(6): e2012270–e2012270. DOI: [10.1001/jamanetworkopen.2020.12270](https://doi.org/10.1001/jamanetworkopen.2020.12270).
- Sun L, Surya S, Le AN, et al. (2021) Rates of COVID-19–Related Outcomes in Cancer Compared With Noncancer Patients. *JNCI Cancer Spectrum* 5(1): pkaa120. DOI: [10.1093/jncics/pkaa120](https://doi.org/10.1093/jncics/pkaa120).
- Talavera B, García-Azorín D, Martínez-Pías E, et al. (2020) Anosmia is associated with lower in-hospital mortality in COVID-19. *Journal of the Neurological Sciences* 419: 117163. DOI: [10.1016/j.jns.2020.117163](https://doi.org/10.1016/j.jns.2020.117163).
- Tao P-Y, Leng L, Liu K, et al. (2020) Determination of risk factors for predicting the onset of symptoms in asymptomatic COVID-19 infected patients. *Int J Med Sci* 17: 2187–2193. DOI: [10.7150/ijms.47576](https://doi.org/10.7150/ijms.47576).
- Tardif J-C, Bouabdallaoui N, L’Allier PL, et al. (2021) Efficacy of Colchicine in Non-Hospitalized Patients with COVID-19. DOI: [10.1101/2021.01.26.21250494](https://doi.org/10.1101/2021.01.26.21250494).
- Targher G, Mantovani A, Wang XB, et al. (2020) Patients with diabetes are at higher risk for severe illness from COVID-19. *Diabetes & Metabolism*. DOI: [10.1016/j.diabet.2020.05.001](https://doi.org/10.1016/j.diabet.2020.05.001).
- Tavakol Z, Ghannadi S, Tabesh MR, et al. (2021) Relationship between physical activity, healthy lifestyle and COVID-19 disease severity; a cross-sectional study. *J. public health NA*: 1–9. DOI: [10.1007/s10389-020-01468-9](https://doi.org/10.1007/s10389-020-01468-9).
- Team RC (2013) The R Project for Statistical Computing.: 1–12.
- Thakur K, Sagayaraj A, Prasad KC, et al. (2021) Olfactory Dysfunction in COVID-19 Patients: Findings from a Tertiary Rural Centre. *Indian j. otolaryngol. head neck surg NA*: 1–7. DOI: [10.1007/s12070-021-02364-8](https://doi.org/10.1007/s12070-021-02364-8).
- Thiabaud A, Iten A, Balmelli C, et al. (2020) SARS-CoV-2/COVID-19 hospitalised patients in Switzerland: A prospective cohort profile. preprint, December. *Epidemiology*. DOI: [10.1101/2020.12.10.20246884](https://doi.org/10.1101/2020.12.10.20246884).
- Thompson JV, Meghani N, Powell BM, et al. (2020) Patient characteristics and predictors of mortality in 470 adults admitted to a district general hospital in England with Covid-19. *medRxiv*: 2020.07.21.20153650. DOI: [10.1101/2020.07.21.20153650](https://doi.org/10.1101/2020.07.21.20153650).
- Torres-Macho J, Ryan P, Valencia J, et al. (2020) The PANDEMYC Score. An Easily Applicable and Interpretable Model for Predicting Mortality Associated With COVID-19. *Journal of Clinical Medicine* 9(10): 3066. DOI: [10.3390/jcm9103066](https://doi.org/10.3390/jcm9103066).
- Tricco AC, Antony J, Zarin W, et al. (2015) A scoping review of rapid review methods. *BMC Medicine* 13(1): 224. DOI: [10.1186/s12916-015-0465-6](https://doi.org/10.1186/s12916-015-0465-6).
- Trubiano JA, Vogrin S, Smibert OC, et al. (2020) COVID-MATCH65 - A prospectively derived clinical decision rule for severe acute respiratory syndrome coronavirus 2. *medRxiv*: 2020.06.30.20143818. DOI: [10.1101/2020.06.30.20143818](https://doi.org/10.1101/2020.06.30.20143818).
- Ugur Chousein EG, Cortuk M, Cinarka H, et al. (2020) Is there any effect of smoking status on severity and mortality of

- hospitalized patients with COVID-19 pneumonia? *Tuberk. Toraks* 68: 371–378. DOI: [10.5578/tt.70352](https://doi.org/10.5578/tt.70352).
- Ullah AZMD, Sivapalan L, Chelala C, et al. (2020) COVID-19 in patients with hepatobiliary and pancreatic diseases in East London: A single-centre cohort study. *medRxiv*: 2020.09.07.20189621. DOI: [10.1101/2020.09.07.20189621](https://doi.org/10.1101/2020.09.07.20189621).
- Vahidy FS, Pan AP, Ahnstedt H, et al. (2021) Sex differences in susceptibility, severity, and outcomes of coronavirus disease 2019: Cross-sectional analysis from a diverse US metropolitan area. *PloS one* 16. DOI: [10.1371/journal.pone.0245556](https://doi.org/10.1371/journal.pone.0245556).
- Valenti L, Bergna A, Pelusi S, et al. (2020) SARS-CoV-2 seroprevalence trends in healthy blood donors during the COVID-19 Milan outbreak. *medRxiv*: 2020.05.11.20098442. DOI: [10.1101/2020.05.11.20098442](https://doi.org/10.1101/2020.05.11.20098442).
- Valenzuela O, Ibanez SE, Poli M, et al. (2020) First report of tocilizumab use in a cohort of Latin American patients hospitalized for severe COVID-19 pneumonia. *medRxiv*: 2020.08.12.20173104. DOI: [10.1101/2020.08.12.20173104](https://doi.org/10.1101/2020.08.12.20173104).
- Valle DMD, Kim-schulze S, Hsin-hui H, et al. (2020) An inflammatory cytokine signature helps predict COVID-19 severity and death. *medRxiv*: 2020.05.28.20115758. DOI: [10.1101/2020.05.28.20115758](https://doi.org/10.1101/2020.05.28.20115758).
- Vaquero LM, Barrado MES, Escobar D, et al. (2020) C-Reactive protein and SOFA score as early predictors of critical care requirement in patients with COVID-19 pneumonia in Spain. *medRxiv*: 2020.05.22.20110429. DOI: [10.1101/2020.05.22.20110429](https://doi.org/10.1101/2020.05.22.20110429).
- Vardavas CI and Nikitara K (2020) COVID-19 and smoking: A systematic review of the evidence. *Tob Induc Dis* 18: 20. DOI: [10.18332/tid/119324](https://doi.org/10.18332/tid/119324).
- Veras FP, Pontelli M, Silva C, et al. (2020) SARS-CoV-2 triggered neutrophil extracellular traps (NETs) mediate COVID-19 pathology. *medRxiv*: 2020.06.08.20125823. DOI: [10.1101/2020.06.08.20125823](https://doi.org/10.1101/2020.06.08.20125823).
- Vila-Corcoles A, Satue-Gracia E, Vila-Rovira A, et al. (2020) *COVID-19 TARRACO Cohort Study: Development of a predictive prognostic rule for early assessment of COVID-19 patients in primary care settings*. preprint, December. Infectious Diseases (except HIV/AIDS). DOI: [10.1101/2020.12.11.20247932](https://doi.org/10.1101/2020.12.11.20247932).
- Vila-Corcoles A, Satue-Gracia E, Vila-Rovira A, et al. (2021) COVID19-related and all-cause mortality among middle-aged and older adults across the first epidemic wave of SARS-COV-2 infection in the region of Tarragona, Spain: Results from the COVID19 TARRACO Cohort Study, March-June 2020. DOI: [10.1101/2021.02.02.21251028](https://doi.org/10.1101/2021.02.02.21251028).
- Vila-Córcoles A, Ochoa-Gondar O, Satué-Gracia EM, et al. (2020) Influence of prior comorbidities and chronic medications use on the risk of COVID-19 in adults: A population-based cohort study in Tarragona, Spain. *BMJ Open* 10(12): e041577. DOI: [10.1136/bmjopen-2020-041577](https://doi.org/10.1136/bmjopen-2020-041577).
- Villar-Garcia J, Vivanco-Hidalgo RM, Cleries M, et al. (2020) Risk factors for SARS-CoV-2 infection, hospitalisation, and death in Catalonia, Spain: A population-based cross-sectional study. *medRxiv*: 2020.08.26.20182303. DOI: [10.1101/2020.08.26.20182303](https://doi.org/10.1101/2020.08.26.20182303).
- Wan S, Xiang Y, Fang W, et al. (2020) Clinical features and treatment of COVID-19 patients in northeast Chongqing. *Journal of Medical Virology* 92(7): 797–806. DOI: [10.1002/jmv.25783](https://doi.org/10.1002/jmv.25783).
- Wang A-L, Zhong X and Hurd Y (2020) Comorbidity and Sociodemographic determinants in COVID-19 Mortality in an US Urban Healthcare System. *medRxiv*: 2020.06.11.20128926. DOI: [10.1101/2020.06.11.20128926](https://doi.org/10.1101/2020.06.11.20128926).
- Wang B, Oekelen OV, Mouhieddine T, et al. (2020) A tertiary center experience of multiple myeloma patients with COVID-19: Lessons learned and the path forward. *medRxiv*: 2020.06.04.20122846. DOI: [10.1101/2020.06.04.20122846](https://doi.org/10.1101/2020.06.04.20122846).

- Wang JG, Liu B, Percha B, et al. (2020) Cardiovascular disease and severe hypoxemia associated with higher rates of non-invasive respiratory support failure in COVID-19. DOI: [10.1101/2020.09.27.20202747](https://doi.org/10.1101/2020.09.27.20202747).
- Wang R, Pan M, Zhang X, et al. (2020) Epidemiological and clinical features of 125 Hospitalized Patients with COVID-19 in Fuyang, Anhui, China. *International Journal of Infectious Diseases* 95: 421–428. DOI: [10.1016/j.ijid.2020.03.070](https://doi.org/10.1016/j.ijid.2020.03.070).
- Wang Z, Zheutlin A, Kao Y-H, et al. (2020) Hospitalised COVID-19 patients of the Mount Sinai Health System: A retrospective observational study using the electronic medical records. *BMJ Open* 10(10): e040441. DOI: [10.1136/bmjopen-2020-040441](https://doi.org/10.1136/bmjopen-2020-040441).
- Wang Z-H, Shu C, Ran X, et al. (2020) Critically Ill Patients with Coronavirus Disease 2019 in a Designated ICU: Clinical Features and Predictors for Mortality. *Risk Management and Healthcare Policy* Volume 13: 833–845. DOI: [10.2147/RMHP.S263095](https://doi.org/10.2147/RMHP.S263095).
- Ward H, Atchison CJ, Whitaker M, et al. (2020) Antibody prevalence for SARS-CoV-2 in England following first peak of the pandemic: REACT2 study in 100,000 adults. *medRxiv*: 2020.08.12.20173690. DOI: [10.1101/2020.08.12.20173690](https://doi.org/10.1101/2020.08.12.20173690).
- Weerahandi H, Hochman KA, Simon E, et al. (2020) Post-discharge health status and symptoms in patients with severe COVID-19. *medRxiv*: 2020.08.11.20172742. DOI: [10.1101/2020.08.11.20172742](https://doi.org/10.1101/2020.08.11.20172742).
- Wei W, Ortwine JK, Mang NS, et al. (2020) Limited Role for Antibiotics in COVID-19: Scarce Evidence of Bacterial Coinfection. *medRxiv*: 2020.06.16.20133181. DOI: [10.1101/2020.06.16.20133181](https://doi.org/10.1101/2020.06.16.20133181).
- Williamson EJ, Walker AJ, Bhaskaran K, et al. (2020) Factors associated with COVID-19-related death using OpenSAFELY. *Nature* 584(7821): 430–436. DOI: [10.1038/s41586-020-2521-4](https://doi.org/10.1038/s41586-020-2521-4).
- Woolcott OO and Castilla-Bancayán JP (2020) *Diabetes and Mortality Among 1.6 Million Adult Patients Screened for SARS-CoV-2 in Mexico*. preprint, November. *Infectious Diseases (except HIV/AIDS)*. DOI: [10.1101/2020.11.25.20238345](https://doi.org/10.1101/2020.11.25.20238345).
- Woolford SJ, D'Angelo S, Curtis EM, et al. (2020) COVID-19 and associations with frailty and multimorbidity: A prospective analysis of UK Biobank participants. *Aging Clinical and Experimental Research* 32(9): 1897–1905. DOI: [10.1007/s40520-020-01653-6](https://doi.org/10.1007/s40520-020-01653-6).
- Wu MA, Fossali T, Pandolfi L, et al. (2020) COVID-19: The key role of pulmonary capillary leakage. An observational cohort study. *medRxiv*: 2020.05.17.20104877. DOI: [10.1101/2020.05.17.20104877](https://doi.org/10.1101/2020.05.17.20104877).
- Xie Y, Chen S, Wang X, et al. (2020) Early Diagnosis and Clinical Significance of Acute Cardiac Injury - Under the Iceberg: A Retrospective Cohort Study of 619 Non-critically Ill Hospitalized COVID-19 Pneumonia Patients. *medRxiv*: 2020.07.06.20147256. DOI: [10.1101/2020.07.06.20147256](https://doi.org/10.1101/2020.07.06.20147256).
- Xu H, Hou K, Xu H, et al. (2020) Acute Myocardial Injury of Patients with Coronavirus Disease 2019. *medRxiv*: 2020.03.05.20031591. DOI: [10.1101/2020.03.05.20031591](https://doi.org/10.1101/2020.03.05.20031591).
- Yadaw AS, Li Y-c, Bose S, et al. (2020) Clinical features of COVID-19 mortality: Development and validation of a clinical prediction model. *The Lancet Digital Health* 2(10): e516–e525. DOI: [10.1016/S2589-7500\(20\)30217-X](https://doi.org/10.1016/S2589-7500(20)30217-X).
- Yan H, Valdes AM, Vijay A, et al. (2020) Role of Drugs used for chronic disease management on Susceptibility and Severity of COVID-19: A Large Case-Control Study. *medRxiv*: 2020.04.24.20077875. DOI: [10.1101/2020.04.24.20077875](https://doi.org/10.1101/2020.04.24.20077875).
- Yang X, Yu Y, Xu J, et al. (2020) Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: A single-centered, retrospective, observational study. *The Lancet Respiratory Medicine* 8(5): 475–481. DOI: [10.1016/S2213-2600\(20\)30079-5](https://doi.org/10.1016/S2213-2600(20)30079-5).

- Yanover C, Mizrahi B, Kalkstein N, et al. (2020) What factors increase the risk of complications in SARS-CoV-2 positive patients? A cohort study in a nationwide Israeli health organization. *medRxiv*: 2020.05.07.20091652. DOI: [10.1101/2020.05.07.20091652](https://doi.org/10.1101/2020.05.07.20091652).
- Yao K, Hasegawa S, Tagashira Y, et al. (2021) Experience of 101 patients with coronavirus infectious disease 2019 (COVID-19) at a tertiary care center in Japan. *Journal of Infection and Chemotherapy* 27(2): 413–417. DOI: [10.1016/j.jiac.2020.11.024](https://doi.org/10.1016/j.jiac.2020.11.024).
- Yao Q, Wang P, Wang X, et al. (n.d.) A retrospective study of risk factors for severe acute respiratory syndrome coronavirus 2 infections in hospitalized adult patients. DOI: [10.20452/pamw.15312](https://doi.org/10.20452/pamw.15312).
- Yin R, Yang Z, Wei Y, et al. (2020) Clinical characteristics of 106 patients with neurological diseases and co-morbid coronavirus disease 2019: A retrospective study. *medRxiv*: 2020.04.29.20085415. DOI: [10.1101/2020.04.29.20085415](https://doi.org/10.1101/2020.04.29.20085415).
- Yoo E, Percha B, Tomlinson M, et al. (2020) Development and calibration of a simple mortality risk score for hospitalized COVID-19 adults. *medRxiv*: 2020.08.31.20185363. DOI: [10.1101/2020.08.31.20185363](https://doi.org/10.1101/2020.08.31.20185363).
- Yu T, Cai S, Zheng Z, et al. (2020) Association Between Clinical Manifestations and Prognosis in Patients with COVID-19. *Clinical Therapeutics* 42(6): 964–972. DOI: [10.1016/j.clinthera.2020.04.009](https://doi.org/10.1016/j.clinthera.2020.04.009).
- Zacharioudakis IM, Prasad PJ, Zervou FN, et al. (2020) Association of SARS-CoV-2 Genomic Load with COVID-19 Patient Outcomes. *medRxiv*: 2020.07.02.20145151. DOI: [10.1101/2020.07.02.20145151](https://doi.org/10.1101/2020.07.02.20145151).
- Zeng H, Zhang T, He X, et al. (2020) Impact of Chronic Comorbidities on Progression and Prognosis in Patients with COVID-19: A Retrospective Cohort Study in 1031 Hospitalized Cases in Wuhan, China. *medRxiv*: 2020.06.14.20125997. DOI: [10.1101/2020.06.14.20125997](https://doi.org/10.1101/2020.06.14.20125997).
- Zhan T, Liu M, Tang Y, et al. (2020) Retrospective analysis of clinical characteristics of 405 patients with COVID-19. *Journal of International Medical Research* 48(8): 030006052094903. DOI: [10.1177/0300060520949039](https://doi.org/10.1177/0300060520949039).
- Zhan Z, Yang X, Du H, et al. (2020) Early Improvement of Acute Respiratory Distress Syndrome in Patients with COVID-19: Insights from the Data of ICU Patients in Chongqing, China. *medRxiv*: 2020.07.15.20154047. DOI: [10.1101/2020.07.15.20154047](https://doi.org/10.1101/2020.07.15.20154047).
- Zhang J, Cao Y, Tan G, et al. (2020) Clinical, radiological, and laboratory characteristics and risk factors for severity and mortality of 289 hospitalized COVID-19 patients. *Allergy*: all.14496. DOI: [10.1111/all.14496](https://doi.org/10.1111/all.14496).
- Zhang J-j, Dong X, Cao Y-y, et al. (2020) Clinical characteristics of 140 patients infected with SARS-CoV-2 in Wuhan, China. *Allergy* 75(7): 1730–1741. DOI: [10.1111/all.14238](https://doi.org/10.1111/all.14238).
- Zhang Q, Wang Z, Lv Y, et al. (2021) Clinical features and prognostic factors of patients with COVID-19 in Henan Province, China. *Hum Cell* NA. DOI: [10.1007/s13577-021-00499-y](https://doi.org/10.1007/s13577-021-00499-y).
- Zhang X, Cai H, Hu J, et al. (2020) Epidemiological, clinical characteristics of cases of SARS-CoV-2 infection with abnormal imaging findings. *International Journal of Infectious Diseases* 94: 81–87. DOI: [10.1016/j.ijid.2020.03.040](https://doi.org/10.1016/j.ijid.2020.03.040).
- Zhang X, Li X, Sun Z, et al. (2020) Physical activity and COVID-19: An observational and Mendelian randomisation study. *Journal of Global Health* 10(2): 020514. DOI: [10.7189/jogh-10-020514](https://doi.org/10.7189/jogh-10-020514).
- Zhang Y, Yang H, Li S, et al. (2021) Association analysis framework of genetic and exposure risks for COVID-19 in middle-aged and elderly adults. *Mechanisms of Ageing and Development* 194: 111433. DOI: [10.1016/j.mad.2021.111433](https://doi.org/10.1016/j.mad.2021.111433).
- Zhao Z, Chen A, Hou W, et al. (2020) Prediction model and risk scores of ICU admission and mortality in COVID-19. *PLOS*

ONE 15(7): e0236618. DOI: [10.1371/journal.pone.0236618](https://doi.org/10.1371/journal.pone.0236618).

Zheng KI, Gao F, Wang X-B, et al. (2020) Letter to the Editor: Obesity as a risk factor for greater severity of COVID-19 in patients with metabolic associated fatty liver disease. *Metabolism* 108: 154244. DOI: [10.1016/j.metabol.2020.154244](https://doi.org/10.1016/j.metabol.2020.154244).

Zheng Y, Xiong C, Liu Y, et al. (2020) Epidemiological and clinical characteristics analysis of COVID-19 in the surrounding areas of Wuhan, Hubei Province in 2020. *Pharmacological Research* 157: 104821. DOI: [10.1016/j.phrs.2020.104821](https://doi.org/10.1016/j.phrs.2020.104821).

Zhong R, Chen L, Zhang Q, et al. (2021) Which Factors, Smoking, Drinking Alcohol, Betel Quid Chewing, or Underlying Diseases, Are More Likely to Influence the Severity of COVID-19? *Frontiers in Physiology* 11: 623498. DOI: [10.3389/fphys.2020.623498](https://doi.org/10.3389/fphys.2020.623498).

Zhou F, Yu T, Du R, et al. (2020) Clinical course and risk factors for mortality of adult inpatients with COVID-19 in Wuhan, China: A retrospective cohort study. *The Lancet* 395(10229): 1054–1062. DOI: [10.1016/S0140-6736\(20\)30566-3](https://doi.org/10.1016/S0140-6736(20)30566-3).

Zhou J, Ma Y, Liu Y, et al. (2021) A Correlation Analysis between the Nutritional Status and Prognosis of COVID-19 Patients. *J Nutr Health Aging* 25: 84–93. DOI: [10.1007/s12603-020-1457-6](https://doi.org/10.1007/s12603-020-1457-6).

Zhou K, Sun Y, Li L, et al. (2020) Eleven Routine Clinical Features Predict COVID-19 Severity. *medRxiv*: 2020.07.28.20163022. DOI: [10.1101/2020.07.28.20163022](https://doi.org/10.1101/2020.07.28.20163022).

Zhou W, Song L, Wang X, et al. (2020) Cardiac injury prediction and lymphocyte immunity and inflammation analysis in hospitalized patients with coronavirus disease 2019 (COVID-19). *International Journal of Cardiology*: S016752732034002X. DOI: [10.1016/j.ijcard.2020.10.049](https://doi.org/10.1016/j.ijcard.2020.10.049).

Zhou W, Qin X, Hu X, et al. (2020) Prognosis models for severe and critical COVID-19 based on the Charlson and Elixhauser comorbidity indices. *International Journal of Medical Sciences* 17(15): 2257–2263. DOI: [10.7150/ijms.50007](https://doi.org/10.7150/ijms.50007).

Zhou Y, He Y, Yang H, et al. (2020) Exploiting an early warning Nomogram for predicting the risk of ICU admission in patients with COVID-19: A multi-center study in China. *Scandinavian Journal of Trauma, Resuscitation and Emergency Medicine* 28(1): 106. DOI: [10.1186/s13049-020-00795-w](https://doi.org/10.1186/s13049-020-00795-w).

Zhou Y, He X, Zhang J, et al. (2020) Prolonged SARS-CoV-2 Viral Shedding in Patients with COVID-19 was Associated with Delayed Initiation of Arbidol Treatment: A retrospective cohort study. *medRxiv*: 2020.06.09.20076646. DOI: [10.1101/2020.06.09.20076646](https://doi.org/10.1101/2020.06.09.20076646).

Zhu S, Gao Q, Yang L, et al. (2020) Prevalence and risk factors of disability and anxiety in a retrospective cohort of 432 survivors of Coronavirus Disease-2019 (Covid-19) from China. *PLOS ONE* Federici S (ed.) 15(12): e0243883. DOI: [10.1371/journal.pone.0243883](https://doi.org/10.1371/journal.pone.0243883).

Ziehr DR, Alladina J, Petri CR, et al. (2020) Respiratory Pathophysiology of Mechanically Ventilated Patients with COVID-19: A Cohort Study. *American Journal of Respiratory and Critical Care Medicine* 201(12): 1560–1564. DOI: [10.1164/rccm.202004-1163LE](https://doi.org/10.1164/rccm.202004-1163LE).

Zinellu A, Arru F, De Vito A, et al. (2021) The De Ritis ratio as prognostic biomarker of in-hospital mortality in COVID-19 patients. *European Journal of Clinical Investigation* 51(1). DOI: [10.1111/eci.13427](https://doi.org/10.1111/eci.13427).

Zobairy H, Shamsoddin E, Rasouli MA, et al. (2020) Association of olfactory dysfunction with hospitalization for COVID-19: A multicenter study in Kurdistan. *medRxiv*: 2020.07.26.20158550. DOI: [10.1101/2020.07.26.20158550](https://doi.org/10.1101/2020.07.26.20158550).

Zuo Y, Yalavarthi S, Shi H, et al. (2020) Neutrophil extracellular traps in COVID-19. *JCI Insight*. DOI: [10.1172/jci.insight.138999](https://doi.org/10.1172/jci.insight.138999).

Zuo Y, Warnock M, Harbaugh A, et al. (2020) *Plasma tissue plasminogen activator and plasminogen activator inhibitor-1 in hospitalized COVID-19 patients*. preprint, September. Infectious Diseases (except HIV/AIDS). DOI: [10.1101/2020.08.29.20184358](https://doi.org/10.1101/2020.08.29.20184358).

Zuo Y, Estes SK, Gandhi AA, et al. (2020) Prothrombotic antiphospholipid antibodies in COVID-19. *medRxiv*: 2020.06.15.20131607. DOI: [10.1101/2020.06.15.20131607](https://doi.org/10.1101/2020.06.15.20131607).