

Viscosity Limits for Zeroth-Order Pseudodifferential Operators

JEFFREY GALKOWSKI

University College London

MACIEJ ZWORSKI

University of California, Berkeley

Abstract

Motivated by the work of Colin de Verdière and Saint-Raymond on spectral theory for zeroth-order pseudodifferential operators on tori, we consider viscosity limits in which zeroth-order operators, P , are replaced by $P + i\nu\Delta$, $\nu > 0$. By adapting the Helffer–Sjöstrand theory of scattering resonances, we show that, in a complex neighbourhood of the continuous spectrum, eigenvalues of $P + i\nu\Delta$ have limits as the viscosity ν goes to 0. In the simplified setting of tori, this justifies claims made in the physics literature. © 2021 The Authors. *Communications on Pure and Applied Mathematics* published by Wiley Periodicals LLC.

1 Introduction

Spectral properties of zeroth-order pseudodifferential operators arise naturally in the problems of fluid mechanics—for an early example, see Ralston [28]. Recently Colin de Verdière and Saint-Raymond [3,4] investigated such operators under natural dynamical conditions motivated by the study of (linearized) internal waves—see the review article of Dauxois et al. [6] and the introduction to [4] for a physics perspective and references. Dyatlov–Zworski [12] provided proofs of the results of [4] based on the analogy to scattering theory—see Melrose–Zworski [26], Hassell–Melrose–Vasy [17], and [11]. This analogy was developed further by Wang [38], who defined and described a scattering matrix in this setting. Tao [36] constructed an example of an embedded eigenvalue.

Motivated by the physics literature—see, for instance, Rieutord and Valdetaro [29]—we consider here operators with a viscosity term

$$P_\nu := P + i\nu\Delta,$$

where P is a zeroth-order pseudodifferential operator on the torus (1.1) satisfying (1.2) and the dynamical assumption (1.3). The operator Δ is the usual Laplacian on the torus. The assumption (1.3) guarantees continuity of the spectrum at 0 [4, 12]. We then show that as $\nu \rightarrow 0+$ the eigenvalues of P_ν in a complex neighbourhood of 0 tend to a discrete set associated to P alone—see Figure 1.1 for

Communications on Pure and Applied Mathematics, Vol. LXXV, 1798–1869 (2022)

© 2021 The Authors. *Communications on Pure and Applied Mathematics* published by Wiley Periodicals LLC.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.




FIGURE 1.1. We display the resonances of P as red stars (a full explanation using the deformed operator P_θ is given in Appendix B). The paths of the eigenvalues of $P + i\nu\Delta$ as $\nu \rightarrow 0+$ are shown by the green curves with the arrows denoting the direction of the path as ν decreases. P is chosen as in (B.4) with $V_a = \frac{1}{2}(\xi^3 - 1)e^{-\xi^2}$ and $V_m = (1 + (e - 1)(\xi - 2)^2)e^{-(\xi - 2)^2}$. For an animated version of this figure see https://math.berkeley.edu/~zworski/vis_dynam.mov.

a numerical illustration. This justifies claims seen in related models of the physics literature.[†] Our approach is again based on analogy to scattering theory, in this case to the general approach to scattering resonances due to Helffer–Sjöstrand [18].

To state our results precisely, we start with the class of pseudodifferential operators:

$$(1.1) \quad Pu(y) := \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{i(y-y',\eta)} p(y, \eta) u(y') dy' d\eta$$

where $p \in S^m(T^*\mathbb{T}^n)$, $\mathbb{T}^n := \mathbb{R}^n/2\pi\mathbb{Z}^n$, has an analytic continuation from $T^*\mathbb{T}^n$ satisfying

$$(1.2) \quad |p(z, \zeta)| \leq M \quad \text{for } |\text{Im } z| \leq a, \quad |\text{Im } \zeta| \leq b\langle \text{Re } \zeta \rangle.$$

The integral in the definition (1.1) of Pu is considered in the sense of oscillatory integrals (see, for instance, [40, sec. 5.3]), and we extend both $y \mapsto u(y)$ and $y \mapsto p(y, \eta)$ to periodic functions on \mathbb{R}^n .

[†] For example, a claim from [29]: ‘The aim of this paper is to present what we believe to be the asymptotic limit of inertial modes in a spherical shell when viscosity tends to zero.’

The dynamical assumption is formulated using an *escape function*:

$$(1.3) \quad \exists G \in S^1(T^*\mathbb{T}^n), C > 0, H_p G(x, \xi) > 0 \\ \text{for } (x, \xi) \in \{p = 0\} \cap \{|\xi| > C\}.$$

(For the definition of the symbol class $S^1(T^*\mathbb{T}^n)$, see (3.1) and [11, Section E.1], and for a discussion of escape functions [11, Section 6.4].) We make our assumption at $p = 0$ but the value 0 can be replaced by any real number λ by changing the operator to $P - \lambda$. We could also replace p in (1.3) by the principal symbol of P . Examples of operators satisfying our assumptions are given in Appendix B (see also [12] and [36]).

We denote by $\Delta = \sum_{j=1}^n \partial_{x_j}^2$ the usual Laplacian on \mathbb{T}^n and state a precise version of our main result:

THEOREM 1.1. *Suppose that P is given by (1.1) with p satisfying (1.2) and (1.3). Then there exist an open neighbourhood of 0 in \mathbb{C} , U , and a set*

$$\mathcal{R}(P) \subset \{\operatorname{Im} \omega \leq 0\} \cap U$$

such that for every $K \Subset U$, $\mathcal{R}(P) \cap K$ is discrete, and

$$(1.4) \quad \operatorname{spec}_{L^2}(P + i\nu\Delta) \longrightarrow \mathcal{R}(P), \quad \nu \rightarrow 0+,$$

uniformly on K .

Numerical illustrations of this theorem are presented in Appendix B.

Another way to state the theorem is to say that $\mathcal{R}(P) = \{\omega_j\}_{j=1}^N$ (where $N = \infty$ is allowed) and $\operatorname{spec}_{L^2}(P + i\nu\Delta) = \{\omega_j(\nu)\}_{j=1}^\infty$; then (after suitable re-ordering)

$$\omega_j(\nu) \rightarrow \omega_j, \quad \nu \rightarrow 0+,$$

uniformly on compact sets and with agreement of multiplicities. In fact, the proof gives a more precise statement implying smoothness of projectors acting on spaces \mathcal{X} of Theorem 1.2—see [9, prop. 5.3]. Since the statement is essentially the same, we do not reproduce it here.

The Laplacian Δ can be replaced by any second order (or any order) elliptic differential operator with analytic coefficients, and the set $\mathcal{R}(P)$ is independent of that choice. The next theorem shows that $\mathcal{R}(P)$ is defined intrinsically for operators satisfying our assumptions:

THEOREM 1.2. *Suppose that P satisfies the assumptions of Theorem 1.1, and U is the open set presented there. Then there exists a Hilbert space \mathcal{X} such that for $\omega \in U$,*

$$P - \omega : \mathcal{X} \rightarrow \mathcal{X} \quad \text{is a Fredholm operator}$$

and $\mathcal{R}(\omega) := (P - \omega)^{-1} : \mathcal{X} \rightarrow \mathcal{X}$ forms a meromorphic family of operators with poles of finite rank. The set of these poles in U is the set $\mathcal{R}(P)$ in Theorem 1.1 (with inclusion according to multiplicity). Moreover,

$$\mathcal{R}(P) \cap \mathbb{R} = \operatorname{spec}_{\text{pp}, L^2}(P) \cap U.$$

The space $\mathcal{X} = H_\Lambda$ is defined in Section 4 and for some $\delta > 0$,

$$\mathcal{A}_\delta \subset \mathcal{X} \subset \mathcal{A}_{-\delta},$$

where, for $s \in \mathbb{R}$, the spaces \mathcal{A}_s are given by formal Fourier series with Fourier coefficients bounded by $e^{-|n|s}$, $n \in \mathbb{Z}^n$. Hence \mathcal{X} contains the space of analytic functions extending to a sufficiently large complex neighbourhood of \mathbb{T}^n and is contained in the dual of such space—see (4.2) for precise definitions.

We briefly recall similar results in different settings. In Dyatlov–Zworski [9], they showed that if X is the generator of an Anosov flow on a compact manifold and Q is a self-adjoint second-order elliptic operator, then the eigenvalues of $X + i\nu Q$ converge to the *Pollicott–Ruelle resonances* of the Anosov flow. These resonances appear in expansions of correlations—see [9] for a discussion and references. Drouot [8] proved an analogue of this result for *kinetic Brownian motion* in which X is a generator of an Anosov geodesic flow and Q is the ‘spherical Laplacian’ on the fibers. Dang–Rivière [5] showed that for Morse–Smale gradient flows, the eigenvalues of $\mathcal{L}_{\nabla_g} f + i\nu \Delta_g$ (which agree with the eigenvalues of the *Witten Laplacian*) converge to the Pollicott–Ruelle resonance of the gradient flow. That generalized a result of Frenkel–Losev–Nekrasov [14], who, motivated by quantum field theory, considered the case of the height function on the sphere.

The *complex absorbing potential* method (see [11, Section 4.9] for a description and references) is also related to viscosity limits: to obtain discrete complex spectrum, a complex potential, say $-i\epsilon|x|^2$, is added to a Schrödinger operator. In cases where scattering resonances can be defined, the spectrum of this new operator converges to the resonances—see [39, 41].

The essential ingredient in the proof of Theorems 1.1 and 1.2 is the theory of complex microlocal deformations inspired by works of Sjöstrand [31, 33] and Helffer–Sjöstrand [18]. The starting object is a Fourier–Bros–Iagolnitzer (FBI) transform considered with an additional asymptotic parameter $h \rightarrow 0$, which in this paper will be eventually fixed and sufficiently small. In our case we need an FBI transform that respects the analytic structure of the underlying compact analytic manifold. Hence, if M is a compact analytic manifold, we define (using a measure on M coming from a real analytic metric)

$$(1.5) \quad Tu(x, \xi, h) := h^{-\frac{3n}{2}} \int_M K(x, \xi, y, h)u(y)dy,$$

where

$$(x, \xi, y) \rightarrow K(x, \xi, y, h)$$

is holomorphic in a fixed complex conic neighbourhood of $T^*M \times M$, and uniformly in that neighbourhood,

$$(1.6) \quad \begin{aligned} K(x, \xi, y, h) &= \chi(x, y)a(x, \xi, y, h)e^{\frac{i}{h}\varphi(x, \xi, y, h)} + \mathcal{O}(e^{-(\operatorname{Re} \xi)/Ch}), \\ \varphi(x, \xi, x) &= 0, \quad d_y \varphi(x, \xi, y)|_{y=x} = -\xi, \\ \operatorname{Im} d_y^2 \varphi(x, \xi, y)|_{y=x} &\sim \langle \operatorname{Re} \xi \rangle I. \end{aligned}$$

Denoting by \tilde{M} a complex neighbourhood of M , χ satisfies

$$\chi \in C^\infty(U), \quad \chi|_V \equiv 1,$$

$$V \Subset U \subset \tilde{M} \times \tilde{M} \text{ are small neighbourhoods of } \Delta(\tilde{M}),$$

and a is an analytic symbol of order $n/4$ in ξ . (Here $\Delta(\tilde{M})$ denotes the diagonal $\{(x, x) : x \in \tilde{M}\}$.)

Existence of such kernels K can be obtained by choosing a real analytic metric with exponential map $T_x M \ni (x, v) \mapsto \exp_x(v)$, and then putting

$$\varphi(x, \xi, y) = -\xi(\exp_x^{-1}(y)) + \frac{i}{2}\langle \xi \rangle d(x, y)^2.$$

We can then solve the $\bar{\partial}$ -equation with the right-hand side given by $\bar{\partial}_{x,y}$ applied to the first term on the right-hand side of (1.6).

In this paper, in view of our applications and for the sake of clarity, we consider an explicit $K(x, \xi, y, h)$ available in the case of tori, $\mathbb{T}^n := \mathbb{R}^n / (2\pi\mathbb{Z})^n$:

$$(1.7) \quad K(x, \xi, y, h) = c_n \langle \xi \rangle^{\frac{n}{4}} \sum_{k \in \mathbb{Z}} e^{\frac{i}{h}(\langle (x-y-2\pi k, \xi) \rangle + \frac{1}{2}\langle \xi \rangle (x-y-2\pi k)^2)}.$$

Although the analysis works in the more general setting of analytic compact manifolds and FBI transforms satisfying (1.5) and (1.6), we can avoid additional complications such as the study of analytic symbols when the inverse of T is not exact (see Proposition 2.3) and of operators annihilating Tu that do not commute exactly (see Proposition 5.1) by using (1.7). One motivation for this project was to present the theory of *exponential weights*, which are not compactly supported—see Section 4. The expository article [15] is intended as an introduction to these methods in the simpler setting of compactly supported weights; see also Martinez [24] and Nakamura [27] for a very clear approach to compactly supported weights in \mathbb{R}^n (or more generally weights ψ satisfying $\partial^\alpha \psi \in L^\infty$ for $|\alpha| > 0$).

In an independent development Guedes Bonthonneau–Jézéquel [16] presented a similar theory in a more general setting of Gevrey regularity and arbitrary real analytic compact manifolds. Their motivation came from a microlocal study of dynamical zeta functions and trace formulas for Anosov flows; see [10, 23] and references given there.

The paper is organized as follows:

- In Section 2 we define an FBI transform T on tori, and construct its exact left inverse S . The FBI transform takes functions on \mathbb{T}^n to functions on $T^*\mathbb{T}^n$.
- The geometry of complex deformation and their relation to exponential weights is reviewed in Section 3. The complex deformation of our FBI transform, T_Λ , is then investigated in Section 4 where the space $\mathcal{X} = H_\Lambda$ is also defined. Here, Λ is a complex deformation of $T^*\mathbb{T}^n$ associated to G in (1.3) using (3.2).

- Section 5 is motivated by the study of Bergman kernels by Boutet de Monvel–Sjöstrand [2, 33] and of Toeplitz operators by Boutet de Monvel–Guillemin [1]: we construct a parametrix for the orthogonal projector onto the image of \mathcal{X} under T_Λ .
- The action of pseudodifferential operators of the form (1.1) on the space \mathcal{X} is described in Section 6. We also present the compactness and trace class properties needed in our proofs of the Fredholm property and of the viscosity limit for P and $P + i\nu\Delta$.
- Finally, Section 6 is devoted to the proofs of Theorems 1.1 and 1.2.
- Appendix A reviews some aspects of the almost analytic machinery of Melin–Sjöstrand [25]; see also [15, Section 5]. In Appendix B we discuss the (very) special case of escape functions that are linear in ξ . In that case we can use an analogue of the method of complex scaling—see [11, Section 4.5, Section 4.7] and references given there. This method lends itself to numerical experimentation, and some results of that are presented in Appendix B as well.

Comments on notation: We write $f_\alpha = \mathcal{O}_\alpha(g)_H$ for $\|f\|_H \leq C_\alpha g$; that is, we have a bound with constants depending on α . In particular, $f = \mathcal{O}(h^\infty)_H$ means that for any N there exists C_N such that $\|f\|_H \leq C_N h^N$. We denote $\langle \xi \rangle := \sqrt{1 + |\xi|^2}$. The notation $\text{nbhd}_U(\rho)$ means an open neighbourhood of ρ in the space U .

2 A Semiclassical FBI Transform on $\mathbb{T}^n = \mathbb{R}^n / 2\pi\mathbb{Z}^n$

We start by defining an FBI transform on \mathbb{T}^n that respects the *real analytic* structure of \mathbb{T}^n and is invertible with error exponentially small in h and in frequency.

As stated in Section 1 we achieve this with the following transform:

$$(2.1) \quad Tu(x, \xi) := h^{-\frac{3n}{4}} \int_{\mathbb{T}^n} \sum_{k \in \mathbb{Z}^n} e^{\frac{i}{h}\varphi(x, y - 2\pi k, \xi)} \langle \xi \rangle^{\frac{n}{4}} u(y) dy, \quad u \in C^\infty(\mathbb{T}^n),$$

$$\varphi(x, \xi, y) := \langle x - y, \xi \rangle + \frac{i}{2} \langle \xi \rangle (x - y)^2.$$

This sum is rapidly convergent since $\text{Im } \varphi \geq \langle \xi \rangle |x - y|^2 / 2$.

Remark 2.1. As already emphasized in Section 1, the crucial feature of T is the structure of its integral kernel, $K(x, \xi, y)$, which is analytic in all variables and is given by

$$e^{\frac{i}{h}\varphi(x, \xi, y)} a(x, \xi, y) \chi(d(x, y)) + \mathcal{O}(e^{-\langle \xi \rangle / h}),$$

$$\varphi(x, \xi, y) = \langle \exp_y^{-1}(x), \xi \rangle + \frac{i}{2} \langle \xi \rangle d(x, y)^2,$$

where a is a classical analytic symbol and $\chi \in C_c^\infty(\mathbb{R})$ is supported in a small neighbourhood of 0 and is equal to 1 near 0.

Extending u to \mathbb{R}^n as a $2\pi\mathbb{Z}^n$ periodic function, we observe that

$$Tu(x, \xi) = h^{-\frac{3n}{4}} \int_{\mathbb{R}^n} e^{\frac{i}{h}\varphi(x-y, \xi)} \langle \xi \rangle^{\frac{n}{4}} u(y) dy$$

and, moreover, $Tu(x, \xi)$ is $2\pi\mathbb{Z}^n$ periodic in x .

LEMMA 2.2. *The operator $T : C^\infty(\mathbb{T}^n) \rightarrow C^\infty(T^*\mathbb{T}^n)$ extends to an operator*

$$T : L^2(\mathbb{T}^n) \rightarrow L^2(T^*\mathbb{T}^n), \quad \|T\|_{L^2(\mathbb{T}^n) \rightarrow L^2(T^*\mathbb{T}^n)} \leq C,$$

with C independent of h .

PROOF. Suppose that $v \in C_c^\infty(T^*\mathbb{T}^n)$. We extend v periodically in x and consider

$$TT^*v(x, \xi) = h^{-\frac{3n}{2}} \langle \xi \rangle^{\frac{n}{4}} \int_{\mathbb{R}^{3n}} \langle \eta \rangle^{\frac{n}{4}} e^{\frac{i}{h}\Phi} v(y, \eta) dy d\eta dw,$$

where

$$\Phi := \langle x - w, \xi \rangle + \langle w - y, \eta \rangle + \frac{i}{2} (\langle \xi \rangle (x - w)^2 + \langle \eta \rangle (y - w)^2).$$

Completing the square and integrating in w , we then obtain

$$TT^*v(x, \xi) = h^{-n} \int_{T^*\mathbb{T}^n} \frac{\langle \xi \rangle^{\frac{n}{4}} \langle \eta \rangle^{\frac{n}{4}}}{(\langle \xi \rangle + \langle \eta \rangle)^{\frac{n}{2}}} \sum_{k \in \mathbb{Z}^n} e^{\frac{i}{h}\Psi(x-y+2\pi k, \xi, \eta)} v(y, \eta) dy d\eta.$$

where

$$\Psi(z, \xi, \eta) := \frac{i}{2} \frac{(\xi - \eta)^2}{\langle \xi \rangle + \langle \eta \rangle} + \frac{i}{2} \frac{\langle \eta \rangle \langle \xi \rangle z^2}{\langle \xi \rangle + \langle \eta \rangle} + \frac{\langle \eta \rangle \xi + \langle \xi \rangle \eta}{\langle \xi \rangle + \langle \eta \rangle} \cdot z.$$

Schur’s test for boundedness, together with density of $C_c^\infty(T^*\mathbb{T}^n)$ in $L^2(T^*\mathbb{T}^n)$, completes the proof of the lemma. □

Our next goal is to find an inverse for T . To do this, we define

$$(2.2) \quad Sv(y) = h^{-\frac{3n}{4}} \int_{T^*\mathbb{T}^n} \sum_{k \in \mathbb{Z}^n} e^{-\frac{i}{h}\varphi^*(x-2\pi k, y, \xi)} \cdot b(x - y - 2\pi k, \xi) v(x, \xi) dx d\xi$$

Then, as before, extending v periodically in x ,

$$Sv(y) = h^{-\frac{3n}{4}} \int_{T^*\mathbb{R}^n} e^{\frac{i}{h}\varphi^*(x, y, \xi)} b(x - y, \xi) v(x, \xi) dx d\xi.$$

We then have

PROPOSITION 2.3. *Putting*

$$(2.3) \quad b(w, \xi) = 2^{\frac{n}{2}} (2\pi)^{-\frac{3n}{2}} \langle \xi \rangle^{\frac{n}{4}} (1 + \frac{i}{2} \langle w, \xi / \langle \xi \rangle \rangle),$$

in (2.2) gives

$$(2.4) \quad STu = u, \quad u \in L^2(\mathbb{R}^n).$$

PROOF. Using definition (2.1) and (2.2) we have

$$\begin{aligned}
 (2.5) \quad STu &= h^{-\frac{3n}{2}} \int_{\mathbb{R}^n \times \mathbb{R}^n \times \mathbb{R}^n} e^{\frac{i}{h} \langle (x-y, \xi) + \frac{i}{2} \langle \xi \rangle (z-y)^2 + \frac{i}{2} \langle \xi \rangle (x-z)^2 \rangle} \cdot \langle \xi \rangle^{\frac{n}{4}} b(x-z, \xi) u(y) dy dz d\xi \\
 &= h^{-n} \int_{\mathbb{R}^n \times \mathbb{R}^n} e^{\frac{i}{h} \langle x-y, \xi \rangle - \frac{\langle \xi \rangle}{4h} (x-y)^2} a(x, y, \xi; h) u(y) dy d\xi.
 \end{aligned}$$

For our choice of b we have

$$\begin{aligned}
 (2.6) \quad a(x, y, \xi; h) &= h^{-\frac{n}{2}} e^{\frac{\langle \xi \rangle}{4h} (x-y)^2} \int_{\mathbb{R}^n} e^{-\frac{\langle \xi \rangle}{2h} [(x-z)^2 + (z-y)^2]} \cdot \langle \xi \rangle^{\frac{n}{4}} b(x-z, \xi) dz \\
 &= h^{-\frac{n}{2}} e^{\frac{\langle \xi \rangle}{4h} (x-y)^2} \int_{\mathbb{R}^n} e^{-\frac{\langle \xi \rangle}{2h} [(x-w-y)^2 + w^2]} \cdot \langle \xi \rangle^{\frac{n}{4}} b(x-w-y, \xi) dw \\
 &= h^{-\frac{n}{2}} \langle \xi \rangle^{\frac{n}{4}} \int_{\mathbb{R}^n} e^{-\frac{\langle \xi \rangle}{h} v^2} b\left(\frac{1}{2}(x-y) - v, \xi, h\right) dv \\
 &= (2\pi)^{-n} \left(1 + \frac{i}{4} \langle x-y, \xi / \langle \xi \rangle \rangle\right).
 \end{aligned}$$

The proof is now concluded using (2.7) below. □

For the reader’s convenience we include the derivation of Lebeau’s inversion formula used in the proof of Proposition 2.3 (see [20, (9.6.7)]):

LEMMA 2.4. For $u \in C_c^\infty(\mathbb{R}^n)$,

$$(2.7) \quad u(x) = (2\pi h)^{-n} \int_{\mathbb{R}^{2n}} e^{\frac{i}{h} \langle x-y, \xi + ia \langle \xi \rangle (x-y) \rangle} \cdot (1 + ia \langle x-y, \xi / \langle \xi \rangle \rangle) u(y) dy d\xi, \quad a > 0.$$

PROOF. For $u \in C_c^\infty(\mathbb{R}^n)$ the Fourier inversion formula gives

$$u(x) = (2\pi h)^{-n} \lim_{\epsilon \rightarrow 0^+} \int e^{\frac{i}{h} \langle (x-y, \xi) + i\epsilon \langle \xi \rangle \rangle} u(y) dy d\xi,$$

where the integral converges absolutely for $\epsilon > 0$. We deform the contour of integration in ξ to $\Gamma_a(x, y)$ given by

$$\xi \mapsto \eta := \xi + ai \langle \xi \rangle (x-y), \quad \xi \in \mathbb{R}^n, \quad 0 < a \ll 1.$$

This deformation is justified since on Γ ,

$$\text{Im} \langle x-y, \eta \rangle \geq c \langle \eta \rangle (x-y)^2,$$

and for a sufficiently small, $\langle \eta \rangle := (1 + \eta^2)^{\frac{1}{2}}$ has an analytic branch with positive real part. In particular, we have, using that $d \langle \xi \rangle = \sum_i \langle \xi \rangle^{-1} \xi_i d \xi_i$,

$$\begin{aligned} u(x) &= (2\pi h)^{-n} \lim_{\epsilon \rightarrow 0} \int_{\Gamma_a} \int_{\mathbb{R}^n} e^{\frac{i}{h} \langle (x-y, \eta) \rangle + i\epsilon \langle \eta \rangle} u(y) dy d\eta_1 \wedge d\eta_2 \wedge \dots \wedge d\eta_n = \\ &= (2\pi h)^{-n} \lim_{\epsilon \rightarrow 0} \int_{\mathbb{R}^{2n}} e^{\frac{i}{h} \langle (x-y, \xi + ia(x-y)) \rangle + i\epsilon \langle \eta \rangle} \det(\eta_\xi) u(y) dy d\xi \\ &= (2\pi h)^{-n} \int_{\mathbb{R}^{2n}} e^{\frac{i}{h} \langle x-y, \xi + ia(x-y) \rangle} (1 + ia \langle x-y, \xi / \langle \xi \rangle \rangle) u(y) dy d\xi. \end{aligned}$$

Since the right-hand side is analytic in $\{a \in \mathbb{C} : \text{Re } a > 0\}$ it follows that the formula remains valid for all $a > 0$. □

3 Geometry of Complex Deformations

Following [18,33] we will study the FBI transform (2.1) when $T^*\mathbb{T}^n$ is replaced by an I -Lagrangian \mathbb{R} -symplectic manifold submanifold of

$$T^*\widetilde{\mathbb{T}}^n := \{(z, \zeta) \mid z \in \mathbb{C}^n / 2\pi\mathbb{Z}^n, \zeta \in \mathbb{C}^n\} \simeq T^*(\mathbb{C}^n / 2\pi\mathbb{Z}^n).$$

We recall that $T^*\widetilde{\mathbb{T}}^n$ is equipped with the complex symplectic form

$$\sigma := d\zeta \wedge dz := \sum_{j=1}^n d\zeta_j \wedge dz_j = d(\zeta \cdot dz).$$

For a real $2n$ -dimensional submanifold of $T^*\widetilde{\mathbb{T}}^n$, Λ , we say

$$\Lambda \text{ is } I\text{-Lagrangian} \iff \text{Im}(\sigma|_\Lambda) \equiv 0,$$

and that

$$\Lambda \text{ is } \mathbb{R}\text{-symplectic} \iff \text{Re}(\sigma|_\Lambda) \text{ is nondegenerate.}$$

The specific submanifolds used here are as follows: For a function $G(x, \xi) \in C^\infty(T^*\mathbb{T}^n; \mathbb{R})$, assume that for some sufficiently small ϵ_0 (to be chosen in the constructions below),

$$(3.1) \quad \sup_{|\alpha|+|\beta| \leq 2} \langle \xi \rangle^{-1+|\beta|} |\partial_x^\alpha \partial_\xi^\beta G(x, \xi)| \leq \epsilon_0, \quad |\partial_x^\alpha \partial_\xi^\beta G(x, \xi)| \leq C_{\alpha\beta} \langle \xi \rangle^{1-|\beta|}.$$

(The second condition merely states that $G \in S^1(T^*\mathbb{T}^n)$ in the standard notation of [21].) We then define

$$(3.2) \quad \Lambda := \{(x + iG_\xi(x, \xi), \xi - iG_x(x, \xi)) \mid (x, \xi) \in T^*\mathbb{T}^n\} \subset T^*\widetilde{\mathbb{T}}^n.$$

By considering $G(x, \xi)$ as a periodic function of x , we can also think of Λ as a submanifold of $T^*\mathbb{C}^n$.

A submanifold given by (3.2) is always I -Lagrangian:

$$\begin{aligned} \zeta \cdot dz|_\Lambda &= (\xi - iG_x) \cdot d(x + iG_\xi) \\ &= \xi \cdot dx + G_x \cdot G_{\xi\xi} d\xi + G_x \cdot G_{\xi x} dx + i(-G_x \cdot dx + \xi \cdot dG_\xi) \end{aligned}$$

and

$$d(-G_x \cdot dx + \xi \cdot dG_\xi) = d\xi \wedge G_{\xi x} dx + dx \wedge G_{x\xi} d\xi = 0.$$

The smallness of ϵ_0 enters for the first time in guaranteeing that Λ is \mathbb{R} -symplectic:

$$\begin{aligned} & d(\xi \cdot dx + G_x \cdot G_{\xi\xi} d\xi + G_x \cdot G_{\xi x} dx) \\ &= d\xi \wedge dx + G_{xx} dx \wedge G_{\xi\xi} d\xi + G_{x\xi} d\xi \wedge G_{\xi\xi} d\xi \\ &+ G_{x\xi} d\xi \wedge G_{\xi x} dx + G_{xx} dx \wedge G_{\xi x} dx. \end{aligned}$$

The left-hand side is nondegenerate if ϵ_0 in (3.1) is small enough.

Since $\text{Im } \zeta \cdot dz|_\Lambda$ is closed, there exists $H \in C^\infty(\Lambda; \mathbb{R})$ such that

$$(3.3) \quad dH = -\text{Im } \zeta dz|_\Lambda,$$

with the normalization $H \equiv 0$ when $G \equiv 0$. Using the parametrization (3.2), we have the following explicit expression for

$$(3.4) \quad H(x, \xi) = G(x, \xi) - \xi \cdot G_\xi(x, \xi).$$

Any I -Lagrangian and \mathbb{R} -symplectic manifold is automatically maximally totally real in the sense that

$$T_\rho \Lambda \cap iT_\rho \Lambda = \{0\}, \quad \rho \in \Lambda.$$

In fact, suppose that $X, iX \in T_\rho \Lambda$; then for all $Y \in T_\rho \Lambda$,

$$\text{Re } \sigma(Y, iX) = -\text{Im } \sigma(Y, X) = 0$$

as $\text{Im } \sigma$ vanishes on $T_\rho \Lambda$. But then the nondegeneracy of $\text{Re } \sigma$ shows that $X = 0$.

The real symplectic form on Λ defines a natural volume form

$$dm(\alpha) = (\sigma|_\Lambda)^n/n!.$$

If $(z, \zeta) = (x + iG_\xi, \xi - iG_x)$, we sometimes write

$$(3.5) \quad dm_\Lambda(\alpha) = dz d\zeta = d\alpha, \quad \alpha = (z, \zeta) \in \Lambda, \quad \beta = \text{Re } \alpha.$$

Let Γ be a small conic connected neighbourhood of $T^*\mathbb{T}^n$ in $T^*\mathbb{C}^n/\mathbb{Z}^n$, and let $\tilde{G}(z, \zeta)$ be a symbolic almost analytic extension of $G(x, \xi)$ supported in Γ :

$$|\bar{\partial}_z \tilde{G}(z, \zeta)| + \langle \text{Re } \zeta \rangle \bar{\partial}_\zeta \tilde{G}(z, \zeta) \leq \langle \text{Re } \zeta \rangle \mathcal{O}(|\text{Im } z|^\infty + |\text{Im } \zeta / \langle \text{Re } \zeta \rangle|^\infty),$$

$$\sup_{|\alpha|+|\beta| \leq 2} |\partial_z^\alpha \partial_\zeta^\beta \tilde{G}(z, \zeta)| \leq C\epsilon_0 \langle \text{Re } \zeta \rangle^{1-|\beta|},$$

$$|\partial_z^\alpha \partial_\zeta^\beta \tilde{G}_y(z, \zeta)| \leq C_{\alpha\beta} \langle \text{Re } \zeta \rangle^{1-|\beta|},$$

for $(z, \zeta) \in \Gamma$ —see [25, theorem 1.3] (for a brief review of basic concepts of almost analytic machinery; see Appendix A).

We use an almost analytic change of variables in Γ to identify the totally real submanifold Λ with $T^*\mathbb{R}^n$ (on Λ the differentials of that transformation are complex linear): it is the inverse of the map

$$(3.6) \quad F : (z, \zeta) \mapsto (w, \omega) := (z + i\tilde{G}_\zeta(z, \zeta), \zeta - i\tilde{G}_z(z, \zeta)),$$

Using this identification we define

$$(3.7) \quad C_\Lambda(w, \omega) = F(\overline{F^{-1}(w, \omega)}), \quad (w, \omega) \in \Gamma, \quad C_\Lambda|_\Lambda = I_\Lambda.$$

We also denote by σ_Λ the almost analytic extension of $\sigma|_\Lambda$ to Γ .

NOTATION: The different identifications lead to potentially confusing notational issues. We will typically use coordinates

$$\alpha = (\alpha_x, \alpha_\xi) = (x, \xi) \mapsto \beta = (\beta_x, \beta_\xi) = (x + iG_\xi(x, \xi), \xi - iG_x(x, \xi)) \in \Lambda$$

and consider the complexification of α using the identification (3.6). In that case for $\alpha \in \Gamma$, $\bar{\alpha}$ denotes $C_\Lambda(\alpha)$. It is *not* given by taking $(z, \zeta) \mapsto (\bar{z}, \bar{\zeta})$ in the original coordinates on $\widetilde{T}^*\mathbb{T}^n$ (for one thing, it would not be the identity on Λ). Sometimes it is convenient to use $\beta \in \Lambda$ as the variable in formulae and integrations. The choice should be clear from the context.

4 Complex Deformations of the FBI Transform

For Λ given by (3.2) we define an operator T_Λ by prescribing its Schwartz kernel:

$$T_\Lambda(z, \zeta, y) := T((z, \zeta), y)|_{(z, \zeta) \in \Lambda}.$$

We then define an operator S_Λ by

$$S_\Lambda v(y) := \int_\Lambda S(x, \beta)v(\beta)d\beta, \quad \beta = (z, \zeta) \in \Lambda, \quad d\beta = dz \wedge d\zeta|_\Lambda,$$

where $S(x, z, \zeta)$ is the kernel of the operator S :

$$(4.1) \quad S(x, z, \zeta) := h^{-\frac{3n}{4}} \sum_{k \in \mathbb{Z}^n} e^{-\frac{i}{h}\phi^*(z-2\pi k, x, \zeta)} b(z-x-2\pi k, \zeta),$$

with b given in (2.3).

Note that if we parametrize Λ as in (3.2) with $\alpha = (x, \xi)$, we may also write

$$S_\Lambda v(y) := \int_{T^*\mathbb{T}^n} S_\Lambda(x, z(\alpha), \zeta(\alpha))v(\alpha)dm_\Lambda(\alpha),$$

where $dz \wedge d\zeta|_\Lambda = dm_\Lambda(\alpha)$. Finally, we sometimes write $\alpha_x = z(\alpha)$ and $\alpha_\xi = \zeta(\alpha)$.

In order to make sense of the composition $S_\Lambda T_\Lambda$, we start by analyzing T_Λ on a space of analytic functions on \mathbb{T}^n . For $\delta \geq 0$ let

$$(4.2) \quad \mathcal{A}_\delta = \left\{ u \in L^2(\mathbb{T}^n) : \|u\|_{\mathcal{A}_\delta}^2 := \sum_{n \in \mathbb{Z}^n} |\hat{u}(n)|^2 e^{4|n|\delta} < \infty \right\},$$

$$\hat{u}(n) := \frac{1}{(2\pi)^n} \int_{\mathbb{T}^n} u(x)e^{-i\langle x, n \rangle} dx.$$

Let also $\mathcal{A}_{-\delta}$ denote the dual space of \mathcal{A}_δ . Note that $\mathcal{A}_{-\delta}$ is a space of hyperfunctions, but on tori it can be identified with formal Fourier series with coefficients satisfying

$$\sum_{n \in \mathbb{Z}^n} |\widehat{u}(n)|^2 e^{-4|n|\delta} < \infty.$$

(In that case $\widehat{u}(n)$ can be defined using the pairing of the hyperfunction u with the analytic function $x \mapsto e^{-i\langle x, n \rangle} / (2\pi)^n$.) We note that $u \in \mathcal{A}_\delta$ extends to a (periodic) holomorphic function in $|\operatorname{Im} z| < 2\delta$ and (by the Fourier inversion formula and the Cauchy–Schwarz inequality),

$$(4.3) \quad \forall \delta' < 2\delta \exists C \text{ such that for } u \in \mathcal{A}_\delta, \quad \sup_{|\operatorname{Im} z| < \delta'} |u(z)| \leq C \|u\|_{\mathcal{A}_\delta}.$$

LEMMA 4.1. *Define*

$$\Omega_\delta := \{(z, \zeta) \in T^*(\mathbb{C}^n / 2\pi \mathbb{Z}^n) : |\operatorname{Im} \zeta| \leq \delta \langle \operatorname{Re} \zeta \rangle, |\operatorname{Im} z| \leq \delta, |\operatorname{Re} \zeta| \geq 1\}.$$

There exist $c_0, \delta_0 > 0$ such that for $(z, \zeta) \in \Omega_\delta$ and $0 < \delta < \delta_0$,

$$(4.4) \quad |Tu(z, \zeta)| \leq e^{-\delta c_0 |\zeta|/h} \|u\|_{\mathcal{A}_\delta}, \quad |S^t u(z, \zeta)| \leq e^{-c_0 \delta |\zeta|/h} \|u\|_{\mathcal{A}_\delta}.$$

where S^t is defined by

$$S^t u(z, \zeta) := \int_{\mathbb{T}^n} S(y, z, \zeta) u(y) dy$$

where the kernel S is defined in (4.1).

PROOF. Extending u to a periodic function on \mathbb{R}^n , we write

$$Tu(z, \zeta) = h^{-\frac{3n}{4}} \int_{\mathbb{R}^n} e^{\frac{i}{h}(\langle z-y, \zeta \rangle + \frac{i}{2} \langle \zeta \rangle (z-y)^2)} \langle \zeta \rangle^{\frac{n}{4}} u(y) dy.$$

Since u is analytic on $|\operatorname{Im} y| \leq \delta$, we may deform the contour in the y integration to $\Gamma(z, \zeta)$ given by

$$w \mapsto y(w) = w + z - i\delta \frac{\operatorname{Re} \zeta}{\langle \operatorname{Re} \zeta \rangle}, \quad w \in \mathbb{R}^n.$$

Then,

$$Tu(z, \zeta) = h^{-\frac{3n}{4}} \int e^{\frac{i}{h}(\langle -w + i\delta \frac{\operatorname{Re} \zeta}{\langle \operatorname{Re} \zeta \rangle}, \zeta \rangle + \frac{i}{2} \langle \zeta \rangle (w - i\delta \frac{\operatorname{Re} \zeta}{\langle \operatorname{Re} \zeta \rangle})^2)} \langle \zeta \rangle^{\frac{n}{4}} u(y(w)) dw.$$

For $|\operatorname{Im} \zeta| \leq \delta \langle \operatorname{Re} \zeta \rangle$, $|\operatorname{Re} \zeta| \geq 1$, with $\delta < \delta_0$ and δ_0 small enough,

$$\operatorname{Re} \langle \zeta \rangle \geq \frac{1}{2} |\zeta|, \quad |\operatorname{Im} \langle \zeta \rangle| \leq \frac{1}{16} |\zeta|, \quad |\zeta| \geq \frac{1}{2}.$$

Hence for $w \in \mathbb{R}$ and $(z, \zeta) \in \Omega_\delta$ the real part of the phase in the integral above is bounded by

$$-\frac{1}{2} \delta |\zeta| + \frac{1}{16} \delta |w| |\zeta| - \frac{1}{4} (|w|^2 - \delta^2) |\zeta| + \frac{1}{16} \delta |\zeta| \leq -c_0 \delta |\zeta| - c_0 |w|^2, \quad c_0 > 0.$$

In view of (4.3) the integrand is then bounded by $\exp(-c_0(\delta|\zeta| + |w|^2)/h) \|u\|_{\mathcal{A}_\delta}$, which gives the first bound in (4.4). The proof for S^t is identical since the phase agrees with that of T . □

A natural Hilbert space on the FBI transform side is defined by the norm

$$\|v\|_{L^2(\Lambda)}^2 = \int_{\Lambda} |v(\alpha)|^2 e^{-2H(\alpha)/h} d\alpha.$$

The next lemma gives boundedness of S_{Λ} and T_{Λ}^t on exponentially decaying functions on Λ :

LEMMA 4.2. *There exist $\delta_0 > 0$ and $C_0 > 0$ big enough such that for $0 < \epsilon_0 < \delta_0$ in (3.1) we have*

$$S_{\Lambda} : e^{-C_0\delta\langle\xi\rangle/h} L^2(\Lambda) \rightarrow \mathcal{A}_{\delta}, \quad T_{\Lambda}^* : e^{-C_0\delta\langle\xi\rangle/h} L^2(\Lambda) \rightarrow \mathcal{A}_{\delta}.$$

for all $0 < \delta < \delta_0$, where the adjoint T_{Λ}^* is defined using the $L^2(\Lambda, e^{-2H/h})$ inner product.

PROOF. Let $v \in e^{-C_0\langle\xi\rangle/h} L^2(\Lambda)$ with $C_0 > 0$ be chosen large enough and $|\operatorname{Im} y| \leq a\delta$. Then,

$$S_{\Lambda} v(y) = h^{-\frac{3n}{4}} \int_{\Lambda} e^{\frac{i}{h}((y-\alpha_x, \alpha_{\xi}) + \frac{1}{2}(\alpha_{\xi})(\alpha_x - y)^2)} b(y - \alpha_x) v(\alpha) d\alpha.$$

where b is given in (2.3). Therefore, by the Cauchy–Schwarz inequality,

$$|S_{\Lambda} v(y)|^2 \leq C h^{-\frac{3n}{2}} J(y) \|e^{C_0\delta\langle\xi\rangle/h} v\|_{L^2(\Lambda)}^2,$$

where

$$J(y) := \int_{\Lambda} e^{-2\operatorname{Im}((y-\alpha_x, \alpha_{\xi}) + \frac{1}{2}(\alpha_{\xi})(\alpha_x - y)^2)/h + 2H(\alpha)/h} \cdot \langle |y - \alpha_x| \rangle^2 e^{-2C_0\delta\langle|\alpha_{\xi}|\rangle/h} d\alpha$$

Writing $\beta = \operatorname{Re} \alpha$ we now estimate

$$\begin{aligned} -\operatorname{Im}\langle y - \alpha_x, \alpha_{\xi} \rangle &= \langle G_{\xi} - \operatorname{Im} y, \beta_{\xi} \rangle - \langle \beta_x - \operatorname{Re} y, G_x \rangle \\ &\leq (a\delta + \epsilon_0) |\beta_{\xi}| + \epsilon_0 \langle \beta_{\xi} \rangle |\beta_x - \operatorname{Re} y|. \end{aligned}$$

Similarly,

$$\operatorname{Re}\langle \alpha_{\xi} \rangle (\alpha_x - y)^2 \leq -(1 - C\epsilon_0) \langle \beta_{\xi} \rangle (|\beta_x - \operatorname{Re} y|^2 - C\epsilon^2 - Ca^2\delta^2),$$

and $2H(\alpha) \leq C\epsilon_0 \langle \beta_{\xi} \rangle$ (see (3.4)). Hence for $C_0 \gg C$, the phase in $J(y)$ is bounded by

$$-C_1 \delta \langle \beta_{\xi} \rangle \langle \operatorname{Re} y - \beta_x \rangle^2, \quad C_1 > 0.$$

That proves $S_{\Lambda} v$ is analytic and uniformly bounded when $|\operatorname{Im} y| \leq a\delta$. In particular, $S_{\Lambda} v \in \mathcal{A}_{\delta}$. A similar argument applies to T_{Λ}^* . \square

Together, Lemmas 4.1 and 4.2 imply that there are $\delta_1, \delta_2 > 0$ such that $S_{\Lambda} T_{\Lambda}$ as well as an operator $\mathcal{A}_{\delta_1} \rightarrow \mathcal{A}_{\delta_2}$ and as an operator $\mathcal{A}_{-\delta_2} \rightarrow \mathcal{A}_{-\delta_1}$.

We can now show that $S_{\Lambda} T_{\Lambda}$ is the identity on \mathcal{A}_{δ} and $\mathcal{A}_{-\delta}$ for $\delta > 0$ small enough.

PROPOSITION 4.3. *There is $\delta_1 > 0$ such that for all $0 < |\delta| < \delta_1$, S_Λ , and T_Λ as above,*

$$S_\Lambda T_\Lambda = \text{Id} : \mathcal{A}_\delta \rightarrow \mathcal{A}_\delta.$$

PROOF. Assume first that $\delta > 0$ and let $v \in \mathcal{A}_\delta$. Then, by Lemma 4.1 for $\delta > 0$ small enough, $T_\Lambda v \in e^{-c_0\delta|\alpha_\xi|}L^2(\Lambda)$ and is given by

$$T_\Lambda v(\alpha) = \int_{\mathbb{T}^n} T_\Lambda(\alpha, y)v(y)dy.$$

Then, again for $\delta > 0$ small enough, Lemma 4.2 shows that $S_\Lambda T_\Lambda v$ is well-defined and given by

$$(4.5) \quad S_\Lambda T_\Lambda v(x) = \int_\Lambda \int_{\mathbb{T}^n} S_\Lambda(x, \alpha)T_\Lambda(\alpha, y)v(y)dy d\alpha.$$

The decay in $|\alpha_\xi|$ allows a contour deformation in α in (4.5) and then an application of Proposition 2.3. This gives

$$S_\Lambda T_\Lambda v(x) = \int_{T^*\mathbb{T}^n} \int_{\mathbb{T}^n} S(x, \alpha)T(\alpha, y)v(y)dy d\alpha = v(y), \quad v \in \mathcal{A}_\delta.$$

To define $T_\Lambda v$ for $v \in \mathcal{A}_{-\delta}$, $\delta > 0$, we note that Lemma 4.2 shows that if $w \in e^{-C_0\delta\langle\xi\rangle/h}L^2(\Lambda)$, then $T_\Lambda^*w \in \mathcal{A}_\delta$. Therefore,

$$\langle T_\Lambda v, w \rangle_{L^2(\Lambda)} := \langle v, T_\Lambda^*w \rangle_{L^2(\mathbb{T}^n)}$$

is well-defined and $T_\Lambda : \mathcal{A}_{-\delta} \rightarrow e^{C_0\delta\langle\xi\rangle/h}L^2(\Lambda)$.

For $u \in \mathcal{A}_{c_1\delta}$, $c_1 \gg 1$, $c_1\delta < \delta_0$ (with δ_0 of Lemma 4.1), we formally have

$$(4.6) \quad \langle S_\Lambda T_\Lambda v, u \rangle_{L^2(\mathbb{T}^n)} := \langle T_\Lambda v, S_\Lambda^*u \rangle_{L^2(\Lambda)}.$$

Since $S_\Lambda^*u = \overline{S^t|_\Lambda u}e^{2H(\alpha)/h}$ and $H(\alpha) \leq C\epsilon_0\langle\text{Re}\alpha_\xi\rangle$, Lemma 4.1 shows that

$$S_\Lambda^*u \in e^{C\epsilon_0\langle\xi\rangle/h - c_0c_1\delta|\xi|/h}L^2(\Lambda).$$

and hence for $c_1 > 0$ large enough (and δ_1 small enough so that $c_1\delta_1 < \delta_0$), the pairing on the right-hand side of (4.6) is well-defined and

$$\langle S_\Lambda T_\Lambda v, u \rangle = \langle v, T_\Lambda^*S_\Lambda^*u \rangle.$$

We can now deform the contour in the α integral, which gives

$$T_\Lambda^*S_\Lambda^*u(x) = \int_{\mathbb{T}^n} \int_\Lambda T_\Lambda(\alpha, x)S_\Lambda(y, \alpha)u(y)dy d\alpha = u(y).$$

Hence for $v \in \mathcal{A}_\delta$ and $u \in \mathcal{A}_{c_1\delta}$, $\langle S_\Lambda T_\Lambda v, u \rangle_{L^2(\mathbb{T}^n)} = \langle v, u \rangle_{L^2(\mathbb{T}^n)}$. Since $\mathcal{A}_{c_1\delta}$, $c_1 \geq 1$, is dense in \mathcal{A}_δ , the claim follows. \square

We now define natural spaces on which T_Λ, S_Λ act:

DEFINITION 4.4. Let δ_0 be as in Lemma 4.1. We define the Sobolev space of order t adapted to Λ as

$$(4.7) \quad H_\Lambda^t := \overline{\mathcal{A}_{\delta_0}}^{\|\cdot\|_{H_\Lambda^t}}, \quad \|u\|_{H_\Lambda^t}^2 := \int_\Lambda \langle \operatorname{Re} \alpha_\xi \rangle^{2t} |T_\Lambda u(\alpha)|^2 e^{-2H(\alpha)/h} d\alpha$$

where we used the notation from (3.5) and (3.3). We then have an isometry

$$T_\Lambda : H_\Lambda^t \rightarrow \langle \xi \rangle^{-t} L^2(\Lambda),$$

where the notation on the right-hand side is the shorthand for $\langle \operatorname{Re} \alpha_\xi \rangle^{-t}$.

Remarks 4.5.

1. There exists $\delta > 0$ such that

$$\mathcal{A}_\delta \subset H_\Lambda^m \subset \mathcal{A}_{-\delta}.$$

The left inclusion is immediate from the definition. On the other hand, for $u \in H_\Lambda^m$, $T_\Lambda u \in \langle \xi \rangle^m L^2(\Lambda)$, and in particular, by Lemma 4.2 $S_\Lambda T_\Lambda u \in \mathcal{A}_{-\delta}$ for some $\delta > 0$. But $S_\Lambda T_\Lambda u = u$ and hence $u \in \mathcal{A}_{-\delta}$.

2. Let Π_Λ denote the orthogonal projection from $L^2(\Lambda) \rightarrow T_\Lambda(H_\Lambda^0)$. The properties of Π_Λ show that $T_\Lambda(H_\Lambda^t) = \Pi_\Lambda(\langle \xi \rangle^{-t} L^2(\Lambda))$.

Lemmas 4.1 and 4.2 show that (with h -dependent norms and changing c_0 to $c_0/2$),

$$T_\Lambda : \mathcal{A}_\delta \rightarrow e^{-\delta c_0 \langle \xi \rangle} L^2(\Lambda), \quad S_\Lambda : e^{-\delta C_0 \langle \xi \rangle / h} L^2(\Lambda) \mapsto \mathcal{A}_\delta.$$

This means that

$$(4.8) \quad T_\Lambda S_\Lambda : e^{-\delta C_0 \langle \xi \rangle / h} L^2(\Lambda) \rightarrow e^{-\delta c_0 \langle \xi \rangle} L^2(\Lambda).$$

PROPOSITION 4.6. *The operator $T_\Lambda S_\Lambda$ in (4.8) extends to an operator*

$$T_\Lambda S_\Lambda = \mathcal{O}(1) : \langle \xi \rangle^m L^2(\Lambda) \rightarrow \langle \xi \rangle^m L^2(\Lambda).$$

Moreover, there are $k \in S^0(\Lambda \times \Lambda)$, $\alpha, \beta \in \Lambda$, $\chi \in C_c^\infty(\mathbb{R})$ such that for all $\delta > 0$, there is $\epsilon_1 > 0$ such that for G satisfying (3.1) with $\epsilon_0 < \epsilon_1$,

$$T_\Lambda S_\Lambda = K_\Lambda + O_N(e^{-C_\delta/h})_{\langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)},$$

where the Schwartz kernel of K_Λ is given by

$$K_\Lambda(\alpha, \beta) = h^{-n} e^{\frac{i}{h} \Psi(\alpha, \beta)} k(\alpha, \beta) \tilde{\chi}(\alpha, \beta),$$

$$\begin{aligned} \tilde{\chi}(\alpha, \beta) &:= \chi(\delta^{-1} d(\operatorname{Re} \alpha_x, \operatorname{Re} \beta_x)) \\ &\cdot \chi(\delta^{-1} \min(\langle \operatorname{Re} \beta_\xi \rangle, \langle \operatorname{Re} \alpha_\xi \rangle)^{-1} |\operatorname{Re} \alpha_\xi - \operatorname{Re} \beta_\xi|), \end{aligned}$$

and

$$(4.9) \quad \begin{aligned} \Psi &= \frac{i}{2} \frac{(\alpha_\xi - \beta_\xi)^2}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} + \frac{i}{2} \frac{\langle \beta_\xi \rangle \langle \alpha_\xi \rangle (\alpha_x - \beta_x)^2}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} \\ &+ \frac{\langle \beta_\xi \rangle \alpha_\xi + \langle \alpha_\xi \rangle \beta_\xi}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} \cdot (\alpha_x - \beta_x). \end{aligned}$$

We will prove the proposition in two lemmas which for future use are formulated in greater generality. We first study the kernel of the composition $T_\Lambda S_\Lambda$.

LEMMA 4.7. *Let Λ_{G_1} and Λ_{G_2} be given by (3.2) with G_i satisfying (3.1). Then there are $\chi \in C_c^\infty(\mathbb{R})$ and $k \in S^0(\Lambda_{G_2} \times \Lambda_{G_1})$ such that for all $\delta > 0$, there is $\epsilon_1 > 0$ such that for G_1 and G_2 satisfying (3.1) with $\epsilon_0 < \epsilon_1$,*

$$T_{\Lambda_{G_2}} S_{\Lambda_{G_1}} = K + O_N(e^{-C\delta/h})_{\langle \xi \rangle^N L^2(\Lambda_1) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda_2)},$$

where the Schwartz kernel of K is given by

$$h^{-n} e^{\frac{i}{h} \Psi(\alpha, \beta)} k(\alpha, \beta) \chi(\delta^{-1} d(\operatorname{Re} \alpha_x, \operatorname{Re} \beta_x)) \cdot \chi(\delta^{-1} \min(\langle \operatorname{Re} \alpha_\xi \rangle, \langle \operatorname{Re} \beta_\xi \rangle)^{-1} |\operatorname{Re} \alpha_\xi - \operatorname{Re} \beta_\xi|),$$

$(\alpha, \beta) \in \Lambda_{G_2} \times \Lambda_{G_1}$, and where Ψ is as in (4.9).

PROOF. The kernel of $T_{\Lambda_{G_2}} S_{\Lambda_{G_1}}$ (again extending everything to be periodic on \mathbb{R}^n and using integration with respect to $d\beta = (\sigma|_\Lambda)^n/n!$) is given by

$$h^{-\frac{3n}{2}} \int_{\mathbb{R}^n} e^{\frac{i}{h}(\varphi(\alpha, y) - \varphi^*(\beta, y))} \langle \alpha_\xi \rangle^{\frac{n}{4}} b(\beta_x - y, \beta_\xi) dy,$$

where $\alpha \in \Lambda_{G_1}$, $\beta \in \Lambda_{G_2}$, and b is given by (2.3). To analyse it, we first observe that for ϵ_0 small enough

$$\begin{aligned} & \operatorname{Im}(\varphi(\alpha, y) - \varphi^*(\beta, y)) \\ & \geq \frac{1}{4} |\langle \alpha_\xi \rangle| (|\operatorname{Re}(\alpha_x - y)|^2 - |\operatorname{Im}(\alpha_x - y)|^2) \\ & \quad + \frac{1}{4} |\langle \beta_\xi \rangle| (|\operatorname{Re}(\beta_x - y)|^2 - |\operatorname{Im}(\beta_x - y)|^2) \\ & \quad + \operatorname{Im}\langle \alpha_x - y, \alpha_\xi \rangle + \operatorname{Im}\langle y - \beta_x, \beta_\xi \rangle \end{aligned}$$

Now, fix $\delta > 0$, and assume that $|\operatorname{Im} y| \leq \delta$. Then for $\epsilon_0 \leq \delta$ in (3.1), we have

$$\begin{aligned} & \operatorname{Im}(\varphi(\alpha, y) - \varphi^*(\beta, y)) \\ & \geq c |\langle \alpha_\xi \rangle| |\operatorname{Re} \alpha_x - \operatorname{Re} y|^2 + c |\langle \beta_\xi \rangle| |\operatorname{Re} \beta_x - \operatorname{Re} y|^2 \\ & \quad - C\delta^2 (|\langle \alpha_\xi \rangle| + |\langle \beta_\xi \rangle|) + \operatorname{Im}\langle \alpha_x - y, \alpha_\xi \rangle + \operatorname{Im}\langle y - \beta_x, \beta_\xi \rangle. \end{aligned}$$

Therefore, deforming the contour in y using

$$y \mapsto y + \frac{i\delta(\overline{\beta_\xi} - \alpha_\xi)}{\langle \operatorname{Re}(\beta_\xi - \alpha_\xi) \rangle}, \quad y \in \mathbb{R}^n,$$

we have (on the new contour)

$$\begin{aligned} & \operatorname{Im}(\varphi(\alpha, y) - \varphi^*(\beta, y)) \\ & \geq c |\langle \alpha_\xi \rangle| |\operatorname{Re} \alpha_x - \operatorname{Re} y|^2 + c |\langle \beta_\xi \rangle| |\operatorname{Re} \beta_x - \operatorname{Re} y|^2 + \delta \frac{|\alpha_\xi - \beta_\xi|^2}{\langle \operatorname{Re}(\beta_\xi - \alpha_\xi) \rangle} \\ & \quad - C\delta^2 (|\langle \alpha_\xi \rangle| + |\langle \beta_\xi \rangle|) + \operatorname{Im}\langle \alpha_x, \alpha_\xi \rangle - \operatorname{Im}\langle \beta_x, \beta_\xi \rangle. \end{aligned}$$

Using

$$|\operatorname{Im} \beta_x| + |\operatorname{Im} \alpha_x| + |\langle \alpha_\xi \rangle|^{-1} |\operatorname{Im} \alpha_\xi| + |\langle \beta_\xi \rangle|^{-1} |\operatorname{Im} \beta_\xi| \leq C \epsilon_0 \ll \delta,$$

we then obtain

$$\begin{aligned} \operatorname{Im}(\varphi(\alpha, y) - \varphi^*(\beta, y)) &\geq c |\langle \alpha_\xi \rangle| |\operatorname{Re} \alpha_x - y|^2 + c |\langle \beta_\xi \rangle| |\operatorname{Re} \beta_x - y|^2 \\ &\quad + c \delta \frac{|\alpha_\xi - \beta_\xi|^2}{\langle \operatorname{Re}(\beta_\xi - \alpha_\xi) \rangle} - C \delta^2 (|\langle \alpha_\xi \rangle| + |\langle \beta_\xi \rangle|) \end{aligned}$$

In particular, when

$$|\operatorname{Re} \alpha_x - \operatorname{Re} \beta_x| \geq \delta \quad \text{or} \quad |\operatorname{Re} \alpha_\xi - \operatorname{Re} \beta_\xi| \geq 2c\delta \min(\langle \operatorname{Re} \alpha_\xi \rangle, \langle \operatorname{Re} \beta_\xi \rangle) / C,$$

the integrand is bounded by

$$e^{-\langle \operatorname{Re} \alpha_\xi \rangle + \langle \operatorname{Re} \beta_\xi \rangle} (1 + |\operatorname{Re} \alpha_x - \operatorname{Re} \beta_x|) / C h.$$

Therefore, modulo an $O_N(e^{-C/h})_{\langle \xi \rangle^N L^2(\Lambda_1) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda_2)}$ error, the kernel is given by

$$k(\alpha, \beta) := h^{-\frac{3n}{2}} \int_{\mathbb{R}^n} e^{\frac{i}{h}(\varphi(\alpha, y) - \varphi^*(\beta, y))h} k_1(\alpha, \beta, y) \tilde{\chi} dy,$$

$$\tilde{\chi} := \chi(\delta^{-1} d(\alpha_x, \beta_x)) \chi(\delta^{-1} \min(\langle \operatorname{Re} \alpha_\xi \rangle, \langle \operatorname{Re} \beta_\xi \rangle)^{-1} |\alpha_\xi - \beta_\xi|),$$

where χ is a suitable cutoff function and

$$k_1 \in \langle \operatorname{Re} \alpha_\xi \rangle^{\frac{n}{4}} \langle \operatorname{Re} \beta_\xi \rangle^{\frac{n}{4}} S^0(\Lambda_{G_2} \times \Lambda_{G_1} \times \mathbb{R}^n),$$

and the dependence on the last variable is periodic and holomorphic on $|\operatorname{Im} y| \leq c$.

We claim that $k(\alpha, \beta)$ is given by

$$(4.10) \quad h^{-n} e^{\frac{i}{h} \Psi(\alpha, \beta)} k(\alpha, \beta) \tilde{\chi},$$

where $k \in S^0(\Lambda_{G_2} \times \Lambda_{G_2})$. To see this, we note that the critical point in y is given by

$$y_c = \frac{i(\beta_\xi - \alpha_\xi) + \langle \alpha_\xi \rangle \alpha_x + \langle \beta_\xi \rangle \beta_x}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle}.$$

We then deform the contour to $y \mapsto y + y_c$. The phase becomes

$$\begin{aligned} (\alpha_x - \beta_x) \frac{\langle \alpha_\xi \rangle \beta_\xi + \alpha_\xi \langle \beta_\xi \rangle}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} + \frac{i(\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle)}{2} y^2 \\ + \frac{i}{2} \frac{\langle \alpha_\xi \rangle \langle \beta_\xi \rangle (\beta_x - \alpha_x)^2}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} + \frac{i}{2} \frac{(\beta_\xi - \alpha_\xi)^2}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle}. \end{aligned}$$

and the method of steepest descent gives (4.10). \square

The next lemma gives the first part of Proposition 4.6:

LEMMA 4.8. *For all $m \in \mathbb{R}$, there are $C, h_0 > 0$ such that for $0 < h < h_0$,*

$$\|T_\Lambda S_\Lambda\|_{\langle \xi \rangle^m L^2(\Lambda) \rightarrow \langle \xi \rangle^m L^2(\Lambda)} \leq C.$$

PROOF. By Lemma 4.7, we need to show uniform boundedness of K with the kernel given by

$$K(\alpha, \beta) = h^{-n} e^{\frac{i}{h}\Psi(\alpha, \beta)} k(\alpha, \beta) \chi(\delta^{-1} d(\operatorname{Re} \alpha_x, \operatorname{Re} \beta_x)) \cdot \chi\left(\frac{|\operatorname{Re} \alpha_\xi - \operatorname{Re} \beta_\xi|}{\delta \min(\langle \operatorname{Re} \alpha_\xi \rangle, \langle \operatorname{Re} \beta_\xi \rangle)}\right).$$

where Ψ is as in (4.9), $k \in S^0$.

In particular, conjugating by $\langle \operatorname{Re} \alpha_\xi \rangle^m e^{H(\alpha)/h}$, we need to show that the operator with the kernel

$$h^{-n} e^{\frac{i}{h}(\Psi(\alpha, \beta) - iH(\beta) + iH(\alpha))} \tilde{k}(\alpha, \beta)$$

is bounded on $L^2(\Lambda, dm(\alpha))$ where

$$\tilde{k}(\alpha, \beta) := \left(\frac{\langle \operatorname{Re} \alpha_\xi \rangle}{\langle \operatorname{Re} \beta_\xi \rangle}\right)^m k(\alpha, \beta) \chi(\delta^{-1} |\operatorname{Re} \alpha_x - \operatorname{Re} \beta_x|) \cdot \chi\left(\frac{|\operatorname{Re} \alpha_\xi - \operatorname{Re} \beta_\xi|}{\delta \min(\langle \operatorname{Re} \alpha_\xi \rangle, \langle \operatorname{Re} \beta_\xi \rangle)}\right).$$

To establish this we define

$$(4.11) \quad \Phi(\alpha, \beta) := \Psi(\alpha, \beta) - iH(\beta) + iH(\alpha),$$

where we see that $\Phi(\alpha, \alpha) = 0$. Next, we note that

$$(4.12) \quad d_\alpha \Phi|_{\alpha=\beta} = \alpha_\xi d\alpha_x + i d_\alpha H.$$

Therefore (see (3.3)), $\operatorname{Im} d_\alpha \Phi|_{\alpha=\beta} = 0$. Similarly, $\operatorname{Im} d_\beta \Phi|_{\alpha=\beta} = 0$ and hence $\operatorname{Im} \Phi$ vanishes quadratically at $\alpha = \beta$.

In the case of no deformation (that is, for $\Lambda = T^*\mathbb{T}^n$)

$$\operatorname{Im} \Phi \geq c \langle \alpha_\xi \rangle |\alpha_x - \beta_x|^2 + c \langle \alpha_\xi \rangle^{-1} |\alpha_\xi - \beta_\xi|^2, \quad \alpha, \beta \in T^*\mathbb{T}^n.$$

Since Λ is a small conic perturbation of $T^*\mathbb{T}^n$, this remains true on Λ . Hence,

$$\begin{aligned} |K(\alpha, \beta)| &\leq C h^{-n} e^{(c \langle \operatorname{Re} \alpha_\xi \rangle |\operatorname{Re} \alpha_x - \operatorname{Re} \beta_x|^2 + c \langle \operatorname{Re} \alpha_\xi \rangle^{-1} |\operatorname{Re} \alpha_\xi - \operatorname{Re} \beta_\xi|^2)/h} \\ &\quad \cdot \langle \operatorname{Re} \alpha_\xi \rangle^{\frac{n}{4}} \langle \operatorname{Re} \beta_\xi \rangle^{\frac{n}{4}} \tilde{\chi}, \\ \tilde{\chi} &= \chi(\delta^{-1} d(\operatorname{Re} \alpha_x, \operatorname{Re} \beta_x)) \chi(\delta^{-1} \min(\langle \operatorname{Re} \alpha_\xi \rangle, \langle \operatorname{Re} \beta_\xi \rangle))^{-1} \\ &\quad \cdot |\operatorname{Re} \alpha_\xi - \operatorname{Re} \beta_\xi|. \end{aligned}$$

The Schur's test for boundedness on L^2 then shows that K is uniformly bounded on $L^2(\Lambda)$. □

The following lemma shows that compact changes of the Lagrangian Λ change the norm on $L^2(\Lambda)$ but not the elements in the space.

LEMMA 4.9. *Let G_1 and G_2 satisfy (3.1). Then, for all $M, N > 0$,*

$$\mathbb{1}_{|\xi| \leq M} T_{\Lambda_{G_2}} S_{\Lambda_{G_1}} = \mathcal{O}_h(1) : \langle \xi \rangle^N L^2(\Lambda_{G_1}) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda_{G_2}).$$

PROOF. By Lemma 4.7 we only need to show that the operator $\mathbb{1}_{|\xi| \leq M} K$ is bounded. However, the structure of the Schwartz kernel described in that lemma shows that the kernel of $\mathbb{1}_{|\xi| \leq M} K$ is smooth and compactly supported. Except for a loss in the constant due to different weights, the boundedness follows. \square

5 Asymptotic Description of the Projector

The main part of this section consists of a construction of a parametrix for the orthogonal projector onto the (closure of the) image of T_Λ . It is inspired by [33, Section 1], which in turn followed ideas of [1, 2, 18, 25]. A detailed presentation in a simpler case of compactly supported weights can be found in [15, sec. 6], and it can be used as a guide to the more notationally involved case at hand. We then use the argument from [1, 33] to relate the parametrix to the exact projector.

5.1 The structure of the parametrix

We seek an operator of the following form:

$$(5.1) \quad \begin{aligned} B_\Lambda u(\alpha) &= h^{-n} \int_{T^*\mathbb{T}^n} e^{i\psi(\alpha, \beta)/h - 2H(\beta)/h} a(\alpha, \beta, h) u(\beta) dm_\Lambda(\beta), \\ dm_\Lambda(\beta) &:= (\sigma|_\Lambda)^n / n! = d\alpha, \quad \beta = \operatorname{Re} \alpha, \quad \alpha \in \Lambda, \end{aligned}$$

where ψ and a satisfy (for all $k, k', \ell, \ell' \in \mathbb{N}^n$)

$$(5.2) \quad \begin{aligned} \operatorname{supp} \psi, \operatorname{supp} a &\subset \{(\alpha, \beta) : d(\alpha_x, \beta_x) \leq \epsilon, \quad |\alpha_\xi - \beta_\xi| \leq \langle \alpha_\xi \rangle \epsilon\}, \\ \partial_{\alpha_x}^k \partial_{\alpha_\xi}^\ell \partial_{\beta_x}^{k'} \partial_{\beta_\xi}^{\ell'} \psi(\alpha, \beta) &= \mathcal{O}(\langle \alpha_\xi \rangle^{1-|\ell|-|\ell'|}), \quad \overline{\psi(\alpha, \beta)} = -\psi(\beta, \alpha), \end{aligned}$$

and

$$(5.3) \quad \begin{aligned} a(\alpha, \beta, h) &\sim \sum_{j=0}^\infty (\langle \alpha_\xi \rangle^{-1} h)^j a_j(\alpha, \beta), \quad \overline{a(\alpha, \beta)} = a(\beta, \alpha), \\ \partial_{\alpha_x}^k \partial_{\alpha_\xi}^\ell \partial_{\beta_x}^{k'} \partial_{\beta_\xi}^{\ell'} a_j(\alpha, \beta) &= \mathcal{O}(\langle \alpha_\xi \rangle^{-|\ell|-|\ell'|}). \end{aligned}$$

The basic properties we need are self-adjointness and idempotence:

$$(5.4) \quad B_\Lambda = B_\Lambda^{*,H}, \quad B_\Lambda \equiv B_\Lambda^2,$$

where $A \equiv B$ means that $A - B = \mathcal{O}(h^N)_{\langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)}$ for all N .

The deeper requirement comes from relating the image of B_Λ to that of T_Λ :

PROPOSITION 5.1. *Suppose that Z_j , differential operators with holomorphic coefficients in Γ , are defined by*

$$Z_j := \langle \zeta \rangle^{-1} (hD_{z_j} - \zeta_j) + \frac{1}{2} \langle \zeta \rangle^{-3} \zeta_j (hD_z - \zeta)^2 - ihD_{\zeta_j} - \frac{n}{4} h \langle \zeta \rangle^{-2} \zeta_j.$$

If

$$(5.5) \quad Z_j^\Lambda = Z_j|_\Lambda$$

in the sense of restriction of holomorphic operators to totally real submanifolds, then for $u \in \mathcal{A}_\delta$,

$$(5.6) \quad Z_j^\Lambda T_\Lambda u(\alpha) = 0, \quad j = 1, \dots, n.$$

PROOF. Putting

$$W_j = \langle \zeta \rangle^{-\frac{n}{4}} Z_j \langle \zeta \rangle^{\frac{n}{4}} = \langle \zeta \rangle^{-1} (hD_{z_j} - \zeta_j) + \frac{1}{2} \langle \zeta \rangle^{-3} \zeta_j (hD_z - \zeta)^2 - i h D_{\zeta_j} - \frac{n}{2} h \langle \zeta \rangle^{-2} \zeta_j,$$

we check that

$$W_j (e^{\frac{i}{h}((z-y+2\pi k, \zeta) + \frac{1}{2} \langle \zeta \rangle (z-y+2\pi k)^2)}) = 0$$

for all $y \in \mathbb{T}^n$ and $k \in \mathbb{Z}^n$. The definition of T_Λ then immediately gives (5.6). \square

We note that Z_j 's commute and hence we also have

$$[Z_j^\Lambda, Z_k^\Lambda] = 0.$$

We write

$$z_j^\Lambda := \langle \zeta \rangle^{-1} (z_j^* - \zeta_j) + \frac{1}{2} \langle \zeta \rangle^{-3} (z_j^* - \zeta)^2 \zeta_j - i \zeta_j^*|_\Lambda, \quad \{z_j^\Lambda, z_k^\Lambda\} = 0,$$

for the principal symbol of Z_j^Λ (in a sense that will be explained after the rescaling below). The vanishing of the Poisson bracket reflects the fact that z_j^Λ vanish on the involutive manifold $\{(\alpha, d_\alpha \varphi(\alpha, y) : \alpha \in \Lambda, y \in \mathbb{T}^n)\}$ —see Lemma 5.4 below.

Since B_Λ is supposed to be a parametrix for a self-adjoint projection onto the image T_Λ , Proposition 5.1 shows that we should have

$$(5.7) \quad Z_j^\Lambda B_\Lambda \equiv 0, \quad B_\Lambda (Z_j^\Lambda)^{*,H} \equiv 0,$$

where the definition of \equiv is given in (5.10) below.

To explore the second condition in terms of the kernel of B_Λ , we denote by A^* the formal adjoint of an operator A on $L^2(\Lambda, dm_\Lambda)$ (no weight). We also define a transpose of A by

$$\int_\Lambda Au(\alpha)v(\alpha)dm_\Lambda(\alpha) = \int_\Lambda u(\alpha)A^t v(\alpha)dm_\Lambda(\alpha).$$

We note the general fact $(A^*)^t = J \circ A \circ J$, $Ju := \bar{u}$. Then, with $K_\Lambda(\alpha, \beta) := h^{-n} e^{i\psi(\alpha, \beta)/h} a(\alpha, \beta, h)$,

$$\begin{aligned} (AB_\Lambda)^{*,H} u(\alpha) &= \int_\Lambda K_\Lambda(\alpha, \beta) A^*(e^{-2H(\bullet)/h} u(\bullet))(\beta) dm_\Lambda(\beta) \\ &= \int_\Lambda (A^*)^t (K_\Lambda(\alpha, \bullet))(\beta) e^{-2H(\beta)/h} u(\beta) dm_\Lambda(\beta) \\ &= \int_\Lambda (J \circ A \circ J)(K_\Lambda(\alpha, \bullet))(\beta) e^{-2H(\beta)/h} u(\beta) dm_\Lambda(\beta). \end{aligned}$$

Using (5.7) and the above calculation with $A = Z_j^\Lambda$ gives

$$(5.8) \quad \tilde{Z}_j^\Lambda (K_\Lambda(\alpha, \bullet)) \equiv 0, \quad \tilde{Z}_j^\Lambda := J \circ Z_j^\Lambda \circ J. \quad Ju := \bar{u},$$

The principal symbols are given by

$$(5.9) \quad \tilde{z}_j^\Lambda(\beta, \beta^*) = \bar{z}_j^\Lambda(\beta, -\beta^*), \quad \bar{z}_j^\Lambda := \overline{z_j^\Lambda(\bar{\beta}, \bar{\beta}^*)},$$

and by almost analytic continuation are defined in Γ .

Remarks 5.2.

1. Here we recall that the complex conjugation of β and β^* is defined as in (3.7).
2. Lemma 5.4 will discuss some properties of z_j^Λ and \bar{z}_j^Λ after a linear rescaling. Here we point out that z_j^Λ is a restriction to Λ of a holomorphic function in Γ but $\bar{z}_j^\Lambda(\alpha, \alpha^*) = \overline{\zeta_j^\Lambda(\alpha, \alpha^*)}$, $(\alpha, \alpha^*) \in T^*\Lambda$, is *not*.

5.2 A general construction

Here we establish the following:

PROPOSITION 5.3. *Let Z_j^Λ and \tilde{Z}_j^Λ be given by (5.5) and (5.8), respectively. Suppose that $b = b(\alpha, h)$ satisfies (5.3) (with no dependence on β).*

Then there exist $\psi(\alpha, \beta)$ and $a(\alpha, \beta, h)$ satisfying (5.2) and (5.3) and such that

$$(5.10) \quad \begin{aligned} \psi(\alpha, \alpha) &= -2iH(\alpha), \quad a_j(\alpha, \alpha) = b_j(\alpha), \\ e^{-\frac{i}{h}\psi(\alpha, \beta)} Z_j^\Lambda(\alpha, hD_\alpha, h) (e^{\frac{i}{h}\psi(\alpha, \beta)} a(\alpha, \beta, h)) &= \mathcal{O}_\infty, \\ e^{-\frac{i}{h}\psi(\alpha, \beta)} \tilde{Z}_j^\Lambda(\beta, hD_\beta, h) (e^{\frac{i}{h}\psi(\alpha, \beta)} a(\alpha, \beta, h)) &= \mathcal{O}_\infty, \end{aligned}$$

where

$$\mathcal{O}_\infty := \mathcal{O}(d(\alpha_x, \beta_x)^\infty + (\langle \alpha_\xi \rangle^{-1} |\alpha_\xi - \beta_\xi|)^\infty + (\langle \alpha_\xi \rangle^{-1} h)^\infty).$$

The phase $\psi(\alpha, \beta)$ and amplitudes $a_j(\alpha, \beta)$ are uniquely determined by $b_j(\alpha)$ up to \mathcal{O}_∞ and

$$(5.11) \quad -H(\alpha) - \text{Im } \psi(\alpha, \beta) - H(\beta) \leq -(d(\alpha_x, \beta_x)^2 + \langle \alpha_\xi \rangle^{-1} |\alpha_\xi - \beta_\xi|^2) / C$$

for some $C > 0$.

We will see that a and ψ are essentially determined by their values on the diagonal in $\Lambda \times \Lambda$. Therefore, the construction of ψ and a can be done locally, and we now work near $\alpha^0 = (x^0, \xi^0) \in \Lambda$, where we identify Λ with $T^*\mathbb{T}^n$ as in (3.6).

For α, β in a conic neighborhood of α^0 , we rescale Z_j using the following change of variables:

$$(5.12) \quad \begin{aligned} \tilde{\alpha}_x &:= \alpha_x - \alpha_x^0, & \tilde{\alpha}_\xi &:= \langle \alpha_\xi^0 \rangle^{-1} (\alpha_\xi - \alpha_\xi^0), \\ \tilde{\beta}_x &:= \beta_x - \alpha_x^0, & \tilde{\beta}_\xi &:= \langle \alpha_\xi^0 \rangle^{-1} (\beta_\xi - \alpha_\xi^0). \end{aligned}$$

In these new coordinates, the operators Z_j^Λ become

$$\begin{aligned}
 Z_j^\Lambda &= \zeta_j^\Lambda(\tilde{\alpha}, \tilde{h}D_{\tilde{\alpha}}) + \tilde{h}\zeta_j^1(\tilde{\alpha}, \tilde{h}D_{\tilde{\alpha}}) + \tilde{h}^2\zeta_j^2(\tilde{\alpha}), \quad \zeta_j^\Lambda = z_j|_\Lambda, \\
 z_j(z, \zeta, z^*, \zeta^*) &:= \lambda(\zeta)(z_j^* - \zeta_j - \theta_j) \\
 &\quad + \frac{1}{2}\lambda(\zeta)^3(z^* - \zeta - \theta)^2(\zeta_j + \theta_j) - i\zeta_j^*, \\
 \tilde{h} &:= \frac{h}{\langle \alpha_\xi^0 \rangle}, \quad \lambda(\zeta) := \frac{\langle \alpha_\xi^0 \rangle}{\langle \langle \alpha_\xi^0 \rangle (\zeta + \theta) \rangle}, \quad \theta := \frac{\alpha_\xi^0}{\langle \alpha_\xi^0 \rangle},
 \end{aligned}
 \tag{5.13}$$

where we still have $\{\zeta_j^\Lambda, \zeta_k^\Lambda\} = 0$. The operators \tilde{Z}_j^Λ are defined using (5.8).

We now define the rescaled phase and amplitudes:

$$\begin{aligned}
 \tilde{\psi}(\tilde{\alpha}, \tilde{\beta}) &:= \langle \alpha_\xi^0 \rangle^{-1} \psi(\alpha, \beta), \quad \tilde{H}(\tilde{\alpha}) := \langle \alpha_\xi^0 \rangle^{-1} H(\alpha), \\
 \tilde{G}(\tilde{\alpha}) &= \langle \alpha_\xi^0 \rangle^{-1} G(\alpha), \\
 \tilde{a}_j(\tilde{\alpha}, \tilde{\beta}) &:= \langle \alpha_\xi^0 \rangle^j a_j(\alpha, \beta), \quad \tilde{b}_j(\tilde{\alpha}) := \langle \alpha_\xi^0 \rangle^j b_j(\alpha),
 \end{aligned}
 \tag{5.14}$$

so that

$$a(\alpha, \beta) \sim \sum_{j=0}^\infty \tilde{h}^j \tilde{a}_j(\tilde{\alpha}, \tilde{\beta}), \quad b(\alpha) \sim \sum_{j=0}^\infty \tilde{h}^j \tilde{b}_j(\tilde{\alpha}).$$

Hence (5.10) becomes

$$\begin{aligned}
 \tilde{\psi}(\tilde{\alpha}, \tilde{\alpha}) &= -2i\tilde{H}(\tilde{\alpha}), \quad \tilde{a}_j(\tilde{\alpha}, \tilde{\alpha}) = \tilde{b}_j(\tilde{\alpha}), \\
 e^{-\frac{i}{\tilde{h}}\tilde{\psi}(\tilde{\alpha}, \tilde{\beta})} Z_j^\Lambda(e^{\frac{i}{\tilde{h}}\tilde{\psi}(\bullet, \tilde{\beta})} \tilde{a}(\bullet, \tilde{\beta}, \tilde{h}))(\tilde{\alpha}) &= \mathcal{O}(|\tilde{\alpha} - \tilde{\beta}|^\infty + \tilde{h}^\infty), \\
 e^{-\frac{i}{\tilde{h}}\tilde{\psi}(\tilde{\alpha}, \tilde{\beta})} \tilde{Z}_j^\Lambda(e^{\frac{i}{\tilde{h}}\tilde{\psi}(\tilde{\alpha}, \bullet)} \tilde{a}(\tilde{\alpha}, \bullet, \tilde{h}))(\tilde{\beta}) &= \mathcal{O}(|\tilde{\alpha} - \tilde{\beta}|^\infty + \tilde{h}^\infty),
 \end{aligned}
 \tag{5.15}$$

where now $\tilde{\psi}$ and \tilde{a}_j are smooth functions in a neighbourhood of $0 \in \mathbb{R}^{2n} \times \mathbb{R}^{2n}$.

(5.16) To simplify notation we now drop the \sim in \tilde{h} , $\tilde{\psi}$, \tilde{H} , \tilde{G} , and \tilde{a} .

This will apply until the end of the construction of the phase and the amplitude.

Eikonal equations

Here we work in the coordinates (5.12) and use the convention (5.16). Hence we assume that Λ is a neighbourhood of 0 in $T^*\mathbb{R}^n$.

Let ζ_j^Λ and $\tilde{\zeta}_j^\Lambda$ be the principal symbols of Z_j^Λ and \tilde{Z}_j^Λ , respectively—see (5.13). The eikonal equations we want to solve are

$$\begin{aligned}
 \zeta_j^\Lambda(\alpha, d_\alpha \psi(\alpha, \beta)) &= \mathcal{O}(|\alpha - \beta|^\infty), \\
 \tilde{\zeta}_j^\Lambda(\beta, d_\beta \psi(\alpha, \beta)) &= \mathcal{O}(|\alpha - \beta|^\infty),
 \end{aligned}
 \tag{5.17}$$

for $\alpha, \beta \in \Lambda$. We also put

$$\bar{\zeta}_j^\Lambda(\alpha, \alpha^*) := \tilde{\zeta}_j^\Lambda(\alpha, -\alpha^*);$$

see (5.8) and (5.9). We note that for $(\alpha, \alpha^*) \in T^*\Lambda$, $\bar{\zeta}_j^\Lambda(\alpha, \alpha^*) = \overline{\zeta_j^\Lambda(\alpha, \alpha^*)}$. The next lemma records the Poisson bracket properties of ζ_j^Λ on Λ .

LEMMA 5.4. *Let $\{\bullet, \bullet\}$ denote the Poisson bracket on $T^*\mathbb{R}^n$ defined using the (real) symplectic form $\sigma_\Lambda := (\sigma_{T^*\mathbb{C}^n})|_\Lambda$ and coordinates (3.6). Let*

$$\Sigma := \{\zeta_j^\Lambda(\rho) = 0 : \rho \in T^*\mathbb{R}^{2n}, \quad |x^* - \xi - \theta| < \lambda(\xi)^{-1}\}, \quad \rho = (x, \xi, x^*, \xi^*),$$

where λ is defined in (5.13).

Then, for ζ_j^Λ defined above we have $\{\zeta_j^\Lambda, \zeta_k^\Lambda\} = 0$, and for $\|G\|_{C^2} \ll 1$,

$$(5.18) \quad \left(\frac{1}{2i}\{\zeta_j^\Lambda, \bar{\zeta}_k^\Lambda\}(\alpha, \alpha^*)\right)_{1 \leq j, k \leq n} \gg cI, \quad c > 0,$$

for $(\alpha, \alpha^*) \in \Sigma \cap \text{nbhd}_{T^*\mathbb{R}^{2n}}(0)$.

The positivity condition in Lemma 5.4 will be used in two places. First, it is used to guarantee that the Lagrangian used to construct the phase solving (5.17) is strictly positive (see (A.16)). Next, when G is only smooth, this condition will be crucial when proving (5.31) (see also [15, (6.29)]) and hence that the Lagrangian we construct is almost analytic. The proof of the lemma will also show that there are solutions to $\zeta_j^\Lambda(\rho) = 0$ with $|x^* - \xi - \theta| \geq \lambda(\xi)$ ($\lambda(\xi) \sim 1$ for ξ in a neighborhood of 0). However, (5.18) may not be satisfied at these points and hence the Lagrangian will *not* be appropriately positive.

PROOF. It is enough to check (5.18) for $G = 0$. In that case Σ is contained in $\{(\alpha, d_\alpha\varphi(\alpha, y) : \alpha \in \mathbb{R}^{2n}, y \in \mathbb{C}^n\}$ where $\varphi(\alpha, y)$ is the rescaled phase of our FBI transform (this follows from the fact that ζ_j^Λ are principal symbols of operators annihilating T). Hence

$$(5.19) \quad \begin{aligned} \Sigma &:= \{(x, \xi, \varphi_x, \varphi_\xi) : y \in \mathbb{C}^n, |\xi + \theta - \varphi_x| < 1, x, \xi \in \mathbb{R}^n\} \\ &\quad \cap T^*\mathbb{R}^{2n} \\ &= \{(x, \xi, \xi + \theta, 0)\}, \\ \varphi &= \varphi(x, \xi, y) := \langle x - y, \xi + \theta \rangle + \frac{i}{2}\lambda(\xi)^{-1}(x - y)^2, \end{aligned}$$

where λ was defined in (5.13). To check this, observe that if $x^* = \varphi_x(x, \xi, y)$, $\xi^* = \varphi_\xi(x, \xi, y)$, then $y = x - i\lambda(\xi + \theta - x^*)$ and

$$\xi^* = i\lambda(\xi + \theta - x^*) + \frac{1}{2}i\partial_\xi\lambda(\xi + \theta - x^*)^2.$$

As x^* and ξ^* are real, we obtain that either $x^* = \xi + \theta$ as claimed or

$$\lambda|\xi + \theta - x^*| = 2\lambda^2|\partial_\xi\lambda|^{-1} = 2\lambda^{-1}|\xi + \theta|^{-1} = 2\frac{\langle \alpha_\xi^0 \rangle(\xi + \theta)}{\langle \alpha_\xi^0 \rangle|\xi + \theta|} \geq 2.$$

which contradicts the condition in (5.19). Hence $\xi = x^* - \theta$ and $y = x$. Here we have used the definition of λ , (5.13)

Since $\{\zeta_j^\Lambda, \zeta_k^\Lambda\} = 0$ we see that

$$\begin{aligned} \frac{1}{2i} \{\zeta_j^\Lambda, \bar{\zeta}_k^\Lambda\} &= \{\text{Im } \zeta_j^\Lambda, \text{Re } \zeta_k^\Lambda\} \\ &= -\partial_{\xi_j} (\lambda(\xi)(x_k^* - \xi_k - \theta_k) + \frac{1}{2}\lambda(\xi)^3(x^* - \xi - \theta)^2(\xi_k + \theta_k)) \\ &= \lambda(\xi)\delta_{jk}, \end{aligned}$$

when evaluated at $x^* = \xi + \theta$. Hence, for $G = 0$, the matrix in (5.18) is given by $\lambda(\xi)I$ and $\lambda(\xi) \sim 1$ for ξ bounded. Hence for G small the matrix stays positive definite. \square

>From the geometric point of view, the framework for construction of the phase is the same as in [15, Section 2.2] (see also [15, Section 6.1] for a presentation in a simpler case). It is convenient to remove the weight by putting

$$\psi_H(\alpha, \beta) := iH(\alpha) + \psi(\alpha, \beta) + iH(\beta).$$

We also define,

$$(5.20) \quad \begin{aligned} \zeta_j^H(\alpha, \alpha^*) &:= \zeta_j^\Lambda(\alpha, \alpha^* - i dH(\alpha)), \\ \bar{\zeta}_j^H(\alpha, \alpha^*) &:= \bar{\zeta}_j^\Lambda(\alpha, \alpha^* + i dH(\alpha)) = \overline{\zeta_j^H(\bar{\alpha}, \bar{\alpha}^*)}. \end{aligned}$$

(Here again the $\bar{\alpha}$ and $\bar{\alpha}^*$ are defined after an identification of Λ with $T^*\mathbb{R}^n$.) Lemma 5.4 remains valid for ζ_j^H .

The eikonal equations (5.17) become

$$(5.21) \quad \begin{aligned} \zeta_j^H(\alpha, d_\alpha \psi_H(\alpha, \beta)) &= \mathcal{O}(|\alpha - \beta|^\infty), \\ \bar{\zeta}_j^H(\beta, -d_\beta \psi_H(\alpha, \beta)) &= \mathcal{O}(|\alpha - \beta|^\infty), \end{aligned}$$

for $\alpha, \beta \in \Lambda$. Since we demand that $\psi(\alpha, \alpha) = -2iH(\alpha)$, it follows that ψ_H vanishes on the diagonal, and by differentiation,

$$(5.22) \quad 0 = d_\alpha(\psi_H(\alpha, \alpha)) = d_\alpha \psi_H(\alpha, \beta)|_{\beta=\alpha} + d_\beta \psi_H(\alpha, \beta)|_{\beta=\alpha}, \quad \alpha \in \Lambda.$$

To construct ψ_H we will construct \mathcal{C}_H , a Lagrangian relation for which ψ_H will be the generating function:

$$(5.23) \quad \begin{aligned} \mathcal{C}_H &= \{(\alpha, d_\alpha \psi_H(\alpha, \beta), \beta, -d_\beta \psi_H(\alpha, \beta)) : \\ &(\alpha, \beta) \in \text{nbhd}_{\mathbb{C}^{4n}}(\text{Diag}(\Lambda \times \Lambda))\}. \end{aligned}$$

We first assume that G , and hence H , are real analytic and have holomorphic extensions.

Writing $\rho = (x, \xi, x^*, \xi^*)$, the eikonal equations require that we should have (up to equivalence of almost analytic manifolds and exactly on $T^*\Lambda$)

$$(5.24) \quad \begin{aligned} \mathcal{C}_H \subset S \times \bar{S}, \quad S &:= \{\rho : \zeta_j^H(\rho) = 0, \rho \in \text{nbhd}_{\mathbb{C}^{4n}}(\mathbb{R}^{4n}), |x^* - \xi - \theta| < 1\}, \\ \bar{S} &:= \{\bar{\rho} : \bar{\rho} \in S\} = \{\rho : \bar{\zeta}_j^H(\rho) = 0, \rho \in \text{nbhd}_{\mathbb{C}^{4n}}(\mathbb{R}^{4n}), |x^* - \xi - \theta| < 1\}. \end{aligned}$$

The condition (5.22) means that

$$(5.25) \quad \mathcal{C}_H \cap \pi^{-1}(\Delta_{\mathbb{C}^{2n} \times \mathbb{C}^{2n}}) = \Delta((S \cap \bar{S}) \times (S \cap \bar{S})),$$

where $\Delta(A \times A) := \{(a, a) : a \in A\}$ and

$$\pi : \mathbb{C}^{2n} \times \mathbb{C}^{2n} \times \mathbb{C}^{2n} \times \mathbb{C}^{2n} \rightarrow \mathbb{C}^{2n} \times \mathbb{C}^{2n}$$

is the natural projection; i.e., $\pi(\alpha, \alpha^*, \beta, \beta^*) = (\alpha, \beta)$. In fact, (5.22) shows that this must be true for $\mathcal{C}_H \cap \pi^{-1}(\Delta_{\mathbb{R}^{2n} \times \mathbb{R}^{2n}})$, and then it follows by analytic continuation (or an equivalence of almost analytic manifolds once we move to the C^∞ category). We have the following additional property, which comes from the choice of the weight H :

LEMMA 5.5. *Let S and \bar{S} be defined in (5.24). Then for H satisfying (3.3), we have*

$$(5.26) \quad (S \cap \bar{S})_{\mathbb{R}} = S_{\mathbb{R}} = \{(\alpha, \operatorname{Re}(zd\zeta|_{\Lambda}) : \alpha \in \operatorname{nbhd}_{\mathbb{R}^{2n}}(0)\}.$$

PROOF. As in the proof of Lemma 5.4, it is useful to go to the origins of the symbols ζ_j^H (5.20): Z_j^Λ 's, with symbols ζ_j^Λ annihilate the phase in T_Λ and that shows that, after switching to ζ_j^H ,

$$\begin{aligned} S_\alpha &:= S \cap T_\alpha^* \Lambda^{\mathbb{C}} = \{(\alpha, d_\alpha \varphi(\alpha, y) + i dH(\alpha)) : y \in \mathbb{C}^n\}, \\ \varphi(\alpha, y) &:= \langle z - y, \zeta + \theta \rangle + \frac{i}{2} \lambda^{-1}(\zeta)(z - y)^2, \\ z &= \alpha_x + i G_\xi(\alpha_x, \alpha_\xi), \quad \zeta = \alpha_\xi - i G_x(\alpha_x, \alpha_\xi). \end{aligned}$$

where $\lambda(\zeta)$ and θ were defined in (5.13).

In the case $G = 0$ (and hence $H = 0$), S_α and $\bar{S}_\alpha := \bar{S} \cap T_\alpha^* \Lambda^{\mathbb{C}}$ intersect transversally in one point. This remains true for a small perturbation induced by G with $\|G\|_{C^2} \ll 1$ (this corresponds to symbolic norms before rescaling). Hence we are looking for a solution to

$$(5.27) \quad d_\alpha \varphi(\alpha, y) + i dH(\alpha) = \overline{d_\alpha \varphi(\alpha, y')} - i dH(\alpha).$$

Now, at $y = y' = \alpha_x$ we have $d_\alpha \varphi(\alpha, y) = \zeta dz|_{\Lambda}$, and in view of the definition of dH in (3.3), (5.27) holds. It follows that for $\alpha \in \Lambda$, that is, for α real,

$$S_\alpha \cap \bar{S}_\alpha = \{(\alpha, \operatorname{Re}(zd\zeta|_{\Lambda}))\} = S_\alpha \cap T^* \Lambda, \quad \Lambda \simeq \operatorname{nbhd}_{T^* \mathbb{R}^n}(0).$$

But this proves (5.26). □

Since

$$\mathcal{C}_H \subset \bigcap_{j=1}^n (\pi_L^* \zeta_j^H)^{-1}(0) \cap (\pi_R^* \bar{\zeta}_j^H)^{-1}(0), \quad \pi_L(\rho, \rho') := \rho, \quad \pi_R(\rho, \rho') := \rho',$$

it follows that the complex vector fields $H_{\pi_L^* \zeta_j^H}$ and $H_{\pi_R^* \bar{\zeta}_j^H}$ are tangent to \mathcal{C}_H . By checking the case of $T^* \Lambda = T^* \mathbb{R}^n$ (no deformation and hence $H \equiv 0$) we have (see [15, Section 2.2]) that $S \cap \bar{S}$ is a symplectic submanifold (with respect to the complex symplectic form) of complex dimension $2n$. The independence of

$H_{\zeta_k^H}, H_{\bar{\zeta}_j^H}$, $j, k = 1, \dots, n$ (again easily seen in the unperturbed case), shows that

$$(5.28) \quad B_{\mathbb{C}^n}(0, \epsilon) \times B_{\mathbb{C}^n}(0, \epsilon) \times (S \cap \bar{S}) \ni (t, s, \rho) \\ \mapsto (\exp\langle t, H_{\zeta^H} \rangle(\rho), \exp\langle s, H_{\bar{\zeta}^H} \rangle(\rho)) \in \mathbb{C}^{8n},$$

is a biholomorphic map to an embedded (complex) $4n$ -dimensional submanifold. This implies that

$$(5.29) \quad \mathcal{C}_H = \{(\exp\langle t, H_{\zeta^H} \rangle(\rho), \exp\langle s, H_{\bar{\zeta}^H} \rangle(\rho)) : \rho \in S \cap \bar{S}, t, s \in B_{\mathbb{C}^n}(0, \epsilon)\},$$

where $\langle t, H_{\bullet^H} \rangle := \sum_{k=1}^n t_k H_{\bullet_k^H}$, $\bullet = \zeta, \bar{\zeta}$. Checking again in the unperturbed case, we have that for $\rho \in S \cap \bar{S}$

$$(5.30) \quad \pi_* : T_\rho \mathcal{C}_H \rightarrow T_{\pi(\rho)} \mathbb{C}^{4n} \quad \text{is onto.}$$

We now explain how to use almost analytic extensions off Λ in the C^∞ case. We first identify Λ with $T^*\mathbb{R}^n$ using (3.6) and extending G almost analytically to \mathbb{C}^{4n} . The symplectic form is now the almost analytic extension of the symplectic form $d\zeta \wedge dz|_\Lambda$. Hence we define (see Appendix A for the definitions)

$$\mathcal{C}_H = \{(\exp \widehat{\langle t, H_{\zeta^H} \rangle}(\rho), \exp \widehat{\langle s, H_{\bar{\zeta}^H} \rangle}(\rho)) : \rho \in S \cap \bar{S}, t, s \in B_{\mathbb{C}^n}(0, \epsilon)\}.$$

We claim that

$$(5.31) \quad |\text{Im} \exp \widehat{\langle t, H_{\zeta^H} \rangle}(\rho)| \geq |t|/C, \quad |\text{Im} \exp \widehat{\langle s, H_{\bar{\zeta}^H} \rangle}(\rho)| \geq |s|/C, \quad \rho \in S \cap \bar{S}.$$

In fact, in view of Lemma 5.4 at $\rho \in T^*\Lambda \cap S$ and for $\|G\|_{C^2}$ small, we can assume $\{\zeta_j^\Lambda, \bar{\zeta}_k^\Lambda\}(\rho)/2i$ is positive definite. The changes of variable leading to ζ_j^H is a symplectomorphism and hence we have the same property for ζ_j^H . By changing ζ_j^H by a linear transformation, we can then assume that $\{\zeta_j^H, \bar{\zeta}_k^H\}(\rho)/2i = \delta_{kj}$. Hence we can make a linear symplectic change of variables at any point of $T^*\Lambda$ giving new variables (x, y, ξ, η) , $x, y, \xi, \eta \in \mathbb{R}^n$, centered at $0 \in \mathbb{R}^{4n}$, such that

$$\zeta_j^H = c(\eta_j + iy_j) + \mathcal{O}(|x|^2 + |y|^2 + |\xi|^2 + |\eta|^2), \quad c > 0.$$

This continues to hold for the almost analytic continuations of ζ_j^H . That means that near 0,

$$(5.32) \quad S \cap \bar{S} = \{(z, 0, \zeta, 0) + F(z, \zeta) : (z, \zeta) \in \text{nbhd}_{\mathbb{C}^{2n}}(0)\}, \quad F = \mathcal{O}(|z|^2 + |\zeta|^2),$$

We also note that for $(z, \zeta) \in \mathbb{R}^{2n}$ (which corresponds to the intersection with $T^*\Lambda$), $S \cap \bar{S}$ is real. This means that in (5.32),

$$\text{Im} F(z, \zeta) = \mathcal{O}((|\text{Im} z| + |\text{Im} \zeta|)(|z| + |\zeta|)).$$

Hence,

$$\begin{aligned} & |\operatorname{Im} \exp \widehat{\langle t, H_{\zeta^H} \rangle}((z, 0, \zeta, 0) + F(z, \zeta))| \\ &= |(\operatorname{Im} z, c \operatorname{Im} t, \operatorname{Im} \zeta, c \operatorname{Re} t)| + \mathcal{O}((|\operatorname{Im} z| + |\operatorname{Im} \zeta| + |t|)(|z| + |\zeta|) + |t|^2) \\ &\geq |t|/C \quad \text{if } |z|, |\zeta| \ll 1, \end{aligned}$$

with the corresponding estimate for $\bar{\zeta}^H$. Lemma A.2 and (5.31) now show the almost analyticity of \mathcal{C}_H , and Lemma A.3 shows that \mathcal{C}_H is Lagrangian in the almost analytic sense:

$$(\pi_L^* \omega_{T^* \mathbb{C}^{2n}} - \pi_R^* \omega_{T^* \mathbb{C}^{2n}})|_{\mathcal{C}_H} \sim 0.$$

(See the appendix for the review of the almost analytic machinery and notation.) Lemma 5.5 shows that $\Delta((S \cap \bar{S}_{\mathbb{R}} \times (S \cap \bar{S})_{\mathbb{R}}) = (\mathcal{C}_H)_{\mathbb{R}}$ is a submanifold, Lemma 5.4 shows that \mathcal{C}_H is therefore a strictly positive almost analytic Lagrangian submanifold, and hence, using (5.30), Lemma A.5 now gives $\psi_H = \psi_H(\alpha, \beta)$ such that

$$d_{\bar{\alpha}, \bar{\beta}} \psi_H(\alpha, \beta) = \mathcal{O}(|\operatorname{Im} \alpha|^\infty + |\operatorname{Im} \beta|^\infty + |\operatorname{Im} \psi_H(\alpha, \beta)|^\infty),$$

and (5.23) holds in the sense of equivalence of almost analytic manifolds (that is, with \sim of (A.2) replacing the equality). In addition, in view of (5.26) and (5.31),

$$(5.33) \quad \begin{aligned} d_\alpha \psi_H(\alpha, \beta)|_{\beta=\alpha} &= \operatorname{Re}(\zeta \cdot dz|_\Lambda), \\ d_\beta \psi_H(\alpha, \beta)|_{\beta=\alpha} &= -\operatorname{Re}(\zeta \cdot dz|_\Lambda), \end{aligned} \quad \alpha \in \operatorname{nbhd}_{\mathbb{R}^{2n}}(0),$$

and $d_\alpha(\psi_H(\alpha, \alpha)) \sim 0$. Hence we can choose $\psi_H(\alpha, \alpha) = 0$. We also see that

$$d_\alpha \operatorname{Im} \psi_H(\alpha, \beta)|_{\beta=\alpha} = 0, \quad d_\beta \operatorname{Im} \psi_H(\alpha, \beta)|_{\beta=\alpha} = 0, \quad \alpha \in \operatorname{nbhd}_{\mathbb{R}^{2n}}(0),$$

which means that $\operatorname{Im} \psi_H(\alpha, \beta) = \mathcal{O}(|\alpha - \beta|^2)$, $\alpha, \beta \in \operatorname{nbhd}_{\mathbb{R}^{2n}}(0)$, and the comparison with the case of $G = 0$ shows that

$$(5.34) \quad \operatorname{Im} \psi_H(\alpha, \beta) \sim |\alpha - \beta|^2.$$

Finally, we return to (5.21) (recall that ζ_j^H are the almost analytic extensions of ζ_j^H from $T^* \Lambda$ and that $\{\zeta_j^H, \zeta_k^H\} \sim 0$):

$$\begin{aligned} \widehat{\langle s, H_{\pi_L^* \zeta^H} \rangle} \pi_L^* \zeta_j^H &= \sum_{k=1}^n (H_{s_k \zeta_k^H} \zeta_j^H + \overline{H_{s_k \zeta_k^H}} \zeta_j^H) \\ &= \mathcal{O}(|\operatorname{Im} Z|^\infty), \quad Z = (\alpha, \beta, \alpha^*, \beta^*), \end{aligned}$$

with similar estimates for $\pi_R^* \bar{\zeta}_j^H$'s. Hence using the definition (A.2). This implies that

$$\pi_L^* \zeta_j^H, \pi_R^* \bar{\zeta}_j^H \sim 0 \quad \text{on } \mathcal{C}_H.$$

In view of the discussion above (\mathcal{C}_H equivalent to the right-hand side of (5.23)) we obtain

$$\zeta_j^H(\alpha, d_\alpha \psi_H(\alpha, \beta)) = \mathcal{O}(|\operatorname{Im} \alpha|^\infty + |\operatorname{Im} \beta|^\infty + |\operatorname{Im} d_\alpha \psi_H|^\infty + |\operatorname{Im} d_\beta \psi_H|^\infty),$$

with the same estimate for $\overline{\zeta_j^H}(\beta, -d_\beta \psi_H(\alpha, \beta))$. This and (5.34) give (5.17).

This completes the construction of the phase needed in Proposition 5.3. The construction of \mathcal{C}_H satisfying (5.26), (5.25), and (5.24) is equivalent, in the almost analytic sense, to the construction of ψ_H satisfying (5.34) and (5.22) that gives uniqueness of ψ_H .

We have achieved more, as the definition of \mathcal{C}_H shows that, in the analytic case (5.29), $\mathcal{C}_H \circ \mathcal{C}_H = \mathcal{C}_H$ (see [15, lemma 2] for a simple linear algebra case). In general, we have $\mathcal{C}_H \circ \mathcal{C}_H \sim \mathcal{C}_H$, which for real values of α and β means that

$$\text{c.v.}\gamma(\psi_H(\alpha, \gamma) + \psi_H(\gamma, \beta)) = \psi_H(\alpha, \beta) + \mathcal{O}(|\alpha - \beta|^\infty).$$

We now return to our original ψ in (5.1), $\psi(\alpha, \beta) = -iH(\alpha) + \psi_H(\alpha, \beta) - iH(\beta)$. Our construction shows that

$$(5.35) \quad (5.17) \text{ holds, } \psi(\alpha, \alpha) = -2iH(\alpha), \quad \psi(\alpha, \beta) = -\overline{\psi(\beta, \alpha)}, \quad \alpha, \beta \in \Lambda.$$

The value of $d_\alpha \psi$ on the diagonal $\zeta \cdot dz|_\Lambda$ is determined by (3.3) and (5.33). In addition, ψ is uniquely determined, up to $\mathcal{O}(|\alpha - \beta|^\infty)$, by (5.35).

Returning to the original problem of solving (5.17) we now record our findings:

PROPOSITION 5.6. *With the convention of (5.16), suppose that H is given by (3.3) and $\zeta_j^\Lambda, \tilde{\zeta}_j^\Lambda$ are defined in (5.13). Then there exists $\psi \in C^\infty(\Lambda \times \Lambda)$, $\Lambda = \text{nbhd}_{\mathbb{R}^{2n}}(0)$, such that (5.17) holds and $\psi(\alpha, \alpha) = -2iH(\alpha)$. The function ψ is uniquely determined modulo $\mathcal{O}(|\alpha - \beta|^\infty)$. Moreover, we have*

$$(5.36) \quad \begin{aligned} \text{c.v.}\gamma(\psi(\alpha, \gamma) + 2iH(\gamma) + \psi(\gamma, \beta)) &= \psi(\alpha, \beta) + \mathcal{O}(|\alpha - \beta|^\infty), \\ -H(\alpha) - \text{Im } \psi(\alpha, \beta) - H(\beta) &\leq -|\alpha - \beta|^2/C, \quad C > 0, \\ (d_\alpha \psi)(\alpha, \alpha) &= \zeta \cdot dz|_\Lambda. \end{aligned}$$

Transport equations

Keeping the convention (5.16), we now solve the transport equations arising from (5.15). We start with a formal discussion (valid when all the objects are analytic). We first note that in view of (5.17) and (5.36) for any $b(\alpha, \beta)$ analytic in a neighbourhood of 0 (in the notation of (5.13) and (5.16)),

$$(5.37) \quad \begin{aligned} Z_j^\Lambda(\alpha, hD_\alpha)(e^{\frac{i}{h}\psi(\alpha, \beta)} b(\alpha, \beta)) &= h e^{\frac{i}{h}\psi(\alpha, \beta)} ((V_j + c_j)b(\alpha, \beta) + \mathcal{O}(h)), \\ \tilde{Z}_j^\Lambda(\beta, hD_\beta)(e^{\frac{i}{h}\psi(\alpha, \beta)} b(\alpha, \beta)) &= h e^{\frac{i}{h}\psi(\alpha, \beta)} ((\tilde{V}_j + \tilde{c}_j)b(\alpha, \beta) + \mathcal{O}(h)). \end{aligned}$$

Here

$$\begin{aligned} V_j &:= \langle V_j(\alpha, \beta), \partial_\alpha \rangle, \quad V_j(\alpha, \beta)_\ell := \partial_{\alpha_\ell} \zeta_j^\Lambda(\alpha, d_\alpha \psi(\alpha, \beta)), \\ c_j(\alpha, \beta) &:= \frac{1}{2} \sum_{\ell=1}^{2n} \partial_{\alpha_\ell} V_j(\alpha, \beta) + \zeta_j^1(\alpha, d_\alpha \psi(\alpha, \beta)) \\ &\quad - i \sum_{k, \ell=1}^{2n} \partial_{\alpha_k \alpha_\ell}^2 \psi(\alpha, \beta) \partial_{\alpha_k^* \alpha_\ell^*}^2 \zeta_j^\Lambda(\alpha, d_\alpha \psi(\alpha, \beta)), \end{aligned}$$

with similar expressions coming from the applications $\tilde{Z}_j^\Lambda(\beta, hD_\beta)$: V_j, c_j , replaced by \tilde{V}_j, \tilde{c}_j , and with the roles of α and β switched.

A key observation here is that the holomorphic vector fields $H_{\xi_j^\Lambda(\alpha)}$ and $H_{\tilde{\xi}_j^\Lambda(\beta)}$ are tangent to

$$\mathcal{C} = \{(\alpha, d_\alpha \psi(\alpha, \beta), \beta, d_\beta \psi(\alpha, \beta)) : \alpha, \beta \in \text{nbhd}_{\mathbb{C}^{2n}}(0)\},$$

and that they commute. In the parametrization of \mathcal{C} by (α, β) , they are given by V_j and $-\tilde{V}_j$, respectively. Hence,

$$(5.38) \quad [V_j, V_k] = 0, \quad [V_j, \tilde{V}_k] = 0, \quad [\tilde{V}_k, \tilde{V}_k] = 0.$$

Hence, we seek a of the form

$$a(\alpha, \beta) \sim \sum_{k=0}^{\infty} h^k a_k(\alpha, \beta),$$

where, we want to solve

$$(5.39) \quad V_j a_k(\alpha, \beta) + c_j(\alpha, \beta) a_k(\alpha, \beta) = F_{k-1}^j(a_0, \dots, a_{k-1})(\alpha, \beta), \quad F_{-1}^j \equiv 0,$$

with the corresponding expression involving \tilde{V}_j .

Solving (5.39) means that

$$(5.40) \quad \begin{aligned} Z_j^\Lambda(\alpha, hD_\alpha) \left(e^{\frac{i}{h}\psi(\alpha, \beta)} \sum_{k=0}^{K-1} h^k a_k(\alpha, \beta) \right) &= h^{K+1} e^{\frac{i}{h}\psi(\alpha, \beta)} F_{K-1}^j(\alpha, \beta), \\ \tilde{Z}_j^\Lambda(\beta, hD_\beta) \left(e^{\frac{i}{h}\psi(\alpha, \beta)} \sum_{k=0}^{K-1} h^k a_k(\alpha, \beta) \right) &= h^{K+1} e^{\frac{i}{h}\psi(\alpha, \beta)} \tilde{F}_{K-1}^j(\alpha, \beta). \end{aligned}$$

Since

$$\begin{aligned} [Z_j^\Lambda(\alpha, hD_\alpha), Z_k^\Lambda(\alpha, hD_\alpha)] &= 0, \quad [\tilde{Z}_j^\Lambda(\beta, hD_\beta), \tilde{Z}_k^\Lambda(\beta, hD_\beta)] = 0, \\ [Z_j^\Lambda(\alpha, hD_\alpha), \tilde{Z}_k^\Lambda(\beta, hD_\beta)] &= 0, \end{aligned}$$

we have from (5.38) and (5.37),

$$(5.41) \quad V_j c_\ell = V_\ell c_j, \quad V_\ell \tilde{c}_j = \tilde{V}_j c_\ell, \quad \tilde{V}_\ell \tilde{c}_j = \tilde{V}_j \tilde{c}_\ell.$$

Similarly, (5.40) gives

$$(5.42) \quad \begin{aligned} (V_j + c_j) F_{K-1}^\ell &= (V_\ell + c_\ell) F_{K-1}^j, \quad (\tilde{V}_j + \tilde{c}_j) \tilde{F}_{K-1}^\ell = (\tilde{V}_\ell + \tilde{c}_\ell) \tilde{F}_{K-1}^j, \\ (V_j + c_j) \tilde{F}_{K-1}^\ell &= (\tilde{V}_\ell + \tilde{c}_\ell) F_{K-1}^j. \end{aligned}$$

Equations (5.41) and (5.42) provide compatibility conditions for solving (5.39):

$$(V_j + c_j) a_k = F_{k-1}^j, \quad (\tilde{V}_\ell + \tilde{c}_\ell) a_k = F_{k-1}^\ell, \quad a_k(\alpha, \alpha) = b_k(\alpha),$$

where the b_k 's are prescribed. In fact, since the V_ℓ 's and \tilde{V}_j 's are independent when $\alpha = \beta$ (as complex vectorfields),

$$\begin{aligned} \mathbb{C}^{2n} \times \mathbb{C}^n \times \mathbb{C}^n &\ni (\rho, t, s) \mapsto (\alpha, \beta) \\ &= (\exp\langle V, t \rangle(\rho), \exp\langle \tilde{V}, s \rangle(\rho)) \in \mathbb{C}^{2n} \times \mathbb{C}^{2n}, \\ \langle V, t \rangle &:= \sum_{j=1}^n t_j V_j, \quad \langle \tilde{V}, s \rangle := \sum_{\ell=1}^n s_j \tilde{V}_\ell, \end{aligned}$$

is a local biholomorphic map onto of $\text{nbhd}_{\mathbb{C}^{4n}}(\text{diag}(\Lambda \times \Lambda))$ (almost analytic in the general case). In view of this and of (5.38) and (5.41), the following integrating factor, $g = g(\alpha, \beta)$, is well-defined (in the analytic case) on $\text{nbhd}_{\mathbb{C}^{4n}}(\text{diag}(\Lambda \times \Lambda))$:

$$g(e^{\langle V, t \rangle}(\rho), e^{\langle \tilde{V}, s \rangle}(\rho)) := - \sum_{j=1}^n \int_0^1 (t_j c_j + s_j \tilde{c}_j)|_{(\alpha, \beta) = (e^{\tau \langle V, t \rangle}(\rho), e^{\tau \langle \tilde{V}, s \rangle}(\rho))} d\tau,$$

and satisfies

$$V_j g(\alpha, \beta) = c_j(\alpha, \beta), \quad \tilde{V}_j g(\alpha, \beta) = \tilde{c}_j(\alpha, \beta), \quad j = 1, \dots, n.$$

We then define $a_k(\alpha, \beta)$ inductively as follows: at $(\alpha, \beta) = (e^{\langle V, t \rangle}(\rho), e^{\langle \tilde{V}, s \rangle}(\rho))$,

$$\begin{aligned} a_k(\alpha, \beta) &= e^{g(\alpha, \beta)} b_k(\rho) \\ &+ e^{g(\alpha, \beta)} \int_0^1 e^{-g(\gamma, \gamma')} (t_j F_{k-1}^j(\gamma, \gamma') \\ &+ s_j \tilde{F}_{k-1}^j(\gamma, \gamma'))|_{(\gamma, \gamma') = (e^{\tau \langle V, t \rangle}(\rho), e^{\tau \langle \tilde{V}, s \rangle}(\rho))} d\tau. \end{aligned}$$

The compatibility relations (5.42) then show that (5.39) holds.

We now modify this discussion to the C^∞ case using almost analytic extensions as in Section A.3 and that provides solutions of (5.39) for $(\alpha, \beta) \in \Lambda \times \Lambda$ valid to infinite order at $\text{diag}(\Lambda \times \Lambda)$ with any initial data on the diagonal.

Hence we have solved (5.15) locally near $(\alpha, \beta) = (0, 0)$. We now return to the original coordinates and note the uniqueness of the local construction gives us ψ and a in (5.10) satisfying (5.2) and (5.3). This completes the proof of Proposition 5.3.

5.3 Projection property

It remains to choose $a|_\Delta$ so that $B_\Lambda^2 \equiv B_\Lambda$. From (5.36) we already know that the phase in (5.1) has the correct composition property and hence we need to find the amplitude $a(\alpha, \beta)$. From Proposition 5.3 it is enough to determine a on the diagonal. For that we consider the kernel of B_Λ^2 on the diagonal:

$$(5.43) \quad K_{B_\Lambda^2}(\alpha, \alpha) = h^{-2n} \int_\Lambda e^{\frac{i}{h}(\psi(\alpha, \beta) + 2iH(\beta) + \psi(\beta, \alpha))} a(\alpha, \beta, h) a(\beta, \alpha, h) dm_\Lambda(\beta).$$

We note that the support property of a in (5.2) implies that the integration takes place over a bounded set $|\beta_\xi| \leq C \langle \alpha_\xi \rangle$. Application of complex stationary phase to (5.43) yields

$$(5.44) \quad K_{B_\Lambda^2} = h^{-n} e^{\frac{i}{h} \psi(\alpha, \beta)} c(\alpha, \beta), \quad c(\alpha, \alpha) \sim \sum_j h^j L_{2j} a(\alpha, \gamma, h) a(\gamma, \alpha, h)|_{\gamma=\alpha},$$

where L_{2j} are differential operators of order $2j$ in γ and

$$L_0|_\Delta = f(\alpha), \quad |f(\alpha)| > 0, \quad \Delta := \Delta(\Lambda \times \Lambda).$$

Since $\psi(\alpha, \beta) = -\overline{\psi(\beta, \alpha)}$, $f(\alpha) \in \mathbb{R}$. (Strictly speaking, we should again proceed with the rescaling (5.12) and we are tacitly using the convention (5.16) here.)

Writing $a \sim \sum_j h^j a_j$, we have

$$c(\alpha, \beta) \sim \sum_j h^j c_j(\alpha, \beta), \quad c_j(\alpha, \alpha) = \sum_{k+\ell+m=j} L_{2k} a_\ell(\alpha, \gamma) a_m(\gamma, \alpha)|_{\gamma=\alpha}.$$

We note that if $a(\alpha, \beta) = \overline{a(\beta, \alpha)}$, then B_Λ is self-adjoint and hence so is B_Λ^2 . That means in particular that $c(\alpha, \alpha)$ is real. Hence if $a_\ell(\alpha, \beta) = \overline{a_\ell(\beta, \alpha)}$ for $\ell \leq M$, then $c_\ell|_\Delta \in \mathbb{R}$ for $\ell \leq M$. Since

$$b_M(\alpha, \alpha) = 2f(\alpha) a_0(\alpha, \alpha) a_M(\alpha, \alpha) + \sum_{\substack{k+\ell+m=M \\ \ell, m < M}} L_{2k} a_\ell(\alpha, \gamma) a_m(\gamma, \alpha)|_{\gamma=\alpha},$$

it follows that

$$(5.45) \quad a_\ell(\alpha, \beta) = \overline{a_\ell(\beta, \alpha)}, \ell < M \implies \sum_{\substack{k+\ell+m=M \\ \ell, m < M}} L_{2k} a_\ell(\alpha, \gamma) a_m(\gamma, \alpha)|_{\gamma=\alpha} \in \mathbb{R}.$$

We iteratively solve the following sequence of equations,

$$(5.46) \quad \sum_{k+\ell+m=j} L_{2k} a_\ell(\alpha, \gamma) a_m(\gamma, \alpha)|_{\gamma=\alpha} = a_j(\alpha, \alpha),$$

with $a_j|_\Delta$ real. Proposition 5.3 then gives us the desired $a(\alpha, \beta)$.

First, let

$$a_0(\alpha, \alpha) = \frac{1}{f(\alpha)} \in C^\infty(T^*\mathbb{R}^n; \mathbb{R})$$

so that $f(\alpha) a_0(\alpha, \alpha)^2 = a_0(\alpha, \alpha)$ (i.e., (5.46) is solved for $j = 0$). The proof of Proposition 5.3 (see Section 5.2) shows that we can then find $a_0(\alpha, \beta)$ so that (5.40) holds with $K = 0$ and $a_0|_\Delta = 1/f(\alpha)$.

Assume now that (5.46) is solved for $j \leq M - 1$. Then, (5.46) with $j = M$ reads

$$\begin{aligned} a_M(\alpha, \alpha) &= \sum_{k+\ell+m=M} L_{2k} a_\ell(\alpha, \gamma) a_m(\gamma, \alpha)|_{\gamma=\alpha} \\ &= 2a_M(\alpha, \alpha) + \sum_{\substack{k+\ell+m=M \\ \ell, m < M}} L_{2k} a_\ell(\alpha, \gamma) a_m(\gamma, \alpha)|_{\gamma=\alpha}. \end{aligned}$$

Putting

$$a_M(\alpha, \alpha) = - \sum_{\substack{k+\ell+m=M \\ \ell, m < M}} L_{2k} a_\ell(\alpha, \gamma) a_m(\gamma, \alpha)|_{\gamma=\alpha},$$

we solve (5.46) for $j = M$. From (5.45) we see that $a_M(\alpha, \alpha)$ is real. The argument in Section 5.2 provides the construction of a_M from $a_\ell, \ell < M$ and $a_k|_\Delta$. Taking an almost analytic continuation with $a_M(\alpha, \beta) = \overline{a_M(\beta, \alpha)}$ then completes the construction of a_M , and hence by induction and the Borel summation lemma we have, in the notation of Proposition 5.3,

$$(5.47) \quad c = a + \mathcal{O}_\infty.$$

This gives the following:

PROPOSITION 5.7. *There exists a unique choice of $b_j(\alpha)$ in Proposition 5.3 for which the operator B_Λ defined by (5.1) satisfies*

$$(5.48) \quad B_\Lambda = B_\Lambda^{*,H}, \quad B_\Lambda = B_\Lambda^2 + \mathcal{O}(h^N)_{(\xi)^N L^2(\Lambda) \rightarrow (\xi)^{-N} L^2(\Lambda)},$$

for all N .

PROOF. In view of (5.47) we need to check that for $r = \mathcal{O}_\infty$ (in the notation of Proposition 5.3), for all N ,

$$\begin{aligned} R &= \mathcal{O}(h^N)_{(\xi)^N L^2(\Lambda) \rightarrow (\xi)^{-N} L^2(\Lambda)}, \\ Ru(\alpha) &:= h^{-n} \int_\Lambda r(\alpha, \beta, h) e^{\frac{i}{h}\psi(\alpha, \beta)} u(\beta) dm_\Lambda(\beta). \end{aligned}$$

But this is an immediate consequence of (5.11) and Schur’s criterion for boundedness on L^2 . □

5.4 Construction of the projector

We now show that the exact orthogonal projector $\Pi_\Lambda : L^2(\Lambda) \rightarrow H(\Lambda)$ satisfies

$$(5.49) \quad \Pi_\Lambda = B_\Lambda + \mathcal{O}(h^\infty)_{(\xi)^N L^2(\Lambda) \rightarrow (\xi)^{-N} L^2(\Lambda)}$$

for all N . For that we follow the proof of [33, prop. 1.1, formula (1.46)], which is related to an earlier construction in [1, step 3, proof of corollary A.4.6].

We start with the exact projector P_Λ :

$$P_\Lambda(L^2_\Lambda(\Lambda)) = H(\Lambda), \quad P_\Lambda^2 = P_\Lambda, \quad P_\Lambda = \mathcal{O}(1)_{L^2(\Lambda) \rightarrow L^2(\Lambda)},$$

given by

$$P_\Lambda = T_\Lambda S_\Lambda.$$

For a real-valued $f \in S(\Lambda)$,

$$f(\alpha) \sim \sum_{k=0}^\infty f_k(\alpha)(h/\langle\alpha_\xi\rangle)^k, \quad f_0(\alpha) > 1/C,$$

we define the following self-adjoint operator:

$$A_f := P_\Lambda f P_\Lambda^{*,H},$$

$$A_f u(\alpha) =: h^{-n} \int_\Lambda e^{\frac{i}{h}\psi_1(\alpha,\beta)} a_f(\alpha, \beta, h) u(\beta) e^{-2H(\beta)/h} dm_\Lambda(\beta),$$

where ψ_1 and a_f were obtained by using the method of complex stationary phase (again it is justified using the rescaling (5.12)).

We claim that $\psi_1 = \psi + \mathcal{O}_\infty$ (in the notation of Proposition 5.3). Indeed, since $A_f^{*,H} = A_f$ and $P_\Lambda = T_\Lambda S_\Lambda$, the arguments leading to (5.17) apply and ψ_1 satisfies the same eikonal equations. Similarly, $a_f(\alpha, \beta, h)$ satisfies transport equations implied by (5.10). Arguing as in the proof of Lemma 4.8, we find the value of $\psi_1|_\Delta$ to be

$$\psi_1(\alpha, \alpha) + 2iH(\alpha) = \text{c.v.}_\beta(\Psi(\alpha, \beta) - \overline{\Psi(\alpha, \beta)}) = 0.$$

We then invoke the uniqueness statement in Proposition 5.3.

If we can choose f so that $a_f|_\Delta = a|_\Delta + \mathcal{O}((h/\langle\alpha_\xi\rangle)^\infty)$, with a in (5.1), then the same uniqueness statement shows that $a_f = a + \mathcal{O}_\infty$. Hence

$$(5.50) \quad a_f|_\Delta = a|_\Delta + \mathcal{O}((h/\langle\alpha_\xi\rangle)^\infty) \implies$$

$$A_f = B_\Lambda + \mathcal{O}(h^\infty)_{(\xi)^N L^2(\Lambda) \rightarrow (\xi)^{-N} L^2(\Lambda)},$$

determined by its value on the diagonal. We find that, using Ψ given in (4.9) and satisfying (4.12),

$$\psi_1(\alpha, \alpha) + 2iH(\alpha) = \text{c.v.}_\beta(\Psi(\alpha, \beta) - \overline{\Psi(\alpha, \beta)} + 2iH(\alpha) - 2iH(\beta)) = 0.$$

But this means that (5.35) holds for ψ_1 and hence

$$\psi_1(\alpha, \beta) = \psi(\alpha, \beta) + \mathcal{O}(|\alpha - \beta|^\infty).$$

We next choose f so that $A_f = B_\Lambda + \mathcal{O}(h^\infty)_{(\xi)^N L^2(\Lambda) \rightarrow (\xi)^{-N} L^2(\Lambda)}$.

Writing $a_f(\alpha, \beta) \sim \sum_{k=0}^\infty (h/\langle\alpha_\xi\rangle)^k a_{f,j}(\alpha, \beta)$, we proceed as in Section 5.2: with different L_{2k} 's, $g := L_0|_\Delta \neq 0$,

$$a_{f,j}(\alpha, \alpha) = \sum_{k+l=j} L_{2k} f_l(\alpha) = g(\alpha) f_j(\alpha) + \sum_{\substack{k+l=j \\ \ell < j}} L_{2k} f_\ell(\alpha).$$

(In our special case, the amplitude in P is constant, which is not the case in generalizations—but the argument works easily just the same.) Using this, solving

$a_{f,j}(\alpha) = a_j(\alpha)$ for f is immediate. As in the construction of the amplitude of B_Λ in Section 5.2, we see that f is real valued and that f_0 is bounded from below.

We can now follow [33] and complete the proof of (5.49). We record this statement as

PROPOSITION 5.8. *Suppose that Π_Λ is an orthogonal projector from $L^2(\Lambda)$ to $H(\Lambda)$ and that B_Λ is given by Proposition 5.7. Then*

$$(5.51) \quad \Pi_\Lambda = B_\Lambda + \mathcal{O}(h^\infty)_{\langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)}$$

for all N .

PROOF. To start we observe that for $u \in H(\Lambda)$, $\|u\|_{L^2(\Lambda)} > 0$,

$$\begin{aligned} \langle A_f u, u \rangle_{L^2(\Lambda)} &= \langle P_\Lambda f P_\Lambda^* u, u \rangle_{L^2(\Lambda)} = \langle f P_\Lambda^* u, P_\Lambda^* u \rangle_{L^2(\Lambda)} \\ &\geq \min_{\alpha \in T^*\mathbb{R}^n} f(\alpha) \|P_\Lambda^* u\|_{L^2(\Lambda)}^2 \geq \frac{|\langle P_\Lambda^* u, u \rangle|^2}{C \|u\|_{L^2(\Lambda)}^2} = \|u\|_{L^2(\Lambda)}^2 / C. \end{aligned}$$

Hence

$$(5.52) \quad \begin{aligned} \|u\|_{L^2(\Lambda)} / C &\leq \|A_f u\|_{L^2(\Lambda)} \leq C \|u\|_{L^2(\Lambda)}, \quad u \in H(\Lambda), \\ A_f u &= 0, \quad u \in H(\Lambda)^\perp, \quad A_f^* = A_f, \end{aligned}$$

and

$$(5.53) \quad \Pi_\Lambda = \frac{1}{2\pi} \int_\gamma (\lambda - A_f)^{-1} d\lambda,$$

where γ is a positively oriented boundary of an open set in \mathbb{C} containing $[1/C, C]$ and excluding 0. From (5.50) and Proposition 5.7, we know that

$$(5.54) \quad A_f = A_f^2 + \mathcal{O}(h^\infty)_{\langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)},$$

and we want to use this property to show that Π_Λ is close to A_f . For that we note that if $A = A^2$, then, at first for $|\lambda| \gg 1$,

$$(\lambda - A)^{-1} = \sum_{j=0}^\infty \lambda^{-j-1} A^j = \lambda^{-1} + \lambda^{-1} \sum_{j=0}^\infty \lambda^{-j} A = \lambda^{-1} + A \lambda^{-1} (\lambda - 1)^{-1}.$$

Hence it is natural to take the right-hand side as the approximate inverse in the case when $A^2 - A$ is small:

$$(\lambda - A_f)(\lambda^{-1} + A_f \lambda^{-1} (\lambda - 1)^{-1}) = I - (A_f^2 - A_f) \lambda^{-2} (\lambda - 1)^{-1}.$$

In view of (5.54) and for h small enough, the right-hand side is invertible for $\lambda \in \gamma$ with the inverse equal to $I + R$, $R = \mathcal{O}(h^\infty)_{\langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)}$. Hence for $\lambda \in \gamma$,

$$(\lambda - A_f)^{-1} = \lambda^{-1} + \lambda^{-1} (\lambda - 1)^{-1} A_f + \mathcal{O}(h^\infty)_{\langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)}.$$

Inserting this identity into (5.53) and using Cauchy's formula gives

$$\Pi_\Lambda = A_f + \mathcal{O}(h^\infty)_{\langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)},$$

which combined with (5.50) implies (5.51). □

6 Deformation of Pseudodifferential Operators

In this section we analyse pseudodifferential operators with analytic symbols acting on spaces H_Λ^m defined in Section 4. That means describing the action on the FBI side of operators P :

$$(6.1) \quad T_\Lambda P u = (T_\Lambda P S_\Lambda)(T_\Lambda u) = (\Pi_\Lambda T_\Lambda P S_\Lambda \Pi_\Lambda)(T_\Lambda u).$$

The class of pseudodifferential operators we consider is given by

$$(6.2) \quad P u(y) := \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} e^{\frac{i}{h}\langle y-y', \eta \rangle} p(y, \eta) u(y') dy' d\eta$$

where $p \in S^m(T^*\mathbb{T}^n)$ has an analytic continuation from $T^*\mathbb{T}^n$ satisfying

$$(6.3) \quad |p(z, \zeta)| \leq M \langle \zeta \rangle^m \quad \text{for } |\text{Im } z| \leq a, |\text{Im } \zeta| \leq b \langle \text{Re } \zeta \rangle.$$

The integral in the definition (6.2) of $P u$ is considered in the sense of oscillatory integrals (see, for instance, [40, sec. 5.3]), and we extend both $y \mapsto u(y)$ and $y \mapsto p(y, \eta)$ to periodic functions on \mathbb{R}^n .

6.1 Pseudodifferential operators as Toeplitz operators

We start with a lemma that describes the middle term in (6.1):

LEMMA 6.1. *Suppose P is defined by (6.2) with p satisfying (6.3). Then, for G satisfying (3.1) with $\epsilon_0 > 0$ small enough, the Schwartz kernel of $T_\Lambda P S_\Lambda$ is given by*

$$(6.4) \quad K_P(\alpha, \beta) = c_0 h^{-n} e^{\frac{i}{h}\Psi(\alpha, \beta)} a_P(\alpha, \beta) + r(\alpha, \beta)$$

where Ψ is as in (4.9),

$$(6.5) \quad a_P \sim \sum_{j=0}^{\infty} h^j \langle \alpha_\xi \rangle^{-j} a_j, \quad a_0(\alpha, \alpha) = p|_\Lambda(\alpha),$$

$a_j \in S^0(\Lambda \times \Lambda)$ is supported in a conic neighbourhood of $\Delta(\Lambda \times \Lambda)$ and

$$(6.6) \quad |r(\alpha, \beta)| \leq e^{-((\text{Re } \alpha_\xi) + \langle \text{Re } \beta_\xi \rangle + \langle \text{Re } \alpha_x - \text{Re } \beta_x \rangle)/Ch}.$$

PROOF. We first note that for each $\beta \in \Lambda$,

$$v_\beta(y') = e^{-\frac{i}{h}\varphi^*(\beta, y')} b(\beta_x - y', \beta_\xi)$$

is a Schwartz function and hence the integral

$$h^{-\frac{3n}{4}} \frac{1}{(2\pi h)^n} \int_{\mathbb{R}^{2n}} e^{\frac{i}{h}(\langle y-y', \eta \rangle - \varphi^*(\beta, y'))} p(y, \eta) b(\beta_x - y', \beta_\xi) dy' d\eta$$

defines a Schwartz function of y . In particular, the kernel of $T_\Lambda P S_\Lambda$ is given by

$$\frac{h^{-\frac{3n}{2}}}{(2\pi h)^n} \int_{\mathbb{R}^{3n}} e^{\frac{i}{h}(\varphi(\alpha, y) + \langle y-y', \eta \rangle - \varphi^*(\beta, y'))} p(y, \eta) b(\beta_x - y', \beta_\xi) \langle \alpha_\xi \rangle^{\frac{n}{4}} dy' d\eta dy.$$

To obtain (6.4) we start by deforming the contour in η : $\eta \mapsto \eta + i \delta_1 \langle \eta \rangle \frac{y-y'}{\langle y-y' \rangle}$. The phase Φ is then given by

$$\begin{aligned} \Phi &= \langle \alpha_x - y, \alpha_\xi \rangle + \frac{i \langle \alpha_\xi \rangle}{2} (\alpha_x - y)^2 + \frac{i \langle \beta_\xi \rangle}{2} (\beta_x - y')^2 \\ &\quad + \langle y' - \beta_x, \beta_\xi \rangle + \langle y - y', \eta \rangle + i \delta_1 \langle \eta \rangle \frac{(y - y')^2}{\langle y - y' \rangle}. \end{aligned}$$

We then deform the contour in y, y' as follows:

$$y \mapsto y + i \delta_1 \frac{\eta - \alpha_\xi}{\langle \eta - \alpha_\xi \rangle}, \quad y' \mapsto y' + i \delta_1 \frac{\beta_\xi - \eta}{\langle \beta_\xi - \eta \rangle}.$$

The phase Φ becomes

$$\begin{aligned} \Phi &= \langle \alpha_x - y, \alpha_\xi \rangle + \frac{i \langle \alpha_\xi \rangle}{2} (\alpha_x - y)^2 + \frac{i \langle \beta_\xi \rangle}{2} (\beta_x - y')^2 + \langle y' - \beta_x, \beta_\xi \rangle \\ &\quad + \langle y - y', \eta \rangle + i \delta_1 \left[\frac{(\alpha_\xi - \eta)^2}{\langle \alpha_\xi - \eta \rangle} + \frac{(\beta_\xi - \eta)^2}{\langle \beta_\xi - \eta \rangle} + \langle \eta \rangle \frac{(y - y')^2}{\langle y - y' \rangle} \right] \\ &\quad + \frac{i \langle \alpha_\xi \rangle}{2} \left[-2i \delta_1 \frac{\langle \alpha_\xi - \eta, \alpha_x - y \rangle}{\langle \alpha_\xi - \eta \rangle} - \delta_1^2 \frac{(\alpha_\xi - \eta)^2}{\langle \alpha_\xi - \eta \rangle^2} \right] \\ &\quad + \frac{i \langle \beta_\xi \rangle}{2} \left[-2i \delta_1 \frac{\langle \eta - \beta_\xi, \beta_x - y' \rangle}{\langle \beta_\xi - \eta \rangle} - \delta_1^2 \frac{(\beta_\xi - \eta)^2}{\langle \beta_\xi - \eta \rangle^2} \right] \\ &\quad + \mathcal{O} \left(\delta_1^2 \langle \eta \rangle \frac{y - y'}{\langle y - y' \rangle} \left[\frac{|\langle \alpha_\xi - \eta \rangle \langle \beta_\xi - \eta \rangle + (\eta - \beta_\xi) \langle \alpha_\xi - \eta \rangle|}{|\langle \alpha_\xi - \eta \rangle \langle \beta_\xi - \eta \rangle|} \right] \right) \\ &\quad + \mathcal{O} \left(\frac{\delta_1^3 \langle \eta \rangle}{\langle y - y' \rangle} \left[\frac{|\langle \alpha_\xi - \eta \rangle \langle \beta_\xi - \eta \rangle + (\eta - \beta_\xi) \langle \alpha_\xi - \eta \rangle|^2}{|\langle \alpha_\xi - \eta \rangle \langle \beta_\xi - \eta \rangle|^2} \right] \right) \end{aligned}$$

We first consider the case when $\langle \text{Re } \alpha_\xi \rangle \geq 2 \langle \text{Re } \beta_\xi \rangle$. Then,

$$|\text{Re } \alpha_\xi - \eta| + |\text{Re } \beta_\xi - \eta| \geq c(\langle \text{Re } \alpha_\xi \rangle + \langle \text{Re } \beta_\xi \rangle + \langle \eta \rangle).$$

and in particular,

$$\text{Im } \Phi \geq c(\langle \text{Re } \alpha_\xi \rangle + \langle \text{Re } \beta_\xi \rangle + \langle \text{Re } \eta \rangle) + c(|\text{Re } \alpha_x - y| + |\text{Re } \beta_x - y'| + |y - y'|),$$

which produces a term that can be absorbed into r satisfying (6.6).

Similar arguments show that we can assume that $\langle \text{Re } \alpha_\xi \rangle$, $\langle \text{Re } \eta \rangle$, and $\langle \text{Re } \beta_\xi \rangle$ are proportional.

We now suppose that

$$\frac{|\text{Re } \alpha_\xi - \text{Re } \beta_\xi|}{\langle \text{Re } \alpha_\xi \rangle + \langle \text{Re } \beta_\xi \rangle} + |\text{Re } \alpha_x - \text{Re } \beta_x| > \delta.$$

Then, the imaginary part of the phase is bounded below by

$$\begin{aligned} \text{Im } \Phi &\geq c(\langle \text{Re } \alpha_\xi \rangle + \langle \text{Re } \beta_\xi \rangle)(1 + |\text{Re } \alpha_x - \text{Re } \beta_x|) \\ &\quad + c(|\text{Re } \alpha_x - y| + |y - y'| + |\text{Re } \beta_x - y| + |\eta - \text{Re } \alpha_\xi| + |\eta - \text{Re } \beta_\xi|). \end{aligned}$$

In particular, when

$$\frac{|\operatorname{Re} \alpha_\xi - \operatorname{Re} \beta_\xi|}{\langle \operatorname{Re} \alpha_\xi \rangle + \langle \operatorname{Re} \beta_\xi \rangle} + |\operatorname{Re} \alpha_x - \operatorname{Re} \beta_x| > \delta.$$

the integral is bounded by $Ce^{-((\operatorname{Re} \alpha_\xi) + \langle \operatorname{Re} \beta_\xi \rangle)(1 + |\operatorname{Re} \alpha_x - \operatorname{Re} \beta_x|)/h}$. Hence, we can insert a cutoff

$$\chi\left(\delta^{-1} \left[\frac{|\operatorname{Re} \alpha_\xi - \operatorname{Re} \beta_\xi|}{\langle \operatorname{Re} \alpha_\xi \rangle + \langle \operatorname{Re} \beta_\xi \rangle} + |\operatorname{Re} \alpha_x - \operatorname{Re} \beta_x| \right]\right)$$

into the integral.

With this cutoff inserted, we deform in y, y' to the critical point

$$y \mapsto y + y_c(\alpha, \beta), \quad y' \mapsto y' + y_c(\alpha, \beta),$$

where

$$y_c(\alpha, \beta) = \frac{\alpha_x \langle \alpha_\xi \rangle + \beta_x \langle \beta_\xi \rangle}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} + i \frac{\beta_\xi - \alpha_\xi}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle}.$$

This contour deformation is justified since the cutoff function guarantees that

$$\operatorname{Re} \left| \frac{\alpha_\xi - \beta_\xi}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} \right| \leq C\delta.$$

The phase is then given by

$$\begin{aligned} \Phi = & \frac{i}{2} \left(\frac{(\beta_\xi - \alpha_\xi)^2 + \langle \alpha_\xi \rangle \langle \beta_\xi \rangle (\alpha_x - \beta_x)^2}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} + \langle \alpha_\xi \rangle y^2 + \langle \beta_\xi \rangle (y')^2 \right) \\ & + \left\langle \alpha_x - \beta_x, \frac{\alpha_\xi \langle \beta_\xi \rangle + \beta_\xi \langle \alpha_\xi \rangle}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} \right\rangle + \langle y - y', \eta - \eta_c(\alpha, \beta) \rangle \end{aligned}$$

with

$$\eta_c(\alpha, \beta) = \frac{\alpha_\xi \langle \beta_\xi \rangle + \beta_\xi \langle \alpha_\xi \rangle}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} + i \frac{\langle \alpha_\xi \rangle \langle \beta_\xi \rangle (\beta_x - \alpha_x)}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle}.$$

We would now like to shift the contour to $\eta \mapsto \eta + \eta_c$. However, p only has an analytic continuation to $|\operatorname{Im} \eta| \leq b \langle \operatorname{Re} \eta \rangle$, and $\operatorname{Im} \eta_c$ is not, in general, bounded. Therefore, when $|\operatorname{Re} \eta| \ll |\operatorname{Re} \eta_c|$, we cannot make this deformation. To finish the proof, we consider two cases.

We first assume that $|\eta_c(\alpha, \beta)| \leq b/2$. Then, the contour deformation $\eta \mapsto \eta + \eta_c$ is justified, and we may perform complex stationary phase to complete the proof.

We now consider the more involved case when

$$|\eta_c| \geq \frac{b}{2} \gg \epsilon_0 > 0,$$

where ϵ_0 is as in (3.1). In that case we use the deformation

$$y \mapsto y + i\delta_1 \frac{(\eta - \eta_c)}{\langle \eta - \eta_c \rangle}, \quad y' \mapsto y' - i\delta_1 \frac{(\eta - \eta_c)}{\langle \eta - \eta_c \rangle},$$

to obtain the phase

$$\begin{aligned} & \left\langle \alpha_x - \beta_x, \frac{\alpha_\xi \langle \beta_\xi \rangle + \beta_\xi \langle \alpha_\xi \rangle}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} \right\rangle \\ & + \frac{i}{2} \left[\frac{(\beta_\xi - \alpha_\xi)^2 + \langle \alpha_\xi \rangle \langle \beta_\xi \rangle (\alpha_x - \beta_x)^2}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} + \langle \alpha_\xi \rangle y^2 + \langle \beta_\xi \rangle (y')^2 \right] \\ & + \left\langle y - y', (\eta - \eta_c) \left(1 - \frac{\delta_1}{\langle \eta - \eta_c \rangle} \right) \right\rangle + 2i\delta_1 \frac{(\eta - \eta_c)^2}{\langle \eta - \eta_c \rangle} \left(1 - \frac{\delta_1}{2\langle \eta - \eta_c \rangle} \right). \end{aligned}$$

Finally, let $\chi \in C_c^\infty((1/2, 2))$ with $\chi \equiv 1$ on $(3/4, 3/2)$, and shift contours

$$\eta \mapsto \eta + \eta_c \chi \left(\frac{|\operatorname{Re} \eta|}{|\operatorname{Re} \eta_c|} \right).$$

Note that this deformation is now justified since $|\operatorname{Re} \eta| \geq c |\operatorname{Re} \eta_c|$ on the deformation and $|\operatorname{Im} \eta_c| \leq c\epsilon_0 |\operatorname{Re} \eta_c|$. The phase is then given by

$$\begin{aligned} & \frac{i}{2} \left[\frac{(\beta_\xi - \alpha_\xi)^2 + \langle \alpha_\xi \rangle \langle \beta_\xi \rangle (\alpha_x - \beta_x)^2}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} + \langle \alpha_\xi \rangle y^2 + \langle \beta_\xi \rangle (y')^2 \right] \\ & + \left\langle \alpha_x - \beta_x, \frac{\alpha_\xi \langle \beta_\xi \rangle + \beta_\xi \langle \alpha_\xi \rangle}{\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle} \right\rangle \\ & + \left\langle y - y', (\eta - (1 - \chi)\eta_c) \left(1 - \frac{\delta_1}{\langle \eta - (1 - \chi)\eta_c \rangle} \right) \right\rangle \\ & + 2i\delta_1 \frac{(\eta - (1 - \chi)\eta_c)^2}{\langle \eta - (1 - \chi)\eta_c \rangle} \left(1 - \frac{\delta_1}{2\langle \eta - (1 - \chi)\eta_c \rangle} \right). \end{aligned}$$

and, since on $|\eta_c| \geq b/2 \gg \epsilon_0$,

$$C\epsilon_0 |\operatorname{Re}(\eta - (1 - \chi)\eta_c)| \geq |\operatorname{Im}(\eta - (1 - \chi)\eta_c)|,$$

we have that the imaginary part of the phase satisfies

$$\begin{aligned} \operatorname{Im} \Phi & \geq \operatorname{Im} \Psi(\alpha, \beta) + c(|\langle \alpha_\xi \rangle| |y|^2 + |\langle \beta_\xi \rangle| |y'|^2) + c\delta_1 |\eta - (1 - \chi)\eta_c| \\ & \quad - |y - y'| |\operatorname{Im}((1 - \chi)\eta_c)| \\ & \geq \operatorname{Im} \Psi(\alpha, \beta) + c(|\langle \alpha_\xi \rangle| |y|^2 + |\langle \beta_\xi \rangle| |y'|^2) + c\delta_1 |\eta - (1 - \chi)\eta_c| \\ & \quad - C\epsilon_1 \frac{|(1 - \chi)\eta_c|^2}{|\langle \alpha_\xi \rangle + \langle \beta_\xi \rangle|} \\ & \geq \operatorname{Im} \Psi(\alpha, \beta) + c(|\langle \alpha_\xi \rangle| |y|^2 + |\langle \beta_\xi \rangle| |y'|^2) + c\delta_1 |\eta - (1 - \chi)\eta_c| \\ & \quad - C\epsilon_1 |(1 - \chi)^2 \eta_c| \\ & \geq \operatorname{Im} \Psi(\alpha, \beta) + c(|\langle \alpha_\xi \rangle| |y|^2 + |\langle \beta_\xi \rangle| |y'|^2) + c\delta_1 |\eta - (1 - \chi)\eta_c|, \end{aligned}$$

where we have used that α_ξ and β_ξ are comparable and taken $\epsilon_0 \ll \delta_1$ small enough. Thus, we may apply the method of complex stationary phase to obtain the result. \square

The next result gives the description of the rightmost term in (6.1). For a simpler case capturing the idea of the proof, see [15, theorem 2].

PROPOSITION 6.2. *Suppose P is defined by (6.2) with p satisfying (6.3). Then, for G satisfying (3.1) with $\epsilon_0 > 0$ small enough,*

$$\Pi_\Lambda T_\Lambda P S_\Lambda \Pi_\Lambda = \Pi_\Lambda b_P \Pi_\Lambda + \mathcal{O}(h^\infty)_{\langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)}$$

where

$$b_P \sim \sum_{j=0}^\infty h^j b_j, \quad b_j \in S^{m-j}, \quad b_0 = p|_\Lambda.$$

PROOF. Lemma 6.1 shows that we need to prove

$$(6.7) \quad \Pi_\Lambda K_P \Pi_\Lambda = \Pi_\Lambda b \Pi_\Lambda + \mathcal{O}(h^\infty)_{\langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)},$$

where K_P is given by (6.4). Propositions 5.7 and 5.8 show that, modulo negligible terms, the Schwartz kernel of the left-hand side is given by

$$\int_\Lambda \int_\Lambda e^{\frac{i}{h}(\psi(\alpha, \gamma) + \Psi(\gamma, \gamma') + \psi(\gamma', \beta) + 2iH(\gamma) + 2iH(\beta))} a(\alpha, \gamma) a_P(\gamma, \gamma') a(\gamma', \beta) d\gamma d\gamma',$$

where the support property of a (see (5.2)) shows that integration is over a compact set. An application of the complex stationary phase theorem produces a phase (with critical values taken for almost analytic continuation—see [25, Theorem 2.3, p. 148])

$$\psi_1(\alpha, \beta) = \text{c.v.}_{\gamma, \gamma'} (\psi(\alpha, \gamma) + \Psi(\gamma, \gamma') + \psi(\gamma', \beta) + 2iH(\gamma)).$$

If we show that $\psi_1(\alpha, \alpha) = -2iH(\alpha)$, then the uniqueness part of Proposition 5.3 shows that (modulo negligible terms) we can take $\psi_1 = \psi$. To see this, we claim that for $\alpha = \beta$ the critical point is given by $\gamma = \gamma' = \alpha$, that is,

$$(6.8) \quad \begin{aligned} d_\gamma(\psi(\alpha, \gamma) + \Psi(\gamma, \gamma') + \psi(\gamma', \alpha) + 2iH(\gamma))|_{\gamma=\gamma'=\alpha=0} &= 0, \\ d_{\gamma'}(\psi(\alpha, \gamma) + \Psi(\gamma, \gamma') + \psi(\gamma', \alpha) + 2iH(\gamma))|_{\gamma=\gamma'=\alpha=0} &= 0. \end{aligned}$$

To see this, we first use the formula (4.9) for Ψ to obtain

$$(6.9) \quad d_\gamma \Psi(\gamma, \gamma')|_{\gamma=\gamma'} = \zeta dz|_\Lambda = -d_{\gamma'} \Psi(\gamma, \gamma')|_{\gamma=\gamma'}.$$

This immediately gives the second equation in (6.8).

We then consider

$$(6.10) \quad \begin{aligned} d_\gamma(\psi(\alpha, \gamma) + \Psi(\gamma, \gamma') + \psi(\gamma', \beta) + 2iH(\gamma))|_{\gamma'=\alpha} &= \\ d_\gamma(\psi(\alpha, \gamma) + 2iH(\gamma) + \psi(\gamma, \gamma') - \psi(\gamma, \gamma') + \Psi(\gamma, \gamma'))|_{\gamma'=\alpha}. \end{aligned}$$

The last line in (5.36) and (6.9) give

$$d_\gamma(-\psi(\gamma, \gamma') + \Psi(\gamma, \gamma'))|_{\gamma=\gamma'} = 0.$$

Therefore, to obtain the first equation in (6.8), it is enough to have

$$d_\gamma(\psi(\alpha, \gamma) + 2iH(\gamma) + \psi(\gamma, \gamma'))|_{\gamma=\gamma'=\alpha} = 0,$$

which follows from the first line of (5.36) together with $\psi(\alpha, \alpha) = -2iH(\alpha)$. Since $\psi(\alpha, \alpha) = -2iH(\alpha)$, the critical value is given by $\psi_1(\alpha, \alpha) = \psi(\alpha, \alpha)$. It follows that

$$\begin{aligned} \Pi_\Lambda K_P \Pi_\Lambda u(\alpha) &= h^{-n} \int_\Lambda e^{\frac{i}{h}\psi(\alpha, \beta)} c(\alpha, \beta, h) e^{-2H(\beta)} u(\beta) d\beta \\ &+ \mathcal{O}(h^\infty \|u\|_{(\xi)^N L^2(\Lambda)} (\xi)^{-N} L^2(\Lambda)), \end{aligned}$$

where c satisfies (5.3) (and the support property in (5.2)). Arguing as in (5.7)–(5.8), we see that the terms in the expansion of c satisfy transport equations of (5.39) and hence are determined by their values on the diagonal.

Assume that we have obtained $b_j, j = 0, \dots, J - 1$ (the case of $J = 0$, that is, no b_j 's, is also allowed as the first step) so that

$$(6.11) \quad \Pi_\Lambda K_P \Pi_\Lambda = \Pi_\Lambda \left(\sum_{j=0}^{J-1} \langle \alpha_\xi \rangle^{-j} h^j b_j \right) \Pi_\Lambda + R_\Lambda^J,$$

where

$$\begin{aligned} R_\Lambda^J u(\alpha) &= h^{J-n} \langle \alpha_\xi \rangle^{-J} \int_\Lambda e^{\frac{i}{h}\psi(\alpha, \beta)} a^J(\alpha, \beta) e^{-2H(\beta)/h} u(\beta) d\beta, \\ a^J &\sim a_0^J + h \langle \alpha_\xi \rangle^{-1} a_1^J + \dots, \end{aligned}$$

with a_k^J satisfying the transport equations of Section 5.2. If we apply the method of stationary phase to the kernel of the first term on the right-hand side of (6.11), we obtain, by the inductive hypothesis, a kernel with the expansion

$$e^{\frac{i}{h}\psi(\alpha, \beta)} (a_0 + \dots + h^{J-1} \langle \alpha_\xi \rangle^{-J+1} a_{J-1} + h^J \langle \alpha_\xi \rangle^{-J} r_0^J + h^{J+1} \langle \alpha_\xi \rangle^{-J-1} r_1^J + \dots),$$

where a_j 's are the same as in (6.5). Again, all the terms satisfy transport equations and hence are uniquely determined from their values on the diagonal. Hence, if we put

$$b_J(\alpha) := r_0^J(\alpha, \alpha) + a_0^J(\alpha, \alpha),$$

we obtain (6.11) with J replaced by $J + 1$. When $J = 0, b_J(\alpha) = a_0(\alpha, \alpha) = p|_\Lambda(\alpha)$. □

PROPOSITION 6.3. *Suppose $p_1 \in S^{m_1}$ and $p_2 \in S^{m_2}$. Then, for G satisfying (3.1) with $\epsilon_0 > 0$ small enough,*

$$\Pi_\Lambda p_1 \Pi_\Lambda p_2 \Pi_\Lambda = \Pi_\Lambda b \Pi_\Lambda + \mathcal{O}(h^\infty)_{(\xi)^N L^2(\Lambda) \rightarrow (\xi)^{-N} L^2(\Lambda)}$$

where

$$b \sim \sum_{j=0}^\infty h^j b_j, \quad c_j \in S^{m_1+m_2-j}, \quad b_0 = p_1 p_2.$$

PROOF. From Propositions 5.7 and 5.8, we have that, modulo negligible terms, the Schwartz kernel of $\Pi_\Lambda b_1 \Pi_\Lambda b_2 \Pi_\Lambda$ is given by

$$h^{-3n} \int_\Lambda \int_\Lambda e^{\frac{i}{h}(\psi(\alpha, \gamma) + 2iH(\gamma) + \psi(\gamma, \gamma') + 2H(\gamma') + \psi(\gamma', \beta) + 2iH(\beta))} p_1(\gamma) a(\alpha, \gamma) a(\gamma, \gamma') p_2(\gamma') a(\gamma', \beta) d\gamma d\gamma'$$

where the support property of a (see (5.2)) shows that integration is over a compact set. As in the proof of Proposition 6.2, we apply complex stationary phase to the integral resulting in the phase

$$\psi_1(\alpha, \beta) = \text{c.v.}_{\gamma, \gamma'}(\psi(\alpha, \gamma) + \psi(\gamma, \gamma') + \psi(\gamma', \beta) + 2iH(\gamma) + 2iH(\gamma')).$$

It follows from (5.36) that, modulo negligible terms, we may take $\psi_1(\alpha, \beta) = \psi(\alpha, \beta)$ and that, when $\alpha = \beta$, the critical point given by $\gamma = \gamma' = \alpha$ and hence that

$$\begin{aligned} \Pi_\Lambda K_P \Pi_\Lambda u(\alpha) &= h^{-n} \int_\Lambda e^{\frac{i}{h}\psi(\alpha, \beta)} c(\alpha, \beta, h) e^{-2H(\beta)} u(\beta) d\beta \\ &\quad + \mathcal{O}(h^\infty \|u\|_{\langle \xi \rangle^N L^2(\Lambda)})_{\langle \xi \rangle^{-N} L^2(\Lambda)}, \end{aligned}$$

where c satisfies (5.3) (and the support property in (5.2)). Arguing as in (5.7)–(5.8) we see that the terms in the expansion of c satisfy transport equations of (5.39) and hence are determined by their values on the diagonal. Arguing as in the last paragraph of the proof of Proposition 6.2 then completes the proof. \square

6.2 Compactness properties of the spaces $H^t(\Lambda)$

We next study the compactness and trace class properties for operators on the spaces $H^m(\Lambda)$.

We start with the following:

LEMMA 6.4. *There is $h_0 > 0$ such that for all $s \in \mathbb{R}$ and $0 < h < h_0$*

$$(6.12) \quad (hD_\alpha)^\gamma \Pi_\Lambda = \mathcal{O}(1) : \langle \xi \rangle^{s-|\gamma|} L^2(\Lambda) \rightarrow \langle \xi \rangle^s L^2(\Lambda)$$

and

$$(6.13) \quad t > s \implies H^t(\Lambda) \hookrightarrow H^s(\Lambda) \text{ is compact.}$$

PROOF. To prove (6.12) we show the equivalent fact that the operator

$$\langle \xi \rangle^{-s} (hD_\alpha)^\gamma \Pi_\Lambda \langle \xi \rangle^{s-|\gamma|} : L^2(\Lambda) \rightarrow L^2(\Lambda)$$

is uniformly bounded. By Proposition 4.6, the kernel of this operator is given, modulo acceptable errors, by

$$\begin{aligned} &h^{-n} e^{\frac{i}{h}\Psi(\alpha, \beta)} ((\partial_\alpha \Psi)^\gamma k(\alpha, \beta) + \mathcal{O}(\langle \text{Re } \alpha_\xi \rangle^{|\gamma|-1})) \langle \text{Re } \beta_\xi \rangle^{s-|\gamma|} \langle \text{Re } \alpha_\xi \rangle^{-s} \tilde{\chi}, \\ \tilde{\chi} &:= \chi(|\text{Re } \alpha_x - \text{Re } \beta_x|) \chi(\min(\langle \text{Re } \beta_\xi \rangle, \langle \text{Re } \alpha_\xi \rangle)^{-1} |\text{Re } \alpha_\xi - \text{Re } \beta_\xi|), \quad \chi \in C_c^\infty(\mathbb{R}), \end{aligned}$$

where Ψ is defined in (4.9). Now, on the support of the integrand, $c \langle \text{Re } \alpha_\xi \rangle \leq \langle \text{Re } \beta_\xi \rangle \leq C \langle \text{Re } \beta_\xi \rangle$ and therefore, $|\partial_\alpha \Psi| \leq C \langle \text{Re } \beta_\xi \rangle$. In particular, after conjugation by $e^{H/h}$, the kernel is bounded by

$$Ch^{-n} e^{c(\langle \text{Re } \alpha_\xi \rangle |\text{Re } \alpha_x - \text{Re } \beta_x|^2 + \langle \text{Re } \alpha_\xi \rangle^{-1} |\text{Re } \alpha_\xi - \text{Re } \beta_\xi|^2)/h} \tilde{\chi},$$

and hence, by Schur’s test for L^2 -boundedness, the operator is uniformly bounded on L^2 .

We prove a slightly stronger statement, namely that

$$T_\Lambda(H^t(\Lambda)) \hookrightarrow \langle \xi \rangle^{-s} L^2(\Lambda)$$

is compact, to see (6.13). For that we observe that for $u \in T_\Lambda(H^t(\Lambda))$, $u = \Pi_\Lambda u$, and (6.12) shows that for $m \in \mathbb{Z}$ and $k \in \mathbb{N}$,

$$\Pi_\Lambda : \langle \xi \rangle^{-m} L^2(\Lambda) \rightarrow H^{k,m-k}(\Lambda), \quad H^{r,s}(\Lambda) := \langle hD_{x,\xi} \rangle^{-r} \langle \xi \rangle^{-s} L^2(\Lambda).$$

Hence, by interpolation,

$$\Pi_\Lambda : \langle \xi \rangle^{-t} L^2(\Lambda) \rightarrow H^{r,t-r}(\Lambda), \quad r \geq 0, t \in \mathbb{R}.$$

Setting $r = (t-s)/2 > 0$ we obtain continuity of $T_\Lambda(H^t(\Lambda)) \hookrightarrow H^{r, \frac{s+t}{2}}(\Lambda)$. The lemma then follows from Rellich’s theorem: $H^{r,s+t}(\Lambda) \hookrightarrow \langle \xi \rangle^s L^2(\Lambda)$, $r > 0$, is compact. \square

The next lemma provides trace class properties needed in the study of determinants:

LEMMA 6.5. *For $t > 3n + s$ the inclusion $H^t(\Lambda) \hookrightarrow H^s(\Lambda)$ is of trace class.*

PROOF. First, note that for all $r \in \mathbb{R}$, $m_r(\alpha, \alpha^*) := \langle \alpha_\xi \rangle^{\frac{r}{2}} \langle \alpha^* \rangle^r$ is an order function in the sense of [40, sec. 4.4.1] and for $r < -2n$

$$\int_{T^*\Lambda} m_r(\alpha, \alpha^*) d\alpha d\alpha^* < \infty.$$

Therefore (see [40, (C.3.6)] or [7, chap. 8]) if $\langle \xi \rangle^{-s} A \langle \xi \rangle^s \in \Psi(m_r)$ for $r < -2n$, then $A : \langle \xi \rangle^{-s} L^2(\Lambda) \rightarrow \langle \xi \rangle^{-s} L^2(\Lambda)$ is of trace class.

On the other hand, Lemma 6.4 shows that

$$A := \langle \alpha_\xi \rangle^{-\frac{r}{2}} \langle hD_\alpha \rangle^{-\frac{r}{2}} = \mathcal{O}(1) : T(H^t(\Lambda)) \rightarrow \langle \xi \rangle^{-s} L^2(\Lambda), \quad r = \frac{2(s-t)}{3}.$$

Also, $A \in \Psi(m_{-r})$ is elliptic and invertible and hence $A^{-1} \in \Psi(m_r)$. Therefore A^{-1} is of trace class if $r = \frac{2(s-t)}{3} < -2n$, that is, when $t > 3n + s$. We conclude that

$$\begin{aligned} & \|\text{Id}\|_{\mathcal{L}^1(T(H^t(\Lambda)), \langle \xi \rangle^{-s} L^2(\Lambda))} \\ & \leq \|A^{-1}\|_{\mathcal{L}^1(\langle \xi \rangle^{-s} L^2(\Lambda), \langle \xi \rangle^{-s} L^2(\Lambda))} \|A\|_{T(H^t(\Lambda)) \rightarrow \langle \xi \rangle^{-s} L^2(\Lambda)} < \infty, \end{aligned}$$

where \mathcal{L}^1 denotes the trace class. \square

7 Zeroth-Order Operators and Viscosity Limits

Recall that the constructions in the previous sections depend only on finitely many S^1 norms of G determining

$$\Lambda = \Lambda_G = \{(x + iG_\xi, \xi - iG_x) \mid (x, \xi) \in T^*\mathbb{T}^n\}.$$

(Unless we worked with different G 's, we suppress the dependence on G in Λ_G .) Therefore, we start by fixing $h > 0, \epsilon_0 > 0$ small enough and $N_0 > 0$ large enough such that if

$$(7.1) \quad \sup_{|\alpha|+|\beta|\leq N_0} \langle \xi \rangle^{1-|\beta|} |\partial_x^\alpha \partial_\xi^\beta G| \leq \epsilon_0,$$

the constructions of T_Λ, S_Λ , are valid and

$$B_\Lambda - \Pi_\Lambda : \langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda) \quad \text{for all } N \in \mathbb{N}.$$

7.1 Elliptic regularity in deformed spaces

We begin with the following preliminary elliptic regularity lemma.

LEMMA 7.1. *Suppose that $G \in S^1(T^*\mathbb{T}^n)$ satisfies (7.1) and*

$$(7.2) \quad \sup_{|\alpha|+|\beta|\leq 1} |\langle \xi \rangle^{1-|\beta|} \partial_x^\alpha \partial_\xi^\beta G| \leq \epsilon_1$$

for a fixed ϵ_1 . Suppose also that E is given by (6.2) with e (replacing p) satisfying (6.3) and

$$|e(z, \zeta)| \geq c_1 |\zeta|^m, \quad |\zeta| \geq C, \quad |\text{Im } z| \leq \epsilon_1, \quad |\text{Im } \zeta| \leq \epsilon_1 \langle \zeta \rangle.$$

Then $E : H_\Lambda^s \rightarrow H_\Lambda^{s-m}$ is a Fredholm operator and there exists

$$C_1 = C_1(s, \epsilon_1, E, N) > 0 \quad \text{such that} \quad \frac{1}{2} c_1 \|u\|_{H_\Lambda^s} \leq \|Eu\|_{H_\Lambda^{s-m}} + C_1 \|u\|_{H_\Lambda^{-N}}.$$

PROOF. The assumptions on e guarantee that

$$|e|_\Lambda(\alpha) \geq c_1 |\alpha_\xi|^m, \quad |\alpha_\xi| \geq C, \quad \alpha \in \Lambda.$$

Proposition 6.2 then shows

$$\Pi_\Lambda T_\Lambda E S_\Lambda \Pi_\Lambda = \Pi_\Lambda \tilde{b}_E \Pi_\Lambda + R'$$

with

$$\tilde{b}_E \sim \sum_j \tilde{b}_j, \quad \tilde{b}_j \in S^{m-j}, \quad \tilde{b}_0 = e|_\Lambda$$

and

$$\|R'\|_{\langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-s+m} L^2(\Lambda)} \leq C'_2 = C'_2(s, \epsilon_1, E, N).$$

Next, by Proposition 6.3,

$$\Pi_\Lambda \overline{\tilde{b}_E} \Pi_\Lambda \tilde{b}_E \Pi_\Lambda = \Pi_\Lambda b_E \Pi_\Lambda + R''$$

with

$$b_E \sim \sum_j b_j, \quad b_j \in S^{m-j}, \quad b_0 = |e|_\Lambda|^2,$$

and

$$\|R''\|_{(\xi)^N L^2(\Lambda) \rightarrow (\xi)^{-s+m} L^2(\Lambda)} \leq C_2'' = C_2''(s, \epsilon_1, E, N).$$

Since $|b_0| \geq c_1^2 |\alpha_\xi|^{2m}$ on $|\alpha_\xi| \geq C$, there is $b_\infty \in S^{-\infty}$ such that with $b := b_E + b_\infty$,

$$|b| \geq \frac{2}{3} c_1^2 \langle |\alpha_\xi| \rangle^{2m}$$

and

$$\begin{aligned} (\Pi_\Lambda T_\Lambda E S_\Lambda \Pi_\Lambda)^* (\Pi_\Lambda T_\Lambda E S_\Lambda \Pi_\Lambda) &= \Pi_\Lambda b \Pi_\Lambda + R, \\ \|R\|_{(\xi)^N L^2(\Lambda) \rightarrow (\xi)^{-s+m} L^2(\Lambda)} &\leq C_2 = C_2(s, \epsilon_1, E, N). \end{aligned}$$

For $u \in H^s(\Lambda)$ we compute

$$\begin{aligned} \|Eu\|_{H_\Lambda^{s-m}}^2 &= \langle \Pi_\Lambda T_\Lambda E S_\Lambda \Pi_\Lambda T_\Lambda u, \Pi_\Lambda T_\Lambda E S_\Lambda \Pi_\Lambda T_\Lambda u \rangle_{(\xi)^{-s+m} L^2} \\ &= \langle \Pi_\Lambda (b + R) T_\Lambda u, T_\Lambda u \rangle_{(\xi)^{-s+m} L^2} \\ &\geq \langle b T_\Lambda u, T_\Lambda u \rangle_{(\xi)^{-s+m} L^2} - C_2 \|u\|_{H_\Lambda^{-N}}^2 \\ &\geq \frac{1}{4} c_1^2 \|u\|_{H_\Lambda^s}^2 - C_1^2 \|u\|_{H_\Lambda^{-N}}^2 \end{aligned}$$

with $C_1 = C_1(s, \epsilon_1, E, N)$. Therefore,

$$(7.3) \quad \frac{1}{2} c_1 \|u\|_{H_\Lambda^s} \leq \|Eu\|_{H_\Lambda^{s-m}} + C_1 \|u\|_{H_\Lambda^{-N}}.$$

We now note that for all s ,

$$\begin{aligned} \langle E^* u, v \rangle_{H_\Lambda^s} &= \langle u, E v \rangle_{H_\Lambda^s} = \langle T_\Lambda u, \Pi_\Lambda T_\Lambda E S_\Lambda \Pi_\Lambda T_\Lambda v \rangle_{(\xi)^{-s} L^2(\Lambda)} \\ (7.4) \quad &= \langle T_\Lambda u, (b + R) T_\Lambda v \rangle_{(\xi)^{-s} L^2(\Lambda)} \\ &= \langle (\bar{b} + R^*) T_\Lambda u, T_\Lambda v \rangle_{(\xi)^{-s} L^2(\Lambda)}. \end{aligned}$$

Using (7.4), we obtain

$$\begin{aligned} \|E^* u\|_{H_\Lambda^{s-m}}^2 &= \langle E^* u, E^* u \rangle_{H_\Lambda^{s-m}} = \langle (\bar{b} + R^*) T_\Lambda u, T_\Lambda u \rangle_{(\xi)^{-s+m} L^2} \\ &\geq \frac{1}{4} c_1^2 \|u\|_{H_\Lambda^s}^2 - C_1^2 \|u\|_{H_\Lambda^{-N}}^2. \end{aligned}$$

Therefore

$$(7.5) \quad \frac{1}{2} c_1 \|u\|_{H_\Lambda^s} \leq \|E^* u\|_{H_\Lambda^{s-m}} + C_1 \|u\|_{H_\Lambda^{-N}}.$$

Combining (7.3) and (7.5) with s replaced by $m - s$ and applying Lemma 6.4, we have for $N > m - s$ that $H_\Lambda^{s-m} \rightarrow H_\Lambda^{-N}$ is compact. Thus, we have proved that $E : H_\Lambda^s \rightarrow H_\Lambda^{s-m}$ is a Fredholm operator. \square

7.2 Zeroth-order operators on deformed spaces

We now work in the setting of Theorem 1.2. Let $P \in \Psi^0$ satisfy the assumptions there, $G_0 \in S^1(T^*\mathbb{T}^n)$, and $C > 0$ such that

$$(7.6) \quad H_p G_0 > 0, \quad \{|\xi| > C\} \cap \{p = 0\}.$$

Define the \mathbb{R} -symplectic I -Lagrangian submanifold $\Lambda_\theta \subset \widetilde{T^*\mathbb{T}^n}$ by

$$\Lambda_\theta = \{(x + i\theta\partial_\xi G_0, \xi - i\theta\partial_x G_0) \mid (x, \xi) \in T^*\mathbb{T}^n\}.$$

We work with the spaces $H_{\Lambda_\theta}^m$ as defined in (4.7). Observe that for $|\theta|$ small enough, θG_0 satisfies (7.1). To avoid cumbersome notation, we will suppress the dependence of Λ_θ on θ .

For $u \in H_{\Lambda}^m$ we have $T_\Lambda P u = \Pi_\Lambda T_\Lambda P S_\Lambda T_\Lambda u$. By Proposition 6.2

$$(7.7) \quad \Pi_\Lambda T_\Lambda P S_\Lambda \Pi_\Lambda = \Pi_\Lambda b_P \Pi_\Lambda + R_1, \quad \Pi_\Lambda T_\Lambda \Delta S_\Lambda \Pi_\Lambda = \Pi_\Lambda a_\Delta \Pi_\Lambda + R_2,$$

where $R_i : \langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)$,

$$b_P - p|_\Lambda \in S^{-1}, \quad a_\Delta + (\zeta)^2|_\Lambda \in S^{-1}.$$

Now,

$$p|_\Lambda = p(x, \xi) - i\theta H_p G_0 + \mathcal{O}(\theta^2)_{S^0}.$$

In particular, by (7.6), there are $c, C > 0$ such that for $\theta > 0$ small enough,

$$\operatorname{Im} p|_\Lambda \leq (-c + C\langle \xi \rangle^{-1})\theta, \quad \text{on } |\operatorname{Re} p|_\Lambda| \leq c.$$

In particular, there exists $b_\infty \in S^{-\infty}$ such that for $e_0 := b_P + b_\infty$, there are $c_0, C_0 > 0$ satisfying

$$(7.8) \quad |e_0| > c_0\theta > 0, \quad \operatorname{Im} e_0 \leq -c_0\theta \text{ on } |\operatorname{Re} e_0| < c_0, \quad |\operatorname{Im} e_0| \leq C_0(\theta + \langle \xi \rangle^{-1}).$$

By (7.7), we also have

$$(7.9) \quad \Pi_\Lambda T_\Lambda P S_\Lambda \Pi_\Lambda = \Pi_\Lambda e_0 \Pi_\Lambda + R_0$$

with $R_0 : \langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)$ uniformly over $0 \leq \theta \leq \theta_0$.

To analyse the contribution of the Laplacian we note that

$$\operatorname{Re}(\zeta)^2|_\Lambda \geq (1 - C\theta^2)|\xi|^2, \quad |\operatorname{Im}(\zeta)^2|_\Lambda| \leq C\theta|\xi|^2.$$

Therefore, for $\theta > 0$ small enough, we can find $a_\infty \in S^{-\infty}$ such that

$$(7.10) \quad \operatorname{Re}(a_\Delta + a_\infty) \geq \frac{1}{2}\langle \xi \rangle^2, \quad |\operatorname{Im}(a_\Delta + a_\infty)| \leq C\theta|\xi|^2 + C|\xi|,$$

and we have

$$(7.11) \quad \Pi_\Lambda T_\Lambda (P + i\nu\Delta) S_\Lambda \Pi_\Lambda = \Pi_\Lambda e_\nu \Pi_\Lambda + R_\nu, \quad e_\nu := e_0 - i\nu(a_\Delta + a_\infty),$$

where, by (7.7) and (7.9), $R_\nu : \langle \xi \rangle^N L^2(\Lambda) \rightarrow \langle \xi \rangle^{-N} L^2(\Lambda)$ uniformly in $0 \leq \nu \leq 1, 0 \leq \theta \leq \theta_0$.

The next lemma gives us crucial properties of e_ν :

LEMMA 7.2. *There exist $c_1, \theta_0, \nu_0 > 0$ such that for all $0 \leq \nu \leq \nu_0$ and for all $0 < \theta \leq \theta_0$*

$$(7.12) \quad \begin{aligned} &|e_\nu| > c_1\theta(1 + \nu|\xi|^2) > 0, \\ &\operatorname{Im} e_\nu \leq -c_1\theta(1 + \nu|\xi|^2) \quad \text{on } |\operatorname{Re} e_\nu| < c_1(1 + \nu|\xi|^2). \end{aligned}$$

PROOF. We consider two cases. First, suppose $|\xi| \geq M\nu^{-1/2}$. Then, by (7.10) and (7.8), there are $c_2, C_2 > 0$ such that

$$\operatorname{Im} e_\nu \leq -c_2(M^2 + \nu|\xi|^2) + C_2(\theta + M^{-1}\nu^{1/2}).$$

Therefore, setting

$$M = \max(1, 2\sqrt{C_2/c_2}),$$

(7.12) holds on $|\xi| \geq M\nu^{-1/2}$ (uniformly in $0 \leq \nu \leq 1$ and $0 \leq \theta \leq 1$).

We next consider the case $|\xi| \leq M\nu^{-1/2}$. If $c_0 \geq 2|\operatorname{Re} e_\nu|$, then

$$\frac{c_0}{2} \geq |\operatorname{Re} e_\nu| \geq |\operatorname{Re} e_0| - C\theta M^2 - C\nu^{1/2}M.$$

Choosing θ_0 and ν_0 small enough, we obtain $|\operatorname{Re} e_0| \leq c_0$ and hence

$$\operatorname{Im} e_\nu \leq \operatorname{Im} e_0 \leq -c_0\theta,$$

which completes the proof of (7.12). □

7.3 Fredholm properties and meromorphy of the resolvent

We add a localized absorbing potential to $P + i\nu\Delta$ to obtain invertibility. That is, for $q \in C_c^\infty(\Lambda; [0, \infty))$ we define

$$(7.13) \quad P_{q,\nu} := P + i\nu\Delta - iQ, \quad Q := S_\Lambda \Pi_\Lambda q \Pi_\Lambda T_\Lambda.$$

This family includes the operator $P = P_{0,0}$ and the viscous operator $P + i\nu\Delta = P_{0,\nu}$. We note that

$$(7.14) \quad \begin{aligned} \Pi_\Lambda T_\Lambda Q S_\Lambda \Pi_\Lambda &= \Pi_\Lambda T_\Lambda S_\Lambda \Pi_\Lambda q \Pi_\Lambda T_\Lambda S_\Lambda \Pi_\Lambda \\ &= \Pi_\Lambda P_\Lambda \Pi_\Lambda q \Pi_\Lambda P_\Lambda \Pi_\Lambda = \Pi_\Lambda q \Pi_\Lambda, \end{aligned}$$

where we recall that $P_\Lambda = T_\Lambda S_\Lambda$ satisfies $\Pi_\Lambda = P_\Lambda \Pi_\Lambda$. We record the following lemma for use later.

LEMMA 7.3. *The adjoint of $P : H^s(\Lambda) \rightarrow H^s(\Lambda)$ satisfies*

$$\langle P^*u, v \rangle_{H^s(\Lambda)} = \langle (\Pi_\Lambda \bar{e}_0 \Pi_\Lambda + R_0^*)Tu, Tv \rangle_{\langle \xi \rangle^{-s}L^2(\Lambda)},$$

and the adjoint of $P_{q,\nu} : H^{s+2}(\Lambda) \rightarrow H^s(\Lambda)$, $\nu > 0$, satisfies

$$\langle P_{q,\nu}^*u, v \rangle_{H^s(\Lambda)} = \langle (\Pi_\Lambda (\bar{e}_\nu - iq) \Pi_\Lambda + R_\nu^*)Tu, Tv \rangle_{\langle \xi \rangle^{-s}L^2(\Lambda)}.$$

PROOF. The lemma follows from (7.4). □

We start by proving that $P - \omega$ is Fredholm on H_Λ^s . In particular, the next lemma proves the first part of Theorem 1.2 with $\mathcal{X} = H_\Lambda^s$.

LEMMA 7.4. *There is $\omega_0 > 0$ such that for $\text{Im } \omega > -\omega_0\theta$ and $|\text{Re } \omega| < \omega_0$,*

$$P - \omega : H_\Lambda^s \rightarrow H_\Lambda^s$$

is a Fredholm operator. For $\text{Im } \omega \gg 1$, $P - \omega$ is invertible with inverse $\mathcal{R}(\omega)$ satisfying

$$\|\mathcal{R}(\omega)\|_{H_\Lambda^s \rightarrow H_\Lambda^s} \leq \frac{C}{\text{Im } \omega}, \quad \text{Im } \omega \geq C, \quad |\text{Re } \omega| < \omega_0.$$

In particular, $\mathcal{R}(\omega) : H_\Lambda^s \rightarrow H_\Lambda^s$ is a meromorphic family of operators for $\omega \in (-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$.

PROOF. First, observe that by (7.11) with $\nu = 0$ and using Proposition 6.3

$$\begin{aligned} & \Pi_\Lambda \langle \xi \rangle^{2s} (\Pi_\Lambda T_\Lambda (P - \omega) S_\Lambda \Pi_\Lambda)^* \langle \xi \rangle^{-2s} (\Pi_\Lambda T_\Lambda (P - \omega) S_\Lambda \Pi_\Lambda) \\ &= \Pi_\Lambda |e_0 - \omega|^2 \Pi_\Lambda + \tilde{R}_0 \end{aligned}$$

where

$$\tilde{R}_0 : \langle \xi \rangle^s L^2(\Lambda) \rightarrow \langle \xi \rangle^{s+1} L^2(\Lambda).$$

By (7.12), there is $c_1 > 0$ such that $\text{Im } e_0 \leq -c_1\theta$ on $|\text{Re } e_0| \leq c_1$. Therefore, on $|\text{Re } e_0| \leq c_1$,

$$|e_0 - \omega|^2 \geq (c_1\theta + \text{Im } \omega)^2 \geq \frac{c_1^2\theta^2}{4} + \max(\text{Im } \omega, 0)^2,$$

where we have taken $\omega_0 = \frac{c_1}{2}$ and $\text{Im } \omega \geq -\omega_0\theta$. Then, using $|\text{Re } \omega| \leq \omega_0$, on $|\text{Re } e_0| \geq c_1$, there is $C > 0$ such that

$$|e_0 - \omega|^2 \geq \frac{c_1^2}{4} + \min(|\text{Im } \omega|^2 - C, 0) \geq c_1(1 + |\text{Im } \omega|^2).$$

In particular, for $u \in H_\Lambda^s$,

$$\begin{aligned} & \|(P - \omega)u\|_{H_\Lambda^s}^2 \\ &= |\langle (P - \omega)u, (P - \omega)u \rangle_{H_\Lambda^s}| \\ &= |\langle \Pi_\Lambda T_\Lambda (P - \omega) S_\Lambda \Pi_\Lambda T_\Lambda u, \Pi_\Lambda T_\Lambda (P - \omega) S_\Lambda \Pi_\Lambda T_\Lambda u \rangle_{\langle \xi \rangle^{-s} L^2}| \\ &\geq |\langle |e_0 - \omega|^2 T_\Lambda u, T_\Lambda u \rangle_{\langle \xi \rangle^{-s} L^2}| - |\langle \tilde{R}_0 T_\Lambda u, T_\Lambda u \rangle_{\langle \xi \rangle^{-s} L^2}| \\ &\geq c_1(1 + |\text{Im } \omega|^2) \|u\|_{H_\Lambda^s}^2 - C \|u\|_{H_\Lambda^{s-1/2}}^2. \end{aligned}$$

In particular, iterating this argument, we have for any N that there is $C_N > 0$ such that

$$(7.15) \quad \|u\|_{H_\Lambda^s}^2 \leq C(1 + |\text{Im } \omega|^2)^{-1} [\|(P - \omega)u\|_{H_\Lambda^s}^2 + C_N \|u\|_{H_\Lambda^{-N}}^2].$$

By almost exactly the same argument, using Lemma 7.3, we obtain

$$(7.16) \quad \|u\|_{H_\Lambda^s}^2 \leq C(1 + |\text{Im } \omega|^2)^{-1} [\|(P^* - \bar{\omega})u\|_{H_\Lambda^s}^2 + C_N \|u\|_{H_\Lambda^{-N}}^2].$$

Next, by Lemma 6.4, for $N > -s$, the embedding $H_\Lambda^s \rightarrow H_\Lambda^{-N}$ is compact. Therefore, $P - \omega : H_\Lambda^s \rightarrow H_\Lambda^s$ is Fredholm.

Finally, taking $\text{Im } \omega \gg 1$, we may absorb the H_Λ^{-N} error into the left-hand sides of (7.15) and (7.16) to obtain that $P - \omega : H_\Lambda^s \rightarrow H_\Lambda^s$ is invertible with the desired estimate. The meromorphic Fredholm theorem (see, e.g., [11, theorem C.9]) then shows that $\mathcal{R}(\omega)$ is a meromorphic family of operators on $(-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$. \square

We next study the meromorphy of the inverse of $P_{q,v} - \omega$, where $P_{q,v}$ is given in (7.13).

LEMMA 7.5. *There exists $\epsilon_0 > 0$ such that the following holds: For all $s \in \mathbb{R}$, $K \in \mathbb{N}$, $\omega \in \mathbb{C}$, $q \in C_c^\infty(T^*\mathbb{T}^n)$, and $v > 0$ there are $C_0 = C_{s,v,K}$ and $C_1 = C_{s,v,K,N}$ such that for all G satisfying (7.1),*

$$(P_{q,v} - \omega)^K : H_\Lambda^s \rightarrow H_\Lambda^{s-2K}$$

is a Fredholm operator and

$$(7.17) \quad \|u\|_{H_\Lambda^s} \leq C_0 \|(P_{q,v} - \omega)^K u\|_{H_\Lambda^{s-2K}} + C_1 \|u\|_{H_\Lambda^{-N}}.$$

Moreover, $P_{q,v} - \omega : H_\Lambda^s \rightarrow H_\Lambda^{s-2}$ is invertible for $\text{Im } \omega \gg 1$ with inverse $\mathcal{R}_{q,v}(\omega)$ satisfying

$$\|\mathcal{R}_{q,v}(\omega)\|_{H_\Lambda^s \rightarrow H_\Lambda^{s+2}} \leq C v^{-1}, \quad \text{Im } \omega \geq C(1 + v).$$

In particular, for all $v > 0$, $\mathcal{R}_{q,v}(\omega) : H_\Lambda^s \rightarrow H_\Lambda^{s+2}$ is a meromorphic family of operators for $\omega \in \mathbb{C}$.

PROOF. We first note that

$$|\sigma(\Delta)(z, \zeta)| = |\zeta^2| \geq |\text{Re } \zeta|^2 - |\text{Im } \zeta|^2$$

and hence Δ^K satisfies the hypotheses of Lemma 7.1 for any $\epsilon_1 < 1$. In particular, by that lemma $(i v \Delta)^K : H_\Lambda^s \rightarrow H_\Lambda^{s-2K}$ is a Fredholm operator and satisfies

$$(7.18) \quad \|u\|_{H_\Lambda^s} \leq C_1^K v^{-K} \|(i v \Delta)^K u\|_{H_\Lambda^{s-2K}} + v^{-K} C_{s,N,K} \|u\|_{H_\Lambda^{-N}}$$

for any G satisfying (7.1).

Next, observe that $(P_{q,v} - \omega)^K - (i v \Delta)^K : H_\Lambda^s \rightarrow H_\Lambda^{s-2K+2}$ and $H_\Lambda^{s-2K+2} \rightarrow H_\Lambda^{s-2K}$ is compact by Lemma 6.4. Therefore, $(P_{q,v} - \omega)^K : H_\Lambda^s \rightarrow H_\Lambda^{s-2K}$ is a Fredholm operator.

Finally, using (7.18),

$$\begin{aligned} \|u\|_{H_\Lambda^s} &\leq C_{v,K} \|(i v \Delta)^K u\|_{H_\Lambda^{s-2K}} + C_{s,v,N,K} \|u\|_{H_\Lambda^{-N}} \\ &\leq C_{v,K} \|(P_{q,v} - \omega)^K u\|_{H_\Lambda^{s-2K}} + C_{s,v,K} \|u\|_{H_\Lambda^{s-2}} + C_{s,v,N,K} \|u\|_{H_\Lambda^{-N}}. \end{aligned}$$

Estimating $\|u\|_{H_\Lambda^{s-2}}$ by $\|(P_{q,v} - \omega)^K u\|_{H_\Lambda^{s-2K-2}}$ and iterating we obtain (7.17).

For invertibility, let $\text{Im } \omega \gg 1$ and consider

$$\begin{aligned}
 & \| (P_{q,v} - \omega)u \|_{H_\Lambda^{s-2}} \| u \|_{H_\Lambda^{s-2}} \\
 (7.19) \quad & \geq -\text{Im} \langle (P_{q,v} - \omega)u, u \rangle_{H_\Lambda^{s-2}} \\
 & \geq -\text{Im} \langle \Pi_\Lambda (iva - iq - \omega) \Pi_\Lambda T_\Lambda u, T_\Lambda u \rangle_{(\xi)^{-s+2}L^2(\Lambda)} - C_N \| u \|_{H_\Lambda^{-N}}^2
 \end{aligned}$$

where $a = a_\Delta + a_\infty$ is as in (7.10). In particular, for $\text{Im } \omega > 0$,

$$\text{Im } iva - iq - \omega \leq -cv|\xi|^2 + Cv|\xi| - \text{Im } \omega,$$

and for $\text{Im } \omega \geq C_N + 1 + Cv$,

$$\text{Im } iva - iq - \omega \leq -C_N - 1 - cv|\xi|^2.$$

Using this in (7.19), we obtain

$$\| (P_{q,v} - \omega)u \|_{H_\Lambda^{s-2}} \geq Cv \| u \|_{H_\Lambda^s}.$$

This same argument implies that

$$\| (P_{q,v}^* - \bar{\omega})u \|_{H_\Lambda^{s-2}} \geq Cv \| u \|_{H_\Lambda^s}.$$

and hence $P_{q,v} - \omega$ is invertible with inverse as claimed. The meromorphic Fredholm theorem (see, e.g., [11, theorem C.9]) then shows that $\mathcal{R}_{q,v}(\omega)$ is a meromorphic family of operators for $\omega \in \mathbb{C}$. □

7.4 A parametrix for the resolvent of $P_{0,v} - \omega$

We next find q so that the compact perturbation $P_{q,v}$ of $P_{0,v}$ is invertible. This inverse will be used to approximate the inverse of $P_{0,v}$.

LEMMA 7.6. *There are $\omega_0, \nu_0, \theta_0 > 0$ so that for all $\epsilon > 0$ and $\theta \in (0, \theta_0)$, there is $q = q(\epsilon, \theta) \in C_c^\infty(\Lambda; [0, \infty))$ such that for all $\nu \in (0, \nu_0]$ and $\omega \in (-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$, the operators*

$$P_{q,v} - \omega : H_\Lambda^{s+2} \rightarrow H_\Lambda^s \quad \text{and} \quad P_{q,0} - \omega : H_\Lambda^s \rightarrow H_\Lambda^s$$

are invertible with inverse $\mathcal{R}_{q,v}(\omega) := (P_{q,v} - \omega)^{-1}$ satisfying

$$\| \mathcal{R}_{q,v}(\omega) \|_{H_\Lambda^s \rightarrow H_\Lambda^{s-\epsilon}} \leq 1.$$

PROOF. We assume without loss of generality that $\epsilon < \frac{1}{2}$, and observe that by (7.11) and (7.14)

$$\Pi_\Lambda T_\Lambda P_{q,v} S_\Lambda \Pi_\Lambda = \Pi_\Lambda (e_\nu - iq) \Pi_\Lambda + R_\nu.$$

and thus, by Proposition 6.3,

$$\begin{aligned}
 & \Pi_\Lambda \langle \xi \rangle^{2s} (\Pi_\Lambda T_\Lambda (P_{q,v} - \omega) S_\Lambda \Pi_\Lambda)^* \langle \xi \rangle^{-2s} \Pi_\Lambda T_\Lambda (P_{q,v} - \omega) S_\Lambda \Pi_\Lambda \\
 & = \Pi_\Lambda |e_\nu - iq - \omega|^2 \Pi_\Lambda + \tilde{R}_\nu
 \end{aligned}$$

where for all $s \in \mathbb{R}$,

$$\| \tilde{R}_\nu v \|_{H_\Lambda^s} \leq C_0 (\| v \|_{H_\Lambda^{s-1}} + \nu \| v \|_{H_\Lambda^{s+1}} + \nu^2 \| v \|_{H_\Lambda^{s+3}}), \quad 0 \leq \nu \leq 1.$$

Therefore,

$$\begin{aligned}
 & \| (P_{q,v} - \omega)u \|_{H_\Lambda^s}^2 \\
 & \geq | \langle \langle \xi \rangle^\epsilon |e_v - iq - \omega|^2 \langle \xi \rangle^\epsilon \langle \xi \rangle^{-\epsilon} T_\Lambda u, \langle \xi \rangle^{-\epsilon} T_\Lambda u \rangle_{\langle \xi \rangle^{-s} L^2} | \\
 & \quad - | \langle \tilde{R}_v \langle \xi \rangle^\epsilon \langle \xi \rangle^{-\epsilon} T_\Lambda u, \langle \xi \rangle^\epsilon \langle \xi \rangle^{-\epsilon} T_\Lambda u \rangle_{\langle \xi \rangle^{-s} L^2} | \\
 (7.20) \quad & \geq | \langle \langle \xi \rangle^\epsilon |e_v - iq - \omega|^2 \langle \xi \rangle^\epsilon \langle \xi \rangle^{-\epsilon} T_\Lambda u, \langle \xi \rangle^{-\epsilon} T_\Lambda u \rangle_{\langle \xi \rangle^{-s} L^2} | \\
 & \quad - C_0 \left(\| \langle \xi \rangle^{-\epsilon} T_\Lambda u \|_{\langle \xi \rangle^{s-\frac{1}{2}+\epsilon} L^2(\Lambda)}^2 + \nu \| \langle \xi \rangle^{-\epsilon} T_\Lambda u \|_{\langle \xi \rangle^{s+\frac{1}{2}+\epsilon} L^2(\Lambda)}^2 \right. \\
 & \quad \left. + \nu^2 \| \langle \xi \rangle^{-\epsilon} T_\Lambda u \|_{\langle \xi \rangle^{s+\frac{3}{2}+\epsilon} L^2(\Lambda)}^2 \right)
 \end{aligned}$$

Let θ_0, ν_0 , and c_1 be as in Lemma 7.2 and fix $\chi = \chi_\epsilon \in C_c^\infty(\Lambda; [0, 1])$ with

$$\chi \equiv 1 \quad \text{on } \langle \xi \rangle^{2\epsilon} < \frac{8 \max(C_0, 1)}{c_1^2 \min(1, \theta^2)}.$$

Then, let $q = M\chi_\epsilon$ for M to be chosen later and $\omega_0 \leq c_1/2$. On $\text{supp}(1 - \chi_\epsilon)$,

$$\begin{aligned}
 & \langle \xi \rangle^{2\epsilon} |e_v - iq - \omega|^2 \\
 & \geq \langle \xi \rangle^{2\epsilon} (|\text{Re } e_v - \text{Re } \omega|^2 + |\text{Im } e_v - q - \text{Im } \omega|^2) \\
 & \geq \langle \xi \rangle^{2\epsilon} (\min((c_1(1 + \nu|\xi|^2) - |\text{Re } \omega|)^2, |c_1\theta(1 + \nu|\xi|^2) + \text{Im } \omega|^2) \\
 & \geq \frac{1}{4} \langle \xi \rangle^{2\epsilon} c_1^2 \min(1, \theta^2) (1 + \nu|\xi|^2)^2 \\
 & \geq (1 + C_0)(1 + \nu|\xi|^2)^2.
 \end{aligned}$$

On $\chi_\epsilon \equiv 1$, we have

$$\langle \xi \rangle^{2\epsilon} |e_v - iq - \omega|^2 \geq \langle \xi \rangle^{2\epsilon} (M^2 - 4(\omega_0\theta + |e_v|)^2) \geq \frac{1}{4} (M^2 - C)$$

for some $C > 0$ independent of $\nu, \omega_0, \theta \in [0, 1]$. Thus, for $\omega_0 := \min(c_1/2, 1)$ and

$$M^2 := C + 4(1 + C_0) \left(1 + \frac{8 \max(C_0, 1)}{c_1^2 \min(1, \theta^2)} \right)^2,$$

we have

$$\inf_\Lambda \langle \xi \rangle^{2\epsilon} |e_v - iq - \omega|^2 \geq (1 + C_0)(1 + \nu|\xi|^2)^2, \quad 0 \leq \nu \leq 1.$$

In particular, using this in (7.20) yields

$$(7.21) \quad \|u\|_{H_\Lambda^{s-\epsilon}} \leq \| (P_{q,v} - \omega)u \|_{H_\Lambda^s}.$$

As in the proofs of Lemma 7.4, an identical argument using $\| (P_{q,v}^* - \bar{\omega})u \|_{H_\Lambda^s}^2$ implies

$$(7.22) \quad \|u\|_{H_\Lambda^{s-\epsilon}} \leq \| (P_{q,v}^* - \bar{\omega})u \|_{H_\Lambda^s}.$$

Since $P_{0,v} - P_{q,v} : H_\Lambda^{s+2} \rightarrow H_\Lambda^N$ for any N , $P_{q,v} - \omega$ is a Fredholm operator. In particular, (7.21) and (7.22) imply that $\mathcal{R}_{q,v}(\omega)$ exists and satisfies the requisite bounds. \square

7.5 Convergence of the poles of $\mathcal{R}_{0,v}(\omega)$

We now finish the proof of Theorem 1.1.

PROOF OF THEOREM 1.1. First, note that by Lemma 7.6 for $\omega \in (-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$ and $v \in [0, v_0]$, the inverse $\mathcal{R}_{q,v} : H_\Lambda^s \rightarrow H_\Lambda^s$ exists and satisfies

$$(\text{Id} + i\mathcal{R}_{q,v}(\omega)Q) = \mathcal{R}_{q,v}(\omega)(P_{0,v} - \omega).$$

Moreover, by Lemmas 7.4 and 7.5, there is $C_v > 0$ such that for $\omega \in (-\omega_0, \omega_0) + i(C_v, \infty)$, $\mathcal{R}_{0,v}(\omega) : H_\Lambda^s \rightarrow H_\Lambda^s$ exists. Therefore, for ω in this region, the inverse

$$(\text{Id} + i\mathcal{R}_{q,v}(\omega)Q)^{-1} = \mathcal{R}_{0,v}(\omega)(P_{q,v} - \omega) : H_\Lambda^s \rightarrow H_\Lambda^s$$

exists.

Now, for any $N > 0$, $Q : H_\Lambda^s \rightarrow H_\Lambda^{s+N+\epsilon}$ and $\mathcal{R}_{q,v}(\omega) : H_\Lambda^{s+N+\epsilon} \rightarrow H_\Lambda^{s+N}$, with uniform bounds in $v \geq 0$. Therefore, Lemma 6.5 implies that for any s

$$\mathcal{R}_{q,v}(\omega)Q : H_\Lambda^s \rightarrow H_\Lambda^s$$

is trace class with uniformly bounded trace class norm. In particular, for $\omega \in (-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$ the operator

$$\text{Id} + i\mathcal{R}_{q,v}(\omega)Q : H_\Lambda^s \rightarrow H_\Lambda^s$$

is Fredholm with index 0. Thus, by the meromorphic version of Fredholm analyticity (see, for instance, [11, theorem C.10])

$$(\text{Id} + i\mathcal{R}_{q,v}(\omega)Q)^{-1} : H_\Lambda^s \rightarrow H_\Lambda^s$$

is a meromorphic family of operators satisfying

$$(7.23) \quad \mathcal{R}_{0,v}(\omega) = (\text{Id} + i\mathcal{R}_{q,v}(\omega)Q)^{-1}\mathcal{R}_{q,v}(\omega).$$

For q chosen in Lemma 7.6, $\mathcal{R}_{q,v}(\omega)$ is analytic in $(-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$. Hence, the eigenvalues of $P_{0,v}$ on H_Λ^0 agree, with multiplicity, with the zeroes of

$$f_v(\omega) := \det_{H_\Lambda^0}(\text{Id} + i\mathcal{R}_{q,v}(\omega)Q).$$

LEMMA 7.7. *We have*

$$f_v(\omega) \xrightarrow{v \rightarrow 0} f_0(\omega)$$

uniformly on compact subsets of $\omega \in (-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$.

PROOF. First, note that

$$v^{-1}[(\mathcal{R}_{q,v}(\omega) - \mathcal{R}_{q,0}(\omega))Q] = -i\mathcal{R}_{q,v}(\omega)\Delta\mathcal{R}_{q,0}(\omega)Q.$$

Since $Q : H_\Lambda^s \rightarrow H_\Lambda^{s+N}$ for any N , and, by Lemma 7.6, $\mathcal{R}_{q,v} : H_\Lambda^s \rightarrow H_\Lambda^{s-\epsilon}$ with uniform bounds in v , $\mathcal{R}_{q,v}(\omega)\Delta\mathcal{R}_{q,0}(\omega)Q : H_\Lambda^s \rightarrow H_\Lambda^{s+N}$ is uniformly bounded in v for any N . In particular, Lemma 6.5 implies

$$v^{-1}\|(\mathcal{R}_{q,v}(\omega) - \mathcal{R}_{q,0}(\omega))Q\|_{\mathcal{L}^1(H_\Lambda^s \rightarrow H_\Lambda^s)} \leq C.$$

By [11, prop. B.29]

$$\begin{aligned} & \left| \det_{H_\Lambda^s} (I + A) - \det_{H_\Lambda^s} (I + B) \right| \\ & \leq \|A - B\|_{\mathcal{L}^1(H_\Lambda^s \rightarrow H_\Lambda^s)} e^{1 + \|A\|_{\mathcal{L}^1(H_\Lambda^s \rightarrow H_\Lambda^s)} + \|B\|_{\mathcal{L}^1(H_\Lambda^s \rightarrow H_\Lambda^s)}}. \end{aligned}$$

Therefore, since $\mathcal{R}_{q,v} Q : H_\Lambda^s \rightarrow H_\Lambda^N$ is uniformly bounded in v for any N , the lemma is proved. \square

Finally, we show that the eigenvalues of $P_{0,v}$ on H_Λ^s agree with those on L^2 . Together with Lemma 7.7, this will complete the proof of Theorem 1.1.

LEMMA 7.8. *Let $v > 0$ and G_0, G_1 satisfy (7.1). Suppose that $u \in H_{\Lambda_{G_1}}^s$ and*

$$(7.24) \quad (P_{0,v} - \omega)^K u = 0.$$

Then $u \in H_{\Lambda_{G_0}}^k$ for any k . In particular, the spectrum of $P_{0,v}$ on $L^2(\mathbb{T}^n)$ agrees with that on H_Λ^s .

PROOF. Let $G_\epsilon = (1 - \chi(\epsilon|\xi|))G_1 + \chi(\epsilon|\xi|)G_0$ with $\chi \in C_c^\infty(\mathbb{R})$, $\chi \equiv 1$ on $[-1/2, 1/2]$ and $\text{supp } \chi \subset (-1, 1)$. Note that G_ϵ satisfies (7.1) and $\lim_{\epsilon \rightarrow 0} G_\epsilon = G_0$ pointwise. In particular, $G_\epsilon = G_0$ on $2|\xi| < \epsilon^{-1}$ and $G_\epsilon = G_1$ on $|\xi| > \epsilon^{-1}$.

Suppose that $u \in H_{\Lambda_{G_1}}^s$ satisfies (7.24). Then,

$$\begin{aligned} & \|u\|_{H_{\Lambda_{G_\epsilon}}^s} \\ & = \|\langle \xi \rangle^s T_{\Lambda_{G_\epsilon}} u\|_{L^2(\Lambda_{G_\epsilon})} \\ & \leq \|1_{|\xi| \leq \epsilon^{-1}} \langle \xi \rangle^k T_{\Lambda_{G_\epsilon}} u\|_{L^2(\Lambda_{G_\epsilon})}^2 + \|1_{|\xi| > \epsilon^{-1}} \langle \xi \rangle^s T_{\Lambda_{G_1}} u\|_{L^2(\Lambda_{G_1})}^2 \\ & \leq \|1_{|\xi| \leq \epsilon^{-1}} \langle \xi \rangle^s T_{\Lambda_{G_\epsilon}} S_{\Lambda_{G_1}} T_{\Lambda_{G_1}} u\|_{L^2(\Lambda_{G_\epsilon})}^2 + \|1_{|\xi| > \epsilon^{-1}} \langle \xi \rangle^s T_{\Lambda_{G_1}} u\|_{L^2(\Lambda_{G_1})}^2 \\ & \leq C_\epsilon \|\langle \xi \rangle^s T_{\Lambda_{G_1}} u\|_{L^2(\Lambda_{G_1})}, \end{aligned}$$

where in the last line we use Lemma 4.9. In particular, $u \in H^s(\Lambda_{G_\epsilon})$ for each fixed $\epsilon > 0$.

Since $u \in H_{\Lambda_{G_\epsilon}}^s$, we can apply (7.17) together with (7.24) to obtain

$$\|u\|_{H_{\Lambda_{G_\epsilon}}^k} \leq C_{k,N,v} \|u\|_{H_{\Lambda_{G_\epsilon}}^{-N}}.$$

where $C_{s,N,v}$ does not depend on ϵ . Writing this on the FBI transform side, we have

$$\begin{aligned} & \|\langle \xi \rangle^k T_{\Lambda_{G_\epsilon}} u\|_{L^2(\Lambda_{G_\epsilon})} \\ & \leq C_{k,N,v} \|\langle \xi \rangle^{-N} T_{\Lambda_{G_\epsilon}} u\|_{L^2(\Lambda_{G_\epsilon})} \\ & \leq C_{k,N,v} (\|1_{|\xi| \leq M} T_{\Lambda_{G_\epsilon}} u\|_{L^2(\Lambda_{G_\epsilon})} + M^{k-N} \|\langle \xi \rangle^k T_{\Lambda_{G_\epsilon}} u\|_{L^2(\Lambda_{G_\epsilon})}). \end{aligned}$$

Now, choosing $M \geq (2C_{k,N,v})^{1/(N-k)}$ large enough, and moving the last term to the left-hand side, we obtain

$$\begin{aligned} \|\langle \xi \rangle^k T_{\Lambda_{G_\epsilon}} u\|_{L^2(\Lambda_{G_\epsilon})} &\leq C_{k,N,v} \|1_{|\xi| \leq M} T_{\Lambda_{G_\epsilon}} u\|_{L^2(\Lambda_{G_\epsilon})} \\ &= C_{k,N,v} \|1_{|\xi| \leq M} T_{\Lambda_{G_0}} u\|_{L^2(\Lambda_{G_0})} \\ &= C_{k,N,v} \|1_{|\xi| \leq M} T_{\Lambda_{G_0}} S_{\Lambda_{G_1}} T_{\Lambda_{G_1}} u\|_{L^2(\Lambda_{G_0})} \\ &\leq C_{k,s,N,v} \|\langle \xi \rangle^s T_{\Lambda_{G_1}} u\|_{L^2(\Lambda_{G_1})}, \end{aligned}$$

where in the last line we apply Lemma 4.9.

In particular, sending $\epsilon \rightarrow 0^+$, we have that

$$\limsup_{\epsilon \rightarrow 0} \|\langle \xi \rangle^k T_{\Lambda_{G_\epsilon}} u\|_{L^2(\Lambda_{G_\epsilon})} \leq C \|u\|_{H_{\Lambda_1}^s}.$$

Finally, by Fatou’s lemma together with the fact that $G_\epsilon \rightarrow G_0$, this implies

$$\|u\|_{H_{\Lambda_{G_0}}^k} = \|\langle \xi \rangle^k T_{\Lambda_{G_0}} u\|_{L^2(\Lambda_{G_0})} \leq C \|u\|_{H_{\Lambda_1}^s},$$

and in particular, $u \in H_{\Lambda_{G_0}}^k$ as claimed. □

This completes the proof of Theorem 1.1. □

7.6 Poles of the resolvent $\mathcal{R}_{0,0}(\omega)$ in the upper half-plane

Finally, we study the behavior of $\mathcal{R}_{0,0}(\omega) = \mathcal{R}(\omega)$ for $\text{Im } \omega \geq 0$ and complete the proof of Theorem 1.2. (That resolvent was defined in Lemma 7.4.)

LEMMA 7.9. *Suppose that*

$$\omega_1 \in \{|\text{Re } \omega| < \omega_0, \text{Im } \omega \geq 0\} \setminus \text{spec}_{\text{pp},L^2}(P).$$

Then, $\mathcal{R}(\omega)$ is analytic near ω_1 . Moreover, for $s \in \mathbb{R}$, $\text{Im } \omega > 0$, and $u \in L^2(\mathbb{T}^n) \cap H_{\Lambda}^s$, $\mathcal{R}(\omega)u = \mathcal{R}^{L^2}(\omega)u$ where \mathcal{R}^{L^2} denotes the L^2 resolvent for P .

Conversely, if $\omega_1 \in (-\omega_0, \omega_0) \cap \text{spec}_{\text{pp},L^2}(P)$, then ω_1 is a pole of $\mathcal{R}(\omega)$ and

$$\mathcal{R}(\omega) = A(\omega) + \frac{\Pi_{\omega_1}}{\omega - \omega_1},$$

where $A(\omega) : H_{\Lambda}^s \rightarrow H_{\Lambda}^s$ is analytic near ω_1 and Π_{ω_1} is the orthogonal projection onto the L^2 eigenspace of P at ω_1 .

PROOF. For $\text{Im } \omega > 0$, the spectral theorem shows that the resolvent of P ,

$$\mathcal{R}^{L^2}(\omega) := (P - \omega)^{-1} : L^2(\mathbb{T}^n) \rightarrow L^2(\mathbb{T}^n),$$

exists and is analytic. Also, Lemma 7.4 shows that $\mathcal{R}(\omega) = \mathcal{R}_{0,0}(\omega)$ is a meromorphic family of operators in $(-\omega_0, \omega_0) + i(-c_0\theta, \infty)$.

Lemma B.7 implies that for $u \in \mathcal{A}_\delta$ (defined in (4.2)) and $\text{Im } \omega \gg 1$,

$$R^{L^2}(\omega)u \in \mathcal{A}_\delta.$$

Since $P : H_\Lambda \rightarrow H_\Lambda$ and $\mathcal{A}_\delta \subset H_\Lambda$, $\mathcal{R}(\omega)(P - \omega)|_{\mathcal{A}_\delta} = \text{Id}_{\mathcal{A}_\delta}$. Therefore, for $u \in \mathcal{A}_\delta$,

$$\mathcal{R}^{L^2}(\omega)u = [\mathcal{R}(\omega)(P - \omega)]\mathcal{R}^{L^2}u = \mathcal{R}(\omega)[(P - \omega)\mathcal{R}^{L^2}(\omega)]u = \mathcal{R}(\omega)u.$$

Since \mathcal{A}_δ are dense in both $L^2(\mathbb{T}^n)$ and H_Λ^s ,

$$\mathcal{R}(\omega)u = \mathcal{R}^{L^2}(\omega)u, \quad \text{Im } \omega > 0, \quad \text{Re } \omega \in (-\omega_0, \omega_0), \quad u \in L^2(\mathbb{T}^n) \cap H_\Lambda^s.$$

This proves the first part of the lemma.

To prove the second part, let $\omega_1 \in (-\omega_0, \omega_0)$ and $\Pi_{\omega_1} : L^2(\mathbb{T}^n) \rightarrow L^2(\mathbb{T}^n)$ be the orthogonal projection onto the ω_1 eigenspace for P . (Note that Π_{ω_1} may be the zero operator if ω_1 is not an embedded eigenvalue for P .) By [12, lemma 3.2], the ω_1 eigenfunctions of P are smooth and hence $\tilde{P} = P + \Pi_{\omega_1}$ has the same symbol as P and no embedded eigenvalue at ω_1 . Moreover, we may choose $\epsilon > 0$ so small that ω_1 is the only embedded eigenvalue for P in $|\omega - \omega_1| < \epsilon$. Then, for $0 < |\omega - \omega_1| < \epsilon$, $\text{Im } \omega > 0$,

$$\begin{aligned} \mathcal{R}^{L^2}(\omega) &= (\tilde{P} - \omega)^{-1} + (P - \omega)^{-1}\Pi_{\omega_1}(\tilde{P} - \omega)^{-1} \\ &= (\tilde{P} - \omega)^{-1} + \frac{\Pi_{\omega_1}}{(\omega_1 - \omega)(1 + \omega_1 - \omega)}. \end{aligned}$$

Note that by [12, lemma 3.3] for $\omega \in (\omega_1 - \epsilon, \omega_1 + \epsilon)$, the limiting absorption resolvent $(\tilde{P} - \omega - i0)^{-1} : H^{1/2+0} \rightarrow H^{-1/2-0}$ exists.

The meromorphy of $\mathcal{R}(\omega) : H_\Lambda^s \rightarrow H_\Lambda^s$ (Lemma 7.4) gives

$$(7.25) \quad \mathcal{R}(\omega) = A(\omega) + \sum_{j=1}^K \frac{B_j}{(\omega - \omega_1)^j}$$

where $A : H_\Lambda^s \rightarrow H_\Lambda^s$ is holomorphic near ω_1 . Therefore, for $|\omega - \omega_1| < \epsilon$, $\text{Im } \omega > 0$, and $u \in L^2(\mathbb{T}^n) \cap H_\Lambda^s$,

$$\begin{aligned} (7.26) \quad &(\omega - \omega_1)^K A(\omega)u + \sum_{j=1}^K (\omega - \omega_1)^{K-j} B_j u \\ &= (\omega - \omega_1)^K \left[(\tilde{P} - \omega)^{-1} - \frac{\Pi_{\omega_1}}{1 + \omega_1 - \omega} \right] u - (\omega - \omega_1)^{K-1} \Pi_{\omega_1} u. \end{aligned}$$

Let $u \in \mathcal{A}_\delta \subset H_\Lambda^s \cap H^{(1/2)+0}$. Then, using (7.26) with $\omega = \omega_1 + ir$, we obtain

$$\begin{aligned} B_K u &= \lim_{r \rightarrow 0^+} \left[(ir)^K A(\omega_1 + ir)u + \sum_{j=1}^K (ir)^{K-j} B_j u \right] \\ &= \lim_{r \rightarrow 0^+} (ir)^K \left[(\tilde{P} - \omega_1 - ir)^{-1} - \frac{\Pi_{\omega_1}}{1 - ir} \right] u - (ir)^{K-1} \Pi_{\omega_1} u \\ &= \delta_{K1} \Pi_{\omega_1} u. \end{aligned}$$

Since \mathcal{A}_δ is dense in both $H^{(1/2)+0}$ and H_Λ^s , $B_K = \delta_{K1} \Pi_{\omega_1}$. In particular, we may write (7.25) with $K = 1$ and by the same argument obtain $B_1 = \Pi_{\omega_1}$. \square

Appendix A Review of Some Almost Analytic Constructions

Here we include some facts about almost analytic functions and manifolds. For an in-depth presentation see [25, secs. 1–3] and [37, chap. X].

A.1 Almost analytic manifolds

Let U be an open subset of \mathbb{C}^m and let $U_{\mathbb{R}} := U \cap \mathbb{R}^m$. We define an almost analytic function as follows:

$$f \in C^{aa}(U) \iff \partial_{\bar{z}} f(z) = \mathcal{O}_K(|\text{Im } z|^\infty), \quad z \in K \Subset U.$$

This definition is nontrivial only for $U_{\mathbb{R}} \neq \emptyset$. We write $f \sim 0$ in U if $f(z) = \mathcal{O}_K(|\text{Im } z|^\infty)$, $z \in K \Subset U \subset \mathbb{C}^m$. We note that (see [37, lemma X.2.2]) for $f \in C^\infty$ that implies $\partial^\alpha f \sim 0$ in U .

We also need the notion of an almost analytic manifold. Let $\Lambda \subset \mathbb{C}^m$ be a smooth manifold and $\Lambda_{\mathbb{R}} := \Lambda \cap \mathbb{R}^m$. We say that Λ is almost analytic if near any point $z_0 \in \Lambda_{\mathbb{R}}$, there exist a neighbourhood U of z_0 in \mathbb{C}^m and functions $f_1, \dots, f_k \in C^\infty(\mathbb{C}^m)$ such that

$$\begin{aligned} \Lambda \cap U &= \{z : f_j(z) = 0, 1 \leq j \leq k\}, \quad \partial_z f_j(z_0) \text{ are linearly independent,} \\ |\partial_{\bar{z}} f_j(z)| &= \mathcal{O}(|\text{Im } z|^\infty + \left| \sup_{1 \leq \ell \leq k} f_\ell(z) \right|^\infty); \end{aligned}$$

see [25, theorem 1.4].

A special case is given by

$$(A.1) \quad f_j(z) = z_j - h_j(z'), \quad z' := (z_{k+1}, \dots, z_n).$$

Equivalence of two almost analytic manifolds can be defined as follows (see [25, def. 1.6, prop. 1.7]): suppose $\Lambda_1 \cap \mathbb{R}^m = \Lambda_2 \cap \mathbb{R}^m$ and that Λ_k is defined by (A.1) with $h = h_k$, $k = 1, 2$, respectively. Then

Λ_1 and Λ_2 are equivalent as almost analytic submanifolds (denoted $\Lambda_1 \sim \Lambda_2$)

if and only if, on compact subsets,

$$(A.2) \quad \begin{aligned} |h_1(z') - h_2(z')| &= \mathcal{O}(|\operatorname{Im} h_1(z')|^\infty) \\ &\text{or, equivalently,} \\ |h_1(z') - h_2(z')| &= \mathcal{O}(|\operatorname{Im} z'|^\infty + |\operatorname{Im} h_1(z')|^\infty). \end{aligned}$$

We now consider *almost analytic vector fields*:

$$V = \sum_{j=1}^m a_j(z) \partial_{z_j}, \quad a_j \in C^{\text{aa}}(\mathbb{C}^n),$$

which we identify with *real vector fields* \widehat{V} such that for u holomorphic $\widehat{V}f = V$:

$$\begin{aligned} \widehat{V} &:= V + \bar{V} = 2 \operatorname{Re} V \\ &= \sum_{j=1}^m \operatorname{Re} a_j(z) (\partial_{z_j} + \partial_{\bar{z}_j}) + i \operatorname{Im} a_j(z) (\partial_{z_j} - \partial_{\bar{z}_j}) \\ &= \sum_{j=1}^m \operatorname{Re} a_j(z) \partial_{\operatorname{Re} z_j} + \operatorname{Im} a_j(z) \partial_{\operatorname{Im} z_j}. \end{aligned}$$

Example A.1. Suppose $M \subset \mathbb{C}^m$, $\dim_{\mathbb{R}} M = 2k$, is almost analytic. Then vector fields tangent to M are spanned by almost analytic vector fields, $V_j = a_j(z) \cdot \partial_z$, $\partial_{\bar{z}} a_j(z) = \mathcal{O}(|\operatorname{Im} z|^\infty)$, $z \in M$, $j = 1, \dots, k$. In fact, using [25, theorem 1.4, 3°] we can write M locally near any $z \in M \cap \mathbb{R}^m$ as $\{(z', h(z')) : z' \in \mathbb{C}^k\}$, $h = (h_{k+1}, \dots, h_m) : \mathbb{C}^k \rightarrow \mathbb{C}^{m-k}$, $\partial_{\bar{z}} h = \mathcal{O}(|\operatorname{Im} z'|^\infty + |\operatorname{Im} h(z')|^\infty)$. We then put

$$(A.3) \quad V_j = \partial_{z_j} + \sum_{\ell=k+1}^m \partial_{z_j} h_\ell(z') \partial_{z_\ell}.$$

The real vector fields \widehat{V}_j then span the vector fields tangent to M .

Following [25, 30] we define the (small complex time) flow of V as follows for $s \in \mathbb{C}$, $|s| \leq \delta$:

$$(A.4) \quad \Phi_s(z) := \exp s \widehat{V}(z).$$

The right-hand side is the flow out at time 1 of the real vector field $s \widehat{V}$, unless the coefficients in V are holomorphic $[\widehat{V}, i \widehat{V}] \neq 0$, which means that $\exp(s + t)V \neq \exp sV \exp tV$ for $s, t \in \mathbb{C}$. However, we still have $[i \widehat{V}, \widehat{V}] \sim 0$.

LEMMA A.2. *Suppose that $\Gamma \in \mathbb{C}^m$ is an embedded almost analytic submanifold of real dimension $2k$ and V is an almost analytic vector field. Assume that*

$$(A.5) \quad \widehat{V}, i \widehat{V} \text{ are linearly independent with span transversal to } \Gamma,$$

and that, in the notation of (A.4),

$$(A.6) \quad |\operatorname{Im} \Phi_t(z)| \geq |t|/C_K, \quad z \in K \Subset \Gamma.$$

Then for any $U \Subset \mathbb{C}^m$, there exists δ such that

$$\Lambda := \{\exp i\widehat{V}(\rho) : \rho \in \Gamma \cap U, |t| < \delta, t \in \mathbb{C}\}$$

is an almost analytic manifold, $\Lambda_{\mathbb{R}} = \Gamma_{\mathbb{R}}$ and $\dim_{\mathbb{R}} \Lambda = 2k + 2$.

We will use the following geometric lemma:

LEMMA A.3. Suppose $Z_j \in C^\infty(\mathbb{R}^m; T^*\mathbb{R}^m)$, $j = 1, \dots, J$, are smooth vector fields and, for $s \in \mathbb{R}^J$,

$$\langle s, Z \rangle := \sum_{j=1}^J s_j Z_j \in C^\infty(\mathbb{R}^m; T^*\mathbb{R}^m).$$

Then for $f \in C^\infty(\mathbb{R}^m)$

$$(A.7) \quad f(e^{\langle s, Z \rangle}(\rho)) = \sum_{p=1}^P \frac{1}{p!} (\langle s, Z \rangle)^p f(\rho) + \mathcal{O}_K(|s|^{P+1}), \quad \rho \in K \Subset \mathbb{R}^m.$$

while for $Y \in C^\infty(\mathbb{R}^m; T^*\mathbb{R}^m)$,

$$(A.8) \quad e_*^{\langle s, Z \rangle} Y(\rho) = \sum_{p=1}^P \frac{1}{p!} \operatorname{ad}_{\langle s, Z \rangle}^p Y(\rho) + \mathcal{O}_K(|s|^{P+1}), \quad \rho \in K \Subset \mathbb{R}^m.$$

For a proof, see [22, app. A]. We recall that $F_* Y(F(\rho)) := dF(\rho)Y(\rho)$.

PROOF OF LEMMA A.2. Let $\iota : \Gamma \hookrightarrow \mathbb{C}^m$ be the inclusion map. Then

$$\partial \exp(t_1 \widehat{V} + t_2 i\widehat{V}) \circ \iota(\rho) : T_{(0,\rho)}(\mathbb{R}_t^2 \times \Gamma) \rightarrow T_\rho \mathbb{C}^m$$

is given by $(T, X) \mapsto T_1 \widehat{V} + T_2 i\widehat{V} + \iota_* X$, which, thanks to our assumptions, is surjective onto a $(2k + 2)$ (real)-dimensional subspace of $T^*\mathbb{C}^m$. Hence, by the implicit function theorem, Λ is a $(2k + 2)$ dimensional embedded submanifold of \mathbb{C}^m .

To fix ideas we start with the simplest case of $\Gamma = \{0\} \subset \mathbb{C}^n$. In that case $\{\Lambda = \{\Phi_t(0) : t \in \mathbb{C}, |t| < \delta\}$, and from our assumption

$$|\operatorname{Im} \Phi_t(0)| \sim |t_1 \widehat{V} + t_2 i\widehat{V}| \sim |t|.$$

The tangent space is given by

$$T_{\Phi_t(0)} \Lambda = \{\partial_t \Phi_t(0)T + \partial_{\bar{t}} \Phi_t(0)\bar{T} : T \in \mathbb{C}\} \subset \mathbb{C}^2.$$

If we show that

$$(A.9) \quad \partial_{\bar{t}} \Phi_t(0) = \mathcal{O}(|t|^\infty),$$

then $d(T_{\Phi_t(0)} \Lambda, iT_{\Phi_t(0)} \Lambda) = \mathcal{O}(|t|^\infty)$ and almost analyticity of Λ follows from [25, theorem 1.4, 1°]. The estimate (A.9) will follow from showing that for any

holomorphic function f , $\partial_{t_1}^{\alpha_1} \partial_{t_2}^{\alpha_2} \partial_{\bar{t}} f(\Phi_t(0))|_{t=0} = 0$. But this follows from (A.7) and the fact that $[\widehat{V}, i\widehat{V}] \sim 0$ at 0. Indeed,

$$\begin{aligned}
 & \partial_{t_1}^{\alpha_1} \partial_{t_2}^{\alpha_2} \partial_{\bar{t}} f(\Phi_t(0))|_{t=0} \\
 &= \partial_{t_1}^{\alpha_1} \partial_{t_2}^{\alpha_2} \partial_{\bar{t}} \left(\sum_{k=0}^{\infty} \frac{1}{k!} (t_1 \widehat{V} + t_2 i \widehat{V})^k f(0) \right) |_{t=0} \\
 \text{(A.10)} \quad &= \partial_{t_1}^{\alpha_1} \partial_{t_2}^{\alpha_2} \left(\sum_{k=0}^{\infty} \frac{1}{k!} (t_1 \widehat{V} + t_2 i \widehat{V})^k (\widehat{V} + i i \widehat{V}) f(0) \right) |_{t=0} \\
 &= \widehat{V}^{\alpha_1} i \widehat{V}^{\alpha_2} (\widehat{V} + i i \widehat{V}) f(0) = \widehat{V}^{\alpha_1} i \widehat{V}^{\alpha_2} (V - V) f(0) = 0.
 \end{aligned}$$

The fact that \widehat{V} and $i\widehat{V}$ commute to infinite order at 0 was crucial in this calculation. Holomorphy of f was used to have $\widehat{W}f = Wf$.

We now move to the general case. For $z \in \Gamma$, $T_{\Phi_t(z)}\Lambda$ is spanned by

$$\text{(A.11)} \quad \partial_t \Phi_t(z)T + \partial_{\bar{t}} \Phi_t(z)\bar{T}, \quad T \in \mathbb{C}, \quad d\Phi_t(z)X, \quad X \in T_z\Gamma.$$

We can repeat the calculation (A.10) with 0 replaced by z to see that, using the assumption (A.6) and the fact that $\text{Im } \Phi_t(z) = \text{Im } z + \mathcal{O}(t)$,

$$\text{(A.12)} \quad \partial_{\bar{t}} \Phi(z) = \mathcal{O}(|t|^\infty + |\text{Im } z|^\infty) = \mathcal{O}(|\text{Im } \Phi_t(z)|^\infty).$$

To consider $d\Phi_t(z)X = (\Phi_t)_*Y(\Phi_t(z))$, we choose a vector field tangent to Γ , $Y, Y_c(z) = X$. We choose

$$\text{(A.13)} \quad Y_c = \widehat{W}_c, \quad W_c = \sum_{j=1}^k c_j V_j, \quad c \in \mathbb{C}^k,$$

a constant coefficient linear combination of vector fields (A.3). Then $d\Phi_t(z)X = (\Phi_t)_*Y_c(\Phi_t(z))$, and we want to show that

$$\text{(A.14)} \quad c \mapsto (\Phi_t)_*Y_c(\Phi_t(z)) \quad \text{is complex linear modulo errors } \mathcal{O}(|\text{Im } \Phi_t(z)|^\infty).$$

In view of (A.11), which shows that $d(T_{\Phi_t(z)}\Lambda, iT_{\Phi_t(z)}\Lambda) = \mathcal{O}(|\text{Im } \Phi_t(z)|^\infty)$ and from [25, theorem 1.4, 1°], we conclude that Λ is almost analytic.

To establish (A.14) we use (A.8) with $\langle s, X \rangle = s_1 \widehat{V} + s_2 i \widehat{V}$, $s_1 = \text{Re } t$, $s_2 = \text{Im } t$. Since $[\widehat{V}, i\widehat{V}] \sim 0$ and $\widehat{V} \sim i\widehat{V}/i$ at $\text{Im } w = 0$, we see that

$$\text{(A.15)} \quad (\Phi_t)_*Y_c(w) = \sum_{p=0}^{\infty} \frac{t^p}{p!} \text{ad}_{\widehat{V}}^p W_c(w) + \mathcal{O}(|t|^{K+1} + |\text{Im } w|^\infty).$$

Because of the form of W_c (see (A.3) and (A.13))

$$\text{ad}_{\widehat{V}}^p W_c(w) = \widehat{\text{ad}}_{\widehat{V}}^p W_c(w) + \mathcal{O}(|\text{Im } w'|^\infty + |\text{Im } h(w')|^\infty)$$

and

$$c \mapsto \text{ad}_{\widehat{V}}^p W_c(w) \quad \text{is complex linear.}$$

Since $w = \Phi_t(z)$, $z \in \Gamma$,

$$\begin{aligned} |\operatorname{Im} w'| + |\operatorname{Im} h(w')| &= \mathcal{O}(|\operatorname{Im} z'| + |\operatorname{Im} h(z')| + |t|) \\ &= \mathcal{O}(|\operatorname{Im} z| + |t|) = \mathcal{O}(|\operatorname{Im} w| + |t|) = \mathcal{O}(|\operatorname{Im} w|), \end{aligned}$$

since $|\operatorname{Im} w| = |\operatorname{Im} \Phi_t(z)| \geq |t|/C$. Combining these estimates with (A.15) gives (A.14). \square

A.2 Almost analytic generating functions

We now recall how to obtain generating functions for almost analytic strictly positive Lagrangian manifolds. We recall that an almost analytic submanifold of $T^*\mathbb{C}^n$, Λ , is Lagrangian if

$$(\omega_{\mathbb{C}})|_{\Lambda} \sim 0, \quad \omega_{\mathbb{C}} := \sum_{j=1}^n d\zeta_j \wedge dz_j.$$

In addition, we say that Λ is *strictly positive* ([25, def. 3.3]) if $\Lambda_{\mathbb{R}}$ is a submanifold of $T^*\mathbb{R}^m$ and for all $\rho \in \Lambda_{\mathbb{R}}$,

$$(A.16) \quad \frac{1}{i}\sigma(V, \bar{V})(\rho) > 0 \quad \text{for all } V \in T_{\rho}\Lambda \setminus (T_{\rho}\Lambda_{\mathbb{R}})^{\mathbb{C}}.$$

LEMMA A.4. *Suppose that Λ is a strictly positive almost analytic Lagrangian submanifold of $\operatorname{nbhd}_{T^*\mathbb{C}^m}(0)$ given by*

$$\{(z, \zeta(z)) \mid z \in \operatorname{nbhd}_{\mathbb{C}^m}(0)\}.$$

Then there is $C > 0$ such that for $x \in \operatorname{nbhd}_{\mathbb{R}^m}(0)$,

$$(A.17) \quad \frac{1}{C}d(x, \pi(\Lambda_{\mathbb{R}})) \leq |\operatorname{Im} \zeta(x)| \leq Cd(x, \pi(\Lambda_{\mathbb{R}})).$$

PROOF. Since $\Lambda_{\mathbb{R}}$ is a submanifold of $T^*\mathbb{R}^m$, we may choose real coordinates on \mathbb{R}^m such that (near 0)

$$\Lambda_{\mathbb{R}} = \{(x', 0, \zeta(x', 0)) \mid (x', x'') \in \mathbb{R}^k \times \mathbb{R}^{m-k}\}.$$

Then, with $\rho = (x', 0, \zeta(x', 0))$, $(T_{\rho}\Lambda_{\mathbb{R}})^{\mathbb{C}} = \{(\delta_{z'}, 0, \partial_{x'}\zeta(x', 0)\delta_{z'}) \mid \delta_{z'} \in \mathbb{C}^k\}$, and it follows that for all $\delta_{x''} \in \mathbb{R}^{m-k}$,

$$(0, \delta_{x''}, \partial_{x''}\zeta(x', 0)\delta_{x''}) \in T_{\rho}\Lambda \setminus (T_{\rho}\Lambda_{\mathbb{R}})^{\mathbb{C}}.$$

Strict positivity of Λ then implies that

$$\frac{1}{i}\sigma((0, \delta_{x''}, \partial_{x''}\zeta(x', 0)\delta_{x''}), \overline{(0, \delta_{x''}, \partial_{x''}\zeta(x', 0)\delta_{x''}})) = 2\langle \operatorname{Im} \partial_{x''}\zeta \delta_{x''}, \delta_{x''} \rangle > 0.$$

Since in our coordinates,

$$\frac{1}{C}|x''| \leq d(x, \pi(\Lambda_{\mathbb{R}})) \leq C|x''|,$$

(A.17) follows. \square

With this lemma in place we can find generating functions in the almost analytic setting:

LEMMA A.5. *Suppose that Λ is a strictly positive almost analytic Lagrangian submanifold of $\text{nbhd}_{T^*\mathbb{C}^m}(0)$ and that $\pi_* : T_{(0,0)}\Lambda \rightarrow T_0\mathbb{C}^m$ is onto. Then there exists $\Psi \in C^\infty(\text{nbhd}_{\mathbb{C}^m}(0))$ satisfying*

$$(A.18) \quad \partial_{\bar{z}}\Psi = \mathcal{O}(|\text{Im } z|^\infty + |\text{Im } \Psi(z)|^\infty)$$

such that, as almost analytic manifolds,

$$(A.19) \quad \Lambda \sim \{(z, \Psi_z(z)) : |z| < \epsilon\}, \quad \Psi_z(0) = 0.$$

PROOF. Since Λ is an almost analytic Lagrangian, we have $\sigma|_\Lambda \sim 0$ (vanishes to infinite order at $\Lambda_{\mathbb{R}}$) while the projection property shows that, near $z = 0$, $\Lambda = \{(z, \zeta(z)) : z \in \mathbb{C}^m\}$, $\zeta(0) = 0$. Hence $d(\zeta(z)dz) \sim 0$ and (see [25, theorem 1.4, 3°])

$$\partial_{\bar{z}}\zeta(z) = \mathcal{O}(|\text{Im } z|^\infty + |\text{Im } \zeta(z)|^\infty).$$

We note that for $z = x \in \mathbb{R}^n$, Lemma A.4 together with the strict positivity at $\Lambda_{\mathbb{R}} = \{(0, 0)\}$ shows that

$$(A.20) \quad |x''|/C \leq |\text{Im } \zeta(x)| \leq C|x''|, \quad x \in \mathbb{R}^n, \quad |x| < \epsilon.$$

where $\pi(\Lambda_{\mathbb{R}})$ is given by $\{|x''| = 0\}$. We now see that

$$0 \sim \sigma|_\Lambda = \sum_{j=1}^n \partial_z \zeta_j(z) \wedge dz_j + \mathcal{O}(|\text{Im } z|^\infty + |\text{Im } \zeta(z)|^\infty)_{C^\infty(\mathbb{C}^n; \wedge^{2n} \mathbb{C}^n)},$$

and in view of (A.20)

$$\partial_{z_k} \zeta_j(x) - \partial_{z_j} \zeta_k(x) = \mathcal{O}(|x''|^\infty), \quad x \in \mathbb{R}^n, \quad |x| < \epsilon.$$

For $x \in \mathbb{R}^n$, define Ψ by a simple version of the Poincaré lemma: $\Psi(x) = \int_0^1 \zeta(tx) \cdot x dt$. Then

$$\begin{aligned} \partial_{x_j} \Psi(x) &= \int_0^1 \left(\sum_{k=1}^n t z_k \partial_{x_j} \zeta_k(tx) + \zeta_j(tx) \right) dt \\ (A.21) \quad &= \int_0^1 \left(\sum_{k=1}^n t z_k \partial_{x_k} \zeta_j(tx) + \zeta_j(tx) \right) dt + \mathcal{O}(|x''|^\infty) \\ &= \int_0^1 \partial_t (t \zeta_j(tx)) dt + \mathcal{O}(|x''|^\infty) = \zeta_j(x) + \mathcal{O}(|\text{Im } \zeta(x)|^\infty); \end{aligned}$$

in the last argument we used (A.20) again. We now define $\Psi(z)$ as an almost analytic extension of Ψ . From [25, prop. 1.7(ii)] we obtain (A.19). \square

A.3 Integration of almost analytic vector fields

Here we show how to solve transport equations arising in Section 5.2. For clarity we present a simpler case (see also [15, sec. 5.2.2]). Thus we assume that V is an almost analytic vector field on \mathbb{C}^n ($w = (w_1, w') \in \mathbb{C}^n$, $w_1 \in \mathbb{C}$, $w' \in \mathbb{C}^{n-1}$) satisfying

$$\begin{aligned} |\operatorname{Im} \exp(\widehat{tV})(0, w')| &\geq |t|/C, \quad w' \in B_{\mathbb{C}^{n-1}}(0, \epsilon), \\ t \in \mathbb{C}, \quad |t| < \epsilon, \quad dw_1(V) &\neq 0. \end{aligned}$$

Then $(t, w') \mapsto \exp(\widehat{tV})(0, w') =: z(t, w')$ is a diffeomorphism for ϵ small enough. We solve

$$(A.22) \quad Va \sim b, \quad a(0, w') = a_0(w'),$$

by putting

$$\begin{aligned} a(z) &:= a_1(z) + a_2(z), \quad a_1(z) = a_0(w'(z)), \\ a_2(z(t, w')) &:= \int_0^1 tb(z(ts, w'))ds. \end{aligned}$$

We calculate the action of V on a_1 using almost analyticity of b , the properties of $z(t, w')$ and (A.7):

$$\begin{aligned} (Va_2)(z(t, w')) &\sim \int_0^1 \sum_{k=0}^{\infty} \frac{s^k}{k!} V \widehat{tV}^k tb(z(0, w'))ds + \mathcal{O}(|t|^\infty) \\ &\sim \int_0^1 \sum_{k=0}^{\infty} \frac{s^k}{k!} \widehat{tV}^{k+1} b(z(0, w'))ds + \mathcal{O}(|t|^\infty) + \mathcal{O}(|\operatorname{Im} z(0, w')|^\infty) \\ &\sim \sum_{k=0}^{\infty} \frac{1}{(k+1)!} \widehat{tV}^{k+1} b(z(0, w')) + \mathcal{O}(|t|^\infty) + \mathcal{O}(|\operatorname{Im} z(0, w')|^\infty) \\ &= b(z(t, w')) + \mathcal{O}(|t|^\infty + |\operatorname{Im} z(0, w')|^\infty) \\ &= b(z(t, w')) + \mathcal{O}(|\operatorname{Im} z|^\infty). \end{aligned}$$

Similarly, $Va_1 \sim 0$ and we obtain (A.22).

Appendix B Physical Deformations and Numerical Results

The purpose of this appendix is to illustrate our results by numerical examples. We have not yet implemented the general theory numerically. However, in some circumstances, it is enough to consider physical deformations of \mathbb{T}^n rather than the more complicated phase space deformations. In particular, this is possible when

there exists $G(x, \xi)$ linear in ξ satisfying $H_p G > 0$ on $\{p = 0\} \cap \{|\xi| \geq C\}$. This type of deformation is analogous to the method of *complex scaling* (rediscovered as the method of *perfectly matched layers* in numerical analysis)—see [11, secs. 4.5, 4.7] for an introduction and references.

B.1 Deformations of analytic pseudodifferential operators

For $u \in \mathcal{D}'(\mathbb{T}^n)$ we extend u to be $2\pi\mathbb{Z}^n$ periodic on \mathbb{R}^n . We consider (B.1)

$$(Pu)(x) = \lim_{\epsilon \rightarrow 0^+} \lim_{\delta \rightarrow 0^+} \frac{1}{(2\pi)^n} \int e^{i\langle x-y, \xi \rangle - \epsilon|\xi|^2 - \delta|x-y|^2} p(x, \xi) u(y) dy d\xi,$$

$$|p(z, \zeta)| \leq C \langle \text{Re } \zeta \rangle^m, \quad |\text{Im } z| \leq a, \quad |\text{Im } \zeta| \leq b \langle \text{Re } \zeta \rangle,$$

$$p(x, \xi) = p(x + 2\pi k, \xi), \quad k \in \mathbb{Z}^n.$$

and $G(x, \xi) \in S^1(T^*\mathbb{T}^n)$ such that

$$G(x, \xi) = \langle G_0(x), \xi \rangle, \quad G_0 \in C^\infty(\mathbb{T}^n; \mathbb{R}^n).$$

Remark B.1. Observe that (B.1) agrees with the definition of the standard left quantization of the symbol p as in (6.2).

We consider the complex deformed operator, P_θ , defined by the property that when u is analytic in a sufficiently large neighbourhood of \mathbb{T}^n (or simply for u being a trigonometric polynomial),

$$P_\theta(u|_{\Gamma_\theta}) = (Pu)|_{\Gamma_\theta}, \quad \Gamma_\theta := \{x + i\theta G_0(x) \mid x \in \mathbb{T}^n\}.$$

We start by deriving a formula for the kernel of P_θ :

LEMMA B.2. *Suppose $u \in C^\omega(\mathbb{T}^n)$ extends analytically to $|\text{Im } z| < a$. Then, for $|\text{Im } z| < a$, the limit*

$$v(z) = \lim_{\epsilon \rightarrow 0^+} \lim_{\delta \rightarrow 0^+} \frac{1}{(2\pi)^n} \int e^{i\langle z-y, \xi \rangle - \epsilon|\xi|^2 - \delta|z-y|^2} p(z, \xi) u(y) dy d\xi$$

exists, and moreover, $v(z)$ is the analytic continuation of Pu .

PROOF. For each fixed ϵ, δ , the resulting function of z is manifestly analytic in a neighbourhood of \mathbb{T}^n (or \mathbb{R}^n if we think of periodic functions). Therefore, in order to see that v itself is analytic, we need only show that the limit exists and the convergence is uniform on compact subsets of $|\text{Im } z| < a$. For this, we deform the contour in y to

$$\Gamma(z) : y \mapsto y + i \text{Im } z,$$

so that we have

$$v(z) = \lim_{\epsilon \rightarrow 0^+} \lim_{\delta \rightarrow 0^+} \frac{1}{(2\pi)^n} \iint_{\Gamma(z)} e^{i\langle z-y, \xi \rangle - \epsilon|\xi|^2 - \delta|z-y|^2} p(z, \xi) u(y) dy d\xi$$

$$= \lim_{\epsilon \rightarrow 0^+} \lim_{\delta \rightarrow 0^+} \frac{1}{(2\pi)^n} \int e^{i\langle \text{Re } z-y, \xi \rangle - \epsilon|\xi|^2 - \delta|\text{Re } z-y|^2} p(z, \xi) u(y + i \text{Im } z) dy d\xi.$$

This contour deformation is justified for each fixed δ since the integrand is super exponentially decaying in y . Now, integrating by parts in ξ using

$$L_\epsilon = \frac{1 + \langle \operatorname{Re} z - y - 2i\epsilon, D_\xi \rangle}{1 + |\operatorname{Re} z - y|^2 + 4\epsilon|\xi|^2},$$

we have for any $N > 0$

$$v(z) = \lim_{\epsilon \rightarrow 0^+} \lim_{\delta \rightarrow 0^+} \frac{1}{(2\pi)^n} \int e^{i\langle \operatorname{Re} z - y, \xi \rangle - \epsilon|\xi|^2 - \delta|\operatorname{Re} z - y|^2} \cdot (L_\epsilon^t)^N (p(z, \xi)) u(y + i \operatorname{Im} z) dy d\xi.$$

In particular, since $|\partial_\xi^\alpha p| \leq C_\alpha \langle \operatorname{Re} \xi \rangle^{m-|\alpha|}$, the integrand is bounded by

$$C_N \langle \operatorname{Re} z - y \rangle^{-N} \langle \xi \rangle^m e^{-\epsilon|\xi|^2}$$

uniformly in $\delta > 0$ and compact subsets of $|\operatorname{Im} z| < a$. Hence, for N large enough, the limit in δ exists and is uniform in compact subsets of $|\operatorname{Im} z| < a$. It is given by

$$v(z) = \lim_{\epsilon \rightarrow 0^+} \frac{1}{(2\pi)^n} \int e^{i\langle \operatorname{Re} z - y, \xi \rangle - \epsilon|\xi|^2} (L_\epsilon^t)^N [p(z, \xi)] u(y + i \operatorname{Im} z) dy d\xi.$$

Defining $B := (1 + \langle \xi, D_y \rangle) / (1 + |\xi|^2)$ we have, for any $N > 0$,

$$v(z) = \lim_{\epsilon \rightarrow 0^+} \frac{1}{(2\pi)^n} \int_\Gamma e^{i\langle \operatorname{Re} z - y, \xi \rangle - \epsilon|\xi|^2} (B^t)^N \cdot [(L_\epsilon^t)^N [p(z, \xi)] u(y + i \operatorname{Im} z)] dy d\xi.$$

Since $|\partial_y^\alpha u(y + i \operatorname{Im} z)| \leq C_\alpha$ and $|\partial_x^\alpha \partial_\xi^\beta p(x, \xi)| \leq C_{\alpha\beta} \langle \operatorname{Re} \xi \rangle^{m-|\beta|}$, the integrand is bounded by

$$C_N \langle \operatorname{Re} z - y \rangle^{-N} \langle \xi \rangle^{m-N},$$

uniformly in $\epsilon > 0$ and compact subsets of $|\operatorname{Im} z| < a$. In particular, the limit in ϵ exists, is uniform in compact subsets of $|\operatorname{Im} z| < a$, and is given by

$$v(z) = \frac{1}{(2\pi)^n} \int_\Gamma e^{i\langle \operatorname{Re} z - y, \xi \rangle} (B^t)^N [(L_0^t)^N [p(z, \xi)] u(y + i \operatorname{Im} z)] dy d\xi.$$

Thus, we see that v is analytic on $|\operatorname{Im} z| < a$ and agrees with Pu on $\operatorname{Im} z = 0$. In particular, by uniqueness of analytic continuation, $v(z) = (Pu)(z)$. \square

We now move to the representation for the Schwartz kernel of P_θ .

LEMMA B.3. *The kernel of P_θ acting on $2\pi \mathbb{Z}^n$ periodic functions on \mathbb{R}^n is given by*

$$(B.2) \quad K_\theta(x, y) = \frac{1}{(2\pi)^n} \int e^{i\langle x - y, \xi \rangle} p(\gamma_\theta(x), (e_\theta(x, y)^{-1})^t \xi) \frac{\det(\partial_y \gamma_\theta(y))}{\det e_\theta(x, y)} d\xi,$$

where the integral is interpreted as an oscillatory integral,

$$\gamma_\theta := x + i\theta G_0(x),$$

and $e_\theta(x, y)$ satisfies

$$e_\theta(x, y)(x - y) = \gamma_\theta(x) - \gamma_\theta(y).$$

In particular, $P_\theta \in \Psi^m$ and its principal symbol is given by

$$\sigma(P_\theta) = p(\gamma_\theta(x), (\partial\gamma_\theta(x)^{-1})^t \xi).$$

Remark B.4. Note that the symbol in (B.2) is not $2\pi\mathbb{Z}^n$ periodic in x . However, it is of the form $a(x, x - y, \xi)$ where a is $2\pi\mathbb{Z}^n$ periodic in the first variable. Therefore, it still maps periodic functions to periodic functions.

PROOF. By Lemma B.2, for u analytic on $|\operatorname{Im} z| < a$ and θ small enough

$$\begin{aligned} & (P_\theta u|_{\Gamma_\theta})(\gamma_\theta(x)) \\ &= \lim_{\epsilon \rightarrow 0^+} \lim_{\delta \rightarrow 0^+} \frac{1}{(2\pi)^n} \int e^{i(\gamma_\theta(x)-y, \xi) - \epsilon|\xi|^2 - \delta(\gamma_\theta(x)-y)^2} p(\gamma_\theta(x), \xi) u(y) dy d\xi. \end{aligned}$$

Now, since for each fixed $\delta > 0$, the integrand is super exponential decaying in y , we may deform the contour in y to Γ_θ , to obtain

$$\begin{aligned} & (P_\theta u|_{\Gamma_\theta})(\gamma_\theta(x)) \\ \text{(B.3)} \quad &= \lim_{\epsilon \rightarrow 0^+} \lim_{\delta \rightarrow 0^+} \frac{1}{(2\pi)^n} \int e^{i(\gamma_\theta(x)-\gamma_\theta(y), \xi) - \epsilon|\xi|^2 - \delta(\gamma_\theta(x)-\gamma_\theta(y))^2} \\ & \quad \cdot p(\gamma_\theta(x), \xi) \det(\partial_y \gamma_\theta(y)) u(\gamma_\theta(y)) dy d\xi. \end{aligned}$$

Next, using that for each fixed $\epsilon > 0$, the integrand is super exponentially decaying in ξ and that, with $e_\theta(x, y)(x - y) = \gamma_\theta(x) - \gamma_\theta(y)$, we have $e_\theta = \operatorname{Id} + O(\theta(x - y)^{-1})$, we can deform the contour in ξ to $\Gamma_1 = \xi \mapsto (e_\theta(x, y)^t)^{-1} \xi$, to obtain

$$\begin{aligned} & (P_\theta u|_{\Gamma_\theta})(\gamma_\theta(x)) \\ &= \lim_{\epsilon \rightarrow 0^+} \lim_{\delta \rightarrow 0^+} \frac{1}{(2\pi)^n} \int e^{i(x-y, \xi) - \epsilon((e_\theta^t)^{-1} \xi)^2 - \delta(\gamma_\theta(x)-\gamma_\theta(y))^2} \\ & \quad \cdot p(\gamma_\theta(x), (e_\theta(x, y)^{-1})^t \xi) \frac{\det(\partial_y \gamma_\theta(y))}{\det e_\theta(x, y)} u(\gamma_\theta(y)) dy d\xi. \end{aligned}$$

Now, integrating by parts as in the proof of Lemma B.2 results in the formula (B.2).

To prove the final claim, let $\chi \in C_c^\infty(-1, 1)$ with $\chi \equiv 1$ near 0. Then, for any $\delta > 0$, we have by (B.2) that with

$$P'_\theta = \frac{1}{(2\pi)^n} \int e^{i(x-y, \xi)} p(\gamma_\theta(x), (e_\theta(x, y)^{-1})^t \xi) \frac{\det(\partial_y \gamma_\theta(y))}{\det e_\theta(x, y)} \chi(\delta^{-1}|x - y|) d\xi,$$

the error $P_\theta - P'_\theta$ is smoothing and maps periodic functions to periodic functions. In particular, P_θ is a pseudodifferential operator on \mathbb{T}^n with symbol

$$\sigma(P_\theta) = p(\gamma_\theta(x), (\partial\gamma_\theta(x)^{-1})^t \xi),$$

proving the last claim in the lemma. □

B.2 The resolvent of the deformed operator

We now consider the setting of Theorem 1.1. Namely, we assume that P is a self-adjoint zeroth-order pseudodifferential operator and study the properties of P_θ .

PROPOSITION B.5. *Suppose that $P \in \Psi^0$ is self-adjoint and satisfies (B.1) and that $G = \langle G_0, \xi \rangle \in S^1$ has $H_p G > 0$ on $\{p = 0\} \cap \{|\xi| > C\}$. Then there are $\omega_0, \theta_0 > 0$ such that for $0 < \theta < \theta_0$, $\omega \in (-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$, and all $s \in \mathbb{R}$,*

$$\mathcal{R}_\theta(\omega) := (P_\theta - \omega)^{-1} : H^s(\mathbb{T}^n) \rightarrow H^s(\mathbb{T}^n)$$

is meromorphic with finite rank poles.

PROOF. First, note that there is $\omega_1 > 0$ such that

$$H_p G > c > 0 \quad \text{on } \{|p| < \omega_0\} \cap \{|\xi| > C\}.$$

We compute

$$\begin{aligned} \sigma(P_\theta)(x, \xi) &= p(x, \xi) + i\theta \langle \partial_x p(x, \xi), G_0(x) \rangle \\ &\quad + \langle \partial_\xi p(x, \xi), (\partial \gamma_\theta(x)^{-1})^t - \text{Id} \rangle \xi + O(\theta^2) \\ &= p(x, \xi) + i\theta (\langle \partial_x p(x, \xi), G_0(x) \rangle - \langle \partial_\xi p(x, \xi), (\partial_x G_0)^t(x) \rangle \xi) + O(\theta^2) \\ &= p(x, \xi) - i\theta H_p G(x, \xi) + O(\theta^2). \end{aligned}$$

Therefore, for $\theta > 0$ small enough, and $\omega_0 = \min(c, \omega_1)$, $P_\theta - \omega$ is elliptic when $\text{Im } \omega \geq -\omega_0\theta$, $|\text{Re } \omega| < \omega_0$.

In particular, $P_\theta - \omega : H^s \rightarrow H^s$ is Fredholm for $\omega \in [-\omega_0, \omega_0] + i(-\omega_0\theta, \infty)$. Moreover, since $P_\theta : H^s \rightarrow H^s$ is bounded, if $\text{Im } \omega \gg 1$, $P_\theta - \omega$ is invertible by Neumann series and hence $P_\theta - \omega$ has index 0. By the meromorphic Fredholm theorem [11, theorem C.9], its inverse $\mathcal{R}_\theta(\omega) = (P_\theta - \omega)^{-1} : H^s \rightarrow H^s$ is a meromorphic family of operators with finite rank poles for $\omega \in (-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$. \square

PROPOSITION B.6. *Let P and G be as in Proposition B.5. There are $\theta_0, \omega_0 > 0$ such that for $0 < \theta < \theta_0$, the poles of $\mathcal{R}_\theta(\omega)$ for $\omega \in (-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$ agree with multiplicity with those of $\mathcal{R}^{H^\Lambda}(\omega)$ where $\mathcal{R}^{H^\Lambda}(\omega)$ is the resolvent of P on H_Λ^s from Lemma 7.4.*

We will need the following lemma.

LEMMA B.7. *Suppose that P is as in (B.1) and $\mathcal{R}^{L^2}(\omega)$ is the resolvent of P on L^2 . Then, there are $C > 0$ and $\delta > 0$ such that for $\text{Im } \omega \geq C$,*

$$\mathcal{R}^{L^2}(\omega) : \mathcal{A}_\delta \rightarrow \mathcal{A}_\delta.$$

PROOF. We start by showing that $P : \mathcal{A}_\delta \rightarrow \mathcal{A}_\delta$ is bounded. For this, note that for $j \in \mathbb{Z}^n$,

$$\widehat{Pu}(j) = \sum_{k \in \mathbb{Z}^n} \widehat{u}(k) \widehat{p}(j - k, k)$$

where $\widehat{p}(j, \xi)$ denotes the Fourier series for $p(x, \xi)$ in the x -variable. Note that by (B.1), there is $C > 0$ such that

$$|\widehat{p}(k, \xi)| \leq C \langle \xi \rangle^m e^{-a|k|}.$$

Therefore, for $\delta < \frac{a}{2}$,

$$\begin{aligned} \|Pu\|_{\mathcal{A}_\delta}^2 &= \sum_j \left| \sum_k \widehat{u}(k) \widehat{p}(j - k, k) \right|^2 e^{4|j|\delta} \\ &\leq \sum_j \left(\sum_k |\widehat{u}(k)|^2 e^{4|k|\delta} \right) \left(\sum_k |\widehat{p}(j - k, k)|^2 e^{4(|j|-|k|)\delta} \right) \\ &= \|u\|_{\mathcal{A}_\delta}^2 \left(\sum_{k,j} e^{-2a|j-k|} e^{4(|j|-|k|)\delta} \right) \leq C \|u\|_{\mathcal{A}_\delta}^2 \left(\sum_j e^{(4\delta - 16\frac{a\delta}{4\delta+2a})|j|} \right) \\ &\leq C \|u\|_{\mathcal{A}_\delta}^2. \end{aligned}$$

Since $\text{Im } \omega > \|P\|_{\mathcal{A}_\delta \rightarrow \mathcal{A}_\delta}$, $\mathcal{R}^{L^2}(\omega) = -\sum_{k=0}^\infty \omega^{-k-1} P^k$, the proof is complete. \square

PROOF OF PROPOSITION B.6. Let ω_0 be the minimum of ω_0 from Proposition B.5 and Lemma 7.4, and suppose that $\omega \in (\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$ with $\text{Im } \omega \gg 1$. Then, $\mathcal{R}_\theta(\omega) : H^s(\Gamma_\theta) \rightarrow H^s(\Gamma_\theta)$ and $\mathcal{R}^{L^2}(\omega) : \mathcal{A}_\delta \rightarrow \mathcal{A}_\delta$.

Let $u \in \mathcal{A}_\delta$. Then we have $\mathcal{R}^{L^2}(\omega)u \in \mathcal{A}_\delta$ and

$$(P_\theta - \omega)([\mathcal{R}^{L^2}(\omega)u]|_{\Gamma_\theta}) = ((P - \omega)\mathcal{R}^{L^2}(\omega)u)|_{\Gamma_\theta} = u|_{\Gamma_\theta}.$$

In particular, $[\mathcal{R}^{L^2}(\omega)u]|_{\Gamma_\theta} = \mathcal{R}_\theta(\omega)(u|_{\Gamma_\theta})$.

For $u, v \in \mathcal{A}_\delta$, and $\text{Im } \omega \gg 1$,

$$\langle \mathcal{R}_\theta(\omega)u|_{\Gamma_\theta}, v|_{\Gamma_\theta} \rangle_{L^2(\Gamma_\theta)} = \langle [\mathcal{R}^{L^2}(\omega)u|_{\mathbb{T}^n}]_{\Gamma_\theta}, v|_{\Gamma_\theta} \rangle_{L^2(\Gamma_\theta)}.$$

By Lemma 7.9, when $\text{Im } \omega > 0$ and $u \in \mathcal{A}_\delta$, we have $\mathcal{R}^{H^\Lambda}(\omega)u = \mathcal{R}^{L^2}(\omega)u$. Then, deforming the contour of integration in the inner product to \mathbb{T}^n ,

$$\begin{aligned} &\langle \mathcal{R}_\theta(\omega)u|_{\Gamma_\theta}, v|_{\Gamma_\theta} \rangle_{L^2(\Gamma_\theta)} \\ &= \langle [\mathcal{R}^{L^2}(\omega)u|_{\mathbb{T}^n}]|_{\Gamma_\theta}, v|_{\Gamma_\theta} \rangle_{L^2(\Gamma_\theta)} \\ &= \langle \mathcal{R}^{L^2}(\omega)u, v \rangle_{L^2(\mathbb{T}^n)} \\ &= \langle \mathcal{R}^{H^\Lambda}(\omega)u, v \rangle_{L^2(\mathbb{T}^n)} = \langle \mathcal{R}^{H^\Lambda}(\omega)u, v \rangle_{\mathcal{A}_{-\delta}(\mathbb{T}^n), \mathcal{A}_\delta(\mathbb{T}^n)}. \end{aligned}$$

Since $\mathcal{A}_\delta \subset H_\Lambda^0 \subset \mathcal{A}_{-\delta}$, both sides of this equality continue meromorphically from $\text{Im } \omega \gg 1$ to $\omega \in (-\omega_0, \omega_0) + i(-\omega_0\theta, \infty)$, and the equality continues




FIGURE B.1. We display the eigenvalues of P_θ acting on $e^{inx_2}L^2_{x_1}$ for three different values of θ and $-20 \leq n \leq 20$. P is chosen as in (B.4) with $V_a = 0$ and $V_m = ((1 - \xi^2) + e\xi^2)e^{-\xi^2}$. These choices guarantee the existence of an embedded eigenvalue at 0 [36, example 1]. Note that once the eigenvalues emerge from the continuous spectrum, they are independent of the choice of θ .

to hold for ω in this set. Finally, since \mathcal{A}_δ is dense in H^s_Λ and $\mathcal{A}_\delta|_{\Gamma_\theta}$ is dense in $L^2(\Gamma_\theta)$, this equality implies that the poles of \mathcal{R}_θ and \mathcal{R}^{H_Λ} coincide with an agreement of multiplicities. \square

B.3 Numerical examples and discretization

In our numerical study, we consider operators of the form

$$(B.4) \quad P = \langle D \rangle^{-1} D_{x_2} + \sin(x_1)(\text{Id} - V_m(D_{x_1})) + (\text{Id} - V_m(D_{x_1})) \sin(x_1) + V_a(D_{x_1}),$$

with $V_\bullet(\xi_1)$, $\bullet = a, m$, satisfying

$$(B.5) \quad |V_\bullet(\xi_1)| \leq C e^{-c|\text{Re } \xi_1|^2}, \quad |\text{Im } \xi_1| < b|\text{Re } \xi_1|, \quad \bullet = a, m.$$

Then, P satisfies the assumptions of Proposition B.5 with $G_0 = (-2 \cos(x_1), 0)$.

Since the deformation G does not involve x_2 , we may decompose

$$L^2(\mathbb{T}^2) = \bigoplus_{n=-\infty}^{\infty} e^{inx_2} L^2(x_1)$$

and use that $D_{x_2}|_{e^{inx_2}L^2_{x_1}} = n \text{Id}$.




FIGURE B.2. We display the eigenvalues of P_θ on $e^{inx_2}L^2_{x_1}$ with $\theta = 0.8$ as well as those for $P + i\nu\Delta$ for five values of ν approaching 0 and $-20 \leq n \leq 20$. P is chosen as in (B.4) with $V_m = ((1 - \xi^2) + e\xi^2)e^{-\xi^2}$ and $V_a = 0$. These choices guarantee the existence of an embedded eigenvalue at 0 [36, example 1].

In order to discretize the operator $P_\theta|_{e^{inx_2}L^2_{x_1}}$, we replace \mathbb{T}_{x_1} by $\frac{2\pi}{N}(\mathbb{Z}/N\mathbb{Z})$ and denote by $X_1, Y_1 \in [\frac{2\pi}{N}(\mathbb{Z}/N\mathbb{Z})]^N$ vectors with the j^{th} entry $X_1(j) = \frac{2\pi j}{N}$. We will represent the Fourier dual to $\frac{2\pi}{N}(\mathbb{Z}/N\mathbb{Z})$ as $-\frac{N}{2} + \mathbb{Z}/N\mathbb{Z}$ and index vectors on the Fourier side with $K \in \{-\frac{N}{2}, -\frac{N}{2} + 1, \dots, \frac{N}{2} - 1\}$. We then compute a matrix that approximates the action of P_θ on the Fourier series side.

Letting $\gamma_\theta(x) := x + i\theta G_0(x)$ and \mathcal{F}_N the matrix for the discrete Fourier transform,

$$(\mathcal{F}_N)_{K,j} := -\frac{e^{-2\pi ijK/N}}{\sqrt{N}}, \quad -\frac{N}{2} \leq K \leq \frac{N}{2} - 1, \quad 0 \leq j \leq N - 1,$$

we obtain for a vector $u \in \mathbb{C}^N$,

$$\mathcal{F}_N \frac{u}{\gamma'_\theta} = \mathbf{M} \mathcal{F}_N u, \quad \mathbf{M} := \mathcal{F}_N \text{Diag} \left(\frac{1}{\gamma'_\theta(X_1)} \right) \mathcal{F}_N^*,$$

so that

$$[\mathcal{F}_N (D_{x_1})_\theta u] \approx [\mathbf{MK} \mathcal{F}_N u]$$

where

$$\mathbf{K} = \text{Diag}(K).$$

At this point we discretize $(\langle D \rangle^{-1/2})_\theta$ using the functional calculus of [35, sec. 4]. In particular, writing

$$\langle D \rangle_\theta = (1 + n^2)\text{Id} + (\mathbf{MK})^2,$$

we have

$$\mathcal{F}_N[(\langle D \rangle^{-\frac{1}{2}})_\theta u] \approx (\sqrt{\langle \mathbf{D} \rangle_\theta})^{-1} \mathcal{F}_N u$$

where the square root is taken in the sense of matrices and all eigenvalues are taken with nonnegative real part.

Remark B.8. Note that the functional calculus definition of $(\langle D \rangle^{-1/2})_\theta$ agrees with the definition (B.3) with $p = (1 + n^2 + \xi^2)^{-\frac{1}{2}}$. To see this, observe that at $\theta = 0$, the two operators agree and hence their analytic extensions to Γ_θ agree.

The operators $[V_{a/m}(D)]_\theta$ are computed by using (B.3) to write

$$(V_{a/m}(D))_\theta u(x) = \sum_k \frac{1}{2\pi} \int e^{i(\gamma_\theta(x) - \gamma_\theta(y))k} V_{a/m}(k) u(y) dy.$$

This sum converges since $V_{a/m}$ satisfies (B.5). We then evaluate x and y on $\frac{2\pi}{N}(\mathbb{Z}/N\mathbb{Z})$ in the above kernel to obtain the matrix approximation on the Fourier transform side. More precisely, putting

$$(\widehat{V}_\bullet^\theta)_{ij} := \sum_k \frac{1}{2\pi} \int e^{i(\gamma_\theta(X_1(i)) - \gamma_\theta(Y_1(j)))k} V_\bullet(k), \quad \bullet = a, m,$$

we have

$$\mathcal{F}_N V_\bullet(D)_\theta u \approx \mathbf{V}_\bullet^\theta \mathcal{F}_N u, \quad \mathbf{V}_\bullet^\theta = \mathcal{F}_N \widehat{V}_\bullet^\theta \mathcal{F}_N^*.$$

Note that this approximation is valid since we take $|V(\xi)| \leq C e^{-C|\xi|}$ and therefore the sum in k converges rapidly. Finally, writing

$$\mathbf{S}^\theta = \mathcal{F}_N \text{Diag}(\sin(\gamma_\theta(X_1))) \mathcal{F}_N^*,$$

Our total operator is then approximated by

$$P_\theta|_{e^{inx_2} L^2_{x_1}} \approx \mathcal{F}_N^* \mathbf{P}_N^\theta \mathcal{F}_N,$$

$$\mathbf{P}_N^\theta := n(\sqrt{\langle \mathbf{D} \rangle_\theta})^{-1} + \mathbf{S}^\theta (\text{Id} - \mathbf{V}_m^\theta) + (\text{Id} - \mathbf{V}_m^\theta) \mathbf{S} + \mathbf{V}_a^\theta.$$

Since \mathcal{F}_N is unitary, we compute the eigenvalues of \mathbf{P}_N^θ to approximate the eigenvalues of $P_\theta|_{e^{inx_2} L^2_{x_1}}$.

When approximating $P + i\nu\Delta$, these computations are much simpler and we use the standard Fourier series approximations

$$P + i\nu\Delta \approx \mathcal{F}_N^* \mathbf{P}_N^\nu \mathcal{F}_N, \quad \mathbf{P}_N^\nu := n(\mathbf{D})^{-1/2} + \mathbf{S}(\text{Id} - \mathbf{V}_m) + (\text{Id} - \mathbf{V}_m) \mathbf{S} + \mathbf{V}_a - i\nu \mathbf{K}^2,$$

where

$$\langle \mathbf{D} \rangle^{-1/2} := \text{Diag}(\langle K \rangle^{-1/2}), \quad \mathbf{V}_\bullet := \text{Diag}(V_\bullet(K)),$$

$$\mathbf{S} = \mathcal{F}_N \text{Diag}(\sin(X_1)) \mathcal{F}_N^*.$$

The results of several numerical experiments are displayed in Figures 1.1, B.1, and B.2.

Acknowledgments. We would like to thank Semyon Dyatlov for many enlightening discussions and Johannes Sjöstrand for helpful comments on the first version of [15]. Partial support by National Science Foundation grants DMS-1900434 and DMS-1502661 (JG) and DMS-1500852 (MZ) is also gratefully acknowledged.

Bibliography

- [1] Boutet de Monvel, L.; Guillemin, V. *The spectral theory of Toeplitz operators*. Annals of Mathematics Studies, 99. Princeton University Press, Princeton, NJ; University of Tokyo Press, Tokyo, 1981. doi:10.1515/9781400881444
- [2] Boutet de Monvel, L.; Sjöstrand, J. Sur la singularité des noyaux de Bergman et de Szegő. *Journées: Équations aux Dérivées Partielles de Rennes (1975)*, 123–164. *Astérisque*, **34-35**. Soc. Math. France, Paris, 1976.
- [3] Colin de Verdière, Y. Spectral theory of pseudo-differential operators of degree 0 and applications to forced linear waves. *Anal. PDE* **13** (2020), no. 5, 1521–1537. doi:10.2140/apde.2020.13.1521
- [4] Colin de Verdière, Y.; Saint-Raymond, L. Attractors for two dimensional waves with homogeneous Hamiltonians of degree 0. *Comm. Pure Appl. Math.* **73** (2020), no. 2, 421–462. doi:10.1002/cpa.21845
- [5] Dang, N.; Riviere, R. Pollicott-Ruelle spectrum and Witten Laplacians. *J. Eur. Math. Soc. (JEMS)* **23** (2021), no. 6, 1797–1857. doi:10.4171/JEMS/1044
- [6] Dauxois, T.; Joubaud, S.; Odier, P.; Venaille, A. Instabilities of internal gravity wave beams. *Annual review of fluid mechanics*. Vol. 50, 131–156. Annu. Rev. Fluid Mech., 50. Annual Reviews, Palo Alto, CA, 2018.
- [7] Dimassi, M.; Sjöstrand, J. *Spectral asymptotics in the semi-classical limit*. London Mathematical Society Lecture Note Series, 268. Cambridge University Press, Cambridge, 1999. doi:10.1017/CBO9780511662195
- [8] Drouot, A. Stochastic stability of Pollicott-Ruelle resonances. *Comm. Math. Phys.* **356** (2017), 357–396.
- [9] Dyatlov, S.; Zworski, M. Stochastic stability of Pollicott-Ruelle resonances. *Nonlinearity* **28** (2015), no. 10, 3511–3533. doi:10.1088/0951-7715/28/10/3511
- [10] Dyatlov, S.; Zworski, M. Dynamical zeta functions for Anosov flows via microlocal analysis. *Ann. Sci. Éc. Norm. Supér. (4)* **49** (2016), no. 3, 543–577. doi:10.24033/asens.2290
- [11] Dyatlov, S.; Zworski, M. *Mathematical theory of scattering resonances*. Graduate Studies in Mathematics, 200. American Mathematical Society, Providence, RI, 2019. doi:10.1090/gsm/200
- [12] Dyatlov, S.; Zworski, M. Microlocal analysis of forced waves. *Pure Appl. Anal.* **1** (2019), no. 3, 359–394. doi:10.2140/paa.2019.1.359
- [13] Faure, F.; Sjöstrand, J. Upper bound on the density of Ruelle resonances for Anosov flows. *Comm. Math. Phys.* **308** (2011), no. 2, 325–364. doi:10.1007/s00220-011-1349-z
- [14] Frenkel, E.; Losev, A.; Nekrasov, N. Instantons beyond topological theory. I, *J. Inst. Math. Jussieu* **10** (2011), no. 3, 463–565. doi:10.1017/S1474748011000077
- [15] Galkowski, J.; Zworski, M. Introduction to complex microlocal deformations. Preprint, 2019. arXiv:1912.09845 [math.AP]
- [16] Guedes Bonthonneau, Y.; Jézéquel, M. FBI transform in Gevrey classes and Anosov flows. Preprint, 2020. arXiv:2001.03610 [math.AP]
- [17] Hassell, A.; Melrose, R.; Vasy, A. Spectral and scattering theory for symbolic potentials of order zero. *Adv. Math.* **181** (2004), no. 1, 1–87. doi:10.1016/S0001-8708(03)00020-3
- [18] Helffer, B.; Sjöstrand, J. Resonances en limite semiclassique, *Bull. Soc. Math. France* **114**, no. 24–25, 1986.

- [19] Hitrik, M.; Sjöstrand, J. Two minicourses on analytic microlocal analysis. *Algebraic and analytic microlocal analysis*, 483–540. Springer Proc. Math. Stat., 269. Springer, Cham, 2018. doi:10.1007/978-3-030-01588-6_10
- [20] Hörmander, L. *The analysis of linear partial differential operators. I. Distribution theory and Fourier analysis*. Grundlehren der mathematischen Wissenschaften, 256. Springer, Berlin, 1983. doi:10.1007/978-3-642-96750-4
- [21] Hörmander, L. *The analysis of linear partial differential operators. III. Pseudodifferential operators*. Grundlehren der mathematischen Wissenschaften, 274. Springer, Berlin, 1985.
- [22] Jean, F. *Control of nonholonomic systems: from sub-Riemannian geometry to motion planning*. SpringerBriefs in Mathematics. Springer, Cham, 2014. doi:10.1007/978-3-319-08690-3
- [23] Jézéquel, M. Global trace formula for ultra-differentiable Anosov flows. *Comm. Math. Phys.* **385** (2021), no. 3, 1771–1834. doi:10.1007/s00220-020-03930-x
- [24] Martinez, A. *An introduction to semiclassical and microlocal analysis*. Universitext. Springer, New York, 2002. doi:10.1007/978-1-4757-4495-8
- [25] Melin, A.; Sjöstrand, J. *Fourier integral operators with complex-valued phase functions. Fourier integral operators and partial differential equations (Colloq. Internat., Univ. Nice, Nice, 1974)*, 120–223. Lecture Notes in Math., 459. Springer, Berlin, 1975.
- [26] Melrose, R.; Zworski, M. Scattering metrics and geodesic flow at infinity. *Invent. Math.* **124** (1996), no. 1-3, 389–436. doi:10.1007/s002220050058
- [27] Nakamura, S. On Martinez’ method of phase space tunneling. *Rev. Math. Phys.* **7** (1995), no. 3, 431–441. doi:10.1142/S0129055X95000189
- [28] Ralston, J. V. On stationary modes in inviscid rotating fluids. *J. Math. Anal. Appl.* **44** (1973), 366–383. doi:10.1016/0022-247X(73)90065-6
- [29] Rieutord M.; Valdetaro, L. Axisymmetric inertial modes in a spherical shell at low Ekman numbers. *J. Fluid Mech.* **844** (2018), 597–634. doi:10.1017/jfm.2018.201
- [30] Sjöstrand, J. *Applications of Fourier distributions with complex phasefunctions. Fourier integral operators and partial differential equations (Colloq. Internat., Univ. Nice, Nice, 1974)*, 255–282. Lecture Notes in Math., 459. Springer, Berlin, 1975.
- [31] Sjöstrand, J. Singularités analytiques microlocales. *Astérisque* **95**, 1–166. Soc. Math. France, Paris, 1982.
- [32] Sjöstrand, J. Geometric bounds on the density of resonances for semiclassical problems. *Duke Math. J.* **60** (1990), no. 1, 1–57. doi:10.1215/S0012-7094-90-06001-6
- [33] Sjöstrand, J. Density of resonances for strictly convex analytic obstacles. *Canad. J. Math.* **48** (1996), no. 2, 397–447. doi:10.4153/CJM-1996-022-9
- [34] Sjöstrand, J. *Lectures on resonances*. Version préliminaire. Printemps 2002. Available at: <https://sjostrand.perso.math.cnrs.fr/Coursgbg.pdf>
- [35] Sjöstrand, J.; Zworski, M. Complex scaling and the distribution of scattering poles. *J. Amer. Math. Soc.* **4** (1991), no. 4, 729–769. doi:10.2307/2939287
- [36] Tao, Z. 0-th order pseudodifferential operators on the circle. Preprint, 2019. arXiv:1909.06316 [math.AP]
- [37] Trèves, F. *Introduction to pseudodifferential and Fourier integral operators. Vol. 2. Fourier integral operators*. University Series in Mathematics. Plenum Press, New York–London, 1980.
- [38] Wang, J. The scattering matrix for 0th order pseudodifferential operators. Preprint, 2019. arXiv:1909.06484 [math.AP]
- [39] Xiong, H. Resonances as viscosity limits for exterior dilation analytic potentials. Preprint, 2020. arXiv:2002.12490 [math-ph]
- [40] Zworski, M. *Semiclassical analysis*. Graduate Studies in Mathematics, 138. American Mathematical Society, Providence, RI, 2012. doi:10.1090/gsm/138
- [41] Zworski, M. Scattering resonances as viscosity limits. *Algebraic and analytic microlocal analysis*, 635–654. Springer Proc. Math. Stat., 269. Springer, Cham, 2018. doi:10.1007/978-3-030-01588-6_14

JEFFREY GALKOWSKI
Department of Mathematics
University College
London WC1H 0AY
UNITED KINGDOM
E-mail: j.galkowski@ucl.ac.uk

MACIEJ ZWORSKI
Department of Mathematics
University of California, Berkeley
Berkeley, CA 94720
USA
E-mail: zworski@
math.berkeley.edu

Received May 2021.