

The 2020 edition of the GEISA spectroscopic database

T. Delahaye ¹, R. Armante ¹, A. Chédin ¹, L. Crépeau ¹, C. Crevoisier ¹, V. Douet ², N. Jacquinet-Husson ¹, A. Perrin ¹, N.A. Scott ¹, A. Barbe ³, V. Boudon ⁴, A. Campargue ⁵, L.H. Coudert ⁶, V. Ebert ^{14,15}, J.-M. Flaud ⁷, R.R. Gamache ⁹, D. Jacquemart ¹⁰, A. Jolly ⁷, F. Kwabia Tchana ⁷, A. Kyuberis ²², G. Li ¹⁴, O.M. Lyulin ¹¹, L. Manceron ^{10,16}, S. Mikhailenko ¹¹, N. Moazzen-Ahmadi ²³, H.S.P. Müller ¹², O.V. Naumenko ¹¹, A. Nikitin ¹¹, V.I. Perevalov ¹¹, C. Richard ⁴, E. Starikova ¹¹, S.A. Tashkun ¹¹, V.I.G. Tyuterev ^{3,17}, J. Vander Auwera ¹³, B. Vispoel ^{9,18,19}, A. Yachmenev ^{20,21}, S. Yurchenko ⁸.

¹Laboratoire de Météorologie Dynamique/IPSL, CNRS, Ecole Polytechnique, Université Paris-Saclay, 91128, Palaiseau, France

²Institut Pierre Simon Laplace, Université Pierre et Marie Curie, 75252 Paris, France

³ Université de Reims-Champagne-Ardenne, Groupe de Spectrométrie Moléculaire et Atmosphérique, 51062 Reims, France

⁴ Laboratoire Interdisciplinaire Carnot de Bourgogne, UMR 6303 CNRS-Univ. Bourgogne Franche-Comté, 9 Avenue Alain Savary, BP 47 870, F-21078 DIJON Cedex, France

⁵ Univ. Grenoble Alpes, CNRS, LIPhy, 38000 Grenoble, France

⁶ Université Paris-Saclay, CNRS, Institut des Sciences Moléculaires d'Orsay, 91405 Orsay, France

⁷ Laboratoire Interuniversitaire des Systèmes Atmosphériques (LISA), UMR CNRS 7583, Université de Paris, Université Paris-Est Créteil, Institut Pierre-Simon Laplace, 61 avenue du Général de Gaulle, 94010 Créteil Cedex, France

⁸ Department of Physics and Astronomy, University College London, London WC1E 6BT, United Kingdom

⁹ University of Massachusetts Lowell, Department of Environmental Earth and Atmospheric Sciences, Lowell, MA 01854, USA

¹⁰ Sorbonne Université, CNRS, De la MOlécule aux NANO-objets : Réactivité, Interactions et Spectroscopies, MONARIS, 75005 Paris, France

¹¹ V.E. Zuev Institute of Atmospheric Optics, SB, Russian Academy of Sciences, 1, Academician Zuev square, 634055, Tomsk, Russia

¹² I. Physikalisches Institut, Universität zu Köln, 50937 Köln, Germany

¹³ Spectroscopy, Quantum Chemistry and Atmospheric Remote Sensing (SQUARES), C.P. 160/09, Université Libre de Bruxelles, 50 avenue F.D. Roosevelt, B-1050 Brussels, Belgium.

¹⁴ Physikalisch-Technische Bundesanstalt, Bundesallee 100, 38116 Braunschweig, Germany

¹⁵ Reactive Flows and Diagnostics, Technische Universität Darmstadt, Otto-Berndt-Str. 3 64287 Darmstadt, Germany

¹⁶ Synchrotron Soleil Ligne AILES, BP 48, 91192 Cedex Gif-sur-Yvette, France

¹⁷ QUAMER laboratory, Tomsk State University, 36 Lenin Ave., Tomsk, Russia 634050

¹⁸ Research Unit Lasers and Spectroscopies (LLS), Institute of Life, Earth and Environment (ILEE), University of Namur (UNamur), 61 rue de Bruxelles, B-5000, Namur, Belgium

¹⁹ Royal Belgian Institute for Space Aeronomy (BIRA-IASB), 3 Avenue Circulaire, 1180, Brussels, Belgium

²⁰ Center for Free-Electron Laser Science, Deutsches Elektronen-Synchrotron DESY, Notkestraße 85, 22607 Hamburg, Germany

²¹ Center for Ultrafast Imaging, Universität Hamburg, Luruper Chaussee 149, 22761 Hamburg, Germany

²² Van Swinderen Institute for Particle Physics and Gravity, University of Groningen, Nijenborgh 4, 9747 Groningen, The Netherlands

²³ Department of Physics and Astronomy, University of Calgary, 2500 University Drive North West, Calgary, Alberta T2N 1N4, Canada

Number of pages for text:	55
Number of pages for references	13
Number of Tables	11
Number of Figures	13

Highlights

- GEISA-2020 database release: 6,746,987 entries in the line parameters database
- 23 molecules updated and 6 new molecules added (HONO, COFCl, CH₃F, CH₃I, RuO₄, H₂C₃H₂ (isomer of C₃H₄))
- Evaluation of the spectroscopic parameters through radiative transfer simulations compared to atmospheric spectra (SPARTE chain)

Corresponding author

Dr. Raymond Armante

Phone: +33169335111

E-mail: raymond.armante@lmd.ipsl.fr

ABSTRACT

This paper describes the 2020 release of the GEISA database (Gestion et Etude des Informations Spectroscopiques Atmosphériques: Management and Study of Atmospheric Spectroscopic Information), developed and maintained at LMD since 1974. GEISA is the reference database for several current or planned Thermal and Short-Wave InfraRed (TIR and SWIR) space missions IASI (Infrared Atmospheric Sounding Interferometer), IASI-NG (IASI New Generation), MicroCarb (Carbon Dioxide Monitoring Mission), Merlin (Methane Remote sensing Lidar mission). It is actually a compilation of three databases: the “line parameters database”, the “cross-section sub-database” and the “microphysical and optical properties of atmospheric aerosols sub-database”. The new edition concerns only the line parameters dataset, with significant updates and additions implemented using the best available spectroscopic data.

The GEISA-2020 line parameters database involves 58 molecules (145 isotopic species) and contains 6,746,987 entries, in the spectral range from 10^{-6} to 35877 cm^{-1} . In this version, 23 molecules have been updated (with 10 new isotopic species) and 6 new molecules have been added (HONO, COFCl, CH₃F, CH₃I, RuO₄, H₂C₃H₂ (isomer of C₃H₄)) corresponding to 15 isotopic species. The compilation can be accessed through the AERIS data and services center for the atmosphere website (<https://geisa.aeris-data.fr/>), with the development of a powerful graphical tool and convenient searching, filtering, and plotting of data using modern technologies (PostgreSQL database, REST API, VueJS, Plotly).

Based on four examples (H₂O, O₃, O₂ and SF₆), this paper also shows how the LMD in house validation algorithm SPARTE (Spectroscopic Parameters And Radiative Transfer Evaluation) helps to evaluate, correct, reject or defer the input of new spectroscopic data into GEISA and this, thanks to iterations with researchers from different communities (spectroscopy, radiative transfer).

Key words: molecular spectroscopic database ; line parameters ; earth and planetary radiative transfer ; atmospheric absorption ; spectroscopic parameters evaluation

1 Introduction¹:

From the mid 1960's, various scientific communities (Remote Sensing, Astrophysics, Atmospheric Physics, Metrology, Climate and Chemistry) have required access to databases detailing the spectral characteristics of atmospheric molecular absorption and atmospheric particle scattering. The purpose of spectroscopic databases like GEISA [1] and HITRAN [2] is to give, to each community, access to the best information at a given time.

In France, archiving spectroscopic data to be used in conjunction with remote sensing applications was first initiated in the 70's. The first release of the GEISA database [1] included major absorbers (N_2 , O_2 , H_2O , CO_2 , O_3 , N_2O , CO , CH_4) of the earth atmosphere as well as complementary species (e.g. NH_3 , HNO_3 , PH_3 , C_2H_4 , C_2H_2 , HC_3N , $HCOOH$, NO , SO_2 , NO_2). Some molecules, mainly related to planetary atmospheres (e.g. within the frame of the Voyager experiment and observations of the giant planets) like GeH_4 , C_3H_8 , C_2N_2 , C_3H_4 , HNC , C_6H_6 , and C_2HD were also included. Since its first publication, (Chédin *et al.* (1982) [3]), including spectroscopic data and management softwares, GEISA has undergone numerous updates (Husson *et al.* (1992) [4], Jacquinet-Husson *et al.* (1999, 2008, 2009, 2016) [1], [5]–[7]).

One of the differences of GEISA, in comparison with other databases such as HITRAN, has been to consider, when necessary, isotopologues of a species with a symmetry property different from that of the main isotopologue as an independent molecular species. This was initiated in the mid 70's with CH_3D and CH_4 within the frame of the Voyager experiment and later on with HDO and H_2O , C_2HD and C_2H_2 . Another difference is that some species are specifically considered by only one database (for example O and H_2 for HITRAN, C_3H_8 and C_3H_4 for GEISA).

Among the important number of information updated, a large part has been done with the valuable and sustained support of the international spatial agencies. Especially, since the launch of [Metop-A](#) (24 October 2006), GEISA became the official reference spectroscopic database used by the international working group (ISSWG) in charge of the [IASI](#) instrument. GEISA is also now involved in the definition of 3 future space missions: [IASI-NG](#), [Merlin](#) and [MicroCarb](#).

¹ Acronyms used in the text are documented in Appendix A

GEISA and its associated management software facilities are implemented and distributed (in the same way as GEISA-2011) via the atmospheric pole AERIS. It is used on-line by more than 350 laboratories working in various fields including atmospheric physics, planetary science, astronomy, astrophysics.

As in previous editions of GEISA, new data have been provided by different groups. They are listed in Column 3 of Table 1 together with their contribution.

Since the beginning of the 2000's, a specific effort has been made in GEISA and HITRAN to undertake a more systematic evaluation of spectroscopic parameters when selecting, introducing or replacing data. This has been enabled by the evolving capacity of several space-borne, ground-based or laboratory observations that have reached a level of precision sufficient to evaluate a large range of spectroscopic parameters produced by laboratory measurements or theory. Concerning GEISA, use is made of the now well-established evaluation method (called SPARTE, Spectroscopic Parameters And Radiative Transfer Evaluation) [8] that was developed and first used during the former GEISA release (called GEISA-2015 in the following). Using high resolution and high signal-to-noise ratio space-borne or ground-based measurements (like the one provided in the Near and Short-Wave Infrared by ground-based spectrometers of the Total Carbon Column Observing Network (TCCON) [9] and in the Thermal Infrared by the Infrared Atmospheric Sounding Interferometer (IASI) SPARTE enables to identify spectroscopic issues by investigating the differences between observations and simulations performed using the 4A/OP radiative transfer code [10].

This paper provides, in section 2, a detailed description of the data newly implemented or corrected in the latest 2020 GEISA release, for the line parameters sub-database only (the corresponding cross-section and aerosol will be updated and documented in a forthcoming paper). In this GEISA-2020 edition, the format is unchanged from the previous one. The inclusion of additional spectroscopic parameters describing refined collisional effects such as the speed dependence and Dicke narrowing effects is planned for a forthcoming version.

Examples of spectroscopic evaluations performed using, among others, the SPARTE algorithm are presented in Section 3. Focusing on H₂O, O₂, O₃ and SF₆ molecules, we illustrate how the evaluation guided our selection of sources of line parameters.

2 GEISA-2020 Line parameters database description

General Overview

The GEISA-2020 edition is based on the release of spectroscopic data for 23 molecules in the range of 10⁻⁶ to 35,877 cm⁻¹, and on the addition of 6 molecules in the range 296 to 1913 cm⁻¹

(i.e. ‘HONO, COFCl, CH₃I, CH₃F, RuO₄ and H₂C₃H₂ (allene, isomer of C₃H₄)), whose molecule code number (parameter I) has been fixed to 53 to 58, respectively. The molecules which were updated are listed in Table 1, together with their molecule code in the GEISA notation and indication about the involved research groups.

Molecule ID.	Code	Contributors
H₂O	1	A. Campargue, S. Mikhailenko, Y. Yurchenko, B. Gamache (CA6, CA9, K19, Y20)
CO₂	2	V. Perevalov, S. Tashkun (P19)
O₃	3	A.Barbe, S.Mikhailenko, E.Starikova, V. Tyuterev (T19, M19)
N₂O	4	V. Ebert, G. Li (L19)
CO	5	G. Li, V. Ebert (L19)
CH₄	6	A. Nikitin, V.Tyuterev, E.Starikova (N19, S19)
SO₂	9	O. Naumenko (1N9, 2N9, 3N9, 4N9, 5N9)
NO₂	10	A. Perrin, L. Manceron (P19)
NH₃	11	N. Maaroufi, F. Kawbia (M19)
PH₃	12	A. Nikitin(N19)
HNO₃	13	A. Perrin (H ¹⁵ NO ₃ New isotopologue, P19)
HF, HBr, HI	15,17,18	G. Li, I. Gordon (L19)
HCl	16	G. Li, V. Ebert (L19)
OCS	20	S. Galalou; F. Kwabia (G19)
C₂H₆	22	N. Moazzen-Ahmadi (M19)
C₂H₂	24	D. Jacquemart, A. Campargue, V. Perevalov, O. Lyulin (J19, J20, L19, C19)
C₂H₄	25	J. Vander Auwera (A19)
GeH₄	26	V. Boudon (B19)
NO⁺	45	H.S.P. Müller (M18)
CF₄	49	V. Boudon (B19)
HDO	51	L. Coudert, A. Kyuberis (CO9, K19)
New molecules		
HONO	53	A. Perrin (P18)
COFCl	54	A. Perrin (P19)
CH₃I	55	F. Kwabia, A. Perrin (P19)
CH₃F	56	D. Jacquemart (J19)
RuO₄	57	V. Boudon (B19)
H₂C₃H₂, Allene (C₃H₄ isomer)	58	A. Jolly (J19)

Table 1: Updated molecular species in the GEISA-2020 edition. The internal GEISA code for the identification of the source of the data is indicated within parentheses (parameter G).

To have a more precise idea of the update, Table 2 gives the main differences between GEISA-2015 and GEISA-2020 (in term of spectral range, number of lines and integrated intensity range).

		GEISA-2015				GEISA-2020			
Mol.	ID	Spectral range (cm ⁻¹)	# lines	Exponent of the intensity (cm molecule ⁻¹) at 296 K.		Spectral range (cm ⁻¹)	# lines	Exponent of the intensity (cm molecule ⁻¹) at 296 K.	
				Max.	Min.			Max.	Min.
				H ₂ O	1			0.052 -25336.949	191846
CO ₂	2	5.890 -14075.298	534227	-18	-30	345.936 - 14075.301	532533	-18	-30
O ₃	3	0.026 - 6996.681	405919	-20	-31	0.026 - 6996.681	464288	-20	-31
N ₂ O	4	0.838 - 7796.633	50633	-17	-25	0.838 - 7796.633	50633	-18	-26
CO	5	3.414 - 8464.882	13515	-18	-77	3.402 - 14477.377	14985	-19	-78
CH ₄	6	0.001 -11501.877	421811	-19	-39	0.001 - 11501.877	447004	-19	-39
O ₂	7	10-6 -15927.804	16197	-24	-54	0.000 - 15927.804	16197	-24	-54
NO *	8	10-6 - 9273.214	105079	-19	-94	0.000 - 9273.214	105079	-20	-95
SO ₂	9	0.017 -4092.948	83668	-20	-30	0.017 - 4159.945	561008	-20	-30
NO ₂	10	0.498 - 3074.152	104223	-18	-27	0.498 - 4775.314	185965	-19	-28
NH ₃	11	0.058 -6999.429	46414	-19	-39	0.058 - 6999.429	46414	-19	-39
PH ₃	12	17.805 - 3601.652	20364	-18	-27	8.904 - 3659.266	34542	-19	-28
HNO ₃	13	0.012 -1769.982	691161	-20	-28	0.012- 1769.982	738171	-20	-28
OH *	14	0.005 - 35877.031	42866	-16	-84	0.005 - 35877.031	42866	-17	-85
HF	15	41.111 - 11535.570	107	-16	-25	13.620 - 32351.592	20010	-17	-99
HCl	16	20.240 -13457.841	533	-18	-25	5.342 - 20231.245	53436	-19	-99
HBr	17	16.231 - 9758.564	1293	-18	-32	7.656 - 16033.492	8980	-19	-71
HI	18	12.509 - 8487.305	806	-19	-29	5.888 - 13907.689	4751	-20	-64
ClO *	19	0.015 - 1207.639	7230	-20	-29	0.015 - 1207.639	7230	-21	-30
OCS	20	0.381 - 4199.671	33809	-17	-27	0.381 - 7821.109	37479	-18	-28
H ₂ CO *	21	0.000 -3099.958	44611	-20	-39	0.000 - 3099.958	44611	-20	-39
C ₂ H ₆	22	706.601 -3070.971	53803	-20	-29	225.045 - 3070.971	65741	-20	-38
CH ₃ D *	23	7.760 -6510.324	58763	-23	-30	7.760 - 6510.324	58763	-23	-30
C ₂ H ₂	24	604.774 -9889.038	12969	-18	-28	13.624 - 9889.038	70107	-18	-31
C ₂ H ₄	25	614.740 -3242.172	53227	-20	-37	614.740 - 3242.172	100100	-20	-37
GeH ₄	26	1937.371 -224.570	824	-18	-21	1928.902 - 2265.678	32372	-19	-23
HCN *	27	0.018 -17581.009	138103	-19	-32	0.019 - 17581.009	138103	-19	-32
C ₃ H ₈ *	28	700.015 - 799.930	8983	-21	-23	700.015 - 799.930	8983	-22	-24
C ₂ N ₂ *	29	200.817 -2181.690	71954	-20	-24	200.818 - 2181.690	71954	-20	-24
C ₄ H ₂ *	30	189.422 -1302.217	417540	-19	-24	189.423 - 1302.217	417540	-19	-24
HC ₃ N *	31	463.604 - 759.989	179347	-19	-23	463.604 - 759.989	179347	-20	-24
HOCl *	32	0.0236 - 3799.682	17862	-19	-27	0.024 - 3799.682	17862	-20	-28
N ₂ *	33	1992.63 -2625.497	120	-27	-33	1992.628 - 2625.497	120	-28	-34
CH ₃ Cl *	34	0.872 -3197.961	83043	-21	-32	0.873 - 3197.961	83043	-21	-32
H ₂ O ₂ *	35	0.043 - 1730.371	126983	-19	-28	0.043 - 1730.371	126983	-20	-29
H ₂ S *	36	1.030 -1329.780	58650	-19	-30	1.031 - 11329.780	58650	-19	-30
HCOOH *	37	10.018 - 1889.334	62684	-19	-25	10.018 - 1889.334	62684	-20	-26
COF ₂ *	38	725.005 -2001.348	70904	-19	-23	725.005 - 2001.348	70904	-20	-24
SF ₆ *	39	588.488 - 975.787	92398	-19	-23	588.488 - 975.787	92398	-20	-24
C ₃ H ₄ *	40	288.913 - 673.479	19001	-19	-23	288.913 - 673.479	19001	-20	-24
HO ₂ *	41	0.173 - 3675.819	38804	-19	-25	0.173 - 3675.819	38804	-20	-26
ClONO ₂ *	42	0.636 - 797.741	356899	-21	-27	0.636 - 797.741	356899	-22	-28
CH ₃ Br *	43	794.403 -1705.612	36911	-21	-27	794.403 - 1705.612	36911	-21	-27
CH ₃ OH *	44	0.019 - 1407.206	19897	-19	-34	0.019 - 1407.206	19897	-20	-35
NO ⁺	45	1634.83 -2530.462	1206	-18	-80	3.938 - 4742.008	1333	-19	-81
HNC *	46	0.145 -4692.098	75554	-18	-30	0.146 - 4692.098	75554	-18	-30

		GEISA-2015				GEISA-2020			
Mol.	ID	Spectral range (cm ⁻¹)	# lines	Exponent of the intensity (cm molecule ⁻¹) at 296 K.		Spectral range (cm ⁻¹)	# lines	Exponent of the intensity (cm molecule ⁻¹) at 296 K.	
				Max.	Min.			Max.	Min.
				C ₆ H ₆ *	47			642.427 - 705.262	9797
C ₂ HD *	48	416.785 - 3421.864	15512	-22	-28	416.785 - 3421.864	15512	-23	-29
CF ₄	49	594.581 - 1312.647	60033	-19	-23	1.173 - 1329.697	258208	-20	-30
CH ₃ CN *	50	890.052 - 1650.000	17172	-19	-37	890.052 - 1650.000	17172	-20	-38
HDO	51	0.007 - 17080.098	63641	-22	-33	0.000 - 19935.167	77599	-22	-39
SO ₃ *	52	0.477 - 2824.347	10881	-19	-31	0.478 - 2824.347	10881	-19	-31
HONO	53					722.534 - 996.281	26041	-20	-25
COFCL	54					734.997 - 1912.632	215639	-20	-24
CH ₃ I	55					693.022 - 1125.255	70291	-21	-31
CH ₃ F	56					1067.375 - 1290.255	1499	-21	-24
RUO ₄	57					889.987 - 948.138	30205	-20	-23
H ₂ C ₃ H ₂	58					296.316 - 1192.446	31686	-20	-27
Total # lines			5,067,351			6,746,987			

*: Molecule which has not been updated in the GEISA-2020

Table 2. Contents of the GEISA-2020 line parameters database. Details per molecule of the evolution of GEISA contents since its 2015 edition. Reference temperature is 296 K.

Description of updates per individual molecular species

This description is given in sub-sections 2.2.1 to 2.2.25, for each molecular species identified by its formula associated with its GEISA identification code.

2.1.1 H₂O and HDO

2.1.1.1 H₂O (molecule 1)

New H₂O data implemented in GEISA-2020 consist in three main parts which are: (i) an updated line list for the main H₂¹⁶O isotopologue between 5850 and 8340 cm⁻¹, (ii) the addition of electric-quadrupole transitions up to 10000 cm⁻¹, and (iii) a major update of doubly deuterated D₂O isotopologues up to 18,000 cm⁻¹.

GEISA-2020 was updated using line intensities, air-broadening and -shifting coefficients provided from Mikhailenko *et al.* [11], whereas the line positions were refined using the most recent version of their line list [12]. Note that local discrepancies due to erroneous broadening values of some H₂O lines were found in these line lists and in GEISA-2015 (see for example residuals of the H₂O line at 7804.6092 cm⁻¹ in section 3.1.1), for which calculated air-broadening parameters from Vispoel *et al.* [13], [14] were used to correct the incriminated lines (see details below). As a result, 40,980 lines were updated in GEISA-2020 (from which 7572 are new lines not referenced in GEISA-2015). Note that we present in section 3.1.1 a quantitative evaluation of these H₂O data.

The internal GEISA code for the data identification has been set to [CA6](#) and [CA9](#).

Up to now, similarly to all other databases, the GEISA H₂O database was limited to ro-vibrational electric-dipole transitions. Electric-quadrupole transitions do also exist, but they are typically a million times weaker than electric-dipole transitions, making their observation extremely challenging. The recent detection of weak quadrupole transitions in the spectrum of water vapor [15], [16] has shown that in their present status the spectroscopic databases are not complete for polyatomic molecules. Actually, databases provide lists of electric-dipole transitions above an intensity cut-off (*e.g.* 10^{-36} cm/molecule for water in the GEISA database) largely below intensity values of the strongest electric-quadrupole lines (*e.g.* up to 10^{-26} cm/molecule for water vapor [16]). The laboratory measurements of Refs [15], [16] (in total, about twenty quadrupole lines near 5.4, 2.5 and 1.3 μm) have validated the *ab initio* intensities computed in Ref. [15] within the experimental error bars. Consequently, in the present release, the *ab initio* quadrupole line list of Ref. [15] up to $10,000\text{ cm}^{-1}$ (6429 lines in total above a 10^{-30} cm/molecule intensity cut-off) has been incorporated in the water vapor line list for the main isotopologue H₂¹⁶O. Figure 1 shows an overview of the electric-dipole and electric-quadrupole transitions listed in GEISA 2020, up to 9500 cm^{-1} . In the calculations of Ref. [15], the variational values of the lower and upper energy levels of H₂¹⁶O were replaced by the corresponding empirical values from IUPAC [17]. The resulting line position accuracy is thus expected to be of the order of 10^{-3} cm^{-1} for most of the quadrupole lines.




Figure 1: Overview of the H_2^{16}O absorption line list in GEISA-2020 up to 9000 cm^{-1} . The electric-quadrupole transitions (red dots) are superimposed to the electric-dipole transitions (black dots).

The internal GEISA code for the data identification has been set to [Y20](#).

As line shape parameters are critical for atmospheric applications, an updated calculation for $\text{H}_2\text{O}-\text{H}_2\text{O}$, $\text{H}_2\text{O}-\text{N}_2$ and $\text{H}_2\text{O}-\text{O}_2$ broadenings and shifts, along with the temperature dependence of these parameters was provided by Gamache and Vispoel (unpublished data, 2020). These parameters were calculated for H_2^{16}O , H_2^{18}O , H_2^{17}O (but also HD^{17}O , HD^{18}O , D_2^{16}O , D_2^{18}O , and D_2^{17}O species), and include the H_2^{16}O quadrupole transitions. From values calculated using the Modified Complex Robert Bonamy (MCRB) model by Vispoel *et al.* [13] for $\text{H}_2\text{O}-\text{N}_2$ and $\text{H}_2\text{O}-\text{O}_2$ (Vispoel and Gamache, unpublished data, 2020), the pressure broadening information for $\text{H}_2\text{O}-\text{air}$ was produced considering $\gamma(\text{air}) = 0.79\gamma(\text{N}_2) + 0.21\gamma(\text{O}_2)$, with a similar formula for the line shift. Note that the calculations were performed for 13 temperatures from 200 K to 3000 K, considering 10,782 rotational transitions in the rotational band and for 1 to 4 vibrational quanta changes in the ν_1 , ν_2 , and ν_3 bands, giving updated profile parameters for about 140,000 transitions.

Pressure-broadening and -shifting parameters of more than 300,000 dipole transitions needed to be carefully selected, evaluated and systematically compared to previous data of the

database. This specific process being a task of major importance, it will be part of a future dedicated publication on H₂O. As a result, only line-shape parameters for quadrupole transitions were considered for this update of GEISA, except for some (electric-dipole) isolated lines known to be problematic in GEISA-2015 (cf. section 3.1.1 for a detailed example of H₂O data validation). For GEISA-2020, line-shape parameters of 6475 lines were updated in this way.

Besides the main isotopologue H₂¹⁶O, another update was made in GEISA-2020 for the doubly deuterated D₂O isotopologues of water vapor, based on the recent work of Kyuberis *et al.* [18]. A total of 156,375 new lines were added and 5860 lines updated with respect to the previous version of the database. In particular, data for the isotopologue D₂¹⁷O, which were missing in all previous versions of GEISA, were implemented. These *ab initio* calculations were obtained using the base PES (Potential Energy Surface) of Shirin *et al.* [19] which was explicitly developed for D₂O. More details can be found in Ref. [18]. Half-width and pressure-shift data were taken from the measurement database of Gamache and Hartmann [20] for 228, 227, 133 (119, 119, 88 for line shifts) transitions of D₂¹⁶O, D₂¹⁸O, and D₂¹⁷O, respectively. For the remaining transitions, data from Ref. [18] were retained.

The internal GEISA code for the data identification has been set to **K19**.

2.1.1.2 HDO (molecule 51)

Following the data update made for H₂O (cf. previous section), the HDO linelist was also updated using line intensities and air-broadening and -shifting coefficients provided by Mikhailenko *et al.* [11], and line positions which were refined using the most recent version of their line list [12]. As a result, 17,015 lines were updated in GEISA-2020 (from which 4468 are new lines not referenced in GEISA-2015).

The internal GEISA code for the data identification has been set to **CA6** and **CA9**.

Moreover, recent data from Coudert [21], based on the line positions and line strengths analyses spanning the 0 to 2000 cm⁻¹ spectral range, were added in the new version of GEISA. The line list contains rovibrational transitions within the ground and (010) states as well as transitions belonging to the ν_2 band. The maximum value of the rotational quantum number J was set to 22 and an intensity cut-off of 10⁻²⁸ cm⁻¹ / (molecule cm⁻²) was used for a temperature of 296 K and an isotopic abundance [22] of 3.107×10⁻⁴. In total 5813 lines were updated, from which 260 lines were not referenced in GEISA-2015.

The internal GEISA code for the data identification has been set to **CO9**.

Finally, *ab initio* predictions from Kyuberis *et al.* [18] were also considered. The most abundant deuterated water species HD¹⁶O required the most extensive variational calculations. The base potential was the fitted HDO07 PES due to Yurchenko *et al.* [23]. Calculations were performed for J up to 23 but intensities were limited to transitions below 20 000 cm⁻¹. Concerning HD¹⁷O and HD¹⁸O, isotopic abundance factors imply that only strong lines are important for atmospheric studies, which raises fewer issues with unstable intensities. For these species line lists generated by Down were used (see Ref. [18] for more information). In total 67,928 lines were updated, from which 9230 lines were not referenced in GEISA-2015.

The internal GEISA code for the data identification has been set to **K19**.

2.1.2 CO₂ (molecule 2)

The GEISA-2020 carbon dioxide line list is based on the updated version of the CDS-296 databank [24]. Compared to the previous version of the CDS-296 [25] on which GEISA-2015 was based [1], in the new version, the $\Delta P=9$ region for all isotopologues was updated [$P=2V_1+V_2+3V_3$ is the polyad number (V_i being the vibrational quantum numbers)]. Moreover, in the previous version of the CDS-296, the line intensities of some asymmetric isotopologues were incorrectly recalculated from the effective dipole moment parameters of more abundant ones. In the new version, the respective line intensities have been corrected. After the publication of the updated version of the CDS-296 [24], some updates were made [26]. First, the line intensities for the $\Delta P=6$ series of transitions in the ¹⁶O¹²C¹⁸O isotopologue were recalculated using the new set of the effective dipole moment parameters. Secondly, 226 lines with $\Delta \ell_2=4$ for the ¹²C¹⁶O₂, ¹³C¹⁶O₂ and ¹⁶O¹²C¹⁸O isotopologues were added. They were absent in the CDS-296 because of the used $\Delta \ell_2 < 4$ cut-off.

The 01111-00001 band and the respective hot bands of the ¹⁶O¹²C¹⁸O isotopologue were excluded from the last version of the CDS-296 because the effective dipole moment parameters responsible for the line intensities of these bands were fitted to the line intensities retrieved from the Venus spectra obtained by the Venus Express satellite [27]. These line intensities have been considered as not sufficiently precise. Considering its importance for the astronomer's community [28], the specific 01111-00001 band has been added to the GEISA-2020 release: the corresponding values are from the GEISA2011 version.

All these updates are now included in GEISA-2020 and have been referenced with the identifier **P19** (J internal code parameter).

2.1.3 O_3 (molecule 3)

Due to the importance of the ozone molecule for atmospheric applications, measurements and calculations of ozone spectra undergo continuous extension and improvement [29]–[32] (and references therein). The current GEISA-2020 update corresponds essentially to the S&MPO-2020 version of the “Spectroscopy and Molecular Properties of Ozone” information system [33] jointly developed by Reims and Tomsk groups. The linelists were produced using effective Hamiltonian and effective dipole transition moment models derived from analyses of experimental Fourier Transform spectra in the infrared as described in [33], [34].

(i) Main ozone isotopologue $^{16}O_3$

Eighteen bands of the main ozone isotopologue $^{16}O_3$ were newly included in the GEISA databank. In the range between 4384 and 4826 cm^{-1} the hot bands corresponding to the upper tetrad of strongly coupled vibrational states (213)/(080)/(114)/(321) were included. Cold bands of the (023)/(122)/(400) triad of interacting states (4268 – 4455 cm^{-1}) were calculated using unpublished parameters of Barbe and Mikhailenko available at the S&MPO web site (<https://smo.iao.ru/transitions/params>). The resonance coupling parameters were given in the representation of ladder angular momentum components as defined in [33]. The transition moment parameters for the line intensities of the hot bands between 3564 and 3752 cm^{-1} are those of Ref. [30]. The hot bands $5\nu_3-\nu_2$, $3\nu_1+\nu_2+\nu_3-\nu_2$ and $\nu_1+4\nu_3-\nu_2$ (in the region 4128 – 4229 cm^{-1}) were calculated using Effective Hamiltonian (EH) parameters [35] and unpublished transition moment parameters obtained by Barbe. In addition, five previously missing hot bands between 598 and 1542 cm^{-1} belonging to the tetrad of interacting (002)/(101)/(030)/(200) vibrational states were added. Unpublished EH parameters of the upper states as well as published parameters of the lower states [36]–[38] were used for line position calculations. Line intensity calculations used unpublished parameters for the $2\nu_1-\nu_2$ band and parameters of Refs. [39], [40] for the $3\nu_2-2\nu_2$, $\nu_1+\nu_3-2\nu_2$, $3\nu_2-\nu_1$, and $3\nu_2-\nu_3$ bands.

Twenty-four bands previously included in the GEISA line list between 281 and 4122 cm^{-1} were updated:

- The $\nu_3-\nu_2$ and $\nu_1-\nu_2$ hot bands (281 – 446 cm^{-1}) were calculated using EH parameters of Flaud *et al.* [36] and dipole moment transition parameters of Birk *et al.* [41].
- Three hot bands (002)/(101)/(030) – (010) in the 1302 – 1476 cm^{-1} range were calculated using unpublished upper state EH parameters by Barbe and Tyuterev and the dipole moment transition parameters of S&MPO-2015. The lower state parameters were taken from [37].

- The same upper state EH parameters were used for six hot bands (002)/(101)/(200) – (001)/(100) (899 – 1235 cm⁻¹) with the dipole moment transition parameters of [40]. The lower state parameters were taken from [36].
- Two cold bands 2ν₂+ν₃ and ν₁+2ν₂ (2336 – 2617 cm⁻¹) were calculated using parameters by Barbe *et al.* with updates by Sulakshina [42].
- The line positions of the hot (ν₂+3ν₃-ν₂ and ν₁+ν₂+2ν₃-ν₂ (2899 – 3122 cm⁻¹)) and cold (ν₂+3ν₃ and ν₁+ν₂+2ν₃ (3584 – 3831 cm⁻¹)) bands were calculated using EH parameters from [36], [37], [43]. The dipole moment parameters of [43] were used for the calculations of the line intensities of cold bands. The intensities of hot bands were calculated using unpublished parameters by Mikhailenko and Barbe.
- The results of a re-investigation of the spectral range from 3600 to 4300 cm⁻¹ led to a large improvement of the analysis for the strongest ν₁+3ν₃ band in this range [30]. A recent observation of the hot band 4ν₃-ν₃ [30] enabled to identify the strong anharmonic resonance with the “dark” 3ν₁+ν₂ band. As the upper (310) state is also in interaction with the (211) state, it was necessary to include seven vibration states to correctly reproduce 3389 transitions with the root mean square deviation of 3.54×10⁻³ cm⁻¹ for line positions and 1295 intensities within 8.6%. Seven calculated bands in this range were included in the compilation, for a total number of 13536 updated transitions (from which 7447 are new lines in GEISA) between 2816 and 4122 cm⁻¹. Newly included bands are summarized in the **Table 3**. Figure 2 gives an overview of the added lines in the considered spectral range.

The internal GEISA code for the data described above was set to *M19*.

Band	Num. Tra.	Region / cm ⁻¹	S _v [*] / cm/mol
030 – 020	2558	598.26 – 804.86	1.928E-21
101 – 020	6	656.12 – 725.68	4.902E-25
030 – 100	20	956.13 – 1170.40	1.902E-23
030 – 001	15	957.33 – 1201.66	2.238E-24
200 – 010	102	1366.62 – 1541.76	5.267E-24
023 – 010	1355	3564.72 – 3657.05	3.650E-22
122 – 010	620	3584.75 – 3751.75	3.505E-23
400 – 010	4	3623.87 – 3653.87	6.695E-25
005 – 010	848	4128.33 – 4228.24	2.354E-23
311 – 010	1018	4129.14 – 4228.62	3.420E-23
104 – 010	66	4146.09 – 4228.28	6.265E-25
023 – 000	1192	4268.37 – 4357.21	1.914E-22
122 – 000	724	4284.15 – 4454.93	4.328E-23
400 – 000	8	4325.56 – 4398.58	5.854E-25
213 – 100	63	4384.72 – 4423.50	1.407E-24

321 – 100	126	4429.78 – 4466.82	5.709E-24
114 – 001	306	4452.15 – 4508.72	1.304E-23
213 – 010	503	4757.08 – 4825.80	2.329E-23
Total	9534	598.26 – 4825.80	2.6928E-21

*: Line intensities are given for 99.29% $^{16}\text{O}_3$ abundance.

Table 3. The list of new $^{16}\text{O}_3$ ozone bands in the GEISA-2020 release. Num. Tra.: Number of transitions.

Recent *ab initio* calculations by Tyuterev *et al.* [29], using the dipole moment functions of [44], have suggested that absolute intensities of the strongest bands in most of spectroscopic databases [1], [2], [33] should be corrected to achieve the consistency between microwave and infrared 5 μm and 10 μm spectral ranges. The recommended intensity scaling corrections with respect to the HITRAN2016 data were +3%, +3% and +3.5% for the dominant rotational, ν_3 and $\nu_1+\nu_3$ bands. These corrections have been found in agreement [29] with accurate Stark-effect measurements and the JPL catalogue based on these measurements in the microwave, as well with the most recent infrared FTS measurements by Barbe *et al.* [29], and by Jacquemart *et al.* (unpublished data) for 5 μm and 10 μm . This was also consistent with DRL FTS measurements in the 10 μm range reported by Wagner *et al.* (unpublished data) within less than 1% for strong lines. The GEISA-2015 data being somewhat different from HITRAN2016, particularly in the 5 μm range, the *ab initio* scaling factors of [29] with respect to GEISA-2015 were thus adapted to the three corresponding strongest bands as given in #: internal data identification set to **T19**

Table 4.

Spectral Region	Band	Intensity scaling factor with respect to GEISA- 2015	Nb affected lines [#]
Microwave	000-000	1.030	5493
10 μm	001-000	1.027	7498
5 μm	101-000	1.015	6878

[#]: internal data identification set to **T19**

Table 4. Scaling factors applied to $^{16}\text{O}_3$ intensities and number of affected lines in GEISA-2020 for the three spectral regions considered.

In total, 81773 (sum of **M19** and **T19**) transitions of $^{16}\text{O}_3$ were inserted in GEISA-2020 that correspond to the integrated absorption (sum of line intensities) $S_v = 1.5983\text{E-}17$ cm/molecule at 296 K.

(ii) *Ozone isotopic species*

A significant update was also carried out for the ozone isotopologues $^{16}\text{O}^{16}\text{O}^{18}\text{O}$, $^{16}\text{O}^{16}\text{O}^{17}\text{O}$, and $^{16}\text{O}^{17}\text{O}^{16}\text{O}$, using linelists obtained from the analyses of Reims FTS spectra [45], [46] with the internal data identification set to **T19**.

For the asymmetric ^{18}O enriched isotopologues $^{16}\text{O}^{16}\text{O}^{18}\text{O}$ the 49148 lines of thirteen bands in the spectral range 961 – 3164 cm^{-1} are concerned by these changes with the

integrated band intensity (sum of all lines) $S_v = 6.101 \cdot 10^{-20}$ cm/molecule at 296 K for the natural isotopic abundance [33]. Six bands are new with respect to GEISA-2015: these are $2\nu_2$ and the second dyad bands $\nu_2+\nu_3/\nu_1+\nu_2$ in the range 1342 – 1894 cm^{-1} , as well as $3\nu_3/\nu_1+2\nu_3/2\nu_1+\nu_3$ in the range 2930 – 3164 cm^{-1} . The contributions for new bands of 668 are collected in **Table 5**.

Band	Num. Tra.	Region / cm^{-1}	S_v^* / cm/mol
020 – 000	505	1342.895 – 1398.807	7.706E-25
011 – 000	2474	1644.367 – 1720.541	2.033E-22
110 – 000	4188	1663.383 – 1894.671	6.376E-23
003 – 000	1562	2930.525 – 3011.532	4.316E-22
102 – 000	3241	2965.774 – 3123.188	1.221E-22
201 – 000	1940	3103.035 – 3164.883	3.782E-23

* Line intensities are given for the natural abundance of 0.00398194.

Table 5. New bands of ozone $^{16}\text{O}^{16}\text{O}^{18}\text{O}$ species [46] in GEISA-2020.

Another seven bands of $^{16}\text{O}^{16}\text{O}^{18}\text{O}$ were updated with respect to the previous GEISA release. This includes the first dyad ν_1/ν_3 (961 – 1187 cm^{-1}) and the first triad bands $2\nu_3/\nu_1+\nu_3/2\nu_1$ (1897 – 2274 cm^{-1}), where the resonance interaction parameters have been predicted by contact transformation method using MOL_CT code [47] from the potential energy surface in order to separate resonance contributions in the A-type and B-type sub-bands [46]. The line position and intensity values were updated for 4076 transitions of cold $\nu_1+\nu_2+\nu_3$ (2701 – 2767 cm^{-1}) and of hot $\nu_1+\nu_2+\nu_3-\nu_2$ (2015 – 2084 cm^{-1}) bands.

Four bands of ^{17}O enriched species were updated according new analyses of Ref. [45]. For the asymmetric $^{16}\text{O}^{16}\text{O}^{17}\text{O}$ isotopologue 2135 transitions of the $\nu_1+\nu_3$ (2045 – 2121 cm^{-1}) were updated, whereas for the symmetric $^{16}\text{O}^{17}\text{O}^{16}\text{O}$ species 2148 transitions of two fundamentals ν_1/ν_3 (968 – 1054 cm^{-1}) and the strongest combination band $\nu_1+\nu_3$ (2029 – 2101 cm^{-1}) were updated. The integrated absorption of these inserted lines of ^{17}O enriched species corresponds to $S_v = 5.768\text{E-}21$ at 296 K for the natural isotopic abundance.




Figure 2 Overview of the updated $^{16}\text{O}_3$ lines added to GEISA 2020 (red circles), compared to the previous O_3 linelist of GEISA2015 (blue diamonds).

2.1.4 N_2O (molecule 4)

Since GEISA-2011 (no update in the GEISA-2015 version), two contributions have been studied. The first one comes from the work of Tashkun *et al* [48], [49]. This update concerns lines with intensities at least 3 orders of magnitude lower than those in the former database. With such low values of the intensities, it is difficult to make an evaluation of the line list with the SPARTE chain. Combined to the fact, as mentioned in HITRAN 2016, that “*For the line intensities from these line lists, the uncertainty code 3 (> 20%) is adopted.* », we have decided not to retain these data for GEISA-2020 and to wait for eventual new data in 2021.

Finally, only a small update has been performed: line intensities, Einstein-A coefficients, self- and air-induced broadening and shift parameters of the R-branch lines of the $00^0_2-00^0_0$ band of $^{14}\text{N}^{14}\text{N}^{16}\text{O}$ have been updated using the values in references [50], [51] and [52]. All the lines have been replaced, with a pressure shift that has been kept unchanged from former version of GEISA.

The corresponding internal code J has been set to [L19](#).

2.1.5 CO (molecule 5)

The new CO line list originates from the work of Li *et al.* [53]. In this paper, a small incompatibility issue between the definition of potential curve parameters in the LEVEL 8.0 program [54] and in the work of Coxon *et al.* [55] causes drifts in line positions of up to 0.03 cm^{-1} for high J lines of the high overtone and hot bands. This issue has been fixed, and the calculated line positions now agree very well with the experimental values. A few other minor changes to the original Li *et al.* line list [53] have been discussed and described in

HITRAN2016 paper [2] and are also adopted here, resulting in an identical CO line list between HITRAN2016 and GEISA-2020. It corresponds to a total of 5381 lines with 3911 that were already with the former database.

The internal GEISA code for the data identification has been set to *L19*.

2.1.6 *CH₄ (molecule 6)*

The new list of CH₄ lines in the 5300-5850 cm⁻¹ range of the tetradecad, based on the work of Nikitin *et al.* [56]-[57], contains more than 15000 transitions, a significant part of which includes quantum identifications. Line positions and intensities have been retrieved by least square curve-fitting procedures from observed spectra recorded for different temperatures and pressures. The spectra in the 5695-5859 cm⁻¹ range [56] were recorded in Grenoble (France) at room temperature and 80K, by differential absorption spectroscopy (DAS) with a noise equivalent absorption of about $\alpha_{\min} \approx 1.5 \times 10^{-7}$ cm⁻¹ (except the 5718.8-5724.25 and 5792.36-5814.29 cm⁻¹ intervals). In the 5300- 5695 cm⁻¹ range [58] [57], long path (202, 602, 1604 and 1804 m) room temperature spectra of natural methane samples at different pressures were recorded with a Fourier transform spectrometer in Reims (France). One cold spectrum of enriched ¹²CH₄ at 80 K has been recorded in Jet Propulsion Laboratory (JPL, Pasadena, USA). This spectrum was useful for low-*J* line positions. Another cold spectrum at 80 K of enriched ¹³CH₄ from JPL was used to identify the isotopic species lines. Quantum assignments were made using the effective Hamiltonian based on high order contact transformation [59] and the effective dipole moment expressed in terms of irreducible tensor operators adapted to spherical top molecules. Self-broadening and air-broadening coefficient from [60], [61] were added to the final line list.

The corresponding internal code J has been set to *N19*.

New data for ¹³CH₄ were also implemented, based on the work of Starikova *et al.* [62]. This line list was generated using the line list retrieved from DAS spectra recorded at 80 K and room temperature. Assignments were transferred from the cold spectrum analysis of Ref [62]. The broadening coefficients were calculated according to the methodology used for the GOSAT project [32][34]. When no quantitative information was determined, a default value of the lower energy level of -0.9999 cm⁻¹ is given (cf. Appendix B) while corresponding default values for the air- and self- broadening coefficients are 0.08100 cm⁻¹atm⁻¹ and 0.06030 cm⁻¹atm⁻¹, respectively, and an air-shifting coefficient of -0.012 cm⁻¹atm⁻¹ is adopted.

The corresponding internal code J has been set to *S19*.




Figure 3: Comparison of $^{13}\text{CH}_4$ lines between 5800 and 6200 cm^{-1} involved in GEISA-2015 and GEISA-2020.

2.1.7 O_2 (molecule 7)

No major update was made for the O_2 molecule, the main reason being that recent studies mainly considered non-Voigt profiles for the line-shape determination. However, the work from Tran *et al.* [64] in the $a^1\Delta_g - X^3\Sigma_g^- (0 - 0)$ band of O_2 centered at 1.27 μm , and tested by comparison with high-resolution ground-based atmospheric TCCON measurements, highlighted systematic errors using air-broadened line-shift parameters from the GEISA-2015 or HITRAN 2016 databases. Therefore, we decided to partially update the GEISA-2020 database, considering only the new line intensities, line positions and air-broadened line-shift parameters determined in the work of Tran *et al.* [65] (cf. section 3.1.2 for an illustrative validation of these O_2 data using TCCON measurements). In this study, parameters for 85 air-broadened transitions, from 7784.8 to 7915.8 cm^{-1} , were retrieved from spectra measured by a cavity ring down spectrometer referenced to a frequency comb, for a large range of pressure values, from 6.7 to 94 kPa (50 to 700 Torr). The data published in the work of Tran *et al.* were originally determined using the quadratic speed dependent Nelkin-Ghatak (qsdNG) profile, but for the GEISA-2020 update the authors also refitted these lines using Voigt profiles from which the new air-broadened line-shift parameters are derived (unpublished data).

The corresponding internal code J has been set to *T19*.

2.1.8 *SO₂ (molecule 9)*

The ³²S¹⁶O₂ empirical list is mainly based on the experimental energy levels recently derived in [66] and variational intensities presented in [67]. In addition to experimental energy levels derived in [66], the new experimental energy levels obtained from the analysis of Fourier transform spectra of SO₂ [68] were also used. In parallel with pure experimental energy levels, the predicted energy levels obtained from the effective Hamiltonian calculations performed in [66] were used for evaluation of the line positions.

The spectrum between 0.025 and 4159 cm⁻¹ consists of 549,200 transitions with 1.0e⁻²⁷ cm/molecule intensity cutoff. Most part of the line positions (63%) have an estimated accuracy between 0.0001 and 0.001 cm⁻¹ in accordance with the uncertainties of the corresponding experimental energy levels reported in [66]. For 185,637 transitions (34%) derived from the calculated energy levels the positions accuracy corresponds to the 0.001 - 0.01 cm⁻¹ error range. In case of the (110)-(000), (011)-(000), (101)-(000), (111)-(010) bands the calculated intensities [67] are substituted by the recent experimental data [69],[70] when available. We estimate the quality of variational intensities as 5-10% on average.

The resulting empirical list of ³²S¹⁶O₂ transitions was compared both with the HITRAN 2016 database and a recent study [71] performed by the AMES group at NASA where the similar empirical lists for SO₂ isotopologues have been reported based on the experimental energy levels [66] and new variational calculations. It turned out that the HITRAN 2016 database among 72,469 transitions in total attributed to ³²S¹⁶O₂ molecule contains a large number of simulated data. Disagreements between GEISA and HITRAN's line positions amount up to 0.01 - 0.16 cm⁻¹ for 3308 lines. The calculated intensities [67] adopted in our empirical list also may differ from the HITRAN data by up to 40% for relatively strong lines and disagree by up to 60 -100% for weakest lines. A comparison of our list with HITRAN and Ames data [71] is shown in Figure 4. It is obvious from the figure that the presently generated empirical list for ³²S¹⁶O₂ molecule includes a considerably large number of lines, including very strong ones (up to 5e⁻²⁰ cm/molecule).




Figure 4: Comparison of the new empirical list for $^{32}\text{S}^{16}\text{O}_2$ molecule with the HITRAN 2016 and Ames [71] data.

The internal GEISA codes for the data identification have been set to:

- **1N9:** positions are determined from the upper and lower experimental energy levels, intensities are calculated (variational) from [67].
- **2N9:** positions are determined from the upper and lower experimental energy levels, intensities are experimental from [69].
- **3N9:** positions are determined from the upper and lower experimental energy levels, intensities are experimental from [70].
- **4N9:** one of the two energy levels (or both) used for position determination is predicted by the effective Hamiltonian calculation, intensities are calculated (variational) from [67].
- **5N9:** pure experimental value is taken for most accurate position.

2.1.9 NO_2 (molecule 10)

For nitrogen dioxide the previous GEISA compilation included over 100,000 $^{14}\text{N}^{16}\text{O}_2$ lines covering 0–3074 cm^{-1} . The new list includes 185,965 lines and covers the 0-4776 cm^{-1} spectral range. This new list includes lines from both the $^{14}\text{N}^{16}\text{O}_2$ and $^{15}\text{N}^{16}\text{O}_2$ isotopic species.

In two recent studies [72], [73], a list of line positions, line intensities and line shape parameters of NO_2 covering the 0 to 4750 cm^{-1} spectral range was generated. Except for the far infrared and the 13.3 μm regions, all bands in all spectral regions are significantly affected by this update. For the 6.2 μm and 3.4 μm spectral regions, the new list involves the main

cold bands (ν_1 , $2\nu_2$ and ν_3 and $\nu_1 + \nu_3$, respectively) together with their associated first hot bands (originating from the (0,1,0) first vibrational state), and, whenever possible, hotter bands (involving the (1,0,0), (0,2,0) and (0,0,1), (1,1,0), (2,0,0) or (0,0,2) states as lower states). Also several weak cold bands ($\nu_1 + \nu_2$, $\nu_2 + \nu_3$, $2\nu_1$, $2\nu_2 + \nu_3$, $2\nu_3$, $2\nu_2 + \nu_3$, $\nu_1 + \nu_2 + \nu_3$, $2\nu_1 + \nu_3$, and $3\nu_3$) were added for the first time in the GEISA database. Finally we included the ν_3 [74] and $\nu_1 + \nu_3$ [75] bands for $^{15}\text{N}^{16}\text{O}_2$ which is the second most abundant isotopologue of NO_2 [22]. These new lists were generated using existing literature line positions or intensity parameters. When necessary, these parameters were updated using experimental data issued from high resolution Fourier transform spectra recorded at SOLEIL at 296K for this purpose [72], [73]. Also, the line broadening parameters were computed using the line shape parameters achieved for the ν_3 band by Benner *et al.* [76].

Figure 5 gives an overview of cross sections calculated from the list of NO_2 line parameters in GEISA-2020 for the 1000- 4780 cm^{-1} spectral range and compares it to the status of HITEMP-2019 [77] and of HITRAN-2016 or GEISA-2015.




Figure 5: Calculated absorption cross sections of NO_2 at 296 K using line lists from HITRAN-2016, GEISA-2015, HITEMP-2019 and GEISA-2020.

The validation of this new update, which concerns the 1153- 4776 cm^{-1} region, was performed through a careful comparison between computed and observed spectra using laboratory spectra recorded either during recent investigations [73], [72] or during previous analyses of the literature (see Refs. [73], [72] for detailed references). In parallel, similar inter comparisons at $T=296\text{K}$ were also performed using the most recent versions (HITRAN-2020 and HITEMP-2019) of the HITRAN and HITEMP databases [77]. The results of these checks

prove that, for each vibrational bands, the individual lines in GEISA-2020 are more precise than those included in HITRAN-2020 or HITEMP-2019 (or those of previous versions). Concerning the number of vibrational bands, the GEISA linelist is also more extended than HITRAN-2020, but less than HITEMP-2019 which is generated for hot temperature conditions and includes lines involving very high rotational quantum numbers (up to $N=100$) and a significant number of additional hot bands when compared to GEISA-2020.

The internal GEISA code for the data identification has been set to **P19**.

2.1.10 NH_3 (molecule 11)

GEISA-2020 includes, for the first time, a line list at $3.0 \mu\text{m}$ for the first-most abundant isotopologue of ammonia, $^{14}NH_3$. The $2\nu_4$ vibrational band for this isotopologue was added using a high resolution Fourier transform investigation performed at $3.0 \mu\text{m}$ by Maaroufi *et al.* [78]. Absolute line intensities and self-broadening and shift coefficients were measured at room temperature, for rovibrational lines ranging from $1 \leq J \leq 11$ and $0 \leq K \leq 11$, and located in the spectral range $3078.80407 - 3532.50519 \text{ cm}^{-1}$. This represents a total of 298 entries. The lines were fitted with a single spectrum non-linear least squares fitting procedure with a Voigt profile suited to fit the experimental line shapes without the need of taking into account line mixing effects, in the pressure range used. Average absolute accuracies of the measurements are estimated to $\pm 4\%$ for line intensities and self-broadening coefficients and to $\pm 13\%$ for self-shift coefficients.

The J and K dependences of the self-broadening coefficients (γ_0) were observed and modeled using an empirical polynomial expression suggested by Nemtchinov *et al.* [79]:

$$\gamma_0(m, K) = \beta_0 + \beta_1 m + \beta_2 K + \beta_3 m^2 + \beta_4 K^2 + \beta_5 mK$$

where $m = | -J, J, J+1 |$ for P, Q and R branches, and β_i are the polynomial coefficients. The polynomial expression reproduced the observed self-broadening coefficients within an average accuracy of 9.6%, leading to accurate coefficients constants (β_i) for the $2\nu_4$ band of $^{14}NH_3$ [78].

The internal GEISA code for the data identification has been set to **M19**.

2.1.11 PH_3 (molecule 12)

The PH_3 line list in the $2733\text{-}3660 \text{ cm}^{-1}$ range is based on an improved analysis of positions and intensities of phosphine spectral lines from Nikitin *et al.* [80]. Some 5768 positions and

1752 intensities have been modeled with RMS deviations of 0.00185 cm^{-1} and 10.9%, respectively. Based on an *ab initio* potential energy surface, the full Hamiltonian of phosphine nuclear motion was reduced to an effective Hamiltonian using a high-order Contact Transformations method adapted to polyads of symmetric top AB_3 -type molecules with a subsequent empirical optimization of parameters. The line list contains more than 2000 new ro-vibrational assignments for all 13 vibrational octad sublevels. A comparison of this line list with experimental spectra of PNNL [81] showed that the new set of line parameters from this work permits better simulations of observed cross-sections than HITRAN 2016 linelist [2]. Vibrational assignments were added to current GEISA line list.

The internal GEISA code for the data identification has been set to *N17*.

2.1.12 HNO_3 (molecule 13)

A new investigation of laboratory spectra of nitric acid in the $7.6 \mu\text{m}$ region was performed using a suitable theoretical model which involves six interacting states, instead of two in the previous computation [82]. In this way, an improved set of line positions and intensities have been generated for the $7.6 \mu\text{m}$ spectral region. This new list, which includes also the $\nu_3+\nu_9-\nu_9$ for the first time was validated using atmospheric spectra recorded by the Michelson Interferometer for Passive Atmospheric Sounding (MIPAS) limb emission radiances in the 11 and $7.6 \mu\text{m}$ domains [83]. Table 6 describes the new list for HNO_3 and compares with the list included in GEISA-2015.

Table 6: HNO_3 line parameters in the $7.6 \mu\text{m}$ region [NB is the number of lines, σ_{\min} and σ_{\max} (cm^{-1}) are the lower and upper limits, S_{\min} and S_{\max} are the smallest and largest line intensity ($\text{cm}^{-1}/\{\text{molecule cm}^2\}$ at 296 K) and S_{tot} is the sum of the line intensities]

Band	NB	S_{tot} (10^{-18})	σ_{\min}	σ_{\max}	S_{\min} (10^{-25})	S_{\max} (10^{-21})
<i>GEISA-2020 database</i>						
ν_3	16408	24.94	1252.010	1387.081	4.910	32.0
ν_4	18105	9.834	1238.929	1387.561	4.020	21.4
$2\nu_6$	2451	0.1194	1243.465	1348.275	4.624	3.660
$\nu_5+\nu_9$	13817	0.7163	1246.929	1390.071	2.081	3.543
$\nu_7+\nu_8$	11125	0.7615	1246.422	1395.679	2.314	5.017

$3\nu_9$	13894	1.177	1233.107	1388.497	4.582	2.378
Sum		37.55				
$\nu_3+\nu_9-\nu_9$	12106	1.408	1271.050	1394.899	5.285	1.798
GEISA-2015 database						
ν_3	21308	25.37	1098.376	1387.849	1.037	31.33
ν_4	19584	12.78	1229.867	1387.561	1.037	18.67
Sum		38.15				

The Internal GEISA code for these data identification has been set to [P19](#).

2.1.13 Hydrogen Halides HF, HCl, HBr, HI (resp. molecule 15, 16, 17 and 18)

The new hydrogen halides line lists originate from refs. [84] and [85], with higher overtone bands removed due to abnormalities when intensities were extrapolated outside of experimental temperature and pressure ranges. Except for the 2-0 band of HCl, the HX line lists are identical to those in HITRAN2016. For P10 to R10 lines of the HCl 2-0 band, line intensities, Einstein A-coefficients and self-induced broadenings have been updated with reference [86], whilst air-induced broadenings and shifts have been updated with reference [87].

The Internal GEISA code for these data identification has been set to [L19](#).

2.1.14 OCS (molecule 20)

The 5 μ m spectral region is the most often used to monitor the most abundant sulfur-containing species OCS [88]. In the last years, Cavity Ring Down Spectroscopy made the near infrared spectral range more and more attractive, especially for the in situ detection of trace gases [89].

First, the line list used in the HITRAN-2016 database [2] has been adopted in the new GEISA-2020 release. This line list was derived from Fourier transform spectra of OCS recorded in the 6170 – 6680 cm^{-1} (dominated by the $3\nu_1+2\nu_3$ band) and 7700 – 8160 cm^{-1} (dominated by the $2\nu_1+3\nu_3$ band) spectral ranges using a femto/OPO laser absorption source and cavity enhanced spectroscopy [90]. It contains 3670 lines in the ranges 6484 – 6660 cm^{-1} and 7728 – 7821 cm^{-1} . The accuracies of the line positions and intensities (parameters A' and B', see appendix B) have been estimated to be at best 0.001 cm^{-1} and 10 to 20%, respectively.

Secondly, 104 new spectroscopic line parameters for the main isotopologue $^{16}\text{O}^{12}\text{C}^{32}\text{S}$ have been included in GEISA-2020. The update focuses on the broadening and shift coefficients in the $4\nu_2$ band of OCS perturbed by N_2 , O_2 and Air, measured and calculated at room temperature, in the spectral region $2087.4280 - 2130.7584 \text{ cm}^{-1}$. These data, which were derived from high resolution Fourier transform spectra, are rather accurate: the uncertainties can be estimated to be $\sim 3\%$ for the broadening coefficients and $\sim 10\%$ for the shift coefficients [91]. In addition, theoretical calculations of broadening and line shift coefficients were performed, using the semi classical Robert and Bonamy formalism that reproduces rather well the experimental m ($m = -J$ for $P(J)$ lines and $m = J + 1$ for $R(J)$ lines) quantum number dependence of the N_2 and O_2 broadening coefficients.

The Internal GEISA code for these data identification has been set to [A16](#).

2.1.15 C_2H_6 (molecule 22)

The data from Moazzen *et al.* [92] was received too late to be incorporated in GEISA-2015. For this reason, this work, which has already been included in HITRAN2016 [2], is now implemented in GEISA-2020.

The internal GEISA code for these data identification has been set to [M19](#).

2.1.16 C_2H_2 (molecule 24)

The GEISA-2020 list of acetylene has been significantly extended compared to the GEISA-2015 version [1] (see Figure 6). The changes concern four spectral regions. Three line lists in the far-infrared and infrared domains ($\Delta P = 0$ between 13 and 248 cm^{-1} , $\Delta P = 1$ between 389 and 893 cm^{-1} and $\Delta P = 6$ between 3738 and 3993 cm^{-1}) were generated based on the global model developed by Lyulin and Perevalov [93][94]. Here P is a polyad number defined as $P = 5V_1 + 3V_2 + 5V_3 + V_4 + V_5$ where $V_i = 1-5$ are the main quantum numbers of the C_2H_2 harmonic oscillator. The predictive ability of the model in the $\Delta P = 0, 1$ regions was checked experimentally in [95], [96] leading to line lists of calculated positions and intensities for the $\Delta P = 0$ region (5479 transitions) and for the $\Delta P = 1$ region (41021 transitions). A complete description of the line lists in $\Delta P = 0$ and 1 regions can be found in [95], [96]. For the $\Delta P = 6$ region around 3900 cm^{-1} , the strong Q branch around 3882 cm^{-1} as well as other weaker structures were missing in GEISA-2015. The 2 strong cold bands already present in GEISA have been updated and 8 bands have been added based on the global model [93][94] for line positions and intensities. A summary of updated/included bands is given in Table 7. Note that the vibrational assignment from the global model [93][94] (linear combination of vibrational

basis functions) has been reformulated in order to have a unique vibrational assignment for a series of rotational transitions.




Figure 6: Overview of the GEISA-2015 (solid red circles) and GEISA-2020 (open black circles) databases for acetylene $^{12}\text{C}_2\text{H}_2$ and $^{12}\text{C}^{13}\text{CH}_2$ isotopologues.

The fourth line list is the empirical database (EDB17, hereafter) elaborated by Lyulin and Campargue [97] for the wide $5850\text{--}9415\text{ cm}^{-1}$ near-infrared region. The EDB17 covers the range of the $\Delta P = 9\text{--}14$ series excluding the $6341\text{--}7000\text{ cm}^{-1}$ interval roughly corresponding to the $\Delta P = 10$ series. The EDB17 was constructed by gathering results of three studies by Fourier transform spectroscopy [98]–[100] and three studies by cavity ring down spectroscopy (CRDS) [101]–[103]. In the regions of low absorption, the sensitivity of the CRDS technique enabled the detection of weak lines with intensities on the order of $10^{-29}\text{ cm/molecule}$. The assignment of the experimental spectra relies in great part on the predicted data of the Acetylene Spectroscopic Databank (ASD) [94]. For each branch of a given band, the line positions and line intensities were calculated for all the transitions up to a maximum J value exceeding by 1 the largest J value of the observations in the considered branch. Such an approach enabled smoothing the measurement errors, completing the experimental line list by interpolation and extending slightly beyond the observations. Compared to the HITRAN2016 [2] and GEISA-2015 [1] lists in the region, the number of bands and lines was increased by more than a factor of ten (see Figure 6). As a result, the EDB includes a total of 10973 lines

belonging to 146 bands of $^{12}\text{C}_2\text{H}_2$ and 29 bands of $^{12}\text{C}^{13}\text{CH}_2$. For comparison the GEISA-2015 database in the same region includes 869 lines of 14 bands, all belonging to $^{12}\text{C}_2\text{H}_2$.

For all $^{12}\text{C}_2\text{H}_2$ and $^{12}\text{C}^{13}\text{CH}_2$ isotopologues, the vibrational assignment has been changed to $V_1 V_2 V_3 V_4 V_5 \ell_4 \ell_5$ from $V_1 V_2 V_3 V_4 V_5 \ell$, where ℓ is the sum of ℓ_4 and ℓ_5 . Indeed, the ℓ_4 and ℓ_5 quantum numbers are necessary to assign the various bands that include strong resonance interactions between ro-vibrational states [95]–[103] used to update the GEISA-2020 database. Based on the new vibrational assignment and the symmetry E/F of the rotational levels, the symmetries $+/-$ have been generated for all transitions of both isotopologues, and symmetry u/g have been generated only for symmetric $^{12}\text{C}_2\text{H}_2$ isotopologue. For all transitions, Einstein-A coefficients have been calculated from line intensities at 296 K using Eq. 20 of [104], state-independent weight $g_i = 1$ and 8 respectively for $^{12}\text{C}_2\text{H}_2$ and $^{12}\text{C}^{13}\text{CH}_2$ isotopologues, and state-dependent weight $g_s = 1$ or 3 for $^{12}\text{C}_2\text{H}_2$ (depending on symmetries) and $g_s = 1$ for $^{12}\text{C}^{13}\text{CH}_2$. Total internal partition functions [105] equal to 412.45 and 1656.2 at 296 K have been used for the $^{12}\text{C}_2\text{H}_2$ and $^{12}\text{C}^{13}\text{CH}_2$ isotopologues respectively. Smoothed values of air-broadening coefficients (Table 5 of [106]) already in previous GEISA editions have been used for new entries since no vibrational dependence of the widths is observed.

Vibrational Assignment		N	Sig_min	Sig_max	Sum_S	S_min	S_max	J_min	J_max	
Upper	Lower									
00110 1 0	00000 0 0	155	3738.098	3995.170	5.96E-20	1.3E-27	3.0E-21	0	53	GEISA-2015
01031 3-1	00010 1 0	263	3744.789	3968.111	5.80E-21	1.0E-27	1.4E-22	1	49	
01022 2-2	00001 0 1	251	3749.044	3958.870	1.18E-21	1.2E-27	3.6E-23	1	45	
01021 2-1	00000 0 0	153	3752.906	3980.295	5.54E-20	1.2E-27	2.7E-21	0	53	GEISA-2015
01022 2 0	00001 0 1	220	3756.621	3962.215	1.31E-21	1.2E-27	3.7E-23	1	45	
00120 2 0	00010 1 0	250	3770.399	3983.995	5.76E-21	1.2E-27	1.5E-22	1	48	
00111 1 1	00001 0 1	240	3772.294	3979.654	2.03E-21	1.3E-27	5.6E-23	1	45	
00111 1-1	00001 0 1	233	3780.389	3974.747	1.63E-21	1.2E-27	3.8E-23	1	46	
01031 1-1	00010 1 0	205	3794.438	3976.946	3.58E-21	1.4E-27	2.4E-22	1	40	
00120 0 0	00010 1 0	101	3817.941	3974.507	2.37E-21	2.8E-27	1.5E-22	1	35	

Table 7: Summary of bands updated or included in GEISA-2020 in the $\Delta P = 6$ region. The first column corresponds to the vibrational labeling used for series of transitions (bands). N is the total number of transitions belonging to a given band. Minimum and maximum wavenumbers of calculated transitions belonging to the band are given in cm^{-1} , Sum_S is the sum of calculated line intensities (see text) in $\text{cm}^{-1}/(\text{molecule}\cdot\text{cm}^{-2})$ at 296K for natural abundances. S_min and S_max are respectively the minimum and maximum calculated line intensities in the band. Minimum and maximum values of J in the band are given.²

² Note: The vibrational labeling for upper and lower state corresponds to $V_1 V_2 V_3 V_4 V_5 \ell_4 \ell_5$. GEISA-2015 refers to the two strongest bands already present in GEISA-2015 but updated in GEISA-2020.

For self-broadening coefficients, an improved set of parameters (given in Table 8) based on the work performed in [106] has been used for all C₂H₂ transitions in GEISA-2020. For J values greater than 36 self-broadening coefficients have been fixed to 0.078 cm⁻¹/atm. Default values have been used for the temperature dependence of air-broadening coefficients (equal to 0.75) and for air-shifting coefficients (equal to -0.001 cm⁻¹/atm).

m	$\gamma_{self}(296K)$	m	$\gamma_{self}(296K)$	m	$\gamma_{self}(296K)$
1	0.2055	13	0.1438	25	0.1088
2	0.1925	14	0.1407	26	0.1059
3	0.1823	15	0.1376	27	0.1030
4	0.1744	16	0.1347	28	0.0999
5	0.1683	17	0.1319	29	0.0967
6	0.1635	18	0.1291	30	0.0934
7	0.1598	19	0.1264	31	0.0898
8	0.1568	20	0.1225	32	0.0861
9	0.1543	21	0.1198	33	0.0831
10	0.1520	22	0.1171	34	0.0812
11	0.1499	23	0.1143	35	0.0800
12	0.1471	24	0.1117	36	0.0780

Table 8: Self-broadening coefficients used in GEISA-2020.

The internal GEISA code for the data identification has been set to [L20](#) and [J20](#).

2.1.17 C₂H₄ (molecule 25)

The ¹²C₂H₄ update in GEISA-2020 was generated using data for the ν_{10} , ν_7 and ν_4 bands from the corrected line list of Ref. [107], however ignoring lines having an intensity smaller than 10⁻³⁰ cm⁻¹/(molecule cm⁻²) at 296 K. Note that errors in the degeneracies provided for the rotational levels in the 10¹ and 7¹ vibrational levels in the linelists of Refs. [107] and [108] were corrected (I. Gordon is acknowledged for bringing them to our attention). The positions and intensities of the ν_{12} band lines were taken from GEISA-2015, excluding 187 lines having an intensity smaller than 10⁻³⁰ cm⁻¹/(molecule cm⁻²) at 296 K to be consistent with the information provided for the 3 other bands. For all the lines, the self-broadening coefficients were calculated using the empirical expressions reported in Ref. [109], the air-broadening coefficients were generated as described in [110], and the temperature dependence exponent n was set to 0.76, determined as described in Ref. [108].

The internal GEISA code for the data identification has been set to [A19](#).

2.1.18 GeH₄ (molecule 26)

Germane (GeH₄) is a molecule that has been detected in giant gas planets like Jupiter and Saturn [111], [112]. The current [Juno](#) mission has renewed its spectroscopy interest especially

in the 4.5 – 5.0 μm window where an accurate model is needed for radiative transfer computations. A first complete analysis of the ν_1/ν_3 stretching dyad region for all natural five isotopologues ($^{70}\text{GeH}_4$, $^{72}\text{GeH}_4$, $^{73}\text{GeH}_4$, $^{74}\text{GeH}_4$ and $^{76}\text{GeH}_4$) was recently performed by Boudon *et al.* [113] for line positions and absolute line intensities. This study led to a set of 32,378 calculated lines, covering the range 1929 cm^{-1} to 2266 cm^{-1} , that have been introduced in the new GeCaSDa [114][115] database in Dijon and in the present version of GEISA. The former version of GEISA only contained lines for $^{74}\text{GeH}_4$.

The Internal GEISA code for the data identification has been set to **B19**.

2.1.19 NO^+ (molecule 45)

Nitrosylium, NO^+ , was detected in the Earth's ionosphere as an important trace species [116] and tentatively in the interstellar medium [117]. The rovibrational transitions have been introduced into GEISA-2020, based on version 2 of the CDMS [118] entry (tag 030512). Rotational transition frequencies [117],[119] were fitted together with rovibrational data [116],[120]. The dipole moments are 0.368 D and 0.380 D in $v = 0$ and 1, respectively [121].

The Internal GEISA code for the data identification has been set to **M18**.

2.1.20 CF_4 (molecule 49)

Carbon tetrafluoride (CF_4) is a strong greenhouse gas of both natural and anthropogenic origin whose concentration is monitored using, among other techniques, solar occultation measurements from space [122]. It has been shown that, up to now, the line-by-line databases were quite incomplete because of the lack of hot band lines for this quite heavy species [123]. Recently, a systematic study of many low-lying energy levels was undertaken by the SOLEIL/AILES and Dijon groups to deal with this problem, resulting in a global analysis that enabled to simulate the main hot bands in the strongly absorbing ν_3 C-F stretching region [124]. Moreover, line positions and intensities for purely rotational transitions in the $\nu_3 = 1$ state were recorded and fitted in the far infrared (THz) region, thanks to synchrotron radiation [125]. These last two studies are used for the present update of $^{12}\text{CF}_4$ lines in GEISA. We also adapted the line intensity thresholds in each spectral region to avoid unwanted inaccurate extrapolated lines of very low intensity. These line lists contain 258,208 lines and are included in the new TFMeCaSDa [115], [126] Dijon's database, as well as in the present version of GEISA.

The Internal GEISA code for the data identification has been set to **B19**.

2.1.21 HONO (new molecule 53)

Using spectroscopic data collected in the literature, a linelist of positions and intensities has been generated for the 11 μm region of HONO [127] which correspond to the ν_6 bands located at 790.117 and 851.943 cm^{-1} for the Cis- and Trans- isomers, respectively.

For the computation of line positions, the vibrational energies and rotational constants quoted in Refs. [128] and [129] were used for the upper and lower state energies. On the other hand the line intensities were calibrated relatively to band intensities measured by Kagann and Maki [130]. The problem of the uncertainties associated to these intensities was discussed in details in Ref. [127]. Indeed, in usual laboratory conditions nitrous acid exists only in the form of an equilibrium mixture with other species like NO, NO₂ and H₂O, together with smaller quantities of N₂O₃, N₂O₅, and HNO₃ [130]. Due to these difficulties, one may estimate an overall uncertainty on the absolute intensities between 20% to 30%, while, on the relative scale, one may expect the intensities of the Trans- and Cis species at 296K to be rather consistent, with an uncertainty of 5% on average, and of 10% for the weaker lines.

To our knowledge, line shape parameters are absent in the literature for HONO. As the permanent dipole moment of Trans-HONO and Cis-HONO have values which are rather similar to those of water, it was decided to use for the HONO line shape parameters “a priori” values similar to those of water. More explicitly,

$$\gamma_{\text{Air}}=0.1 \text{ cm}^{-1}/\text{Atm}, \gamma_{\text{Self}}=0.4 \text{ cm}^{-1}/\text{Atm}, \text{ and } n_{\text{Air}}=0.7$$

were implemented for the air-broadened half width, self-broadened half-width and for the n-temperature dependent coefficient, respectively.

Species	Nb	Sig_min	Sig_max	Int_Max [#] $\times 10^{-17}$	Total_A [#] $\times 10^{-17}$	Total_B [#] $\times 10^{-17}$	Total_Int [#] $\times 10^{-17}$
Trans- HONO	7621	724.39	838.57	0.166	1.218	0.0	1.218
Cis-HONO	18420	722.53	996.28	0.148	1.115	0.1338	1.249
All	26041						2.467

[#] All intensities in $\text{cm}^{-1}/(\text{molecule} \cdot \text{cm}^{-2})$ at 296K.

Table 9: The HONO linelist in GEISA. The following abbreviations are used: Nb: number of lines; Sig_Min/Sig_Max; minimum and maximum sigma values (in cm^{-1}); Int_Max: Maximum intensity; Total-A/Total_B: sum of the intensities for A- type and B- type

transitions, respectively; Total int: sum of the individual intensity for the Trans and Cis species, $\text{TransInt}(\nu_4, T=296\text{K})$ and $\text{CisInt}(\nu_4, T=296\text{K})$, respectively.

The minimum of intensity retained is $0.5 \cdot 10^{-24} \text{ cm}^{-1}/(\text{molecule}\cdot\text{cm}^{-2})$. The internal GEISA code for the data identification has been set to **PI8**.

2.1.22 COFCl (new molecule 54)

Carbonylchlorofluoride (COFCl) is a new molecule identified as molecule 54 in the GEISA notation. A list of line positions and of line intensities was implemented for the ν_1 , ν_2 , and ν_3 fundamental bands of the $^{12}\text{C}^{16}\text{O}^{35}\text{FCl}$ and $^{12}\text{C}^{16}\text{O}^{37}\text{FCl}$ isotopologues of carbonylchlorofluoride, located at 5.3, 9.1, and 13.1 μm , respectively [131]. In addition, for the most abundant isotopologue $^{12}\text{C}^{16}\text{O}^{35}\text{FCl}$, this line list includes also the contributions from the first two associated hot bands. The parameters included in this database were generated by combining the results of previous experimental analyses [132] and *ab initio* calculations [133]. To our knowledge, no line broadening or shifting parameters data exists in the literature for COFCl. Therefore, default values for the self and air-broadening coefficients of $0.1 \text{ cm}^{-1} \text{ atm}^{-1}$ and temperature dependence of $n=0.75$ are given in the present database. This COFCl linelist was generated in order to improve the quality of remote sensing of the atmosphere in the mid-IR. Analyses of atmospheric solar occultation spectra measured by the JPL MkIV interferometer [131] show that the new line list not only improves the quality of retrievals of COFCl, but also of several other overlapping gases.

The internal code for the data identification has been set to **PI9**.

2.1.23 CH₃I (new molecule 55)

Methyl iodide (CH₃I) is a new molecule identified as molecule 55 in the GEISA notation. The line list in GEISA involves the ν_6 and $2\nu_3$ interacting bands at 11 μm . For the computation of the line positions and intensities, the hyperfine structure due to the iodine nuclear quadrupole moment was accounted for explicitly, together with the vibration-rotation resonances. The work dealing with line positions and line intensities is described in Refs. [134] and [135], respectively. In addition, detailed self- N₂ [136] and O₂ [137] line shape parameters were measured.

Band	Number of lines	Band intensity	Int _{Max}	Sigma Min	Sigma Max	J _{Max}	K _{Max}
		$\times 10^{-19}$	$\times 10^{-23}$				
ν_6	49603	12.3	220	693.02	1125.25	88	24

2v ₃	20688	0.181	3.3	985.67	1076.68	88	21
Total	70291	12.5					

Table 10: Linelist for Methyl iodide (¹²CH₃I) in the 11 μm region for GEISA-2020. “Hyp” and “No Hyp” identify the number of lines generated when accounting and not accounting for the hyperfine structure, respectively. “Band intensity” and “IntMax” (maximum value of the computed line intensities) are in cm⁻¹/(molecule cm⁻²). “Sigma Min” and “Sigma Max” are the minimum and maximum values of the computed line positions (in cm⁻¹), respectively.

The Internal GEISA code for the data identification has been set to **P19**.

2.1.24 CH₃F (new molecule 56)

Methyl fluoride (CH₃F) is present as traces in the Earth atmosphere. A line list for the v₆ band around 1200 cm⁻¹ is now included for the first time in GEISA, based on the work of Papoušek *et al.* [138] for line positions, on the work of Jacquemart and Guinet [139] for line intensities and self-broadening coefficients, and on the work of Barbouchi Ramani *et al.* [140] for N₂-broadening coefficients.

The internal GEISA code for data identification has been set to **J19**.

2.1.25 RuO₄ (new molecule 57)

Ruthenium tetroxide is a molecule of importance for nuclear power plant security and other industrial applications (see [141], [142], [143] and references therein). In natural abundance, it possesses 7 isotopologues, due to the different ruthenium isotopes (we do not consider here oxygen isotopes): ⁹⁷RuO₄, ⁹⁸RuO₄, ⁹⁹RuO₄, ¹⁰⁰RuO₄, ¹⁰¹RuO₄, ¹⁰²RuO₄ (the main one) and ¹⁰⁴RuO₄. There are also two radioactive, short-lived, isotopologues, ¹⁰³RuO₄ and ¹⁰⁶RuO₄. In Ref. [141], the v₃ band for all observable isotopologues was analyzed, and the parameters for the radioactive ones were extrapolated. Absolute line intensities were also analyzed in Ref. [143] for ¹⁰²RuO₄. Assuming that the dipole moment derivative parameter does not change significantly for the different isotopologues, this allowed to calculate line lists for all 9 isotopologues. These line lists contain 30,205 lines and are included in the new RuCaSDa [115], [144] Dijon’s database, as well as in the present version of GEISA.

For the broadening parameters, in general, a default value is given (as 0.1 cm⁻¹/atm for the air). In this update, and to specifically indicate that these values are unknown, users have to take care of the fact that the air and self-broadening have been set to respectively -.9999 and -9.9999.

The Internal GEISA code for the data identification has been set to **B19**.

2.1.26 $H_2C_3H_2$ (new molecule 58, Allene, isomer of C_3H_4)

$H_2C_3H_2$ (Allene, isomer of C_3H_4) is the last new molecule added to GEISA-2020, and identified as molecule 58 in the GEISA notation. This linelist includes the ν_{11} far infrared band near 352 cm^{-1} as well as the ν_{10}/ν_9 region peaking at 845 cm^{-1} . It is a result of the search for allene (also called propadiene) in Titan's atmosphere using the CIRS (Composite InfraRed Spectrometer) instrument on board the Cassini-mission [145] that was positively concluded by a clear detection using high resolution observations from the ground [146]. No allene linelist was present neither in HITRAN nor in GEISA but the linelist of propyne (C_3H_4) which is an isomer of allene has been provided by GEISA for a long time [1].

Coustenis *et al.* [147] started to investigate the detectability of allene in Titan's atmosphere using spectroscopic parameters by Chazelas *et al.* [148] for the ν_{10} band and from Pliva and Kauppinen for ν_{11} [149]. Line intensities could be obtained from band intensity measurements by Koga *et al.* [150]. Later Jolly *et al.* [151] used the same spectroscopic study but also added spectroscopic parameters obtained by high resolution studies by Nissen *et al.* [152] and Hegelund *et al.* [153] which also include spectroscopic parameters for the hot bands. The hot band contribution is particularly important at room temperature and needs to be included in order to compare the calculated line lists with room temperature spectra. No new high-resolution measurements were necessary for establishing this new linelist. It is essentially based on the sub-band analysis from previously mentioned authors but has also been validated against new measurements.

As described in Lombardo *et al.* [145], calculated band profiles have been compared to room and high temperature cross section measurements recorded at 0.08 cm^{-1} resolution in the ν_{10}/ν_9 wavenumber range [154]. All sub-band intensities could be fitted using those intensity calibrated measurements. Concerning the far infrared region where the ν_{11} band of allene can be observed, room temperature spectra were recorded at the resolution of 0.1 cm^{-1} at the SOLEIL-AILES synchrotron beamline as described in Jolly *et al.* [151]. Band intensities were found to be in agreement with Koga *et al.* [151]

This linelist that we have adopted is the same that was used for the search and detection of allene on Titan. It provides for each line the position, the intensity and the lower energy but does not provide any quantum assignment. Pressure shift is fixed to $0.1\text{ cm}^{-1}/\text{atm}$ and self-broadening to 0.71 using the same values as for allene's isomere C_3H_4 (propyne).

The internal GEISA code for the data identification has been set to **J19**.

3 Quality assessment of the spectroscopic parameters

The process of updating or expanding spectroscopic databases in order to provide the users with a database that insures an optimal characterization of spectral properties of molecular absorption is challenging. To help making the most relevant choices and updates, the Spectroscopic Parameters And Radiative Transfer Evaluation (SPARTE) chain [8] was developed at LMD and applied for the first time to the GEISA-2015 release. Figure 7 presents the general principles of the SPARTE chain.

SPARTE relies on the comparison between forward radiative transfer simulations made by the 4A radiative transfer code [155], [156], using a given spectroscopic database as input, and various remote sensing observations for several thousands of well-characterized atmospheric situations. Assuming that observed spectra come from instruments characterized by high radiometric and spectral stabilities, each individual residual can be the sum of a random errors coming from the imperfect description of the atmospheric state or from the radiometric noise of the instruments, with systematic errors linked in general to the spectra calculation (spectroscopic parameters, line shape, line mixing, continua, etc.). Averaging the resulting ‘calculated-observed’ spectral residuals thus minimizes the random errors and enables the detection of error features that may come from an imperfect knowledge of lines parameters (see Figure 7 of Ref. [8] for the typical shapes of the ‘calculated-observed’ residuals originating from an error in a given line parameter).

In the following, ‘clear-sky’ observations from two well established instruments have been used: the space-borne IASI instrument (15 to 3.7 μm) and ground-based Fourier Transform Spectrometers (FTS) instruments at the Park Falls Total Carbon Column Observing Network [9] (TCCON, 2.5 to 0.7 μm). For IASI, more than 15 000 numbers of observations, spanning the period 2019/01-2019/12, have been used. Atmospheric state determinations that are collocated with IASI observations are from Analyzed RadioSoundings Archive (ARSA) developed and maintained at LMD (<https://ara.lmd.polytechnique.fr/index.php?page=arsa>). Clear sky observations – *i.e.* « free from clouds and aerosols » – are selected through a sophisticated procedure combining microwave and infrared radiances measured by AMSU-A and IASI, both on Metop satellites, respectively. For TCCON, clear-sky observations have been selected to each PI’s station. Here, a set of more than 300 spectra from Park Falls facility, selected based on a good representation of the airmass (*i.e.* solar zenith angle) as well as period of the year, are used. Atmospheric states that are collocated with TCCON observations are coming from the TCCON procedure and delivered together with the spectra.

When possible (*i.e.* for matching spectral coverage and a significant spectral signature of the spectroscopic change), the SPARTE chain was used for the update of GEISA-2020 in a 3-step procedure: (i) to identify improperly described lines (position, intensity, width, pressure shift, etc.); (ii) to provide feedback and suggest revised values to the contributing spectroscopy research teams; and (iii) to finalize the validation of the updated parameters.




Figure 7. The general principle of the SPARTE chain.

Four applications and impact on the GEISA-2020 version

3.1.1 H_2O evaluation in the 6600–6616 cm^{-1} and 7550-7575 cm^{-1} spectral region

As stated in section 2.1.1.1, the major update on $H_2^{16}O$ was based on the work of Mikhailenko *et al.* [11] (referenced as Mikhailenko 2016 in the following), who published a linelist of H_2O parameters between 5830 and 8340 cm^{-1} based on a collection of several data. More recently they published refined line positions and shifts with respect to their former linelist [12] (included in GEISA-2020 update), but other parameters like line intensities or air-broadenings remained unchanged. Although much of these new data helped to improve line parameters in the GEISA database, the validation process using SPARTE based on TCCON observations highlighted a few remaining issues or artefacts in the SWIR domain. In the following, we present a selection of three representative cases that illustrate the benefit of a careful further evaluation of the spectroscopic parameters as nominally delivered by the spectroscopist research team.

The first case is the most straightforward as it concerns ‘wide-range’ issues, *i.e.* systematic errors on one or several line parameters that affect wide ranges of the TCCON spectrum.

Figure 8 summarizes the comparisons of the ‘calculated-observed’ residuals between TCCON and 4A/OP [10] when three different spectroscopic databases are used: GEISA-2015 (red), Mikhailenko 2016 (blue) and GEISA-2020 (green). The upper panel shows the spectrum of atmospheric transmission measured by TCCON in the regions 6600–6616 cm^{-1} and 7550–7575 cm^{-1} . The corresponding residuals (differences of transmissions averaged on the 325 TCCON spectra) are shown in the lower panels. In the left figure, the calculations using the line list from Mikhailenko 2016 results in unexpectedly bad ‘calculated-observed’ values: negative (left panel) or positive (right panel). This wide range behavior, affecting several lines over tens of cm^{-1} , combined with the typical ‘line intensity’ residual shape, suggests systematic line-intensity issues in the data reported in the Mikhailenko 2016 list. This is also supported by the fact that the GEISA-2015 residuals - which did not include Mikhailenko 2016 results – not only were smaller but also did not exhibit similar behavior. An analysis of the affected lines showed that they were all coming from Oudot *et al.* [157] work. Removing all the occurrence of this work in the Mikhailenko 2016 linelist enabled to solve these issues. A work is in progress to assess the origin of these systematic intensity-like issues in Oudot *et al.* data.




Figure 8: Comparison between GEISA-2015, Mikhailenko 2016 [11] and GEISA-2020. Upper panel: Transmission spectrum observed at Park Falls TCCON site (average over 325 spectra). Lower panel: ‘Calculated-observed’ residuals generated by the SPARTE chain for GEISA-2015 (red), Mikhailenko 2016 (blue) and GEISA-2020 (green). Left: 6600–6616 cm^{-1} . Right: 7550–7575 cm^{-1} .

Figure 9 illustrates a similar case. It shows comparisons of the ‘calculated-observed’ residuals obtained from simulations using the three different spectroscopic line lists in the 6615-6616 cm^{-1} , 6669-6672 cm^{-1} and 6710-6712 cm^{-1} regions, and the corresponding TCCON observations. Significant deviations are observed on the residuals when using Mikhailenko 2016 linelist on some isolated lines of the spectrum, with a deterioration of the situation with regard to GEISA-2015. Careful study of the line parameters gathered in

Position cm^{-1}	Intensity cm/atm			Air-broadening cm/cm^{-1}		
	GEISA-2015	Ref. [11]	GEISA-2020	GEISA-2015	Ref. [11]	GEISA-2020
6031.899200	7.13E-26	7.13E-26	7.13E-26	0.0380	0.0078	0.0300
6615.451320	1.76E-25	2.49E-25	1.76E-25	0.0729	0.0729	0.0729
6670.485430	5.04E-24	2.52E-23	5.04E-24	0.0360	0.0360	0.0360
6711.331920	1.04E24	1.32E-24	1.04E-24	0.0330	0.0330	0.0330
7804.608830	6.60E-26	6.87E-26	6.87E-26	0.0092	0.0092	0.0478
7851.277170	3.90E-26	4.13E-26	4.13E-26	0.0089	0.0089	0.0300

Table 11 shows that the main differences come from line intensities of some strong lines. For these lines, the data from Romanini *et al.* [158] were considered in Mikhailenko 2016. However, the CRDS setup used in [158], being very efficient for the measurement of weak lines but less suitable for the measurement of strong lines, leads to significant uncertainties on these line intensities. As a consequence, the choice has been made in GEISA-2015 and GEISA-2020 to use the line intensities determined by Toth *et al.* [159] using a Fourier Transform spectrometer. Other parameters were updated using the data from references [11] and [12].




Figure 9: Comparison between GEISA-2015, Mikhailenko 2016 [11] and GEISA-2020. Upper panel: Transmission spectrum observed by TCCON, as input (average over 325 spectra). Lower panel: ‘Calculated-observed’ residuals generated by the SPARTE chain for GEISA-2015 (red), Mikhailenko 2016 (blue) and GEISA-2020 (green).

Finally, Figure 10 presents comparisons of the ‘calc-obs’ residuals obtained from the three different spectroscopic line lists in the 6631.5-6632.5 cm^{-1} , 7804-7805 cm^{-1} and 7850.5-7851.5 cm^{-1} regions, and the corresponding TCCON observations. Again, significant deviations are observed on the residuals when using Mikhailenko 2016 linelist on some isolated lines of the spectrum, but also using GEISA-2015, a notable exception being the line in 6032.9 cm^{-1} for which GEISA-2015 is far better. This time, the residual shapes indicates a significant difference between the air-broadening coefficients of the three line lists, as illustrated in

Position cm^{-1}	Intensity cm/atm			Air-broadening cm/cm^{-1}		
	GEISA-2015	Ref. [11]	GEISA-2020	GEISA-2015	Ref. [11]	GEISA-2020
6031.899200	7.13E-26	7.13E-26	7.13E-26	0.0380	0.0078	0.0300

6615.451320	1.76E-25	2.49E-25	1.76E-25	0.0729	0.0729	0.0729
6670.485430	5.04E-24	2.52E-23	5.04E-24	0.0360	0.0360	0.0360
6711.331920	1.04E24	1.32E-24	1.04E-24	0.0330	0.0330	0.0330
7804.608830	6.60E-26	6.87E-26	6.87E-26	0.0092	0.0092	0.0478
7851.277170	3.90E-26	4.13E-26	4.13E-26	0.0089	0.0089	0.0300

Table 11 (a slight difference on line intensities is also seen for the two lines at 7804.61 and 7851.28 cm^{-1} , but cannot explain the large deviations seen in Figure 10). The values of these parameters reported in GEISA-2015 and Mikhailenko 2016, and determined in a former work of Gamache *et al.* (updated version of Refs. [160], [161]), prove to be too weak, leading to large errors in the ‘calc-obs’ residuals. In this particular case, we gave a feedback to Gamache and coworkers so that they can rework on a revised set of data. The new line parameters were used to correct the problematic lines in GEISA-2020, as stated in section 2.1.1.1. Thus, Figure 10 shows that the residuals obtained using the GEISA-2020 release give much better results than the two other line lists. Also note that SPARTE was able to detect inconsistencies on parameters of very weak lines (e.g. the line centered at 7804.5 cm^{-1}), detecting features smaller than the instrumental noise.




Figure 10: Comparison between GEISA-2015, Mikhailenko 2016 [11] and GEISA-2020. Upper panel: Transmission spectrum observed by TCCON, as input (average over 325 spectra). Lower panel: ‘Calculated-observed’ residuals generated by the SPARTE chain for GEISA-2015 (red), Mikhailenko 2016 (blue) and GEISA-2020 (green).

Position cm ⁻¹	Intensity cm/atm			Air-broadening cm/cm ⁻¹		
	GEISA-2015	Ref. [11]	GEISA-2020	GEISA-2015	Ref. [11]	GEISA-2020
6031.899200	7.13E-26	7.13E-26	7.13E-26	0.0380	0.0078	0.0300
6615.451320	1.76E-25	2.49E-25	1.76E-25	0.0729	0.0729	0.0729
6670.485430	5.04E-24	2.52E-23	5.04E-24	0.0360	0.0360	0.0360
6711.331920	1.04E24	1.32E-24	1.04E-24	0.0330	0.0330	0.0330
7804.608830	6.60E-26	6.87E-26	6.87E-26	0.0092	0.0092	0.0478
7851.277170	3.90E-26	4.13E-26	4.13E-26	0.0089	0.0089	0.0300

Table 11: Line parameters contained in the 3 spectroscopic databases GEISA-2015, Mikhailenko 2016 and GEISA-2020 for several absorption lines located in the spectral range considered in Figure 9 and Figure 10. Parameters written in italic-bold are those identified as responsible for strong departures in the ‘calculated-observed’ residuals (see text for details).

3.1.2 *O₂ evaluation*

As stated in section 2.2.7, no major update of the O₂ database was made in GEISA-2020. However, a recent work from Tran *et al.* [65] highlighted systematic errors in the 1.27 μ m band using ¹⁶O₂ air-broadened line-shift parameters from GEISA-2015 or HITRAN 2016 databases. To illustrate these errors, Figure 11 summarizes the comparisons of the ‘calculated-observed’ residuals between the 325 TCCON spectra recorded at Park Falls and 4A/OP when three different spectroscopic databases are used: GEISA-2015 taking into account the Pressure Shift (red, marked GEISA-2015 WPS), GEISA-2015 with ¹⁶O₂ pressure shifts set to zero (blue, marked GEISA-2015 WOPS) and GEISA-2020 (green). Figure 11a shows the atmospheric transmission spectrum measured by TCCON in the region 7785–7915 cm⁻¹. The corresponding ‘calc-obs’ residuals are shown in Figure 11b, c, d, for GEISA-2015 WPS, GEISA-2015 WOPS and GEISA-2020, respectively. Comparisons of the residuals between GEISA-2015 WPS and GEISA-2015 WOPS shows that systematic deviations seen in GEISA-2015 are due to erroneous values of the ¹⁶O₂ pressure shifts, as they are mainly addressed when setting these parameters to zero. Figure 11d shows that the new data from Tran *et al.* [65] included in GEISA-2020 (see section 2.1.7) greatly improves these ‘calculated-observed’ deviations.




Figure 11: Comparison between GEISA-2015 (WPS), GEISA-2015 with ¹⁶O₂ pressure shifts set to zero (GEISA-2015 WOPS) and GEISA-2020. (a) Transmission spectrum observed by TCCON, as input. (b) ‘Calculated-observed’ residuals generated by the SPARTE chain for GEISA-2015 WPS (red), (c) GEISA-2015 WOPS (blue) and GEISA-2020 (green).

3.1.3 O₃ evaluation

As described in section 2.1.3, a significant update has been made for the 2020 version of GEISA, notably by the application of intensity scaling factors in three vibrational bands, namely the 000-000, 001-000 and 101-000 bands, respectively in the microwave, 10μm and 5μm spectral regions. Impact of such scaling factors may be evaluated by comparison of simulated spectra generated by the SPARTE chain with observed IASI spectra.

The impact of the release has been studied on the whole IASI spectral range, but a significant impact has been only observed for the 9.6 μm band only. Figure 12 shows the comparisons of ‘calculated-observed’ residuals obtained with the SPARTE chain applied on IASI spectra using the two databases GEISA-2015 (in blue) and GEISA-2020 (in red), in this corresponding spectral range 975-1075 cm⁻¹, for the ‘Tropical’ atmospheres collocated with IASI. Figure 12 shows significantly worse results with GEISA-2020 than those obtained with GEISA-2015. When solicited, the spectroscopists have largely documented and justified (by private communications) the changes made in this spectral region, so the reason for such a

behavior displayed in Figure 12 can only come from SPARTE. Since the same fixed version of 4A/OP has been used for the results displayed in Figure 13, the reason of this deterioration of the residuals using GEISA-2020 can only come from the ozone profiles. So far, as described in [8], O₃ atmospheric profiles originate from the ERA Interim reanalysis [162] that are space and time collocated with each individual quality controlled radiosonde profile retained in the ARSA database. As a consequence, we decided to study the impact of a different set of ozone profiles, especially the new ECMWF reanalysis product, ERA5 [163], which is an improvement of the former ERA Interim product. Because the ERA5 database is large (0.25 x 0.25 degrees, every hour), as a first step, we estimated the differences between both in terms of bias for the whole 2019 period, and for the ‘tropical’ atmospheres only, as shown in Figure 12. The most significant change between the two ERA products being a bias of 8% between 2 and 25 mbars, we evaluated the impact on SPARTE by shifting the ERA Interim profiles by 8% in the pressure range considered in SPARTE. The corresponding residuals are shown on the Figure 13 in green.

As can be easily seen, the previous bias obtained using ERA Interim profiles along with GEISA-2020 has been largely removed. The remaining spectral residuals can be explained by the fact that the best approach is to collocate each ARSA radiosounding with the corresponding ERA5 ozone profile, an adaptation of SPARTE that will be done in the coming weeks. But, from this evaluation, we estimated that the demonstration of the good quality of the new ozone line list is made here, and thus we decided to keep this new version of ozone spectroscopic parameters in GEISA-2020.




Figure 12: Comparison of ‘calculated-observed’ IASI residuals in the 975-1075 cm-1 spectral range and for tropical atmospheres, obtained using GEISA-2015 and GEISA-2020 O₃ line parameters and with different sources of ozone profiles used in the SPARTE chain: GEISA-2015 and ERA in blue, GEISA-2020 and ERA Interim in red and GEISA-2020 and ERA-5 in green).

3.1.4 Discussion on SF₆

.Contrary to the species describe in the section 2, SF₆ has not been updated in the current version of GEISA, although new data were provided by Ke *et al.* from the Dijon’s database SheCaSDa [164]. Indeed, recent data from Nikitin *et al.* [165] consisting of a complete line list based of extensive *ab initio* calculations (from TheoReTS database [166]) leads to significantly different results. The direct comparison of the different available databases with experimental absorption cross section spectra recorded at PNNL by Sharpe *et al.* [167] justify why we made the choice not to update the GEISA-2020 database with new data from Dijon.

Figure 13 shows a comparison of SF₆ cross sections at room temperature (296K, upper panel) and at cold temperature (180K, lower panel). In the upper panel, the experimental PNNL spectrum [81], [167] (recorded at T=29K and N₂ pressure of 760 Torr with a medium resolution of 0.112 cm⁻¹) is plotted along with simulated spectra derived from GEISA-2015 [1], HITRAN 2016 [2], Dijon SheCaSDa [115] and TheoReTS [165] databases. The lower

panel compares only experimental PNNL spectrum at 180K with corresponding calculation from TheoReTS. It appears that *ab initio* cross sections agree much better with the observations than the simulations using SF₆ line lists constructed from effective models as available in HITRAN 2016, SheCaSDa and GEISA-2015 databases. The main reason relies on the fact that effective models, generally derived from cold spectra over a limited set of ground vibrational states, fail to extrapolate data at room temperature which requires an account of more significant contributions of hot bands involving higher vibrational states. On the other hand, *ab initio* calculations succeed in giving a qualitative agreement with experimental data, both at 180K where the comparison is very good, and at 296K for which it is slightly worse (Nikitin *et al.* mentioned that their calculation should be extended [165]).




Figure 13: Comparison of SF₆ cross-sections. Upper panel: simulation at 296 K from the line list of TheoReTS [165] (green_a), the experimentally observed PNNL spectrum [167] (black_b), simulation from line list of GEISA-2015 [1] (blue_c), the theoretical calculation of Dijon SheCaSDa [164] (red_d), and simulation from line list of HITRAN 2016 database [2] (yellow_e). Lower panel: simulation at 180 K from the line list of TheoReTS (green_a) and the experimental observed spectrum from PNNL (black_b).

4 Conclusion and perspectives

Since its creation in the mid-1970s, and in order to meet the needs of researchers and international space agencies, GEISA has undergone 6 updates. This was done by collecting,

archiving, and distributing the most up to date as well as accurate spectroscopic information and by concomitantly evolving the associated management softwares and quality control tools. Accordingly, the GEISA-2020 edition reported in this paper contains important updates and additions to the line parameters sub-database. With the addition of 6 new molecules (HONO, COFCl, CH₃F, CH₃I, RuO₄, H₂C₃H₂), it now involves 58 molecular species - 145 isotopologues (which 27 are new entries) – representing 6,746,987 entries in the database, which span the spectral range from 10⁻⁶ to 35877 cm⁻¹. In close relation with space missions (IASI, IASI-NG, Merlin, MicroCarb) and taking advantage of ground-based observations (here TCCON FTS), noticeable efforts have been made to improve the validation of the updates for species relevant for the Earth's atmosphere, such as H₂O, O₂, O₃, CO₂ and CH₄. Part of this work was enabled by the systematic use of the validation chain SPARTE to evaluate individual line parameters, detect problematic line parameters and consolidate the choices made by the GEISA team.

While the update presented in this paper highly upgrades the previous GEISA2015 release, there is still significant room for improvement. One of the most important aspects of our work is to make available the results of the quality controls we made during the phase of acquisition and delivery processes of new or updated spectroscopic data. The SPARTE chain proves to be a powerful tool to validate and select data for major species of the Earth's atmosphere, but it is still limited for evaluating more exotic species or species having too small spectral signatures. The improvement of some parts of the SPARTE chain through improvement of underlying models (4A/OP), reference spaceborne observations (ACE-FTS limb measurements) or databases (ARSA, ECMWF Reanalyses...) is under progress. In parallel, we plan to generalize, when possible, the use of *ab initio* data, in collaboration with other teams, both as a point of comparison with experimentally derived effective models traditionally used in spectroscopic databases, but also to improve or complete existing data.

Considering the line parameter database, a crucial path of improvement is the consideration of non-Voigt line-shape parameters. Even though the amount of measurements and calculations involving the recommended Hartmann-Tran profile [168] remains limited so far, the level of precision achieved by the sophisticated non-Voigt profiles is more and more required by ground-based or spaceborne instruments. The future releases of GEISA will be designed to gather such high-level information.

The current GEISA-2020 as well as the GEISA-2015 compilations can be accessed through <https://geisa.aeris-data.fr/>. Thanks to an updated user-friendly interface, the operations of

selection, extraction, sorting, filtering, graphical representation, statistical analysis of the most characteristic spectroscopic parameters are more attractive than in the previous web version.

5 Acknowledgements.

IASI has been developed and built under the responsibility of CNES. It is flown onboard the Metop satellites as part of the EUMETSAT Polar System. The IASI L1 data are received through the EUMETCast near real time data distribution service. We particularly wish to thank the Mésocentre ESPRI services for their help in getting IASI data. TCCON data were obtained from the TCCON Data Archive, hosted by the Carbon Dioxide Information Analysis Center (CDIAC) - tcon.onrl.gov. Calculations were performed using the resources of IDRIS, the computing centre of CNRS and of the IPSL data and computing centre Mésocentre ESPRI. This work has been funded by CNES. The authors thank the CNES MicroCarb team for support and fruitful discussions. The authors are thankful to ECMWF for making the ERA-I and ERA5 outputs available through the ECMWF Data Server.

We also thank the AERIS/IPSL/CNES atmospheric data and computing centre for ensuring a friendly access to the GEISA international community and for managing the website.

Pr. R.R. Gamache is pleased to acknowledge support of this research by the National Science Foundation through Grant No. AGS-1156862.

The researches from V.E. Zuev Institute of Atmospheric Optics of Siberian Branch of Russian Academy of Sciences were supported by the Ministry of Science and Higher Education of the Russian Federation.

The financial support of the Natural Sciences and Engineering Research Council of Canada is gratefully acknowledged by Pr. N. Moazzen-Ahmadi.

The work of Tomsk group on ozone spectroscopy was supported by the Russian Science Foundation RNF grant no. 19-12-00171. GSMA Reims and LiPhy Grenoble acknowledge a support from the French-Russian collaboration program LIA CNRS "SAMIA". V. Ebert and G. Li acknowledge support of the work by the EUMETRISPEC project (www.eumetrispec.org) within the European Metrology Research Program (EMRP).

6 Appendix A. List of acronyms

4A	Atlas Automatisé des Absorptions Atmosphériques ; Automatized Atmospheric Absorption Atlas
4A/OP	4A/Operational release
AERIS	Atmosphere and service data pole (CNES, CNRS, IPSL), France
AFGL	Air Force Geophysics Laboratory

ARA/ABC(t)	Atmospheric Radiation Analysis/Atmosphère-Biosphère-Climat (télédétection)
ASD	Acetylene Spectroscopic Databank
CAL/VAL	Calibration/Validation
CDMS	Cologne Database for Molecular Spectroscopy
CDS	Carbon Dioxide Spectroscopic Databank
CIRS	Composite InfraRed Spectrometer
CNRS	Centre National de la Recherche Scientifique (France)
CNES	Centre National d'Etudes Spatiales (France)
CRB	Complex Robert-Bonamy
CRDS	Cavity ring-down spectroscopy
CVRQD	name of the adiabatic semiglobal PESs of water isotopologues
DAS	Differential laser Absorption Spectroscopy
DMS	Dipole Moment Surface-3D
DVR-3D	Discrete Variable Representation
FT-CEAS	Continuous Wave-Cavity Absorption Spectroscopy
EH	Effective Hamiltonian
ENVISAT	ENVironmental SATellite
EUMETSAT	European Organisation for the Exploitation of Meteorological Satellites
FTIR	Fourier Transformed InfraRed spectroscopy
FTS	Fourier Transform Spectrometer
GEISA	Gestion et Etude des Informations Spectroscopiques Atmosphériques; Management and study of Atmospheric Spectroscopic Information
GOSAT	Greenhouse Observing SATellite project
GS	Ground State
GSMA	Groupe de Spectroscopie Moléculaire et Atmosphérique (France)
HITRAN	HIGH-resolution TRANsmision molecular absorption database
HWHM	Half Width at Half Maximum
IAO	Institute of Atmospheric Optics (Russia)
IASI	Infrared Atmospheric Sounder Interferometer
IASI/NG	Infrared Atmospheric Sounder Interferometer/New Generation
ICB	Institut Carnot de Bourgogne
ID	Identification code
IDRIS	Institut du développement et des ressources en informatique scientifique
INSU	Institut National des Sciences de l'Univers (France)
IPSL	Institut Pierre Simon LaplaceIR InfraRed
ISSWG	IASI Sounding Science Working Group
JPL	Jet Propulsion Laboratory (USA)
KIT	Institute for Meteorology and Climate Research Centre Karlsruhe, (Germany)
LIPhy	Laboratoire Interdisciplinaire de Physique (France)
LISA	Laboratoire Inter-Universitaire des Systèmes Atmosphériques (France)
LMD	Laboratoire de Météorologie Dynamique (France)
MCRB	Modified Complex Robert-Bonamy
MERLIN	Methane Remote Sensing Lidar Mission
MICROCARB	MICROsatellite mission to monitor the fluxes of CARBOn dioxide
MIPAS	Michelson Interferometer for Passive Atmospheric Sounding
Metop	Meteorological operational satellite
NASA	National Aeronautics and Space Administration (USA)
NCAR	National Center for Atmospheric research (USA)

NIR	Near-InfraRed
PES	Potential Energy Surface
SPARTE	Spectroscopic Parameters And Radiative Transfer Evaluation
S&MPO	Spectroscopy & molecular properties of Ozone
SWIR	ShortWave Infrared
TIR	Thermal InfraRed
UV	Ultra Violet
VAMDC	Virtual Atomic and Molecular Data Centre
WN	Wave Number (cm ⁻¹)

7 Appendix B. Description of the format used for the line parameters stored in the GEISA-2020 edition:

The format of each entry is described in the following Table 14.

Each entry is a 252 character record to describe the 31 spectroscopic line parameters.

First line of Table 14: the 31 Spectroscopic line parameters are listed in the 31 columns and their description is given below .

Line 2 and line 3 display the field length and the FORTAN format descriptor, respectively.

Line 4 displays the standard_default values associated to each parameter.

Line 5 displays a cumulative index indicating the position of the last character of the record associated to each of the 31 spectroscopic line parameters.

The standard default values for fields «O'», « T » and « T' », have been changed and set to “zero”. This modification was made to avoid potential misunderstanding and thus improper use of these parameters in some applications especially related to forward radiative transfer.- Value in field “M” is documented in GEISA only if it is directly provided by the author of the spectroscopic line entry.

Table 14

Format of each entry in GEISA-2015

Parameter	A	B	C	D	E1	E2	E3	E4	F	G	I	J
Field length	12	11	6	10	25	25	15	15	4	3	3	3
Fortran descriptor	F12.6	1PD11.4	0PF6.4	F10.4	A25	A25	A15	A15	F4.2	I3	I3	A3
Undefined values	NR	-9.9999D-01	-.9999	-0.9999	*	*	*	*	-.99	-.99	-.99	*
Record counting	12	23	29	39	64	89	104	119	123	126	129	132

K	L	M	N	O	R	A'	B'	C'	F'
2	1	10	7	9	6	10	11	6	4
I2	I1	1PE10.3	0PF7.4	F9.6	F6.4	F10.6	1PD11.4	0PF6.4	F4.2
-9	0	-9.999E-01	-9.9999	0.000000	-.9999	-0.999999	-9.9999D-01	-.9999	-.99
134	135	145	152	161	167	177	188	194	198

O'	R'	N'	S	S'	T	T'	U	U'
9	6	7	4	4	8	8	4	4
F9.6	F6.4	F7.4	F4.2	F4.2	F8.6	F8.6	F4.2	F4.2
0.000000	-.9999	-9.9999	-.99	-.99	0.000000	0.000000	-.99	-.99
207	213	220	224	228	236	244	248	252

A : wave number (cm^{-1}) of the line

B : intensity of the line in ($\text{cm}^{-1}/(\text{molecule} \cdot \text{cm}^{-2})$) at 296K

C : Air broadening pressure halfwidth (HWHM) ($\text{cm}^{-1}\text{atm}^{-1}$) at 296K

D : Energy of the lower transition level (cm^{-1})

E_i (i=1,2,3,4) : Transition quantum identifications for the lower and upper state of the transition

E1 : upper state vibrational identification E2 : lower state vibrational identification

E3 : upper state rotationnal identification E4 : lower state rotationnal identification

F : temperature dependence coefficient n of the air broadening halfwidth
 G : identification code for isotopologue as in GEISA
 I : identification code for molecule as in GEISA
 J : Internal GEISA code for the data identification
 K : Molecule number as in HITRAN
 L : isotopologue number (1=most abundant, 2= second...etc) as in HITRAN
 M : Einstein A-coefficient
 N : self broadening pressure halfwidth (HWHMself) ($\text{cm}^{-1}\text{atm}^{-1}$) at 296K
 O : air pressure shift of the line transition (cm^{-1}) at 296K
 R : temperature dependence coefficient of the air pressure shift
 A' : estimated accuracy (cm^{-1}) on the line position
 B' : estimated accuracy on the intensity of the line in ($\text{cm}^{-1}/(\text{molecule}\cdot\text{cm}^{-2})$)
 C' : estimated accuracy on the air collision halfwidth (HWHM) ($\text{cm}^{-1}\text{atm}^{-1}$)
 F' : estimated accuracy on the temperature dependence coefficient of the air broadening halfwidth
 O' : estimated accuracy on the air pressure shift of the line transition (cm^{-1}) at 296K
 R' : estimated accuracy on the temperature dependence coefficient of the air pressure shift
 N' : estimated accuracy on the self broadened (HWHM) ($\text{cm}^{-1}\text{atm}^{-1}$) at 296K
 S : temperature dependence coefficient of the self broadening halfwidth
 S' : estimated accuracy on the temperature dependence coefficient of the self broadening halfwidth
 T : self pressure shift of the line transition (cm^{-1}) at 296K
 T' : estimated accuracy on the self pressure shift of the line transition (cm^{-1}) at 296K
 U : temperature dependence coefficient of the self pressure shift
 U' : estimated accuracy on the temperature dependence coefficient of the self pressure shift

As shown in line 4 of Table 14 GEISA undefined values are attributed to the line parameter entries when no value is available from the data provider (missing data).

8 Appendix C. Molecules and isotopologues in GEISA-2015

Description of molecule and isotopologue codes in GEISA-2020 are given in Table 15. The molecule names and associated codes are in the two first columns; for each molecule, the isotopologue codes and the corresponding detailed formula are in columns 3 and 4 respectively. New molecules are in red and new isotopologues are in purple. The abundance used in GEISA is given in t 5th column.

Table 15 Description of the molecule and isotopologue codes in GEISA-2020

Molecule	Molecule Code	I Code	Formula	Abundance
H ₂ O	1	161	H ¹⁶ OH	0.997317
		181	H ¹⁸ OH	0.002000

Molecule	Molecule Code	I Code	Formula	Abundance
		171	H ¹⁷ OH	3.718840 × 10 ⁻⁴
		262	D ₂ ¹⁶ O	2.419700 × 10 ⁻⁸
		282	D ₂ ¹⁸ O	4.852080 × 10 ⁻¹¹
		272	D ₂ ¹⁷ O	9.022841 × 10 ⁻¹²
CO ₂	2	626	¹⁶ O ¹² C ¹⁶ O	0.9842
		636	¹⁶ O ¹³ C ¹⁶ O	1.106 x 10 ⁻²
		628	¹⁶ O ¹² C ¹⁸ O	3.947 x 10 ⁻³
		627	¹⁶ O ¹² C ¹⁷ O	7.339 x 10 ⁻⁴
		638	¹⁶ O ¹³ C ¹⁸ O	4.434 x 10 ⁻⁵
		637	¹⁶ O ¹³ C ¹⁷ O	8.246 x 10 ⁻⁶
		828	¹⁸ O ¹² C ¹⁸ O	3.957 x 10 ⁻⁶
		728	¹⁷ O ¹² C ¹⁸ O	1.472 x 10 ⁻⁶
		727	¹⁷ O ¹² C ¹⁷ O	1.430 x 10 ⁻⁷
		838	¹⁸ O ¹³ C ¹⁸ O	4.446 x 10 ⁻⁸
		738	¹⁷ O ¹³ C ¹⁸ O	1.654 x 10 ⁻⁸
		737	¹⁷ O ¹³ C ¹⁷ O	1.55 x 10 ⁻⁹
O ₃	3	666	¹⁶ O ¹⁶ O ¹⁶ O	0.992901
		668	¹⁶ O ¹⁶ O ¹⁸ O	0.003982
		686	¹⁶ O ¹⁸ O ¹⁶ O	0.001991
		667	¹⁶ O ¹⁶ O ¹⁷ O	7.404750 × 10 ⁻⁴
		676	¹⁶ O ¹⁷ O ¹⁶ O	3.702370 × 10 ⁻⁴
N ₂ O	4	446	¹⁴ N ¹⁴ N ¹⁶ O	0.990333
		456	¹⁴ N ¹⁵ N ¹⁶ O	0.003641
		546	¹⁵ N ¹⁴ N ¹⁶ O	0.003641
		448	¹⁴ N ¹⁴ N ¹⁸ O	0.001986
		458	¹⁴ N ¹⁴ N ¹⁸ O	7.300807 × 10 ⁻⁴
		548	¹⁵ N ¹⁴ N ¹⁸ O	7.300807 × 10 ⁻⁴
		447	¹⁴ N ¹⁴ N ¹⁷ O	3.692800 × 10 ⁻⁴
		556	¹⁵ N ¹⁵ N ¹⁶ O	1.338574 × 10 ⁻⁵
CO	5	26	¹² C ¹⁶ O	0.986544
		36	¹³ C ¹⁶ O	0.011084
		28	¹² C ¹⁸ O	0.001978
		27	¹² C ¹⁷ O	3.678670 × 10 ⁻⁴
		38	¹³ C ¹⁸ O	2.222500 × 10 ⁻⁵
		37	¹³ C ¹⁷ O	4.132920 × 10 ⁻⁶
CH ₄	6	211	¹² CH ₄	0.988274
		311	¹³ CH ₄	0.011103

Molecule	Molecule Code	I Code	Formula	Abundance
O ₂	7	66	¹⁶ O ¹⁶ O	0.995262
		68	¹⁶ O ¹⁸ O	0.003991
		67	¹⁶ O ¹⁷ O	7.422350 × 10 ⁻⁴
NO	8	46	¹⁴ N ¹⁶ O	0.993974
		56	¹⁵ N ¹⁶ O	0.003654
		48	¹⁴ N ¹⁸ O	0.001993
SO ₂	9	626	³² S ¹⁶ O ₂	0.945678
		646	³⁴ S ¹⁶ O ₂	0.041950
NO ₂	10	646	¹⁴ N ¹⁶ O ₂	0.991616
		656	¹⁵ N ¹⁶ O ₂	0.003646
NH ₃	11	411	¹⁴ NH ₃	0.995872
		511	¹⁵ NH ₃	0.003661
PH ₃	12	131	³¹ PH ₃	0.999533
HNO ₃	13	146	H ¹⁴ N ¹⁶ O	0.989110
		156	H ¹⁵ N ¹⁶ O	0.003636
OH	14	61	¹⁶ OH	0.997473
		81	¹⁸ OH	0.002000
		62	¹⁶ OD	1.553710 × 10 ⁻⁴
HF	15	19	H ¹⁹ F	0.999844
		29	D ¹⁹ F	1.557410 × 10 ⁻⁴
HCl	16	15	H ³⁵ Cl	0.757587
		17	H ³⁷ Cl	0.242257
		25	D ³⁵ Cl	1.180050 × 10 ⁻⁴
		27	D ³⁷ Cl	3.773500 × 10 ⁻⁵
HBr	17	19	H ⁷⁹ Br	0.506781
		11	H ⁸¹ Br	0.493063
		29	D ⁷⁹ Br	7.893840 × 10 ⁻⁵
		21	D ⁸¹ Br	7.680160 × 10 ⁻⁵
HI	18	17	H ¹²⁷ I	0.999844
		27	D ¹²⁷ I	1.557410 × 10 ⁻⁴
ClO	19	56	³⁵ Cl ¹⁶ O	0.755908
		76	³⁷ Cl ¹⁶ O	0.241720
OCS	20	622	¹⁶ O ¹² C ³² S	0.937395
		624	¹⁶ O ¹² C ³⁴ S	0.041583
		632	¹⁶ O ¹³ C ³² S	0.010531
		623	¹⁶ O ¹² C ³³ S	0.007399
		822	¹⁸ O ¹² C ³² S	0.001880
		634	¹⁶ O ¹³ C ³⁴ S	4.671757 × 10 ⁻⁴
H ₂ CO	21	126	H ₂ ¹² C ¹⁶ O	0.986237
		136	H ₂ ¹³ C ¹⁶ O	0.011080
		128	H ₂ ¹² C ¹⁸ O	0.001978
C ₂ H ₆	22	226	¹² C ₂ H ₆	0.976990
		236	¹² C ¹³ CH ₆	0.021953

Molecule	Molecule Code	I Code	Formula	Abundance
CH ₃ D	23	212	¹² CH ₃ D	6.157510×10^{-4}
		312	¹³ CH ₃ D	6.917850×10^{-6}
C ₂ H ₂	24	221	¹² C ₂ H ₂	0.977599
		231	¹² C ¹³ CH ₂	0.021966
C ₂ H ₄	25	211	¹² C ₂ H ₄	0.977294
		311	¹² C ¹³ CH ₄	0.021959
GeH ₄	26	411	⁷⁴ GeH ₄	1.000000
		211	⁷² GeH ₄	1.000000
		011	⁷⁰ GeH ₄	1.000000
		311	⁷³ GeH ₄	1.000000
		611	⁷⁶ GeH ₄	1.000000
HCN	27	124	H ¹² C ¹⁴ N	0.985114
		134	H ¹³ C ¹⁴ N	0.011068
		125	H ¹² C ¹⁵ N	0.003622
		224	D ¹² C ¹⁴ N	1.534456×10^{-4}
C ₃ H ₈	28	221	¹² C ₃ H ₈	0.965835
C ₂ N ₂	29	224	¹² C ₂ ¹⁴ N ₂	0.970752
C ₄ H ₂	30	211	¹² C ₄ H ₂	0.955998
HC ₃ N	31	124	H ¹² C ₃ ¹⁴ N	0.963346
HOCl	32	165	H ¹⁶ O ³⁵ Cl	0.755790
		167	H ¹⁶ O ³⁷ Cl	0.241683
N ₂	33	44	¹⁴ N ¹⁴ N	0.992687
CH ₃ Cl	34	215	¹² CH ₃ ³⁵ Cl	0.748937
		217	¹² CH ₃ ³⁷ Cl	0.239491
H ₂ O ₂	35	166	H ₂ ¹⁶ O ¹⁶ O	0.994952
H ₂ S	36	121	H ₂ ³² S	0.949884
		141	H ₂ ³⁴ S	0.042137
		131	H ₂ ³³ S	0.007498
HCOOH	37	261	H ¹² C ¹⁶ O ¹⁶ OH	0.983898
COF ₂	38	269	¹² C ¹⁶ O ¹⁹ F ₂	0.986544
SF ₆	39	29	³² S ¹⁹ F ₆	0.950180
C ₃ H ₄	40	341	¹² C ₃ H ₆	0.966587
HO ₂	41	166	H ¹⁶ O ₂	0.995107
ClONO ₂	42	564	¹⁵ Cl ¹⁶ O ¹⁴ N ¹⁶ O ₂	0.749570
		764	¹⁷ Cl ¹⁶ O ¹⁴ N ¹⁶ O ₂	0.239694
CH ₃ Br	43	79	¹² CH ₃ ⁷⁹ Br	0.500995
		81	¹² CH ₃ ⁸¹ Br	0.487433
CH ₃ OH	44	216	¹² CH ₃ ¹⁶ OH	0.985930
NO ⁺	45	46	¹⁴ N ¹⁶ O ⁺	0.993974
HNC	46	142	H ¹⁴ N ¹² C	0.985114
C ₆ H ₆	47	266	¹² C ₆ H ₆	0.934291
C ₂ HD	48	122	¹² C ₂ HD	3.045500×10^{-4}
CF ₄	49	291	¹² C ¹⁹ F ₄	0.988890

Molecule	Molecule Code	I Code	Formula	Abundance
CH ₃ CN	50	234	¹² CH ₃ ¹² C ¹⁴ N	0.973866
HDO	51	162	H ¹⁶ OD	3.106930 × 10 ⁻⁴
		182	H ¹⁸ OD	6.230030 × 10 ⁻⁷
		172	H ¹⁷ OD	1.158530 × 10 ⁻⁷
SO ₃	52	26	³² S ¹⁶ O ₃	0.943400
HONO	53	646	H ¹⁶ O ¹⁴ N ¹⁶ O	0.9937102
COFCl	54	265	¹² C ¹⁶ O ¹⁶ F ³⁵ Cl	1.000000
		267	¹² C ¹⁶ O ¹⁶ F ³⁷ Cl	1.000000
CH ₃ I	55	217	¹² CH ₃ ¹²⁷ I	1.000000
CH ₃ F	56	219	¹² CH ₃ ¹⁹ F	1.000000
RuO ₄	57	102	¹⁰² Ru ¹⁶ O ₄	1.000000
		104	¹⁰⁴ Ru ¹⁶ O ₄	1.000000
		101	¹⁰¹ Ru ¹⁶ O ₄	1.000000
		99	⁹⁹ Ru ¹⁶ O ₄	1.000000
		100	¹⁰⁰ Ru ¹⁶ O ₄	1.000000
		97	⁹⁷ Ru ¹⁶ O ₄	1.000000
		98	⁹⁸ Ru ¹⁶ O ₄	1.000000
		106	¹⁰⁶ Ru ¹⁶ O ₄	1.000000
		103	¹⁰³ Ru ¹⁶ O ₄	1.000000
H ₂ C ₃ H ₂	58	121	H ₂ ¹² C ₃ H ₂	1.000000

Bibliography

- [1] N. Jacquinet-Husson *et al.*, “The 2015 edition of the GEISA spectroscopic database,” *J. Mol. Spectrosc.*, vol. 327, 2016, doi: 10.1016/j.jms.2016.06.007.
- [2] I. E. Gordon *et al.*, “The HITRAN2016 molecular spectroscopic database,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 203, pp. 3–69, 2017, doi: 10.1016/j.jqsrt.2017.06.038.
- [3] A. Chedin, N. Husson, and N. Scott, “Une banque de données pour l’étude des phénomènes de transfert radiatif dans les atmosphères planétaires: la banque GEISA,” *Bull. d’Information du Cent. Données Stellaires, Fr.*, vol. 22, pp. 21–121, 1982.
- [4] N. Husson, B. Bonnet, N. A. Scott, and A. Chedin, “Management and study of spectroscopic information: The GEISA program,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 48, no. 5, pp. 509–518, 1992, doi: [https://doi.org/10.1016/0022-4073\(92\)90116-L](https://doi.org/10.1016/0022-4073(92)90116-L).
- [5] N. Jacquinet-Husson *et al.*, “The 1997 spectroscopic GEISA databank,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 62, no. 2, pp. 205–254, 1999, doi: [https://doi.org/10.1016/S0022-4073\(98\)00111-3](https://doi.org/10.1016/S0022-4073(98)00111-3).
- [6] N. Jacquinet-Husson *et al.*, “The GEISA spectroscopic database: Current and future archive for Earth and planetary atmosphere studies,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 109, no. 6, pp. 1043–1059, 2008, doi: <https://doi.org/10.1016/j.jqsrt.2007.12.015>.
- [7] N. Jacquinet-Husson *et al.*, “The 2009 edition of the GEISA spectroscopic database,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 112, no. 15, pp. 2395–2445, 2011, doi: <https://doi.org/10.1016/j.jqsrt.2011.06.004>.
- [8] R. Armante *et al.*, “Evaluation of spectroscopic databases through radiative transfer simulations compared to observations. Application to the validation of GEISA 2015 with IASI and TCCON,” *J. Mol. Spectrosc.*, vol. 327, pp. 180–192, Sep. 2016, doi: 10.1016/j.jms.2016.04.004.
- [9] D. Wunch *et al.*, “The total carbon column observing network,” *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*, vol. 369, no. 1943, pp. 2087–2112, 2011, doi: 10.1098/rsta.2010.0240.
- [10] A. Scott, N.A. and Chédin, “A Fast Line-by-Line Method for Atmospheric Absorption Computations: The Automatized Atmospheric Absorption Atlas,” *J. Appl. Meteorol.*, vol. 20, pp. 802–812, 1981.
- [11] S. N. Mikhailenko, S. Kassi, D. Mondelain, R. R. Gamache, and A. Campargue, “A spectroscopic database for water vapor between 5850 and 8340 cm^{-1} ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 179, pp. 198–216, 2016, doi: 10.1016/j.jqsrt.2016.03.035.
- [12] S. N. Mikhailenko, S. Kassi, D. Mondelain, and A. Campargue, “Water vapor absorption between 5690 and 8340 cm^{-1} : Accurate empirical line centers and validation tests of calculated line intensities,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 245, p. 106840, 2020, doi: 10.1016/j.jqsrt.2020.106840.
- [13] B. Vispoel, J. H. Cavalcanti, and R. R. Gamache, “Modified complex Robert-Bonamy calculations of line shape parameters and their temperature dependence for water vapor in collision with N_2 ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 228, pp. 79–89, 2019, doi: 10.1016/j.jqsrt.2019.02.023.
- [14] B. Vispoel, J. H. Cavalcanti, E. T. Paige, and R. R. Gamache, “Vibrational dependence, temperature dependence, and prediction of line shape parameters for the $\text{H}_2\text{O}-\text{N}_2$ collision

system,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 253, p. 107030, 2020, doi: <https://doi.org/10.1016/j.jqsrt.2020.107030>.

- [15] A. Campargue, S. Kassi, A. Yachmenev, A. A. Kyuberis, J. Küpper, and S. N. Yurchenko, “Observation of electric-quadrupole infrared transitions in water vapor,” *Phys. Rev. Res.*, vol. 2, no. 2, p. 23091, 2020, doi: 10.1103/PhysRevResearch.2.023091.
- [16] A. Campargue, A. M. Solodov, A. A. Solodov, A. Yachmenev, and S. N. Yurchenko, “Detection of electric-quadrupole transitions in water vapour near 5.4 and 2.5 μm ,” *Phys. Chem. Chem. Phys.*, vol. 22, no. 22, pp. 12476–12481, 2020, doi: 10.1039/D0CP01667E.
- [17] J. Tennyson *et al.*, “IUPAC critical evaluation of the rotational–vibrational spectra of water vapor. Part I—Energy levels and transition wavenumbers for H₂O and H₂¹⁸O,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 110, no. 9, pp. 573–596, 2009, doi: <https://doi.org/10.1016/j.jqsrt.2009.02.014>.
- [18] A. A. Kyuberis *et al.*, “Room temperature line lists for deuterated water,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 203, pp. 175–185, 2017, doi: 10.1016/j.jqsrt.2017.06.026.
- [19] S. V. Shirin, N. F. Zobov, and O. L. Polyansky, “Theoretical line list of D₂O up to 16,000 cm^{-1} with an accuracy close to experimental,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 109, no. 4, pp. 549–558, 2008, doi: 10.1016/j.jqsrt.2007.07.010.
- [20] R. R. Gamache and J.-M. Hartmann, “An intercomparison of measured pressure-broadening and pressure-shifting parameters of water vapor,” *Can. J. Chem.*, vol. 82, no. 6, pp. 1013–1027, 2004, doi: 10.1139/v04-069.
- [21] L. H. Coudert, “The bending potential energy function of HDO obtained from high-resolution data,” *J. Mol. Spectrosc.*, vol. 330, pp. 112–119, 2016, doi: 10.1016/j.jms.2016.07.008.
- [22] P. Bièvre, M. Gallet, N. E. Holden, and I. L. Barnes, “Isotopic Abundances and Atomic Weights of the Elements,” *J. Phys. Chem. Ref. Data*, vol. 13, no. 3, pp. 809–891, 1984, doi: 10.1063/1.555720.
- [23] S. N. Yurchenko *et al.*, “Potential energy surface of HDO up to 25 000 cm^{-1} ,” *J. Chem. Phys.*, vol. 128, no. 4, pp. 1–13, 2008, doi: 10.1063/1.2806165.
- [24] S. A. Tashkun, V. I. Perevalov, R. R. Gamache, and J. Lamouroux, “CDS-296, high-resolution carbon dioxide spectroscopic databank: An update,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 228, pp. 124–131, May 2019, doi: 10.1016/J.JQSRT.2019.03.001.
- [25] S. A. Tashkun, V. I. Perevalov, R. R. Gamache, and J. Lamouroux, “CDS-296, high resolution carbon dioxide spectroscopic databank: Version for atmospheric applications,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 152, pp. 45–73, Feb. 2015, doi: 10.1016/J.JQSRT.2014.10.017.
- [26] “<ftp://ftp.iao.ru/pub/CDS-296/>.”
- [27] V. Wilquet *et al.*, “Line parameters for the 01111–00001 band of ¹²C¹⁶O¹⁸O from SOIR measurements of the Venus atmosphere,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 109, no. 6, pp. 895–905, Apr. 2008, doi: 10.1016/J.JQSRT.2007.12.021.
- [28] G. L. Villanueva, M. J. Mumma, R. E. Novak, and T. Hewagama, “Identification of a new band system of isotopic CO₂ near 3.3 μm : Implications for remote sensing of biomarker gases on Mars,” *Icarus*, vol. 195, no. 1, pp. 34–44, 2008, doi: 10.1016/j.icarus.2007.11.014.

- [29] V. G. Tyuterev, A. Barbe, D. Jacquemart, C. Janssen, S. N. Mikhailenko, and E. N. Starikova, “Ab initio predictions and laboratory validation for consistent ozone intensities in the MW, 10 and 5 μ m ranges,” *J. Chem. Phys.*, vol. 150, no. 18, 2019, doi: 10.1063/1.5089134.
- [30] S. Mikhailenko and A. Barbe, “High resolution infrared spectrum of 16O₃: The 3600–4300 cm⁻¹ range reinvestigated,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 244, 2020, doi: 10.1016/j.jqsrt.2019.106823.
- [31] V. Kokoouline, D. Lapierre, A. Alijah, and V. Tyuterev, “Localized and delocalized bound states of the main isotopologue 48O₃ and of 18O-enriched 50O₃ isotopomers of the ozone molecule near the dissociation threshold,” *Phys. Chem. Chem. Phys.*, vol. 22, no. 28, pp. 15885–15899, 2020, doi: 10.1039/D0CP02177F.
- [32] S. Vasilchenko *et al.*, “Detection and assignment of ozone bands near 95% of the dissociation threshold: Ultrasensitive experiments for probing potential energy function and vibrational dynamics,” *Phys. Rev. A*, vol. 102, no. 5, p. 52804, Nov. 2020, doi: 10.1103/PhysRevA.102.052804.
- [33] Y. L. Babikov, S. N. Mikhailenko, A. Barbe, and V. G. Tyuterev, “S&MPO – An information system for ozone spectroscopy on the WEB,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 145, pp. 169–196, 2014, doi: <https://doi.org/10.1016/j.jqsrt.2014.04.024>.
- [34] A. Barbe *et al.*, “Ozone spectroscopy in the electronic ground state: High-resolution spectra analyses and update of line parameters since 2003,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 130, pp. 172–190, 2013, doi: <https://doi.org/10.1016/j.jqsrt.2013.06.007>.
- [35] J. M. Flaud, A. Barbe, C. Camy-Peyret, and J. J. Plateaux, “High Resolution Analysis of the 5v₃, 3v₁+ v₂+ v₃, and v₁+ 4v₃ Bands of 16O₃: Line Positions and Intensities,” *J. Mol. Spectrosc.*, vol. 177, no. 1, pp. 34–39, 1996, doi: <https://doi.org/10.1006/jmsp.1996.0114>.
- [36] J.-M. Flaud, C. Camy-Peyret, V. M. Devi, C. P. Rinsland, and M. A. H. Smith, “The v₁ and v₃ bands of 16O₃: Line positions and intensities,” *J. Mol. Spectrosc.*, vol. 124, no. 1, pp. 209–217, 1987, doi: [https://doi.org/10.1016/0022-2852\(87\)90135-4](https://doi.org/10.1016/0022-2852(87)90135-4).
- [37] J.-M. Flaud, C. Camy-Peyret, C. P. Rinsland, M. A. H. Smith, and V. Malathy Devi, “Line parameters for 16O₃ bands in the 7- μ m region,” *J. Mol. Spectrosc.*, vol. 134, no. 1, pp. 106–112, 1989, doi: [https://doi.org/10.1016/0022-2852\(89\)90132-X](https://doi.org/10.1016/0022-2852(89)90132-X).
- [38] A. Barbe, A. Chichery, V. Tyuterev, S. Taskhun, and S. Mikhailenko, “The 2v₂ and 3v₂–v₂ bands of ozone,” *Spectrochim. Acta Part A Mol. Biomol. Spectrosc.*, vol. 54, no. 12, pp. 1935–1945, 1998, doi: [https://doi.org/10.1016/S1386-1425\(98\)00156-5](https://doi.org/10.1016/S1386-1425(98)00156-5).
- [39] H. M. Pickett *et al.*, “The vibrational and rotational spectra of ozone for the (0, 1, 0) and (0, 2, 0) states,” *J. Mol. Spectrosc.*, vol. 128, no. 1, pp. 151–171, 1988, doi: [https://doi.org/10.1016/0022-2852\(88\)90214-7](https://doi.org/10.1016/0022-2852(88)90214-7).
- [40] J. M. Flaud *et al.*, “Ozone absorption around 10 μ m,” *J. Geophys. Res. Atmos.*, vol. 108, no. D9, 2003, doi: <https://doi.org/10.1029/2002JD002755>.
- [41] M. Birk, G. Wagner, J. M. Flaud, and D. Hausamann, “Linestrengths in the v₃–v₂ Hot Band of Ozone,” *J. Mol. Spectrosc.*, vol. 163, no. 1, pp. 262–275, 1994, doi: <https://doi.org/10.1006/jmsp.1994.1022>.
- [42] A. Barbe, O. Sulakshina, J. J. Plateaux, A. Hamdouni, and S. Bouazza, “High-Resolution

- Infrared Spectra of Ozone in the 2300–2600 cm^{-1} Region,” *J. Mol. Spectrosc.*, vol. 170, no. 1, pp. 244–250, 1995, doi: <https://doi.org/10.1006/jmsp.1995.1068>.
- [43] S. Bouazza, S. Mikhailenko, B. A. L. Regalia, V. Tyuterev, and P. J.J, “The $\nu_1 + \nu_2 + 2\nu_3$ and $\nu_2 + 3\nu_3$ Bands of $^{16}\text{O}_3$,” *J. Mol. Spectrosc.*, vol. 174, pp. 510–519, 1995.
- [44] V. G. Tyuterev, R. V Kochanov, and S. A. Tashkun, “Accurate ab initio dipole moment surfaces of ozone: First principle intensity predictions for rotationally resolved spectra in a large range of overtone and combination bands,” *J. Chem. Phys.*, vol. 146, no. 6, p. 64304, 2017, doi: [10.1063/1.4973977](https://doi.org/10.1063/1.4973977).
- [45] A. Barbe, E. Starikova, and M.-R. De Backer, “High resolution infrared spectra of the $^{16}\text{O}^{16}\text{O}^{17}\text{O}$ and the $^{16}\text{O}^{17}\text{O}^{16}\text{O}$ ozone isotopic species. The 5 and 10 micron spectral ranges revisited,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 203, pp. 293–299, 2017, doi: <https://doi.org/10.1016/j.jqsrt.2017.03.034>.
- [46] A. Barbe, E. Starikova, M. R. De Backer, and V. G. Tyuterev, “Analyses of infrared FT spectra of asymmetric ozone isotopologue $^{16}\text{O}^{16}\text{O}^{18}\text{O}$ in the range 950–3850 cm^{-1} ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 218, pp. 231–247, 2018, doi: <https://doi.org/10.1016/j.jqsrt.2018.06.022>.
- [47] V. G. Tyuterev, S. A. Tashkun, and H. Seghir, “High-order contact transformations: general algorithm, computer implementation, and triatomic tests,” in *Proceedings of SPIE - The International Society for Optical Engineering*, 2004, vol. 5311, pp. 164–175, doi: [10.1117/12.545641](https://doi.org/10.1117/12.545641).
- [48] S. A. Tashkun, V. I. Perevalov, A. W. Liu, and S. M. Hu, “Global modeling of the $^{15}\text{N}^{216}\text{O}$ line positions within the framework of the polyad model of effective Hamiltonian and a room temperature $^{15}\text{N}^{216}\text{O}$ line list,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 175, pp. 1–7, 2016, doi: [10.1016/j.jqsrt.2016.01.038](https://doi.org/10.1016/j.jqsrt.2016.01.038).
- [49] S. A. Tashkun, V. I. Perevalov, E. V. Karlovets, S. Kassi, and A. Campargue, “High sensitivity cavity ring down spectroscopy of N_2O near 1.22 μm : (II) $^{14}\text{N}^{216}\text{O}$ line intensity modeling and global fit of $^{14}\text{N}^{218}\text{O}$ line positions,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 176, pp. 62–69, 2016, doi: [10.1016/j.jqsrt.2016.02.020](https://doi.org/10.1016/j.jqsrt.2016.02.020).
- [50] V. Werwein, J. Brunzendorf, A. Serdyukov, O. Werhahn, and V. Ebert, “First measurements of nitrous oxide self-broadening and self-shift coefficients in the 0002–0000 band at 2.26 μm using high resolution Fourier transform spectroscopy,” *J. Mol. Spectrosc.*, vol. 323, pp. 28–42, May 2016, doi: [10.1016/J.JMS.2016.01.010](https://doi.org/10.1016/J.JMS.2016.01.010).
- [51] V. Werwein, J. Brunzendorf, G. Li, A. Serdyukov, O. Werhahn, and V. Ebert, “High-resolution Fourier transform measurements of line strengths in the 00^02 – 00^00 main isotopologue band of nitrous oxide,” *Appl. Opt.*, vol. 56, no. 11, p. E99, Apr. 2017, doi: [10.1364/AO.56.000E99](https://doi.org/10.1364/AO.56.000E99).
- [52] V. Werwein, G. Li, A. Serdyukov, J. Brunzendorf, O. Werhahn, and V. Ebert, “High-resolution Fourier transform measurements of air-induced broadening and shift coefficients in the 0002–0000 main isotopologue band of nitrous oxide,” *J. Mol. Spectrosc.*, vol. 348, pp. 68–78, Jun. 2018, doi: [10.1016/J.JMS.2017.07.002](https://doi.org/10.1016/J.JMS.2017.07.002).
- [53] G. Li *et al.*, “ROVIBRATIONAL LINE LISTS FOR NINE ISOTOPOLOGUES OF THE CO MOLECULE IN THE $X^1\Sigma^+$ GROUND ELECTRONIC STATE,” *Astrophys. J. Suppl. Ser.*, vol. 216, no. 1, p. 15, Jan. 2015, doi: [10.1088/0067-0049/216/1/15](https://doi.org/10.1088/0067-0049/216/1/15).

- [54] R. J. Le Roy, "LEVEL: A computer program for solving the radial Schrödinger equation for bound and quasibound levels," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 186, pp. 167–178, 2017, doi: <https://doi.org/10.1016/j.jqsrt.2016.05.028>.
- [55] J. A. Coxon and P. G. Hajigeorgiou, "Direct potential fit analysis of the X1 Σ^+ ground state of CO," *J. Chem. Phys.*, vol. 121, no. 7, pp. 2992–3008, Aug. 2004, doi: 10.1063/1.1768167.
- [56] M. Ghysels, D. Mondelain, S. Kassı, A. V. Nikitin, M. Rey, and A. Campargue, "The methane absorption spectrum near 1.73 μm (5695–5850 cm^{-1}): Empirical line lists at 80 K and 296 K and rovibrational assignments," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 213, pp. 169–177, 2018, doi: 10.1016/j.jqsrt.2018.04.007.
- [57] A. V. Nikitin *et al.*, "Assignment and modelling of 12CH₄ spectra in the 5550–5695, 5718–5725 and 5792–5814 cm^{-1} regions," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 219, pp. 323–332, 2018, doi: 10.1016/j.jqsrt.2018.08.006.
- [58] A. V. Nikitin *et al.*, "Measurements and modeling of long-path 12CH₄ spectra in the 5300–5550 cm^{-1} region," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 202, pp. 255–264, 2017, doi: 10.1016/j.jqsrt.2017.07.030.
- [59] V. Tyuterev, S. Tashkun, M. Rey, R. Kochanov, A. Nikitin, and T. Delahaye, "Accurate spectroscopic models for methane polyads derived from a potential energy surface using high-order contact transformations," *J. Phys. Chem. A*, vol. 117, no. 50, 2013, doi: 10.1021/jp408116j.
- [60] O. M. Lyulin *et al.*, "Measurements of N₂- and O₂-broadening and shifting parameters of methane spectral lines in the 5550–6236 cm^{-1} region," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 110, no. 9–10, pp. 654–668, 2009, doi: 10.1016/j.jqsrt.2009.02.012.
- [61] A. V. Nikitin *et al.*, "GOSAT-2014 methane spectral line list," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 154, no. 2015, pp. 63–71, 2015, doi: 10.1016/j.jqsrt.2014.12.003.
- [62] E. Starikova *et al.*, "Assignment and modeling of the absorption spectrum of 13CH₄ at 80 K in the region of the 2v₃ band (5853–6201 cm^{-1})," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 177, pp. 170–180, 2016, doi: 10.1016/j.jqsrt.2015.12.023.
- [63] A. V. Nikitin *et al.*, "GOSAT-2009 methane spectral line list in the 5550–6236 cm^{-1} range," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 111, no. 15, pp. 2211–2224, 2010, doi: <https://doi.org/10.1016/j.jqsrt.2010.05.010>.
- [64] D. D. Tran, V. T. Sironneau, J. T. Hodges, R. Armante, J. Cuesta, and H. Tran, "Prediction of high-order line-shape parameters for air-broadened O₂ lines using requantized classical molecular dynamics simulations and comparison with measurements," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 222–223, 2019, doi: 10.1016/j.jqsrt.2018.10.013.
- [65] D. D. Tran, H. Tran, S. Vasilchenko, S. Kassı, A. Campargue, and D. Mondelain, "High sensitivity spectroscopy of the O₂ band at 1.27 μm : (II) air-broadened line profile parameters," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 240, no. II, p. 106673, 2020, doi: 10.1016/j.jqsrt.2019.106673.
- [66] R. Tóbiás *et al.*, "Critical evaluation of measured rotational–vibrational transitions of four sulphur isotopologues of S16O₂," *J. Quant. Spectrosc. Radiat. Transf.*, vol. 208, pp. 152–163, 2018, doi: <https://doi.org/10.1016/j.jqsrt.2018.01.006>.
- [67] D. S. Underwood *et al.*, "ExoMol molecular line lists - XIV. The rotation-vibration spectrum

of hot SO₂,” *Mon. Not. R. Astron. Soc.*, vol. 459, no. 4, pp. 3890–3899, 2016, doi: 10.1093/mnras/stw849.

- [68] I. Kovacs, *High Resolution Molecular Spectroscopy*, vol. 7, no. 12. 1976.
- [69] O. N. Ulenikov, E. S. Bekhtereva, O. V Gromova, V.-M. Horneman, C. Sydow, and S. Bauerecker, “High resolution FTIR spectroscopy of sulfur dioxide in the 1550–1950cm⁻¹ region: First analysis of the $\nu_1+\nu_2/\nu_2+\nu_3$ bands of 32S16O18O and experimental line intensities of ro-vibrational transitions in the $\nu_1+\nu_2/\nu_2+\nu_3$ bands of 32S16O2, 34S16O2, 32S18O2 and,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 203, pp. 377–391, 2017, doi: <https://doi.org/10.1016/j.jqsrt.2017.02.005>.
- [70] Y. G. Borkov *et al.*, “CO₂-broadening and shift coefficients of sulfur dioxide near 4 μm ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 225, pp. 119–124, 2019, doi: <https://doi.org/10.1016/j.jqsrt.2018.12.030>.
- [71] X. Huang, D. W. Schwenke, and T. J. Lee, “Quantitative validation of Ames IR intensity and new line lists for 32/33/34S16O₂, 32S18O₂ and 16O32S18O,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 225, pp. 327–336, 2019, doi: <https://doi.org/10.1016/j.jqsrt.2018.11.039>.
- [72] A. Perrin, L. Manceron, and F. Kwabia Tchana, “New line positions analysis of the $2\nu_1$ and $\nu_1+\nu_3$ bands of NO₂ at 3637.848 and 2906.070 cm⁻¹,” *Mol. Phys.*, vol. 8976, 2020, doi: 10.1080/00268976.2019.1711235.
- [73] A. Perrin *et al.*, “The new nitrogen dioxide (NO₂) linelist in the GEISA database and first identification of the $\nu_1+2\nu_3-\nu_3$ band of 14N16O₂,” *J. Mol. Spectrosc.*, vol. 376, p. 111394, Dec. 2020, doi: 10.1016/j.jms.2020.111394.
- [74] A. Perrin, G. Toon, and J. Orphal, “Detection of atmospheric 15NO₂ in the ν_3 spectral region (6.3 μm),” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 154, no. 2, pp. 91–97, 2015, doi: 10.1016/j.jqsrt.2014.12.006.
- [75] S. Miljanic, A. Perrin, and J. Orphal, “New high-resolution analysis of the $\{\nu_1 + \nu_3\}$ band of the 15N16O₂ isotopomer of nitrogen dioxide: Line positions and intensities,” *J. Mol. Spectrosc.*, vol. 242, no. 2, pp. 176–181, 2007, doi: 10.1016/j.jms.2007.02.024.
- [76] D. C. Benner, T. A. Blake, L. R. Brown, V. M. Devi, M. A. H. Smith, and R. A. Toth, “Air-broadening parameters in the ν_3 band of 14N 16O₂ using a multispectrum fitting technique,” *J. Mol. Spectrosc.*, vol. 228, no. 2 SPEC. ISS., pp. 593–619, 2004, doi: 10.1016/j.jms.2004.07.006.
- [77] R. J. Hargreaves *et al.*, “Spectroscopic line parameters of NO, NO₂, and N₂O for the HITEMP database,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 232, no. 2, pp. 35–53, 2019, doi: 10.1016/j.jqsrt.2019.04.040.
- [78] N. Maaroufi, C. Jalleli, F. K. Tchana, X. Landsheere, and H. Aroui, “Absolute line intensities and first measurements of self-collisional broadening and shift coefficients in the $2\nu_4$ band of NH₃,” *J. Mol. Spectrosc.*, vol. 354, pp. 24–31, 2018, doi: <https://doi.org/10.1016/j.jms.2018.09.003>.
- [79] V. Nemtchinov, K. Sung, and P. Varanasi, “Measurements of line intensities and half-widths in the 10- μm bands of 14NH₃,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 83, no. 3, pp. 243–265, 2004, doi: [https://doi.org/10.1016/S0022-4073\(02\)00354-0](https://doi.org/10.1016/S0022-4073(02)00354-0).
- [80] A. V. Nikitin *et al.*, “Analysis of PH₃ spectra in the Octad range 2733–3660 cm⁻¹,” *J.*

Quant. Spectrosc. Radiat. Transf., vol. 203, pp. 472–479, 2017, doi: 10.1016/j.jqsrt.2017.04.032.

- [81] T. J. Johnson, R. L. Sams, and S. W. Sharpe, “The PNNL quantitative infrared database for gas-phase sensing: a spectral library for environmental, hazmat, and public safety standoff detection,” *Chem. Biol. Point Sensors Homel. Def.*, vol. 5269, p. 159, 2004, doi: 10.1117/12.515604.
- [82] A. Perrin, “New analysis of the ν_3 and ν_4 bands of HNO₃ in the 7.6 μm region,” *J. Phys. Chem. A*, vol. 117, no. 50, pp. 13236–13248, 2013, doi: 10.1021/jp401979v.
- [83] A. Perrin, J. M. Flaud, M. Ridolfi, J. Vander Auwera, and M. Carlotti, “MIPAS database: New HNO₃ line parameters at 7.6 μm validated with MIPAS satellite measurements,” *Atmos. Meas. Tech.*, vol. 9, no. 5, pp. 2067–2076, 2016, doi: 10.5194/amt-9-2067-2016.
- [84] G. Li *et al.*, “Reference spectroscopic data for hydrogen halides. Part I: Construction and validation of the ro-vibrational dipole moment functions,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 121, pp. 78–90, May 2013, doi: 10.1016/J.JQSRT.2013.02.005.
- [85] G. Li, P. G. Hajigeorgiou, J. A. Coxon, and L. S. Rothman, “Reference spectroscopic data for hydrogen halides, Part II: The line lists,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 130, pp. 284–295, Nov. 2013, doi: 10.1016/J.JQSRT.2013.07.019.
- [86] G. Li, A. Serdyukov, M. Gisi, O. Werhahn, and V. Ebert, “FTIR-based measurements of self-broadening and self-shift coefficients as well as line strength in the first overtone band of HCl at 1.76 μM ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 165, pp. 76–87, Nov. 2015, doi: 10.1016/J.JQSRT.2015.06.021.
- [87] R. E. Asfin, A. V. Domanskaya, and C. Maul, “Broadening and shifting coefficients of rotation–vibrational lines in the fundamental and first overtone bands of HCl and HBr induced by oxygen and air,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 130, pp. 296–303, Nov. 2013, doi: 10.1016/J.JQSRT.2013.07.014.
- [88] B. Lejeune *et al.*, “Optimized approach to retrieve information on atmospheric carbonyl sulfide (OCS) above the Jungfraujoch station and change in its abundance since 1995,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 186, pp. 81–95, 2017, doi: <https://doi.org/10.1016/j.jqsrt.2016.06.001>.
- [89] N. G. Phillips *et al.*, “Mapping urban pipeline leaks: Methane leaks across Boston,” *Environ. Pollut.*, vol. 173, pp. 1–4, 2013, doi: <https://doi.org/10.1016/j.envpol.2012.11.003>.
- [90] D. Golebiowski, X. de Ghellinck d’Elsegheem Vaernewijck, M. Herman, J. Vander Auwera, and A. Fayt, “High sensitivity (femto-FT-CEAS) spectra of carbonyl sulphide between 6200 and 8200 cm^{-1} , and new energy pattern in the global rovibrational analysis of 16O12C32S,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 149, pp. 184–203, 2014, doi: <https://doi.org/10.1016/j.jqsrt.2014.07.005>.
- [91] S. Galalou, K. Ben Mabrouk, H. Aroui, F. Kwabia Tchana, F. Willaert, and J. M. Flaud, “N₂ and O₂ pressure broadening and pressure shift in the 4 ν_2 band of 16O12C32S,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 112, no. 18, pp. 2750–2761, 2011, doi: 10.1016/j.jqsrt.2011.08.008.
- [92] N. Moazzen-Ahmadi *et al.*, “An intensity study of the torsional bands of ethane at 35 μm ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 151, pp. 123–132, Jan. 2015, doi: 10.1016/j.jqsrt.2014.09.016.

- [93] O. M. Lyulin and V. I. Perevalov, “Global modeling of vibration-rotation spectra of the acetylene molecule,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 177, pp. 59–74, 2016, doi: 10.1016/j.jqsrt.2015.12.021.
- [94] O. M. Lyulin and V. I. Perevalov, “ASD-1000: High-resolution, high-temperature acetylene spectroscopic databank,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 201, pp. 94–103, 2017, doi: 10.1016/j.jqsrt.2017.06.032.
- [95] D. Jacquemart, O. Lyulin, and V. I. Perevalov, “Recommended acetylene line list in the 20–240 cm^{-1} and 400–630 cm^{-1} regions: New measurements and global modeling,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 203, pp. 440–453, 2017, doi: 10.1016/j.jqsrt.2017.03.008.
- [96] D. Jacquemart, P. Soulard, and O. M. Lyulin, “Recommended acetylene $12\text{C}_2\text{H}_2$ line list in 13.6 μm spectral region: New measurements and global modeling,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 256, p. 107200, 2020, doi: 10.1016/j.jqsrt.2020.107200.
- [97] O. M. Lyulin and A. Campargue, “An empirical spectroscopic database for acetylene in the regions of 5850–6341 cm^{-1} and 7000–9415 cm^{-1} ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 203, pp. 461–471, 2017, doi: 10.1016/j.jqsrt.2017.01.036.
- [98] O. M. Lyulin, J. Vander Auwera, and A. Campargue, “The Fourier transform absorption spectrum of acetylene between 8280 and 8700 cm^{-1} ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 177, pp. 234–240, 2016, doi: 10.1016/j.jqsrt.2015.11.026.
- [99] O. M. Lyulin, J. Vander Auwera, and A. Campargue, “The Fourier transform absorption spectrum of acetylene between 7000 and 7500 cm^{-1} ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 160, pp. 85–93, 2015, doi: 10.1016/j.jqsrt.2015.03.018.
- [100] S. Béguier, O. M. Lyulin, S. M. Hu, and A. Campargue, “Line intensity measurements for acetylene between 8980 and 9420 cm^{-1} ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 189, pp. 417–420, 2017, doi: 10.1016/j.jqsrt.2016.12.020.
- [101] O. M. Lyulin, A. Campargue, D. Mondelain, and S. Kassi, “The absorption spectrum of acetylene by CRDS between 7244 and 7918 cm^{-1} ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 130, pp. 327–334, 2013, doi: 10.1016/j.jqsrt.2013.04.028.
- [102] O. M. Lyulin, D. Mondelain, S. Béguier, S. Kassi, J. Vander Auwera, and A. Campargue, “High-sensitivity CRDS absorption spectroscopy of acetylene between 5851 and 6341 cm^{-1} ,” *Mol. Phys.*, vol. 112, no. 18, pp. 2433–2444, 2014, doi: 10.1080/00268976.2014.906677.
- [103] S. Kassi, O. M. Lyulin, S. Béguier, and A. Campargue, “New assignments and a rare peculiarity in the high sensitivity CRDS spectrum of acetylene near 8000 cm^{-1} ,” *J. Mol. Spectrosc.*, vol. 326, pp. 106–114, 2016, doi: 10.1016/j.jms.2016.02.013.
- [104] M. Šimečková, D. Jacquemart, L. S. Rothman, R. R. Gamache, and A. Goldman, “Einstein A-coefficients and statistical weights for molecular absorption transitions in the HITRAN database,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 98, no. 1, pp. 130–155, 2006, doi: <https://doi.org/10.1016/j.jqsrt.2005.07.003>.
- [105] R. R. Gamache *et al.*, “Total internal partition sums for 166 isotopologues of 51 molecules important in planetary atmospheres: Application to HITRAN2016 and beyond,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 203, pp. 70–87, 2017, doi: 10.1016/j.jqsrt.2017.03.045.
- [106] O. M. Lyulin, D. Jacquemart, N. Lacome, V. I. Perevalov, and J. Y. Mandin, “Line parameters of acetylene in the 1.9 and 1.7 μm spectral regions,” *J. Quant. Spectrosc. Radiat.*

Transf., vol. 109, no. 10, pp. 1856–1874, 2008, doi: 10.1016/j.jqsrt.2007.11.016.

- [107] A. Alkadrou, M.-T. Bourgeois, M. Rotger, V. Boudon, and J. Vander Auwera, “Corrigendum to ‘Global frequency and intensity analysis of the $\nu_{10}/\nu_7/\nu_4/\nu_{12}$ band system of $12\text{C}_2\text{H}_4$ at $10\mu\text{m}$ using the D2h top data system’ [J. Quant. Spectrosc. Radiat. Transf. 182 (2016) 158–171],” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 190, p. 88, 2017, doi: <https://doi.org/10.1016/j.jqsrt.2017.01.007>.
- [108] A. Alkadrou, M.-T. Bourgeois, M. Rotger, V. Boudon, and J. Vander Auwera, “Global frequency and intensity analysis of the $\nu_{10}/\nu_7/\nu_4/\nu_{12}$ band system of $12\text{C}_2\text{H}_4$ at $10\mu\text{m}$ using the D2h Top Data System,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 182, pp. 158–171, 2016, doi: <https://doi.org/10.1016/j.jqsrt.2016.05.024>.
- [109] J. Vander Auwera *et al.*, “Self-broadening coefficients and improved line intensities for the ν_7 band of ethylene near $10.5\mu\text{m}$, and impact on ethylene retrievals from Jungfraujoch solar spectra,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 148, pp. 177–185, 2014, doi: <https://doi.org/10.1016/j.jqsrt.2014.07.003>.
- [110] M. Rotger, V. Boudon, and J. Vander Auwera, “Line positions and intensities in the ν_{12} band of ethylene near 1450cm^{-1} : An experimental and theoretical study,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 109, no. 6, pp. 952–962, 2008, doi: <https://doi.org/10.1016/j.jqsrt.2007.12.005>.
- [111] U. Fink, H. P. Larson, and R. R. Treffers, “Germane in the atmosphere of Jupiter,” *Icarus*, vol. 34, no. 2, pp. 344–354, May 1978, doi: 10.1016/0019-1035(78)90172-0.
- [112] K. S. Noll, R. F. Knacke, T. R. Geballe, and A. T. Tokunaga, “Evidence for Germane in Saturn,” *Icarus*, vol. 75, no. 3, pp. 409–422, Sep. 1988, doi: 10.1016/0019-1035(88)90154-6.
- [113] V. Boudon *et al.*, “Line positions and intensities for the ν_3 band of 5 isotopologues of germane for planetary applications,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 205, pp. 174–183, Jan. 2018, doi: 10.1016/J.JQSRT.2017.10.017.
- [114] “<http://vamdc.icb.cnrs.fr/PHP/gecasda.php>.” .
- [115] C. Richard, V. Boudon, and M. Rotger, “Calculated spectroscopic databases for the VAMDC portal: New molecules and improvements,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 251, p. 107096, 2020, doi: 10.1016/j.jqsrt.2020.107096.
- [116] M. López-Puertas, J.-M. Flaud, J. Peralta-Calvillo, B. Funke, and S. Gil-López, “NO⁺ fundamental and first hot ro-vibrational line frequencies from MIPAS/Envisat atmospheric spectra,” *J. Mol. Spectrosc.*, vol. 237, no. 2, pp. 218–224, 2006, doi: <https://doi.org/10.1016/j.jms.2006.03.015>.
- [117] J. Cernicharo *et al.*, “Tentative detection of the nitrosylium ion in space,” *Astrophys. J.*, vol. 795, no. 1, pp. 6–11, 2014, doi: 10.1088/0004-637X/795/1/40.
- [118] C. P. Endres, S. Schlemmer, P. Schilke, J. Stutzki, and H. S. P. Müller, “The Cologne Database for Molecular Spectroscopy, CDMS, in the Virtual Atomic and Molecular Data Centre, VAMDC,” *J. Mol. Spectrosc.*, vol. 327, pp. 95–104, 2016, doi: <https://doi.org/10.1016/j.jms.2016.03.005>.
- [119] W. C. Bowman, E. Herbst, and F. C. De Lucia, “Millimeter and submillimeter spectrum of NO⁺,” *J. Chem. Phys.*, vol. 77, no. 8, pp. 4261–4262, Oct. 1982, doi: 10.1063/1.444307.
- [120] W. C. Ho, I. Ozier, D. T. Cramb, and M. C. L. Gerry, “Diode laser spectroscopy of the

vibrational fundamental of NO⁺,” *J. Mol. Spectrosc.*, vol. 149, no. 2, pp. 559–561, 1991, doi: [https://doi.org/10.1016/0022-2852\(91\)90311-W](https://doi.org/10.1016/0022-2852(91)90311-W).

- [121] R. Polák and J. Fišer, “A comparative icMRCI study of some NO⁺, NO and NO[−] electronic ground state properties,” *Chem. Phys.*, vol. 303, no. 1, pp. 73–83, 2004, doi: <https://doi.org/10.1016/j.chemphys.2004.04.027>.
- [122] C. P. Rinsland *et al.*, “Long-term stratospheric carbon tetrafluoride (CF₄) increase inferred from 1985-2004 infrared space-based solar occultation measurements,” *Geophys. Res. Lett.*, vol. 33, no. 2, pp. 2–5, 2006, doi: [10.1029/2005GL024709](https://doi.org/10.1029/2005GL024709).
- [123] E. Mahieu *et al.*, “Spectrometric monitoring of atmospheric carbon tetrafluoride (CF₄) above the Jungfraujoch station since 1989: Evidence of continued increase but at a slowing rate,” *Atmos. Meas. Tech.*, vol. 7, no. 1, pp. 333–344, 2014, doi: [10.5194/amt-7-333-2014](https://doi.org/10.5194/amt-7-333-2014).
- [124] M. Carlos *et al.*, “High-resolution spectroscopy and global analysis of CF₄ rovibrational bands to model its atmospheric absorption,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 201, pp. 75–93, 2017, doi: <https://doi.org/10.1016/j.jqsrt.2017.06.039>.
- [125] V. Boudon, M. Carlos, C. Richard, and O. Pirali, “Pure rotation spectrum of CF₄ in the v₃=1 state using THz synchrotron radiation,” *J. Mol. Spectrosc.*, vol. 348, pp. 43–46, 2018, doi: <https://doi.org/10.1016/j.jms.2017.07.010>.
- [126] “<http://vamdc.icb.cnrs.fr/PHP/tfmecasda.php>.”
- [127] R. Armante, A. Perrin, F. Kwabia-Tchana, and L. Manceron, “The v₄ linelist at 11 μm: linelists for the Trans- and Cis- conformer forms of nitrous acid, (HONO), in the 2019 version of the GEISA database,” *Mol. Phys.*, vol. submitted, 2021.
- [128] I. Kleiner, J. M. Guilmot, M. Carleer, and M. Herman, “The v₄ fundamental bands of trans- and cis-HNO₂,” *J. Mol. Spectrosc.*, vol. 149, no. 2, pp. 341–347, 1991, doi: [https://doi.org/10.1016/0022-2852\(91\)90290-Q](https://doi.org/10.1016/0022-2852(91)90290-Q).
- [129] A. Dehayem-Kamadjeu, O. Pirali, J. Orphal, I. Kleiner, and P.-M. Flaud, “The far-infrared rotational spectrum of nitrous acid (HONO) and its deuterated species (DONO) studied by high-resolution Fourier-transform spectroscopy,” *J. Mol. Spectrosc.*, vol. 234, no. 1, pp. 182–189, 2005, doi: <https://doi.org/10.1016/j.jms.2005.09.006>.
- [130] R. H. Kagann and A. G. Maki, “Infrared absorption intensities of nitrous acid (HONO) fundamental bands,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 30, no. 1, pp. 37–44, 1983, doi: [https://doi.org/10.1016/0022-4073\(83\)90071-7](https://doi.org/10.1016/0022-4073(83)90071-7).
- [131] A. Perrin, J. Demaison, and G. Toon, “The v₁, v₂, and v₃ bands of carbonyl chlorofluoride (COFCl) at 5.3, 9.1, and 13.1 μm: position and intensity parameters and their use for atmospheric studies,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 112, no. 8, pp. 1266–1279, 2011, doi: <https://doi.org/10.1016/j.jqsrt.2011.01.003>.
- [132] A. Perrin, J.-M. Flaud, H. Bürger, G. Pawelke, S. Sander, and H. Willner, “First High-Resolution Analysis of the Six Fundamental Bands v₁, v₂, v₃, v₄, v₅, and v₆ of COF₃₅Cl in the 340- to 2000-cm^{−1} Region,” *J. Mol. Spectrosc.*, vol. 209, no. 1, pp. 122–132, 2001, doi: <https://doi.org/10.1006/jmsp.2001.8415>.
- [133] J. Demaison, A. Perrin, and H. Bürger, “Ab initio anharmonic force field and equilibrium structure of carbonyl chlorofluoride,” *J. Mol. Spectrosc.*, vol. 221, no. 1, pp. 47–56, 2003, doi: [https://doi.org/10.1016/S0022-2852\(03\)00169-3](https://doi.org/10.1016/S0022-2852(03)00169-3).

- [134] A. Perrin, I. Haykal, F. KwabiaTchana, L. Manceron, D. Doizi, and G. Ducros, “New analysis of the ν_6 and $2\nu_3$ bands of methyl iodide (CH_3I),” *J. Mol. Spectrosc.*, vol. 324, pp. 28–35, 2016, doi: <https://doi.org/10.1016/j.jms.2016.04.014>.
- [135] F. Kwabia-Tchana, Y. Attafi, L. Manceron, D. Doizi, J. Vander Auwera, and A. Perrin, “Line intensities for the ν_6 and $2\nu_3$ bands of methyl iodide ($12\text{CH}_3\text{I}$),” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 222–223, pp. 130–137, 2019, doi: <https://doi.org/10.1016/j.jqsrt.2018.10.001>.
- [136] Y. Attafi *et al.*, “Self and N_2 collisional broadening of rovibrational lines in the ν_6 band of methyl iodide ($12\text{CH}_3\text{I}$) at room temperature: The J and K dependence,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 231, pp. 1–8, 2019, doi: <https://doi.org/10.1016/j.jqsrt.2019.04.017>.
- [137] Y. Attafi *et al.*, “Oxygen broadening and shift coefficients in the ν_6 band of methyl iodide ($12\text{CH}_3\text{I}$) at room temperature,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 239, p. 106679, 2019, doi: <https://doi.org/10.1016/j.jqsrt.2019.106679>.
- [138] D. Papoušek, P. Pracna, M. Winnewisser, S. Klee, and J. Demaison, “Simultaneous Rovibrational Analysis of the ν_2 , ν_3 , ν_5 , and ν_6 Bands of H_3^{12}CF ,” *J. Mol. Spectrosc.*, vol. 196, no. 2, pp. 319–323, 1999, doi: <https://doi.org/10.1006/jmsp.1999.7875>.
- [139] D. Jacquemart and M. Guinet, “Line parameters measurements and modeling for the ν_6 band of CH_3F : Generation of a complete line list for atmospheric databases,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 185, pp. 58–69, 2016, doi: <https://doi.org/10.1016/j.jqsrt.2016.08.010>.
- [140] A. B. Ramchani, D. Jacquemart, P. Soulard, and M. Guinet, “Measurements and modeling of N_2 -broadening coefficients for the ν_6 band of CH_3F , comparison with CH_3Cl and CH_3Br molecules,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 203, pp. 480–489, 2017, doi: <https://doi.org/10.1016/j.jqsrt.2017.06.013>.
- [141] S. Reymond-Laruinaz, V. Boudon, L. Manceron, L. Lago, and D. Doizi, “Infrared spectroscopy of ruthenium tetroxide and high-resolution analysis of the ν_3 band,” *J. Mol. Spectrosc.*, vol. 315, pp. 46–54, Sep. 2015, doi: [10.1016/J.JMS.2015.02.008](https://doi.org/10.1016/J.JMS.2015.02.008).
- [142] S. Reymond-Laruinaz, M. Faye, V. Boudon, D. Doizi, and L. Manceron, “High-resolution infrared spectroscopy and analysis of the ν_2/ν_4 bending dyad of ruthenium tetroxide,” *J. Mol. Spectrosc.*, vol. 336, pp. 29–35, Jun. 2017, doi: [10.1016/J.JMS.2017.04.010](https://doi.org/10.1016/J.JMS.2017.04.010).
- [143] J. Vander Auwera, S. Reymond-Laruinaz, V. Boudon, D. Doizi, and L. Manceron, “Line intensity measurements and analysis in the ν_3 band of ruthenium tetroxide,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 204, pp. 103–111, Jan. 2018, doi: [10.1016/J.JQSRT.2017.09.016](https://doi.org/10.1016/J.JQSRT.2017.09.016).
- [144] “<http://vamdc.icb.cnrs.fr/PHP/rucasda.php>.”
- [145] N. A. Lombardo *et al.*, “Spatial and seasonal variations in C_3H_x hydrocarbon abundance in Titan’s stratosphere from Cassini CIRS observations,” *Icarus*, vol. 317, pp. 454–469, 2019, doi: <https://doi.org/10.1016/j.icarus.2018.08.027>.
- [146] N. A. Lombardo *et al.*, “Detection of Propadiene on Titan,” *Astrophys. J.*, vol. 881, no. 2, p. L33, 2019, doi: [10.3847/2041-8213/ab3860](https://doi.org/10.3847/2041-8213/ab3860).
- [147] A. Coustenis *et al.*, “Modeling Titan’s Thermal Infrared Spectrum for High-Resolution Space Observations,” *Icarus*, vol. 102, no. 2, pp. 240–260, 1993, doi: <https://doi.org/10.1006/icar.1993.1047>.
- [148] J. Chazelas, J. Plíva, A. Valentin, and L. Henry, “Analysis of the $\nu_9\nu_{10}$ band system of

allene,” *J. Mol. Spectrosc.*, vol. 110, no. 2, pp. 326–338, 1985, doi: [https://doi.org/10.1016/0022-2852\(85\)90298-X](https://doi.org/10.1016/0022-2852(85)90298-X).

- [149] J. Plíva and J. Kauppinen, “High-resolution Fourier transform study of the perpendicular band ν_{11} of allene at 353 cm^{-1} ,” *J. Mol. Spectrosc.*, vol. 111, no. 1, pp. 93–101, 1985, doi: [https://doi.org/10.1016/0022-2852\(85\)90072-4](https://doi.org/10.1016/0022-2852(85)90072-4).
- [150] Y. Koga, S. Kondo, T. Nakanaga, and S. Saëki, “Infrared absorption intensities of allene,” *J. Chem. Phys.*, vol. 71, no. 6, pp. 2404–2411, 1979, doi: 10.1063/1.438645.
- [151] A. Jolly, Y. Benilan, and L. Manceron, “Search for evidence of Butane on Titan with new spectroscopic data,” *EPSC Abstr.*, vol. 11, pp. 10–11, 2017.
- [152] S. Nissen, F. Hegelund, M. S. Johnson, and B. Nelander, “High-Resolution Infrared Study of the ν_{11} Band of Allene,” *J. Mol. Spectrosc.*, vol. 216, no. 2, pp. 197–202, 2002, doi: <https://doi.org/10.1006/jmsp.2002.8596>.
- [153] F. Hegelund, N. Andresen, and M. Koivusaari, “A High-Resolution Infrared Study of the $\nu_9 + \nu_{11} - \nu_{11}$, $\nu_{10} + \nu_{11} - \nu_{11}$ Hot Band System in Allene,” *J. Mol. Spectrosc.*, vol. 159, no. 1, pp. 230–248, 1993, doi: <https://doi.org/10.1006/jmsp.1993.1120>.
- [154] E. Es-sebbar, A. Jolly, Y. Benilan, and A. Farooq, “Quantitative mid-infrared spectra of allene and propyne from room to high temperatures,” *J. Mol. Spectrosc.*, vol. 305, pp. 10–16, 2014, doi: <https://doi.org/10.1016/j.jms.2014.09.004>.
- [155] N. A. Scott and A. Chedin, “A Fast Line-by-Line Method for Atmospheric Absorption Computations: The Automatized Atmospheric Absorption Atlas,” *J. Appl. Meteorol. Climatol.*, vol. 20, no. 7, pp. 802–812, doi: 10.1175/1520-0450(1981)020<0802:AFLBLM>2.0.CO;2.
- [156] F. Cheruy, N. A. Scott, R. Armante, B. Tournier, and A. Chedin, “Contribution to the development of radiative transfer models for high spectral resolution observations in the infrared,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 53, no. 6, 1995, doi: 10.1016/0022-4073(95)00026-H.
- [157] L. Regalia *et al.*, “Water vapor line parameters from 6450 to 9400 cm^{-1} ,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 136, pp. 119–136, 2014, doi: 10.1016/j.jqsrt.2013.11.019.
- [158] P. Macko *et al.*, “High sensitivity CW-cavity ring down spectroscopy of water in the region of the $1.5\text{ }\mu\text{m}$ atmospheric window,” *J. Mol. Spectrosc.*, vol. 227, no. 1, pp. 90–108, 2004, doi: 10.1016/j.jms.2004.05.020.
- [159] R. a Toth, “Line lists of water vapor parameters from 500 to 8000 cm^{-1} ,” *JPL Nasa*, vol. (to be pub, no. 818, 2013.
- [160] R. R. Gamache and A. L. Laraia, “ N_2 -, O_2 -, and air-broadened half-widths, their temperature dependence, and line shifts for the rotation band of H_2O ,” *J. Mol. Spectrosc.*, vol. 257, no. 2, pp. 116–127, 2009, doi: <https://doi.org/10.1016/j.jms.2009.07.004>.
- [161] R. R. Gamache, “Lineshape parameters for water vapor in the 3.2 – $17.76\text{ }\mu\text{m}$ region for atmospheric applications,” *J. Mol. Spectrosc.*, vol. 229, no. 1, pp. 9–18, 2005, doi: <https://doi.org/10.1016/j.jms.2004.08.004>.
- [162] D. P. Dee *et al.*, “The ERA-Interim reanalysis: configuration and performance of the data assimilation system,” *Q. J. R. Meteorol. Soc.*, vol. 137, no. 656, pp. 553–597, 2011, doi: <https://doi.org/10.1002/qj.828>.

- [163] H. Hersbach *et al.*, “The ERA5 global reanalysis,” *Q. J. R. Meteorol. Soc.*, vol. 146, no. 730, pp. 1999–2049, 2020, doi: <https://doi.org/10.1002/qj.3803>.
- [164] H. Ke, V. Boudon, C. Richard, V. Madhur, M. Faye, and L. Manceron, “Analysis and modeling of combination bands of sulfur hexafluoride $^{32}\text{SF}_6$ based on global fits. Update of the SHeCaSDa database,” *J. Mol. Spectrosc.*, vol. 368, pp. 1–14, 2020, doi: [10.1016/j.jms.2020.111251](https://doi.org/10.1016/j.jms.2020.111251).
- [165] A. V. Nikitin, M. Rey, I. S. Chizhmakova, and V. G. Tyuterev, “First Full-Dimensional Potential Energy and Dipole Moment Surfaces of SF_6 ,” *J. Phys. Chem. A*, vol. 124, no. 35, pp. 7014–7023, 2020, doi: [10.1021/acs.jpca.0c02733](https://doi.org/10.1021/acs.jpca.0c02733).
- [166] M. Rey, A. V. Nikitin, Y. L. Babikov, and V. G. Tyuterev, “TheoReTS – An information system for theoretical spectra based on variational predictions from molecular potential energy and dipole moment surfaces,” *J. Mol. Spectrosc.*, vol. 327, pp. 138–158, 2016, doi: [10.1016/j.jms.2016.04.006](https://doi.org/10.1016/j.jms.2016.04.006).
- [167] S. W. Sharpe, T. J. Johnson, R. L. Sams, P. M. Chu, G. C. Rhoderick, and P. A. Johnson, “Gas-Phase Databases for Quantitative Infrared Spectroscopy,” *Appl. Spectrosc.*, vol. 58, no. 12, pp. 1452–1461, 2004.
- [168] N. H. Ngo, D. Lisak, H. Tran, and J. M. Hartmann, “An isolated line-shape model to go beyond the Voigt profile in spectroscopic databases and radiative transfer codes,” *J. Quant. Spectrosc. Radiat. Transf.*, vol. 129, pp. 89–100, 2013, doi: [10.1016/j.jqsrt.2013.05.034](https://doi.org/10.1016/j.jqsrt.2013.05.034).