



Importance of access to epilepsy monitoring units during the COVID-19 pandemic: Consensus statement of the International League against epilepsy and the International Federation of Clinical Neurophysiology [☆]



Sándor Beniczky ^{a,*}, Aatif Husain ^{b,c}, Akio Ikeda ^d, Haifa Alabri ^e, J. Helen Cross ^f, Jo Wilmschurst ^g, Margitta Seeck ^h, Niels Focke ⁱ, Patricia Braga ^j, Samuel Wiebe ^k, Stephan Schuele ^l, Eugen Trinka ^{m,n,o}

^a Department of Clinical Neurophysiology, Neurology, Aarhus University Hospital, Aarhus and Danish Epilepsy Centre, Dianalund, Denmark

^b Department of Neurology, Duke University Medical Center, Durham, NC, USA

^c Neurodiagnostic Center, Veterans Affairs Medical Center, Durham, NC, USA

^d Department of Epilepsy, Movement Disorders and Physiology, Kyoto University Graduate School of Medicine, Kyoto, Japan

^e Neurology Unit, Department of Medicine, Sultan Qaboos University Hospital, Sultan Qaboos University, Oman

^f UCL NIHR BRC Great Ormond Street Institute of Child Health, Great Ormond Street Hospital for Children, London, Member of ERN EpiCARE, and Young Epilepsy Lingfield, UK

^g Department of Paediatric Neurology, Red Cross War Memorial Children's Hospital, Neuroscience Institute, University of Cape Town, South Africa

^h EEG & Epilepsy Unit, University Hospital of Geneva, Medical Faculty of the University of Geneva, Switzerland

ⁱ Department of Neurology, University Medical Center, Göttingen, Germany

^j Institute of Neurology, Hospital de Clínicas, Facultad de Medicina, Universidad de la República, Uruguay

^k Department of Clinical Neurosciences, University of Calgary, Canada

^l Department of Neurology, Northwestern University, Feinberg School of Medicine, Chicago, USA

^m Department of Neurology, Christian Doppler University Hospital, Paracelsus Medical University and Centre for Cognitive Neuroscience, Salzburg, Austria

ⁿ Affiliated EpiCARE Partner, Neuroscience Institute, Christian Doppler University Hospital, Salzburg, Austria

^o Department of Public Health, Health Services Research and Health Technology Assessment, UMIT – University for Health Sciences, Medical Informatics and Technology, Hall in Tirol, Austria

ARTICLE INFO

Article history:

Available online 21 July 2021

Keywords:

COVID-19

Epilepsy

Epilepsy monitoring unit

Video-EEG

HIGHLIGHTS

- The COVID-19 pandemic has led to lockdown of many Epilepsy Monitoring Units.
- Postponing video-EEG monitoring of patients had detrimental effects.
- Access to video-EEG monitoring be given high priority.

ABSTRACT

Restructuring of healthcare services during the COVID-19 pandemic has led to lockdown of Epilepsy Monitoring Units (EMUs) in many hospitals. The ad-hoc taskforce of the International League Against Epilepsy (ILAE) and the International Federation of Clinical Neurophysiology (IFCN) highlights the detrimental effect of postponing video-EEG monitoring of patients with epilepsy and other paroxysmal events. The taskforce calls for action to continue functioning of Epilepsy Monitoring Units during emergency situations, such as the COVID-19 pandemic. Long-term video-EEG monitoring is an essential diagnostic service. Access to video-EEG monitoring of the patients in the EMUs must be given high priority. Patients should be screened for COVID-19, before admission, according to the local regulations. Local policies for COVID-19 infection control should be adhered to during the video-EEG monitoring. In cases of differential diagnosis where reduction of antiseizure medication is not required, consider home video-EEG monitoring as an alternative in selected patients.

© 2021 International Federation of Clinical Neurophysiology Inc. and International League Against Epilepsy. Published by Elsevier B.V. All rights reserved. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

[☆] This article also appears in issue 23/4 of the ILAE's journal, *Epileptic Disorders*, published by John Libbey Eurotext.

* Corresponding author at: Visby Allé 5, 4293 Dianalund, Denmark.

E-mail address: sbz@filadelfia.dk (S. Beniczky).

1. Background and justification

The International League Against Epilepsy (ILAE) and the International Federation of Clinical Neurophysiology (IFCN) appointed an ad-hoc taskforce to provide a rapid response to the challenges concerning video-EEG monitoring, encountered during the current pandemic, caused by the coronavirus SARS-CoV-2 disease (COVID-19). Through consensus discussions, review of the published evidence and experience of the experts representing the two societies, the ad-hoc taskforce elaborated this statement.

During the restructuring of healthcare services due to the current pandemic, many hospitals closed video-EEG monitoring facilities, referred to in this document as Epilepsy Monitoring Units (EMU) (Krysl et al., 2020). A European survey showed that in most centers, inpatient video-EEGs monitoring had been stopped (61.7% for adults, 36.2% for children) or was restricted (38.3% for adults, 53.2% for children) (Krysl et al., 2020), with detrimental effects on patients with complex and severe epilepsy and other paroxysmal events (Krysl et al., 2020), such as lack of optimizing the medical treatment and lack of evaluation for epilepsy surgery. There is limited likelihood of triggering seizure emergencies in patients with epilepsy and neurological complications. The measures of closing EMUs were adopted by healthcare providers to focus on re-allocation of resources to services considered more important, more immediately required, and to prevent spreading the disease. Long-term video-EEG monitoring in EMUs was regarded by the healthcare providers as an elective procedure that could be postponed without significant consequences, a categorization that we challenge as incorrect for the following reasons.

Long-term video-EEG monitoring is an essential diagnostic tool in patients with complex and severe epilepsy (Kobulashvili et al., 2018; Tatum et al., 2018). The main indications are: diagnostic and presurgical evaluation (Kobulashvili et al., 2018; Tatum et al., 2018). While video-EEG monitoring is diagnostic, it has direct implications on treatment of epileptic seizures, comorbidities and important differential diagnoses (arrhythmia and cardiac death, psychogenic non-epileptic seizures and the risk of suicide).

Reasoning for continuing the diagnostic monitoring: approximately one third of patients referred to specialized centers on suspicion of drug-resistant epilepsy, do not have epilepsy (Chadwick & Smith, 2002; McBride et al., 2002; Asano et al., 2005; Uldall et al., 2006). Persistent misdiagnosis of paroxysmal events, often cardiac or psychogenic in origin, has severe consequences for them (Ferrie, 2006; LaFrance & Benbadis, 2006; Nightscales et al., 2020). In patients with drug resistant epilepsy, misclassification of the seizure-types can lead to inadequate choice of antiseizure medication (Kobulashvili et al., 2018). Video-EEG recording of the patients' habitual clinical episodes is the diagnostic gold standard for patients with unclear paroxysmal events (Kobulashvili et al., 2018; Tatum et al., 2018).

Reasoning for continuing presurgical evaluation: epilepsy surgery is the evidence-based treatment in patients with drug-resistant focal epilepsy (Wiebe et al., 2001; Engel et al., 2012; Dwivedi et al., 2017). This requires video-EEG recording of the seizure and in around one third of patients invasive monitoring (Kobulashvili et al., 2018; Tatum et al., 2018). Failure to proceed towards surgery, unnecessarily exposes the patients to further seizures, injuries associated with seizures and the risk of Sudden Unexpected Death in Epilepsy (SUDEP) (Devinsky et al., 2016). The appropriate and unrestricted utilization of EMUs in comprehensive epilepsy centers has been shown to reduce mortality of patients with epilepsy (Lowerison et al., 2019).

High quality epilepsy care, including video-EEG monitoring has decreased morbidity and mortality (Lowerison et al., 2019; Hargreaves et al., 2019; Granbichler et al., 2017). Hence, increasing

waiting times can cause considerable problems, with increasing morbidity and mortality. These patients often have worsening epilepsy, co-morbidities, and prioritizing care with restricted resources becomes more and more challenging. Some EMUs managed to continue video-EEG monitoring during the pandemic (Krysl et al., 2020). Using measures of prevention and protection generally adopted in the hospitals, these EMUs were able to continue this important diagnostic function, without causing local outbreaks (Lowerison et al., 2019; Krysl et al., 2020; Joint proposals, 2020; Gélisse et al., 2020). Recommendations for neurophysiology staff with risk factors for COVID-19, and for mental health of the staff have been proposed by the Latin American chapter of the IFCN Task Force – COVID-19 (San-Juan et al., 2020).

2. Summary statements

The ILAE-IFCN ad-hoc taskforce issues the following statement, related to functioning of the EMUs, during the COVID-19 pandemic:

1. Long-term video-EEG monitoring is an essential diagnostic service.
2. Access to video-EEG monitoring of the patients in the EMUs must be given high priority.
3. Patients should be screened for COVID-19, before admission, according to the local regulations.
4. Local policies for COVID-19 infection control should be adhered to during the video-EEG monitoring.
5. In cases of differential diagnosis where reduction of antiseizure medication is not required, consider home video-EEG monitoring as an alternative in selected patients.

3. Conclusion

The ILAE-IFCN ad-hoc taskforce calls for action to ensure that healthcare providers understand the importance of providing diagnostic services for patients with epilepsy and paroxysmal events, and that EMUs continue functioning during emergency situations like the COVID-19 pandemic, while adhering to local healthcare policies.

4. Disclaimer

This report was written by experts selected by the International League Against Epilepsy (ILAE) and the International Federation of Clinical Neurophysiology (IFCN) and was approved for publication by the ILAE and IFCN. Opinions expressed by the authors, however, do not necessarily represent the policy or position of the ILAE and IFCN.

5. Conflict of interest statement

Sándor Beniczky reports speaker honoraria and personal fees from: Natus Neuro, Philips EGI, Epihunter, UCB Pharma, Eisai and Bial-Portela, outside this work.

Aatif Husain has received personal compensation for consulting from: UCB Pharma, Jazz Pharma, BlackThorn Therapeutics, Eisai Pharmaceuticals, Marinus Pharmaceuticals, Neurelis Pharmaceuticals, and Merck. He has also received royalties from Wolters Kluwer, Springer and Demos Publishers. Additionally, he has received stipend for editorship from American Clinical Neurophysiology Society.

Akio IKEDA belongs to Department of Epilepsy, Movement Disorders and Physiology which is the Industry-Academia Collaboration Courses, supported by Eisai Co., Ltd., Nihon Kohden Corpora-

tion, Otsuka Pharmaceutical Co., and UCB Japan Co., Ltd. He also receives honorarium from Eisai Co., Ltd and UCB Japan Co., Ltd

J. Helen Cross has acted as an investigator for studies with GW Pharma, Zogenix, Vitaflor and Marinus. She has been a speaker and on advisory boards for GW Pharma, Zogenix, and Nutricia; all remuneration has been paid to her department. Her research is supported by the National Institute of Health Research (NIHR) Biomedical Research Centre at Great Ormond Street Hospital. She holds an endowed chair at UCL Great Ormond Street Institute of Child Health; she holds grants from NIHR, EPSRC, GOSH Charity, ERUK, and the Waterloo Foundation

Jo Wilmshurst receives an honorarium for her role of associate editor for *Epilepsia*.

M Seck received speaker's fees of EGI Philips, holds shares of Epilog and received consulting fees from the Wyss Foundation.

N. K. Focke received speaker bureaus and consultancy fees from Bial, Eisai and EGI/Phillips outside the submitted work.

Patricia Braga reports no conflicts of interest.

Samuel Wiebe, through his institution, has received unrestricted educational grants from UCB Pharma, Sunovion and Eisai.

Dr. Schuele has received personal compensation in form of a stipend from Wolters Kluwer as Associate Editor of the *Journal of Clinical Neurophysiology*. He receives royalties from a book published from Demos Publishers Inc., Springer Publishing on Stereoecephalography. Honoraria for speaking engagements have been received from the American Academy of Neurology, American Epilepsy Society, American Clinical Neurophysiology Society, Sunovion Inc., SK Life Science, Neurelis, and Greenwich.

Eugen Trinka reports personal fees from EVER Pharma, Marinus, Arvelle, Argenix, Medtronic, Bial-Portela & C^o, NewBridge, GL Pharma, GlaxoSmithKline, Boehringer Ingelheim, LivaNova, Eisai, UCB, Biogen, Genzyme Sanofi, and Actavis; his institution received grants from Biogen, UCB Pharma, Eisai, Red Bull, Merck, Bayer, the European Union, FWF Österreichischer Fond zur Wissenschaftsförderung, Bundesministerium für Wissenschaft und Forschung, and Jubiläumsfond der Österreichischen Nationalbank outside the submitted work.

References

Asano E, Pawlak C, Shah A, et al. The diagnostic value of initial video-EEG monitoring in children - review of 1,000 cases. *Epilepsy Res* 2005;66:129–35.

Chadwick D, Smith D. The misdiagnosis of epilepsy. *BMJ* 2002;324:495–6.

Devinsky O, Hesdorffer DC, Thurman DJ, Lhatoo S, Richerson G. Sudden unexpected death in epilepsy: epidemiology, mechanisms, and prevention. *Lancet Neurol*. 2016;15(10):1075–88. [https://doi.org/10.1016/S1474-4422\(16\)30158-2](https://doi.org/10.1016/S1474-4422(16)30158-2).

Dwivedi R, Ramanujam B, Chandra PS, Sapra S, Gulati S, Kalaivani M, Garg A, Bal CS, Tripathi M, Dwivedi SN, Sagar R, Sarkar C, Tripathi M. Surgery for drug-resistant epilepsy in children. *N Engl J Med*. 2017;377(17):1639–47. <https://doi.org/10.1056/NEJMoa1615335>.

Engel Jr J, McDermott MP, Wiebe S, Langfitt JT, Stern JM, Dewar S, Sperling MR, Gardiner I, Erba G, Fried I, Jacobs M, Vinters HV, Mintzer S, Kieburz K. Early Randomized Surgical Epilepsy Trial (ERSET) Study Group. Early surgical therapy for drug-resistant temporal lobe epilepsy: a randomized trial. *JAMA*. 2012;307(9):922–30. <https://doi.org/10.1001/jama.2012.220>.

Ferrie CD. Preventing misdiagnosis of epilepsy. *Arch Dis Child* 2006;91:206–9.

Gélisse P, Rossetti AO, Genton P, Crespel A, Kaplan PW. How to carry out and interpret EEG recordings in COVID-19 patients in ICU? *Clin Neurophysiol*. 2020;131:2023–31. <https://doi.org/10.1016/j.clinph.2020.05.006>.

Granbichler CA, Oberaigner W, Kuchukhidze G, Ndayisaba JP, Ndayisaba A, Taylor A, Bauer G, Luef G, Trinka E. Decrease in mortality of adult epilepsy patients since 1980: lessons learned from a hospital-based cohort. *Eur J Neurol*. 2017;24:667–72.

Hargreaves DS, Arora S, Viveiro C, Hale DR, Ward JL, Sherlaw-Johnson C, Viner RM, Dunkley C, Cross JH. Association of quality of paediatric epilepsy care with mortality and unplanned hospital admissions among children and young people with epilepsy in England: a national longitudinal data linkage study. *Lancet Child Adolesc Health*. 2019;3:627–35.

Joint proposals by the Japan Epilepsy Society, Japanese Society of Clinical Neurophysiology, The Japan Neurosurgical Society, Japanese Society of Neurology, Japanese Society of Child Neurology, and Japanese Society of Psychiatry and Neurology: Alert for EEG examinations to prevent COVID-19 infection. <https://www.ilae.org/files/dmfile/JES-Alert-for-EEG-examinations.pdf>. Latest accessed: February 6, 2020.

Kobulashvili T, Kuchukhidze G, Brigo F, Zimmermann G, Höfler J, Leitinger M, Dobesberger J, Kalss G, Rohrachner A, Neuray C, Wakonig A, Ernst F, Braun KPJ, Mouthaan BE, Van Eijsden P, Rylvlin P, Cross JH, Trinka E. E-PILEPSY consortium. Diagnostic and prognostic value of noninvasive long-term video-electroencephalographic monitoring in epilepsy surgery: a systematic review and meta-analysis from the E-PILEPSY consortium. *Epilepsia*. 2018;59:2272–83. <https://doi.org/10.1111/epi.14598>.

Krysl D, Beniczky S, Franceschetti S, Arzimanoglou A. The COVID-19 outbreak and approaches to performing EEG in Europe. *Epileptic Disord*. 2020;22(5):548–54. <https://doi.org/10.1684/epd.2020.1208>.

Lafrance Jr WC, Benbadis SR. Avoiding the costs of unrecognized psychological nonepileptic seizures. *Neurology* 2006;13(66):1620–1.

Lowerison MW, Josephson CB, Jetté N, Sajobi TT, Patten S, Williamson T, Deardon R, Barkema HW, Wiebe S. Association of levels of specialized care with risk of premature mortality in patients with epilepsy. *JAMA Neurol*. 2019;76(11):1352–8. <https://doi.org/10.1001/jamaneurol.2019.2268>.

McBride AE, Shih TT, Hirsch LJ. Video-EEG monitoring in the elderly: a review of 94 patients. *Epilepsia* 2002;43:165–9.

Nightscales R, McCartney L, Auvrez C, Tao G, Barnard S, Malpas CB, Perucca P, McIntosh A, Chen Z, Sivathamboo S, Ignatiadis S, Jones S, Adams S, Cook MJ, Kwan P, Velakoulis D, D'Souza W, Berkovic SF, O'Brien TJ. Mortality in patients with psychogenic nonepileptic seizures. *Neurology*. 2020;95(6):e643–52.

San-Juan D, Jiménez CR, Camilli CX, de la Cruz Reyes LA, Galindo EGA, Burbano GER, Penela MM, Perassolo MB, Valdéz AT, Godoy JG, Moreira AL, Kimaid PAT. Guidance for clinical neurophysiology examination throughout the COVID-19 pandemic. Latin American chapter of the IFCN task force - COVID-19. *Clin Neurophysiol*. 2020;131:1589–98. <https://doi.org/10.1016/j.clinph.2020.04.011>.

Tatum WO, Rubboli G, Kaplan PW, Mirsafari SM, Radhakrishnan K, Gloss D, Caboclo LO, Drislane FW, Koutroumanidis M, Schomer DL, Kasteleijn-Nolst Trenite D, Cook M, Beniczky S. Clinical utility of EEG in diagnosing and monitoring epilepsy in adults. *Clin Neurophysiol*. 2018;129(5):1056–82. <https://doi.org/10.1016/j.clinph.2018.01.019>.

Uldall P, Alving J, Hansen LK, et al. The misdiagnosis of epilepsy in children admitted to a tertiary epilepsy centre with paroxysmal events. *Arch Dis Child* 2006;91:219–21.

Wiebe S, Blume WT, Girvin JP, Eliasziw M. Effectiveness and Efficiency of Surgery for Temporal Lobe Epilepsy Study Group. A randomized, controlled trial of surgery for temporal-lobe epilepsy. *N Engl J Med*. 2001;345(5):311–8. <https://doi.org/10.1056/NEJM200108023450501>.