



$W\gamma$ production at NNLO+PS accuracy in GENEVA

Thomas Cridge^a, Matthew A. Lim^{b,*}, Riccardo Nagar^b

^a Department of Physics and Astronomy, University College London, London, WC1E 6BT, UK

^b Università degli Studi di Milano-Bicocca & INFN Sezione di Milano-Bicocca, Piazza della Scienza 3, Milano 20126, Italy

ARTICLE INFO

Article history:

Received 3 June 2021

Received in revised form 21 December 2021

Accepted 19 January 2022

Available online 21 January 2022

Editor: G.F. Giudice

ABSTRACT

We present an event generator for the process $pp \rightarrow \ell\nu_\ell\gamma$ at next-to-next-to-leading order (NNLO) in QCD and matched to the PYTHIA8 parton shower. The calculation makes use of the GENEVA framework, which combines a resummed calculation obtained via Soft-Collinear Effective Theory (SCET) with the fixed-order result. We validate the NNLO accuracy of our results by comparing predictions for inclusive quantities with an independent fixed-order calculation, and then present the resummed partonic 0-jettiness spectrum at next-to-next-to-leading logarithmic (NNLL') accuracy. Finally, we compare our predictions against data collected by the ATLAS experiment at the Large Hadron Collider during its 7 TeV run, finding good agreement.

© 2022 The Author(s). Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The process $pp \rightarrow \ell\nu_\ell\gamma$ is particularly interesting among diboson processes measurable at the Large Hadron Collider (LHC). Due to the presence of a radiation zero at leading order,¹ the next-to-leading order (NLO) corrections in the strong coupling are artificially large. This motivates the inclusion of higher order (next-to-next-to-leading, or NNLO) corrections in theoretical predictions for this process. It is also sensitive to the value of the non-Abelian electroweak gauge coupling, making it important for studies which seek to stress-test the gauge structure of the Standard Model (SM) and probe the effects of additional higher-dimensional operators in the SM effective field theory (SMEFT).

The $W\gamma$ final state was originally observed in $p\bar{p}$ collisions at the Tevatron [2–5], and has since been measured by both the ATLAS [6–8] and CMS [9–11] collaborations at centre-of-mass energies of 7 and 13 TeV. These measurements have been used to constrain anomalous $WW\gamma$ couplings and, in the case of Ref. [11], to set limits on SMEFT Wilson coefficients.

Fixed order calculations for this process at NNLO accuracy first appeared in Ref. [12]. These were made publicly available in the MATRIX program [13], which employs the q_T slicing method [14] to regularise infrared divergences. Electroweak corrections to this process at NLO are also known [15], and have been combined with

NLO [16] and NNLO [17] QCD corrections. Very recently, a new calculation of the process at NNLO QCD accuracy appeared [18], which utilised the N -jettiness slicing method [19] and analytic expressions for the multi-loop matrix elements. An NLO+PS event generator was also implemented in the POWHEG framework [20].

There has been much recent activity in the field of matching calculations at NNLO to parton showers (PS), with four main approaches extant [21–28]. Event generators at NNLO+PS accuracy for several diboson processes are already available – in the GENEVA framework, diphoton production [29], ZZ production [30], and Higgsstrahlung [31] have all been implemented, while processes such as WW production [32,33] and $Z\gamma$ [34] are also available at NNLO+PS via the MINNLO_{PS} method and the VH processes have been calculated using the MINLO' approach [35,36].²

In this work, we present an event generator at NNLO+PS accuracy for the $W\gamma$ process. The GENEVA approach, which we follow in this work, relies on a consistent matching of calculations resummed in a jet resolution variable at a high logarithmic accuracy with fixed order predictions at NNLO. The events thus produced are then fed to a parton shower program, which creates final states of high multiplicity and can be compared to experimental data. The framework is in principle fully general, in the sense that it does not depend on a particular choice of resolution variable or a particular resummation formalism, nor is it restricted in its application to a specific class of processes. For this case, we make use of the 0-jettiness resolution variable and its resummation in Soft-Collinear Effective Theory (SCET), which have been success-

* Corresponding author.

E-mail address: matthew.lim@unimib.it (M.A. Lim).

¹ The tree-level $W\gamma$ amplitude vanishes exactly at a centre-of-mass scattering angle of $\cos\theta^* = 1/3$ due to the diagram featuring the non-Abelian coupling, suppressing the leading order partonic cross section [1].

² Other processes not directly related to diboson production have also been implemented at NNLO+PS accuracy, see Refs. [37–39].

fully employed in previous GENEVA calculations. We shower the partonic events with PYTHIA8, which also simulates hadronisation.

The Letter is organised as follows. In sec. 2 we define the process and provide a brief recap of the GENEVA formulae, paying particular attention to the caveats that are needed for processes such as ours which are divergent at Born level. In sec. 3 we present a validation against the MATRIX result at NNLO and compare our predictions against ATLAS data. We conclude in sec. 4. In Appendix A, we discuss the impact of missing power corrections on our results and the reweighting procedure that we use to limit this.

2. Process definition

We apply the GENEVA formalism to the process $pp \rightarrow \ell \nu_\ell \gamma + X$. The presence of a photon in the final state requires the introduction of criteria which ensure that the photon is produced in the hard interaction, rather than being the result of fragmentation from a quark or gluon. To this end, we employ the smooth-cone isolation procedure introduced by Frixione [40] which limits the amount of transverse hadronic energy around the photon in an infrared-safe manner. Explicitly, the procedure requires that a continuous series of sub-cones with radius $r \leq R_{\text{iso}}$ satisfy the criterion $E_T^{\text{had}}(r) \leq E_T^{\text{max}} \chi(r; R_{\text{iso}})$, where a standard choice for the isolation function is

$$\chi(r; R_{\text{iso}}) = \left(\frac{1 - \cos r}{1 - \cos R_{\text{iso}}} \right)^n, \quad (1)$$

though other choices are possible. In this way, hadronic activity is smoothly reduced as one approaches the photon direction until at $r = 0$ it is completely absent.

The smooth-cone isolation procedure has several theoretical advantages such as infrared safety. It is, however, somewhat at odds with the approach taken in collider experiments, where the discretised nature of the detector means that the smooth sub-cone concept is not easily realised in practice, and it is instead common to consider a cone of fixed size R_{iso} . For this reason, Refs. [41,42] investigated the possibility of using a hybrid isolation procedure in theoretical predictions. In this approach, events are generated subject to a smooth-cone isolation with only loose cuts being placed (i.e. a small R_{iso} is used) and subsequently tighter cuts are imposed at analysis level using a cone of fixed size. In this way one combines the advantages of both approaches and makes the direct comparison to experimental data possible. We make use of this approach in our comparison to experimental data in sec. 3.

In addition to the photon isolation, it is also necessary to impose other process-defining cuts to ensure that the Born-level cross section is free of QED divergences. Specifically, one needs to separate in some way the photon from the initial-state partons and also from the charged lepton produced in the W decay. We achieve these separations by placing requirements on the transverse momentum of the photon with respect to the beam direction $p_T^\gamma \geq p_{T,\text{cut}}^\gamma$ and on the $\Delta R \equiv \sqrt{(\Delta\phi)^2 + (\Delta y)^2}$ of the lepton-photon pair $\Delta R^{\ell\gamma} \geq \Delta R_{\text{cut}}^{\ell\gamma}$.

The Monte Carlo (MC) cross sections of various multiplicity in GENEVA are defined to correspond directly to the weights of physical and IR-finite events at a given perturbative accuracy. This is achieved by partitioning the phase space into regions with different numbers of resolved emissions classified according to some N -jet resolution variables $\mathcal{T}_N$, and then mapping IR-divergent final states with $M \geq N$ partons to IR-finite final states with N partonic jets. This ensures the cancellation of IR singularities on an event-by-event basis. The implication is that the GENEVA MC cross sections $d\sigma_N^{\text{MC}}$ receive contributions from configurations with both N and M partons, where the additional emissions lie below a cut $\mathcal{T}_N^{\text{cut}}$ and are considered unresolved. By resumming the resolution

parameters at a high logarithmic accuracy, the dependence on the partition boundaries $\mathcal{T}_N^{\text{cut}}$ is then substantially mitigated.

Although the GENEVA method does not rely on any specific choice for the resolution variables $\mathcal{T}_N$ (and indeed several different choices are possible), thus far most implementations have relied on the 0- and 1-jettiness variables, defined as [43]

$$\mathcal{T}_N = \sum_k \min \left\{ \hat{q}_a \cdot p_k, \hat{q}_b \cdot p_k, \hat{q}_1 \cdot p_k, \dots, \hat{q}_N \cdot p_k \right\}, \quad (2)$$

with $N = 0$ or 1 , $q_{a,b}$ representing the beam directions, $q_{1\dots N}$ any final-state massless directions that minimise $\mathcal{T}_N$, and p_k the final-state partonic momenta. These separate the 0- and 1-jet exclusive from the 2-jet inclusive cross sections.

The GENEVA formulae, which have been presented in full in several papers [21,31], are slightly complicated by the presence of process-defining cuts at Born level. Since the additional complications which arise have been discussed in detail in Ref. [29], in this Letter we simply state the final results. The differential cross sections in the presence of process-defining cuts $\Theta^{\text{PS}}(\Phi_N)$ are given by

$$\frac{d\sigma_0^{\text{MC}}}{d\Phi_0}(\mathcal{T}_0^{\text{cut}}) = \quad (3)$$

$$\begin{aligned} & \frac{d\sigma^{\text{NNLL}'}}{d\Phi_0}(\mathcal{T}_0^{\text{cut}}) - \left[\frac{d\sigma^{\text{NNLL}'}}{d\Phi_0}(\mathcal{T}_0^{\text{cut}}) \right]_{\text{NNLO}_0} \\ & + (B_0 + V_0 + W_0)(\Phi_0) \Theta^{\text{PS}}(\Phi_0) \\ & + \int \frac{d\Phi_1}{d\Phi_0} (B_1 + V_1)(\Phi_1) \Theta^{\text{PS}}(\Phi_1) \\ & \quad \times \Theta^{\text{proj}}(\tilde{\Phi}_0) \theta(\mathcal{T}_0(\Phi_1) < \mathcal{T}_0^{\text{cut}}) \\ & + \int \frac{d\Phi_2}{d\Phi_0} B_2(\Phi_2) \Theta^{\text{PS}}(\Phi_2) \theta(\mathcal{T}_0(\Phi_2) < \mathcal{T}_0^{\text{cut}}), \\ & \frac{d\sigma_1^{\text{MC}}}{d\Phi_1}(\mathcal{T}_0 \leq \mathcal{T}_0^{\text{cut}}; \mathcal{T}_1^{\text{cut}}) = \quad (4) \end{aligned}$$

$$\begin{aligned} & B_1(\Phi_1) \Theta^{\text{PS}}(\Phi_1) \left[\bar{\Theta}_{\text{iso}}^{\text{proj}}(\tilde{\Phi}_0) + \bar{\Theta}_{\text{map}}^{\text{FKS}}(\Phi_1) \right] \theta(\mathcal{T}_0 < \mathcal{T}_0^{\text{cut}}), \\ & \frac{d\sigma_1^{\text{MC}}}{d\Phi_1}(\mathcal{T}_0 > \mathcal{T}_0^{\text{cut}}; \mathcal{T}_1^{\text{cut}}) = \quad (5) \end{aligned}$$

$$\begin{aligned} & \Theta^{\text{PS}}(\Phi_1) \left\{ \left[\frac{d\sigma^{\text{NNLL}'}}{d\Phi_0 d\mathcal{T}_0} - \frac{d\sigma^{\text{NNLL}'}}{d\Phi_0 d\mathcal{T}_0} \right]_{\text{NLO}_1} \right\} \mathcal{P}(\Phi_1) \Theta^{\text{proj}}(\tilde{\Phi}_0) \\ & + [B_1 + V_1^C](\Phi_1) \left\{ U_1(\Phi_1, \mathcal{T}_1^{\text{cut}}) \theta(\mathcal{T}_0 > \mathcal{T}_0^{\text{cut}}) \right. \\ & + \int \left[\frac{d\Phi_2}{d\Phi_1} B_2(\Phi_2) \Theta^{\text{PS}}(\Phi_2) \Theta^{\text{proj}}(\tilde{\Phi}_1) \theta(\mathcal{T}_0(\Phi_2) > \mathcal{T}_0^{\text{cut}}) \right. \\ & \quad \times \theta(\mathcal{T}_1 < \mathcal{T}_1^{\text{cut}}) - \frac{d\Phi_2}{d\Phi_1^C} C_2(\Phi_2) \Theta^{\text{PS}}(\Phi_1) \theta(\mathcal{T}_0 > \mathcal{T}_0^{\text{cut}}) \left. \right] \\ & \left. - B_1(\Phi_1) U_1^{(1)}(\Phi_1, \mathcal{T}_1^{\text{cut}}) \Theta^{\text{PS}}(\Phi_1) \theta(\mathcal{T}_0 > \mathcal{T}_0^{\text{cut}}), \right. \\ & \frac{d\sigma_{\geq 2}^{\text{MC}}}{d\Phi_2}(\mathcal{T}_0 > \mathcal{T}_0^{\text{cut}}, \mathcal{T}_1 \leq \mathcal{T}_1^{\text{cut}}) = \quad (6) \end{aligned}$$

$$\begin{aligned} & B_2(\Phi_2) \Theta^{\text{PS}}(\Phi_2) \left[\bar{\Theta}^{\text{proj}}(\tilde{\Phi}_1) + \bar{\Theta}_{\text{map}}^{\mathcal{T}}(\Phi_2) \right] \\ & \quad \times \theta(\mathcal{T}_1 < \mathcal{T}_1^{\text{cut}}) \theta(\mathcal{T}_0(\Phi_2) > \mathcal{T}_0^{\text{cut}}), \\ & \frac{d\sigma_{\geq 2}^{\text{MC}}}{d\Phi_2}(\mathcal{T}_0 > \mathcal{T}_0^{\text{cut}}, \mathcal{T}_1 > \mathcal{T}_1^{\text{cut}}) = \quad (7) \\ & \Theta^{\text{PS}}(\Phi_2) \left\{ \left[\frac{d\sigma^{\text{NNLL}'}}{d\Phi_0 d\mathcal{T}_0} - \frac{d\sigma^{\text{NNLL}'}}{d\Phi_0 d\mathcal{T}_0} \right]_{\text{NLO}_1} \right\} \mathcal{P}(\tilde{\Phi}_1) \Theta^{\text{proj}}(\tilde{\Phi}_0) \end{aligned}$$

$$\begin{aligned}
& + (B_1 + V_1^C)(\tilde{\Phi}_1) \left\{ U_1'(\tilde{\Phi}_1, \mathcal{T}_1) \theta(\mathcal{T}_0 > \mathcal{T}_0^{\text{cut}}) \right\} \Big|_{\tilde{\Phi}_1 = \Phi_1^{\mathcal{T}}(\Phi_2)} \\
& \times \Theta^{\text{proj}}(\tilde{\Phi}_1) \mathcal{P}(\Phi_2) \theta(\mathcal{T}_1 > \mathcal{T}_1^{\text{cut}}) \\
& + \Theta^{\text{PS}}(\Phi_2) \left\{ B_2(\Phi_2) \theta(\mathcal{T}_1 > \mathcal{T}_1^{\text{cut}}) - B_1(\Phi_1^{\mathcal{T}}) U_1^{(1)'}(\tilde{\Phi}_1, \mathcal{T}_1) \right. \\
& \times \mathcal{P}(\Phi_2) \Theta^{\text{proj}}(\tilde{\Phi}_1) \Theta(\mathcal{T}_1 > \mathcal{T}_1^{\text{cut}}) \left. \right\} \theta(\mathcal{T}_0(\Phi_2) > \mathcal{T}_0^{\text{cut}}).
\end{aligned}$$

The B_j , V_j and W_j are the 0-, 1- and 2-loop matrix elements for j QCD partons in the final state; we denote by $N^k\text{LO}_n$ a quantity with n additional partons in the final state computed at $N^k\text{LO}$ accuracy.

The formulae above require one to evaluate the resummed and resummed-expanded terms on phase space points resulting from a projection from a higher to a lower multiplicity. We denote such projected phase space points by $\tilde{\Phi}_N$. It is vital that these projected configurations also satisfy the process-defining cuts. We denote by $\Theta^{\text{proj}}(\tilde{\Phi}_N)$ the set of restrictions acting on the higher dimensional Φ_{N+1} phase space due to the cuts on $\tilde{\Phi}_N$. In practice, this means that when a term in the cross section, evaluated at a Φ_{N+1} phase space point, is multiplied by $\Theta^{\text{proj}}(\tilde{\Phi}_N)$, we perform the projection $\Phi_{N+1} \rightarrow \tilde{\Phi}_N$ and apply the cuts to $\tilde{\Phi}_N$. If the projected point fails the cuts, the initial Φ_{N+1} configuration is excluded and is instead assigned to either of the nonsingular bins in eqs. (4) and (6).

We use the shorthand

$$\frac{d\Phi_M}{d\Phi_N^{\mathcal{O}}} = d\Phi_M \delta[\Phi_N - \Phi_N^{\mathcal{O}}(\Phi_M)] \Theta^{\mathcal{O}}(\Phi_M) \quad (8)$$

to indicate an integration over the portion of the Φ_M phase space which can be reached from a fixed value of Φ_N while keeping some observable $\mathcal{O}$ also fixed, with $N \leq M$. The $\Theta^{\mathcal{O}}(\Phi_M)$ term additionally limits the integration to the phase space points belonging to the singular contribution for the given observable $\mathcal{O}$. For example, when generating 1-body events we use

$$\frac{d\Phi_2}{d\Phi_1^{\mathcal{T}}} \equiv d\Phi_2 \delta[\tilde{\Phi}_1 - \Phi_1^{\mathcal{T}}(\Phi_2)] \Theta^{\mathcal{T}}(\Phi_2), \quad (9)$$

where the map used by the $1 \rightarrow 2$ splitting has been constructed to preserve $\mathcal{T}_0$, i.e.

$$\mathcal{T}_0(\Phi_1^{\mathcal{T}}(\Phi_2)) = \mathcal{T}_0(\Phi_2), \quad (10)$$

and $\Theta^{\mathcal{T}}(\Phi_2)$ guarantees that the Φ_2 point is reached from a genuine QCD splitting of the Φ_1 point. The use of a $\mathcal{T}_0$ -preserving mapping is necessary to ensure that the point-wise singular $\mathcal{T}_0$ dependence is alike among all terms in eqs. (5) and (7) and that the cancellation of said singular terms is guaranteed on an event-by-event basis.³

The non-projectable regions of Φ_1 and Φ_2 , on the other hand, belong to the cross sections in eqs. (4) and (6). These events are entirely nonsingular in nature. We denote the constraints due to the choice of map by Θ_{map} ⁴ and use Θ^{proj} to indicate the restrictions due to both the isolation cuts and the flavour structure of the underlying Born configuration (see e.g. Ref. [31] for more details).

The term V_1^C denotes the soft-virtual contribution of a standard NLO local subtraction:

$$V_1^C(\Phi_1) = V_1(\Phi_1) + \int \frac{d\Phi_2}{d\Phi_1^C} C_2(\Phi_2), \quad (11)$$

³ We remark that this differentiates our approach from the slicing method used in Ref. [18] – the use of a $\mathcal{T}_0$ -preserving mapping ensures that our subtraction, though not fully local, results in point-wise cancellation of divergences in this singular variable.

⁴ We use the FKS map [44] for the $\Phi_1 \rightarrow \tilde{\Phi}_0$ projection and, as mentioned above, our own $\mathcal{T}_0$ -preserving map for the $\Phi_2 \rightarrow \tilde{\Phi}_1$ projection.

with C_2 a singular approximation of B_2 ; in practice we use the subtraction counterterms which we integrate over the radiation variables $d\Phi_2/d\Phi_1^C$ using the singular limit C of the phase space mapping.

U_1 is an NLL Sudakov factor which resums large logarithms of $\mathcal{T}_1$, and U_1' its derivative with respect to $\mathcal{T}_1$; the $\mathcal{O}(\alpha_s)$ expansions of these quantities are denoted by $U_1^{(1)}$ and $U_1^{(1)'}$ respectively.

We extend the differential dependence of the resummed terms from the N -jet to the $(N+1)$ -jet phase space using a normalised splitting probability $\mathcal{P}(\Phi_{N+1})$ which satisfies

$$\int \frac{d\Phi_{N+1}}{d\Phi_N d\mathcal{T}_N} \mathcal{P}(\Phi_{N+1}) = 1. \quad (12)$$

The two extra variables are chosen to be an energy ratio z and an azimuthal angle ϕ . The functional forms of the $\mathcal{P}(\Phi_{N+1})$ are based on the Altarelli-Parisi splitting kernels, weighted by parton distribution functions (PDFs) where appropriate.

In principle, the GENEVA method is compatible with any formalism which can provide the resummed spectrum at NNLL' accuracy. In practice, we obtain this from a factorisation theorem derived in SCET and in the case of colour singlet production write

$$\begin{aligned}
\frac{d\sigma^{\text{SCET}}}{d\Phi_0 d\mathcal{T}_0} &= \sum_{ij} H_{ij}(\Phi_0, \mu) \int dt_a dt_b B_i(t_a, x_a, \mu) \\
& B_j(t_b, x_b, \mu) S(\mathcal{T}_0 - \frac{t_a+t_b}{Q}, \mu), \quad (13)
\end{aligned}$$

where the sum runs over all possible $q\bar{q}$ flavours. The cross section has been factorised into hard H_{ij} , soft S and beam $B_{i,j}$ functions which achieves a separation of scales: a judicious scale choice for each component will therefore result in the absence of large logarithms. Resummation is then achieved via renormalisation group evolution to a common scale μ , viz.

$$\begin{aligned}
\frac{d\sigma^{\text{NNLL}'}}{d\Phi_0 d\mathcal{T}_0} &= \sum_{ij} H_{ij}(\Phi_0, \mu_H) U_H(\mu_H, \mu) \\
& \{ [B_i(t_a, x_a, \mu_B) \otimes U_B(\mu_B, \mu)] \\
& \times [B_j(t_b, x_b, \mu_B) \otimes U_B(\mu_B, \mu)] \} \\
& \otimes [S(\mu_S) \otimes U_S(\mu_S, \mu)], \quad (14)
\end{aligned}$$

where the U_i denote the evolution kernels and the convolutions are written in a schematic form. The cusp (non-cusp) anomalous dimensions up to 3-(2-)loop order needed for the evolution at NNLL' have been available for some time [45–48,19]. Other inputs to the factorisation theorem also appear in the literature – the beam and soft functions have been computed in Refs. [49–51], while the helicity amplitudes necessary for the extraction of the hard function were first computed in Ref. [52].

The events thus produced are then fed to the PYTHIA8 parton shower [53] which adds radiation to produce final states of a high multiplicity. This restores the radiation that was integrated over in the construction of the 0- and 1-jet cross sections, and adds additional particles to the inclusive 2-jet bin. The consequence is that, while quantities which are inclusive over the radiation are expected to remain unmodified by the action of the shower, the distributions of more exclusive observables will be modified. We thus retain NNLO accuracy for inclusive quantities.

For further details on the topics discussed above, we refer the reader to Refs. [21,29].

3. Results and comparison to LHC data

We begin by summarising the process-specific features of our implementation. At Born level, the process receives contributions




Fig. 1. Comparison between the MATRIX and GENEVA predictions at NNLO accuracy. Top row: the rapidity of the electron (left), the transverse momentum of the electron (centre) and the rapidity of the photon (right). Bottom row: the invariant mass (left) and the rapidity (right) of the colour singlet system. Scale uncertainty bands include 3-point renormalisation and factorisation scale variations. Statistical errors connected to the Monte Carlo integration are shown as vertical error bars.

from diagrams with three different resonance structures. Two of these are akin to the Drell-Yan process, but with the photon emitted either from the vector boson line or from the charged lepton, while the third is a t -channel process in which the photon is emitted from the quark line. In order to efficiently sample the phase spaces associated with these different resonance structures, we make use of the tunnel between GENEVA and MUNICH [54], which was first constructed in Ref. [30]. We rely on OPENLOOPS2 [55,56] for the calculation of tree level and 1-loop matrix elements. We make use of the 2-loop hard function implemented for this process in MATRIX [52,12] and use a diagonal CKM matrix, though this could in principle be trivially extended.⁵

In Fig. 1 we validate the NNLO accuracy of our predictions by comparison with MATRIX. We consider the electronic decay channel of the W^- boson at a pp centre-of-mass energy of $\sqrt{S} = 7$ TeV. We require a separation $\Delta R_{e\gamma} > 0.7$ between the electron and the photon, and place a transverse momentum cut on the photon $p_{T,\text{cut}}^\gamma = 15$ GeV. We employ a smooth-cone isolation procedure with parameters $E_T^{\text{max}} = 4$ GeV, $R = 0.4$, $n = 1$ and use the MSHT20_as118 sets [58] from LHAPDF6 [59]. We set $\alpha_s(M_Z) = 0.118$ and $\alpha(M_Z) = 7.55638 \times 10^{-3}$. We take $M_W = 80.385$ GeV and $\Gamma_W = 2.085$ GeV – for the purposes of this comparison alone, we do not perform any resummation of the logarithms of T_1 . We show here only plots involving the negatively charged W , but those for the W^+ channel are similar.




Fig. 2. The partonic T_0 spectrum at NNLL'+NNLO₀ accuracy (blue) compared to results after showering (orange) and hadronisation (green). The sum over both charges of the intermediate W boson is shown. A semi-logarithmic scale is used for T_0 , which is linear up to 20 GeV and logarithmic beyond.

⁵ When performing our validation, we noticed discrepancies between the hard function at 1-loop order as implemented in the public version 1.0.5 of MATRIX and the results of OPENLOOPS2, due to the use of an outdated version of the `hplog` package [57] in the former. We have corrected this in our implementation.




Fig. 3. Comparison between GENEVA predictions and ATLAS measurements at 7 TeV (4.6 fb^{-1}). Selection cuts, bin widths and observable definitions are defined in the original ATLAS [8] publication.

In general, we observe a good agreement between the central values from MATRIX and GENEVA for all rapidity distributions. In the transverse momentum distribution, we notice deviations from the fixed order prediction similar to those observed in Ref. [29]. These are related to ‘leptonic’ fiducial power corrections [60] which arise when measuring properties of the electroweak final state particles. The fact that we evaluate our resummed $\mathcal{T}_0$ spectrum and splitting function on a Φ_1 point means that the hadronic recoil and thereby the fiducial power corrections are correctly accounted for in the majority of the cross section where $\mathcal{T}_0 > \mathcal{T}_0^{\text{cut}}$, and hence GENEVA treats these fiducial power corrections in a physical manner.⁶ Nevertheless, the two predictions agree within the scale variation bands. We also observe a good agreement within scale uncertainties for the whole range of $M_{ev\gamma}$, with GENEVA predictions slightly higher above the peak. Examining the bands, we observe a difference in size between MATRIX and GENEVA which is particularly noticeable in the rapidity distribution of the electron. We remark that the reweighting procedure described in Appendix A is constructed to recover the value of the fixed order total cross section and its variations correctly, while the kinematic dependence of power corrections is still omitted. The effect on the scale bands is particularly noticeable in this case due to the accidentally small size of the scale uncertainties, which are a result of adopting a correlated variation with $\mu_R = \mu_F$ (see Ref. [12]).

The NNLL+NNLO accurate $\mathcal{T}_0$ spectrum at partonic level is shown in Fig. 2, where we also show the effects of showering and hadronisation with PYTHIA8. In this plot and those following, we have adopted the scale choice $\mu = M_{ev\gamma}^T$ where

$$(M_{ev\gamma}^T)^2 \equiv \left(\sqrt{M_{e\gamma}^2 + p_{T,e\gamma}^2} + p_{T,e\gamma}^{\text{miss}} \right)^2 - p_{T,e\gamma}^{\text{miss}2} \quad (15)$$

where $p_{T,e\gamma}^{\text{miss}}$ is the transverse momentum of the electron, photon and $p_{T,e\gamma}^{\text{miss}}$ system.

In order to simplify the analysis, we have deactivated both the QED part of the shower and also the simulation of multiparton interactions – in the case of the former, we would not expect this to have a major impact on the $\mathcal{T}_0$ spectrum. In the case of the latter, this can result in significant shifts in the $\mathcal{T}_0$ spectrum (see e.g. Ref. [31]) due to the definition of the observable, but should have a minimal effect on inclusive distributions. Comparing the partonic with the showered result, we note that the distribution is changed very little by the shower in the peak region (up to $\sim 10 \text{ GeV}$) where resummation effects are most important. The effect of hadronisation is to shift the peak of the distribution to higher values of $\mathcal{T}_0$ – its impact is lessened in the tail of the distribution, where nonperturbative effects become increasingly less relevant.

Finally, in Fig. 3 we compare the GENEVA predictions to data collected by the ATLAS experiment at the LHC in $\sqrt{S} = 7 \text{ TeV}$ collisions, with a total integrated luminosity of 4.6 fb^{-1} [8]. We generate events with a set of isolation cuts which are looser than those used in the experimental analysis, shower and hadronise, and then pass the resulting events through RIVET [61] – this amounts to using a hybrid-type photon isolation. The sum over events with both signs of the intermediate W boson is considered. We show the distribution of the transverse momentum of the photon p_γ^T , and the normalised distribution of the transverse mass of the colour singlet system $M_{ev\gamma}^T$, the latter with a harder cut on the photon of $p_\gamma^T > 40 \text{ GeV}$. After showering, we expect our predictions for these distributions to have an accuracy no worse than that provided by the standalone PYTHIA8. The agreement with data is reasonably good and of a similar quality to the NNLO results provided by MATRIX in Ref. [12]. In particular, for the case of the $M_{ev\gamma}^T$ distribution we notice that the predictions tend to undershoot the data at high values. We anticipate that the inclusion of electroweak corrections would improve the agreement in this region, where the contribution due to photon-initiated processes is known to be large (though partially mitigated by correspondingly large and negative corrections to the $q\bar{q}$ channel) [16].

⁶ Methods for incorporating the correct treatment of fiducial power corrections in a slicing calculation at fixed order are also discussed in Ref. [60].

4. Conclusions

In this work we have presented the first event generator at NNLO accuracy for the diboson production process $pp \rightarrow \ell\nu\ell\gamma$ matched to a parton shower. This has been implemented in the GENEVA framework, which matches a resummed calculation at NNLL' in the resolution variable $\mathcal{T}_0$ to fixed order predictions at NNLO and allows the resulting events to be passed to PYTHIA8. We validated the NNLO accuracy of our predictions for inclusive observables by comparing with the fixed order code MATRIX, finding that the kinematic effects of missing inclusive nonsingular power corrections on differential distributions are mild at worst. We then examined the effect of the shower on the $\mathcal{T}_0$ distribution, finding that the NNLL' accuracy was numerically well-preserved in the peak region, and observed that the effect of hadronisation is limited to a shift at low values where nonperturbative effects are particularly relevant. The final logarithmic accuracy of our results is at least as good as the standalone PYTHIA8 parton shower.

Lastly, we compared to data collected at the 7 TeV LHC by the ATLAS experiment in 2011. We found good agreement, consistent with previous studies at fixed order in QCD, with small deviations seen in regions where EW effects are likely to be important.

It would be important to consider the inclusion of these electroweak corrections to this process, especially given their numerical relevance at the 13 TeV LHC [18]. A study of the differences with respect to $Z\gamma$ production and the effect of the inclusion of a non-diagonal CKM matrix may also prove interesting – we leave these issues to a future work.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We thank our collaborators Simone Alioli, Alessandro Broggio, Alessandro Gavardi, Stefan Kallweit, and Davide Napoletano for useful discussions. We thank Stefan Kallweit for help with MATRIX and for providing the necessary MUNICH libraries. We are grateful to Johannes Michel for conversations regarding the impact of leptonic fiducial power corrections. The work of MAL and RN is supported by the ERC Starting Grant REINVENT-714788. TC is supported by the Science and Technology Facilities Council (STFC) via grant awards ST/P000274/1 and ST/T000856/1. We acknowledge the CINECA award under the ISCRA initiative for the availability of the high performance computing resources needed for this work.

Appendix A. Nonsingular power corrections

The contribution to the 0-jet bin below the resolution cut-off $\mathcal{T}_0^{\text{cut}}$ (eq. (3)) would require a local NNLO subtraction in order to be properly implemented. Instead, we substitute the expression with

$$\begin{aligned} \frac{\widehat{d\sigma}_0^{\text{MC}}}{d\Phi_0}(\mathcal{T}_0^{\text{cut}}) &= \frac{d\sigma^{\text{NNLL}'}}{d\Phi_0}(\mathcal{T}_0^{\text{cut}}) - \left[\frac{d\sigma^{\text{NNLL}'}}{d\Phi_0}(\mathcal{T}_0^{\text{cut}}) \right]_{\text{NLO}_0} \\ &+ (B_0 + V_0)(\Phi_0) \Theta^{\text{PS}}(\Phi_0) \\ &+ \int \frac{d\Phi_1}{d\Phi_0} B_1(\Phi_1) \Theta^{\text{PS}}(\Phi_1) \\ &\times \Theta^{\text{proj}}(\tilde{\Phi}_0) \theta(\mathcal{T}_0(\Phi_1) < \mathcal{T}_0^{\text{cut}}), \end{aligned} \quad (\text{A.1})$$

and so neglect the contribution




Fig. A.4. Size of the neglected power corrections $\Sigma_{\text{ns}}^{(2)}$ as a function of the resolution parameter $\mathcal{T}_0^{\text{cut}}$. The target result σ_{NNLO} is the fixed-order cross section as computed by MATRIX. The error band shows the size of the numerical fluctuations. On the right axis we also show the relative size of the corrections as a fraction of the MATRIX NNLO cross section.

$$\frac{d\Sigma_{\text{ns}}^{(2)}}{d\Phi_0}(\mathcal{T}_0^{\text{cut}}) = \frac{d\sigma_0^{\text{MC}}}{d\Phi_0}(\mathcal{T}_0^{\text{cut}}) - \frac{\widehat{d\sigma}_0^{\text{MC}}}{d\Phi_0}(\mathcal{T}_0^{\text{cut}}). \quad (\text{A.2})$$

These neglected terms are power corrections of $\mathcal{O}(\alpha_s^2)$. They therefore contribute at the same order as the power corrections which arise from evaluating higher multiplicity configurations on projected phase space points, which is an issue common to any method of event generation. The integral of all these terms is shown in Fig. A.4 as a function of $\mathcal{T}_0^{\text{cut}}$. We observe that the behaviour is indeed nonsingular so that the size of the corrections is reduced at smaller values of $\mathcal{T}_0^{\text{cut}}$. For the purposes of this Letter we set

$$\mathcal{T}_0^{\text{cut}} = 0.5 \text{ GeV}. \quad (\text{A.3})$$

This choice is the smallest value of $\mathcal{T}_0^{\text{cut}}$ for which we could maintain a positive σ_0^{MC} . Despite the larger size of power corrections at this value, we observe that the effect on the total cross section is still $< 2\%$. This receives contributions both from ‘inclusive’ power corrections, which would be present in GENEVA even for processes without fiducial cuts (such as inclusive Drell-Yan, for example) and also fiducial power corrections, which arise when e.g. isolation criteria are introduced. We are able to recover the effect of these missing contributions on the total cross section and its variations by reweighting the events below $\mathcal{T}_0^{\text{cut}}$. As with any event generator, however, we miss the $\mathcal{O}(\alpha_s^2)$ nonsingular kinematic dependence of the inclusive power corrections for this class of events. The comparisons in Fig. 1 attest to the fact that this omission does not significantly affect our predictions for differential distributions.

References

- [1] K.O. Mikaelian, M.A. Samuel, D. Sahdev, The magnetic moment of weak bosons produced in pp and p anti- p collisions, Phys. Rev. Lett. 43 (1979) 746, <https://doi.org/10.1103/PhysRevLett.43.746>.
- [2] S. Abachi, et al., Measurement of the $WW\gamma$ gauge boson couplings in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV, Phys. Rev. Lett. 75 (1995) 1034–1039, <https://doi.org/10.1103/PhysRevLett.75.1034>, arXiv:hep-ex/9505007.
- [3] D. Acosta, et al., Measurement of $W\gamma$ and $Z\gamma$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, Phys. Rev. Lett. 94 (2005) 041803, <https://doi.org/10.1103/PhysRevLett.94.041803>, arXiv:hep-ex/0410008.
- [4] V.M. Abazov, et al., Measurement of the $p - \bar{p} \rightarrow W\gamma + X$ cross section at $\sqrt{s} = 1.96$ -TeV and $WW\gamma$ anomalous coupling limits, Phys. Rev. D 71 (2005) 091108, <https://doi.org/10.1103/PhysRevD.71.091108>, arXiv:hep-ex/0503048.

- [5] V.M. Abazov, et al., $W\gamma$ production and limits on anomalous $WW\gamma$ couplings in $p\bar{p}$ collisions, Phys. Rev. Lett. 107 (2011) 241803, <https://doi.org/10.1103/PhysRevLett.107.241803>, arXiv:1109.4432.
- [6] G. Aad, et al., Measurement of $W\gamma$ and $Z\gamma$ production in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector, J. High Energy Phys. 09 (2011) 072, [https://doi.org/10.1007/JHEP09\(2011\)072](https://doi.org/10.1007/JHEP09(2011)072), arXiv:1106.1592.
- [7] G. Aad, et al., Measurement of $W\gamma$ and $Z\gamma$ production cross sections in pp collisions at $\sqrt{s} = 7$ TeV and limits on anomalous triple gauge couplings with the ATLAS detector, Phys. Lett. B 717 (2012) 49–69, <https://doi.org/10.1016/j.physletb.2012.09.017>, arXiv:1205.2531.
- [8] G. Aad, et al., Measurements of $W\gamma$ and $Z\gamma$ production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC, Phys. Rev. D 87 (11) (2013) 112003, <https://doi.org/10.1103/PhysRevD.87.112003>, Erratum: Phys. Rev. D 91 (2015) 119901, arXiv:1302.1283.
- [9] S. Chatrchyan, et al., Measurement of $W\gamma$ and $Z\gamma$ production in pp collisions at $\sqrt{s} = 7$ TeV, Phys. Lett. B 701 (2011) 535–555, <https://doi.org/10.1016/j.physletb.2011.06.034>, arXiv:1105.2758.
- [10] S. Chatrchyan, et al., Measurement of the $W\gamma$ and $Z\gamma$ inclusive cross sections in pp collisions at $\sqrt{s} = 7$ TeV and limits on anomalous triple gauge boson couplings, Phys. Rev. D 89 (9) (2014) 092005, <https://doi.org/10.1103/PhysRevD.89.092005>, arXiv:1308.6832.
- [11] A.M. Sirunyan, et al., Measurement of $W\gamma$ production cross section in proton-proton collisions at $\sqrt{s} = 13$ TeV and constraints on effective field theory coefficients, arXiv:2102.02283.
- [12] M. Grazzini, S. Kallweit, D. Rathlev, $W\gamma$ and $Z\gamma$ production at the LHC in NNLO QCD, J. High Energy Phys. 07 (2015) 085, [https://doi.org/10.1007/JHEP07\(2015\)085](https://doi.org/10.1007/JHEP07(2015)085), arXiv:1504.01330.
- [13] M. Grazzini, S. Kallweit, M. Wiesemann, Fully differential NNLO computations with MATRIX, Eur. Phys. J. C 78 (7) (2018) 537, <https://doi.org/10.1140/epjc/s10052-018-5771-7>, arXiv:1711.06631.
- [14] S. Catani, M. Grazzini, An NNLO subtraction formalism in hadron collisions and its application to Higgs boson production at the LHC, Phys. Rev. Lett. 98 (2007) 222002, <https://doi.org/10.1103/PhysRevLett.98.222002>, arXiv:hep-ph/0703012.
- [15] E. Accomando, A. Denner, S. Pozzorini, Electroweak correction effects in gauge boson pair production at the CERN LHC, Phys. Rev. D 65 (2002) 073003, <https://doi.org/10.1103/PhysRevD.65.073003>, arXiv:hep-ph/0110114.
- [16] A. Denner, S. Dittmaier, M. Hecht, C. Pasold, NLO QCD and electroweak corrections to $W+\gamma$ production with leptonic W -boson decays, J. High Energy Phys. 04 (2015) 018, [https://doi.org/10.1007/JHEP04\(2015\)018](https://doi.org/10.1007/JHEP04(2015)018), arXiv:1412.7421.
- [17] M. Grazzini, S. Kallweit, J.M. Lindert, S. Pozzorini, M. Wiesemann, NNLO QCD + NLO EW with Matrix+OpenLoops: precise predictions for vector-boson pair production, J. High Energy Phys. 02 (2020) 087, [https://doi.org/10.1007/JHEP02\(2020\)087](https://doi.org/10.1007/JHEP02(2020)087), arXiv:1912.00068.
- [18] J.M. Campbell, G. De Laurentis, R.K. Ellis, S. Seth, The $pp \rightarrow W(\rightarrow l\nu) + \gamma$ process at next-to-next-to-leading order, arXiv:2105.00954.
- [19] J. Gaunt, M. Stahlhofen, F.J. Tackmann, J.R. Walsh, N-jettiness subtractions for NNLO QCD calculations, J. High Energy Phys. 09 (2015) 058, [https://doi.org/10.1007/JHEP09\(2015\)058](https://doi.org/10.1007/JHEP09(2015)058), arXiv:1505.04794.
- [20] L. Barze, M. Chiesa, G. Montagna, P. Nason, O. Nicrosini, F. Piccinini, V. Prospero, $W\gamma$ production in hadronic collisions using the POWHEG+MiNLO method, J. High Energy Phys. 12 (2014) 039, [https://doi.org/10.1007/JHEP12\(2014\)039](https://doi.org/10.1007/JHEP12(2014)039), arXiv:1408.5766.
- [21] S. Alioli, C.W. Bauer, C. Berggren, F.J. Tackmann, J.R. Walsh, Drell-Yan production at NNLL+NNLO matched to parton showers, Phys. Rev. D 92 (9) (2015) 094020, <https://doi.org/10.1103/PhysRevD.92.094020>, arXiv:1508.01475.
- [22] S. Alioli, C.W. Bauer, A. Broggio, A. Gavardi, S. Kallweit, M.A. Lim, R. Nagar, D. Napoletano, L. Rottoli, Matching NNLO to parton shower using N^3 LL colour-singlet transverse momentum resummation in GENEVA, arXiv:2102.08390.
- [23] K. Hamilton, P. Nason, E. Re, G. Zanderighi, NNLOPS simulation of Higgs boson production, J. High Energy Phys. 1310 (2013) 222, [https://doi.org/10.1007/JHEP10\(2013\)222](https://doi.org/10.1007/JHEP10(2013)222), arXiv:1309.0017.
- [24] P.F. Monni, P. Nason, E. Re, M. Wiesemann, G. Zanderighi, MiNNLO_{PS}: a new method to match NNLO QCD to parton showers, J. High Energy Phys. 05 (2020) 143, [https://doi.org/10.1007/JHEP05\(2020\)143](https://doi.org/10.1007/JHEP05(2020)143), arXiv:1908.06987.
- [25] P.F. Monni, E. Re, M. Wiesemann, MiNNLO_{PS}: optimizing $2 \rightarrow 1$ hadronic processes, Eur. Phys. J. C 80 (11) (2020) 1075, <https://doi.org/10.1140/epjc/s10052-020-08658-5>, arXiv:2006.04133.
- [26] S. Höche, Y. Li, S. Prestel, Drell-Yan lepton pair production at NNLO QCD with parton showers, Phys. Rev. D 91 (7) (2015) 074015, <https://doi.org/10.1103/PhysRevD.91.074015>, arXiv:1405.3607.
- [27] S. Höche, Y. Li, S. Prestel, Higgs-boson production through gluon fusion at NNLO QCD with parton showers, Phys. Rev. D 90 (5) (2014) 054011, <https://doi.org/10.1103/PhysRevD.90.054011>, arXiv:1407.3773.
- [28] Y. Hu, C. Sun, X.-M. Shen, J. Gao, Hadronic decays of Higgs, arXiv:2101.08916.
- [29] S. Alioli, A. Broggio, A. Gavardi, S. Kallweit, M.A. Lim, R. Nagar, D. Napoletano, L. Rottoli, Precise predictions for photon pair production matched to parton showers in GENEVA, J. High Energy Phys. 04 (2021) 041, [https://doi.org/10.1007/JHEP04\(2021\)041](https://doi.org/10.1007/JHEP04(2021)041), arXiv:2010.10498.
- [30] S. Alioli, A. Broggio, A. Gavardi, S. Kallweit, M.A. Lim, R. Nagar, D. Napoletano, Next-to-next-to-leading order event generation for Z boson pair production matched to parton shower, arXiv:2103.01214.
- [31] S. Alioli, A. Broggio, S. Kallweit, M.A. Lim, L. Rottoli, Higgsstrahlung at NNLL+NNLO matched to parton showers in GENEVA, Phys. Rev. D 100 (9) (2019) 096016, <https://doi.org/10.1103/PhysRevD.100.096016>, arXiv:1909.02026.
- [32] E. Re, M. Wiesemann, G. Zanderighi, NNLOPS accurate predictions for W^+W^- production, J. High Energy Phys. 12 (2018) 121, [https://doi.org/10.1007/JHEP12\(2018\)121](https://doi.org/10.1007/JHEP12(2018)121), arXiv:1805.09857.
- [33] D. Lombardi, M. Wiesemann, G. Zanderighi, W^+W^- production at NNLO+PS with MiNNLO_{PS}, arXiv:2103.12077.
- [34] D. Lombardi, M. Wiesemann, G. Zanderighi, Advancing MiNNLO_{PS} to diboson processes: $Z\gamma$ production at NNLO+PS, arXiv:2010.10478.
- [35] W. Astill, W. Bizon, E. Re, G. Zanderighi, NNLOPS accurate associated HW production, J. High Energy Phys. 06 (2016) 154, [https://doi.org/10.1007/JHEP06\(2016\)154](https://doi.org/10.1007/JHEP06(2016)154), arXiv:1603.01620.
- [36] W. Astill, W. Bizon, E. Re, G. Zanderighi, NNLOPS accurate associated HZ production with $H \rightarrow b\bar{b}$ decay at NLO, J. High Energy Phys. 11 (2018) 157, [https://doi.org/10.1007/JHEP11\(2018\)157](https://doi.org/10.1007/JHEP11(2018)157), arXiv:1804.08141.
- [37] W. Bizon, E. Re, G. Zanderighi, NNLOPS description of the $H \rightarrow b\bar{b}$ decay with MiNLO, J. High Energy Phys. 06 (2020) 006, [https://doi.org/10.1007/JHEP06\(2020\)006](https://doi.org/10.1007/JHEP06(2020)006), arXiv:1912.09982.
- [38] S. Alioli, A. Broggio, A. Gavardi, S. Kallweit, M.A. Lim, R. Nagar, D. Napoletano, L. Rottoli, Resummed predictions for hadronic Higgs boson decays, J. High Energy Phys. 04 (2021) 254, [https://doi.org/10.1007/JHEP04\(2021\)254](https://doi.org/10.1007/JHEP04(2021)254), arXiv:2009.13533.
- [39] J. Mazzeitelli, P.F. Monni, P. Nason, E. Re, M. Wiesemann, G. Zanderighi, Next-to-next-to-leading order event generation for top-quark pair production, Phys. Rev. Lett. 127 (6) (2021) 062001, <https://doi.org/10.1103/PhysRevLett.127.062001>, arXiv:2012.14267.
- [40] S. Frixione, Isolated photons in perturbative QCD, Phys. Lett. B 429 (1998) 369–374, [https://doi.org/10.1016/S0370-2693\(98\)00454-7](https://doi.org/10.1016/S0370-2693(98)00454-7), arXiv:hep-ph/9801442.
- [41] X. Chen, T. Gehrmann, N. Glover, M. Höfer, A. Huss, Isolated photon and photon+jet production at NNLO QCD accuracy, J. High Energy Phys. 04 (2020) 166, [https://doi.org/10.1007/JHEP04\(2020\)166](https://doi.org/10.1007/JHEP04(2020)166), arXiv:1904.01044.
- [42] F. Siegert, A practical guide to event generation for prompt photon production with Sherpa, J. Phys. G 44 (4) (2017) 044007, <https://doi.org/10.1088/1361-6471/aa5f29>, arXiv:1611.07226.
- [43] I.W. Stewart, F.J. Tackmann, W.J. Waalewijn, N-jettiness: an inclusive event shape to veto jets, Phys. Rev. Lett. 105 (2010) 092002, <https://doi.org/10.1103/PhysRevLett.105.092002>, arXiv:1004.2489.
- [44] S. Frixione, P. Nason, C. Oleari, Matching NLO QCD computations with parton shower simulations: the POWHEG method, J. High Energy Phys. 11 (2007) 070, <https://doi.org/10.1088/1126-6708/2007/11/070>, arXiv:0709.2092.
- [45] A. Idilbi, X.-d. Ji, F. Yuan, Resummation of threshold logarithms in effective field theory for DIS, Drell-Yan and Higgs production, Nucl. Phys. B 753 (2006) 42–68, <https://doi.org/10.1016/j.nuclphysb.2006.07.002>, arXiv:hep-ph/0605068.
- [46] T. Becher, M. Neubert, B.D. Pecjak, Factorization and momentum-space resummation in deep-inelastic scattering, J. High Energy Phys. 01 (2007) 076, <https://doi.org/10.1088/1126-6708/2007/01/076>, arXiv:hep-ph/0607228.
- [47] A. Hornig, C. Lee, I.W. Stewart, J.R. Walsh, S. Zuberi, Non-global structure of the $O(\alpha_s^2)$ dijet soft function, J. High Energy Phys. 08 (2011) 054, [https://doi.org/10.1007/JHEP08\(2011\)054](https://doi.org/10.1007/JHEP08(2011)054), arXiv:1105.4628.
- [48] D. Kang, O.Z. Labun, C. Lee, Equality of hemisphere soft functions for e^+e^- , DIS and pp collisions at $O(\alpha_s^2)$, Phys. Lett. B 748 (2015) 45–54, <https://doi.org/10.1016/j.physletb.2015.06.057>, arXiv:1504.04006.
- [49] R. Kelley, M.D. Schwartz, R.M. Schabinger, H.X. Zhu, The two-loop hemisphere soft function, Phys. Rev. D 84 (2011) 045022, <https://doi.org/10.1103/PhysRevD.84.045022>, arXiv:1105.3676.
- [50] P.F. Monni, T. Gehrmann, G. Luisoni, Two-loop soft corrections and resummation of the thrust distribution in the dijet region, J. High Energy Phys. 08 (2011) 010, [https://doi.org/10.1007/JHEP08\(2011\)010](https://doi.org/10.1007/JHEP08(2011)010), arXiv:1105.4560.
- [51] J.R. Gaunt, M. Stahlhofen, F.J. Tackmann, The quark beam function at two loops, J. High Energy Phys. 04 (2014) 113, [https://doi.org/10.1007/JHEP04\(2014\)113](https://doi.org/10.1007/JHEP04(2014)113), arXiv:1401.5478.
- [52] T. Gehrmann, L. Tancredi, Two-loop QCD helicity amplitudes for $q\bar{q} \rightarrow W^\pm\gamma$ and $q\bar{q} \rightarrow Z^0\gamma$, J. High Energy Phys. 02 (2012) 004, [https://doi.org/10.1007/JHEP02\(2012\)004](https://doi.org/10.1007/JHEP02(2012)004), arXiv:1112.1531.
- [53] T. Sjöstrand, S. Mrenna, P.Z. Skands, A brief introduction to PYTHIA 8.1, Comput. Phys. Commun. 178 (2008) 852–867, <https://doi.org/10.1016/j.cpc.2008.01.036>, arXiv:0710.3820.
- [54] MUNICH is the abbreviation of “MULTI-channel Integrator at Swiss (CH) precision”—an automated parton level NLO generator by S. Kallweit, in preparation.

- [55] F. Buccioni, S. Pozzorini, M. Zoller, On-the-fly reduction of open loops, *Eur. Phys. J. C* 78 (1) (2018) 70, <https://doi.org/10.1140/epjc/s10052-018-5562-1>, arXiv:1710.11452.
- [56] F. Buccioni, J.-N. Lang, J.M. Lindert, P. Maierhöfer, S. Pozzorini, H. Zhang, M.F. Zoller, OpenLoops 2, *Eur. Phys. J. C* 79 (10) (2019) 866, <https://doi.org/10.1140/epjc/s10052-019-7306-2>, arXiv:1907.13071.
- [57] T. Gehrmann, E. Remiddi, Numerical evaluation of harmonic polylogarithms, *Comput. Phys. Commun.* 141 (2001) 296–312, [https://doi.org/10.1016/S0010-4655\(01\)00411-8](https://doi.org/10.1016/S0010-4655(01)00411-8), arXiv:hep-ph/0107173.
- [58] S. Bailey, T. Cridge, L.A. Harland-Lang, A.D. Martin, R.S. Thorne, Parton distributions from LHC, HERA, Tevatron and fixed target data: MSHT20 PDFs, *Eur. Phys. J. C* 81 (4) (2021) 341, <https://doi.org/10.1140/epjc/s10052-021-09057-0>, arXiv:2012.04684.
- [59] A. Buckley, J. Ferrando, S. Lloyd, K. Nordström, B. Page, M. Rüfenacht, M. Schönherr, G. Watt, LHAPDF6: parton density access in the LHC precision era, *Eur. Phys. J. C* 75 (3) (2015) 132, <https://doi.org/10.1140/epjc/s10052-015-3318-8>, arXiv:1412.7420.
- [60] M.A. Ebert, J.K.L. Michel, I.W. Stewart, F.J. Tackmann, Drell-Yan q_T resummation of fiducial power corrections at N^3 LL, *J. High Energy Phys.* 04 (2021) 102, [https://doi.org/10.1007/JHEP04\(2021\)102](https://doi.org/10.1007/JHEP04(2021)102), arXiv:2006.11382.
- [61] C. Bierlich, et al., Robust independent validation of experiment and theory: rivet version 3, *SciPost Phys.* 8 (2020) 026, <https://doi.org/10.21468/SciPostPhys.8.2.026>, arXiv:1912.05451.