

# Macrophages as determinants and regulators of fibrosis in systemic sclerosis

Yehya Al-Adwi<sup>1</sup>, Johanna Westra<sup>2</sup>, Harry van Goor<sup>3</sup>, Janette K. Burgess<sup>3</sup>, Christopher P. Denton<sup>4,5</sup>,  
Douwe J. Mulder<sup>1\*</sup>

<sup>1</sup>University of Groningen, University Medical Centre Groningen, Department of Internal Medicine, Division of Vascular Medicine, Groningen, The Netherlands.

<sup>2</sup>University of Groningen, University Medical Centre Groningen, Department of Rheumatology and Clinical Immunology, The Netherlands.

<sup>3</sup>University of Groningen, University Medical Centre Groningen, Department of Pathology and Medical Biology, Groningen, The Netherlands.

<sup>4</sup>UCL Division of Medicine, University College London, London, UK.

<sup>5</sup>UCL Centre for Rheumatology and Connective Tissue Diseases, Royal Free Hospital, London, UK.

\*Corresponding author: Douwe J Mulder, MD, PhD, Department of Internal Medicine, Division of Vascular Medicine, University Medical Centre Groningen, University of Groningen, Groningen, The Netherlands.

Postal address: Hanzeplein 1, 9700RB, Groningen, The Netherlands.

E-mail: d.j.mulder@umcg.nl

ORCID: 0000-0003-3715-6474


© The Author(s) 2022. Published by Oxford University Press on behalf of the British Society for Rheumatology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

Abstract

Systemic Sclerosis (SSc) is a multiphase autoimmune disease with a well-known triad of clinical manifestations including vasculopathy, inflammation and fibrosis. Although a plethora of drugs has been suggested as potential candidates to halt SSc progression, nothing has proven clinically efficient. In SSc, both innate and adaptive immune systems are abnormally activated fuelling fibrosis of the skin and other vital organs. Macrophages have been implicated in the pathogenesis of SSc and are thought to be a major source of immune dysregulation. Due to their plasticity, macrophages can initiate and sustain chronic inflammation when classically activated while, simultaneously or parallelly, when alternatively activated they are also capable of secreting fibrotic factors. Here, we briefly explain the polarization process of macrophages. Subsequently, we link the activation of macrophages and monocytes to the molecular pathology of SSc, and illustrate the interplay between macrophages and fibroblasts. Finally, we present recent/near-future clinical trials and discuss novel targets related to macrophages/monocytes activation in SSc.

Graphical Abstract

Created with BioRender.com



51  
52  
53  
54  
55

Key words: scleroderma, systemic sclerosis, macrophages, monocytes, fibrosis, potential targeted therapeutics.

56

**Key messages:**

- Systemic sclerosis (SSc) is a heterogeneous disease and monocytes/macrophages are central within this heterogeneity.
- Plasticity of monocytes/macrophages allow them to reflect and affect all disease phases of SSc.
- There is an urgent unmet need for personalized medicine to treat SSc patients.

56

**Systemic Sclerosis**

58 Systemic sclerosis (SSc) is an autoimmune disease that involves microangiopathy, early inflammation  
59 and progressive fibrosis of the skin and internal organs (1–6). Limited cutaneous (lcSSc), diffuse  
60 cutaneous (dcSSc) and sine SSc forms of SSc can present as stable conditions but can also progress to  
61 severe disease modes with increased morbidity and mortality (7–9). Although advances have been  
62 made in biomarker discovery for SSc, it is still difficult to predict which patients are going to progress  
63 to severe fibrotic disease for which no disease-modifying treatment is currently available (9,10).  
64 Interstitial lung disease, a result of inflammatory and fibrosing processes, is the leading cause of  
65 mortality in SSc patients who develop this complication (~50%) (10,11). Therefore, new treatment  
66 strategies are urgently needed to attenuate progression and potentially modify the disease course.  
67 Recently, macrophages have captured the interest of the SSc scientific community. This interest is due  
68 to the abundance of these cells in affected tissues and their potential in driving both inflammatory, as  
69 well as fibrotic processes (12). Macrophages possess great plasticity which allows them to adopt  
70 different polarization states.

71

72 This review focuses on the polarization dynamics of macrophages and their precursors (monocytes) in  
73 SSc, the influence of monocytes and macrophages on the disease course and the role of cytokines in  
74 the activation of macrophages in SSc patients. We also discuss the interplay between activated  
75 macrophages and fibroblasts. Furthermore, we describe the heterogeneity of the disease and what a  
76 multi-phased disease means for the clinic when it comes to treatment. Finally, we discuss how  
77 focussing on macrophage polarization could potentially facilitate novel targeted therapy discovery for  
78 SSc patients.

79

**Macrophage in health and disease**

81 Macrophages play a principal role in maintaining (physiological) homeostasis by engulfing, degrading  
82 and clearing of cellular debris, dead cells, and cancer cells (13). Additionally, macrophages function as

1  
2  
3 83 reparatory machines, playing an essential role in the wound healing process, allowing quicker post-  
4 84 insult recovery (14). Their capacity of releasing chemoattractants and cytokines to recruit other  
5 85 effector immune cells, make them crucial in terms of host defence response (15).  
6  
7  
8

9 86

10  
11 87 Macrophage tissue infiltration is a known phenomenon in most autoimmune diseases including SSc  
12 88 (16–24). Accumulating evidence has revealed a crucial role of innate immune reactions in driving not  
13 89 only disease flare ups (causing progression) but also contributing to the ignition processes in such  
14 90 diseases (25,26).  
15  
16  
17  
18

19 91

### 20 21 92 Macrophage polarization

22  
23 93 Monocytes are circulating cells, composing around 10% of the cells in healthy peripheral blood, that  
24 94 are known to be precursors of macrophages and dendritic cells forming the mononuclear phagocytic  
25 95 system. CD14 is expressed on the surface of monocytes and is used as a marker to identify them. A  
26 96 complex network of stimuli including cytokines, chemokines and inter-cellular signalling is coordinated  
27 97 in a healthy individual to regulate the differentiation of monocytes to macrophages (27,28).  
28  
29  
30  
31

32 98

33  
34 99 Historically, macrophages were classified according to their activation pattern into classically activated  
35 100 macrophages (M1) or alternatively activated macrophages (M2). Certain cytokines will  
36 101 polarize/differentiate macrophages into a pro-inflammatory phenotype (classical) which handles  
37 102 pathogen destruction. On the other side of the spectrum, another set of cytokines, chemokines and  
38 103 hormones skew the activation of macrophages into a healing/regenerative phenotype (alternative)  
39 104 which are (pro)fibrotic, anti-inflammatory and in charge of tissue repair (29,30). The contribution of  
40 105 specific factors will be explored later in this review. Recent scientific observations on macrophage  
41 106 classification confirmed that the earlier nomenclature is based on *in vitro* experimentation and does  
42 107 not represent *in vivo* scenarios. Thus, macrophage polarization is rather considered as a continuum  
43 108 than two distinct populations where classical activation is at one end and alternative activation is at  
44 109 the other. Therefore, macrophage phenotypes can slide across this spectrum of the  
45 110 classical/alternative paradigm of activations (31).  
46  
47  
48  
49

50  
51  
52 111

53  
54  
55  
56  
57 112 Classically activated macrophages maintain inflammation as a defence mechanism to ward off  
58 113 intruders. To be able to function in that manner, classically activated macrophages express a distinct

1  
2  
3 114 set of surface receptors (see table.1) that allow them to respond adequately to specific stimuli (27,30).  
4  
5 115 An important stimulus for macrophage polarization towards the classical phenotype is interferon-  
6  
7 116 gamma (IFN- $\gamma$ ) through its ability to directly activate effector genes including antiviral proteins,  
8  
9 117 microbicidal molecules, phagocytic receptors, chemokines and cytokines (32). Additionally, IFN- $\gamma$  can  
10  
11 118 indirectly activate macrophages by enhancing their reaction to other stimuli through what is known  
12  
13 119 as “priming” (32). When macrophages are stimulated with IFN- $\gamma$ , the result is Janus kinase 1 (JAK 1)  
14  
15 120 and JAK 2 activation, and signal transducer and activator of transcription 1 (STAT1)/interferon  
16  
17 121 regulatory factors (IRF) signalling, leading to differentiation to the classical phenotype as the product  
18  
19 122 (33). This activation pathway is not the only means to produce classical macrophages. During bacterial  
20  
21 123 infections, lipopolysaccharide (LPS) is present abundantly in the body which is well-recognized for  
22  
23 124 stimulating the classical polarization through its binding with toll-like receptors 2 and 4 (TLRs) which  
24  
25 125 in turn initiates nuclear factor-light-chain-enhancer of activated B cells (NF- $\kappa$ B), activator protein-1  
26  
27 126 (AP-1), IRF and STAT1 signalling (34). Finally, granulocyte-macrophage colony-stimulating factor (GM-  
28  
29 127 CSF) is capable of inducing the classical phenotype through the activation of the JAK2 pathway (35)  
30  
31 128 (Fig.1).

32  
33 129  
34  
35 130 On the other side of the spectrum lies the alternatively activated macrophage phenotype. The central  
36  
37 131 aim of alternative macrophages is to release anti-inflammatory cytokines and recruit specific tissue-  
38  
39 132 regenerating cells (36,37). Activated adaptive immune cells such as mast cells, basophils and type 2 T  
40  
41 133 helper ( $T_{H2}$ ) cells release IL-4/IL-13 which, in turn, stimulate alternative polarization through the JAK 1  
42  
43 134 and JAK3/STAT6 pathway. This pathway is considered to be the canonical pathway for alternative  
44  
45 135 activation (38). However, to have a more discrete nomenclature within the alternative phenotype, the  
46  
47 136 IL-4/IL-13-induced activation of macrophages is named M2a (39). Other specific alternative subtypes  
48  
49 137 can be induced by other stimuli such as immune complexes and TLR ligands (40). Such interactions  
50  
51 138 shut down the proinflammatory cytokine IL-12 release and substitute it with the profibrotic cytokine  
52  
53 139 IL-10. This macrophage-activation state involves spleen tyrosine kinase (Syk) and phosphoinositide 3-  
54  
55 140 kinase (PI3K) activation and is known as M2b. The M2b macrophages can release both pro-  
56  
57 141 inflammatory and anti-inflammatory cytokines (41) (see table.1). Both glucocorticoids and IL-10 are  
58  
59 142 able to induce the third subtype of alternative macrophages; M2c through the glucocorticoid receptor  
60  
143 (GRC) or IL-10R, respectively. The M2c subtype has a strong fibroproliferative cytokine signature  
144 releasing IL-10 and TGF- $\beta$  cytokines. The fourth subtype is M2d macrophages which are activated  
145 through the binding of TLR agonists to the Adenosine 2 receptor. Consequently, significant  
146 suppression of pro-inflammatory cytokine release and promotion of anti-inflammatory cytokines

1  
2  
3 147 production occurs (42) (Fig.1). The detailed macrophage polarization pathways, different cytokine  
4  
5 148 signatures and distinct surface markers have all been previously described elsewhere (43,44).  
6  
7 149

### 10 150 Monocyte/Macrophage signature in Systemic Sclerosis

11 151 Plasticity allows macrophages to influence all phases of SSc. Although limited in number, several  
12 152 studies have investigated the role of macrophage polarization in SSc pathogenesis.  
13  
14 153

15 154 Higashi-Kuwata et al. utilized flow cytometry and immunohistochemistry (IHC) techniques to show  
16 155 that the blood and skin of SSc, respectively, have higher expression of macrophage fibrotic markers  
17 156 (CD163 and CD204) compared to healthy controls. Flow cytometry was used on isolated peripheral  
18 157 blood mononuclear cells (PBMCs) where CD14 surface marker was used to gate for monocytes and  
19 158 CD163 and CD204 surface markers were used to detect the fibrogenic phenotype. Skin biopsies were  
20 159 stained with antibodies against the pan-macrophage surface markers CD68, CD163 and CD204. They  
21 160 found an enhanced expression of CD163 and CD204 on the PBMCs and in skin biopsies from SSc  
22 161 patients compared to controls. Consequently, the authors suggested that the activation status of  
23 162 monocytes/macrophages in SSc patients is profibrotic compared to controls. However, they did not  
24 163 investigate M1 markers to detect classical polarization. (45). Mathai and colleagues showed that CD14  
25 164 monocytes isolated from PBMCs of SSc-associated interstitial lung disease (SSc-ILD) patients show  
26 165 higher expression of CD163 compared to controls. Interestingly, when CD14+ monocytes were  
27 166 isolated from PBMCs of SSc-ILD patients and treated *in vitro* with LPS, a classical activation inducer,  
28 167 these monocytes were skewed into a more profibrotic pattern in contrast to a proinflammatory  
29 168 profile, with more CD163+, CCL18 and IL-10 expression compared to controls (46). However, as the  
30 169 primary question in this study was whether monocytes from SSc-ILD patients have higher expression  
31 170 of profibrotic markers, inflammatory markers were not studied.  
32  
33 171

34 172 In a monocyte-derived macrophage (MDMs) *in vitro* transcriptomic study, Morano-Moral et al  
35 173 identified 602 genes that were differentially regulated in SSc patients compared to controls.  
36 174 Upregulated genes were related to hypoxia, glycolysis and mTOR pathways while IFN- $\gamma$  response  
37 175 pathways were downregulated. This study also highlighted gasdermin A specific variant as an SSc risk  
38 176 factor when upregulated, suggesting that MDMs could be the reason behind dysregulated pyroptosis  
39 177 in SSc. This study robustly links SSc pathogenesis with genetic changes and presents a transcriptomic  
40 178 signature in MDMs of SSc patients (47).  
41  
42 179

1  
2  
3 180 It is known that the affected skin of SSc patients is infiltrated with immune cells, especially T cells and  
4  
5 181 macrophages. To understand the recruitment of macrophages to the skin, researchers have studied  
6  
7 182 the chemokine gene expression in affected skin from SSc patients. RT-qPCR data of homogenised skin  
8  
9 183 showed a higher expression of CCL2, CCL5, CCL18, CCL19 and CXCL13 in dcSSc patients when  
10  
11 184 compared to control skin. Skin biopsies from dcSSc patients exposed a colocalization of CD163+  
12  
13 185 macrophage subset with CCL19, strongly suggesting the release of CCL19 from the CD163+  
14  
15 186 macrophage subset. Moreover, not only was CCL2, an important macrophage recruiting chemokine,  
16  
17 187 expression positively correlated with skin thickening in dcSSc skin but also serum levels of CCL2 were  
18  
19 188 elevated in these patients. These localized and systemic correlations are strongly suggestive of the  
20  
21 189 involvement of the alternative macrophage phenotype in the development of skin lesions in SSc.  
22  
23 190 Another group studied patients with SSc-ILD compared to those SSc patients without ILD. There were  
24  
25 191 increased mixed classic (CD80, CD86, TLR2 and TLR4) and alternative (CD206, CD204 and CD163)  
26  
27 192 circulating monocytes in patients with SSc-ILD. (48). Additionally, they found that these markers were  
28  
29 193 significantly elevated on PBMCs isolated from SSc patients compared to healthy controls (49). These  
30  
31 194 results point out that the circulating monocytes from SSc patients have enriched classic and alternative  
32  
33 195 markers compared to controls while this enrichment is even greater when ILD is present.  
34  
35 196

36  
37 197 On a single-cell level, RNA-sequencing of SSc-ILD lung tissue revealed several monocyte/macrophage  
38  
39 198 subgroups in which SPP1<sup>hi</sup> proliferating macrophages were more predominant in SSc-ILD lungs  
40  
41 199 compared to controls (50). SPP1 macrophages have been attributed to lung fibrosis through the  
42  
43 200 activation of myofibroblasts in idiopathic pulmonary fibrosis (51) which could be having the same role  
44  
45 201 in SSc-ILD. On the same level, RNA-sequencing of dcSSc skin tissues revealed innate immune system  
46  
47 202 activation(52). Specifically, macrophages highly expressing Fcy receptor IIIA were only associated with  
48  
49 203 dcSSc skin but not in healthy skin. Importantly, the proliferating macrophages were exclusively  
50  
51 204 detected in dcSSc skin but not in healthy ones (52). Thus, it is plausible that proliferating macrophages  
52  
53 205 are fundamental to skin and lung disease progression in SSc.  
54  
55 206

56  
57 207 Due to their plasticity, monocytes/macrophages could be an important link for the transition from the  
58  
59 208 inflammatory to the fibrotic phase in SSc pathology. Perhaps, monocyte/macrophage polarization  
60  
209 shifts along the classic/alternative spectrum of activation to a more fibrotic state over time due to  
210 intracellular changes and differential presence of cytokines and chemokines in their environment. The  
211 change in the cytokine and chemokine profile can be attributed to reactive B cells (IL-6 release) and  
212 activated CD4+ T<sub>H2</sub> cells (IL-13 and IL-4 release) (53,54). Consequently, monocytes/macrophages  
213 become profibrotic and start to release fibrotic factors that lead to the activation of more

1  
2  
3 214 monocyte/macrophages (and of fibroblasts) into the profibrotic phenotype generating an autocrine  
4  
5 215 loop. Moreover, the mounting recruitment of fibrogenesis-effector cells such as fibrocytes and  
6  
7 216 fibroblasts into affected tissues, and their activation by the released fibroproliferative chemokines are  
8  
9 217 key events in SSc- related tissue fibrosis.

10  
11 218

### 12 13 219 The interplay between fibroblasts and macrophages

14 220 Activation of monocytes/macrophages is crucial for stimulation of the fibrosis effector cells  
15 221 (fibroblasts) in affected tissues. Bhandari et al. (55) showed that the activation of fibroblasts is  
16 222 dependent on SSc plasma-differentiated macrophages. SSc plasma significantly activated monocytes  
17 223 from both control and SSc groups into the profibrotic (alternative) phenotypes when compared to  
18 224 monocytes cultured with control plasma. Moreover, significantly higher mRNA and protein expression  
19 225 and production of CCL2, IL-6 and TGF- $\beta$  were reported in SSc plasma-cultured compared to control-  
20 226 cultured monocytes. This experiment illustrated that SSc plasma can differentiate control monocytes  
21 227 into SSc phenotype macrophages. Additionally, RT-qPCR data from dermal fibroblasts revealed  
22 228 overexpression of  $\alpha$ -SMA in SSc fibroblasts co-cultured with SSc plasma-differentiated macrophages,  
23 229 compared to healthy dermal fibroblasts co-cultured with SSc plasma-differentiated macrophages.  
24 230 These data indicate that SSc macrophages induce and activate dermal fibroblasts into becoming  
25 231 fibrogenic cells through fibroblast to myofibroblast transdifferentiation (55).

26  
27 232

28 233 In *in vivo* scenarios, macrophages have to be in close proximity to fibroblasts to stimulate them to  
29 234 myofibroblasts (56). Pakshir et al (57) described how mechanosensation and integrins help  
30 235 myo/fibroblasts attract macrophages to the vicinity of the fibrotic niche. Fibroblasts establish ECM  
31 236 cues through remodeling collagens to form deformation fields in the collagen mesh which guides  
32 237 macrophages to come closer to myo/fibroblasts. Importantly, these ECM alterations have more far-  
33 238 reaching effects than chemotaxis.

34  
35  
36 239 Thus, the interaction between macrophages and myo/fibroblasts is necessary to establish a  
37 240 progressive fibrotic niche.

38  
39 241

40  
41 242

42  
43 243

### 44 244 Emerging role of oxidative stress in monocyte/macrophage polarization in SSc

45  
46 245 Antioxidant/oxidant imbalance is thought to be connected to SSc pathogenesis (58) The nuclear  
47 246 factor erythroid 2 (NF-E2)-related factor 2 (Nrf-2) is an important cellular sensing protein for



1  
2  
3 247 oxidative stress which- in turn- can stimulate the transcription of antioxidants including glutathione  
4 248 (GSH). In an SSc mouse model, Nrf<sup>-/-</sup> knockdown and wild-type (control) mice were intradermally  
5 249 injected with hypochloric acid (a substance to induce oxidative stress). The Nrf<sup>-/-</sup> mice showed more  
6 250 severe inflammation and fibrosis than controls. Importantly, in the skin of hypochloric acid-treated  
7 251 mice, the Nrf<sup>-/-</sup> type had a more pronounced M2 polarization marker profile than the wild-type  
8 252 mice. This indicates that oxidative stress induces a shift towards M2 polarization and suggests a  
9 253 strong link between Nrf-2 function, alternative polarization, fibroblasts activation and fibrogenesis  
10 254 (59).

11 255

12 256 **Systemic Sclerosis is a multi-phase disease – Interventional remarks focusing on targeting**  
13 257 **monocytes/macrophages** (summary of current and prospective/promising SSc therapeutics  
14 258 is stated in table.2)

15 259

16 260 SSc-investigated targeted therapies

17 261 In SSc patients who present early in their disease course, inflammation is generally the predominant  
18 262 process activated, especially in progressive dcSSc. As elaborated, M1 phenotype macrophage  
19 263 activation may be central at this stage, and it would be reasonable to introduce drugs that target the  
20 264 effector pathways early. In patients with SSc-ILD, monocytes are known to produce higher amounts  
21 265 of IL-6 compared to healthy controls (60). This high production is strongly associated with SSc  
22 266 pathogenesis since it leads to the activation, differentiation and proliferation of T lymphocytes.  
23 267 Tocilizumab (anti-IL-6R monoclonal antibody) is of growing interest and use in clinical practice. It is  
24 268 FDA-approved to slow the rate of decline of lung function in adult patients with SSc-ILD (61,62).  
25 269 Moreover, IL-6 is abundantly produced by activated B cells expressing CD20. The DESIRES RCT showed  
26 270 that using rituximab, an anti-CD20 monoclonal antibody, in SSc patients resulted in a significant  
27 271 reduction in mRSS compared to the placebo arm (63). Rituximab's positive results can be indirectly  
28 272 attributed to the blockage of macrophage polarization leading to mitigated skin and lung disease  
29 273 (64,65).

30 274

31 275 An IFN type I serum profile is related to higher mRSS and HAQ-Disability Index. Researchers have  
32 276 indirectly targeted IFN type I by targeting CD52. In a translational study, transcriptomic analyses of  
33 277 circulating CD14<sup>+</sup> monocytes obtained from SSc patients revealed enhanced expression of IFN I-  
34 278 related genes compared to healthy controls. These monocytes also displayed down-regulated CD52

1  
2  
3 279 expression which is an important T cell inhibitory antigen. Interestingly, when healthy monocytes were  
4  
5 280 treated with an anti-CD52 antibody, enhanced activation of IFN I pathways were achieved.  
6  
7 281 Consequently, targeting the CD52-IFN I pathway is a promising approach in early SSc patients (66).  
8

9 282

10  
11 283 Targeting the profibrotic cytokines IL-4 and IL-13 to prevent further activation of  
12  
13 284 monocytes/macrophages to the profibrotic forms has shown promising results. Romilkimab was  
14  
15 285 developed as a humanized bispecific mAB against both IL-4 and IL-13. When neutralizing these serum  
16  
17 286 elevated cytokines, the paracrine and autocrine activation loops of macrophages are blocked. Indeed,  
18  
19 287 a phase 2 RCT in early dcSSc was performed where romilkimab efficacy was tested vs placebo. After  
20  
21 288 24 weeks, patients who have been treated with romilkimab had a significant improvement in their  
22  
23 289 mRSS compared to the placebo group (67).

24 290

25  
26 291 Although it is known to work as a multi tyrosine kinase inhibitor, nintedanib also functions by  
27  
28 292 disturbing the expression of surface markers, and/or the chemokine and cytokine signature of  
29  
30 293 monocyte-derived macrophages. It also inhibits the phosphorylation of the colony-stimulating factor  
31  
32 294 1 receptor in monocyte/macrophages which is essential in activation and polarization of these cells.  
33  
34 295 When monocytes were stimulated to polarize to classical or alternative macrophages *-in vitro*-  
35  
36 296 subsequent to treatment with nintedanib, several alterations were observed. First, classical  
37  
38 297 macrophages continued to express classical surface markers at the same level as untreated  
39  
40 298 macrophages but released significantly less proinflammatory cytokines. Second, the alternatively-  
41  
42 299 stimulated macrophages had a significant decrease in their M2 markers while their profibrotic  
43  
44 300 cytokines and chemokines release remained comparable to untreated alternative macrophages  
45  
46 301 (68,69). Clinically, Azuma et al. (70) performed a phase 3 RCT where SSc patients with at least 10%  
47  
48 302 lung fibrosis on HRCT were included. The primary endpoint was the annual rate of decline in FVC. After  
49  
50 303 52 weeks, the annual rate of decline in FVC was significantly higher in the placebo arm than in the  
51  
52 304 treatment arm. Based on these data, nintedanib was the first drug to be approved for treating SSc-ILD  
53  
54 305 (71–73).

55 306

56  
57 307 Pirfenidone has shown inhibitory effects on rat alternatively-activated lung macrophages cultured *in*  
58  
59 308 *vitro*. This was demonstrated by a significant reduction of TGF- $\beta$  release and lower expression of M2  
60  
61 309 surface markers when macrophages were treated with pirfenidone. When the supernatant of the  
62  
63 310 pirfenidone-treated macrophages was used to treat rat lung fibroblasts, suppressed proliferation, and  
64  
65 311 collagen mRNA expression and production were observed in these fibroblasts (74). In light of these

1  
2  
3 312 data, Khanna et al. (75) and Sharma et al. (76) performed phase 2 clinical trials to assess the efficacy  
4 313 of pirfenidone in SSc-ILD patients. Although the data from these trials have not shown significant  
5 314 differences in lung function decline between the treatment and placebo groups, pirfenidone was well  
6 315 tolerable and appeared likely to maintain lung function better than the placebo.  
7  
8  
9

10 316

11 317 Current studies

12  
13 318 Several studies are currently investigating pharmaceutical agents that could hinder the activation  
14 319 and/or the release of cytokines/chemokines from monocytes/macrophages. For example, in the “Hit  
15 320 hard and early” study [NCT03059979], very early diagnosed SSc (VEDOSS) patients are being treated  
16 321 with high dose methylprednisolone, potentially preventing early vasculopathy by forcing attenuation  
17 322 of inflammation (77). It is also highly plausible that the mechanism behind this strategy is mitigating  
18 323 the polarization of macrophages towards the classical inflammatory phenotype, as this is a known  
19 324 effect of prednisolone (78,79). Another strategy is being investigated using upfront autologous  
20 325 hematopoietic stem cell transplantation (AHSCT) in early dcSSc patients with the aim of resetting the  
21 326 immune system (the UPSIDE study; [NCT04464434]) (80). Monocytes derived from myeloid  
22 327 progenitors are activated and play a role in igniting and perpetuating the inflammatory, and thereafter  
23 328 the fibrosis processes, in SSc patients. Studies show that dcSSc patients have higher expression of  
24 329 CD16+ monocytes which are known to have enhanced pro-inflammatory activation (81). The upfront  
25 330 depletion of these cells, coupled with replacing them with “normal/undiseased” precursors through  
26 331 AHSCT could yield monocytes that can suppress the pathogenetic pathways of SSc by enhancing T<sub>regs</sub>  
27 332 cell production, inhibiting fibroblast to myofibroblast transdifferentiation and suppressing CD4 T cells  
28 333 proliferation (82).  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39

40 334

41  
42  
43 335 The SCLERO JAK [NCT04206644] study is investigating the efficacy of the JAK 1/2 inhibitor ruxolitinib  
44 336 in SSc patients. one of the outcomes aims to gather a greater understanding of the impact of this drug  
45 337 on the activation states of monocyte-derived macrophages obtained from SSc patients. It is  
46 338 hypothesized that blocking the JAK-STAT pathway would attenuate the profibrotic properties of  
47 339 monocyte-derived macrophages. This will be tested in an *in vitro* model by measuring CCL18 levels in  
48 340 the culture media of ruxolitinib-treated SSc macrophages compared to untreated (as a primary  
49 341 outcome). In addition, macrophage surface markers studies will be performed.  
50  
51  
52  
53  
54  
55  
56  
57

58 342

59 343 GSK2330811 is a humanized monoclonal antibody against the oncostatin M (OSM) protein, which is  
60 344 implicated in inflammation, fibrosis and vasculopathy, typical features of SSc pathogenesis. Activated

1  
2  
3 345 monocytes/macrophages are known to produce OSM which alters fibroblasts -among other  
4 346 connective tissue cells- production of cytokines and chemokines such as MCP-1 and IL-6 which in turn  
5 347 affects the polarization of macrophages (paracrine activation loop) (83,84). After showing a well-  
6 348 tolerated safety profile in phase I clinical trial (85), proof of mechanism phase II randomized clinical  
7 349 trial [NCT03041025] in dcSSc patients is currently being undertaken and it is hoped the data will be  
8 350 available soon.

13  
14 351

15  
16 352 Future perspectives – promising therapeutics/pathways

17  
18 353 SSc upstream processes involve both the innate and adaptive immune systems. repurposing drugs  
19 354 from other medical fields such as oncology and haematology to the field of autoimmune diseases is  
20 355 not unusual. Therefore, we are suggesting the following potential drugs abide the recent  
21 356 understanding of SSc pathogenesis.

22  
23 357

24  
25 358 Targeting purinergic signalling may ameliorate fibrosis. As explained above, adenosine can skew  
26 359 macrophages towards the alternative phenotype. Degradation of adenosine using pegylated  
27 360 recombinant adenosine deaminase reduced fibrogenesis in SSc preclinical models. Adenosine  
28 361 deaminase has also shown promising results regarding vasculopathy and inflammation in a mouse  
29 362 model of SSc (86). The effects of such a drug should be examined in a clinical trial to better  
30 363 comprehend its potential efficacy in SSc patients.

31  
32 364

33 365 Direct targeting of proliferating monocyte progenitor cells without affecting other progenitor cells or  
34 366 mature monocytes could be an approach to diminish monocytes' contribution to pathophysiology in  
35 367 SSc patients. Using dimeric pyrrolobenzodiazepine (dPBD)-conjugated anti-CD64 antibody (anti-CD64-  
36 368 dPBD), Izumi et al (87) were able to selectively induce apoptosis in proliferating human monocyte  
37 369 progenitors.

38  
39 370

40 371 Targeting the migration of inflammatory monocytes to sites of injury using small interfering RNA  
41 372 (siRNA) is another promising approach that could benefit SSc patients. CCR2 chemokine receptor is  
42 373 known to be over-expressed on inflammatory monocytes. Targeting cells with high levels of this  
43 374 receptor with nanoparticles-containing anti-CCR2 siRNA showed promising results in several  
44 375 inflammatory diseases in preclinical settings. In these preclinical models, anti-CCR2 siRNA was able to  
45 376 silence CCR2 mRNA of inflammatory monocytes and consequently reduced migration as well as  
46 377 numbers of monocyte-derived macrophages without affecting other healing, physiologically essential  
47 378 functions of monocytes and associated macrophages (88).

379

380 Non-specific memory of the innate immune system, also as known as “Trained immunity”, is thought  
381 to be part of the enhanced and continuity of cytokines and chemokines production by monocytes in  
382 SSc patients. Pharmacological blocking of upstream processes of trained immunity using NOD2 and  
383 dectin 1 inhibitors, GSK669 or laminarin, respectively could be beneficial (89). Additionally,  
384 nanomedicine could offer another novel approach for directly targeting and skewing localized  
385 inflammatory monocytes to a less inflammatory phenotype through limiting epigenetic and metabolic  
386 changes (89). Statin-encapsulated reconstituted high-density lipoprotein (HDL) nanobiologic is a  
387 promising tool targeting inflammatory monocytes and macrophages. Such a drug has shown  
388 promising results in inflammatory atherosclerotic plaques (90) but has not yet been applied to  
389 autoimmune diseases including SSc.

390

391 Reprogramming alternatively-activated macrophages towards apoptosis or classical polarization is a  
392 well-characterized strategy in tumour research. RP-182 and RP-832c, host immune peptides, can  
393 target CD206 alternative MDMs in lung fibrosis leading to alleviation of fibrogenesis. This mechanism  
394 could be beneficial for SSc patients, especially for those who are suffering from dermal and lung  
395 fibrosis (91,92).

396

### 397 Conclusion

398 SSc is a multi-organ, multi-phase disease with various potential pharmaceutical interventions. In order  
399 to combat its complications, it is first necessary to identify the phase of the disease. Interpretations of  
400 literature and previous research highlight monocytes/macrophages as promising biomarkers that  
401 dynamically change according to disease progression reflecting disease status. They can also be  
402 considered as potential therapeutic targets through modulation of their polarization.

403

404 Due to the heterogeneity of SSc pathogenesis, examination of SSc patients must recognise that each  
405 patient is a unique case. This is a unique opportunity to address the unmet need for personalized  
406 medicine in treating SSc patients. Most SSc complications share similar phenotypical and molecular  
407 characteristics, however, several important differences have been observed when it comes to  
408 progression and initiating factors. Finally, although this personalized approach is still under  
409 development, each SSc patient requires a special set of therapeutics according to their disease phase,  
410 active pathogenesis pathway, and number and type of complications.

411

### 412 Acknowledgements:

1  
2  
3 413 All figures have been created using BioRender.com  
4

5 414 [Funding:](#)

6  
7 415 No specific funding was received from any bodies in the public, commercial or not-for-profit sectors  
8 416 to carry out the work described in this article.

9  
10 417 [Conflict of interest:](#)

11  
12 418 JKB receives unrestricted research funding from Boehringer Ingelheim. All other co-authors have  
13 419 declared no conflicts of interest.

14  
15 420 [Data availability statement](#)

16  
17 421 Data are available upon reasonable request.  
18  
19 422  
20  
21 423  
22  
23 424  
24  
25 425  
26  
27 426

28 427 *Table.1:* Differences in surface markers and cytokine signatures in classical and alternative  
29 428 (phenotypes) macrophages  
30  
31 429  
32  
33 430  
34  
35 431

Macrophage phenotype	Surface markers	Cytokine signature
Classical	CD86, CD68, CD80, MHC-II, TLR-2, TLR-4, IL-1R	IL-1 $\beta$ , IL-6, IL-12, IL-18, IL-23, TNF- $\alpha$
Alternative (M2a)	CD206, CD36, CD163, IL-1R	IL-10, TGF- $\beta$
Alternative (M2b)	CD86, MHC-II	TNF- $\alpha$ , IL-1 $\beta$ , IL-6, IL-10
Alternative (M2c)	CD206, TLR-1, TLR-8, CD163	IL-10, TGF- $\beta$
Alternative (M2d)	CD206, CD204, CD163	IL-10, VEGF

36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48 432  
49  
50 433  
51  
52 434  
53  
54 435  
55  
56 436

57 437 *Table.2:* SSc-investigated targeted therapies and suggested novel monocyte/macrophage targeted  
58 438 therapies in the treatment of systemic sclerosis and systemic sclerosis-associated interstitial lung  
59  
60 439 disease

1  
2  
3 440  
4  
5 441  
6  
7 442  
8  
9 443  
10  
11 444

Therapeutic	Target (action)
AHSCT	resetting myeloid progenitor cells including monocytes
Tocilizumab	Anti-IL-6 receptor $\alpha$ -subunit (attenuates monocyte downstream effects)
Rituximab	Anti-CD20 "B cells depletion" (attenuates downstream macrophage polarization)
Romilkimab	Anti- IL-4 and IL-13 cytokines (blocks alternative activation)
Nintedanib	Multi-tyrosine kinase inhibitor (disturbs classical and alternative activations)
Pirfenidone	Blocks alternative activation
Ruxolitinib	JAK 1/2 inhibitor (proposed to attenuate alternative activation)
GSK2330811	Anti-oncostatin M protein (attenuates monocytes downstream effects)
Pegylated adenosine deaminase	adenosine molecules (blocks alternative polarization)
anti-CD64-dPBD	proliferating monocyte progenitors
anti-CCR2 siRNA nanotherapy	inflammatory monocytes migration
Statin-encapsulated HDL nanobiologic	inflammatory monocytes systemically and inflammatory macrophages locally
RP-182 and RP-832c	CD206+ cells "alternatively-activated macrophages"

445  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

446




Figure 1: Macrophage polarization.

Monocytes are attracted to injury sites (e.g., infected tissue) by chemoattractants. Thereafter, macrophages are activated dependent on the cytokines/stimuli in the milieu to M1 (classic activation) or M2 (alternative activation). Polarization state of macrophages is rather a dynamic process where macrophages can shift along a polarization spectrum. Activated macrophages express a specific set of (surface) markers and release particular collection of cytokines according to the activation pattern which in turn affects the milieu. CD: Cluster of Differentiation, IL: Interleukin, TGF- $\beta$ : Transforming Growth Factor-beta, TLR: Toll-Like Receptor, IFN: Interferon, MHC II: Major Histocompatibility Complex class II, VEGF: Vascular Endothelial Growth Factor, TNF- $\alpha$ : Tumor Necrosis Factor-alpha, LPS: Lipopolysaccharide, GM-CSF: Granulocyte-Macrophage Colony-Stimulating Factor. Created with BioRender.com



463 References:

- 464 1. Herrick AL. Systemic sclerosis : clinical features and management Key points. *Medicine*  
465 [Internet]. 2018;46(2):131–9. Available from: <https://doi.org/10.1016/j.mpmed.2017.11.007>
- 466 2. Pavec L, Girgis RE, Lechtzin N, Mathai SC, Launay D, Hummers LK, et al. Systemic Sclerosis –  
467 Related Pulmonary Hypertension Associated With Interstitial Lung Disease. *Arthritis and*  
468 *Rheumatism*. 2011;63(8):2456–64.
- 469 3. McFarlane IM, Bhamr MS, Kreps A, Iqbal S, Al-Ani F, Saladini-Aponte C, et al. Gastrointestinal  
470 Manifestations of Systemic Sclerosis. *Rheumatology (Sunnyvale)*. 2018;8(1):1–35.
- 471 4. Chrabaszc M, Małyszko J, Sikora M, Stochmal A, Rudnicka L. Renal Involvement in Systemic  
472 Sclerosis : An Update. *Kidney & Blood Pressure Research*. 2020;45:532–48.
- 473 5. Lambova S. Cardiac manifestations in systemic sclerosis. *World Journal of Cardiology*.  
474 2014;6(9):993–1005.
- 475 6. Denton CP, Khanna D. Seminar Systemic sclerosis. *The Lancet* [Internet].  
476 2017;390(10103):1685–99. Available from: [http://dx.doi.org/10.1016/S0140-6736\(17\)30933-9](http://dx.doi.org/10.1016/S0140-6736(17)30933-9)  
477 9
- 478 7. Gester F, Seny D de, Bonhomme O, Moermans C, Struman I, Louis R, et al. Biomarkers in  
479 systemic sclerosis-associated interstitial lung disease : review of the literature.  
480 *Rheumatology*. 2019;58(July):1534–46.
- 481 8. Diab S, Dostrovsky N, Hudson M, Tatibouet S, Fritzier MJ, Baron M, et al. Systemic Sclerosis  
482 Sine Scleroderma : A Multicenter Study of 1417 Subjects. *The Journal of Rheumatology*.  
483 2009;2179–85.
- 484 9. Distler O, Assassi S, Cottin V, Cutolo M, Danoff SK, Denton CP, et al. Predictors of progression  
485 in systemic sclerosis patients with interstitial lung disease. *European Respiratory Journal*  
486 [Internet]. 2020;55. Available from: <http://dx.doi.org/10.1183/13993003.02026-2019>
- 487 10. Elizabeth R. Volkman, Aryeh Fischer. Update on Morbidity and Mortality in Systemic  
488 Sclerosis-Related Interstitial Lung Disease. *Journal of Scleroderma and Related Disorders*.  
489 2021;6(1):11–20.
- 490 11. Cappelli S, Randone SB, Camiciottoli G, Paulis A de, Guiducci S, Matucci-cerinic M. Interstitial  
491 lung disease in systemic sclerosis : where do we stand ? *European Respiratory Review*  
492 [Internet]. 2015;24(137):411–9. Available from:  
493 <http://dx.doi.org/10.1183/16000617.00002915>
- 494 12. Toledo M D, Pioli A P. Macrophages in Systemic Sclerosis: Novel Insights and Therapeutic  
495 Implications. *Current Rheumatological Reports*. 2020;21(7).
- 496 13. Hirayama D, Iida T. The Phagocytic Function of Macrophage-Enforcing Innate Immunity and  
497 Tissue Homeostasis. *International Journal of Molecular Sciences*. 2018;19(92).
- 498 14. Ghobrial RM, Kloc M, Jacek L. Macrophage functions in wound healing. *Journal of Tissue*  
499 *Engineering and Regenerative Medicine*. 2019;9(November):99–109.
- 500 15. Cassetta L, Cassol E, Poli G. Macrophage Polarization in Health and Disease. *The Scientific*  
501 *World Journal*. 2011;11:2391–402.

- 1  
2  
3 502 16. Laria A, Mazzocchi D, Scarpellini M. The macrophages in rheumatic diseases. *Journal*  
4 503 *Inflammation Research*. 2016;9:1–11.  
5  
6 504 17. U. Sack, P. Stichl GG. Distribution of macrophages in rheumatoid synovial membrane and its  
7 505 association with basic activity. *Rheumatology International*. 1993;13:181–6.  
8  
9 506 18. Wildenberg ME, Bootsma H, Vissink A, Rooijen N van, Merwe JP van de, Drexhage HA, et al.  
10 507 Increased frequency of CD16 + monocytes and the presence of activated dendritic cells in  
11 508 salivary glands in primary Sjogren syndrome. *Annals of Rheumatic Diseases*. 2009;68:420–6.  
12  
13 509 19. Zhou D, Chen Y ting, Chen F, Gallup M, Vijmasi T, Bahrami AF, et al. Critical Involvement of  
14 510 Macrophage Infiltration in the. *The American Journal of Pathology* [Internet].  
15 511 2012;181(3):753–60. Available from: <http://dx.doi.org/10.1016/j.ajpath.2012.05.014>  
16  
17 512 20. Mathew J, Hines JE, Toole K, Johnson SJ, Burt OFWJAD. Quantitative analysis of macrophages  
18 513 and perisinusoidal cells in primary biliary cirrhosis. *Histopathology*. 1994;25:65–70.  
19  
20 514 21. Orme J, Mohan C. Macrophage subpopulations in systemic lupus erythematosus. *Discov Med*.  
21 515 2012 Feb;13(69):151–8.  
22  
23 516 22. Kuhl AA, Erben U, Kredel LI, Siegmund B. Diversity of intestinal Macrophages in inflammatory  
24 517 Bowel Diseases. *Frontiers in Immunology*. 2015;6(December):1–7.  
25  
26 518 23. Ishikawa O, Ishikawa H. Macrophage infiltration in the skin of patients with systemic sclerosis.  
27 519 *J Rheumatol*. 1992 Aug;19(8):1202–6.  
28  
29 520 24. López-cacho JM, Gallardo S, Posada M, Aguerri M, Calzada D, Mayayo T, et al. Association of  
30 521 Immunological Cell Profiles with Specific Clinical Phenotypes of Scleroderma Disease.  
31 522 *Biomedical Research International*. 2014;2014.  
32  
33 523 25. Ma W tao, Gao F, Gu K, Chen D kun. The Role of Monocytes and Macrophages in  
34 524 Autoimmune Diseases : A Comprehensive Review. *Frontiers in Immunology*. 2019;10(May):1–  
35 525 24.  
36  
37 526 26. Ma W tao, Chang C, Gershwin ME, Lian Z xiong. Development of autoantibodies precedes  
38 527 clinical manifestations of autoimmune diseases : A comprehensive review. *Journal of*  
39 528 *Autoimmunity* [Internet]. 2017;83:95–112. Available from:  
40 529 <http://dx.doi.org/10.1016/j.jaut.2017.07.003>  
41  
42 530 27. Chávez-galán L, Olleros ML, Vesin D, Garcia I, Harris RA. Much more than M1 and M2  
43 531 macrophages , there are also CD169 + and TCR + macrophages. *Frontiers in Immunology*.  
44 532 2015;6(May):1–15.  
45  
46 533 28. Yang J, Zhang L, Yu C, Yang X feng, Wang H. Monocyte and macrophage differentiation :  
47 534 circulation inflammatory monocyte as biomarker for inflammatory diseases. *Biomarker*  
48 535 *Research*. 2014;2(1):1–9.  
49  
50 536 29. Ricardo SD, Goor H van, Eddy AA, Ricardo SD, Goor H van, Eddy AA. Macrophage diversity in  
51 537 renal injury and repair *Science in medicine Macrophage diversity in renal injury and repair*.  
52 538 *The Journal of Clinical Investigation*. 2008;118(11):3522–30.  
53  
54 539 30. Viola A, Munari F, Sánchez-rodríguez R, Scolaro T, Castegna A. The Metabolic Signature of  
55 540 Macrophage Responses. *Frontiers in Immunology*. 2019;10(July):1–16.  
56  
57  
58  
59  
60

- 1  
2  
3 541 31. Bertani FR, Mozetic P, Fioramonti M, Iuli M, Ribelli G, Pantano F, et al. Classification of M1 /  
4 542 M2-polarized human macrophages by label-free hyperspectral reflectance confocal  
5 543 microscopy and multivariate analysis. *Nature Scientific Reports*. 2017;7(February):1–9.
- 7 544 32. Hu X, Ivashkiv LB. Cross-regulation of Signaling Pathways by Interferon- $\gamma$ : Implications for  
8 545 Immune Responses and Autoimmune Diseases. *Immunity* [Internet]. 2009;31(4):539–50.  
9 546 Available from: <https://www.sciencedirect.com/science/article/pii/S1074761309004063>
- 11 547 33. Pace JL, Russell SW, Schreibert RD, Altmant A, Katzt DH. Macrophage activation : Priming  
12 548 activity from a T-cell hybridoma is attributable to interferon- $\gamma$  Immunology : Proc Natl Acad  
13 549 Sci U S A. 1983;80(June):3782–6.
- 16 550 34. Grassin-delyle S, Salvator H. The Role of Toll-Like Receptors in the Production of Cytokines by  
17 551 Human Lung Macrophages. *Journal of Innate Immunity*. 2020;12(December):63–73.
- 19 552 35. Hamilton TA, Zhao C, Jr PGP, Datta S. Myeloid colony-stimulating factors as regulators of  
20 553 macrophage polarization. *frontiers in Immunology*. 2014;5(November):1–6.
- 22 554 36. Martinez FO, Helming L, Gordon S. Alternative Activation of Macrophages : An Immunologic  
23 555 Functional Perspective. *Annual Review of Immunology*. 2009;27:451–83.
- 25 556 37. Wager CML, Jr FLW. Classical versus alternative macrophage activation : the Ying and the  
26 557 Yang in host defense against pulmonary fungal infections. *Mucosal Immunology*.  
27 558 2014;7(5):1023–35.
- 29 559 38. Ferrante CJ, Leibovich SJ. Regulation of Macrophage Polarization and Wound Healing.  
30 560 *Advances in Wound Care*. 2012;1(1):10–6.
- 32 561 39. Stein BM, Keshav S, Harris N, Gordon S. Interleukin 4 Potently Enhances Murine Macrophage  
33 562 Mannose Receptor Activity: A Marker of Alternative Immunologic Macrophage Activation.  
34 563 *The Journal of External Medicine*. 1992;176(July):287–92.
- 37 564 40. Wang LX, Zhang SX, Wu HJ, Rong XL, Guo J. M2b macrophage polarization and its roles in  
38 565 diseases. *J Leukoc Biol*. 2019 Aug;106(2):345–58.
- 40 566 41. Shapouri-moghaddam Abbas, Saeed Mohammadian, Vazini Hossein, Taghadosi Mahdi,  
41 567 Esmaeili Seyed-Alireza, Mardani Fatemeh, et al. Macrophage plasticity , polarization , and  
42 568 function in health and disease. *Journal of Cellular Physiology*. 2018;9:6425–40.
- 44 569 42. Ferrante CJ, Pinhal-enfield G, Elson G, Cronstein BN, Hasko G, Outram S, et al. The Adenosine-  
45 570 Dependent Angiogenic Switch of Macrophages to an M2-Like Phenotype is Independent of  
46 571 Interleukin-4 Receptor Alpha (IL-4R $\alpha$ ) Signaling. *Inflammation*. 2014;36(4):921–31.
- 48 572 43. Mosser DM, Edwards JP. Exploring the full spectrum of macrophage activation. *Nature*  
49 573 *Publishing Group* [Internet]. 2008;8(12):958–69. Available from:  
50 574 <http://dx.doi.org/10.1038/nri2448>
- 53 575 44. Martinez FO, Gordon S. The M1 and M2 paradigm of macrophage activation : time for  
54 576 reassessment. *F1000Prime Reports*. 2014;13(March):1–13.
- 56 577 45. Higashi-Kuwata N, Jinnin M, Makino T, Fukushima S, Inoue Y, Muchemwa FC, et al.  
57 578 Characterization of monocyte/macrophage subsets in the skin and peripheral blood derived  
58 579 from patients with systemic sclerosis. *Arthritis Res Ther*. 2010;12(4):R128.

- 1  
2  
3 580 46. Mathai SK, Gulati M, Peng X, Russell T, Shaw A, Murray LA, et al. Circulating Monocytes from  
4 581 Systemic Sclerosis Patients with Interstitial Lung Disease Show an Enhanced  
5 582 Profibrotic Phenotype. *Laboratory Investigation*. 2013;90(6):812–23.
- 7 583 47. Moreno-Moral A, Bagnati M, Koturan S, Ko JH, Fonseca C, Harmston N, et al. Changes in  
8 584 macrophage transcriptome associate with systemic sclerosis and mediate GSDMA  
9 585 contribution to disease risk. *Annals of the Rheumatic Diseases*. 2018 Apr 1;77(4):596–601.
- 11 586 48. Trombetta AC, Soldano S, Contini P, Tomatis V, Ruaro B, Paolino S, et al. A circulating cell  
12 587 population showing both M1 and M2 monocyte / macrophage surface markers characterizes  
13 588 systemic sclerosis patients with lung involvement. *Respiratory Research*. 2018;19(186):1–12.
- 16 589 49. Soldano S, Trombetta A, Contini P et al. Increase in circulating cells coexpressing M1 and M2  
17 590 macrophage surface markers in patients with systemic sclerosis. *Annals of Rheumatic  
18 591 Diseases*. 2018;77(12):1842–5.
- 20 592 50. Valenzi E, Bulik M, Tabib T, Morse C, Sembrat J, Trejo Bittar H, et al. Single-cell analysis  
21 593 reveals fibroblast heterogeneity and myofibroblasts in systemic sclerosis-associated  
22 594 interstitial lung disease. *Annals of the Rheumatic Diseases [Internet]*. 2019 Oct 1;78(10):1379.  
23 595 Available from: <http://ard.bmj.com/content/78/10/1379.abstract>
- 26 596 51. Morse C, Tabib T, Sembrat J, Buschur KL, Bittar HT, Valenzi E, et al. Proliferating SPP1/MERTK-  
27 597 expressing macrophages in idiopathic pulmonary fibrosis. *European Respiratory Journal  
28 598 [Internet]*. 2019 Aug 1;54(2):1802441. Available from:  
29 599 <http://erj.ersjournals.com/content/54/2/1802441.abstract>
- 31 600 52. Xue D, Tabib T, Morse C, Yang Y, Domsic RT, Khanna D, et al. Expansion of Fcγ Receptor IIIa–  
32 601 Positive Macrophages, Ficolin 1–Positive Monocyte-Derived Dendritic Cells, and Plasmacytoid  
33 602 Dendritic Cells Associated With Severe Skin Disease in Systemic Sclerosis. *Arthritis &  
34 603 Rheumatology [Internet]*. 2022 Feb 1;74(2):329–41. Available from:  
35 604 <https://doi.org/10.1002/art.41813>
- 38 605 53. Pillai S. T and B lymphocytes in fibrosis and systemic sclerosis. *Current Opinion*.  
39 606 2019;31(6):576–81.
- 41 607 54. Perelas A, Silver RM, Arrossi A v, Highland KB. Series Systemic sclerosis 2 Systemic sclerosis-  
42 608 associated interstitial lung disease. *The Lancet Respiratory [Internet]*. 2020;8(3):304–20.  
43 609 Available from: [http://dx.doi.org/10.1016/S2213-2600\(19\)30480-1](http://dx.doi.org/10.1016/S2213-2600(19)30480-1)
- 45 610 55. Bhandari R, Ball MS, Martyanov V, Popovich D, Schaafsma E, Han S, et al. Profibrotic  
46 611 Activation of Human Macrophages in Systemic Sclerosis. *Arthritis & rheumatology  
47 612 (Hoboken, NJ) [Internet]*. 2020;72(7):1160–9. Available from:  
48 613 <http://europepmc.org/abstract/MED/32134204>
- 51 614 56. Monika L, Elizabeth C, M KH, Pardis P, Brian W, Stellar B, et al. Cadherin-11–mediated  
52 615 adhesion of macrophages to myofibroblasts establishes a profibrotic niche of active TGF-β.  
53 616 *Science Signaling [Internet]*. 2019 Jan 15;12(564):eaao3469. Available from:  
54 617 <https://doi.org/10.1126/scisignal.aao3469>
- 56 618 57. Pakshir P, Alizadehgiashi M, Wong B, Coelho NM, Chen X, Gong Z, et al. Dynamic fibroblast  
57 619 contractions attract remote macrophages in fibrillar collagen matrix. *Nature  
58 620 Communications*. 2019 Dec 1;10(1).

- 1  
2  
3 621 58. Doridot L, Jeljeli M, Chêne C, Batteux F. Implication of oxidative stress in the pathogenesis of  
4 622 systemic sclerosis via inflammation, autoimmunity and fibrosis. Vol. 25, Redox Biology.  
5 623 Elsevier B.V.; 2019.
- 7 624 59. Kavian N, Mehlal S, Jeljeli M, Saidu NEB, Nicco C, Cerles O, et al. The Nrf2-antioxidant  
8 625 response element signaling pathway controls fibrosis and autoimmunity in scleroderma.  
9 626 Frontiers in Immunology. 2018 Aug 16;9(AUG).
- 11 627 60. Crestani B, Seta N, de Bandt M, Soler P, Rolland C, Dehoux M, et al. Interleukin 6 Secretion by  
12 628 Monocytes and Alveolar Macrophages in Systemic Sclerosis with Lung Involvement.
- 15 629 61. Khanna D, Lin CJF, Furst DE, Goldin J, Kim G, Kuwana M, et al. Articles Tocilizumab in systemic  
16 630 sclerosis : a randomised , double- blind , placebo-controlled , phase 3 trial. The Lancet  
17 631 Respiratory Medicine. 2020;8(October):963–74.
- 19 632 62. Rudnik M, Rolski F, Jordan S, Mertelj T, Stellato M, Distler O, et al. Regulation of Monocyte  
20 633 Adhesion and Type I Interferon Signaling by CD52 in Patients With Systemic Sclerosis. Arthritis  
21 634 and Rheumatology. 2021;73(9):1720–30.
- 23 635 63. Ebata S, Yoshizaki A, Oba K, Kashiwabara K, Ueda K, Uemura Y, et al. Safety and efficacy of  
24 636 rituximab in systemic sclerosis (DESIREs): a double-blind, investigator-initiated, randomised,  
25 637 placebo-controlled trial. The Lancet Rheumatology. 2021 Jul 1;3(7):e489–97.
- 28 638 64. Boonstra M, Meijs J, Dorjée AL, Marsan NA, Schouffoer A, Ninaber MK, et al. Rituximab in  
29 639 early systemic sclerosis. RMD Open. 2017 Jul 1;3(2).
- 31 640 65. Sircar G, Goswami RP, Sircar D, Ghosh A, Ghosh P. Intravenous cyclophosphamide vs  
32 641 rituximab for the treatment of early diffuse scleroderma lung disease: Open label,  
33 642 randomized, controlled trial. Rheumatology (United Kingdom). 2018 Dec 1;57(12):2106–13.
- 35 643 66. Rudnik M, Rolski F, Jordan S, Mertelj T, Stellato M, Distler O, et al. Regulation of Monocyte  
36 644 Adhesion and Type I Interferon Signaling by CD52 in Patients With Systemic Sclerosis. Arthritis  
37 645 and Rheumatology. 2021;73(9):1720–30.
- 39 646 67. Allanore Y, Wung P, Soubrane C, Esperet C, Marrache F, Bejuit R, et al. A randomised, double-  
40 647 blind, placebo-controlled, 24-week, phase II, proof-of-concept study of romilkimab  
41 648 (SAR156597) in early diffuse cutaneous systemic sclerosis. Annals of the Rheumatic Diseases  
42 649 [Internet]. 2020 Dec 1;79(12):1600 LP – 1607. Available from:  
43 650 <http://ard.bmj.com/content/79/12/1600.abstract>
- 46 651 68. Bellamri N, Morzadec C, Joannes A, Lecureur V, Wollin L, Jouneau S, et al. Alteration of  
47 652 human macrophage phenotypes by the anti-fibrotic drug nintedanib. International  
48 653 Immunopharmacology. 2019 Jul 1;72:112–23.
- 50 654 69. Bellamri N, Morzadec C, Lecureur V, Joannes A, Wollin L, Jouneau S, et al. Effects of  
51 655 Nintedanib on the M1 and M2a polarization of human macrophages. European Respiratory  
52 656 Journal [Internet]. 2018;52(suppl 62). Available from:  
53 657 [https://erj.ersjournals.com/content/52/suppl\\_62/PA5250](https://erj.ersjournals.com/content/52/suppl_62/PA5250)
- 56 658 70. Azuma A, Fischer A, Mayes MD, Stowasser S, Tetzlaff K, Kuwana M, et al. Nintedanib for  
57 659 Systemic Sclerosis– Associated Interstitial Lung Disease. NEJM. 2019;380:2518–28.
- 59 660 71. Bruni T, Varone F. The adoption of nintedanib in systemic sclerosis : the SENSCIS study.  
60 661 Breathe. 2020;16(2):1–4.

- 1  
2  
3 662 72. Campochiaro C, Allanore Y. An update on targeted therapies in systemic sclerosis based on a  
4 663 systematic review from the last 3 years. *Arthritis Research & Therapy*. 2021;5:1–14.
- 6 664 73. Rivera-ortega P, Hayton C, Blaikley J, Leonard C, Chaudhuri N. Nintedanib in the management  
7 665 of idiopathic pulmonary fibrosis : clinical trial evidence and real-world experience. 2018;1–13.
- 9 666 74. Toda M, Mizuguchi S, Minamiyama Y, Yamamotoooka H, Aota T, Kubo S, et al. Pirfenidone  
10 667 suppresses polarization to M2 phenotype macrophages and the fibrogenic activity of rat lung  
11 668 fibroblasts. *J Clin Biochem Nutr*. 2018;63(1):58–65.
- 14 669 75. Khanna D, Albera C, Fischer A, Seibold J, Raghu G, Khalidi N, et al. Safety and tolerability of  
15 670 pirfenidone (PFD) in patients with systemic sclerosis-associated interstitial lung disease (SSc-  
16 671 ILD)—The LOTUSS study. *European Respiratory Journal [Internet]*. 2015 Sep 1;46(suppl  
17 672 59):OA4489. Available from:  
18 673 [http://erj.ersjournals.com/content/46/suppl\\_59/OA4489.abstract](http://erj.ersjournals.com/content/46/suppl_59/OA4489.abstract)
- 20 674 76. Acharya N, Sharma SK, Mishra D, Dhooria S, Dhir V, Jain S. Efficacy and safety of pirfenidone  
21 675 in systemic sclerosis-related interstitial lung disease—a randomised controlled trial.  
22 676 *Rheumatology International [Internet]*. 2020;40(5):703–10. Available from:  
23 677 <https://doi.org/10.1007/s00296-020-04565-w>
- 26 678 77. van den Hombergh WMT, Kersten BE, Knaapen-Hans HKA, Thurlings RM, van der Kraan PM,  
27 679 van den Hoogen FHJ, et al. Hit hard and early: Analysing the effects of high-dose  
28 680 methylprednisolone on nailfold capillary changes and biomarkers in very early systemic  
29 681 sclerosis: Study protocol for a 12-week randomised controlled trial. *Trials*. 2018 Aug 22;19(1).
- 31 682 78. Ehrchen JM, Roth J, Barczyk-Kahlert K. More than suppression: Glucocorticoid action on  
32 683 monocytes and macrophages. Vol. 10, *Frontiers in Immunology*. Frontiers Media S.A.; 2019.
- 34 684 79. Xie Y, Tolmeijer S, Oskam JM, Tonkens T, Meijer AH, Schaaf MJM. Glucocorticoids inhibit  
35 685 macrophage differentiation towards a pro-inflammatory phenotype upon wounding without  
36 686 affecting their migration. *DMM Disease Models and Mechanisms*. 2019 May 1;12(5).
- 39 687 80. Spierings J, van Rhenen A, Welsing PMW, Marijnissen ACA, de Langhe E, del Papa N, et al. A  
40 688 randomised, open-label trial to assess the optimal treatment strategy in early diffuse  
41 689 cutaneous systemic sclerosis: The UPSIDE study protocol. *BMJ Open*. 2021 Mar 18;11(3).
- 43 690 81. Mukherjee R, Kanti Barman P, Kumar Thatoi P, Tripathy R, Kumar Das B, Ravindran B. Non-  
44 691 Classical monocytes display inflammatory features: Validation in Sepsis and Systemic Lupus  
45 692 Erythematosus. *Scientific Reports*. 2015 Sep 11;5.
- 47 693 82. Servaas NH, Spierings J, Pandit A, van Laar JM. The role of innate immune cells in systemic  
48 694 sclerosis in the context of autologous hematopoietic stem cell transplantation. Vol. 201,  
49 695 *Clinical and Experimental Immunology*. Blackwell Publishing Ltd; 2020. p. 34–9.
- 51 696 83. Stawski L, Trojanowska M. Oncostatin M and its role in fibrosis. Vol. 60, *Connective Tissue  
52 697 Research*. Taylor and Francis Ltd; 2019. p. 40–9.
- 54 698 84. Richards CD, Botelho F. Oncostatin M in the regulation of connective tissue cells and  
55 699 macrophages in pulmonary disease. Vol. 7, *Biomedicines*. MDPI AG; 2019.
- 57 700 85. Reid J, Zamuner S, Edwards K, Rumley SA, Nevin K, Feeney M, et al. In vivo affinity and target  
58 701 engagement in skin and blood in a first-time-in-human study of an anti-oncostatin M  
59 702 monoclonal antibody. *British Journal of Clinical Pharmacology*. 2018 Oct 1;84(10):2280–91.

- 1  
2  
3 703 86. Zhang Y, Zhu H, Layritz F, Luo H, Wohlfahrt T, Chen CW, et al. Recombinant Adenosine  
4 704 Deaminase Ameliorates Inflammation, Vascular Disease, and Fibrosis in Preclinical Models of  
5 705 Systemic Sclerosis. *Arthritis & Rheumatology* [Internet]. 2020 Aug 1;72(8):1385–95. Available  
6 706 from: <https://doi.org/10.1002/art.41259>
- 7  
8  
9 707 87. Izumi Y, Kanayama M, Shen Z, Kai M. An Antibody-Drug Conjugate That Selectively Targets  
10 708 Human Monocyte Progenitors for Anti-Cancer Therapy. 2021;12(February):1–15.
- 11  
12 709 88. Leuschner F, Dutta P, Gorbatov R, Novobrantseva TI, Donahoe JS, Courties G, et al.  
13 710 Therapeutic siRNA silencing in inflammatory monocytes in mice. *Nature Biotechnology*. 2011  
14 711 Nov;29(11):1005–10.
- 15  
16 712 89. Mulder WJM, Ochando J, Joosten LAB, Fayad ZA, Netea MG. Therapeutic targeting of trained  
17 713 immunity. *Nature Reviews Drug Discovery*. 2019 Jul 1;18(7):553–66.
- 18  
19 714 90. Duivenvoorden R, Tang J, Cormode DP, Mieszawska AJ, Izquierdo-Garcia D, Ozcan C, et al. A  
20 715 statin-loaded reconstituted high-density lipoprotein nanoparticle inhibits atherosclerotic  
21 716 plaque inflammation. *Nature Communications*. 2014;5.
- 22  
23 717 91. Ahmed Abdi B, Lopez H, Martin G, Garvin C, Jaynes J, Stanway J, et al. Novel Therapeutic  
24 718 Peptides Which Target CD206 Inhibit Macrophage Dependent Fibroblast Activation in  
25 719 Scleroderma. [https://acrabstracts.org/abstract/novel-therapeutic-peptides-which-target-](https://acrabstracts.org/abstract/novel-therapeutic-peptides-which-target-cd206-inhibit-macrophage-dependent-fibroblast-activation-in-scleroderma/)  
26 720 [cd206-inhibit-macrophage-dependent-fibroblast-activation-in-scleroderma/](https://acrabstracts.org/abstract/novel-therapeutic-peptides-which-target-cd206-inhibit-macrophage-dependent-fibroblast-activation-in-scleroderma/). Accessed March  
27 721 30, 2022. 2018.
- 28  
29 722 92. Ghebremedhin A, Salam A bin, Adu-Addai B, Noonan S, Stratton R, Ahmed MSU, et al. A  
30 723 Novel CD206 Targeting Peptide Inhibits Bleomycin Induced Pulmonary Fibrosis in Mice.  
31 724 bioRxiv [Internet]. 2020 Jul 29; Available from:  
32 725 <http://www.ncbi.nlm.nih.gov/pubmed/32766584>

726