

CODE-EHR best practice framework for the use of structured electronic healthcare records in clinical research

Dipak Kotecha^{1–3}, Folkert W Asselbergs^{3–5}, Stephan Achenbach⁶, Stefan D Anker⁷, Dan Atar^{8,9}, Colin Baigent^{10,11}, Amitava Banerjee^{5,12}, Birgit Beger¹³, Gunnar Brobert¹⁴, Barbara Casadei¹⁵, Cinzia Ceccarelli¹⁶, Martin R Cowie^{17,18}, Filippo Crea^{19,20}, Maureen Cronin²¹, Spiros Denaxas^{5,22,23}, Andrea Derix²⁴, Donna Fitzsimons²⁵, Martin Fredriksson²⁶, Chris P Gale^{27–29}, Georgios V Gkoutos^{2,30}, Wim Goettsch^{31,32}, Harry Hemingway⁵, Martin Ingvar^{33,34}, Adrian Jonas³⁵, Robert Kazmierski³⁶, Susanne Løgstrup¹³, R Thomas Lumbers^{5,37}, Thomas F Lüscher^{38–40}, Paul McGreavy⁴¹, Ileana L Piña^{42,43}, Lothar Roessig²⁴, Carl Steinbeisser^{24,44}, Mats Sundgren⁴⁵, Benoît Tyl⁴⁶, Ghislaine van Thiel⁴⁷, Kees van Bochove⁴⁸, Panos E Vardas^{49,50}, Tiago Villanueva⁵¹, Marilena Vrana¹³, Wim Weber⁵¹, Franz Weidinger⁵², Stephan Windecker⁵³, Angela Wood⁵⁴, and Diederick E Grobbee⁵⁵ on behalf of the Innovative Medicines Initiative BigData@Heart Consortium, European Society of Cardiology, CODE-EHR international consensus group

¹Institute of Cardiovascular Sciences, University of Birmingham, Medical School, Birmingham, UK; ²University Hospitals Birmingham NHS Foundation Trust and Health Data Research UK Midlands, Birmingham, UK; ³Department of Cardiology, Division of Heart and Lungs, University Medical Centre Utrecht, University of Utrecht, Utrecht, Netherlands; ⁴Institute of Cardiovascular Science, Faculty of Population Health Sciences, University College London, London, UK; ⁵Health Data Research UK and Institute of Health Informatics, University College London, London, UK; ⁶Friedrich-Alexander-Universität Erlangen-Nürnberg (FAU), Erlangen, Germany; ⁷Department of Cardiology and Berlin Institute of Health Centre for Regenerative Therapies, German Centre for Cardiovascular Research (DZHK) partner site Berlin; Charité Universitätsmedizin Berlin, Germany; ⁸Department of Cardiology, Oslo University Hospital, Ulleval, Oslo, Norway; ⁹University of Oslo, Institute of Clinical Medicine, Oslo, Norway; ¹⁰MRC Population Health Research Unit, Nuffield Department of Population Health, Oxford, UK; ¹¹Clinical Trial Service Unit and Epidemiological Studies Unit, University of Oxford, Oxford, UK; ¹²University College London Hospitals NHS Trust, London, UK; ¹³European Heart Network, Brussels, Belgium; ¹⁴Bayer AB, Stockholm, Sweden; ¹⁵Division of Cardiovascular Medicine, John Radcliffe Hospital, University of Oxford NIHR Oxford Biomedical Research Centre, Oxford, UK; ¹⁶European Society of Cardiology, Sophia Antipolis, France; ¹⁷Royal Brompton Hospital, Division of Guy's St Thomas' NHS Foundation Trust, London, UK; ¹⁸School of Cardiovascular Medicine Sciences, King's College London, London, UK; ¹⁹Department of Cardiovascular and Pulmonary Sciences, Catholic University of the Sacred Heart, Rome, Italy; ²⁰European Heart Journal, Oxford University Press, University of Oxford, Oxford, UK; ²¹Vifor Pharma, Glattbrugg, Switzerland and Ava AG, Zurich, Switzerland; ²²Alan Turing Institute, London, UK; ²³British Heart Foundation Data Science Centre, London, UK; ²⁴Bayer AG, Leverkusen, Germany; ²⁵School of Nursing and Midwifery, Queen's University Belfast, Northern Ireland; ²⁶Late Clinical Development, Cardiovascular, Renal and Metabolism (CVRM), Biopharmaceuticals RD, AstraZeneca, Gothenburg, Sweden; ²⁷Leeds Institute of Cardiovascular and Metabolic Medicine and Leeds Institute for Data Analytics, University of Leeds, Leeds, UK; ²⁸Leeds Institute for Data Analytics, University of Leeds, Leeds, UK; ²⁹Department of Cardiology, Leeds Teaching Hospitals NHS Trust, Leeds, UK; ³⁰College of Medical and Dental Sciences, Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, UK; ³¹National Health Care Institute (ZIN), Diemen, Netherlands; ³²Division of Pharmacoepidemiology and Clinical Pharmacology, Utrecht Institute for Pharmaceutical Sciences, Utrecht University, Utrecht, Netherlands; ³³Department of Clinical Neuroscience, Karolinska Institutet, Solna, Sweden; ³⁴Department of Neuroradiology, Karolinska University Hospital Stockholm, Stockholm, Sweden; ³⁵Data and Analytics Group, National Institute for Health and Care Excellence, London, UK; ³⁶Office of Cardiovascular Devices, US Food and Drug Administration, Silver Spring, MD, USA; ³⁷Barts Health NHS Trust and University College London Hospitals NHS Trust; ³⁸Centre for Molecular Cardiology, University of Zurich, Zurich, Switzerland; ³⁹Research, Education & Development, Royal Brompton and Harefield Hospitals, London, UK; ⁴⁰Faculty of Medicine, Imperial College London, London, UK; ⁴¹European Society of Cardiology Patient Forum, European Society of Cardiology, Brussels, Belgium; ⁴²Central Michigan University College of Medicine, Midland, MI, USA; ⁴³Centre for Devices and Radiological Health, US Food and Drug Administration, Silver Spring, MD, USA; ⁴⁴Steinbeisser Project Management, Munich, Germany; ⁴⁵Data Science AI, Biopharmaceuticals RD, AstraZeneca, Gothenburg, Sweden; ⁴⁶Centre for Therapeutic Innovation, Cardiovascular and Metabolic Disease, Institut de Recherches Internationales Servier, Suresnes, France; ⁴⁷Julius Centre for Health Sciences and Primary Care, University Medical Centre Utrecht, Utrecht University, Utrecht, Netherlands; ⁴⁸The Hyve, Utrecht, Netherlands; ⁴⁹Hygeia, Mitera, Hospitals Hellenic Health Group, Athens, Greece; ⁵⁰European Heart Agency, European Society of Cardiology, Brussels, Belgium; ⁵¹The BMJ, London, UK; ⁵²Rudolfstiftung Hospital, Vienna, Austria; ⁵³Department of Cardiology, Inselspital, University Hospital Bern, Bern, Switzerland; ⁵⁴Cardiovascular Epidemiology Unit, Department of Public Health and Primary Care, University of Cambridge, Cambridge, UK; and ⁵⁵Department of Epidemiology, University Medical Centre Utrecht, Division Julius Centrum, Utrecht, Netherlands

Accepted 21 June 2022

Correspondence: D Kotecha d.kotecha@bham.ac.uk

This article has been co-published with permission in The BMJ, the *Lancet Digital Health*, and the *European Heart Journal* © the Authors 2022.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited.

The articles are identical except for minor stylistic and spelling differences in keeping with each journal's style. When citing this article, a citation from any of the journals listed is appropriate.

Abstract

Big data is central to new developments in global clinical science aiming to improve the lives of patients. Technological advances have led to the routine use of structured electronic healthcare records with the potential to address key gaps in clinical evidence. The covid-19 pandemic has demonstrated the potential of big data and related analytics, but also important pitfalls. Verification, validation, and data privacy, as well as the social mandate to undertake research are key challenges. The European Society of Cardiology and the BigData@Heart consortium have brought together a range of international stakeholders, including patient representatives, clinicians, scientists, regulators, journal editors and industry. We propose the CODE-EHR Minimum Standards Framework as a means to improve the design of studies, enhance transparency and develop a roadmap towards more robust and effective utilisation of healthcare data for research purposes.

In the context of ageing populations and increasing multimorbidity in all disease areas,^{1–3} large scale, real world data provide an opportunity to better understand the epidemiology of rare and common conditions, and to improve prevention strategies and treatment stratification.⁴ Tailored management for individual patients has become even more essential to constrain healthcare costs and provide patient centred care that can improve a patient's quality of life and prognosis. Embedding controlled trials within the real world setting, either within registries or routine clinical practice, is now possible and could provide more generalisable results to the population at large.⁵

Health data science has undergone rapid development in the past decade, including the common adoption of electronic healthcare record (EHR) systems that condense clinical episodes into a set of coded, structured labels.⁶ However, concerns over quality, data privacy, transparency, and comparability of these systems have limited the use of the evidence generated with structured healthcare data. These issues have also restricted acceptance by regulators, reimbursement authorities, and guideline task forces. Despite the availability of numerous reporting standards, consensus has not been met on how to realise the Findable, Accessible, Interoperable, and Reusable (FAIR) principles⁷ in the context of structured healthcare data. Existing reporting checklists ask authors to indicate where in their paper particular design issues have been discussed. For example, STROBE (Strengthening the Reporting of Observational studies in Epidemiology) for observational studies,⁸ RECORD (Reporting of studies Conducted using Observational Routinely-collected Data) for routinely collected health data,⁹ and CONSORT-AI (Consolidated Standards of Reporting Trials-Artificial Intelligence) for artificial intelligence interventions.¹⁰ However, these checklists are often lengthy, no minimum standards are specified, and adherence does not relate to study quality or even the quality of transparency for that domain.¹¹ Although checklists can benefit research quality, they are often used for box ticking to facilitate journal publication. In a study of radiology journals, only 15% (120/821) of surveyed authors used the reporting guideline when designing their study.¹² With a proliferation of reporting checklists for every scenario, authors and readers of such reports are increasingly confused about the value of these checklists. As of 14 February 2022, 488 reporting checklists were registered with EQUATOR (Enhancing the QUALity and Transparency Of health Research) and 111 were in development.

In the case of observational and randomised clinical research using EHRs and other structured data, the source of data, its manipulation, and underpinning governance are of critical importance to

extrapolating results. Clarity is needed from a broad stakeholder perspective, providing a quality framework to enhance the design and application of clinical research that increasingly depends on these crucial new sources of data. This article reflects the joint work of a wide range of international stakeholders with a remit to improve the use of structured healthcare data. The programme was coordinated by the European Society of Cardiology, a non-profit organisation of healthcare professionals, and the BigData@Heart Consortium, a public-private partnership funded by the European Union Innovative Medicines Initiative. Our aim was to navigate opportunities and limitations, and to develop a framework for a broad audience of global stakeholders across all disease areas. The CODE-EHR framework seeks to realise the exciting opportunity that digitisation of health data affords to increase efficiency of healthcare systems, and improve the lives and wellbeing of patients.

Summary points

- Research using routinely collected structured healthcare data has the potential for major clinical impact but this requires a clear and transparent approach to describe data sources, linkage protocols, coding definitions, and validation of methods and results
- A social license and public mandate are essential components of big data research that can provide societal benefit, addressing the concerns of participants, and ensuring data privacy and integrity
- This paper describes the output of international stakeholder meetings for the use of structured healthcare data for research purposes, including patient representatives, clinicians, scientists, regulators, journal editors, and industry representatives
- The CODE-EHR checklist provides a minimum standards framework to enhance research design and enable more effective use and dissemination of routine healthcare data for clinical research

Stakeholder development of the CODE-EHR framework

A full range of stakeholders participated, including regulators (US Food and Drug Administration, European Medicines Agency), governmental agencies (European Commission, the UK National Institute for Health and Care Excellence, Innovative Medicines Initiative), leading medical journals (*The BMJ*, *European Heart Journal*, *The Lancet*, *The Lancet Digital Health*), and patient advocacy

groups (European Heart Network, ESC Patient Forum), in addition to representatives from the pharmaceutical industry, payers, leading academic institutions, and professional societies (see acknowledgments). Development of the CODE-EHR framework was centred on two stakeholder meetings (7 July 2020 and 26 October 2020), consisting of presentations from key opinion leaders, followed by breakout sessions and plenaries to formulate statements on key topic areas. An iterative process with virtual work was used to achieve consensus positions, with a further meeting on 10 March 2022 to finalise this report.

We aimed to develop pragmatic advice for the use of structured healthcare data within trials and observational studies that is not dependent on particular diseases, and that meets the expectations of stakeholders and the general public. Our objectives were to provide new direction to this increasingly important field in medicine, thereby enhancing the value of routinely collected data to improve future patient wellbeing. Detailed text on the current state-of-the-art for research using healthcare data, in addition to key challenges and limitations, was developed by the stakeholder group, supported by a writing committee. See appendix 1 in which we address the need for common standards to appraise the digital landscape (e.g. coding systems and the vital aspect of linkage), expand on current and future opportunities for the use of structured healthcare data, and show how a social license can lead to co-creation of research with a public health benefit. The key challenges and pathways for improvement are outlined in figure 1, which presents the process of structured healthcare data from initial notation to their potential use to enhance research and subsequently improve clinical practice.

The output of the stakeholder meetings and iterative discussions were condensed into four core central themes: technical process and data stewardship; data security and privacy; publications using structured healthcare data; and addressing the needs of regulators, reimbursement authorities, and clinical practice guidelines. Key statements and advisories from the consensus meetings are summarised in table 1.

Patient and public involvement

The CODE-EHR consensus approach has benefited from patient and public engagement throughout the development process, including representation from the European Society of Cardiology Patient Council and the European Heart Network, an alliance of foundations and associations supporting patients and representing patient interests. We describe a potential method for engagement of the public in future research that can constructively benefit research using big data (fig 2).

CODE-EHR reporting framework

The path from structured healthcare data to clinical research output is complex. To support further development in a transparent way, stakeholder delegates reached consensus of the need for a set of minimum standards that authors could use as a tool to enhance design, reporting, and research output. The CODE-EHR Minimum Standards Framework presented in table 2 allows authors to report on how structured healthcare data were used in their research study

(either in patient identification, disease phenotyping, or outcome derivation). Preferred standards indicate high level attainment of quality and can be used as a tool to improve the future trajectory of research. The checklist was created through an iterative process based on the stakeholder proposals and covers five key areas of enhanced transparency: how and why coding was performed; the process of constructing and linking datasets; clear definitions of both diseases and outcomes; the approach to analysis, including any computational methods; and demonstrating good data governance.


The framework aims to improve the quality of studies using structured healthcare data and to give confidence in their use for clinical decision making. See appendix 2 for a step-by-step approach to completion of the CODE-EHR reporting checklist, with relevant best practice examples. We also present a detailed description of the workflow that led to the checklist in appendix 3. Form versions of the checklist are provided in appendix 4 (word version) and appendix 5 (pdf version).

Discussion

Technological progress has led to rapid evolution in health data systems with immediate impact in daily clinical practice. The potential for improving patient care and outcomes are clear, as are the challenges and limitations to achieving this objective.²² Big data analytics now support large scale (and cost efficient) clinical research, with trials based within registries or the EHR itself now heralding a new era in evidence generation. These processes can be further developed by an accompanying social license and upskilling of knowledge for all stakeholders. Co-creation and shared decision making with patients and the public²³ is an important way to ensure appropriate data stewardship and privacy, leading to clinical impact through robust publications, regulatory decision making, and practice guidelines. In this paper, we have reported on a global multistakeholder process to develop a framework for researchers to use in the design and reporting of studies that include structured or coded healthcare data.

Digital health records are confusing for most researchers, with varying access to a myriad of different coding systems and classifications, and considerable differences across (and within) countries. Linkage of different health sources is often a core component of research based on structured healthcare data, and yet, this aspect is frequently overlooked when reporting such studies. Data privacy and the license for research can be severely compromised if linkage is not secure; hence, our focus is on transparency about how data are coded and linked, and how these approaches are openly discussed and documented. The stakeholder consensus meetings highlighted this area as a key concern for future research, supported by evidence that very few studies have provided sufficient detail to understand the research process.^{24,25} The advent of registry and EHR-based randomised controlled trials^{21,26,27} reinforces the imperative to see improvements in these areas, and to define new concepts for quality research. With the development of robust analytics supported by machine learning algorithms,²⁸ similar approaches have already been used to support artificial intelligence in healthcare.²⁹

A lack of transparency has a direct impact on the value of research using coded records, with issues arising for medical journals, regulators, clinical guideline writers, and more generally clinicians and the



public. Bringing together the full range of these stakeholders, we aimed to take full advantage of recent technical developments to use structured healthcare data for research, to approach limitations directly, and to provide a framework across all medical fields where coded data can be used to improve patient care. A number of other overlapping themes emerged from the discussions, including the generation and retainment of public trust and confidence, and the need for coherent plans to deal with data security failures. Forethought about dealing with the harmonisation of data and the requirement for embedded validation methods were highlighted as key factors for future successful research. Similarly, education and communication are crucial for patients, citizens and healthcare professionals to effectively use the results from structured healthcare data studies.

The covid-19 pandemic has illustrated the need for rapid access to routine healthcare data to guide and monitor clinical care, and a clinical trial infrastructure to allow for immediate deployment. The digitalisation of healthcare, in particular the use of EHRs, offered the clinical community a unique opportunity to develop a learning healthcare system that could efficiently address the effects of covid-19. For example, information about the relationship between covid-19 and cardiovascular disease through linked EHR data that has combined primary care data, hospital data, death records, and covid-19 testing in more than 54 million people.³⁰ However, the pandemic also made clear the obstacles within various systems that restricted the sharing of data in almost real time that could direct care and help design clinical trials. Established governance, security, interoperability (system

Table 1 Output from the stakeholder consensus meetings

Workshop theme	Key consensus statements and advisories
1. Technical process and data stewardship	<p><i>Research using structured healthcare data conducted according to the FAIR data principles of findability, accessibility, interoperability, and reusability.</i>¹³ Important considerations are transparency of who performed the coding, the coding system used, and the purpose of coding (reimbursement, diagnosis, etc.).</p> <p><i>Clear and consistent identification and description of the sources of EHR data.</i> Code lists and phenotyping algorithms can be described in detail and published, ideally before a study commences (for example on a coding repository or open-source archive). The minimum data required to meet the definitions will depend on the use case and can be reported to enhance transparency, in addition to the rationale for why certain decisions were made (for example, why one code was chosen over another, or what the effect would be if data collection periods were changed).</p> <p><i>Validation at local, regional and global levels.</i> Evidence demonstrating how algorithms have been externally validated, and also what quality assessment was performed on the research findings, for example on the accuracy, completeness and timeliness of the data.¹⁴ Data quality rules can be used to assess coded data and allow comparisons across institutions and countries.¹⁵</p> <p><i>Reporting of methods used for data pre-processing and data linkage.</i> This includes the methods used to assess the quality of linkage and the results of any data pre-processing and linkage (with provision of false positive and false negative rates, comparisons of linked and unlinked data, and any sensitivity analyses).¹⁶ A flow diagram showing the processes for cleaning and linking different coding sources and datasets can aid understanding of the study design.</p> <p><i>Reporting of the governance framework underpinning the study from a technical/data stewardship standpoint.</i> This includes a clear purpose for data gathering and the parameters and time limit of consent, clear mechanisms for data processing ("what happens with my data"), and a description of what the data can and cannot be used for (i.e. the mandate given for research).</p>
2. Data security and privacy	<p><i>Working towards a new, sustainable mandate from the public and patients to use their health data may require moving away from abstract rules and regulations and towards more constructive governance, in which trust is a central concept.</i> The trust of patients and the public in research institutions and in science is pivotal because of the liberties they give to researchers to use their data, which are the product of a social licence based on this trust.</p> <p><i>Gaining this trust would benefit from understanding what society and stakeholders expect from scientists conducting health data research, with engagement of stakeholders from the concept stage.</i> Co-creation of data governance based on inclusion of patient/public communities and dialogue with researchers is crucial for ethical and sustainable governance, and to translate expectations into scientific research and scientific output.</p> <p><i>Researchers and big data consortia have to be mindful that trustworthiness comes with the duty to act in ethically-responsible ways.</i> This concerns two areas; first is the competence in data handling (meaning that systems are in place to ensure data protection and there is a framework of rules and regulations for data sharing), and second is what motivates the data analysis. Ongoing dialogue can ensure that public values continue to be aligned with the governance structures of health data research projects. Questions arise as to how to measure success at implementing public values into research, and what levels of public support are sufficient to grant a mandate for data usage.</p> <p><i>Complex organisational structures may be less important for this trust than is often asserted.</i> Complicated rules and regulations may do more harm than good in establishing the conditions for public trust in big data health research to flourish, and as a result be counterproductive especially when a social license has not been adequately achieved.¹⁷</p> <p><i>Embracing values such as transparency, reciprocity, inclusivity and service to the common good.</i> These values can be embedded into the governance framework of big data health research.¹⁸ This calls for constructing a narrative that researchers and research consortia can be held accountable so that patients and the wider public are willing, and consistently willing, to place their trust in health research projects.</p> <p><i>Governance could be aided by developing a framework for accountability.</i> This includes clear distinctions between anonymised, pseudonymised and aggregate data along with plain language explanation to participants and users, and discrimination between primary and secondary use of data sources.</p>
3. Publications using structured healthcare data	<p><i>Accountability for the source of data and how the data have been collected (traceability).</i> As with data security, a framework of accountability would enable editorial teams in medical journals to be aware of the technical processes prior to data analysis.</p> <p><i>Sharing of data, codes and algorithms used to analyse datasets.</i> Similar to the requirement for pre-registration of clinical trials and pre-publication of protocols, journals could restrict publication where the coding within a study is not shared.</p> <p><i>Demonstration of data validity and robust analysis.</i> The FDA and EMA already suggest independent checking or accreditation of data sources; this accreditation could be provided to editors to increase their confidence in data quality.</p>

Continued

Table 1 Continued

Workshop theme	Key consensus statements and advisories
	<p><i>Balancing the speed of publication against requirements for data validation.</i> Prompt publication (for example of results with immediate public health implication) needs to be balanced against validation of data sources to ensure authenticity.</p> <p><i>Scientific advice committees with experts in big data analytics to aid journal editorial teams.</i> The skill-set required in editors and reviewers for studies using structured healthcare data is not the same as having statistical or clinical trials experience; expertise in EHR data and respective coding systems could add value to the journal review process.</p> <p><i>Widening gap between the knowledge of physicians and the advanced methodologies used in big data papers.</i> Medical/graduate students and practising clinicians, as well as hospital managers and leadership, need training in health data management and analysis. This is important to build a digital workforce with increased capacity and capability to translate publications using new approaches to improve patient care.¹⁹</p>
4. Addressing the needs of regulators, reimbursement authorities and clinical practice guidelines	<p><i>EHR-based trials have the potential to generate reliable and cost-efficient results.</i> Each type of trial and each type of clinical question is considered in an individual context, including under what circumstances a particular type of EHR process could assist in answering questions about a particular intervention, and with what limitations.</p> <p><i>Further research may help explore cases in which EHR studies produce valuable evidence, and when they might be flawed.</i> This will generate confidence in regulators for future EHR studies, and for guideline taskforces to appropriately appraise evidence.</p> <p><i>Quality standards will help to ensure that the information recorded in EHR systems represents real events without bias.</i> This will enable confidence that trials using EHRs can produce reliable results on efficacy and safety, and could include examination of the validity of both data sources and data analyses.</p> <p><i>Source data validation to report on appropriate computational phenotypes.</i> This could be supported by an independent adjudication committee to examine a subset of the EHR and confirm outcome events. The use of AI techniques could facilitate larger validation studies by automated extraction of supporting text from clinical notations. Such validation exercises can be pre-registered, for example in the form of a Study-Within-A-Trial.²⁰ Another possibility is for researchers to provide consented and anonymised gold standard cases to benchmark against, or for data from devices used to verify codes (such as lead fractures). The value of synthetic datasets for validation, which mimic real data, needs further exploration.</p> <p><i>Mixed model approaches to collect data on particular endpoints.</i> May be valuable for situations where the EHR does not reliably collect relevant data. For example, where patients and/or clinicians are asked for information, or data are collected via wearable devices or telemonitoring. In some cases, parallel monitoring of patients alongside the EHR study may provide additional confidence (for example to identify serious unexpected adverse events). Technological advances in EHR systems will help, such as the ability to retrieve EHR data on a daily basis to support clinical trials.²¹</p> <p><i>Taking advantage of the many real world data initiatives to support new research.</i> Government agencies, regulators, charities and professional bodies have initiated programmes for better use of real-world data that can support further activity and dissemination.</p>

architecture that spans different EHR systems and healthcare providers), and phenotype definition, among other issues, limited access to routine EHR data especially in the first period of the pandemic.

The CODE-EHR framework is intended to complement available reporting checklists.^{31–34} Although existing checklists are aimed at transparency in the reporting of important methodological components of clinical research, the CODE-EHR framework is designed to ensure that a common set of minimum standards are applied across all research using structured healthcare data. This range includes observational studies and controlled trials, with the preferred standards giving the direction of research design for all future EHR studies. Additionally, the framework supports the wider implementation of good quality real world data research based on the FAIR data principles.¹³ Researchers are advised to use the checklist during the design phase of their study to ensure that key criteria for successful research and research impact are already embedded. The process will aid journal editors, regulators, guideline writers, clinicians, and patients to better appreciate the underpinning value, and also the

limitations of the study. Dissemination plans for CODE-EHR include discussion with journals to request authors complete the checklist when submitting relevant research, attaining full registration with the EQUATOR network,³⁵ and outreach via international digital health groups to engage their members and other relevant stakeholder organisations. After publication, the CODE-EHR framework will undergo a two year evaluation, including discussion with researchers using the approach, with a plan for iterative improvements to adapt to this rapidly developing field of medical research.

Conclusion

The CODE-EHR framework was designed by a multistakeholder panel to improve design and reporting of research studies using structured electronic healthcare data. Research using these data sources is a vital component of future healthcare evaluation and delivery and will take an increasingly important role in decisions by regulatory,

POSITIVE roadmap to research benefiting from Patient & Public Involvement



Figure 2 Patient and public engagement to improve clinical research. POSITIVE steps leading to co-creation with patients and the public, and better research using big data sources. Content adapted from the Consensus Statement on Public Involvement and Engagement with Data-Intensive Health Research³⁶ as used in the DaRe2THINK trial programme.²⁷ Adapted from Bunting et al.³⁷ PPI=patient and public involvement.

governmental, and healthcare agencies, as well as clinicians and patients in every medical specialty. The CODE-EHR checklist asks for clarity on reporting and defines a set of minimum and preferred standards on the processes that underpin coding, dataset construction and linkage, disease and outcome definitions, analysis, and research governance. Iterative updates to this framework are expected to enhance research quality and value and to generate new pathways for impact using routinely collected healthcare data.

Contributors

DK and FWA are joint first authors with equal contribution, and act as guarantors of the work. Correspondence to FWA at f.w.asselbergs@umcutrecht.nl. Each subsequent listed author was involved in the CODE-EHR consensus process and wrote or revised sections of text and revised the final manuscript for intellectual content. The corresponding authors attest that all listed authors meet authorship criteria and that no others meeting the criteria have been omitted.

Supplementary data

Supplementary data is available at *European Heart Journal* online.

Acknowledgments

We are grateful for the work of the CODE-EHR international consensus. Delegates and attendees of the meetings were: Regulatory agencies: European Medicines Agency, Netherlands (Xavier Kurz); US Food and Drug Administration, USA (John Concato, Robert Kazmierski, Jose Pablo Morales, and Ileana Piña). Reimbursement authorities: National Health Care Institute, the Netherlands (Wim Goettsch); National Institute for Health and Care Excellence, UK (Adrian Jonas); Dental and Pharmaceutical Benefits Agency (TLV), Sweden (Niklas Hedberg). Medical Journal Editors: *European Heart Journal*, Switzerland (Filippo Crea and Thomas F Lüscher); *The BMJ*, UK (Wim Weber and Tiago Villanueva); *The Lancet*, UK (Stuart Spencer); *The Lancet Digital Health*, UK (Rupa Sarkar). Industry representatives: AstraZeneca (Martin Fredriksson and Mats Sundgren); Bayer (Andrea Derix, Gunnar Brobert, and Lothar Roessig); Servier (Benoit Tyl); The Hyve (Kees van Bochove); Vifor Pharma (Maureen Cronin). Funders: EU/EFPIA Innovative Medicines Initiative, Belgium (Colm Carroll); European Commission, Belgium (Ceri Thompson). Patient Advocacy: European Heart Network, Belgium (Birgit Beger, Susanne Løgstrup, and Marilena Vrana); European Society of Cardiology Patient Forum, France (Paul McGreavy). Clinical Academic and Professional Societies: European Society of Cardiology, France

Table 2 CODE-EHR framework: best practice checklist to report on the use of structured electronic healthcare records in clinical research

Date of completion:		Study name:		
Item	Objective	Framework standards	Minimum information to provide	Lead author acknowledgment
1. Dataset construction and linkage	To provide an understanding of how the structured healthcare data were identified and used.	Minimum: Flow diagram of datasets used in the study, and description of the processes and directionality of any linkage performed, published within the research report or supplementary documents. Preferred: Provided within a pre-published protocol or open access document.	(a) State the source of any datasets used. (b) Comment on how the observed and any missing data were identified and addressed, and the proportion observed for each variable. (c) Provide data on completeness of follow-up. (d) For linked datasets, specify how linkage was performed and the quality of linkage methods.	Choose one from: (1) Minimum standard not met (2) Minimum standard met OR (3) Preferred standard met
2. Data fit for purpose	To ensure transparency with the approach taken, with respect to coding of the structured healthcare data.	Minimum: Clear unambiguous statements on the process of coding in the methods section of the research report. Preferred: Provided within a pre-published protocol or open access document.	(a) Confirm origin, clinical processes, and the purpose of data. (b) Specify coding systems, clinical terminologies, or classification used and their versions, and any manipulation of the coded data. (c) Provide detail on quality assessment for data capture. (d) Outline potential sources of bias.	Choose one from: (1) Minimum standard not met (2) Minimum standard met OR (3) Preferred standard met
3. Disease and outcome definitions	To fully detail how conditions AND outcome events were defined, allowing other researchers to identify errors and repeat the process in other datasets.	Minimum: State what codes were used to define diseases, treatments, conditions, and outcomes <i>prior to statistical analysis</i> , including those relating to patient identification, therapy, procedures, comorbidities, and components of any composite endpoints. Preferred: Provided within a pre-published protocol or open access document <i>prior to statistical analysis</i> .	(a) Detailed lists of codes used for each aspect of the study. (b) Date of publication and access details for the coding manual (please add to box below). (c) Provide definitions, implementation logic and validation of any phenotyping algorithms used. (d) Specify any processes used to validate the coding scheme or reference to prior work.	Choose one from: (1) Minimum standard not met (2) Minimum standard met OR (3) Preferred standard met
4. Analysis	To fully detail how outcome events were analysed and allow independent assessment of the authenticity of study findings.	Minimum: Describe the process used to analyse study outcomes, including statistical methods and use of any machine learning or algorithmic approaches. Preferred: Provide a statistical analysis plan as a supplementary file, locked before analyses commencing.	(a) Provide details on all statistical methods used. (b) Provide links to any machine code or algorithms used in the analysis, preferably as open source. (c) Specify the processes of testing assumptions, assessing model fit and any internal validation. (d) Specify how generalisability of results was assessed, the replication of findings in other	Choose one from: (1) Minimum standard not met (2) Minimum standard met OR (3) Preferred standard met

Continued

Table 2 Continued

Date of completion:		Study name:		
Item	Objective	Framework standards	Minimum information to provide	Lead author acknowledgment
			datasets, or any external validation.	
5. Ethics and governance	To provide patients, who might or might not have given consent, and regulatory authorities the ability to interrogate the security and provenance of the data.	Minimum: Clear unambiguous statements on how the principles of Good Clinical Practice and Data Protection will be/were met, provided in the methods section of the research report. Preferred: Provided within a pre-published protocol or open access document with evidence of patient and public engagement.	(a) State how informed consent was acquired, or governance if no patient consent. (b) Specify how data privacy was protected in the collection and storage of data. (c) Detail what steps were taken for patient and public involvement in the research study. (d) Provide information on where anonymised source data or code can be obtained for verification and further research.	Choose one from: (1) Minimum standard not met (2) Minimum standard met OR (3) Preferred standard met
6. Coding manual	DOI of publication or website address: Date published:			
7. Comments				
8. Summary declaration	Choose one from: One or more minimum standards not met OR All minimum standards met Number of preferred standards met: / 5			

Directions for use:

Research team: To complete the checklist, authors will need to consider these points during the design of the research to ensure that coding protocols and coding manuals are pre-published. Where applicable, it is advisable that all five minimum standards are met for an individual research study, whether observational or a controlled trial. If any component is not applicable to the study, the corresponding author can indicate why this is the case in the comment box. This checklist can accompany the article as a supplementary file on submission to the journal, with the ability for readers to review responses. A comment on the meeting of standards in the text of the method section is suggested, eg: "this study meets all five of the CODE-EHR minimum framework standards for the use of structured healthcare data in clinical research, with two out of five standards meeting preferred criteria <https://doi.org/10.1093/eurheartj/ehac426>"; OR "this study meets four out of five of the CODE-EHR minimum framework standards for the use of structured healthcare data in clinical research; one of the five minimum standards was not met as coding schemes were not specified prior to analysis <https://doi.org/10.1093/eurheartj/ehac426>." Note, easy to complete form versions of this checklist are available in appendix 4 (word version) and appendix 5 (pdf version), and also at <https://www.escardio.org/bigdata>.

Research appraisers (patients, clinicians, regulators, guideline task forces): Where applicable, it is advisable that all five minimum standards are met for the research study to be considered robust.

(Barbara Casadei, Stephan Achenbach, and Valentina Tursini); European Heart Agency, Belgium (Panos E Vardas); Oslo University Hospital and University of Oslo, Norway (Dan Atar); University of Oxford, UK (Colin Baigent); University of Leeds, UK (Chris P Gale); Queen's University Belfast, UK (Donna Fitzsimons); University Hospital Bern, Switzerland (Stephan Windecker); Charité-Universitätsmedizin Berlin, Germany (Stefan D Anker); Royal Brompton Hospital and Imperial College London, UK (Martin Cowie); University College London, UK (Amitava Banerjee, Harry Hemingway, R Tom Lumbers, Spiros Denaxas); University Medical Centre Utrecht, Netherlands (Folkert W Asselbergs, Rick Grobbee, and Ghislaine Van Thiel); University of Birmingham and University Hospitals Birmingham NHS Trust, UK (Dipak Kotecha and George V Gkoutos); University of Cambridge, UK (Angela Wood); Karolinska Institutet and Karolinska University Hospital Stockholm, Sweden (Martin Ingvar). Administrative: BigData@Heart Project Management, Germany (Carl Steinbeisser

and Ana Petrova); European Society of Cardiology Project Management, France (Cinzia Ceccarelli, Katija Baljevic, and Polyxeni Vairami); Medical Writer, UK (Jennifer Taylor).

Funding

The BigData@Heart project has received funding from the Innovative Medicines Initiative 2 Joint Undertaking under grant agreement No 116074. This Joint Undertaking receives support from the European Union's Horizon 2020 research and innovation programme and European Federation of Pharmaceutical Industries and Associations. The funders had no role in considering the study design or in the collection, analysis, interpretation of data, writing of the report, or decision to submit the article for publication.

Competing interests

All authors have completed the ICMJE uniform disclosure form (www.icmje.org/coi_disclosure.pdf) and declare: DK reports grants

from EU/EFPIA Innovative Medicines Initiative (BigData@Heart 116074), during the study; grants from National Institute for Health Research (NIHR CDF-2015-08-074 RATE-AF; NIHR130280 DaRe2THINK; NIHR132974 D2T-NeuroVascular); British Heart Foundation (PG/17/55/33087, AA/18/2/34218 and FS/CDRF/21/21032); and European Society of Cardiology supported by educational grants from Boehringer Ingelheim, BMS-Pfizer Alliance, Bayer, Daiichi Sankyo, Boston Scientific, the NIHR/University of Oxford Biomedical Research Centre, and British Heart Foundation/University of Birmingham Accelerator Award (STEER-AF NCT04396418); Amomed Pharma and IRCCS San Raffaele/Menarini (Beta-blockers in Heart Failure Collaborative Group NCT0083244), outside of this work; and advisory board personal fees from Bayer, Amomed, Protherics Medicines Development and Myokardia, outside of this work. FWA reports grants from IMI BigData@Heart, during the study. SDA reports grants and personal fees from Vifor Int, Abbott, and Abbott Vascular; personal fees from Bayer, Boehringer Ingelheim, and Servier; personal fees from Cardiac Dimensions, Actimed, Astra Zeneca, Amgen, Bioventrix, Janssen, Respicardia, V-Wave, Brahms, Cordio, and Occlutech, outside of this work. CB reports grants from Medical Research Council, Boehringer Ingelheim, and NIHR, outside of this work. AB reports grants from Astra Zeneca, outside of this work. BB reports grants from European Commission, during the study. GB reports grants from European Commission (IMI project support), during the study; and other from Bayer AB, outside of this work. BC reports non-financial support from Roche Diagnostics and iRhythm, outside of this work. CC reports grants from European Commission, during the study. MRC reports personal fees from AstraZeneca, outside of this work. FC reports personal fees from Amgen, Astra Zeneca, Servier, and BMS; other from GlyCardial Diagnostics, outside of this work. MC reports personal fees from Vifor Pharma, during the study; and personal fees from Vifor Pharma, outside of this work. AD reports other from Bayer AG, outside of this work. MF reports other from AstraZeneca, during the study; other from AstraZeneca, outside of this work; and is employed by AstraZeneca. CPG reports personal fees from AstraZeneca, Amgen, Bayer, Boehringer-Ingelheim, Daiichi Sankyo, Vifor Pharma, Menarini, Wondr Medical, Raisio Group, and Oxford University Press; and grants from BMS, Abbott, British Heart Foundation, NIHR, and ESC, outside of this work. MI reports grants from World Economic Forum, Swedish Innovation Agency, and European Commission, during the study; and other collaboration with Frisq AB, outside of this work. SL reports grants from European Commission, during the study. TFL reports grants from Abbott, Amgen, Novartis, Boehringer Ingelheim, Servier, Vifor, Sanofi, and AstraZeneca; and personal fees from Daiichi Sankyo, Pfizer, and Menarini, outside of this work. LR reports other from Bayer AG, during the study and other from Bayer AG, outside this work. CS reports personal fees from Bayer AG, during the study and personal fees from Bayer AG, outside of this work. BT reports personal fees from Servier, outside of this work. GvT reports grants from IMI, during the study. KvB reports grants from IMI BigData@Heart, during the study. PEV reports personal fees from Hygeia Hospitals Group, HHG group, European Society of Cardiology, and Servier International, outside of this work. TV is working as an editor at *The BMJ* and *Acta Médica Portuguesa* and is

vice president of the European Union of General Practitioners (UEMO). MV reports grants from European Commission, during the study. SW reports grants from Abbott, Amgen, Astra Zeneca, BMS, Bayer, Biotronik, Boston Scientific, Cardinal Health, CardioValve, CSL Behring, Daiichi Sankyo, Edwards Lifesciences, Guerbet, InfraRedx, Johnson & Johnson, Medicure, Medtronic, Novartis, Polares, OrPha Suisse, Pfizer, Regeneron, Sanofi-Aventis, Sinomed, Terumo, and V-Wave, outside of this work; and SW serves as unpaid advisory board member or unpaid member of the steering or executive group of trials funded by Abbott, Abiomed, Amgen, Astra Zeneca, BMS, Boston Scientific, Biotronik, Cardiovalve, Edwards Lifesciences, MedAlliance, Medtronic, Novartis, Polares, Sinomed, V-Wave, and Xeltis, but has not received personal payments by pharmaceutical companies or device manufacturers. SW is also member of the steering or executive committee group of several investigator-initiated trials that receive funding by industry without impact on his personal remuneration. SW is an unpaid member of the Pfizer Research Award selection committee in Switzerland and of the Women as One Awards Committee. SW is also member of the Clinical Study Group of the Deutsches Zentrum für Herz Kreislauf-Forschung and of the Advisory Board of the Australian Victorian Heart Institute. He is chairperson of the ESC Congress Program Committee, former chairperson of the ESC Clinical Practice Guidelines Committee and Deputy Editor of *JACC CV Interventions*. All other authors declare no competing interests.

Dissemination to participants and related patient and public communities: Output will be disseminated to relevant patient and public communities via the European Society of Cardiology, the European Heart Network, and through communication with other relevant international bodies.

Provenance and peer review: Not commissioned; externally peer reviewed.

References

1. Conrad N, Judge A, Tran J, et al. Temporal trends and patterns in heart failure incidence: a population-based study of 4 million individuals. *Lancet* 2018;**391**:572–80.
2. Hernández B, Reilly RB, Kenny RA. Investigation of multimorbidity and prevalent disease combinations in older Irish adults using network analysis and association rules. *Sci Rep* 2019;**9**:14567.
3. Oni T, Youngblood E, Boule A, McGrath N, Wilkinson RJ, Levitt NS. Patterns of HIV, TB, and non-communicable disease multi-morbidity in peri-urban South Africa: a cross sectional study. *BMC Infect Dis* 2015;**15**:20.
4. Studer R, Sartini C, Suzart-Woischnik K, et al. Identification and mapping real-world data sources for heart failure, acute coronary syndrome, and atrial fibrillation. *Cardiology* 2022;**147**:98–106.
5. Wallentin L, Gale CP, Maggioni A, Bardinet I, Casadei B. EuroHeart: European Unified registries on heart care evaluation and randomized trials. *Eur Heart J* 2019;**40**:2745–9.
6. Sherman RE, Anderson SA, Dal Pan GJ, et al. Real-world evidence - what is it and what can it tell us? *N Engl J Med* 2016;**375**:2293–7.
7. John VV, Ian H, Rafael J, et al. Implementation and relevance of FAIR Data principles in biopharmaceutical research and development. *Drug Discov Today* 2019;**24**:933–8.
8. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Lancet* 2007;**370**:1453–7.
9. Nicholls SG, Quach P, von Elm E, et al. The REporting of Studies Conducted Using Observational Routinely-Collected Health Data (RECORD) statement: methods for arriving at consensus and developing reporting guidelines. *PLoS One* 2015;**10**: e0125620.
10. Liu X, Rivera SC, Moher D, Calvert MJ, Denniston AK; SPIRIT-AI and CONSORT-AI Working Group. Reporting guidelines for clinical trial reports for interventions involving artificial intelligence: the CONSORT-AI Extension. *BMJ* 2020;**370**:m3164.

11. Logullo P, McCarthy A, Kirtley S, Collins GS. Reporting guideline checklists are not quality evaluation forms: they are guidance for writing. *Health Sci Rep* 2020;**3**:e165.
12. Dewey M, Levine D, Bossuyt PM, Kressel HY. Impact and perceived value of journal reporting guidelines among Radiology authors and reviewers. *Eur Radiol* 2019;**29**:3986–95.
13. Wilkinson MD, Dumontier M, Aalbersberg IJ, et al. The FAIR guiding principles for scientific data management and stewardship. *Sci Data* 2016;**3**:160018.
14. Chen H, Hailey D, Wang N, Yu P. A review of data quality assessment methods for public health information systems. *Int J Environ Res Public Health* 2014;**11**:5170–207.
15. Peng M, Lee S, D'Souza AG, Doktorchik CTA, Quan H. Development and validation of data quality rules in administrative health data using association rule mining. *BMC Med Inform Decis Mak* 2020;**20**:75.
16. Harron K, Dibben C, Boyd J, et al. Challenges in administrative data linkage for research. *Big Data Soc* 2017;**4**:2053951717745678.
17. Carter P, Laurie GT, Dixon-Woods M. The social licence for research: why care.data ran into trouble. *J Med Ethics* 2015;**41**:404–9.
18. Muller SHA, Kalkman S, van Thiel GJM, Mostert M, van Delden JJM. The social licence for data-intensive health research: towards co-creation, public value and trust. *BMC Med Ethics* 2021;**22**:110.
19. Kotecha D, Bax JJ, Carrera C, et al.; ESC Scientific Document Group; 2016 ESC Education Conference. Roadmap for cardiovascular education across the European Society of Cardiology: inspiring better knowledge and skills, now and for the future. *Eur Heart J* 2019;**40**:1728–38.
20. Treweek S, Bevan S, Bower P, et al. Trial Forge Guidance 1: what is a Study Within A Trial (SWAT)? *Trials* 2018;**19**:139.
21. Wang X, Mobley AR, Tica O, et al. Systematic approach to outcome assessment from coded electronic healthcare records in the DaRe2THINK NHS-embedded randomised trial. *medRxiv* 2022;10.1101/2022.05.24.22275434.
22. Coorevits P, Sundgren M, Klein GO, et al. Electronic health records: new opportunities for clinical research. *J Intern Med* 2013;**274**:547–60.
23. European Patient Innovation Summit. Position paper on digital health from a patient perspective. 2019: <https://www.episummit.net/position-paper.html>.
24. Springate DA, Kontopantelis E, Ashcroft DM, et al. ClinicalCodes: an online clinical codes repository to improve the validity and reproducibility of research using electronic medical records. *PLoS One* 2014;**9**:e99825.
25. Bohensky MA, Jolley D, Sundararajan V, et al. Data linkage: a powerful research tool with potential problems. *BMC Health Serv Res* 2010;**10**:346.
26. Fröbert O, Lagerqvist B, Olivecrona GK, et al.; TASTE Trial. Thrombus aspiration during ST-segment elevation myocardial infarction. *N Engl J Med* 2013;**369**:1587–97.
27. Kotecha D, Shukla D, Clinical Practice Research Datalink. Preventing stroke, premature death and cognitive decline in a broader community of patients with atrial fibrillation using healthcare data for pragmatic research: a randomised controlled trial (DaRe2THINK). Sponsor: University of Birmingham; Funder: National Institute for Health Research 2021;EudraCT 2020-005774-10; ClinicalTrials.gov NCT04700826: <http://www.birmingham.ac.uk/research/cardiovascular-sciences/research/dare2think/index.aspx>.
28. Karwath A, Bunting KV, Gill SK, et al.; card Alc group and the Beta-blockers in Heart Failure Collaborative Group. Redefining β -blocker response in heart failure patients with sinus rhythm and atrial fibrillation: a machine learning cluster analysis. *Lancet* 2021;**398**:1427–35.
29. World Health Organization. Ethics and governance of artificial intelligence for health: WHO guidance. ISBN: 9789240029200 2021: <https://www.who.int/publications/i/item/9789240029200>.
30. Wood A, Denholm R, Hollings S, et al.; CVD-COVID-UK consortium. Linked electronic health records for research on a nationwide cohort of more than 54 million people in England: data resource. *BMJ* 2021;**373**:n826.
31. Schulz KF, Altman DG, Moher D, Group C; CONSORT Group. CONSORT 2010 statement: updated guidelines for reporting parallel group randomised trials. *BMJ* 2010;**340**:c332.
32. von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP; STROBE Initiative. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Ann Intern Med* 2007;**147**:573–7.
33. Benchimol EI, Smeeth L, Guttman A, et al.; RECORD Working Committee. The REporting of studies Conducted using Observational Routinely-collected health Data (RECORD) statement. *PLoS Med* 2015;**12**:e1001885.
34. Chan AW, Tetzlaff JM, Altman DG, et al. SPIRIT 2013 statement: defining standard protocol items for clinical trials. *Ann Intern Med* 2013;**158**:200–7.
35. Simera I, Moher D, Hirst A, Hoey J, Schulz KF, Altman DG. Transparent and accurate reporting increases reliability, utility, and impact of your research: reporting guidelines and the EQUATOR Network. *BMC Med* 2010;**8**:24.
36. Aitken M, Tully MP, Porteous C, et al. Consensus statement on public involvement and engagement with data-intensive health research. *Int J Pop Data Sci* 2019;**4**:586.
37. Bunting KV, Stanbury M, Tica O, Kotecha D. Transforming clinical research by involving and empowering patients- the RATE-AF randomized trial. *Eur Heart J* 2021;**42**:2411–4.