

Boosting peripheral BDNF rescues impaired in vivo axonal transport in CMT2D mice

Authors: James N. Sleight^{1,2,*}, David Villarroel-Campos¹, Sunaina Surana^{1,2}, Tahmina Wickenden^{1†}, Yao Tong³, Rebecca L. Simkin¹, Jose Norberto S. Vargas¹, Elena R. Rhymes¹, Andrew P. Tosolini¹, Steven J. West⁴, Qian Zhang³, Xiang-Lei Yang³, Giampietro Schiavo^{1,2*}

Affiliations:

¹ Department of Neuromuscular Diseases and UCL Queen Square Motor Neuron Disease Centre, UCL Queen Square Institute of Neurology, University College London, London, UK.

² UK Dementia Research Institute, University College London, London, UK.

³ Department of Molecular Medicine, The Scripps Research Institute, La Jolla, California, United States of America.

⁴ Sainsbury Wellcome Centre, University College London, London, UK.

† Current affiliation: UCL Great Ormond Street Institute of Child Health, University College London, London, UK.

* Corresponding authors

Contact:

j.sleigh@ucl.ac.uk/+44(0)2031082722

giampietro.schiavo@ucl.ac.uk/+44(0)2034484334

Conflict-of-interest Statement

The technology described in this work has been protected in the patent GB2303495.2 (patent applicant, UCL Business Ltd., status pending), in which JNS and GS are named as inventors.

The other authors declare no competing interests.

Abstract

Gain-of-function mutations in the housekeeping gene *GARS1*, which lead to the expression of toxic versions of glycyl-tRNA synthetase (GlyRS), cause the selective motor and sensory pathology characterising Charcot-Marie-Tooth disease (CMT). Aberrant interactions between GlyRS mutants and different proteins, including neurotrophin receptor TrkB, underlie CMT type 2D (CMT2D); however, our pathomechanistic understanding of this untreatable peripheral neuropathy remains incomplete. Through intravital imaging of the sciatic nerve, we show that CMT2D mice display early and persistent disturbances in axonal transport of neurotrophin-containing signalling endosomes *in vivo*. We discovered that BDNF-TrkB impairments correlate with transport disruption and overall CMT2D neuropathology, and that inhibition of this pathway at the nerve-muscle interface perturbs endosome transport in wild-type axons. Accordingly, supplementation of muscles with BDNF, but not other neurotrophins, completely restores physiological axonal transport in neuropathic mice. Together, these findings suggest that selectively targeting muscles with BDNF-boosting therapies could represent a viable therapeutic strategy for CMT2D.

Key Words: BDNF, intravital imaging, motor neuron, neuromuscular junction (NMJ), neurotrophic factors, neurotrophin signalling, peripheral neuropathy, signalling endosome, Trk receptors.

Brief Summary: Boosting BDNF in the muscles of mice modelling Charcot-Marie-Tooth disease type 2D rescues *in vivo* deficits in axonal transport of signalling endosomes.

Introduction

Charcot-Marie-Tooth disease (CMT) is an inherited peripheral nerve disorder resulting in life-long disability, with no available disease-modifying therapies (1). Patients usually present with muscle weakness and wasting, as well as sensory deficits, in the hands and feet, indicating that motor and sensory neurons with the longest axons are generally the most susceptible to neuropathy. Given that CMT is caused by mutations in more than 90 genes with diverse functions (2), a single unifying pathomechanism is unlikely; nevertheless, compromised *in vitro* axonal transport has been identified in several CMT subtypes (3, 4) and is hypothesised to be a major driver of axon and synaptic degeneration (5, 6).

CMT type 2D (CMT2D) manifests during adolescence due to dominantly inherited, missense mutations in the widely and constitutively expressed *GARS1* gene, which encodes glycyl-tRNA synthetase (GlyRS), an enzyme essential for protein translation (7). Whereas the mechanisms causing selective motor and sensory nerve pathology in CMT2D remain to be fully resolved (8), *GARS1* mutations have been reported to induce a conformational opening in GlyRS (9, 10), exposing common buried surfaces that enable toxic mis-interactions with several proteins both inside and outside of neurons (11-16). Aberrant binding partners include the neuronal and vascular transmembrane receptor protein neuropilin-1 (11) and the histone deacetylase HDAC6 (14). Enhanced associations with wild-type GlyRS interactors are also possible. Indeed, recent evidence indicates that mutant GlyRS binds but fails to efficiently release tRNA^{Gly}, which leads to ribosome stalling at glycine codons, inhibition of protein synthesis and activation of the integrated stress response (17, 18). Accordingly, tRNA^{Gly} overexpression resulted in a rescue of neuropathy phenotypes in both *Drosophila melanogaster* and mouse models of CMT2D (18). Impaired protein translation is thus a major component of *GARS1* neuropathy, as well as other diseases affecting peripheral nerves (19, 20); however, the driver

of differential vulnerability in neuropathy across motor and sensory neuron subtypes remains to be determined.

We recently discovered that CMT2D-causing, but not wild-type, GlyRS interacts with extracellular domains of tropomyosin receptor kinase (Trk) receptors A, B and C, contributing to the developmental perturbation of sensory neurons in mutant *Gars* mice (13, 21). Via their role as neurotrophin receptors, the Trk receptors are critical to neuronal differentiation and homeostasis (22). Through selective binding and internalisation of secreted neurotrophins at axon terminals, Trk receptors are retrogradely transported within signalling endosomes to somas, where they elicit gene transcription events essential for neuronal survival (23, 24). TrkA preferentially binds to nerve growth factor (NGF), TrkB to both brain-derived neurotrophic factor (BDNF) and neurotrophin-4 (NT-4), and TrkC to neurotrophin-3 (NT-3) (22).

In this study, we thus set out to determine whether *in vivo* disturbances in axonal transport of neurotrophin-containing signalling endosomes contribute to CMT2D pathology.

Results

CMT2D mice display early, persistent perturbations in endosome axonal transport in vivo

Injection of a well-characterised atoxic fluorescent fragment of tetanus neurotoxin (HcT-555) (25) into distal leg muscles permits in vivo imaging and tracking of signalling endosomes within intact sciatic nerve axons of live, anaesthetised mice (Figure 1, A and 1B) (26-28). By imaging thicker axons only, we have previously shown that transport is assessed in motor rather than sensory neurons (29). HcT-555 was therefore injected into wild-type and *Gars*^{C201R/+} mice, which carry a toxic gain-of-function mutation in endogenous *Gars* (30). The C201R mutation causes an array of progressive motor and sensory phenotypes modelling CMT2D that present by one month of age, as well as non-progressive neurodevelopmental alterations (13, 21, 30-33). Coinciding with the onset of neuromuscular junction (NMJ) denervation in the most severely affected hindpaw muscles (31), we found that *Gars*^{C201R/+} mice exhibit an overt decline in in vivo endosome transport speed that manifests between 0.5 and 1 month of age, persisting to at least 3 months (Figure 1, C and D).

To gauge the human relevance of this phenotype, we assessed endosome transport in a second CMT2D mouse model, *Gars*^{ΔETAQ/+}, which carries a deletion in endogenous mouse *Gars* modelling a mutation identified in a patient presenting with severe, early-onset neuropathy (34). First, we confirmed that human GlyRS^{ΔETAQ} aberrantly interacts with the extracellular domain of human TrkB (Supplemental Figure 1, A and B) and that the *Gars*^{ΔETAQ/+} strain displays the same perturbation of sensory neuron fate present in other mutant *Gars* alleles (Supplemental Figure 1, C and D), i.e., that there is a decrease in NF200 and an increase in peripherin levels in lumbar dorsal root ganglia (DRG) (13, 21). Analysing signalling endosome dynamics, we found that 1 month-old *Gars*^{ΔETAQ/+} mice also show impaired in vivo axonal transport early in disease and to a similar extent as *Gars*^{C201R/+} (Supplemental Figure 1, E-L),

indicating that trafficking disruption is a phenotype common to mouse strains modelling CMT2D.

To further link transport disruption to the human neuropathy, we pre-mixed and co-injected HcT-555 with purified human wild-type or CMT2D-causing GlyRS protein into distal leg muscles of wild-type mice. GlyRS^{L129P} and GlyRS^{G240R} were selected because of the strong human genetic evidence supporting their causative role in neuropathy (8, 35). We found that both mutants, but not GlyRS^{WT}, perturbed transport of signalling endosomes in otherwise healthy peripheral nerve axons of wild-type animals (Figure 1, E-H). This result indicates that extracellular exposure of motor nerve terminals to mutant GlyRS is sufficient to alter axonal endosome trafficking, suggesting the presence of a non-cell autonomous component to this phenotype, as previously identified in a *Drosophila* CMT2D model, where muscle-derived mutant GlyRS accumulates at NMJs and coincides with motor neuron degeneration (12, 15).

Together, these findings indicate that neuropathy-causing *GARS1* mutations impair axonal transport of neurotrophin-containing signalling endosomes in motor neurons in vivo and that human mutant GlyRS injected into muscle can induce this phenotype in wild-type mice.

Muscle TrkB levels correlate with denervation in CMT2D

We next sought to identify whether alterations in the BDNF-TrkB pathway correlate with neuropathy in CMT2D. In *Gars*^{C201R/+} mice we have previously identified a spectrum of vulnerability to NMJ denervation across five wholemount muscles with diverse morphological and functional properties (36); the transversus abdominis (TVA), epitrochleoanconeus (ETA), forelimb lumbrical, hindlimb lumbrical and flexor digitorum brevis (FDB) muscles display 0%, 0%, 2.5%, 11.4% and 20.3% complete denervation at 3 months, respectively (33). We

determined that developmental demands and axon length are linked to the loss of neuromuscular connectivity; however, these features do not fully determine susceptibility to disease (33).

We therefore dissected the same five muscles from wild-type mice and probed lysates for TrkB (Figure 2, A-C), the neurotrophin receptor most crucial to NMJ function and stability (37). If aberrant interactions between mutant GlyRS and TrkB disrupt BDNF-TrkB signalling, thereby contributing to peripheral denervation, then levels of TrkB, BDNF and/or GlyRS within muscles and NMJs may correlate with the degree of CMT2D pathology. Wild-type, rather than *Gars*^{C201R/+}, muscles were analysed so that baseline protein availability could be determined without interference from the differential NMJ denervation that occurs in CMT2D mice (33), which probably causes the reduced FL-TrkB levels identified in *Gars*^{C201R/+} muscles (Supplemental Figure 2). Western blotting of the thin, wholemount muscles is technically challenging due to the low protein yield, especially from the lumbrical and FDB muscles. Nevertheless, we identified consistent differences between wild-type muscles in the availability of full-length TrkB (FL-TrkB) and truncated TrkB.T1 (Figure 2, A-C). Levels of FL-TrkB, which is fundamental for pro-survival signalling, correlated with *Gars*^{C201R/+} denervation (Figure 2B), whereas levels of TrkB.T1, which lacks the essential kinase domain, did not (Figure 2C).

Similar western blot analyses were attempted with anti-BDNF antibody, but clear bands representing proBDNF and mature BDNF (mBDNF) at the expected molecular weights of ≈ 35 and 14 kDa, respectively, were not detected. Nevertheless, performing immunofluorescence analyses, we found that total BDNF levels at the wild-type NMJ inversely correlate with the percentage of partially denervated synapses in CMT2D mice (Supplemental Figure 3, A and

B). GlyRS levels did not differ significantly between muscles (Figure 2D and Supplemental Figure 3C), indicating that differences in GlyRS availability are unlikely to be a major driver of NMJ degeneration. Moreover, levels of CHRNA1, an acetylcholine receptor subunit found at the NMJ, were similar across muscles (Figure 2D and Supplemental Figure 3D), indicating that distinctions in number or size of NMJs do not account for the difference in FL-TrkB levels. Notably, there were also no differences in the availability of the motor neuron survival factor glial cell line-derived neurotrophic factor (GDNF) (Supplemental Figure 3, E-G), suggesting that the alterations found in CMT2D are specific for BDNF.

These data support the hypothesis that differences in BDNF-TrkB levels within muscles and at the NMJ may contribute to the selective vulnerability of motor nerve terminals to *GARS1* neuropathy.

Impaired CREB activation and motor neuron size is linked to CMT2D motor degeneration

The decline in endosome speed in *Gars*^{C201R/+} mice will reduce BDNF delivery to motor neuron somas. Provided that endosome flux remains unaffected, the observed 16.8% and 20.5% reductions in speed at 1 and 3 months (Figure 1), respectively, would equate to more than an extra day per week being needed to deliver the same amount of neurotrophins to the spinal cord of CMT2D mice. We hypothesised that this decrease would therefore cause dampened activation of signalling pathways that maintain neuronal health and survival, which could contribute to the slowly progressive motor and sensory neuron degeneration in CMT2D.

To assess this, lumbar spinal cords were dissected from 3 month-old mice and stained for phosphorylated and total CREB (Figure 2, E and F), a pro-survival transcription factor, which is activated in response to BDNF (38). Although differences between genotypes in nuclear p-

CREB or CREB levels did not reach significance in ChAT-positive motor neurons (Supplemental Figure 4A), CREB activation, as indicated by the ratio between p-CREB and CREB levels, was constrained in CMT2D mice (Figure 2G). Moreover, although no loss of these hindlimb-innervating lumbar motor neurons was found (Supplemental Figure 4B), we identified a clear reduction in the area of *Gars*^{C201R/+} motor neuron cell bodies and nuclei (Figure 2H).

A decrease in motor neuron size could result from switching of α motor neuron subtype from large, fast-twitch fatiguable to small, slow-twitch fatigue-resistant motor neurons or even γ motor neurons. However, we have previously shown that there is no change in the proportion of α and γ motor neurons in *Gars*^{C201R/+} spinal cords at 1 month (13), indicating that this is not the cause of the reduced lumbar motor neuron area. We thus attempted spinal cord staining for markers of different α motor neuron subtypes (e.g., chondrolectin, estrogen-related receptor β and matrix metalloproteinase-9), but were unsuccessful. Muscle fibre and α motor neuron subtypes are closely related (39), and denervation can induce changes in motor unit identity (40); thus, by staining for myosin heavy chain (MHC) isoforms, we assessed the proportions of muscle fibre types in transverse sections of tibialis anterior muscles from 3 month-old wild-type and *Gars*^{C201R/+} mice (Supplemental Figure 5A). The tibialis anterior is known to display weakness and NMJ denervation at this age in this mouse strain (30). Type I fibres are the smallest and largely innervated by slow-twitch fatigue-resistant α motor neurons, whereas type II fibres are larger (increasing in size from IIa to IIx to IIb) and innervated by either fast-twitch fatigue-resistant (IIa/IIx) or fast-twitch fatiguable α motor neurons (IIb/IIx) (39).

We first determined that there were no differences between genotypes in the cross-sectional area of each myofiber subtype (Supplemental Figure 5B). We then showed that *Gars*^{C201R/+}

mice possess a lower percentage of MHC type IIb fibres and higher percentages of type IIx and type I fibres (Supplemental Figure 5C). However, by assessing the number of each fibre subtype, we identified that compared to wild-type, *Gars*^{C201R/+} muscles possess fewer type IIb fibres, similar numbers of type IIx fibres and more type I fibres (Supplemental Figure 5D); this results in a reduction in total myofiber number (Supplemental Figure 5E), which may deplete target-derived BDNF, exacerbating transport disruption.

Together, these data suggest that the tibialis anterior muscle of CMT2D mice displays a selective loss of large fast-twitch fatiguable α motor neurons and an increase in slow-twitch fatigue-resistant α motor neurons, likely due to a combination of fibre loss and subtype switching. Along with impaired neurotrophin signalling and dampened protein synthesis (41), this change in subtype proportions will likely contribute to the reduced size of CMT2D lumbar motor neurons.

Dissimilar to the lumbar motor neurons, upper body-innervating thoracic motor neurons of CMT2D mice showed no difference to control in CREB phosphorylation (Supplemental Figure 4, C-E). Like in the lumbar spinal cord, no loss of thoracic motor neurons was observed (Supplemental Figure 4F); however, in contrast to the lumbar motor neurons, soma and nucleus size were unaffected in thoracic motor neurons (Supplemental Figure 4G), indicating that the reduced area of lumbar motor neurons is not caused by the smaller size of CMT2D mice. This was further corroborated by analysing motor neuron areas relative to the area of the spinal cord section from which they were measured (data not shown). These collective data show that impairments in CREB activation and motor neuron size are selective and linked to the more severe motor and sensory pathology observed in CMT2D hindlimbs (21, 33).

Reduced endosome adaptor protein levels correlate with CMT2D pathology

Having assessed distal (NMJs) and proximal (cell bodies) subcellular regions of the motor neuron, we wanted to address whether any axonal disturbances in endosomal transport proteins could be observed in CMT2D mice. Adaptor proteins are required for connecting the retrograde motor cytoplasmic dynein to specific cargoes, such as TrkB-positive signalling endosomes, and their reduced availability could contribute to impaired axonal trafficking (42). To determine whether CMT2D mice display differences in endosome adaptors, we probed lysates from hindlimb-innervating sciatic nerves and forelimb-innervating median and ulnar nerves for the established cytoplasmic dynein binding proteins Snapin (43), Hook1 (44) and RILP (45). In agreement with the peripheral nerve pathology, we found that levels of monomeric Snapin (mSnapin) and RILP are lower in *Gars*^{C201R/+} sciatic nerve (Figure 3A), whereas there was no significant downregulation in median and ulnar nerves (Figure 3B), greatly decreasing the possibility of a systemic phenotype.

Endosome axonal transport is unaffected in forelimb-innervating nerves

Given that neuromuscular pathology is comparatively mild or absent in CMT2D forelimbs, we adapted our *in vivo* imaging approach to perform intravital analysis of signalling endosome transport in median and ulnar nerves (46). To target muscles innervated by motor neurons within these nerves, H_CT-555 was injected into the forepaw of wild-type and *Gars*^{C201R/+} mice at 3 months, representing an age at which axonal transport has been defective in CMT2D sciatic nerves for at least two months (Figure 1, C and D). Consistent with the neuropathology, there was no difference in signalling endosome dynamics between wild-type and *Gars*^{C201R/+} median and ulnar nerves (Figure 3, C-F). Moreover, endosome speeds of both genotypes were similar to those observed in 3 month-old wild-type sciatic nerves (wild-type sciatic 2.51 ± 0.2 ; wild-

type median/ulnar 2.37 ± 0.1 ; *Gars*^{C201R/+} median/ulnar 2.36 ± 0.1 ; $P = 0.699$ one-way ANOVA).

In summary, differences in pro-survival FL-TrkB between muscles strictly correlate with NMJ denervation in CMT2D mice. Furthermore, activation of the pro-survival and BDNF-dependent transcription factor CREB is stunted in *Gars*^{C201R/+} lumbar motor neurons, and this effect is specific, as it is not observed in thoracic motor neurons. Finally, endosome adaptor proteins critical to axonal transport are selectively downregulated in hindlimb, but not forelimb, innervating nerves, correlating with disturbances in endosome axonal transport in motor neurons in vivo. Given the importance of neurotrophin signalling to neuronal health and survival, these phenotypes likely contribute to the selective motor neuropathology observed in CMT2D.

Altering BDNF-TrkB signalling at healthy nerve terminals impairs endosome transport

Our data thus far are commensurate with aberrant mutant GlyRS-TrkB interactions at peripheral nerve terminals underlying impaired BDNF signalling in CMT2D. To test whether dampening BDNF-TrkB signalling at the nerve-muscle interface disrupts endosome transport and thus explains the neuropathic phenotype we observed in vivo, we co-injected HcT-555 with anti-BDNF antibodies into wild-type and *Gars*^{C201R/+} leg muscles. This was performed to sequester extracellular BDNF in the vicinity of peripheral nerve terminals and impede signalling through TrkB. Whilst immunoglobulin Y (IgY) control antibodies had no effect on axonal transport, restricting the availability of BDNF at wild-type NMJs resulted in a slow-down in signalling endosome transport, akin to the CMT2D defect (Figure 4, A and B). This indicates that BDNF availability at nerve terminals modulates axonal endosome transport dynamics. In contrast, anti-BDNF did not exacerbate endosome trafficking in *Gars*^{C201R/+}

mutants (Figure 4, A and B), which is consistent with perturbations in BDNF-TrkB signalling being the cause of the CMT2D axonal transport disruption.

To corroborate this, we co-injected wild-type muscles with HcT-555 and the selective and peripherally-restricted pan-Trk inhibitor, PF-06273340 (47). PF-06273340 binds to the extracellular domain of Trk receptors, thus restricting their interactions with neurotrophins. Trk inhibition slowed transport in healthy axons of wild-type animals (Figure 4, C and D), thus confirming a role for local Trk signalling in maintaining healthy axonal endosome trafficking.

There are several key signalling pathways downstream of BDNF-TrkB, including AKT, PLC γ 1 and ERK1/2 (22). To identify which, if any, of these are affected in CMT2D nerves, we extracted sciatic nerves from wild-type and *Gars*^{C201R/+} mice that received injections of vehicle or BDNF into the tibialis anterior and gastrocnemius muscles. It should be noted that motor neurons innervating these muscles constitute only a small fraction of the total sciatic nerve, hence BDNF-induced signalling changes triggered by this experimental procedure are predicted to be small. Probing for total and phosphorylated forms of key proteins in each of these signalling nodes, we identified that under basal conditions, only ERK1/2 activation was impaired in CMT2D nerves (Supplemental Figure 6), suggesting that this pathway may contribute to the *Gars*^{C201R/+} transport defect. Consistent with this (48), we showed that BDNF treatment enhanced ERK1/2 phosphorylation in both genotypes, but to a lesser extent in CMT2D mice. This was again selective, since AKT and PLC γ 1 phosphorylation were unchanged.

To test the hypothesis that impaired ERK1/2 activation contributes to compromised endosome transport, we injected the ERK1/2 inhibitor, refametinib (49), into wild-type muscles and

analysed the axonal transport of signalling endosomes. We found that ERK1/2 inhibition caused a stark disruption in trafficking (Figure 4, E and F), confirming the importance of ERK1/2 activation for maintaining healthy speeds of signalling endosome axonal transport in motor neurons *in vivo*.

To identify whether a local, rather than a systemic, effect of ERK1/2 inhibition at distal motor neuron terminals causes transport disruption, we co-administered HcT-555 with refametinib into hindlimb muscles on one side of the body and HcT-555 with vehicle control into the other (Supplemental Figure 7A), followed by axonal transport measurement. We found that local ERK1/2 inhibition within the injected muscle perturbs transport, but has no impact on endosome trafficking in contralateral sciatic nerves (Supplemental Figure 7, B-E). This result was corroborated by performing similar experiments using IgY control and anti-BDNF antibodies (Supplemental Figure 7, F-J).

These data in wild-type mice indicate that perturbation of BDNF-TrkB signalling via ERK1/2 inhibition at peripheral nerve terminals is sufficient to disrupt signalling endosome transport in otherwise healthy motor axons, thus replicating the CMT2D phenotype.

*Application of recombinant BDNF to CMT2D muscles restores *in vivo* axonal transport*

BDNF appears crucial for maintaining axonal dynamics of signalling endosomes generated at peripheral nerve terminals. We thus performed intramuscular injections of 25 ng recombinant mBDNF and re-assessed endosome transport in mutant *Gars* mice. In line with the restoration of ERK1/2 activation in CMT2D sciatic nerves (Supplemental Figure 6), BDNF treatment rescued *in vivo* axonal transport in *Gars*^{C201R/+} mice to wild-type levels at both 1 and 3 months of age (Figure 5, A and B). Moreover, we showed that injection of BDNF at the late stage of

13-14 months also enhanced *Gars*^{C201R/+} endosome transport, whereas it had no effect in wild-type mice (Supplemental Figure 8). Importantly, the trafficking deficit observed in 1 month-old *Gars*^{ΔETAQ/+} mice was also fully corrected upon intramuscular injection of BDNF (Figure 5C). We then administered BDNF to ascertain whether it could restrict the non-cell autonomous disruption of endosome transport caused by human mutant GlyRS. Doing so reversed the transport deficit (Figure 5, D and E), suggesting that boosting BDNF levels can overcome the negative, non-cell autonomous effect on axonal endosome mobility caused by mutant GlyRS at peripheral nerve terminals.

To confirm that the Trk receptors are responsible for the observed amelioration of transport caused by BDNF, *Gars*^{C201R/+} mice were treated with a combination of BDNF and pan-Trk inhibitor. The addition of PF-06273340 to the intramuscular injection abolished the BDNF-mediated rescue of *Gars*^{C201R/+} endosome transport (Figure 6, A-D), demonstrating that BDNF regulates axonal trafficking through canonical Trk signalling.

To further assess selectivity of this rescue, we individually injected several different recombinant growth factors into muscles of *Gars*^{C201R/+} mice. We chose vascular endothelial growth factor 165 (VEGF₁₆₅) because it is critical to motor neuron survival and signals through neuropilin-1 (50), a binding partner of mutant, but not wild-type, GlyRS (11). We also tested NT-3 and NT-4, which bind to TrkC and TrkB, respectively. We did not evaluate the impact of NGF, because TrkA is not expressed at the NMJ (37), and NGF administration can cause rapid and direct sensitisation of nociceptors (51), which would cause unnecessary suffering to the treated mice. Unlike BDNF, injection of NT-3 and NT-4 had no effect on *Gars*^{C201R/+} transport, whereas VEGF₁₆₅ exacerbated the CMT2D impairment (Figure 6, E-H), suggesting

that the action of BDNF on axonal endosome trafficking in neuropathic mice is specific and does not extend to all neurotrophic factors.

To evaluate the temporal impact of BDNF treatment, we imaged in vivo axonal transport of *Gars*^{C201R/+} mutants 24 h post-BDNF injection. No difference from vehicle-injected mice was observed (Figure 6, I-L), indicating that the benefit driven by the injected recombinant BDNF is short-lived.

Together, these data indicate that the rescue of endosome axonal trafficking defects by BDNF is specific and occurs via its canonical Trk receptor.

Muscle-specific BDNF gene therapy rescues CMT2D axonal transport

To improve the potential for translation of this discovery, we designed an adeno-associated virus (AAV) to constitutively express BDNF in muscle, thereby avoiding the side-effects of systemic BDNF upregulation (52). AAV serotype 8 (AAV8), which displays efficient muscle tropism (53), was combined with the muscle-specific promoter *tMCK* (54). First, an AAV8-*tMCK*-eGFP control virus was shown to selectively express in skeletal muscles and the heart when injected into the peritoneum of wild-type P2 pups (Supplemental Figure 9, A-C). After confirming muscle-selectivity of our virus/promoter combination, we performed unilateral injections of AAV8-*tMCK*-BDNF into the tibialis anterior and gastrocnemius muscles of P11 wild-type mice at several different doses (Supplemental Figure 9, D-G). At all concentrations tested, we observed a similar robust upregulation of proBDNF in the injected muscles, as well as the adjacent extensor digitorum longus, indicating that our lowest tested dose of 5.0×10^{10} vector genomes (vg) per muscle was sufficient to maximally express the transgene.

Having identified a well-tolerated virus dose that drives observable increases in proBDNF expression, we performed bilateral injections of AAV8-tMCK-BDNF or AAV8-tMCK-eGFP targeting lumbrical/FDB, tibialis anterior and gastrocnemius muscles of P11 wild-type and *Gars*^{C201R/+} mice to analyse the impact on sciatic nerve axonal transport. Successful transgene expression was confirmed in muscles from all treated animals at the experimental end-stage of P38-P41 (Supplemental Figure 9, H-J). Similar to the acute improvement driven by intramuscular injections of recombinant mBDNF, AAV8-tMCK-BDNF successfully rescued axonal transport of endosomes in *Gars*^{C201R/+} mice to wild-type levels (Figure 7, A and B). Moreover, the increased BDNF availability also resulted in faster endosome speeds in wild-type mice compared to treatment with AAV8-tMCK-eGFP (Figure 7B). As this increase was not observed in wild-type animals upon intramuscular injection of BDNF (Supplemental Figure 8), this suggests that the amount of BDNF and length of exposure can impact transport in healthy peripheral nerves. Consistent with the increased transport in CMT2D mice, AAV8-tMCK-BDNF also caused a selective increase in availability of the motor adaptor protein Snapin, but not Hook1 or RILP, in mutant sciatic nerves (Figure 7, C-E).

In conclusion, we have generated and tested a novel muscle-specific gene therapy for CMT2D that effectively augments proBDNF and mBDNF levels in a tissue-specific manner, resulting in a robust correction of the impaired in vivo axonal transport of signalling endosomes.

Discussion

Impaired axonal transport has been reported in diverse neurological diseases, yet it is debated as to whether this is a primary cause of neuropathology or simply a secondary consequence of a degenerating nervous system (55). The data presented here are consistent with early disruption of signalling endosome transport being one of the contributing factors to CMT2D; however, it remains to be experimentally shown whether this pathomechanism drives disease progression. Through intravital imaging of intact sciatic nerves, we report, for the first time, that axonal transport is impaired in mammalian peripheral neuropathy in vivo. The disturbance manifests by one month of age in two different mouse models of CMT2D, coinciding with degeneration of the NMJ in the most severely affected muscles (31). As the uptake of the retrograde axonal transport probe into axonal endosomes is dependent on the presence of motor nerve terminals, these identified trafficking impairments will under-represent the severity of disruption, because NMJs in hindlimb muscles become progressively denervated in mutant *Gars* mice (31, 33). This feature of our assay also provides an explanation as to why the transport deficit does not appear to worsen with age, since at each timepoint we can only assess transport in motor neurons that have a functional connection with the muscle. Further supporting a role in neuropathy, in vivo endosome trafficking in median and ulnar nerve axons was unaffected in *Gars*^{C201R/+}, even at a later disease stage, which is a result consistent with the observation that motor function and NMJ integrity are relatively spared in CMT2D mouse forelimbs (33). For now, it remains to be determined whether transport of other cargoes, such as mitochondria, is disrupted in CMT2D mice in vivo. This is a particularly intriguing possibility, given the recent discovery that endosomes serve as platforms for translation to maintain mitochondrial function within axons (56, 57).

We hypothesise that interactions between mutant GlyRS and the extracellular domain of TrkB cause the endosome trafficking impairment in motor axons, dampening neurotrophin signalling, which may be contributing to the slowly progressive peripheral nerve degeneration characteristic of CMT (Figure 8). After confirming that GlyRS^{ΔETAQ} aberrantly associates with TrkB, we showed that intramuscular injection of two different CMT2D-causing mutants, but not wild-type, GlyRS impair endosome transport in otherwise healthy motor axons of wild-type mice. This is consistent with mutant GlyRS being the pathological driver at the nerve-muscle interface, altering the regulation of axonal transport of signalling endosomes. Enabling this possibility, GlyRS is secreted from several different cell types including muscles (11, 12, 58, 59), and accumulates at the NMJ prior to degeneration in a *Drosophila* CMT2D model (12, 15). TrkB is the principal neurotrophin receptor at the mammalian NMJ, thus it is perhaps no coincidence that its disruption via dominant-negative manipulation (37) or heterozygous knockout (60) disturbs NMJ integrity similar to what is observed in mutant *Gars* mice. Further supporting that impaired neurotrophin signalling plays a role in selective peripheral nerve pathology (33), we found a correlation between the availability of FL-TrkB in wild-type muscles and NMJ denervation in CMT2D mice; muscles displaying the greatest NMJ pathology possessed the highest levels of FL-TrkB and the lowest levels of BDNF at the synapse, which would provide ideal conditions for extracellular mutant GlyRS to impact physiological BDNF-TrkB signalling.

We also evaluated potential BDNF-TrkB signalling impairments by assessing activation of the transcription factor CREB in both lumbar and thoracic motor neurons, as well as levels of endosome adaptor proteins in sciatic and median/ulnar nerves. These anatomical studies were aimed to assess whether CMT2D mouse phenotypes are systemic or are region-specific. We discovered that CREB phosphorylation was selectively disrupted in nuclei of lumbar motor

neurons, and that levels of the endosome adaptors Snapin and RILP were reduced in sciatic nerves alone. Therefore, NMJ degeneration (33), impaired motor function (33), sensory disruption (13, 21), reduced FL-TrkB availability, dampened CREB signalling and diminished endosome adaptor levels are all associated with the deficit in signalling endosome trafficking, suggesting that compromised BDNF-TrkB signalling and disturbed axonal transport are integral features of CMT2D pathology.

While impaired protein translation in motor neuron cell bodies may be a primary driver of disease (17, 18), the extent of these deficits does not change across spinal levels, dissociating the disruption of protein translation from neuropathology. It is therefore possible that aberrant protein interactions at nerve terminals, as well as within axons (14), provide an additional layer of complexity to disease pathogenesis, whereby disturbances in axonal transport drive the selective peripheral degeneration of motor and sensory neurons. In addition, the observation that mutant GlyRS aberrantly associates with HDAC6 to decrease microtubule acetylation (14), suggests that multiple disease mechanisms converge to drive CMT2D pathology.

Acutely increasing the availability of mBDNF in muscle consistently rescued in vivo endosome trafficking in mutant *Gars* axons through its canonical receptor TrkB. Moreover, commensurate with a direct competition between the two proteins at axon terminals, mBDNF injected into wild-type muscles was able to overcome the negative impact of mutant GlyRS on endosome transport. Treatment of CMT2D mice with VEGF₁₆₅, NT-3 or NT-4 did not correct the phenotype, indicating that this effect is specific, and not a generic feature of factors promoting neuron survival. Further supporting the specificity of this rescue, we recently demonstrated that mBDNF was incapable of improving endosome axonal transport defects in SOD1^{G93A} mice modelling amyotrophic lateral sclerosis (61). Crucially, dampening BDNF-

TrkB signalling in wild-type muscles, either through BDNF sequestration or pan-Trk inhibition, slowed endosome transport in otherwise healthy neurons. This confirms a role for the BDNF-TrkB signalling axis in regulating axonal transport of signalling endosomes.

It is currently unknown how perturbations in BDNF-TrkB signalling impair axonal endosome speeds in CMT2D; several non-mutually exclusive possibilities will be addressed in follow-up studies. For instance, BDNF has been shown to facilitate the recruitment of the intermediate chain of cytoplasmic dynein to signalling endosomes, but not mitochondria, in an ERK1/2 phosphorylation-dependent manner (62). Similarly, increased local translation of cytoplasmic dynein and its adaptors (e.g., Snapin) is driven by peripheral neurotrophin signalling (63, 64), perhaps through re-distribution of ribosomes at the axon terminal (65). The blockade of BDNF signalling through TrkB by mutant GlyRS may thus reduce association of the retrograde motor with signalling endosomes, which would be consistent with mutant *Gars* sciatic nerves displaying reduced ERK1/2 phosphorylation and endosome adaptor levels, both of which are rescued by intramuscular administration of BDNF. As multiple dynein motors can interact with and direct the transport of retrograde cargoes for faster movement (66, 67), it is possible that fewer dynein complexes are bound to endosomes in CMT2D axons causing the slowdown of signalling endosomes.

NT-3 was recently shown to alleviate neuropathology in CMT2D mice when expressed in muscles using AAV1 (68). Similarly, lentivirus-mediated delivery of VEGF₁₆₅ into muscle also partially improved mutant *Gars* motor function (11). These findings, coupled with our data showing that neither growth factor was able to acutely improve axonal transport in CMT2D mice, suggest that restoration of axonal endosome trafficking is not required for modest improvements in neuropathy and that CMT2D results from impairments in several distinct

pathways and processes. The beneficial effect observed with VEGF₁₆₅ and NT3 in the absence of axonal transport restoration may be due to systemic effects or modulation of non-neuronal components of the motor unit, e.g., muscles or Schwann cells. Alternatively, the lack of transport correction with NT-3, as well as NT-4, may be due to the dose being too low or treatment too short, such that higher levels delivered for prolonged periods may improve endosome trafficking – as is observed in this study with BDNF treatment of wild-type mice. NT-3 has previously been shown to bind and activate TrkB during sensory neuron development (69), albeit with a lower affinity than BDNF (70, 71), which would be consistent with this hypothesis; however, it is unknown whether the promiscuity of NT-3 in the motor nervous system is preserved in adulthood. On the other hand, NT-4 has a higher affinity for TrkB than NT-3 and even BDNF (71), suggesting that NT-3 and NT-4 simply do not regulate endosome axonal transport. This is consistent with the finding that NT-4 binding to TrkB elicits different effects compared to BDNF (72, 73).

The same lack of response cannot be said for VEGF₁₆₅, which in our hands exacerbated the CMT2D transport defect. We hypothesise that the further reduction in endosome speed may be due to competition between VEGF₁₆₅ and the alternative neuropilin-1 ligand semaphorin 3A, which is known to facilitate axonal transport in vitro (74); by increasing VEGF₁₆₅ availability, semaphorin 3A may be displaced and no longer be able to affect trafficking. Irrespective of the lack of transport improvement, neither NT-3 nor VEGF₁₆₅ treatment resulted in complete rescue of neuropathy (11, 68), indicating that alternative therapeutic strategies are required to treat this disease, and that perhaps a combinatorial approach including BDNF may produce the greatest phenotypic improvements.

The restorative effect of BDNF on transport is short-lived, as it was not observed 24 h post-injection. We therefore developed a gene therapy strategy to constitutively boost mBDNF levels in muscles. This resulted in complete correction of in vivo axonal transport in CMT2D mice up to one month post-treatment, which was associated with an increase in levels of the key endosome adaptor protein Snapin. Further work is underway to determine the impact of enhancing the long-term availability of BDNF in muscles as a therapeutic strategy for CMT2D, taking into consideration that genetic depletion and overexpression of BDNF in muscle have been shown to impact fibre type proportions and muscle function (75).

The approach of augmenting BDNF in muscles may be beneficial to trafficking disruption in other CMT subtypes. Mutations in several different aminoacyl-tRNA synthetase genes cause CMT (8), several of which are associated with structural relaxation of the encoded synthetase and protein mis-interactions (76-78), and share pathomechanistic similarities (17, 79, 80). Additionally, CMT is caused by mutations in genes linked to axonal transport (55), including in the critical signalling endosome protein Rab7 (81), and disruptions in transport are highly prevalent in neuropathy. Moreover, myelinating Schwann cells provide a major source of trophic support to axons (82), and are lost or damaged in type 1/demyelinating CMTs. Thus, boosting the availability of specific growth factors such as BDNF in a spatial and time-dependent manner (Figure 8) may address the therapeutic needs of one of the most widespread forms of human neuropathy.

Materials & Methods

Animals

Mice were maintained under a 12 h light/dark cycle at constant room temperature ($\approx 21^{\circ}\text{C}$) with ad libitum water and food (Teklad global 18% protein rodent diet, Envigo, 2018C). Cages were enriched with nesting material, plastic/cardboard tubes and wooden chew sticks as standard. *Gars*^{C201R/+} (RRID: MGI 3849420) and *Gars* ^{Δ ETAQ/+} mice (provided by Robert W. Burgess, The Jackson Laboratory, Bar Harbor, ME) were maintained as heterozygous (male) \times wild-type (female) breeding pairs on a C57BL/6J background. *Gars*^{C201R/+} mice were genotyped as previously described (30). *Gars* ^{Δ ETAQ/+} mice were genotyped under standard conditions using forward primer 5'-GGTAGTTTACTTGTAACAGGC-3' and reverse primer 5'-TTTCCAATCTGGGCAGCAGC-3'. Both female and male mice were used (Supplemental Table 1) because no differences in pathology have been observed between sexes. Moreover, axonal signalling endosome dynamics do not differ between female and male wild-type C57BL/6J mice (29). Animals sacrificed for 0.5, 1, 3 and 13-14 month timepoints were 15-16, 28-38, 87-104 and 391-426 days old, respectively (Supplemental Table 1). Additional timepoints were used and are detailed where appropriate. Post-natal day 1 (P1) was defined as the day after a litter was first found.

In vivo axonal transport imaging

Live imaging of signalling endosome axonal transport was performed using an atoxic binding fragment of tetanus neurotoxin (H_CT), as described (26-28). For imaging transport in sciatic nerves, H_CT-555 was injected into the right lateral gastrocnemius and tibialis anterior muscles of isoflurane-anaesthetised mice (Figure 1A and 1B). Alternatively, H_CT-555 was administered into the left forepaw (targeting lumbrical muscles, amongst others) for assessment of transport in median and ulnar nerves, as described (46). A 10 μl , 26-gauge Hamilton syringe

(Sigma, 20779) or a pulled, glass micropipette (Drummond Scientific, 5-000-1001-X10) was used for intramuscular injections, whereas micropipettes alone were used for forepaw injections. Muscles of the left leg were also injected in experiments designed to assess the site of action of transport-altering treatments; the side of administration for vehicles and treatments were alternated between mice to limit biases. 5-7 μg of H_CT-555 in phosphate-buffered saline (PBS) was injected per muscle or forepaw in a volume of $\approx 1.5\text{-}2\ \mu\text{l}$, before allowing animals to recover from the anaesthesia. 4-8 h post-injection (unless otherwise stated), nerves were exposed under terminal anaesthesia and imaged on an inverted LSM780 laser scanning microscope (Zeiss) within an environmental chamber pre-warmed to 37°C. Images (1024x1024, 1% laser power) were acquired every $\approx 3\ \text{s}$ using a 63 \times Plan-Apochromat oil immersion objective lens (Zeiss) at 100 \times digital zoom. Faster frame rates have been used to determine that endosome fission/fusion events have no clear impact on transport analyses.

Intramuscular injections

Different substances were pre-mixed and co-administered with H_CT-555 into the gastrocnemius and tibialis anterior muscles: 25 ng recombinant human glycyl-tRNA synthetase (WT/G240R/L129P), 250 ng chicken IgY anti-brain-derived neurotrophic factor (BDNF, R&D Systems, AF248), 250 ng chicken control IgY (R&D Systems, AB-101-C), 13 or 50 nM PF-06273340 (Sigma, PZ0254) with 25 ng recombinant human BDNF (Peprotech, 450-02), dimethyl sulfoxide (DMSO, Sigma, D1435), 50 nM refametinib (Generon, HY-14691), 25 ng recombinant human BDNF, 25 ng recombinant human VEGF₁₆₅ (Peprotech, 100-20), 25 ng recombinant human NT-3 (Peprotech, 450-03) and 25 ng recombinant human NT-4 (Peprotech, 450-04). All amounts are per muscle. Recombinant non-tagged GlyRS was purified as previously described (11). Drug concentrations were selected based on reported IC₅₀ values: 6/2/1 nM PF-06273340 (TrkA/B/C) (47) and 19/47 nM refametinib (ERK1/ERK2)

(49). For western blot analysis of sciatic nerves, similar injections of vehicle or BDNF were performed (where indicated) without the addition of H_cT. Sciatic nerves were dissected 6 h post-injection.

In vivo axonal transport analysis

Confocal .czi files were uploaded to ImageJ (<http://rsb.info.nih.gov/ij/>) and endosome dynamics manually tracked using the TrackMate plugin (83). Only endosomes that could be tracked for ≥ 5 consecutive frames were analysed. Endosomes that paused for ≥ 10 consecutive frames or moved solely anterogradely were excluded. The tracked fraction of endosomes varied depending on how densely populated with fluorescent organelles the axon was (≈ 1 -50%); nevertheless, endosomes were selected for assessment as they entered the field of view, without prior observation of their movements. Endosome frame-to-frame speeds are presented in frequency histograms (581 ± 16 movements per animal, $n = 248$ across the study). Endosomes usually take 5-15 frames to traverse the field of view, with rare slower endosomes reaching ≈ 30 frames. To determine mean endosome speed per animal, individual endosome speeds were averaged (60.1 ± 1.6 endosomes per animal, $n = 248$ across the study). An endosome was determined to have paused if it remained stationary (within $< 0.1 \mu\text{m}$) for two consecutive frames. The ‘% time paused’ is a calculation of the time all tracked endosomes remained stationary. The ‘% pausing endosomes’ defines the proportion of endosomes that paused at least once. A minimum of ten endosomes from at least three individual axons was assessed per animal 5-120 min from initiating anaesthesia (most videos were recorded within 45 min). Thick axons were selected to increase the likelihood of imaging motor, rather than sensory, neurons (29).

Protein extraction and western blotting

Tissues for western blotting were dissected from PBS-perfused and non-perfused mice. Lumbar level 1 (L1) to L5 DRG and wholemount muscles were excised as previously outlined (84-88). Proteins were extracted from tissues as described (13), except that ground tissue samples were incubated on ice in NP-40 lysis buffer (1% [w/v] NP-40, 50 mM NaCl, 50 mM Tris-HCl [pH 8.0]) for 1-2 h rather than 30 min. Several FDB, forelimb lumbrical and hindlimb lumbrical muscles from each side of the body were combined to ensure sufficient protein was extracted. Western blotting was performed following published protocols (13), with primary and secondary antibodies detailed in Supplemental Table 2 and 3, respectively. 20 µg of protein from DRG, 20-40 µg from muscles and 40 µg from sciatic nerves were loaded per well. Densitometric analysis was performed as previously described (89), using GAPDH or total protein stained with 0.1% Coomassie Brilliant Blue R-250 (Thermo Fisher, 20278) as the loading control (90). Hook1 bands at ≈110 kDa and ≈90 kDa were quantified in sciatic and median/ulnar nerves, respectively. Protein phosphorylation levels were calculated relative to the total protein (e.g., p-ERK1/2 relative to ERK1/2). Wholemount muscle protein levels were correlated with previously published *Gars*^{C201R/+} NMJ denervation (*n* = 6-8) (33).

In vitro pull-down assay

The recombinant human TrkB-Fc chimera, consisting of the extracellular domain of TrkB (Cys32-His430; R&D Systems, 688-TK), and control human IgG-Fc (110-HG, R&D Systems) were bound to Dynabeads Protein G (Thermo Fisher, 10003D). Whole cell lysates from NSC-34 cells (CELLutions Biosystems, CLU140) transfected with plasmids encoding human GlyRS^{WT}-V5, GlyRS^{ΔETAQ}-V5, or vector control were added to the beads and incubated overnight at 4°C. The beads were washed twice with wash buffer (25 mM Tris-HCl [pH 7.4], 150 mM NaCl, 1 mM EDTA, 1% [w/v] NP-40, 5% [w/v] glycerol) and treated with sodium dodecyl sulphate loading buffer to elute bound proteins. TrkB-Fc and GlyRS-V5 were analysed

by western blot using anti-human IgG Fc (horseradish peroxidase, pre-adsorbed) and anti-V5 antibodies, respectively (Supplemental Table 2).

Spinal cord dissections and staining

Spinal cords were dissected from 3 month-old mice transcardially-perfused with 4% (w/v) methanol-free formaldehyde (Thermo Fisher, 28908) in PBS. Samples were then post-fixed for 24-48 h at room temperature, before washing with PBS and equilibrating in 30% (w/v) sucrose (Sigma, S7903) in PBS for 24-72 h at 4°C. Using the lumbar enlargement to guide, the L1 to L5 and thoracic level 4 (T4) to T8 segments of the cord were collected and frozen in Tissue-Tek O.C.T. (Sakura Finetek, 4583). Samples were kept at -80°C, until 30 µm sections were cut with an OTF Cryostat (Bright Instruments) and collected onto five parallel series of polysine-coated slides (VWR, 631-0107). For staining, sections were permeabilized for 10 min using 0.3% (w/v) Triton X-100 (Sigma, T8787) in PBS before blocking for 30 min in permeabilization buffer containing 5% (w/v) bovine serum albumin (BSA, Sigma, A2153), and probing overnight at 4°C with primary antibodies (Supplemental Table 2) in block solution. The following day, slides were washed three times with PBS, before incubating with secondary antibodies (Supplemental Table 3). Sections were then washed three times with PBS, mounted in Fluoromount-G (Thermo Fisher, 00-4958-02) and covered with 22 × 50 mm cover glass (VWR, 631-0137). Slides were kept at 4°C to set before imaging.

Immunofluorescent analysis of spinal cords

Motor neurons from L3-L5 and T4-6 spinal cord levels were analysed. Motor neuron counts were performed by assessing the number of ChAT-positive neurons per ventral horn and calculating an average across ten slides. Cell body areas were measured from maximum intensity-projected z-stack images by drawing around the circumferences of ChAT staining

using the freehand tool. 58.0 ± 2.3 cell bodies were assessed per spinal cord segment. To determine average CREB and p-CREB staining intensities, maximum intensity-projected z-stack images were manually thresholded, smoothed and then TDP-43-positive nuclear masks created. These were then used to measure the average fluorescence intensity of CREB (53.8 ± 3.6 nuclei per mouse) and p-CREB (50.0 ± 6.0 nuclei per mouse) in nuclei of ChAT-positive neurons to obtain mean values for each mouse. Secondary antibody only-treated sections were also processed for each animal (17.8 ± 0.8 nuclei per mouse) to generate individual mean values that were subtracted from the mean fluorescence intensities of each mouse to remove background. The resulting values were then used to calculate relative nuclear fluorescence intensities for both CREB and p-CREB separately. To calculate the ratio of p-CREB to CREB, the relative p-CREB value for each mouse was divided by the relative CREB value and multiplied by 100. The areas of motor neuron nuclei were also measured using TDP-43 masks (114.3 ± 6.0 per mouse). TDP-43 was used rather than DAPI, because the fluorescence intensity of the latter is low in motor neurons compared with other cells of the spinal cord, whereas TDP-43 clearly highlights motor nuclei (Supplemental Figure 4C). All sections probed with anti-CREB or anti-p-CREB were processed and analysed in parallel with fluorescence values calculated relative to wild-type.

Immunofluorescent analysis of NMJs

Dissected wholemount muscles were stained as described in detail elsewhere (85). All primary and secondary antibodies are detailed in Supplemental Table 2 and 3, respectively. AlexaFluor647 α -bungarotoxin (α -BTX, Life Technologies, B35450) was used at 1:1,000 to identify post-synaptic acetylcholine receptors. Relative BDNF levels at the NMJ were measured from maximum intensity-projected z-stack images by drawing around the circumferences of α -BTX staining using the freehand tool. Levels of BDNF fluorescence

within the α -BTX mask were assessed using the Integrated Density function in ImageJ. The average background fluorescence was subtracted from mean values per muscle and then expressed as a percentage relative to the ETA muscle. An average of 20 ± 1.2 NMJs per muscle were analysed across three mice. GFP fluorescence was imaged in non-fixed wholemount muscles of AAV-treated mice.

Muscle fibre typing

Tibialis anterior muscles were dissected from PBS-perfused mice and immediately frozen in Tissue-Tek O.C.T. Samples were kept at -80°C , until $30\ \mu\text{m}$ transverse sections were cut with an OTF cryostat and collected onto polysine-coated slides that were then stored at -20°C . Muscle sections were stained with antibodies against MHC isoforms and laminin (Supplemental Table 2 and 3), as previously described (88). Four sections per muscle were imaged at approximately equal positions throughout the muscle. Fibre types and cross-section areas were analysed using the MyoSight plugin for ImageJ (91). Data were averaged across the four sections to get values per animal.

AAV8-tMCK virus production

Self-complementary AAV (scAAV) expression plasmids were created by OXGENE (Oxford, UK). A 745 bp *tMCK* promoter sequence was adapted from those reported by Rodino-Klapac *et al.* (92) and <https://www.addgene.org/105556/> (Supplemental Table 4). The 744 bp coding sequence of human pre-proBDNF was sourced from the European Nucleotide Archive (<https://www.ebi.ac.uk/ena/browser/view/AAA69805>) (Supplemental Table 4). pSF-scAAV-tMCK-eGFP and pSF-scAAV-tMCK-BDNF plasmids were packaged into AAV serotype 8 particles by Charles River Laboratories (previously Vigene Biosciences).

AAV8-tMCK injections

Viruses were kept at -80°C in 0.01% (v/v) pluronic F68 surfactant in PBS and diluted in sterile PBS. Viruses were freeze-thawed a maximum of two times before injection. For assessing tissue-specificity of expression, intraperitoneal AAV8-tMCK-eGFP injections were performed on P2 pups using a Hamilton syringe connected to a 30 gauge needle cannula; 7.0×10^{10} viral genomes (vg) were injected per mouse in a volume of 5 μl . To identify an appropriate dose, intramuscular AAV8 injections were performed with pulled, glass micropipettes directly through the skin into the gastrocnemius and tibialis anterior of P11 pups under isoflurane-induced anaesthesia; 5.0×10^{10} , 1.0×10^{11} or 2.0×10^{11} vg were unilaterally injected into each muscle in a volume of 2.5 μl . To evaluate gene therapy efficacy, bilateral AAV injections targeting the hindpaw lumbrical/FDB, gastrocnemius and tibialis anterior muscles of P11 pups were performed under anaesthesia; each injection of 2.5 μl contained 5.0×10^{10} vg (equating to 3.0×10^{11} vg per animal).

Statistics

Data were assumed to be normally distributed unless evidence to the contrary were provided by the Kolmogorov-Smirnov test for normality, while equal variance between groups was assumed. Normally-distributed data were statistically analysed using unpaired and paired *t*-tests or one-way and two-way analysis of variance (ANOVA) tests followed by Šídák's multiple comparisons test. Non-normally distributed data were analysed using Mann-Whitney *U* tests or Kruskal-Wallis tests followed by Dunn's multiple comparisons tests. Correlation was assessed using Pearson's product moment correlation when data were normally distributed, or Spearman's rank correlation if data were non-parametric. Sample sizes, which were pre-determined using power calculations and previous experience, are reported in figure legends and represent biological replicates (i.e., individual animals). Means \pm SEM are plotted

for all graphs. All tests were two-sided and an α -level of $P < 0.05$ was used to determine significance. GraphPad Prism 9 software (version 9.5.1) was used for statistical analyses and figure production.

Study Approval

Experimentation involving mice was performed under license from the UK Home Office in accordance with the Animals (Scientific Procedures) Act (1986) and was approved by the UCL Queen Square Institute of Neurology Ethical Review Committee.

Author Contributions: JNS and GS conceived the study. JNS, DV-C, SS, TW, YT, RLS, JNSV, ERR, APT, SJW, QZ and GS performed the experiments. JNS wrote the manuscript and produced the figures, with input from all authors. JNS, XLY and GS supervised the experiments and acquired the funding.

Acknowledgements: We thank Amanda Pocratsky (UCL) and Nicholas Mazarakis (Imperial College London) for experimental advice, Robert W. Burgess (The Jackson Laboratory) for providing *Gars* ^{Δ ETAQ/+} mice, personnel of the Denny Brown Laboratory for assistance with mouse colonies (UCL), Alexander M. Rossor (UCL) for commenting on the manuscript and Stuart J. Grice (University of Oxford) for support and encouragement of the work. This work is dedicated to the memory of Amelia Singleton and Althea M. Sleigh. This project was funded by a Medical Research Council award MR/S006990/1 (JNS); a Wellcome Trust Sir Henry Wellcome Postdoctoral Fellowship 103191/A/13/Z (JNS); a Rosetrees Trust grant M806 (JNS, GS); the UCL Neurogenetic Therapies Programme funded by The Sigrid Rausing Trust (JNS, GS); a Human Frontier Science Program Long-Term Fellowship LT000220/2017-L (SS); a Motor Neuron Disease Association Junior Non-Clinical Fellowship Tosolini/Oct20/973-799 (APT); a National Institutes of Health R35 award GM139627 (XLY); Wellcome Trust awards 107116/Z/15/Z and 223022/Z/21/Z (GS); and a UK Dementia Research Institute Foundation award UKDRI-1005 (GS).

Materials & Correspondence: AAV expression plasmids are covered by an MTA with OXGENE (UK). Material requests should be addressed to the corresponding authors.

Data Availability: All data are provided in the main text and supplemental materials.

References

1. Reilly MM, et al. Charcot-Marie-Tooth disease. *J Peripher Nerv Syst.* 2011;16(1):1-14.
2. Pipis M, et al. Next-generation sequencing in Charcot-Marie-Tooth disease: Opportunities and challenges. *Nat Rev Neurol.* 2019;15(11):644-656.
3. Beijer D, et al. Defects in axonal transport in inherited neuropathies. *J Neuromuscul Dis.* 2019;6(4):401-419.
4. Prior R, et al. Defective axonal transport: A common pathological mechanism in inherited and acquired peripheral neuropathies. *Neurobiol Dis.* 2017;105:300-320.
5. Prokop A. A common theme for axonopathies? The dependency cycle of local axon homeostasis. *Cytoskeleton.* 2021;78(2):52-63.
6. Guedes-Dias P, Holzbaur ELF. Axonal transport: Driving synaptic function. *Science.* 2019;366(6462):eaaw9997.
7. Antonellis A, et al. Glycyl tRNA synthetase mutations in Charcot-Marie-Tooth disease type 2D and distal spinal muscular atrophy type V. *Am J Hum Genet.* 2003;72(5):1293-1299.
8. Wei N, et al. Neurodegenerative Charcot-Marie-Tooth disease as a case study to decipher novel functions of aminoacyl-tRNA synthetases. *J Biol Chem.* 2019;294(14):5321-5339.
9. He W, et al. Dispersed disease-causing neomorphic mutations on a single protein promote the same localized conformational opening. *Proc Natl Acad Sci U S A.* 2011;108(30):12307-12312.
10. Childers M, et al. Conformational sampling of CMT-2D associated GlyRS mutations. *Brain Multiphys.* 2022;3:100054.

11. He W, et al. CMT2D neuropathy is linked to the neomorphic binding activity of glycyl-tRNA synthetase. *Nature*. 2015;526(7575):710-714.
12. Grice SJ, et al. Dominant, toxic gain-of-function mutations in *gars* lead to non-cell autonomous neuropathology. *Hum Mol Genet*. 2015;24(15):4397-4406.
13. Sleigh JN, et al. Trk receptor signaling and sensory neuron fate are perturbed in human neuropathy caused by *Gars* mutations. *Proc Natl Acad Sci U S A*. 2017;114(16):E3324-E3333.
14. Mo Z, et al. Aberrant GlyRS-HDAC6 interaction linked to axonal transport deficits in Charcot-Marie-Tooth neuropathy. *Nat Commun*. 2018;9(1):1007.
15. Grice SJ, et al. Plexin-semaphorin signaling modifies neuromuscular defects in a *Drosophila* model of peripheral neuropathy. *Front Mol Neurosci*. 2018;11:55.
16. Cui Q, et al. Diverse CMT2 neuropathies are linked to aberrant G3BP interactions in stress granules. *Cell*. 2023;186(4):803-820.e25.
17. Spaulding EL, et al. The integrated stress response contributes to tRNA synthetase-associated peripheral neuropathy. *Science*. 2021;373(6559):1156-1161.
18. Zuko A, et al. tRNA overexpression rescues peripheral neuropathy caused by mutations in tRNA synthetase. *Science*. 2021;373(6559):1161-1166.
19. Bernabo P, et al. *In vivo* translome profiling in spinal muscular atrophy reveals a role for SMN protein in ribosome biology. *Cell Rep*. 2017;21(4):953-965.
20. Birsa N, et al. FUS-ALS mutants alter FMRP phase separation equilibrium and impair protein translation. *Sci Adv*. 2021;7(30):eabf8660.
21. Sleigh JN, et al. Altered sensory neuron development in CMT2D mice is site-specific and linked to increased GlyRS levels. *Front Cell Neurosci*. 2020;14:232.
22. Chao MV. Neurotrophins and their receptors: A convergence point for many signalling pathways. *Nat Rev Neurosci*. 2003;4(4):299-309.

23. Scott-Solomon E, Kuruvilla R. Mechanisms of neurotrophin trafficking via Trk receptors. *Mol Cell Neurosci.* 2018;91:25-33.
24. Villarroel-Campos D, et al. The many disguises of the signalling endosome. *FEBS Lett.* 2018;592(21):3615-3632.
25. Bercsenyi K, et al. Tetanus toxin entry. Nidogens are therapeutic targets for the prevention of tetanus. *Science.* 2014;346(6213):1118-1123.
26. Gibbs KL, et al. *In vivo* imaging of axonal transport in murine motor and sensory neurons. *J Neurosci Methods* 2016;257:26-33.
27. Sleigh JN, et al. *In vivo* imaging of anterograde and retrograde axonal transport in rodent peripheral nerves. *Methods Mol Biol.* 2020;2143:271-292.
28. Tosolini AP, et al. Expanding the toolkit for *in vivo* imaging of axonal transport. *J Vis Exp.* 2021(178):e63471.
29. Sleigh JN, et al. Mice carrying ALS mutant TDP-43, but not mutant FUS, display *in vivo* defects in axonal transport of signaling endosomes. *Cell Rep.* 2020;30(11):3655-3662.e2.
30. Achilli F, et al. An ENU-induced mutation in mouse glycyl-tRNA synthetase (GARS) causes peripheral sensory and motor phenotypes creating a model of Charcot-Marie-Tooth type 2D peripheral neuropathy. *Dis Model Mech.* 2009;2(7-8):359-373.
31. Sleigh JN, et al. Neuromuscular junction maturation defects precede impaired lower motor neuron connectivity in Charcot-Marie-Tooth type 2D mice. *Hum Mol Genet.* 2014;23(10):2639-2650.
32. Spaulding EL, et al. Synaptic deficits at neuromuscular junctions in two mouse models of Charcot-Marie-Tooth type 2d. *J Neurosci.* 2016;36(11):3254-3267.
33. Sleigh JN, et al. Developmental demands contribute to early neuromuscular degeneration in CMT2D mice. *Cell Death Dis.* 2020;11(7):564.

34. Morelli KH, et al. Allele-specific RNA interference prevents neuropathy in Charcot-Marie-Tooth disease type 2D mouse models. *J Clin Invest.* 2019;129(12):5568-5583.
35. Motley WW, et al. GARS axonopathy: Not every neuron's cup of tRNA. *Trends Neurosci.* 2010;33(2):59-66.
36. Mech AM, et al. Morphological variability is greater at developing than mature mouse neuromuscular junctions. *J Anat.* 2020;237(4):603-617.
37. Gonzalez M, et al. Disruption of Trkb-mediated signaling induces disassembly of postsynaptic receptor clusters at neuromuscular junctions. *Neuron.* 1999;24(3):567-583.
38. Finkbeiner S, et al. CREB: A major mediator of neuronal neurotrophin responses. *Neuron.* 1997;19(5):1031-1047.
39. Stifani N. Motor neurons and the generation of spinal motor neuron diversity. *Front Cell Neurosci.* 2014;8:293.
40. Maggs AM, et al. Nerve-dependent changes in skeletal muscle myosin heavy chain after experimental denervation and cross-reinnervation and in a demyelinating mouse model of Charcot-Marie-Tooth disease type 1A. *Muscle Nerve.* 2008;38(6):1572-1584.
41. Franklin JL, Johnson EM. Control of neuronal size homeostasis by trophic factor-mediated coupling of protein degradation to protein synthesis. *J Cell Biol.* 1998;142(5):1313-1324.
42. Reck-Peterson SL, et al. The cytoplasmic dynein transport machinery and its many cargoes. *Nat Rev Mol Cell Biol.* 2018;19(6):382-398.
43. Cai Q, et al. Snapin-regulated late endosomal transport is critical for efficient autophagy-lysosomal function in neurons. *Neuron.* 2010;68(1):73-86.
44. Olenick MA, et al. Dynein activator Hook1 is required for trafficking of BDNF-signaling endosomes in neurons. *J Cell Biol.* 2019;218(1):220-233.

45. Jordens I, et al. The Rab7 effector protein RILP controls lysosomal transport by inducing the recruitment of dynein-dynactin motors. *Curr Biol.* 2001;11(21):1680-1685.
46. Lang Q, et al. *In vivo* imaging of axonal transport in peripheral nerves of rodent forelimbs. *Neuronal Signal.* 2023; 7(1):NS20220098.
47. Skerratt SE, et al. The discovery of a potent, selective, and peripherally restricted pan-Trk Inhibitor (PF-06273340) for the treatment of pain. *J Med Chem.* 2016;59(22):10084-10099.
48. Delcroix JD, et al. NGF signaling in sensory neurons: Evidence that early endosomes carry NGF retrograde signals. *Neuron.* 2003;39(1):69-84.
49. Iverson C, et al. RDEA119/BAY 869766: A potent, selective, allosteric inhibitor of MEK1/2 for the treatment of cancer. *Cancer Res.* 2009;69(17):6839-6847.
50. Oosthuysen B, et al. Deletion of the hypoxia-response element in the vascular endothelial growth factor promoter causes motor neuron degeneration. *Nat Genet.* 2001;28(2):131-138.
51. Barker PA, et al. Nerve growth factor signaling and its contribution to pain. *J Pain Res.* 2020;13:1223-1241.
52. Binder DK, et al. BDNF and epilepsy: Too much of a good thing? *Trends Neurosci.* 2001;24(1):47-53.
53. Wang Z, et al. Adeno-associated virus serotype 8 efficiently delivers genes to muscle and heart. *Nat Biotechnol.* 2005;23(3):321-328.
54. Wang B, et al. Construction and analysis of compact muscle-specific promoters for AAV vectors. *Gene Ther.* 2008;15(22):1489-1499.
55. Sleigh JN, et al. Axonal transport and neurological disease. *Nat Rev Neurol.* 2019;15(12):691-703.

56. Cioni JM, et al. Late endosomes act as mRNA translation platforms and sustain mitochondria in axons. *Cell*. 2019;176(1-2):56-72.e15.
57. Vargas JNS, et al. Coupling axonal mRNA transport and local translation to organelle maintenance and function. *Curr Opin Cell Biol*. 2022;74:97-103.
58. Park MC, et al. Secreted human glycyl-tRNA synthetase implicated in defense against ERK-activated tumorigenesis. *Proc Natl Acad Sci U S A*. 2012;109(11):E640-E647.
59. Goughnour PC, et al. Extracellular vesicles derived from macrophages display glycyl-tRNA synthetase 1 and exhibit anti-cancer activity. *J Extracell Vesicles* 2020;10(1):e12029.
60. Mantilla CB, et al. TrkB kinase activity maintains synaptic function and structural integrity at adult neuromuscular junctions. *J Appl Physiol*. 2014;117(8):910-920.
61. Tosolini AP, et al. BDNF-dependent modulation of axonal transport is selectively impaired in ALS. *Acta Neuropathol Commun*. 2022;10:121.
62. Mitchell DJ, et al. Trk activation of the ERK1/2 kinase pathway stimulates intermediate chain phosphorylation and recruits cytoplasmic dynein to signaling endosomes for retrograde axonal transport. *J Neurosci*. 2012;32(44):15495-15510.
63. Santos AR, et al. Regulation of local translation at the synapse by BDNF. *Prog Neurobiol*. 2010;92(4):505-516.
64. Villarin JM, et al. Local synthesis of dynein cofactors matches retrograde transport to acutely changing demands. *Nat Commun*. 2016;7:13865.
65. Deng C, et al. Dynamic remodeling of ribosomes and endoplasmic reticulum in axon terminals of motoneurons. *J Cell Sci*. 2021;134(22):jcs258785.
66. Rai AK, et al. Molecular adaptations allow dynein to generate large collective forces inside cells. *Cell*. 2013;152(1-2):172-182.

67. Urnavicius L, et al. Cryo-EM shows how dynactin recruits two dyneins for faster movement. *Nature*. 2018;554(7691):202-206.
68. Ozes B, et al. AAV1.NT-3 gene therapy in a CMT2D model: Phenotypic improvements in *Gars*^{P278KY/+} mice. *Brain Commun*. 2021;3(4):fcab252.
69. Farinas I, et al. Characterization of neurotrophin and Trk receptor functions in developing sensory ganglia: Direct NT-3 activation of TrkB neurons *in vivo*. *Neuron*. 1998;21(2):325-334.
70. Klein R, et al. The trkB tyrosine protein kinase is a receptor for brain-derived neurotrophic factor and neurotrophin-3. *Cell*. 1991;66(2):395-403.
71. Naylor RL, et al. A discrete domain of the human TrkB receptor defines the binding sites for BDNF and NT-4. *Biochem Biophys Res Commun*. 2002;291(3):501-507.
72. Fan G, et al. Knocking the NT4 gene into the BDNF locus rescues BDNF deficient mice and reveals distinct NT4 and BDNF activities. *Nat Neurosci*. 2000;3(4):350-357.
73. Proenca CC, et al. Differential effects of BDNF and neurotrophin 4 (NT4) on endocytic sorting of TrkB receptors. *J Neurochem*. 2016;138(3):397-406.
74. Yamane M, et al. Semaphorin3A facilitates axonal transport through a local calcium signaling and tetrodotoxin-sensitive voltage-gated sodium channels. *Biochem Biophys Res Commun*. 2012;422(2):333-338.
75. Delezie J, et al. BDNF is a mediator of glycolytic fiber-type specification in mouse skeletal muscle. *Proc Natl Acad Sci U S A*. 2019;116(32):16111-20.
76. Bervoets S, et al. Transcriptional dysregulation by a nucleus-localized aminoacyl-tRNA synthetase associated with Charcot-Marie-Tooth neuropathy. *Nat Commun*. 2019;10(1):5045.

77. Blocquel D, et al. CMT disease severity correlates with mutation-induced open conformation of histidyl-tRNA synthetase, not aminoacylation loss, in patient cells. *Proc Natl Acad Sci U S A*. 2019;116(39):19440-19448.
78. Sun L, et al. CMT2N-causing aminoacylation domain mutants enable Nrp1 interaction with AlaRS. *Proc Natl Acad Sci U S A*. 2021;118(13):e2012898118.
79. Ermanoska B, et al. CMT-associated mutations in glycyl- and tyrosyl-tRNA synthetases exhibit similar pattern of toxicity and share common genetic modifiers in *Drosophila*. *Neurobiol Dis*. 2014;68:180-189.
80. Niehues S, et al. Impaired protein translation in *Drosophila* models for Charcot-Marie-Tooth neuropathy caused by mutant tRNA synthetases. *Nat Commun*. 2015;6:7520.
81. Verhoeven K, et al. Mutations in the small GTP-ase late endosomal protein RAB7 cause Charcot-Marie-Tooth type 2B neuropathy. *Am J Hum Genet*. 2003;72(3):722-727.
82. Nave KA. Myelination and the trophic support of long axons. *Nat Rev Neurosci*. 2010;11(4):275-283.
83. Tinevez JY, et al. TrackMate: An open and extensible platform for single-particle tracking. *Methods*. 2017;115:80-90.
84. Murray L, et al. Dissection of the transversus abdominis muscle for whole-mount neuromuscular junction analysis. *J Vis Exp*. 2014(83):e51162.
85. Sleigh JN, et al. Morphological analysis of neuromuscular junction development and degeneration in rodent lumbrical muscles. *J Neurosci Methods* 2014;227:159-165.
86. Sleigh JN, et al. A simple, step-by-step dissection protocol for the rapid isolation of mouse dorsal root ganglia. *BMC Res Notes*. 2016;9:82.
87. Sleigh JN, et al. A video protocol for rapid dissection of mouse dorsal root ganglia from defined spinal levels. *BMC Res Notes*. 2020;13(1):302.

88. Villarroel-Campos D, et al. Dissection, *in vivo* imaging and analysis of the mouse epitrochleoanconeus muscle. *J Anat.* 2022; 241(5):1108-1119.
89. Sleigh JN, et al. Chondrolectin affects cell survival and neuronal outgrowth in *in vitro* and *in vivo* models of spinal muscular atrophy. *Hum Mol Genet.* 2014;23(4):855-869.
90. Welinder C, Ekblad L. Coomassie staining as loading control in Western blot analysis. *J Proteome Res.* 2011;10(3):1416-1419.
91. Babcock LW, et al. MyoSight-semi-automated image analysis of skeletal muscle cross sections. *Skelet Muscle.* 2020;10(1):33.
92. Rodino-Klapac LR, et al. Lack of toxicity of alpha-sarcoglycan overexpression supports clinical gene transfer trial in LGMD2D. *Neurology.* 2008;71(4):240-247.

Figure




Figure 1. Neuropathy-causing *GARS1* mutations impair retrograde axonal transport of signalling endosomes in vivo. (A) Intramuscular injection of a fluorescent fragment of tetanus neurotoxin (H_cT-555), with subsequent exposure of the sciatic nerve, permits in vivo imaging of signalling endosomes in intact peripheral nerve axons of live, anaesthetized mice. (B) Retrogradely transported H_cT-positive endosomes are individually tracked to quantitatively assess their dynamics. Colour-coded arrowheads identify five different endosomes. (C) Endosome frame-to-frame speed histograms of wild-type and CMT2D-modelling *Gars*^{C201R/+} mice aged 0.5, 1 and 3 months. (D) Defective axonal transport manifests in *Gars*^{C201R/+} mice between 0.5 and 1 month of age (genotype $P = 0.033$, age $P = 0.006$, interaction $P = 0.009$ two-way ANOVA). (E) Endosome frame-to-frame speed histograms of 1 month-old wild-type mice receiving intramuscular injections of recombinant human wild-type (left) or CMT2D-

causing (right) GlyRS. (F-H) GlyRS^{L129P} and GlyRS^{G240R}, but not GlyRS^{WT}, cause a non-cell autonomous decrease in signalling endosome speed (F), increased pause time (G) and more pausing endosomes (H) in healthy axons (F-H, $P < 0.001$ one-way ANOVA). For all graphs, * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ Šídák's multiple comparisons test; # $P < 0.05$ unpaired t -test; $n = 5-6$; means \pm SEM plotted. Scale bars = 20 (A) and 10 μm (B). See also Supplemental Figure 1.




Figure 2. Dysfunctional BDNF-TrkB signalling correlates with CMT2D pathology. (A) Representative western blots of lysates from five different wild-type muscles, probed with anti-TrkB. These muscles display a spectrum of vulnerability to NMJ denervation in $Gars^{C201R/+}$, with the transversus abdominis (TVA) and epitrochleoanconeus (ETA) being least impacted and the flexor digitorum brevis (FDB) displaying the greatest degeneration (33). (B and C) Levels of FL-TrkB (** $P = 0.001$ one-way ANOVA) and TrkB.T1 (** $P < 0.001$ Kruskal-Wallis test) differ between muscles. FL-TrkB (** $P < 0.001$ Pearson's product moment correlation), but not TrkB.T1 ($P = 0.267$ Spearman's rank correlation), positively correlates with CMT2D denervation. (D) Representative western blots of lysates from five different wild-type muscles, probed with anti-GlyRS and anti-CHRNA (quantified in Supplemental Figure 3, C and D). (E and F) Representative collapsed z-stack confocal images of lumbar spinal cord ventral horns from wild-type and $Gars^{C201R/+}$ mice stained for ChAT and p-CREB or CREB.

The lower panels show inverted fluorescence images. Arrows highlight motor neuron nuclei. Scale bars = 100 μm . (**G** and **H**) *Gars*^{C201R/+} lumbar motor neurons display reduced p-CREB activation (**G**) and are smaller than their wild-type counterparts (**H**). * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ unpaired t -test. For all graphs, $n = 5-6$; means \pm SEM plotted. Mice were P77-P83 (A-D) and 3 months-old (E-H). See also Supplemental Figures 2-5.




Figure 3. Endosome adaptor protein levels are selectively reduced in hindlimb-innervating nerves. (A) Western blot of dynein adaptor proteins Snapin, Hook1 and RILP in sciatic nerves (hindlimb) from 1 month-old wild-type and *Gars*^{C201R/+} mice. Densitometric analyses show reduced levels of monomeric Snapin (mSnapin, ***P* = 0.002) and RILP (**P* = 0.018) in *Gars*^{C201R/+} sciatic nerves. (B) Western blot of Snapin, Hook1 and RILP in combined median and ulnar nerves (forelimb) from 1 month-old wild-type and *Gars*^{C201R/+} mice. Contrasting with sciatic nerves, densitometric analysis identified an increase in dimeric Snapin (dSnapin, ***P* = 0.004) in median and ulnar nerves. (C) Endosome frame-to-frame speed histograms from 3 month-old wild-type and *Gars*^{C201R/+} median and ulnar nerves. (D-F) Signalling endosome transport speed (D, *P* = 0.954), pause time (E, *P* = 0.910) and pause percentage (F, *P* = 0.810 Mann-Whitney *U* test) in forelimb-innervating nerves are unaffected in *Gars*^{C201R/+} mice. For all graphs, genotypes were compared using unpaired *t*-tests, unless otherwise stated; *n* = 4-5; means ± SEM plotted.




Figure 4. In vivo BDNF depletion and TrkB inhibition impair endosome transport via ERK1/2. (A) Endosome frame-to-frame speed histograms of wild-type and *Gars^{C201R/+}* mice 4-8 h post-treatment with intramuscular injections of anti-BDNF or IgY control antibody. (B) Anti-BDNF slows wild-type endosome transport yet has no effect on *Gars^{C201R/+}* ($P < 0.001$ one-way ANOVA). $***P < 0.001$ Šidák's multiple comparisons test. $n = 5-6$. (C and D) Intramuscular injection of 50 nM pan-Trk inhibitor (PF-06273340) slows signalling endosome transport in wild-type motor axons. $*P < 0.05$ unpaired t -test. $n = 5-6$. (E and F) Intramuscular injection of 50 nM ERK1/2 inhibitor, refametinib, also impairs endosome transport in wild-type mice. $**P < 0.01$ unpaired t -test. $n = 3$. For all graphs, means \pm SEM plotted. Mice were 1 month-old, except panels E and F (P40-P45). See also Supplemental Figures 6 and 7.




Figure 5. Augmenting BDNF in CMT2D muscles restores in vivo axonal transport. (A) Endosome frame-to-frame speed histograms of wild-type and $Gars^{C201R/+}$ mice aged 1 and 3 months 4-8 h post-treatment with intramuscular BDNF injections. (B) BDNF rescues in vivo axonal transport of signalling endosomes in $Gars^{C201R/+}$ mice ($P = 0.875$ one-way ANOVA). (C) BDNF also corrects transport in 1 month-old $Gars^{\Delta ETAQ/+}$ mice ($P < 0.001$ one-way ANOVA). (D and E) The axonal transport impairment induced by human CMT2D-causing GlyRS is prevented by exogenous recombinant BDNF (E, $P < 0.001$ Kruskal-Wallis test). Mice were aged 1 month. For all graphs, $*P < 0.05$, $**P < 0.01$, $***P < 0.001$ Šídák's/Dunn's multiple comparisons test; $n = 4-6$; means \pm SEM plotted. See also Supplemental Figure 8.




Figure 6. BDNF specifically rescues axonal transport in CMT2D mice via a Trk-dependent mechanism. (A) Endosome frame-to-frame speed histograms of one month-old $Gars^{C201R/+}$ mice 4-8 h post-treatment with vehicle, BDNF or BDNF plus pan-Trk inhibitor, (13 nM PF-06273340). (B) Trk inhibition abrogates the rescue effect of BDNF on $Gars^{C201R/+}$ endosome transport speed ($P = 0.023$). (C) Endosomes of $Gars^{C201R/+}$ mice treated with BDNF and PF-06273340 spend as much time paused as vehicle-treated mice ($P < 0.001$ Kruskal-Wallis test). (D) Trk inhibition also abrogates the positive effect of BDNF on pausing endosomes ($P = 0.005$). (E) Endosome frame-to-frame speed histograms of 1 month-old $Gars^{C201R/+}$ mice 4-8 h post-treatment with vehicle, VEGF₁₆₅, NT-3 or NT-4. (F-H) VEGF₁₆₅

further impairs *Gars*^{C201R/+} endosome transport speed, while NT-3 and NT-4 have no effect (F, $P = 0.011$). No significant changes in pausing were observed (G, $P = 0.086$; H, $P = 0.323$). **(I)** Endosome frame-to-frame speed histograms of *Gars*^{C201R/+} mice treated with vehicle or BDNF for 24 h, rather than the usual 4-8 h, prior to imaging. **(J-L)** 24 h post-injection, BDNF no longer rescues *Gars*^{C201R/+} endosome transport speed (J, $P = 0.248$), percentage time paused (K, $P = 0.664$) or the percentage of pausing endosomes (L, $P = 0.966$). Mice were aged P29-P42 (I-L). For all graphs, data were compared using one-way ANOVAs, unless otherwise stated; * $P < 0.05$, ** $P < 0.01$ Šídák's/Dunn's multiple comparisons test; $n = 5-9$; means \pm SEM plotted. N.b., the vehicle treatment data are also presented in Figure 1, C and D, and BDNF treatment data in Figure 5, A and B.




Figure 7. Muscle-specific BDNF gene therapy restores in vivo axonal transport and endosome adaptor levels in CMT2D mice. (A) Endosome frame-to-frame speed histograms of P38-P41 wild-type and *Gars*^{C201R/+} mice treated at P11 with AAV8-tMCK-eGFP or AAV8-tMCK-BDNF. (B) The BDNF gene therapy rescues *Gars*^{C201R/+} in vivo axonal transport and increases wild-type endosome speeds ($P < 0.001$). (C) Representative western blot of sciatic nerve lysates from AAV-treated mice probed with antibodies against dynein adaptors. (D and E) Treatment with AAV8-tMCK-BDNF restores levels of both monomeric (mSnapin; D, $P = 0.016$ Kruskal-Wallis test) and dimeric (dSnapin; E, $P = 0.002$) Snapin in *Gars*^{C201R/+} sciatic nerves, but has little effect on Hook1 ($P = 0.460$, not shown) or RILP ($P = 0.605$, not shown). For all graphs, data were compared using one-way ANOVAs, unless otherwise stated; * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$ Šidák's/Dunn's multiple comparisons test; $n = 6$; means \pm SEM plotted. See also Supplemental Figure 9.




Figure 8. Boosting muscle BDNF rescues impaired axonal transport of signalling endosomes in CMT2D mice. *Top right:* Mutant GlyRS aberrantly interacts with the extracellular domain (ECD) of TrkB, the main neurotrophin receptor found at the NMJ. The availability of full-length TrkB (FL-TrkB) in wild-type muscles correlates with the extent of denervation in CMT2D mice, such that higher FL-TrkB levels are associated with reduced NMJ innervation in neuropathy. *Middle and top left:* CMT2D mice display reduced speed of signalling endosome axonal transport in sciatic, but not median/ulnar, nerves (*middle*), which is associated with reduced ERK1/2 phosphorylation and decreased endosome adaptor levels in sciatic nerves (*middle*), as well as dampened CREB activation (*top left*). *Bottom:* Injection of

mature BDNF (mBDNF) or AAV8-tMCK-BDNF, but not NT-3, NT-4 or VEGF₁₆₅, into CMT2D muscles completely restores physiological axonal transport in vivo. Figure created using <https://www.biorender.com>.