Table 1:

Characteristics	n (%)		
Age (years)			
< 1	3 (2.4)		
1-4	34 (27.0)		
5-9	31 (24.6)		
10-14	43 (34.1)		
15-18	15 (11.9)		
Gender			
Female	58 (46.0)		
Male	68 (54.0)		
World Bank country income level			
High-income	51 (40.5)		
Upper-middle-income	64 (50.8)		
Lower-middle-income	11 (8.7)		
Low-income	0		
CNS tumor diagnosis			
Low grade glioma	25 (19.8)		
High grade glioma	18 (14.3)		
CNS embryonal tumor	41 (32.5)		
Ependymoma	11 (8.7)		
CNS germ cell tumor	8 (6.4)		
Craniopharyngioma	5 (4.0)		
Other intracranial or intraspinal tumor	11 (8.7)		
Missing	7 (5.6)		
CNS tumor treatment phase			
Active therapy	81 (64.3)		
Chemotherapy within 30 days	59 (46.8)		
Palliative therapy (including oral	7 (5.6)		
chemotherapy)			
No active treatment	15 (11.9)		
Treatment completed	22 (17.5)		
Unknown	1 (0.8)		

Table 2:

Characteristics	n (%)	
Symptoms	. ,	
No symptoms	74 (58.7)	
Constitutional	38 (30.2)	
Respiratory	24 (19.0)	
ENT	16 (12.7)	
Neurologic	15 (11.9)	
Gastrointestinal	6 (4.8)	
Dermatologic	1 (0.8)	
Comorbidities		
None	101 (80.2)	
Recent use of high-dose steroids	5 (4.0)	
At least one comorbidity	22 (17.5)	
Imaging findings		
No imaging test done	83 (65.9)	
Normal findings	21 (16.7)	
Abnormal findings	12 (9.5)	
Imaging done, unknown findings	2 (1.6)	
Unknown	8 (6.4)	
ANC (cells/μL)		
ANC > 1000	60 (47.6)	
500 < ANC <= 1000	5 (4.0)	
ANC <= 500	17 (13.5)	
Missing	44 (34.9)	
ALC (cells/μL)		
ALC > 1000	47 (37.3)	
500 < ALC <= 1000	13 (10.3)	
300 < ALC <= 500	2 (1.6)	
100 < ALC <= 300	6 (4.8)	
ALC <= 100	11 (8.7)	
Missing	47 (37.3)	
Admission at presentation		
New admission to hospital	33 (26.2)	
Patient already admitted	31 (24.6)	
No admission required	62 (49.2)	

Table 3:

Characteristics	n (%)		
Treatment modification			
Not receiving cancer-direct therapy	40 (37.4)		
Cancer-directed therapy unmodified	31 (29.0)		
Chemotherapy withheld	25 (23.4)		
Radiation therapy delayed	7 (6.5)		
Neurosurgery was delayed	2 (1.9)		
Unknown	3 (2.8)		
Highest level of respiratory support			
Room air	95 (88.8)		
Nasal cannula/facemask	6 (5.6)		
CPAP/BiPAP	1 (0.9)		
Intubation	2 (1.9)		
Unknown	3 (2.8)		
Level of care			
Not admitted to higher level of care	99 (92.5)		
Admitted to the intensive care unit	5 (4.7)		
Admitted to intermediate care unit	1 (0.9)		
Unknown	2 (1.9)		
Outcomes at 30 days			
SARS-CoV-2 infection cleared	75 (70.1)		
SARS-CoV-2 infection clinically resolved	21 (19.6)		
SARS-CoV-2 still positive, asymptomatic	7 (6.5)		
Expired due to SARS-CoV-2 infection or its complications	1 (0.9)		
Expired, due to other cause	2 (1.9)		
Unknown	1 (0.9)		

Table 4:

Group	Total	Asymptomatic n (%)	Mild/Moderate n (%)	Severe/Critical n (%)	p Value
Chemotherapy timing					
>30 days	6	3 (50)	3(50)	0 (0)	0.76
≤30 days	44	27 (61)	14 (32)	3 (7)	
Comorbidity					
None/Unknown	19	11 (58)	7 (37)	1 (5)	0.92
One or more	97	46 (53)	35 (40)	6 (7)	
ANC					
>500 cells/μL	56	27 (48)	26 (46)	3 (5)	0.04
≤500 cells/μL	9	2 (22)	4 (44)	3 (33)	
ALC					
>300 cells/μL	55	26 (47)	25 (45)	4 (7)	0.85
≤300 cells/μL	8	3 (38)	4 (50)	1 (12)	