



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Mental health in Europe during the COVID-19 pandemic: a systematic review



Nafiso Ahmed, Phoebe Barnett, Anna Greenburgh, Tamara Pemovska, Theodora Stefanidou, Natasha Lyons, Sarah Ikhtabi, Shivangi Talwar, Emma R Francis, Samantha M Harris, Prisha Shah, Karen Machin, Stephen Jeffreys, Lizzie Mitchell, Chris Lynch, Una Foye, Merle Schlief, Rebecca Appleton, Katherine R K Saunders, Helen Baldwin, Sophie M Allan, Luke Sheridan-Rains, Omayya Kharboutly, Ariana Kular, Peter Goldblatt, Robert Stewart, James B Kirkbride, Brynmor Lloyd-Evans, Sonia Johnson

The COVID-19 pandemic caused immediate and far-reaching disruption to society, the economy, and health-care services. We synthesised evidence on the effect of the pandemic on mental health and mental health care in high-income European countries. We included 177 longitudinal and repeated cross-sectional studies comparing prevalence or incidence of mental health problems, mental health symptom severity in people with pre-existing mental health conditions, or mental health service use before versus during the pandemic, or between different timepoints of the pandemic. We found that epidemiological studies reported higher prevalence of some mental health problems during the pandemic compared with before it, but that in most cases this increase reduced over time. Conversely, studies of health records showed reduced incidence of new diagnoses at the start of the pandemic, which further declined during 2020. Mental health service use also declined at the onset of the pandemic but increased later in 2020 and through 2021, although rates of use did not return to pre-pandemic levels for some services. We found mixed patterns of effects of the pandemic on mental health and social outcome for adults already living with mental health conditions.

Introduction

Following the onset of the COVID-19 pandemic on March 11, 2020, mental health was swiftly recognised as an area of concern.¹⁻⁴ Potential consequences of the pandemic and associated social restrictions included increase in psychological distress, increase in new onsets of mental health conditions, and worsening of difficulties already experienced by people living with mental health conditions. Pandemic-related service disruption had the potential to exacerbate such effects on mental health. Many studies have investigated aspects of the pandemic's effect on mental health, but systematic reviews have focused only on early stages of the pandemic,^{5,6} measures of symptoms in the general population,⁷⁻¹⁰ or comparisons between before and after lockdown.^{5,11,12} Our aim is to provide a comprehensive overview of the mental health effect of the pandemic in its first 2 years in one major region of the world, to help to inform planning for the continuing response to the COVID-19 pandemic and future emergencies.

We systematically reviewed evidence regarding mental health epidemiology in high-income countries in Europe. We focused on this region because of the similarities between its countries in timing of COVID-19 waves, health service responses, and social restrictions.^{13,14} We included studies that made comparisons either before and after the onset of the pandemic or between different timepoints during the pandemic, and that addressed the following three questions: (1) what changes have there been in the incidence or prevalence of mental health problems; (2) what changes have there been to mental distress, symptom severity, social functioning, quality of life, suicidal behaviours, and self-harm among people already living with mental health conditions; and (3) what changes have there been in mental health service use?

Methods

We followed PRISMA guidelines.¹⁵ The research questions and protocol were developed and refined through consultation with a stakeholder working group, including experts by experience, health and social care practitioners, and researchers. The protocol was pre-registered on PROSPERO (CRD42022323723). The review was conducted by the National Institute for Health and Care Research Policy Research Unit (funded to deliver evidence to inform health policy making). The research was initiated in response to a policy maker request for an evidence synthesis to guide forecasts of future service needs.

Search strategy and selection criteria

We searched four electronic databases (MEDLINE, PsycINFO, Embase, and CINAHL) for articles published between March 1, 2020, and February 1, 2022, and four pre-print servers (MedRxiv, PsyArXiv, Wellcome Open Research, and JMIR) for articles registered between March 1, 2020, and March 7, 2022. A combination of keyword and subject heading searches was used. Search terms for mental health conditions including psychotic, affective, anxiety, personality, and eating disorders were combined with terms for COVID-19. We included only longitudinal and repeated cross-sectional studies reporting on high-income European countries (using Organisation for Economic Co-operation and Development criteria).^{16,17} No age or language restrictions were applied. We conducted backward reference searching from all included studies, but not forward citation chaining, as doing this with the large number of identified studies would have prevented us from synthesising and delivering evidence promptly. Full search strategies shown in the appendix (pp 1–11).

Lancet Psychiatry 2023

Published Online

June 12, 2023

[https://doi.org/10.1016/S2215-0366\(23\)00113-X](https://doi.org/10.1016/S2215-0366(23)00113-X)

S2215-0366(23)00113-X

National Institute of Health and Care Research Mental Health Policy Research Unit

(N Ahmed PhD, P Barnett MSc, A Greenburgh PhD, T Pemovska MSc, T Stefanidou MSc, N Lyons MSc, M Schlief MSc, R Appleton PhD, H Baldwin MSc, L Sheridan-Rains PhD, A Kular PhD, Prof B Lloyd-Evans PhD, Prof S Johnson DM), **Lived Experience Working Group** (P Shah, K Machin, S Jeffreys BA, L Mitchell, C Lynch), **Division of Psychiatry** (S Ikhtabi MSc, S Talwar MPhil, O Kharboutly BSc, Prof J B Kirkbride PhD), and **Centre for Outcomes Research and Effectiveness** (P Barnett), **Department of Clinical, Educational and Health Psychology, Division of Psychology and Language Sciences** (E R Francis MSc), and **Institute of Health Equity** (Prof P Goldblatt PhD), **University College London, London, UK**; **National Collaborating Centre for Mental Health, Royal College of Psychiatrists, London, UK** (P Barnett); **Department of Health Promotion and Development, Faculty of Psychology, University of Bergen, Bergen, Norway** (S M Harris PhD); **National Institute of Health and Care Research Mental Health Policy Research Unit** (U Foye PhD, K R K Saunders PhD), **Institute of Psychiatry, Psychology and Neuroscience** (Prof R Stewart MD), **King's College London, London, UK**; **Cambridgeshire and Peterborough National Health Service Foundation Trust** (S M Allan ClinPsyD); **South London and Maudsley National Health Service Foundation Trust, London, UK**

(Prof R Stewart); Camden and Islington National Health Service Foundation Trust, London, UK (Prof S Johnson)

Correspondence to: Dr Nafiso Ahmed, National Institute of Health and Care Research Mental Health Policy Research Unit, Division of Psychiatry, University College London, London W1T 7NF, UK nafiso.ahmed@ucl.ac.uk

See Online for appendix

We included longitudinal or repeated cross-sectional studies comparing timepoints during the COVID-19 pandemic versus prior to the pandemic, or between different points in the pandemic. Studies with samples of people without pre-existing mental health conditions were included (research questions one and three), as well as those meeting cutoffs indicating a clinical condition on validated diagnostic instruments or mental health symptoms measures (questions two and three). We excluded studies with samples defined by having a physical health condition or being COVID-19 survivors or health-care professionals due to the unique nature of their experiences. We did not include samples defined by intellectual disability or neurodevelopmental disorders, dementia, or other organic mental disorder, or substance misuse. Included studies reported at least one of the following: incidence or prevalence estimates (either by diagnostic assessment or proportion meeting the clinical threshold on a validated symptom measure); change in mental distress, symptom severity, social functioning, quality of life, suicidal behaviours, or self-harm in people already living with mental health conditions; or change in mental health service use. We included only studies in which the majority of the sample lived in high-income European countries.

Title, abstract, and full-text screening were carried out on EPPI-Reviewer Web.¹⁸ Seven reviewers (ST, SI, UF, RA, ERF, MS, and NL) independently screened titles and abstracts for studies meeting inclusion criteria. Full texts of potentially eligible studies were retrieved and screened independently by the same reviewers. A second reviewer (NA) screened a random 10% of papers at both stages to validate decisions. Disagreements were resolved through team discussion, and steps were taken to improve agreement.

Data extraction and quality appraisal

A data extraction form was developed and piloted on 10% of included studies using EPPI-Reviewer Web.¹⁸ Data were extracted independently by one of 13 reviewers (NA, PB, RA, UF, ERF, ST, KRS, SMH, MS, SI, PS, NL, and LS-R) and checked for accuracy by a second reviewer (NA, PB, SMH, AG, TP, ST, SI, or ERF). We extracted data on the study design, aims and objectives, dataset, country or region, publication status when the data were extracted, sample size, involvement in study of people with relevant lived experience, population, age, gender or sex, ethnicity, comparison group, symptom or condition measured, setting, primary outcome measures, and associated statistical data. Study quality was assessed using the Newcastle-Ottawa Scale for cohort studies and the adapted Newcastle-Ottawa Scale created for cross-sectional studies by Herzog and colleagues (appendix pp 38–45).¹⁹ Certainty of evidence for each outcome was independently assessed by two of four authors (PB, AG, NA, and HB) using the Grading of Recommendations Assessment, Development and Evaluation (GRADE) system.²⁰ GRADE guidance was

adapted for narrative synthesis according to Murad and colleagues,²¹ and it was further adapted according to methodological differences in the studies addressing each research question. Adaptations were discussed and agreed with the working group (appendix pp 46–77).

Data analysis

We analysed changes in prevalence (ascertained through research diagnosis or reaching cutoff score on a clinical measure, a definition used in other reviews focusing on the pandemic)¹² and incidence of mental health problems; mental health symptom severity, social functioning, quality of life, psychosocial outcomes, suicidal behaviours, and self-harm in people with pre-existing mental health conditions; and any mental health service use indicator within crisis and acute mental health services, community mental health and outpatient services, and primary care relating to mental health.

A narrative synthesis was done as the studies were very heterogeneous, especially in terms of timepoints compared and symptoms or services examined. Studies were organised according to their measurement period (pre-pandemic compared with during the pandemic, or different time points over the course of the pandemic), reported outcome, mental health problem, and service type. Studies that measured general psychopathology or mental distress were grouped together as non-specific mental health problems. During the narrative synthesis, we noted whether studies analysed different samples within the same dataset. Data from multiple papers were reported together when the study sample was the same in the different papers.

Results

We identified 7066 records from title and abstract screening. 687 full texts were assessed, of which 149 records met the inclusion criteria. Studies excluded at full-text stage with the reasoning behind exclusions are shown in the appendix (pp 12–37). Pre-print and backward citation searches identified a further 31 records, which gave a total of 177 studies, reported in 180 papers (figure, appendix p 78). 73 (41%) of 177 studies reported changes in prevalence and incidence of mental health problems; 37 (21%) reported studies symptom severity, social functioning, quality of life, suicide behaviours, and self-harm in people with pre-existing mental health conditions; and 76 (43%) mental health service use. Eight (5%) of 177 studies provided information on multiple research questions.

14 (8%) of 177 studies measured mental health outcomes in children and young people aged 6–18 years, and 163 (92%) in adults. Sample sizes ranged from 20 to 24897725, and studies were from 20 European countries: UK (n=46), Italy (n=24), Germany (n=20), Netherlands (n=12), Spain (n=10), France (n=9), Ireland (n=5), Norway (n=6), Austria (n=6), Portugal (n=5), Switzerland (n=5), Sweden (n=3), Belgium (n=3),




Figure: PRISMA flow diagram

OECD=Organisation for Economic Co-operation and Development.

Türkiye (n=3), Denmark (n=2), Poland (n=2), Czech Republic (n=2), Iceland (n=1), Greece (n=1), and Lithuania (n=1). 11 studies reported data from multiple countries. All studies were in English except for three, which were in German.^{22–24}

88 (50%) of 177 studies compared mental health outcomes before the pandemic with those during the pandemic, 55 (31%) measured outcomes at different timepoints over the course of the pandemic, and 34 (19%) had both a pre-pandemic and during pandemic comparator. A repeated cross-sectional design was used in 93 (53%) studies, a longitudinal cohort design in 83 (47%) studies, and one study used an open cohort design. Only four (2%) studies reported involving people with relevant lived experience in designing studies or interpreting results. Of the 180 papers, 112 (62%) were rated as high quality, and 68 (38%) as low quality. GRADE certainty of evidence ratings on 103 outcomes were rated high for 15 (15%), moderate for 22 (21%), and low to very low for 66 (64%) studies. Further details of study characteristics (appendix pp 79–259), quality assessments

(appendix pp 38–45), and GRADE ratings (appendix pp 46–77) are available.

Concerning the prevalence and incidence of mental health problems before versus during the pandemic, and at different timepoints of the pandemic (research question one), there was evidence of high to moderate certainty that the prevalence of depression, generalised anxiety disorder, and non-specific mental health problems was higher during the pandemic (usually restricted to timepoints in 2020) than before the pandemic,^{24–65} with statistically significant increases ranging from 0.25% to 31% (table 1; appendix pp 260–79). Very low-certainty evidence suggested that prevalence of eating disorders and mixed depression and anxiety also increased by 20–21% for eating disorders⁶⁷ and 3.9% for mixed depression and anxiety.⁴¹ We found no relevant evidence on other major conditions such as psychosis or bipolar disorder.

Moderate-certainty evidence suggested that prevalence of depression did not change substantially between the onset of the pandemic and the end of June, 2020.^{35,36,68,69,70,71}

	Number of studies	Countries	Studies showing prevalence or incidence increased	Studies showing prevalence or incidence decreased	Studies showing prevalence or incidence varied over time	Summary	Certainty of evidence
Prevalence pre-pandemic vs during the pandemic							
Depression	30	UK (5), the Netherlands (5), Germany (3), Czech Republic (2), Norway (2), Ireland (2), UK and Ireland (1), Austria (1), Belgium (1), France (1), Iceland (1), Italy (1), Portugal (1), Spain (1), Sweden (1), Switzerland (2)	23/30:17 showed a significant increase (0.25–31%); 3 showed an increase with no statistical testing; 3 showed non-significant increase	7/30: 3 showed significant reduction; 1 showed a reduction with no statistical testing; 3 showed a non-significant decrease	NA	30 studies reported changes in prevalence of depression during the pandemic compared with before the pandemic. Overall, high certainty evidence suggests that prevalence increased, with 23 of 30 studies reporting an increase and 17 studies reporting a significant increase of between 0.25% and 31%. University students: 3 of 3 studies reported a significant increase of between 8.7% and 27%. Children and young people: 2 of 6 studies reported non-significant decreases in prevalence, 1 of 6 reported a non-significant increase in prevalence, and 3 of 6 reported a significant increase of between 0.25% and 14.9%. Pregnant people: 1 of 2 studies reported an increase in prevalence but without any statistical testing, and 1 of 2 reported a non-significant decrease in prevalence. Mothers: 2 of 22 studies reported a significant increase of between 9% and 14%. Disadvantaged immigrants: 1 of 11 studies reported a non-significant increase. Adults older than 50 years: 3 of 3 studies showed a significant rise of between 12% and 16% (2 datasets). ^{34-53,55-56,66}	High
Generalised anxiety disorder	15	UK (2), UK and Ireland (1), Netherlands (5), Germany (2), Ireland (1), Czech Republic (1), Norway (1), Portugal (1), Türkiye (1)	11/15: 7 showed a significant increase (0.28–28.2%), 3 showed an increase with no statistical testing; 1 showed non-significant increase	4/15: 1 showed a significant reduction (6.6%); 1 reported a reduction with no numbers or statistics; 2 showed non-significant reduction	NA	15 studies reported changes in prevalence of generalised anxiety disorder during the pandemic compared with before the pandemic. Overall, high certainty evidence suggests that prevalence increased, with 11 of 15 studies reporting an increase and 7 of these reporting a significant increase of between 0.28% and 28.2%. University students: 1 of 2 reported a significant increase of 19.3%, and 1 of 2 reported a significant decrease of 6.6%. Children and people younger than 18 years: 3 of 3 studies reported a significant increase of between 0.28% and 8.1%. Mothers: 1 study reported a significant increase of 6%. Pregnant people: 1 of 2 studies reported a large increase without statistical testing of 28%, and 1 of 2 reported a non-significant reduction in prevalence. ^{27,29,30,32,35-38,41,45,46,53,55,55,66}	High
Non-specific mental health problem	8 (5 different data sets)	UK (4), the Netherlands (2), Norway (2)	6/8: 5 (3 of 5 datasets) showed a significant increase (0.5–43%); 1 showed a non-significant increase	2/8: 1 showed a significant decrease (1.5%); 1 showed a non-significant decrease	NA	8 studies reported changes in prevalence of non-specific mental health problems or psychological distress. Moderate certainty evidence suggests that prevalence increased, with 6 of 8 studies reporting an increase and 5 of these reporting a significant increase of between 0.5% and 13%. Adolescents aged between 13 and 16 years: 1 study reported a 0.7% increase in prevalence. UK Armed Forces Veterans: 1 study reported a non-significant increase in prevalence. ^{37-59,60-65}	Moderate
Comorbid anxiety and depression	1	UK and Ireland (1)	1/1	0/1	NA	Very low certainty evidence from 1 study suggests that the prevalence of comorbid anxiety and depression increased by 3.9%. University students: 1 study reported a significant increase in prevalence of comorbid anxiety and depression. ⁴¹	Very low
Eating disorders	1	France (1)	1/1	0/1	NA	Very low certainty evidence from 1 study suggests that the prevalence of eating disorders increased by 21.2% in women and 20% in men. University students: 1 study reported a significant increase in prevalence of eating disorders. ⁶⁷	Very low

(Table 1 continues on next page)

	Number of studies	Countries	Studies showing prevalence or incidence increased	Studies showing prevalence or incidence decreased	Studies showing prevalence or incidence varied over time	Summary	Certainty of evidence
(Continued from previous page)							
Prevalence over the course of the pandemic							
Depression							
March to June 2020	5	UK (1), Ireland (1), Germany (1), Italy (1), Spain (1)	1/5 showed non-significant increase	4/5: 1 showed a significant reduction; 1 showed a non-significant reduction; 2 showed a reduction with no statistical testing	0/5	5 studies compared changes in depression over the initial stages of the pandemic. Moderate certainty evidence suggests that there was not a significant reduction in prevalence during this time, although 1 of 5 studies reported a significant reduction of 7%. ^{35,36,68-71}	Moderate
July to December 2020	11	UK (1), UK and Germany (1), UK & Ireland (1), Austria (1), Belgium (1), France (1), Greece (1), EU (1), Norway (1), Sweden (1), Switzerland (1)	4/11 showed a non-significant increase	4/11: 2 showed a significant reduction; 1 showed a reduction with no statistical testing; 1 showed a non-significant decrease	3/11: significantly higher prevalence during lockdown periods and significantly lower prevalence during easing of measures	11 studies compared changes in depression during 2020. Moderate certainty evidence suggests that the prevalence of depression varied throughout 2020 but might have reduced, with 2 of 11 studies reporting significant reductions of around 6%. University students: 1 of 2 studies reported lower prevalence of depression when restrictions were eased, and 1 of 2 studies reported a non-significant increase in prevalence. Pregnant people: 1 study reported lower prevalence of depression when restrictions were eased. ^{33,39,72-80}	Moderate
January to March 2021	8	UK (2), Ireland (1), France (2), Multiple EU countries(1), Poland (1)	2/8: 1 showed significant increase; 1 showed an increase with no statistical testing	3/8: 1 showed a reduction with no statistical testing; 2 showed a significant decrease	3/8: 2 reported lower prevalence during summer when restrictions were eased but the highest during peak COVID-19 periods; 1 reported prevalence was significantly lower in 2021 compared with 2020 but increased during summer 2020	8 studies compared changes in depression between 2020 and early 2021. Very low certainty evidence suggests that prevalence of depression did not significantly change overall, although rates might be slightly higher during later lockdown periods. University students: 1 of 2 studies reported that prevalence was highest during the second lockdown (December to January 2021), and 1 of 2 reported a reduction in prevalence without any statistical testing. ^{33,36,68,67}	Very low
April to July 2021	2	EU (1), Portugal (1)	0/2	2/2: 1 showed a decrease with no statistical testing; 1 showed a non-significant decrease	0/2	2 studies compared changes in depression prevalence during the later stages of 2021. Low certainty evidence provides unclear evidence of how prevalence changed. Mothers: 1 study reported a small decrease in prevalence with no statistical testing. ^{88,89}	Low
Generalised anxiety disorder							
March to June 2020	7	UK (1), Spain (2), Ireland (1), Germany (2), Italy (1)	1/7 showed a significant increase	5/7: 2 showed a significant decrease; 2 showed a decrease with no statistics; 1 showed a non-significant decrease	1/7: prevalence rates were higher immediately after bad news such as personal protective equipment shortages	7 studies compared changes in anxiety over the initial stages of the pandemic. Low certainty evidence suggests that there might have been a decline in prevalence during the start of the pandemic (5 of 7 studies reported reductions with 2 of 7 reporting significant reductions of 2-4-11-5%), although prevalence rates might have increased during the first month of lockdown before reducing. ^{35,36,43,68-71,90}	Low
July to December 2020	11	UK (2), UK and Ireland (1), Austria (1), Belgium (1), Czech Republic (1), France (1), Greece (1), EU (1), Norway (1), Sweden (1)	4/11: 3 showed significant increase; 1 showed non-significant increase	5/11: 2 showed significant reduction; 3 showed non-significant reduction.	2/11: reduced prevalence rates were recorded during non-lockdown periods (summer 2020)	11 studies compared changes in anxiety over the course of 2020. Moderate certainty evidence suggests that there might have been an overall increase in prevalence with 4 of 11 studies reporting an increase and 3 of these reporting a significant increase of 2-3% by November or December, with a decrease in prevalence during the summer of 2020. University Students: 1 study reported increased generalised anxiety disorder prevalence during lockdowns. Pregnant people: 1 reported a non-significant reduction. Parents: one reported a significant decline over the course of 2020. Adults older than 50 years reported a significant increase in prevalence of 2%. ^{33,13-34,72,74-80}	Moderate

(Table 1 continues on next page)

	Number of studies	Countries	Studies showing prevalence or incidence increased	Studies showing prevalence or incidence decreased	Studies showing prevalence or incidence varied over time	Summary	Certainty of evidence
(Continued from previous page)							
January to March 2021	10	UK (2), Ireland (1), France (2), Poland (2), EU (1), Portugal (1), Switzerland (1)	7/10: 4 showed significant increase; 2 showed an increase with no statistics; 1 showed non-significant increase	3/10: 1 showed significant reduction; 2 showed a reduction with no statistics	0/10	10 studies compared changes in generalised anxiety disorder between 2020 and early 2021. Low certainty evidence suggests the prevalence of generalised anxiety disorder was higher during 2021 than in 2020, with 7 of 10 studies reporting an increase and 4 of these reporting a significant increase of 6–19%. University students: 3 of 5 studies reported a significant increase of 8–19%, 1 of 4 studies reported a reduction in prevalence with no statistics, and 1 of 5 reported an increase with no statistics. ^{27,38,39,81-84,89,91}	Low
April to July 2021	1	EU (1)	0/1	1/1 showed a non-significant reduction	0/1	1 study examined changes only between February and June, 2021, and very low certainty evidence from this study suggests that prevalence of generalised anxiety disorder did not change. ⁸⁸	Very low
Non-specific mental health problem							
March to June 2020	2	The Netherlands (1), Spain (1)	0/2	1/2 showed a significant decrease between March and June 2020	1/2 showed increase during April lockdown	2 studies examined changes in prevalence of non-specific mental health problems over the early stages of the pandemic. Very low certainty evidence from these studies suggests that prevalence increased overall although might have varied during this time. ^{64,65,92}	Very low
July to December 2020	4	UK (3), EU (1)	0/4	4/4: 3 (2 of 3 datasets) showed significant reduction; 1 showed non-significant reduction	0/4	4 studies compared changes in non-specific mental health problems over the course of 2020. Low certainty evidence suggests that the prevalence of non-specific mental health problems reduced between early and later stages of 2020, with all 4 studies on 3 datasets reporting a decrease, and 4 studies (2 datasets) reporting a significant decrease of 6–21% (although 1 study comparing prevalence rates across multiple countries reported that Norway did not see a significant decrease when other countries did). ^{29,39-95}	Low
January to March 2021	2	UK (1), Germany (1)	1/2 show an increase but with no statistical testing	0/2	1/2 reported increase during late 2020, and decrease during early 2021	2 studies compared changes in general non-specific mental health problems from 2020 to early 2021. Very low certainty evidence suggests that there was no clear shift in prevalence during this time. ^{86,97}	Very low
PTSD							
July to December 2020	2	UK (1), EU (1)	0/2	2/2	0/2	2 studies compared changes in PTSD over the course of 2020. Low certainty evidence suggests that prevalence might have reduced in late compared with early 2020, with both studies reporting significant decreases of 2–4%. ^{75,98}	Low
January to March 2021	1	Ireland (1)	0/1	0/1	1/1 showed decrease throughout 2020 before increase in 2021	Very low certainty evidence from one study suggests that although prevalence of PTSD might not have differed in 2021 compared with 2020 overall, prevalence might have decreased in later stages of 2020 compared with early stages before increasing in 2021. ^{35,36}	Very low
April to July 2021	1	EU (1)	0/1	1/1	0/1	Very low certainty evidence from one study suggests that prevalence between early and later parts of 2021 decreased significantly by 5%. ⁸⁸	Very low
Panic disorder							
July to December 2020	1	EU (1)	1/1 non-significant report	0/1	0/1	Very low certainty evidence from one study suggests that there was a non-significant reduction in panic disorder symptoms between June and November 2020. ⁷⁵	Very low

(Table 1 continues on next page)

Number of studies	Countries	Studies showing prevalence or incidence increased	Studies showing prevalence or incidence decreased	Studies showing prevalence or incidence varied over time	Summary	Certainty of evidence
(Continued from previous page)						
Incidence before vs during the pandemic						
2	UK (1), Lithuania (1)	0/2	2/2	NA	2 studies compared incidence rates of depression before the pandemic to those during the pandemic. Moderate certainty evidence suggests that there was a significant decline in the incidence of depression diagnoses recorded on medical records, both in comparison to 2018 and 2019 data (n=1, 127-56 per 100 000 reduction), and in comparison to expected rates based on previous years (n=1, 13-8% per 100 000 person-months). ^{88,89}	Moderate
1	UK (1)	0/1	1/1	NA	Low certainty evidence from 1 study suggests that incidence of generalised anxiety disorder reported on clinical records reduced significantly (26-4% per 100 000 person-months less) in 2020, compared with expected rates. ⁸⁸	Low
1	Lithuania (1)	0/1	1/1	NA	Low certainty evidence from one study suggests that the incidence of PTSD diagnoses reported on health-care records reduced significantly in 2020, compared with before the pandemic (1-96 per 100 000 reduction). ⁸⁹	Low
Incidence over the course of the pandemic						
1	UK (1)	1/1 (Northern Ireland, Scotland, and Wales data)	0/1	1/1 (England data)	Low certainty evidence from 1 study suggests that monthly incidence might have reduced in early stages of the pandemic before returning to pre-pandemic rates by September in England, although in the rest of the UK monthly incidence increased. ⁸⁸	Low
1	UK (1)	1/1 (Northern Ireland, Scotland, and Wales data)	0/1	1/1 (England data)	Low certainty evidence from one study suggests that monthly incidence might have reduced in early stages of the pandemic before returning to pre-pandemic rates by September in England, although in the rest of the UK monthly incidence increased. ⁸⁸	Low
1	UK (1)	0/1	1/1	0/1	Low certainty evidence from one study suggests that monthly incidence reduced between April and July 2020, from 28-6% to 9%. ⁸⁹	Low
NA=not applicable. PTSD=post-traumatic stress disorder.						

Table 1: Prevalence and incidence of mental health problems compared with before, and over the course of the pandemic

Low-certainty evidence suggested that generalised anxiety disorder prevalence reduced during this time (2.4–11.5% decline)^{35,36,43,68,69,70,71,90} and very low-certainty evidence suggested prevalence of non-specific mental health problems varied over time but might have increased overall between March and June, 2020.^{64,65,92}

As the pandemic continued, there was moderate-certainty evidence for an overall slight increase (2–3%) in generalised anxiety disorder from the early months of the pandemic to second half of 2020, but with a dip in prevalence during summer months.^{33,52–54,72,74–80} There was low to moderate certainty evidence that depression (around 6% decline), non-specific mental health problems (6–21% decline), and post-traumatic stress disorder (PTSD; 2–4% decline) prevalence reduced between the early pandemic and the end of 2020.^{33,50,57,72–80,93–95}

The few studies comparing later in the pandemic (2021) to earlier (2020) found moderate-certainty evidence of little change in the prevalence of depression or non-specific mental health problems between 2020 and 2021.^{35,36,81–89,96,97} Evidence of low to very low certainty suggested an increase in generalised anxiety disorder rates (6–19%) and decreased PTSD prevalence (5% decrease reported in one study).^{27,35,36,50,81–85,87,88,91}

Studies on sub-populations, including children and young people,^{24,26,37,58} university students,^{27,50,51} and mothers and pregnant people,^{29,33,48,55,56} also reported increased prevalence of all mental health problems during the pandemic compared with before its onset. The exceptions were anxiety disorders in students (one of two studies reported a significant decrease),⁴¹ and depression in children and young people (two out of six studies reported a decrease).^{32,39} Studies of university students reported higher prevalence of depression and generalised anxiety disorder during lockdown periods than when restrictions were eased, both in 2020 and 2021.^{72,85} One study reported that prevalence of depression and generalised anxiety disorder among parents reduced by July 2020, compared with earlier pandemic timepoints.⁷⁷ One study on an older adults group with mean age of 67 years reported a significant increase in prevalence of generalised anxiety disorder by December 2020.⁵⁴

Estimates of incidence based on service data found evidence of low to moderate certainty that the incidence of depression, generalised anxiety disorder, and PTSD showed statistically significant reductions after the onset of the pandemic compared with before the pandemic.^{98,99} Low-certainty evidence also suggested that the monthly incidence recorded by mental health services of depression, generalised anxiety disorder, and non-specific mental health problems reduced further during 2020.^{93,98}

For people already living with mental health conditions (research question two), data are shown in the appendix on changes in symptoms during compared with before the pandemic (pp 279–291) and at different time points of the pandemic (pp 291–308). Among adults living with

mental health conditions at the onset of the pandemic, evidence of moderate to high certainty suggested no significant change in general psychopathology and mental distress symptoms,^{23,57,100,101} significant improvement in depressive symptoms,^{102–106} and mixed findings regarding changes to anxiety and eating disorders symptoms^{100,102,105,106} after the onset of the pandemic. Very low to low certainty evidence suggested significant worsening in PTSD symptoms and mixed findings for changes to schizophrenia and bipolar symptoms among adults during the first half of the pandemic compared with before the pandemic.^{101,103,105}

Comparing timepoints during the pandemic, moderate-certainty evidence suggests no significant change in depressive symptoms, and mixed findings regarding obsessive-compulsive disorder symptoms in adult clinical populations during the first half of 2020.^{70,107–112} Low-certainty evidence suggested no statistically significant change in schizophrenia and bipolar symptoms, but statistically significant improvements in general psychopathology during the early pandemic period.^{108,111–113} Evidence was mixed and of low-certainty on whether eating disorder and anxiety symptoms changed in adult clinical populations during the first half of 2020.^{70,107,108,111,112}

Evidence from studies comparing mental health symptoms among adults between the first and second half of 2020, suggested with moderate to high certainty that general psychopathology and mental distress, and symptoms of depression, schizophrenia, and bipolar disorder did not change significantly.^{103,104,114–117} Findings regarding PTSD were mixed.^{114,115} Low-certainty evidence suggested no significant change in anxiety symptoms over this period, but findings regarding obsessive-compulsive disorder symptoms were mixed.^{116,118}

No statistically significant change in anxiety and eating disorder symptoms among adults was found during the pandemic between June 2020, and January 2021; the evidence was of very low certainty.¹¹⁹ Findings on changes in schizophrenia and bipolar symptoms between October 2020, and February 2021, were mixed and the evidence was of very low certainty.¹²⁰ Moderate-certainty evidence indicated no statistically significant changes in general psychopathology and mental distress, and there was very low to low certainty of no statistically significant changes in anxiety and PTSD symptoms between March or July 2020, and June or July 2021, among adults.^{121,122}

Very low-certainty evidence indicated significant worsening in obsessive-compulsive disorder, general psychopathology, and mental distress among children and young people already living with mental health conditions after the onset of the pandemic.^{123,124} Low-certainty evidence suggested no significant change in symptoms of depression and anxiety among children and young people with mental health conditions in comparisons with pre-pandemic, just after pandemic onset, and December in 2020.³² However, low-certainty evidence also suggested significant worsening in

	Number of studies	Countries	Summary	Certainty
Crisis and acute mental health care before vs during the pandemic				
Inpatient: adults				
March to June 2020	21	Portugal (2), Italy (6), UK (6), Germany (2), Spain (2), Denmark, France, multiple countries	Adult inpatient service use decreased in the early pandemic period compared with before the pandemic. 12 studies reported decrease in measures of admissions (5 reported that these changes were significant), although 8 studies reported increases in measures of admissions (5 reported that these changes were significant). Despite indications that admissions decreased, there was a suggestion that more severe inpatient cases increased during the early pandemic: the types of admissions in which increases were found largely referred to compulsory admissions, hospitalisation after presentation to emergency departments, and psychiatric intensive care unit hospitalisation. Admissions decreased by between 11% and 43%. Only one percentage statistic for increase of admissions was reported (15.7%). ^{22,129,132-150}	High
July to December 2020	9	Germany (2), Italy (2), France, Italy, Denmark, UK, Spain, Türkiye	Adult inpatient service use decreased over 2020, pandemic period compared with before the pandemic. 8 studies found a decrease in admissions (2 reported that decreases were significant, reporting decreases in admissions of 3–41.6%). No study found an increase in the number of admissions overall. Increases were found by 4 studies in length of stay, number of section 136 Mental Health Act assessments, consultations, and proportion of all psychiatric admissions as a function of all inpatient admissions. Unlike earlier in 2020, results were more mixed later in the pandemic for compulsory admissions: 1 study reported a decrease in involuntary admissions, and another an increase in the proportion of admissions that were involuntary. ^{124,130,141,145,151-155}	High
January to March 2021	3	UK, Denmark, multiple countries	There was slightly stronger indication of a reduction than an increase in adult inpatient service use in 2021 compared with pre-pandemic periods. 2 studies reported a decrease in indicators of admissions (rate of admissions, and the perception of change in admissions reported by heads of psychiatry). 1 study reported a significant 5% decrease in the rate of inpatient admissions, but the results were mixed as another study reported an increase in the proportion of referred patients that were admitted and a number of patients on the waitlist. ^{129,143,156}	High
Inpatient: paediatric				
March to June 2020	3	Spain (2), France	Paediatric inpatient service use decreased in the early pandemic compared with before the pandemic. All studies found a decrease in admissions (1 study reported that this was significant; admissions decreased by 18–42%), and 1 found a decrease in length of hospital stay. ¹⁵⁷⁻¹⁵⁹	High
Emergency department and walk-in services: adults				
March to June 2020	27	Portugal (2), Switzerland, Italy (7), UK (9), Belgium, Spain, Portugal, Germany (2), Ireland (2), France, Türkiye	Mental health-related use of adult emergency department and walk-in services decreased in the early pandemic period compared with pre-pandemic rates. 22 studies reported decreases in service use indicators including presentations, consultations, assessments, referrals, liaison psychiatry contacts, and referrals (11 studies reported a significant decrease, and decreases in psychiatric presentations ranged from 13.5% to 58%). The decrease was most pronounced in the early lockdown period (reported by 4 studies), and 1 study including both lockdown and post-lockdown periods in 2020, found the decrease was greater in the lockdown than post-lockdown period. 7 studies reported increases in measures pertaining to service use (4 studies reported that these increases were significant). However only 2 increases were found in psychiatric presentations to emergency departments (0.53–5.6%) and in psychiatric visits as a proportion of all emergency departments visits and the number of repeat visits within 1 month. ^{130,132,134,136-140,142,146,147,150,160-174}	High
July to December 2020	5	Italy (2), Germany, UK, Türkiye	Mental health-related use of adult emergency department and walk-in services decreased during 2020, compared with pre-pandemic periods. 4 studies reported decreases in service use (12–16% reduction in psychiatric presentation numbers to emergency departments; 1 of the decreases reported was reported as significant). Decreases were found in total and daily psychiatric presentations, liaison psychiatry referrals, and consultations. 1 study reported an increase in service use measures, including in the number of urgent psychiatric consultations overall, daily, and by telephone (the daily and telephone measures were reported as significant). ^{22,130,152,175,176}	High
Emergency department and walk-in services: paediatric				
March to June 2020	4	France, Spain (2), UK	Mental health-related use of paediatric emergency department and walk-in services decreased in the early pandemic periods compared with before the pandemic. 3 studies found decreases in psychiatric presentations including for self-harm and suicidality, with decreases ranging from 36% to 61%, but 1 study found a significant increase in presentations by 164.5%. ^{157,177,178,179}	Moderate
July to December 2020	1	France	1 study found an increase in emergency department admissions due to suicide attempts of 80% in November and December, 2020, compared with November and December, 2019. ¹⁷⁷	Very low
January to March 2021	1	France	1 study found an increase of 202% in emergency department admissions due to suicide attempts in March and April 2021, compared with March and April, 2019. ¹⁷⁷	Very low
Community-based crisis care				
March to June 2020	2	UK	Evidence suggests a reduction in use of community-based crisis care in the early pandemic period compared with pre-pandemic rates. 1 study found a decrease of 5–10% in referrals, and another found a 24.9% decrease in total contacts and a 26.4% decrease in overall caseloads, and identified a shift from face-to-face to virtual contacts, with a 102% increase in virtual contacts. ^{148,164}	Moderate

(Table 2 continues on next page)

	Number of studies	Countries	Summary	Certainty
(Continued from previous page)				
Trauma and resuscitation rooms				
March to June 2020	3	Austria (2), UK	All studies identified an increase in mental health-related admissions to trauma and resuscitation rooms in the early pandemic period compared with before the pandemic. These increases were found in the proportion of admissions due to suicide attempts, proportion of admissions with a psychiatric diagnosis, and number of admissions due to suicide attempts. 2 studies reported these increases were significant. ¹⁸⁰⁻¹⁸²	Low
Crisis and acute mental health care over the course of the pandemic				
Inpatient: adults				
March to June 2020	1	Germany	1 study found daily admissions increased in later lockdown compared with early lockdown, and that the length of hospital stay significantly decreased from early to late lockdown. ¹⁸³	Low
July to December 2020	2	Italy, Spain	Evidence was mixed regarding how service use changed after lockdown. 1 study found admissions significantly decreased post-lockdown compared with during lockdown, but the other found they increased, although this change was not significant. ^{145,184}	Very low
January to March 2021	2	Italy, multiple European countries	Studies indicated an increase in inpatients as the pandemic progressed into 2021 (78% in 1 study). A slight decrease in length of hospitalisation was also reported in the second wave vs first wave. ^{141,143}	Very low
Emergency department and walk-in services: adults				
March to June 2020	4	UK (2), Portugal, Spain	2 studies showed visits decreased during March, 2020 (1 study reported this as significant). After the end of March, increases were observed in emergency visits and referrals in 2 studies (1 reported as significant). In comparison with lockdown, 1 study showed visits increased by 21% after lockdown. ^{138,164,171,185}	High
July to December 2020	3	Switzerland, Italy, UK	3 studies reported an increase in service use based on the number of consultations or liaison referrals after lockdown compared with the early lockdown, with the increase ranging from 21% to 56.9%. 1 study reported the proportion of all emergency department activity reported to liaison psychiatry services increased from March to May, but decreased from May to August, 2020. ^{176,184,185}	High
Emergency department and walk-in services: paediatric				
January to March 2021	1	France	There was an increase in emergency department admissions due to suicide attempts from March and April 2020, onwards. By March and April 2021, these admissions (n=48.7) had increased by 524% compared with March and April, 2020 (n=7.8). ¹⁷⁷	Very low
Community mental health and outpatient services before vs during the pandemic				
Adults of working age				
March to June 2020	11	UK (7), Italy, Netherlands, Spain (Balearic Islands), Denmark	Mental health and outpatient services for adults of working age was decreased in the early pandemic compared to pre-pandemic. All studies reported a decrease in service use measures (4 reported these were significant; decreases in referrals ranged from 24% to 75.3%) including number of patients, patients attending follow-up, referral rates, referrals to Improving Access to Psychological Therapies, number of patients accessing these services, face-to-face contacts, home visits, caseloads, assessments, daily caseloads, and self-reported contact with a mental health professional or webpage. Results were mixed as 5 studies also reported an increase in some service use indicator; however, in 3 studies these increases (of 147-157%) were mostly found in virtual contacts, and the only significant increase was found in video consultations, although increases were reported in self-referrals to Improving Access to Psychological Therapies by patients from minoritised backgrounds after the fourth week of lockdown, consultations, and daily caseloads. ^{71,98,129,140,48,349,364,186-188}	High
July to December 2020	7	UK (3), Italy (2), Germany, Denmark	Use of community mental health and outpatient services for adults of working age increased during 2020, compared with pre-pandemic use rates. 6 studies reported increases (1 reported as significant) in measures of service use including online self-referrals, appointments, referrals, and remote consultations. 1 study reported a percentage change in mental health service utilisation (20% increase). 3 studies reported a decrease in service use over the pandemic in 2020, compared with pre-pandemic rates (1 reported as significant), for referrals and self-reported use of mental health professionals. ^{129,152,167,175,186,189,190}	High
January to March 2021	1	Denmark	1 study found referral rate in the second lockdown was non-significantly higher than before lockdown. ¹²⁹	Moderate
Children and adolescents				
March to June 2020	5	UK (3), Italy, Sweden	Use of community mental health and outpatient services for children and adolescents decreased in the early pandemic period compared with before the pandemic. All studies found decreases in indicators of service use (2 reported that decreases were significant), with decreases of contacts ranging from 4.36% to 12%, including in referrals, daily caseloads, total contacts, face-to-face contacts, proportion of patients undergoing neuropsychiatric interventions, and contact with mental health services in children whose schools were closed. 2 studies reported increases: 1 reporting a shift to virtual contacts (126% increase in virtual contacts, 86% decrease in face-to-face contacts), and the other reporting a marginal increase in admissions. ^{125,140,149,164,191}	Moderate
July to December 2020	2	Italy, Sweden	Use of community mental health and outpatient services for children and adolescents was decreased over 2020, compared with before the pandemic. Both studies reported decreases in indicators of service use compared with pre-pandemic rates, both in the proportion of patients undergoing psychiatric and psychopharmacological treatment, and in contact with mental health services in children whose schools had been closed earlier in the pandemic, even after in-person teaching resumed (3.55% decrease). ^{125,191}	Low
January to March 2021	1	Sweden	1 study found a persistent decrease (5.23%) in contacts with mental health services in children whose schools had been closed earlier in the pandemic in 2021, when in-person teaching had been resumed. ¹⁹¹	Low

(Table 2 continues on next page)

	Number of studies	Countries	Summary	Certainty
(Continued from previous page)				
Older adults				
March to June 2020	5	UK (4), Germany	Evidence suggests a decrease in use of community mental health and outpatient services for older adults in the early pandemic compared with before the pandemic. All studies reported a decrease in measures of service use (4 reported that these were significant), including referrals, consultations (5–18% decrease), recognition of incident diseases (16% decrease), daily caseloads (6–14% decrease), total assessments (20% decrease), admissions (27% decrease), and referrals (39% decrease). ^{147,149,154,192,193}	Moderate
July to December 2020	1	Netherlands	1 study found that use of mental health professionals by older adults was significantly decreased in the pandemic over 2020, compared with pre-pandemic rates. ¹⁹⁴	Low
Community mental health and outpatient services over the course of the pandemic				
Adults of working age				
January to March 2021	1	Austria	1 study found psychotherapists reported increased patient numbers overall (by 77.2%), and increased number of patients treated in-person in 2021, compared with the early pandemic in 2020. The number treated over telephone decreased in 2021. ¹⁹⁵	Very low
Children and adolescents				
March to June 2020	1	UK	1 study found a non-significant increase in referrals over the course of lockdown. ¹⁶⁴	Low
July to December 2020	1	Italy	1 study reported a 48% reduction in the number of patients undergoing psychopharmacological treatment in August and September, 2020, compared with during lockdown. ¹²⁵	Very low
January to March 2021	1	Multiple European countries	1 study found that a greater proportion of the heads of children and young people psychiatric services reported that referrals and outpatient numbers had increased in 2021, compared with at the start of the pandemic. However, a greater proportion of them reported that outpatients had decreased in 2021, since the start of the pandemic, than reported that outpatients had increased. ¹⁴³	Very low
Older adults				
March to June 2020	1	UK	1 study found a non-significant increase in referrals over the course of lockdown. ¹⁶⁴	Low
Primary care before vs during the pandemic				
General practitioners				
March to June 2020	1	UK	1 study found mental health-related use of primary care reduced in the early pandemic compared with pre-pandemic rates. First diagnoses of common mental health problems reported by general practitioners decreased significantly by 50%, and related first prescriptions was also lower in this pandemic period. ¹⁹⁶	Very low
July to December 2020	3	UK (3), Norway	Results were mixed for studies investigating mental health-related use of primary care over 2020, compared with pre-pandemic rates. 2 studies found decreases (31–46% significant decrease in consultations, and 20–47% decrease in contacts) and 2 studies reported increases (17% increase in psychological presentations and 6% in prescriptions, both statistically significant). ^{197,198,199}	Moderate
Primary care over the course of the pandemic				
General practitioners				
July to December 2020	2	UK	Both studies found that after the initial drop in contacts and consultations, these rates increased (1–2% per week) after the end of March until the end of the studied periods (July 2020). ^{198,199}	High

Table 2: Service use outcomes compared with before, and over the course of the COVID-19 pandemic

depressive and anxiety symptoms among children and young people from just after pandemic onset to December 2020.³² Overall, findings regarding changes in general psychopathology and mental distress over this timeframe in children and young people with mental health conditions were mixed, with conclusions of very low-certainty.¹²⁵

Moderate-certainty evidence suggested significant worsening in social functioning among adult clinical populations after the pandemic onset versus before the pandemic.^{105,106} Comparing timepoints during the pandemic, there were indications of improvement in psychosocial outcomes—such as psychosocial impairment—during the first of half of 2020 (very low-certainty evidence).¹²⁶ Overall, low-certainty evidence suggested significant improvements in psychosocial burden and negative psychosocial effect of pandemic

restrictions in later 2020 versus just after pandemic onset.^{117,127} Social functioning and quality of life did not change significantly during this time, or between 2020 and 2021 (very low-certainty evidence).^{118,122}

Overall, moderate-certainty evidence indicated that suicidal behaviour (measured by clinical records) did not significantly change between before the pandemic and at various timepoints during it, in 2020 and 2021, among clinical populations of all ages.^{128–130} However, there was low-certainty evidence of a reduction in self-harm measured through clinical records (in a sample aged 10 years and older) early in the pandemic versus pre-pandemic (low-certainty).⁹⁸

Around the onset of the pandemic, self-reported ideation was more frequent, but self-reported self-harm less frequent among clinical populations than at later timepoints before July 2020 (very low-certainty

evidence).^{70,131} Low-certainty evidence comparing pandemic timepoints before July, 2020, with later in 2020, suggested fluctuating rates of case note-reported self-harm.⁹⁸

Concerning service use outcomes during versus before the pandemic, and at different time points during the pandemic, overall, moderate to high-certainty evidence suggests that the use of critical and acute mental health care services decreased in the early pandemic period versus before the pandemic (research question three; table 2, appendix pp 308–325). This decrease included mental health inpatient care (adult admissions: 11–43% reduction, paediatric admissions: 18–42% reduction), mental health presentations to emergency department and walk-in services (adult presentations: 14–58%, paediatric presentations: 36–61%), and community-based crisis care.^{109,129,130,132–149,157–174,177–180,183} A decrease below pre-pandemic service use levels was also found for community mental health and outpatient services for adults of working age (referrals: 24–75% reduction; high-certainty evidence),^{71,98,129,140,147–149,164,186–188} children and young people (total contacts: 4–36–12% reduction; moderate-certainty evidence),^{125,164,140,149,191} and older adults (consultations: 5–18% reduction; moderate-certainty evidence),^{164,147,149,192,193} and for mental-health related contacts in primary care (very low-certainty evidence).¹⁹⁶ Across community mental health services, a shift from face-to-face to remote contacts was reported in the first half of 2020.^{140,148,187} An exception was an increase in mental health-related admissions to trauma and resuscitation units in the first half of 2020, which was three to ten times higher (low-certainty evidence) than before the pandemic.^{180–182} Likewise, despite an overall decrease in use of adult mental health inpatient services, there were indications that more severe difficulties (eg, resulting in compulsory hospitalisations) increased.^{180–182}

After an initial decrease, mental health-related use of adult emergency departments (high-certainty evidence), adult inpatient mental health care (low-certainty evidence), and community mental health services for children and young people and older adults (low-certainty evidence) increased by June, 2020, compared with just after the onset of the pandemic.^{138,164,171,183,200} Mental health-related contacts in emergency departments and primary care was increased in later 2020, compared with earlier in the pandemic (high-certainty evidence).^{160,176,184,198,199} The use of community services for children and young people decreased in 2020, after the pandemic onset but was greater in 2021, than earlier in the pandemic (very low-certainty evidence).^{125,143} Use of all these services remained below pre-pandemic levels at later timepoints in 2020: in adult mental health inpatient care and emergency departments (inpatient admissions: 3–42% lower; mental health eating disorders presentations: 12–16% lower; high-certainty evidence), and in community mental health and outpatient services for children and young people and older adults

(low-certainty evidence).^{22,125,129,130,141,145,151-155,175,176,191,194} Service use was still below pre-pandemic levels in 2021, for adult inpatient care (high-certainty evidence) and children and young people's community services (low-certainty evidence).^{129,143,156,191}

Conversely, the use of community mental health and outpatient services for adults of working age (moderate to high-certainty evidence) and paediatric (very low-certainty evidence) emergency department and walk-in services reached higher levels than pre-pandemic later in 2020 and 2021.^{129,151,152,175,177,186,191,192} Findings concerning primary care service use (moderate-certainty evidence) at later timepoints were mixed.^{197–199}

Discussion

We identified 177 studies from 20 high-income European countries comparing mental health and mental health-service use outcomes either before and during the pandemic, or over the course of the pandemic. Most studies reported that prevalence of mental health problems including depression, anxiety, and non-specific conditions rose after the onset of the pandemic in general population samples. This observation could be interpreted as an acute response to a global event that caused widespread disruption, fear, financial hardship, and grief. Governmental restrictions and lockdowns were most stringent during the beginning of the pandemic.²⁰¹ Lockdowns might have augmented known risk factors for mental ill-health, such as unemployment and social isolation, while disrupting access to face-to-face professional and social support.^{2,202,203}

These rises were modest in most cases, however, and by late 2020, the increase in prevalence of mental health problems appears to have slowed. This is consistent with a meta-analysis of longitudinal studies, which found prevalence was higher than before the pandemic in March and April, 2020, but that it no longer was in July, 2020.¹⁰ That meta-analysis,¹⁰ however, only compared the worldwide prevalence of mental health problems before and early in the pandemic, whereas we have examined changes as the pandemic progressed in more detail in high-income European countries.

Studies using health-care records consistently reported fewer incident diagnoses of mental health problems after the onset of the pandemic than before the pandemic, which is consistent with our previous findings based on data from early in the pandemic¹ and with a 2022 systematic review comparing psychiatric service use before and during the pandemic among children and young people.²⁰⁴ Early in the pandemic, concerns over risk of infection appear to have prevented some from seeking in-person support.^{1,203,205} This disparity between increased prevalence of mental health problems and reduced service use suggests that the treatment gap in addressing mental health problems²⁰⁶ in the population might have increased, with potential long-term repercussions. As well

documented elsewhere,^{207–209} we found that service providers adopted telemental health to ensure continuity of service delivery early in the pandemic. However, for some people, access to these services was challenging, for example due to poor digital connectivity and inadequate private space.^{207,210}

After the initial substantial drop observed, service use began to rise but often remained below pre-pandemic levels. However, later in 2020 and in 2021, service use in paediatric emergency services (mental health presentations) and community mental health services for adults of working age rose above pre-pandemic levels. Given sparse research at timepoints beyond 2020, it is unclear whether these trends continued, and they should be cautiously interpreted given long-term increases in demand for mental services already observed before the pandemic.^{211–213}

We found no clear pattern of change in mental health symptom severity and associated outcomes in adults with pre-existing mental health conditions. Most studies showed either no significant change or different findings for different outcomes, with generally low certainty of evidence, partly as each outcome was the focus of only a few studies. The absence of a clear worsening of most symptoms contrasts with qualitative reports from people with pre-existing mental health conditions on the negative effects of the pandemic on their mental health, for example because of disrupted treatment and routines, and increased social and economic stressors.²¹⁴ Our mixed and sometimes surprising results might be understood in relation to large variations in experiences of people living with mental health conditions, with the effects of the pandemic varying by condition, the extent to which people were able to continue to connect with formal and informal support, and the interactions of the pandemic's effects with pre-existing social isolation and adversity.^{3,215} Some people reported some positive consequences, including a sense of a shared societal experiences, reconnecting with family and friends at the onset of the pandemic, mobilising existing reserves of resilience, peer support, and the absence of some pre-pandemic stressors.^{1,203} An independent commentary on our research process and findings is provided by co-authors of the paper who contributed from the perspective of their experience of living and supporting others with mental health conditions (panel).

In children and young people with some pre-existing mental health conditions, we found evidence that symptoms of obsessive-compulsive disorder, general psychopathology, and mental distress significantly worsened at the start of the pandemic in 2020. This contrasted with reduced service use, suggesting mental health needs were not met. This is in line with findings of a review on increases in depression, anxiety, and psychological distress after the onset of the pandemic,²¹⁷ with three studies reporting these were greater for children and young people who were already living with

a mental health condition before the pandemic. Our Review found some evidence of increased depression and anxiety within this clinical population. Exacerbated symptoms of some mental health problems among children and young people might have been due to school closures, disruptions to daily routines, reduced access to mental health services, and less life-acquired resilience,^{217–219} but the low quality of evidence on these observations needs to be noted.

Panel: Lived experience commentary by Sje, KM, and PS

We have attended the research team's regular meetings, and the lead researcher arranged additional meetings to help us make sense of the large amount of data and conflicting information. With little hands-on involvement in the project, and our limited experience of systematic reviews, we found this study particularly challenging to follow and comment on. Throughout, our questions returned to our own experiences, and those heard through our peer networks and from people interviewed in our previous studies.^{3,214}

Only four of the 177 original studies reported on lived experience involvement in their research. In their haste to understand the COVID-19 response, academic researchers reported data from use of services, but we could not see evidence of them having asked people with lived experience, carers, or their wider communities for input into the design or interpretation of the studies. Involvement in these earlier individual studies might have alerted those researchers to missing perspectives, such as the insights related by peer supporters when they described their work as becoming more serious in response to pandemic-related experiences of loss.²⁰⁹

There are multifarious findings of varying degrees of certainty in the quantitative studies reviewed. However, these studies tell us little about why the changes happened or about the trajectory of people who did not or could not access statutory services. Similarly, reporting of service contacts data is no indicator of the quality or outcomes of those contacts. Our experience is of services becoming inaccessible, and access through emergency departments that could have been avoided if other community services had been available or readily accessible. Alternatives such as digital services did not work for everyone and were often offered with little other choice; pressures in one part of the system created knock-on effects elsewhere. There were very little demographic data reported in the original studies or consideration of differential sub-group effects, despite the recognised disproportionate effect of the pandemic on marginalised communities.

A cautious approach to drawing conclusions from the conflicting and low certainty findings is needed. However, this study is valuable in highlighting the complex effects of the pandemic, uncertainties and gaps in knowledge, points of potential concern, and the necessity of learning from qualitative approaches, experiential perspectives, and the peer support approaches, which filled many of the gaps when traditional services were harder to reach.

To our knowledge, our Review offers the most comprehensive summary of epidemiological patterns in mental health and mental health care in Europe during the COVID-19 pandemic to date. The Review encompasses both the general population and those with pre-existing conditions, and allows a comparison of changes in prevalence of mental health problems and mental health service use. Use of the GRADE framework to assess quality of evidence for each outcome and its integration within our narrative synthesis adds robustness to our conclusions.

Our Review has several limitations. Studies included for research question one often used cutoffs on symptom measures rather than validated diagnostic instruments to measure prevalence, possibly inflating estimates. Measurement of incidence and some other outcomes such as self-harm was generally based on service contacts, with reported results likely to reflect reduced service provision and impediments to seeking help during the pandemic. We looked at broad aggregated shifts across Europe as the pandemic progressed. Commonalities in experiences of the pandemic reported in our Review are likely to be greater than in reviews of global scope. However, variations in the timing of pandemic waves and the extent of social restrictions were still too great for us to examine in detail their relationship with changes in mental health. The numbers of studies per country for any outcome were generally too small for formal comparisons between countries, but we observed no striking between-country differences. Limitations within the included studies also restrict our conclusions. First, although we include more recent studies than other reviews, we found few publications relating to 2021 and beyond, which contributed a low certainty of evidence for findings later in the pandemic. Second, the certainty of evidence regarding some outcomes (eg, incidence and mental health in people with pre-existing mental health conditions) was restricted by small numbers of studies for each outcome. Third, there was considerable variation between studies in the timepoints used, particularly in how far before March, 2020, pre-pandemic data had been collected. We have aimed for clarity regarding the stage of the pandemic at which data were collected, but some loss of detail has occurred in aggregating studies, including in terms of patterns by sex or gender and age, which are beyond the scope of synthesis for this Review. Fourth, there are important groups for whom evidence is so far insufficient, including people with psychosis and bipolar disorder, and groups at particular risk of adverse effects from COVID-19, such as minoritised ethnic groups. Fifth, people with relevant lived experience had rarely been involved in planning, conduct, or interpretation of studies.

Further research using health-care records or qualitative methods could continue to shed light on the effect of the pandemic on mental health and service use, and the experiences underlying these observations.

Further research that carefully distinguishes long-term trajectories in mental health and service use from changes connected to the pandemic is needed to understand the long-term effects of the pandemic on mental health and psychiatric service use. Our study can be repeated on an international scale, including research from low-income and middle-income countries, to provide a fuller picture of how the pandemic affected global mental health. Evidence gaps regarding conditions such as psychosis, bipolar disorder, eating disorders, and people with a personality diagnosis warrant further research, although suitable pre-pandemic comparators are sometimes lacking. More fine-grained quantitative and qualitative investigations of the experiences of groups particularly at risk of adverse outcomes are also warranted and of drivers of variations both in the general population and among people with pre-existing mental health conditions.

Contributors

Sjo and BL-E wrote the original study proposal. NA drafted the study protocol with revisions by PB, NL, CL, KM, PS, SMA, Sje, LM, PG, ST, SI, JK, RS, PG, BL-E, and SJ. PB and NA led the searching and data collection processes. ST, SI, ERF, UF, RA, MS, NL, PS, SMA, KRKS, LSR, SH, OK, AG, and TP have done the screening, data extraction, and quality assessment of papers. PB, AG, and TP led the analysis of the data. NA, TS, CL, KM, PS, Sje, LM, PG, ST, SI, ERF, HB, JK, RS, PG, BL-E, and SJ contributed to the analysis and interpretation of data. JK, PG, RS, Sjo, and BL-E provided subject expertise and methodological guidance. NA, AG, PB, and TP wrote the initial draft of the manuscript. All authors contributed to consecutive drafts and approved the final manuscript.

Declaration of interests

RS reports grants from Janssen Pharmaceuticals and Takeda Pharmaceutical Company to their institution in the past 36 months. RS also reports supervising a part-time PhD of a GSK employee at their institution. All other authors declare no competing interests.

Acknowledgments

This paper presents independent research commissioned and funded by the National Institute for Health and Care Research (NIHR) Policy Research Programme, conducted by the NIHR Policy Research Unit in Mental Health. The views expressed are those of the authors and not necessarily those of the NIHR, the Department of Health and Social Care or its arm's length bodies, or other Government departments. RS is partly funded by the NIHR Biomedical Research Centre at the South London, Maudsley National Health Service Foundation Trust, and King's College London; NIHR Applied Research Collaboration South London at King's College Hospital National Health Service Foundation Trust; the DATAMIND Health Data Research UK Mental Health Data Hub (MRC grant MR/W014386); and the UK Prevention Research Partnership (Violence, Health and Society; MR-VO49879/1), an initiative funded by UK Research and Innovation Councils, the Department of Health and Social Care (England), the UK devolved administrations, and leading health research charities. ERF is funded by the Medical Research Council (MRC Doctoral Training Programme: MR/N013867/1).

References

- 1 Sheridan Rains L, Johnson S, Barnett P, et al. Early impacts of the COVID-19 pandemic on mental health care and on people with mental health conditions: framework synthesis of international experiences and responses. *Soc Psychiatry Psychiatr Epidemiol* 2021; **56**: 13–24.
- 2 Johnson S, Dalton-Locke C, Vera San Juan N, et al. Impact on mental health care and on mental health service users of the COVID-19 pandemic: a mixed methods survey of UK mental health care staff. *Soc Psychiatry Psychiatr Epidemiol* 2021; **56**: 25–37.
- 3 Gillard S, Dare C, Hardy J, et al. Experiences of living with mental health problems during the COVID-19 pandemic in the UK: a coproduced, participatory qualitative interview study. *Soc Psychiatry Psychiatr Epidemiol* 2021; **56**: 1447–57.

- 4 Holmes EA, O'Connor RC, Perry VH, et al. Multidisciplinary research priorities for the COVID-19 pandemic: a call for action for mental health science. *Lancet Psychiatry* 2020; **7**: 547–60.
- 5 Prati G, Mancini AD. The psychological impact of COVID-19 pandemic lockdowns: a review and meta-analysis of longitudinal studies and natural experiments. *Psychol Med* 2021; **51**: 201–11.
- 6 Xiong J, Lipsitz O, Nasri F, et al. Impact of COVID-19 pandemic on mental health in the general population: a systematic review. *J Affect Disord* 2020; **277**: 55–64.
- 7 Akinin LB, De Neve J-E, Dunn EW, et al. Mental health during the first year of the COVID-19 pandemic: a review and recommendations for moving forward. *Perspect Psychol Sci* 2022; **17**: 915–36.
- 8 Sun Y, Wu Y, Fan S, et al. Comparison of mental health symptoms before and during the covid-19 pandemic: evidence from a systematic review and meta-analysis of 134 cohorts. *BMJ* 2023; **380**: e074224.
- 9 Lee Y, Lui LMW, Chen-Li D, et al. Government response moderates the mental health impact of COVID-19: a systematic review and meta-analysis of depression outcomes across countries. *J Affect Disord* 2021; **290**: 364–77.
- 10 Robinson E, Sutin AR, Daly M, Jones A. A systematic review and meta-analysis of longitudinal cohort studies comparing mental health before versus during the COVID-19 pandemic in 2020. *J Affect Disord* 2022; **296**: 567–76.
- 11 Cénat JM, Farahi SMMM, Dalexis RD, et al. The global evolution of mental health problems during the COVID-19 pandemic: a systematic review and meta-analysis of longitudinal studies. *J Affect Disord* 2022; **315**: 70–95.
- 12 Santomauro DF, Herrera AMM, Shadid J, et al. Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *Lancet* 2021; **398**: 1700–12.
- 13 Hale T, Angrist N, Goldszmidt R, et al. A global panel database of pandemic policies (Oxford COVID-19 Government Response Tracker). *Nat Hum Behav* 2021; **5**: 529–38.
- 14 Lionello L, Stranges D, Karki T, et al. Non-pharmaceutical interventions in response to the COVID-19 pandemic in 30 European countries: the ECDC-JRC Response Measures Database. *Euro Surveill* 2022; **27**: 2101190.
- 15 Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *Syst Rev* 2021; **10**: 89.
- 16 Ayiku L, Hudson T, Williams C, Levay P, Jacob C. The NICE OECD countries' geographic search filters: part 2-validation of the MEDLINE and Embase (Ovid) filters. *J Med Libr Assoc* 2021; **109**: 583–89.
- 17 Ayiku L, Levay P, Hudson T. The NICE OECD countries' geographic search filters: part 1-methodology for developing the draft MEDLINE and Embase (Ovid) filters. *J Med Libr Assoc* 2021; **109**: 258–66.
- 18 Thomas J, Graziosi S, Brunton J, et al. EPPI-Reviewer: advanced software for systematic reviews, maps and evidence synthesis. EPPI-Centre. UCL Social Research Institute, University College London, 2022.
- 19 Herzog R, Álvarez-Pasquin MJ, Díaz C, Del Barrio JL, Estrada JM, Gil Á. Are healthcare workers' intentions to vaccinate related to their knowledge, beliefs and attitudes? A systematic review. *BMC Public Health* 2013; **13**: 154.
- 20 Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008; **336**: 924–26.
- 21 Murad MH, Mustafa RA, Schünemann HJ, Sultan S, Santesso N. Rating the certainty in evidence in the absence of a single estimate of effect. *Evid Based Med* 2017; **22**: 85–87.
- 22 Fasshauer JM, Bollmann A, Hohenstein S, et al. Psychiatric emergency admissions and inpatient length of stay before and during the COVID-19 pandemic in Germany. *Psychiatr Prax* 2022; **49**: 271–75 (in Psychiatrische Notfallaufnahmen und stationäre Aufenthaltsdauer vor und während der COVID-19-Pandemie in Deutschland).
- 23 Mergel E, Schützwahl M. COVID-19 and the trajectory of participation, inclusion, and psychological well-being among adults with and without mental disorders—follow-up study from Germany. *Psychiatr Prax* 2021; **48**: 265–68 (in COVID-19 und psychisches Befinden im zweiten Lockdown –Fortsetzung einer Verlaufsuntersuchung).
- 24 Naumann E, von den Driesch E, Schumann A, Thönnissen C. Increase of depressive symptoms among adolescents during the first COVID-19 lockdown in Germany: results from the German family panel pairfam. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz* 2021; **64**: 1533–40 (in Thönnissen C. Anstieg depressiver Symptome bei Jugendlichen und jungen Erwachsenen während des ersten Lockdowns in Deutschland Ergebnisse des Beziehungs- und Familienpanels pairfam).
- 25 Briggs R, McDowell CP, De Looze C, Kenny RA, Ward M. Depressive symptoms among older adults pre- and post-COVID-19 pandemic. *J Am Med Dir Assoc* 2021; **22**: 2251–57.
- 26 Burdzovic Andreas J, Brunborg GS. Self-reported mental and physical health among Norwegian adolescents before and during the COVID-19 pandemic. *JAMA Netw Open* 2021; **4**: e2121934.
- 27 Conceicao V, Rothes I, Gusmao R. The association between changes in the university educational setting and peer relationships: effects in students' depressive symptoms during the COVID-19 pandemic. *Front Psychiatry* 2021; **12**: 783776.
- 28 Dale R, Budimir S, Probst T, Stippl P, Pieh C. Mental health during the COVID-19 lockdown over the Christmas period in Austria and the effects of sociodemographic and lifestyle factors. *Int J Environ Res Public Health* 2021; **18**: 3679.
- 29 Dickerson J, Kelly B, Lockyer B, et al. 'When will this end? Will it end?' The impact of the March–June 2020 UK COVID-19 lockdown response on mental health: a longitudinal survey of mothers in the Born in Bradford study. *BMJ Open* 2022; **12**: e047748.
- 30 Ebrahimi Omid V, Hoffart A, Johnson Sverre U. Physical distancing and mental health during the COVID-19 pandemic: factors associated with psychological symptoms and adherence to pandemic mitigation strategies. *Clin Psychol Sci* 2021; **9**: 489–506.
- 31 Evans S, Alkan E, Bhangoo JK, Tenenbaum H, Ng-Knight T. Effects of the COVID-19 lockdown on mental health, wellbeing, sleep, and alcohol use in a UK student sample. *Psychiatry Res* 2021; **298**: 113819.
- 32 Fischer K, Tieskens JM, Luijten MAJ, et al. Internalizing problems before and during the COVID-19 pandemic in independent samples of Dutch children and adolescents with and without pre-existing mental health problems. *Eur Child Adolesc Psychiatry* 2022; published online May 26. <https://doi.org/10.1007/s00787-022-01991-y>.
- 33 Fransson E, Karalexi M, Kimmel M, et al. Mental health among pregnant women during the pandemic in Sweden—a mixed methods approach using data from the Mom2B mobile application for research. *MedRxiv* 2020; published online Dec 20. <https://doi.org/10.1101/2020.12.18.20248466> (preprint).
- 34 Gosselin A, Melchior M, Carillon S, et al. Deterioration of mental health and insufficient COVID-19 information among disadvantaged immigrants in the greater Paris area. *J Psychosom Res* 2021; **146**: 110504.
- 35 Hyland P, Shevlin M, Murphy J, et al. A longitudinal assessment of depression and anxiety in the Republic of Ireland before and during the COVID-19 pandemic. *Psychiatry Res* 2021; **300**: 113905.
- 36 Hyland P, Vallières F, McBride O, et al. Mental health of adults in Ireland during the first year of the COVID-19 pandemic: results from a nationally representative, longitudinal study. *Psychol Med* 2021; published online Oct 11. <https://doi.org/10.1017/S0033291721004360>.
- 37 Kostev K, Weber K, Riedel-Heller S, von Vultée C, Bohlken J. Increase in depression and anxiety disorder diagnoses during the COVID-19 pandemic in children and adolescents followed in pediatric practices in Germany. *Eur Child Adolesc Psychiatry* 2021; **26**: 1–7.
- 38 Kwong ASF, Pearson RM, Adams MJ, et al. Mental health before and during the COVID-19 pandemic in two longitudinal UK population cohorts. *Br J Psychiatry* 2021; **218**: 334–43.
- 39 Luijten MAJ, van Muilekom MM, Teela L, et al. The impact of lockdown during the COVID-19 pandemic on mental and social health of children and adolescents. *Qual Life Res* 2021; **30**: 2795–804.
- 40 Marmet S, Wicki M, Gmel G, et al. The psychological impact of the COVID-19 crisis on young Swiss men participating in a cohort study. *Swiss Med Wkly* 2021; **151**: w30028.
- 41 McLafferty M, Brown N, McHugh R, et al. Depression, anxiety and suicidal behaviour among college students: comparisons pre-COVID-19 and during the pandemic. *Psychiatry Res Commun* 2021; **1**: 100012.

- 42 Medda E, Toccaceli V, Gigantesco A, Picardi A, Fagnani C, Stazi MA. The COVID-19 pandemic in Italy: depressive symptoms immediately before and after the first lockdown. *J Affect Disord* 2022; **298**: 202–8.
- 43 Muro A, Feliu-Soler A, Castellà J. Psychological impact of COVID-19 lockdowns among adult women: the predictive role of individual differences and lockdown duration. *Women Health* 2021; **61**: 668–79.
- 44 Novotny JS, Gonzalez-Rivas JP, Kunzova S, et al. Risk factors underlying COVID-19 lockdown-induced mental distress. *Front Psychiatry* 2020; **11**: 603014.
- 45 Peters A, Rospleszcz S, Greiser KH, Dallavalle M, Berger K. The impact of the COVID-19 pandemic on self-reported health: early evidence from the German National Cohort. *Dtsch Arztebl Int* 2020; **117**: 861–67.
- 46 Rutland-Lawes J, Wallinheimo AS, Evans SL. Risk factors for depression during the COVID-19 pandemic: a longitudinal study in middle-aged and older adults. *BJPsych Open* 2021; **7**: e161.
- 47 van den Berg YHM, Burk WJ, Cillessen AHN, Roelofs K. Emerging adults' mental health during the COVID-19 pandemic: a prospective longitudinal study on the importance of social support. *Emerg Adulthood* 2021; **9**: 618–30.
- 48 Vatcheva T, Mostaert A, Van Ingelgem V, Henrion E, Legros L. Impact of COVID-19 pandemic on postpartum depression among mothers of extreme and early preterm infants. *Int J Gynaecol Obstet* 2021; **155**: 490–95.
- 49 Vloo A, Alessie RJM, Mierau JO, Lifelines Corona Research Initiative. Gender differences in the mental health impact of the COVID-19 lockdown: longitudinal evidence from the Netherlands. *SSM Popul Health* 2021; **15**: 100878.
- 50 Volken T, Zysset A, Amendola S, et al. Depressive symptoms in Swiss university students during the COVID-19 pandemic and its correlates. *Int J Environ Res Public Health* 2021; **18**: 1458.
- 51 Volken T, Zysset A, Amendola S, von Wyl A, Dratva J. Generalized anxiety among Swiss health professions and non-health professions students: an open cohort study over 14 months in the COVID-19 pandemic. *Int J Environ Res Public Health* 2021; **18**: 10833.
- 52 Winkler P, Formanek T, Mlada K, et al. Increase in prevalence of current mental disorders in the context of COVID-19: analysis of repeated nationwide cross-sectional surveys. *Epidemiol Psychiatr Sci* 2020; **29**: e173.
- 53 Winkler P, Mohrova Z, Mlada K, et al. Prevalence of current mental disorders before and during the second wave of COVID-19 pandemic: an analysis of repeated nationwide cross-sectional surveys. *J Psychiatr Res* 2021; **139**: 167–71.
- 54 Zaninotto P, Iob E, Demakakos P, Steptoe A. Immediate and longer-term changes in the mental health and well-being of older adults in England during the COVID-19 pandemic. *JAMA Psychiatry* 2022; **79**: 151–59.
- 55 Zilver SJM, Broekman BFP, Hendrix YMGA, et al. Stress, anxiety and depression in 1466 pregnant women during and before the COVID-19 pandemic: a Dutch cohort study. *J Psychosom Obstet Gynaecol* 2021; **42**: 108–14.
- 56 Ayaz R, Hocaoglu M, Günay T, Yardımcı OD, Turgut A, Karateke A. Anxiety and depression symptoms in the same pregnant women before and during the COVID-19 pandemic. *J Perinat Med* 2020; **48**: 965–70.
- 57 Daly M, Sutin AR, Robinson E. Longitudinal changes in mental health and the COVID-19 pandemic: evidence from the UK household longitudinal study. *Psychol Med* 2020; **52**: 2549–58.
- 58 Hafstad GS, Sætnen SS, Wentzel-Larsen T, Augusti EM. Adolescents' symptoms of anxiety and depression before and during the Covid-19 outbreak—a prospective population-based study of teenagers in Norway. *Lancet Reg Health Eur* 2021; **5**: 100093.
- 59 Knudsen AKS, Stene-Larsen K, Gustavson K, et al. Prevalence of mental disorders, suicidal ideation and suicides in the general population before and during the COVID-19 pandemic in Norway: a population-based repeated cross-sectional analysis. *Lancet Reg Health Eur* 2021; **4**: 100071.
- 60 Niedzwiedz CL, Green MJ, Benzeval M, et al. Mental health and health behaviours before and during the initial phase of the COVID-19 lockdown: longitudinal analyses of the UK household longitudinal study. *J Epidemiol Community Health* 2021; **75**: 224–31.
- 61 Pierce M, Hope H, Ford T, et al. Mental health before and during the COVID-19 pandemic: a longitudinal probability sample survey of the UK population. *Lancet Psychiatry* 2020; **7**: 883–92.
- 62 Sharp ML, Serfioti D, Jones M, et al. UK veterans' mental health and well-being before and during the COVID-19 pandemic: a longitudinal cohort study. *BMJ Open* 2021; **11**: e049815.
- 63 van der Velden Peter G, Contino C, Das M, Leenen J, Wittmann L. Differences in mental health problems, coping self-efficacy and social support between adults victimised before and adults victimised after the COVID-19 outbreak: population-based prospective study. *Br J Psychiatry* 2022; **220**: 265–71.
- 64 van der Velden PG, Contino C, Das M, van Loon P, Bosmans MWG. Anxiety and depression symptoms, and lack of emotional support among the general population before and during the COVID-19 pandemic. a prospective national study on prevalence and risk factors. *J Affect Disord* 2020; **277**: 540–48.
- 65 van der Velden PG, Hyland P, Contino C, von Gaudecker HM, Muffels R, Das M. Anxiety and depression symptoms, the recovery from symptoms, and loneliness before and after the COVID-19 outbreak among the general population: findings from a Dutch population-based longitudinal study. *PLoS One* 2021; **16**: e0245057.
- 66 Thorisdottir IE, Asgeirsdottir BB, Kristjansson AL, et al. Depressive symptoms, mental wellbeing, and substance use among adolescents before and during the COVID-19 pandemic in Iceland: a longitudinal, population-based study. *Lancet Psychiatry* 2021; **8**: 663–72.
- 67 Tavalacci MP, Ladner J, Déchelotte P. Sharp increase in eating disorders among university students since the COVID-19 pandemic. *Nutrients* 2021; **13**: 28.
- 68 Bendau A, Plag J, Kunas S, Wyka S, Ströhle A, Petzold MB. Longitudinal changes in anxiety and psychological distress, and associated risk and protective factors during the first three months of the COVID-19 pandemic in Germany. *Brain Behav* 2021; **11**: e01964.
- 69 Gullo S, Misici I, Teti A, Liuzzi M, Chiara E. Going through the lockdown: a longitudinal study on the psychological consequences of the coronavirus pandemic. *Res Psychother* 2021; **23**: 494.
- 70 O'Connor RC, Wetherall K, Cleare S, et al. Mental health and well-being during the COVID-19 pandemic: longitudinal analyses of adults in the UK COVID-19 mental health & wellbeing study. *Br J Psychiatry* 2021; **218**: 326–33.
- 71 Ripoll J, Contreras-Martos S, Esteva M, Soler A, Serrano-Ripoll MJ. Mental health and psychological wellbeing during the COVID-19 lockdown: a longitudinal study in the Balearic Islands (Spain). *J Clin Med* 2021; **10**: 20.
- 72 Charbonnier E, Le Vigouroux S, Goncalves A. Psychological vulnerability of French university students during the COVID-19 pandemic: a four-wave longitudinal survey. *Int J Environ Res Public Health* 2021; **18**: 15.
- 73 Daimer S, Mihatsch L, Roman L, Murray GK, Knolle F. Subjective impact of the COVID-19 pandemic on schizotypy and general mental health in Germany and the United Kingdom, for independent samples in May and in October 2020. *Front Psychol* 2021; **12**: 667848.
- 74 Efstathiou V, Michopoulos I, Yotsidi V, et al. Does suicidal ideation increase during the second COVID-19 lockdown? *Psychiatry Res* 2021; **301**: 113990.
- 75 Georgieva I, Lepping P, Bozev V, et al. Prevalence, new incidence, course, and risk factors of PTSD, depression, anxiety, and panic disorder during the COVID-19 pandemic in 11 countries. *Healthcare (Basel, Switzerland)* 202; **9**: 664.
- 76 Jia R, Ayling K, Chalder T, et al. The prevalence, incidence, prognosis and risk factors for symptoms of depression and anxiety in a UK cohort during the COVID-19 pandemic. *BJPsych Open* 2022; **8**: e64.
- 77 Johnson MS, Skjerdingstad N, Ebrahimi OV, Hoffart A, Urnes Johnson S. Mechanisms of parental distress during and after the first COVID-19 lockdown phase: a two-wave longitudinal study. *PLoS One* 2021; **16**: e0253087.
- 78 Myall K, Montero-Marin J, Kuyken W. Anxiety and depression during COVID-19 in elite rugby players: the role of mindfulness skills. *Int J Environ Res Public Health* 2021; **18**: 13.
- 79 Pieh C, Budimir S, Humer E, Probst T. Comparing mental health during the COVID-19 lockdown and 6 months after the lockdown in Austria: a longitudinal study. *Front Psychiatry* 2021; **12**: 625973.
- 80 Van Baelen L, Gremeaux L, Antoine J, et al. COVID-19 and people who use drugs: impact of the pandemic on general anxiety and depressive disorders among adults in Belgium. *J Affect Disord* 2021; **295**: 946–53.

- 81 Hajek A, Sabat I, Neumann-Böhme S, et al. Prevalence and determinants of probable depression and anxiety during the COVID-19 pandemic in seven countries: longitudinal evidence from the European COVID Survey (ECOS). *J Affect Disord* 2022; **299**: 517–24.
- 82 Jakubowski TD, Sitko-Dominik MM. Teachers' mental health during the first two waves of the COVID-19 pandemic in Poland. *PLoS ONE* 2021; **16**: e0257252.
- 83 Jia R, Knight H, Ayling K, et al. Prospective examination of mental health in university students during the COVID-19 pandemic. *MedRxiv* 2021; published online July 31. <https://doi.org/10.1101/2021.07.29.21261196> (preprint).
- 84 Laham S, Bertuzzi L, Deguen S, et al. Impact of longitudinal social support and loneliness trajectories on mental health during the COVID-19 pandemic in France. *Int J Environ Res Public Health* 2021; **18**: 12677.
- 85 Macalli M, Texier N, Schück S, Côté SM, Tzourio C. A repeated cross-sectional analysis assessing mental health conditions of adults as per student status during key periods of the COVID-19 epidemic in France. *Sci Rep* 2021; **11**: 21455.
- 86 Owens M, Townsend E, Hall E, Bhatia T, Fitzgibbon R, Miller-Lakin F. Mental health and wellbeing in young people in the UK during lockdown (COVID-19). *Int J Environ Res Public Health* 2022; **19**: 1132.
- 87 Tang N, McEnery K, Chandler L, et al. Pandemic and student mental health: a cross-sectional and longitudinal analysis of mental health symptoms amongst university students and young adults after the first cycle of lockdown in the UK. *BJPsych Open* 2022; **8**: e138.
- 88 Benatov J, Ochnik D, Rogowska AM, Arzensek A, Bitenc UM. Prevalence and sociodemographic predictors of mental health in a representative sample of young adults from Germany, Israel, Poland, and Slovenia: a longitudinal study during the COVID-19 pandemic. *Int J Environ Res Public Health* 2022; **19**: 1334.
- 89 Fernandes DV, Canavarro MC, Moreira H. The role of mothers' self-compassion on mother-infant bonding during the COVID-19 pandemic: a longitudinal study exploring the mediating role of mindful parenting and parenting stress in the postpartum period. *Infant Ment Health J* 2021; **42**: 621–35.
- 90 Hetkamp M, Schweda A, Bäuerle A, et al. Sleep disturbances, fear, and generalized anxiety during the COVID-19 shut down phase in Germany: relation to infection rates, deaths, and German stock index DAX. *Sleep Med* 2020; **75**: 350–53.
- 91 Rogowska AM, Ochnik D, Kuśniercz C, et al. Changes in mental health during three waves of the COVID-19 pandemic: a repeated cross-sectional study among Polish university students. *BMC Psychiatry* 2021; **21**: 627.
- 92 Ubillos-Landa S, Puente-Martínez A, González-Castro JL. Psychological withdrawal and mental health during the COVID-19 pandemic: a longitudinal study. *Psychol Health* 2021; published online Dec 26. <https://doi.org/10.1080/08870446.2021.2019254>.
- 93 Chandola T, Kumari M, Booker CL, Benzeval M. The mental health impact of COVID-19 and lockdown-related stressors among adults in the UK. *Psychol Med* 2020; **52**: 2997–3006.
- 94 Geirdal AKO, Price D, Schoultz M, et al. The significance of demographic variables on psychosocial health from the early stage and nine months after the COVID-19 pandemic outbreak: A cross-national study. *Int J Environ Res Public Health* 2021; **18**: 20.
- 95 Shevlin M, Butter S, McBride O, et al. Refuting the myth of a 'tsunami' of mental ill-health in populations affected by COVID-19: evidence that response to the pandemic is heterogeneous, not homogeneous. *Psychol Med* 2021; published online April 20. <https://doi.org/10.1017/S0033291721001665>.
- 96 Daly M, Robinson E. Psychological distress associated with the second COVID-19 wave: prospective evidence from the UK household longitudinal study. *J Affect Disord* 2022; **310**: 274–78.
- 97 Liu S, Haucke MN, Heinzel S, Heinz A. long-term impact of economic downturn and loneliness on psychological distress: triple crises of COVID-19 pandemic. *J Clin Med* 2021; **10**: 4596.
- 98 Carr MJ, Steeg S, Webb RT, et al. Effects of the COVID-19 pandemic on primary care-recorded mental illness and self-harm episodes in the UK: a population-based cohort study. *Lancet Public Health* 2021; **6**: e124–35.
- 99 Kazlauskas E, Gelezelyte O, Nomeikaite A, Zelviene P. Posttraumatic stress disorder and adjustment disorder in Lithuanian healthcare in 2018–2020: a nation-wide cohort study of the effects of COVID-19 pandemic. *Healthcare (Basel)* 2021; **9**: 22.
- 100 Castellini G, Cassioli E, Rossi E, et al. The impact of COVID-19 epidemic on eating disorders: a longitudinal observation of pre versus post psychopathological features in a sample of patients with eating disorders and a group of healthy controls. *Int J Eat Disord* 2020; **53**: 1855–62.
- 101 Seitz KI, Bertsch K, Herpertz SC. A prospective study of mental health during the COVID-19 pandemic in childhood trauma-exposed individuals: social support matters. *J Trauma Stress* 2021; **34**: 477–86.
- 102 Giel KE, Schurr M, Zipfel S, Junne F, Schag K. Eating behaviour and symptom trajectories in patients with a history of binge eating disorder during COVID-19 pandemic. *Eur Eat Disord Rev* 2021; **29**: 657–62.
- 103 Koenders M, Mesbah R, Spijker A, et al. Effects of the COVID-19 pandemic in a preexisting longitudinal study of patients with recently diagnosed bipolar disorder: indications for increases in manic symptoms. *Brain Behav* 2021; **11**: e2326.
- 104 Leightley D, Lavelle G, White KM, et al. Investigating the impact of COVID-19 lockdown on adults with a recent history of recurrent major depressive disorder: a multi-centre study using remote measurement technology. *BMC Psychiatry* 2021; **21**: 435.
- 105 Orhan M, Korten N, Paans N, et al. Psychiatric symptoms during the COVID-19 outbreak in older adults with bipolar disorder. *Int J Geriatr Psychiatry* 2021; **36**: 892–900.
- 106 Pan KY, Kok AAL, Eikelenboom M, et al. The mental health impact of the COVID-19 pandemic on people with and without depressive, anxiety, or obsessive-compulsive disorders: a longitudinal study of three Dutch case-control cohorts. *Lancet Psychiatry* 2021; **8**: 121–29.
- 107 Bendau A, Kunas SL, Wyka S, et al. Longitudinal changes of anxiety and depressive symptoms during the COVID-19 pandemic in Germany: the role of pre-existing anxiety, depressive, and other mental disorders. *J Anxiety Disord* 2021; **79**: 102377.
- 108 Caldiroli A, Capuzzi E, Tringali A, et al. The psychopathological impact of the SARS-CoV-2 epidemic on subjects suffering from different mental disorders: an observational retrospective study. *Psychiatry Res* 2022; **307**: 114334.
- 109 Carta MG, Ouali U, Perra A, et al. Living with bipolar disorder in the time of COVID-19: biorhythms during the severe lockdown in Cagliari, Italy, and the moderate lockdown in Tunis, Tunisia. *Front Psychiatry* 2021; **12**: 634765.
- 110 Davide P, Andrea P, Martina O, Andrea E, Davide D, Mario A. The impact of the COVID-19 pandemic on patients with OCD: effects of contamination symptoms and remission state before the quarantine in a preliminary naturalistic study. *Psychiatry Res* 2020; **291**: 113213.
- 111 Gentile A, Torales J, O'Higgins M, et al. Phone-based outpatients' follow-up in mental health centers during the COVID-19 quarantine. *Int J Soc Psychiatry* 2022; **68**: 129–33.
- 112 Nisticò V, Bertelli S, Tedesco R, et al. The psychological impact of COVID-19-related lockdown measures among a sample of Italian patients with eating disorders: a preliminary longitudinal study. *Eat Weight Disord* 2021; **26**: 2771–77.
- 113 Dalkner N, Wagner-Skacel J, Ratzenhofer M, et al. Psychological symptoms during and after Austrian first lockdown in individuals with bipolar disorder? A follow-up control-group investigation. *Int J Bipolar Disord* 2021; **9**: 16.
- 114 Carmassi C, Cordone A, Bertelloni CA, et al. A longitudinal study of post-traumatic stress, depressive, and anxiety symptoms trajectories in subjects with bipolar disorder during the COVID-19 pandemic. *Eur Psychiatry* 2022; **65**: e8.
- 115 Hendrikx LJ, Williamson C, Baumann J, Murphy D. The impact of the COVID-19 pandemic on treatment-seeking veterans in the united kingdom with preexisting mental health difficulties: a longitudinal study. *J Trauma Stress* 2021; **16**: 16.
- 116 Jelinek L, Voderholzer U, Moritz S, Carsten HP, Riesel A, Miegel F. When a nightmare comes true: change in obsessive-compulsive disorder over the first months of the COVID-19 pandemic. *J Anxiety Disord* 2021; **84**: 102493.
- 117 Seethaler M, Just S, Stötzner P, Bermppohl F, Brandl EJ. Psychosocial impact of COVID-19 pandemic in elderly psychiatric patients: a longitudinal study. *Psychiatr Q* 2021; **92**: 1439–57.
- 118 Hennigan K, McGovern M, Plunkett R, Costello S, McDonald C, Hallahan B. A longitudinal evaluation of the impact of the COVID-19 pandemic on patients with pre-existing anxiety disorders. *Ir J Psychol Med* 2021; **38**: 258–65.

- 119 Birgegård A, Abbaspour A, Borg S, et al. Longitudinal experiences and impact of the COVID-19 pandemic among people with past or current eating disorders in Sweden. *Eat Disord* 2022; **30**: 602–17.
- 120 Escolà-Gascón Á. Impact of conspiracist ideation and psychotic-like experiences in patients with schizophrenia during the COVID-19 crisis. *J Psychiatr Res* 2022; **146**: 135–48.
- 121 Murphy D, Hendriks LJ, Williamson C, Baumann J. Longitudinal survey of UK veterans with pre-existing mental health difficulties: mental health during the COVID-19 pandemic. *BMJ Mil Health* 2022; **25**: 25.
- 122 Pedersen MT, Andersen TO, Clotworthy A, et al. Time trends in mental health indicators during the initial 16 months of the COVID-19 pandemic in Denmark. *BMC Psychiatry* 2022; **22**: 25.
- 123 Conti E, Sgandurra G, De Nicola G, et al. Behavioural and emotional changes during COVID-19 lockdown in an Italian paediatric population with neurologic and psychiatric disorders. *Brain Sci* 2020; **10**: 27.
- 124 Tanir Y, Karayagmurlu A, Kaya I, et al. Exacerbation of obsessive compulsive disorder symptoms in children and adolescents during COVID-19 pandemic. *Psychiatry Res* 2020; **293**: 113363.
- 125 Raffagnato A, Iannattone S, Tascini B, et al. The COVID-19 pandemic: a longitudinal study on the emotional-behavioral sequelae for children and adolescents with neuropsychiatric disorders and their families. *Int J Environ Res Public Health* 2021; **18**: 19.
- 126 Machado PPP, Pinto-Bastos A, Ramos R, et al. Impact of COVID-19 lockdown measures on a cohort of eating disorders patients. *J Eat Disord* 2020; **8**: 57.
- 127 Bartels C, Hessmann P, Schmidt U, et al. Medium-term and peri-lockdown course of psychosocial burden during the ongoing COVID-19 pandemic: a longitudinal study on patients with pre-existing mental disorders. *Eur Arch Psychiatry Clin Neurosci* 2021; **25**: 25.
- 128 Berardelli I, Sarubbi S, Rogante E, et al. The impact of the COVID-19 pandemic on suicide ideation and suicide attempts in a sample of psychiatric inpatients. *Psychiatry Res* 2021; **303**: 114072.
- 129 Rømer TB, Christensen RHB, Blomberg SN, Folke F, Christensen HC, Benros ME. Psychiatric admissions, referrals, and suicidal behavior before and during the COVID-19 pandemic in Denmark: a time-trend study. *Acta Psychiatr Scand* 2021; **144**: 553–62.
- 130 Yalçın M, Baş A, Bilici R, et al. Psychiatric emergency visit trends and characteristics in a mental health epicenter in Istanbul during COVID-19 lockdown. *Soc Psychiatry Psychiatr Epidemiol* 2021; **56**: 2299–310.
- 131 John A, Lee SC, Solomon S, et al. Loneliness, coping, suicidal thoughts and self-harm during the COVID-19 pandemic: a repeat cross-sectional UK population survey. *BMJ Open* 2021; **0**: e048123.
- 132 Alves T, Marques M, Carvalho A. The COVID-19 pandemic and its impact on the demand for mental health care—the experience of a hospital centre in Portugal. 2021. <https://pesquisa.bvsalud.org/global-literature-on-novel-coronavirus-2019-ncov/resource/pt/covidwho-1464274> (accessed March 25, 2023).
- 133 Boldrini T, Girardi P, Clerici M, et al. Consequences of the COVID-19 pandemic on admissions to general hospital psychiatric wards in Italy: reduced psychiatric hospitalizations and increased suicidality. *Prog Neuropsychopharmacol Biol Psychiatry* 2021; **110**: 110304.
- 134 Capuzzi E, Di Brita C, Caldiroli A, et al. Psychiatric emergency care during Coronavirus 2019 (COVID 19) pandemic lockdown: results from a Department of Mental Health and Addiction of northern Italy. *Psychiatry Res* 2020; **293**: 113463.
- 135 Clerici M, Durbano F, Spinogatti F, Vita A, de Girolamo G, Micciolo R. Psychiatric hospitalization rates in Italy before and during COVID-19: did they change? An analysis of register data. *Ir J Psychol Med* 2020; **37**: 283–90.
- 136 Gesi C, Grasso F, Dragogna F, et al. How did COVID-19 affect suicidality? Data from a multicentric study in Lombardy. *J Clin Med* 2021; **10**: 29.
- 137 Gómez-Ramiro M, Fico G, Anmella G, et al. Changing trends in psychiatric emergency service admissions during the COVID-19 outbreak: report from a worldwide epicentre. *J Affect Disord* 2021; **282**: 26–32.
- 138 Gonçalves-Pinho M, Mota P, Ribeiro J, Macedo S, Freitas A. The impact of COVID-19 pandemic on psychiatric emergency department visits – a descriptive study. *Psychiatr Q* 2021; **92**: 621–31.
- 139 Montalbani B, Bargagna P, Mastrangelo M, et al. The COVID-19 outbreak and subjects with mental disorders who presented to an Italian psychiatric emergency department. *J Nerv Ment Dis* 2021; **209**: 246–50.
- 140 Nuzum E, Martin E, Broadbent M, Stewart R. Mental health service activity during COVID-19 lockdown among individuals with personality disorders: south London and Maudsley data on services and mortality from January to May 2020. *MedRxiv* 2020; published online Sep 13. <https://doi.org/10.1101/2020.09.13.20193730> (preprint).
- 141 Panariello F, Longobardi S, Cellini L, De Ronchi D, Atti AR. Psychiatric hospitalization during the two SARS-CoV-2 pandemic waves: new warnings for acute psychotic episodes and suicidal behaviors. *World J Psychiatry* 2021; **11**: 1095–105.
- 142 Pignon B, Gourevitch R, Tebeka S, et al. Dramatic reduction of psychiatric emergency consultations during lockdown linked to COVID-19 in Paris and suburbs. *MedRxiv* 2020; published online May 22. <https://doi.org/10.1101/2020.05.19.20095901> (preprint).
- 143 Revet A, Hebebrand J, Anagnostopoulos D, et al. Perceived impact of the COVID-19 pandemic on child and adolescent psychiatric services after 1 year (February/March 2021): ESCAP CovCAP survey. *Eur Child Adolesc Psychiatry* 2021; **29**: 29.
- 144 Riley B, Packer M, Gallier S, Sapey E, Atkin C. Acute, non-COVID related medical admissions during the first wave of COVID-19: a retrospective comparison of changing patterns of disease. *Acute Med* 2020; **19**: 176–82.
- 145 Sánchez-Guarnido AJ, Hidalgo N, Arenas de la Cruz J, Esteban I, Mondón S, Herruzo C. Analysis of the consequences of the COVID-19 pandemic on people with severe mental disorders. *Int J Environ Res Public Health* 2021; **18**: 13.
- 146 Seifert J, Meissner C, Birkenstock A, et al. Peripandemic psychiatric emergencies: impact of the COVID-19 pandemic on patients according to diagnostic subgroup. *Eur Arch Psychiatry Clin Neurosci* 2021; **271**: 259–70.
- 147 Stewart R, Martin E, Bakolis I, Broadbent M, Byrne N, Landau S. Comparison of mental health service activity before and shortly after UK social distancing responses to the COVID-19 pandemic: February–March 2020. *MedRxiv* 2020, published online Sep 26. <https://www.medrxiv.org/content/10.1101/2020.09.26.20202150v1>.
- 148 Stewart R, Martin E, Broadbent M. Mental health service activity during COVID-19 lockdown: south London and Maudsley data on working age community and home treatment team services and mortality from February to mid-May 2020. *MedRxiv* 2020; published online Jun 13. <https://doi.org/10.1101/2020.06.13.20130419> (preprint).
- 149 Tromans S, Chester V, Harrison H, Pankhania P, Booth H, Chakraborty N. Patterns of use of secondary mental health services before and during COVID-19 lockdown: observational study. *BJPsych Open* 2020; **6**: e117.
- 150 Carta S, Ng F. The mental health impact of COVID-19: Salisbury District Hospital. *Psychology (Irvine)* 2021; **12**: 1118–26.
- 151 Bollmann A, Hohenstein S, Pellissier V, et al. Utilization of in- and outpatient hospital care in Germany during the Covid-19 pandemic insights from the German-wide Helios hospital network. *PLOS ONE* 2021; **16**: e0249251.
- 152 Di Lorenzo R, Frattini N, Dragone D, et al. Psychiatric emergencies during the COVID-19 pandemic: 6-month observational study. *Neuropsychiatr Dis Treat* 2021; **17**: 1763–78.
- 153 Fasshauer JM, Bollmann A, Hohenstein S, et al. Impact of COVID-19 pandemic on involuntary and urgent inpatient admissions for psychiatric disorders in a German-wide hospital network. *J Psychiatr Res* 2021; **142**: 140–43.
- 154 Jollant F, Roussot A, Corruble E, et al. Hospitalization for self-harm during the early months of the COVID-19 pandemic in France: a nationwide retrospective observational cohort study. *Lancet Reg Health Eur* 2021; **6**: 100102.
- 155 Sam M, Kelbrick M. Section 136 assessments in the COVID-19 pandemic. *Prog Neurol Psychiatry* 2021; **25**: 21–23.
- 156 Ayton A, Viljoen D, Ryan S, Ibrahim A, Ford D. Risk, demand, capacity and outcomes in adult specialist eating disorder services in south-east of England before and since COVID-19. *BJPsych Bull* 2022; **46**: 89–95.
- 157 Diaz de Neira M, Blasco-Fontecilla H, Garcia M, et al. Demand analysis of a psychiatric emergency room and an adolescent acute inpatient unit in the context of the COVID-19 pandemic in Madrid, Spain. *Fron Psychiatry* 2020; **11**: 557508.

- 158 Graell M, Morón-Nozalea MG, Camarero R, et al. Children and adolescents with eating disorders during COVID-19 confinement: difficulties and future challenges. *Eur Eat Disord Rev* 2020; **28**: 864–70.
- 159 Mourouveye M, Botteman H, Bonny G. Association between suicide behaviours in children and adolescents and the COVID-19 lockdown in Paris, France: a retrospective observational study. *Arch Dis Child, BMJ Journals* 2021; **106**: E42.
- 160 Ambrosetti J, Macheret L, Folliet A, et al. Impact of the COVID-19 pandemic on psychiatric admissions to a large Swiss emergency department: an observational study. *Int J Environ Res Public Health* 2021; **18**: 28.
- 161 Balestrieri M, Rucci P, Amendola D, et al. Emergency psychiatric consultations during and after the COVID-19 lockdown in Italy. A multicentre study. *Front Psychiatry* 2021; **12**: 697058.
- 162 Beghi M, Brandolini R, Casolaro I, et al. Effects of lockdown on emergency room admissions for psychiatric evaluation: an observational study from the AUSL Romagna, Italy. *Int J Psychiatry Clin Pract* 2021; **25**: 135–39.
- 163 Beghi M, Ferraris S, Brandolini R, et al. Effects of lockdown on emergency room admissions for psychiatric evaluation: an observational study from 4 centres in Italy. *Int J Psychiatry Clin Pract* 2022; **26**: 316–320.
- 164 Chen S, She R, Qin P, et al. The medium-term impact of COVID-19 lockdown on referrals to secondary care mental health services: a controlled interrupted time series study. *Front Psychiatry* 2020; **11**: 585915.
- 165 Flament J, Scius N, Zdanowicz N, Regnier M, De Cannière L, Thonon H. Influence of post-COVID-19 deconfinement on psychiatric visits to the emergency department. *Am J Emerg Med* 2021; **48**: 238–42.
- 166 Grimshaw B, Chaudhuri E. Mental-health-related admissions to the acute medical unit during COVID-19. *Clin Med (Lond)* 2021; **21**: e77–79.
- 167 Hawton K, Casey D, Bale E, et al. Self-harm during the early period of the COVID-19 pandemic in England: comparative trend analysis of hospital presentations. *MedRxiv* 2020; published online Nov 25. <https://doi.org/10.1101/2020.11.25.20238030> (preprint).
- 168 Hoyer C, Ebert A, Szabo K, Platten M, Meyer-Lindenberg A, Kranaster L. Decreased utilization of mental health emergency service during the COVID-19 pandemic. *Eur Arch Psychiatry Clin Neurosci* 2021; **271**: 377–79.
- 169 McAndrew J, O'Leary J, Cotter D, et al. Impact of initial COVID-19 restrictions on psychiatry presentations to the emergency department of a large academic teaching hospital. *Ir J Psychol Med* 2021; **38**: 108–15.
- 170 McIntyre A, Tong K, McMahon E, Doherty AM. COVID-19 and its effect on emergency presentations to a tertiary hospital with self-harm in Ireland. *Ir J Psychol Med* 2021; **38**: 116–22.
- 171 Nuzum E, Martin E, Morgan G, et al. Self-harm presentations to emergency departments and place of safety during the 'first wave' of the UK COVID-19 pandemic: south London and Maudsley data on service use from February to June 2020. *MedRxiv* 2020; published online Dec 10. <https://doi.org/10.1101/2020.12.10.20247155> (preprint).
- 172 Olding J, Zisman S, Olding C, Fan K. Penetrating trauma during a global pandemic: changing patterns in interpersonal violence, self-harm and domestic violence in the COVID-19 outbreak. *Surgeon* 2021; **19**: e9–e13.
- 173 Shields C, Bernard J, Mirza OI, Reeves D, Wells A, Heagerty A. Covid-19, lockdown and self-isolation: evaluation of deliberate self-harm admissions. *Front Psychiatry* 2021; **12**: 662885.
- 174 Stein HC, Giordano B, Del Giudice R, Basi C, Gambini O, D'Agostino A. Pre/post comparison study of emergency mental health visits during the COVID-19 lockdown in Lombardy, Italy. *Psychiatry Clin Neurosci* 2020; **74**: 605–07.
- 175 Di Lorenzo R, Fiore G, Bruno A, et al. Urgent psychiatric consultations at mental health center during COVID-19 pandemic: retrospective observational study. *Psychiatr Q* 2021; **92**: 1341–59.
- 176 Sampson EL, Wright J, Dove J, Mukadam N. Psychiatric liaison service referral patterns during the UK COVID-19 pandemic: an observational study. *Eur J Psychiatry* 2022; **36**: 35–42.
- 177 Cousien A, Acquaviva E, Kernéis S, Yazdanpanah Y, Delorme R. Temporal trends in suicide attempts among children in the decade before and during the COVID-19 pandemic in Paris, France. *JAMA Netw Open* 2021; **4**: e2128611.
- 178 Wallis Gómez VG, Apolinario MH, Saavedra Santana P, et al. Evaluation of changes in pediatric healthcare activity during the Covid-19 state of alarm in the Canary Islands. *Public Health Pract (Oxf)* 2021; **2**: 100159.
- 179 Shanmugavadeivel D, Liu JF, Gilhooley C, Elsaadany L, Wood D. Changing patterns of emergency paediatric presentations during the first wave of COVID-19: learning for the second wave from a UK tertiary emergency department. *BMJ Paediatr Open* 2021; **5**: e000967.
- 180 Carlin GL, Baumgartner JS, Moffakhar T, König D, Negrin LL. Impact of COVID-19 lockdown on suicide attempts: a retrospective analysis of the springtime admissions to the trauma resuscitation room at the Medical University of Vienna from 2015–2020. *Wien Klin Wochenschr* 2021; **133**: 915–22.
- 181 Hay D, Jamal MS, Al-Tawil K, et al. The effect of the COVID-19 pandemic on mental health associated trauma, admissions and fractures at a London major trauma centre. *Ann R Coll Surg Engl* 2021; **103**: 114–19.
- 182 Nia A, Popp D, Diendorfer C, et al. Impact of lockdown during the COVID-19 pandemic on number of patients and patterns of injuries at a level I trauma center. *Wien Klin Wochenschr* 2021; **133**: 336–43.
- 183 Fasshauer JM, Bollmann A, Hohenstein S, et al. Emergency hospital admissions for psychiatric disorders in a German-wide hospital network during the COVID-19 outbreak. *Soc Psychiatry Psychiatr Epidemiol* 2021; **56**: 1469–75.
- 184 Capuzzi E, Caldiroli A, Di Brita C, et al. Profile of patients attending psychiatric emergency care during the coronavirus 2019 (COVID 19) pandemic: a comparative cross-sectional study between lockdown and post-lockdown periods in Lombardy, Italy. *Int J Psychiatry Clin Pract* 2022; **26**: 132–138.
- 185 Ambrosetti J, Macheret L, Folliet A, et al. Psychiatric emergency admissions during and after COVID-19 lockdown: short-term impact and long-term implications on mental health. *BMC Psychiatry* 2021; **21**: 465.
- 186 Bauer-Staeb C, Davis A, Smith T, et al. Impact of COVID-19 on primary care mental health services: a descriptive, cross-sectional timeseries of electronic healthcare records. *MedRxiv* 2020; published online Aug 17 <https://doi.org/10.1101/2020.08.15.20175562> (preprint).
- 187 Chow MW, Noorthoorn EO, Wierdsma AI, et al. Impact of the first COVID-19 outbreak on mental health service utilisation at a Dutch mental health centre: retrospective observational study. *BJPsych Open* 2021; **7**: e213.
- 188 Aragona M, Barbato A, Cavani A, Costanzo G, Mirisola C. Negative impacts of COVID-19 lockdown on mental health service access and follow-up adherence for immigrants and individuals in socio-economic difficulties. *Public Health* 2020; **186**: 52–56.
- 189 Hong JS, Sheriff R, Smith K, et al. Impact of COVID-19 on telepsychiatry at the service and individual patient level across two UK NHS mental health Trusts. *Evid Based Ment Health* 2021; **24**: 161–66.
- 190 Howarth A, Munro M, Theodorou A, Mills PR. Trends in healthcare utilisation during COVID-19: a longitudinal study from the UK. *BMJ Open* 2021; **11**: e048151.
- 191 Svaleryd Helena B, Björkegren E, Vlachos J. The impact of the COVID-19 school closure on adolescents' use of mental healthcare services in Sweden. *MedRxiv* 2021; published online Dec 21. <https://doi.org/10.1101/2021.12.12.21267684> (preprint).
- 192 Michalowsky B, Hoffmann W, Bohlken J, Kostev K. Effect of the COVID-19 lockdown on disease recognition and utilisation of healthcare services in the older population in Germany: a cross-sectional study. *Age Ageing* 2021; **50**: 317–25.
- 193 Mueller C, Perera G, Broadbent M, Stewart R, Velayudhan L. A retrospective analysis of patient flow in mental health services for older adults in south London during the COVID-19 pandemic. *Int Psychogeriatr* 2022; **34**: 1–2.
- 194 van der Velden Peter G, Marchand M, Das M, Muffels R, Bosmans M. The prevalence, incidence and risk factors of mental health problems and mental health services use before and 9 months after the COVID-19 outbreak among the general Dutch population. A 3-wave prospective study. *MedRxiv* 2021; published online March 2. <https://doi.org/10.1101/2021.02.27.21251952> (preprint).

- 195 Humer E, Haid B, Schimböck W, et al. Provision of psychotherapy one year after the beginning of the COVID-19 pandemic in Austria. *Int J Environ Res Public Health* 2021; **18**: 5843.
- 196 Williams R, Jenkins DA, Ashcroft DM, et al. Diagnosis of physical and mental health conditions in primary care during the COVID-19 pandemic: a retrospective cohort study. *Lancet Public Health* 2020; **5**: e543–50.
- 197 Hvide HK, Johnsen J. COVID-19 and mental health: a longitudinal population study from Norway. *Eur J Epidemiol* 2022; **37**: 167–72.
- 198 Lemanska A, Hoang U, Jeffreys N, et al. Study into COVID-19 crisis using primary care mental health consultations and prescriptions data. *Stud Health Technol Inform* 2021; **281**: 759–63.
- 199 Mansfield KE, Mathur R, Tazare J, et al. Indirect acute effects of the COVID-19 pandemic on physical and mental health in the UK: a population-based study. *Lancet Digit Health* 2021; **3**: e217–30.
- 200 Hernández-Calle D, Martínez-Alés G, Mediavilla R, Aguirre P, Rodríguez-Vega B, Bravo-Ortiz MF. Trends in psychiatric emergency department visits due to suicidal ideation and suicide attempts during the COVID-19 pandemic in Madrid, Spain. *J Clin Psychiatry* 2020; **81**: 21721.
- 201 European Centre for Disease Prevention and Control. Data on country response measures to COVID-19. European Centre for Disease Prevention and Control, 2022. <https://www.ecdc.europa.eu/en/publications-data/download-data-response-measures-covid-19> (accessed Dec 3, 2022).
- 202 Shankar A, Yu BE, Malvankar-Mehta M. The psychological impact of COVID-19 on socially isolated individuals—a systematic review. *Ment Health Rev (Brighton)* 2021; **26**: 247–57.
- 203 Murphy L, Markey K, O’Donnell C, Moloney M, Doody O. The impact of the COVID-19 pandemic and its related restrictions on people with pre-existent mental health conditions: a scoping review. *Arch Psychiatr Nurs* 2021; **35**: 375–94.
- 204 Wan Mohd Yunus WMA, Kauhanen L, Sourander A, et al. Registered psychiatric service use, self-harm and suicides of children and young people aged 0–24 before and during the COVID-19 pandemic: a systematic review. *Child Adolesc Psychiatry Ment Health* 2022; **16**: 15.
- 205 Thome J, Deloyer J, Coogan AN, et al. The impact of the early phase of the COVID-19 pandemic on mental-health services in Europe. *World J Biol Psychiatry* 2021; **22**: 516–25.
- 206 The Lancet. The widened gap in mental health services during the pandemic. *Lancet Reg Health West Pac* 2021; **15**: 100320.
- 207 Appleton R, Williams J, Vera San Juan N, et al. Implementation, adoption, and perceptions of telemental health during the COVID-19 pandemic: systematic review. *J Med Internet Res* 2021; **23**: e31746.
- 208 Rains LS, Dalton-Locke C, Landau S, Needle JJ, Johnson S. Variations in the uptake of telemental health technologies in community and crisis mental health services during the early pandemic: a survey of mental health professionals in the UK. *BMC Psychiatry* 2022; **22**: 776.
- 209 Lodge AC, Earley J, Peterson HL, Singh P, Manser SS. Evolution of the peer specialist role during COVID-19: challenges and opportunities for innovation beyond the COVID-19 era. *Psychiatr Rehabil J* 2023; published online March 6. <https://doi.org/10.1037/prj0000561>.
- 210 Schlieff M, Saunders KKK, Appleton R, et al. Synthesis of the evidence on what works for whom in telemental health: rapid realist review. *Interact J Med Res* 2022; **11**: e38239.
- 211 Pitchforth J, Fahy K, Ford T, Wolpert M, Viner RM, Hargreaves DS. Mental health and well-being trends among children and young people in the UK, 1995–2014: analysis of repeated cross-sectional national health surveys. *Psychol Med* 2019; **49**: 1275–85.
- 212 Thornicroft G, Chatterji S, Evans-Lacko S, et al. Undertreatment of people with major depressive disorder in 21 countries. *Br J Psychiatry* 2017; **210**: 119–24.
- 213 WHO Regional Office for Europe. The European Mental Health Action Plan 2013–2020. World Health Organization, 2015. Regional Office for Europe. <https://apps.who.int/iris/handle/10665/175672> (accessed Dec 3, 2022).
- 214 Shah P, Hardy J, Birken M, et al. What has changed in the experiences of people with mental health problems during the COVID-19 pandemic: a coproduced, qualitative interview study. *Soc Psychiatry Psychiatr Epidemiol* 2022; **57**: 1291–303.
- 215 Simblett SK, Wilson E, Morris D, et al. Keeping well in a COVID-19 crisis: a qualitative study formulating the perspectives of mental health service users and carers. *J Ment Health* 2021; **30**: 138–47.
- 216 Kauhanen L, Wan Mohd Yunus WMA, Lempinen L, et al. A systematic review of the mental health changes of children and young people before and during the COVID-19 pandemic. *Eur Child Adolesc Psychiatry* 2022; **12**: 1–19.
- 217 Lee J. Mental health effects of school closures during COVID-19. *Lancet Child Adolesc Health* 2020; **4**: 421.
- 218 Golberstein E, Wen H, Miller BF. Coronavirus disease 2019 (COVID-19) and mental health for children and adolescents. *JAMA Pediatr* 2020; **174**: 819–20.
- 219 Gilsbach S, Herpertz-Dahlmann B, Konrad K. Psychological impact of the COVID-19 pandemic on children and adolescents with and without mental disorders. *Front Public Health* 2021; **9**: 679041.

© 2023 Elsevier Ltd. All rights reserved.