



Full length article

## A vision for safer food contact materials: Public health concerns as drivers for improved testing

Jane Muncke<sup>a,\*</sup>, Anna-Maria Andersson<sup>b</sup>, Thomas Backhaus<sup>c</sup>, Scott M. Belcher<sup>d</sup>, Justin M. Boucher<sup>a</sup>, Bethanie Carney Almroth<sup>c</sup>, Terrence J. Collins<sup>e</sup>, Birgit Geueke<sup>a</sup>, Ksenia J. Groh<sup>f</sup>, Jerrold J. Heindel<sup>g</sup>, Frank A. von Hippel<sup>h</sup>, Juliette Legler<sup>i</sup>, Maricel V. Maffini<sup>j</sup>, Olwenn V. Martin<sup>k</sup>, John Peterson Myers<sup>e,l</sup>, Angel Nadal<sup>m</sup>, Cristina Nerin<sup>n</sup>, Ana M. Soto<sup>o,p</sup>, Leonardo Trasande<sup>q</sup>, Laura N. Vandenberg<sup>r</sup>, Martin Wagner<sup>s</sup>, Lisa Zimmermann<sup>a</sup>, R. Thomas Zoeller<sup>r</sup>, Martin Scheringer<sup>t,u,v</sup>

<sup>a</sup> Food Packaging Forum Foundation, Zurich, Switzerland<sup>b</sup> Dept. of Growth and Reproduction, Rigshospitalet and Centre for Research and Research Training in Male Reproduction and Child Health (EDMaRC), Copenhagen University Hospital – Rigshospitalet, Copenhagen, Denmark<sup>c</sup> Dept of Biological and Environmental Sciences, University of Gothenburg, Sweden<sup>d</sup> Dept. of Biological Sciences, North Carolina State University, Raleigh, NC, USA<sup>e</sup> Dept. of Chemistry, Carnegie Mellon University, Pittsburgh, PA, USA<sup>f</sup> Department of Environmental Toxicology, Eawag, Swiss Federal Institute of Aquatic Science and Technology, Dübendorf, Switzerland<sup>g</sup> Healthy Environment and Endocrine Disruptor Strategies, Durham, NC, USA<sup>h</sup> Mel & Enid Zuckerman College of Public Health, University of Arizona, Tucson, AZ, USA<sup>i</sup> Dept. of Population Health Sciences, Faculty of Veterinary Medicine, University of Utrecht, Netherlands<sup>j</sup> Independent Consultant, Frederick, MD, USA<sup>k</sup> Plastic Waste Innovation Hub, Department of Arts and Science, University College London, UK<sup>l</sup> Environmental Health Sciences, Charlottesville, VA, USA<sup>m</sup> IDiBE and CIBERDEM, Miguel Hernández University of Elche, Alicante, Spain<sup>n</sup> Dept. of Analytical Chemistry, I3A, University of Zaragoza, Zaragoza, Spain<sup>o</sup> Department of Immunology, Tufts University School of Medicine, Boston, MA, USA<sup>p</sup> Centre Cavailles, Ecole Normale Supérieure, Paris, France<sup>q</sup> College of Global Public Health and Grossman School of Medicine and Wagner School of Public Service, New York University, New York, NY, USA<sup>r</sup> Department of Environmental Health Sciences, School of Public Health & Health Sciences, University of Massachusetts Amherst, Amherst, MA, USA<sup>s</sup> Dept. of Biology, Faculty of Natural Sciences, Norwegian University of Science and Technology, Trondheim, Norway<sup>t</sup> RECETOX, Masaryk University, Brno, Czech Republic<sup>u</sup> Department of Environmental Systems Science, ETH Zurich, Switzerland

## ARTICLE INFO

Handling Editor: Adrian Covaci

## Keywords:

Food packaging  
Hazard assessment  
Chronic disease  
Chemical safety

## ABSTRACT

Food contact materials (FCMs) and food contact articles are ubiquitous in today's globalized food system. Chemicals migrate from FCMs into foodstuffs, so called food contact chemicals (FCCs), but current regulatory requirements do not sufficiently protect public health from hazardous FCCs because only individual substances used to make FCMs are tested and mostly only for genotoxicity while endocrine disruption and other hazard properties are disregarded. Indeed, FCMs are a known source of a wide range of hazardous chemicals, and they likely contribute to highly prevalent non-communicable diseases. FCMs can also include non-intentionally added substances (NIAS), which often are unknown and therefore not subject to risk assessment. To address these important shortcomings, we outline how the safety of FCMs may be improved by (1) testing the overall migrate, including (unknown) NIAS, of finished food contact articles, and (2) expanding toxicological testing beyond genotoxicity to multiple endpoints associated with non-communicable diseases relevant to human health. To

**Abbreviations:** AOP, Adverse Outcome Pathway; BPA, bisphenol A; CVD, Cardiovascular Disease; FCC, Food Contact Chemical; NCD, Non-Communicable Disease; NIAS, Non-Intentionally Added Substance; PFAS, Per- and polyfluoroalkyl Substances; PFOA, perfluorooctanoic acid; SCOD, Six Clusters of Disease.

\* Corresponding authors at: Food Packaging Forum Foundation, Zurich, Switzerland (J. Muncke). RECETOX, Masaryk University, Brno, Czech Republic (M. Scheringer).

E-mail addresses: [jane.muncke@fp-forum.org](mailto:jane.muncke@fp-forum.org) (J. Muncke), [scheringer@usys.ethz.ch](mailto:scheringer@usys.ethz.ch) (M. Scheringer).

<https://doi.org/10.1016/j.envint.2023.108161>

Received 14 April 2023; Received in revised form 17 August 2023; Accepted 17 August 2023

Available online 26 September 2023

0160-4120/© 2023 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

identify mechanistic endpoints for testing, we group chronic health outcomes associated with chemical exposure into Six Clusters of Disease (SCOD) and we propose that finished food contact articles should be tested for their impacts on these SCOD. Research should focus on developing robust, relevant, and sensitive *in-vitro* assays based on mechanistic information linked to the SCOD, e.g., through Adverse Outcome Pathways (AOPs) or Key Characteristics of Toxicants. Implementing this vision will improve prevention of chronic diseases that are associated with hazardous chemical exposures, including from FCMs.

## 1. Introduction

In today's globalized food system, food contact materials (FCMs) and food contact articles (Fig. 1) such as food packaging, tableware, and food processing equipment are ubiquitous, especially those made of plastic (Chakori et al., 2021; Poças et al., 2009). This increases exposures to food contact chemicals (FCCs) migrating from FCMs (Biryol et al., 2017; Koch and Calafat, 2009; Qian et al., 2018). This widespread, continuous exposure to a wide range of synthetic chemicals requires a more stringent safety assessment of FCMs than the current approaches used in low-, middle- and high-income countries (Neltner et al., 2013a; Maffini et al., 2013; Alger et al., 2013; Grob et al., 2006; Muncke et al., 2017).

FCMs have been studied for over 50 years and are a known source of chemicals that migrate into foodstuffs (Castle et al., 1989; Bradley et al., 2008; Dionisi and Oldring, 2002; Jickells et al., 1993; Sanchis et al., 2017; Nerin and Asensio, 2007; Geueke et al., 2022; Tsochatzis et al., 2021; Oldring et al., 2014). Numerous FCCs, either intentionally used in the manufacture of FCMs or non-intentionally added substances (NIAS) that are present in the finished food contact article and that migrate into foodstuffs (Nerin et al., 2013; Qian et al., 2018; Tisler and Christensen, 2022), are known to be hazardous and implicated with adverse human health impacts (Zimmermann et al., 2022; Groh et al., 2021; Van Bosuyt et al., 2019, 2016; Souton et al., 2017; Bengtstrom et al., 2016; Symeonides et al., 2021).

However, the current approach to chemical risk assessment for FCMs is largely focused on assessing the genotoxicity of single substances used to manufacture FCMs and therefore fails to account for other highly relevant mechanisms of toxicity that are of equal concern as genotoxicity (Muncke et al., 2017) and, what is more, the current approach does not assess NIAS that also migrate from FCMs (Fig. 2) (Muncke et al., 2020; Geueke et al., 2022). Addressing both issues, the limited scope of toxicity testing as well as the lack of testing for all migrating FCCs, is necessary to protect public health, and it can be done in a cost-efficient way.


Indeed, non-cancer non-communicable diseases (NCDs) of increasing prevalence in the global human population have been associated with several widely used FCCs, such as bisphenol A (BPA), bisphenol F (BPF), perchlorate, and di(2-ethyl hexyl) phthalate (DEHP), to name a few. Given that humans are in daily contact with FCMs, those materials are likely a relevant exposure source of hazardous chemicals that contribute to various NCDs.

In this article, we outline an improved assessment scheme for hazard identification of FCCs that captures all exposure-relevant chemicals (known as the *overall migrate*, i.e., all chemicals migrating as a mixture from finished food contact articles into foodstuffs) including (unknown) NIAS, and we present a vision for assessing the safety of FCMs that addresses biological effects linked to the most prevalent NCDs (Muncke, 2021; Jeddi et al., 2021). These include heart disease, stroke, cancer, diabetes, reproductive disorders, immunological disorders, and several neurological conditions. We provide guidance on research and policy actions that should be developed to protect the public from avoidable chronic chemical exposures originating from FCMs and finished food contact articles.

## 2. Problem set-up: Shortcomings of the current approach

### 2.1. Non-communicable diseases are increasingly prevalent and associated with chemical exposures

NCDs are a significant contributor to global mortality (WHO, 2018). However, the impact of NCDs is far greater than mortality alone, especially in low- and middle-income countries where health care is often limited compared to high-income countries. Both mortality and morbidity of selected NCDs have increased substantially over the last 30 years. Premature deaths (<70 years) are primarily associated with cardiovascular disease (17.7 million deaths per year, accounting for 45 % of all NCD deaths), cancer (8.8 million deaths per year, 22 % of all NCD deaths), chronic respiratory disease (3.9 million deaths per year, 10 % of



**Fig. 1.** Illustration of the terms *food contact article* (FCA), *food contact material* (FCM), and *food contact chemical* (FCC). The terms FCA and FCM are often used interchangeably, but only FCAs can be considered “finished” while FCMs oftentimes will be used in combination with other FCMs to make a finished FCA. The term FCC describes any chemical that is present in an FCM or FCA, regardless of whether it was intentionally used, or if it is a non-intentionally added substance (NIAS), such as an impurity, a reaction by-product, a degradation product, or of other origin (e.g., a contaminant from recycling).

all NCD deaths) and diabetes (1.6 million deaths per year, 4 % of all NCD deaths) (WHO, 2018). Expressed in Disability-Adjusted Life Years, cardiovascular diseases have increased by a factor of 1.4 from 1990 to 2017, neoplasms by a factor of 1.5, and diabetes, urogenital, blood and endocrine diseases by a factor of 1.6 (from 1990 to 2016) (Roser et al., 2021) (Fig. S1). Furthermore, among reproductive-age women and men, infertility is now the most prevalent chronic disease (WHO, 2020). Importantly, NCDs incur significant human suffering in addition to their estimated economic costs (Kassotis et al., 2020; Attina et al., 2016; Trasande et al., 2015; Trasande et al., 2016; Obsekov et al., 2022), which further stresses the need for urgent action towards prevention of morbidities associated with NCDs.

Chemical exposures are an important contributor to NCDs, especially when they occur during sensitive stages such as early life, or persist over extended periods of time. Several well-studied types of chemicals such as toxic metals, halogenated aromatics, and some pesticides (Bergman et al., 2013; Rojas-Rueda et al., 2021), as well as some members of the endocrine disrupting compounds (Gore et al., 2015; Goralczyk, 2021; Demeneix and Slama, 2019; Tanner et al., 2020; Chamorro-Garcia et al., 2017) are associated with NCDs such as brain-related disorders, cancers, metabolic disorders, reproductive disorders and cardiovascular disease. Specific FCCs such as BPA and several members of the *ortho*-phthalates group are associated with NCDs such as heart disease, diabetes, and some forms of cancer (Martínez-Ibarra et al., 2021; Svensson et al., 2021) (Table 1). Further, the effects of chemical exposures on risk of NCDs are complex and multifaceted, with some outcomes occurring across generations through transgenerational inheritance (Walker et al., 2018; Feil and Fraga, 2012; Chamorro-Garcia et al., 2017). It is also clear that these effects are not limited to laboratory animals, as mixtures of chemicals including FCCs have been associated with adverse health outcomes in prenatally exposed humans (Svensson et al., 2021; Tanner et al., 2020; Kortenkamp and Faust, 2018; Bornehag et al., 2021; Bornehag et al., 2019; Caporale et al., 2022).


NCDs that are increasingly prevalent in the human population and that are associated with hazardous chemical exposures can be grouped into disease clusters. On this basis, we developed the novel concept of Six Clusters of Disease (SCOD) (Fig. 3). The six clusters are cancers, cardiovascular diseases, reproductive disorders, brain-related disorders, immunological disorders, and metabolic diseases. The SCOD concept

provides a framework for systematically assessing the safety of chemicals in FCMs, with a focus on the prevention of chemical-associated, highly prevalent, and severe NCDs. As such, the SCOD concept expands current efforts for chemical risk assessment of FCCs.

## 2.2. Current risk assessment of food contact chemicals is not sufficiently protective of human health

The universe of known FCCs comprises at least 14,153 substances, and for at least 1,822 FCCs empirical evidence for migration from food contact articles and materials is publicly available (Geueke et al., 2022). Evidence of human exposure exists for hundreds of these chemicals (Barr et al., 2003; Calafat et al., 2005; Silva et al., 2004; Caporale et al., 2022; Correia-Sá et al., 2017; Cortéjade et al., 2017; Koch and Calafat, 2009; Pouech et al., 2015; Rudel et al., 2011; Susmann et al., 2019; Isaacs et al., 2022; Domínguez-Romero et al., 2022; Bil et al., 2023; Jung et al., 2022; Ruan et al., 2019). At least 388 FCCs in use today are known to be carcinogenic, mutagenic or toxic to reproduction, possess endocrine disrupting properties, or have other properties of concern such as persistence, and for at least 127 of these FCCs of concern there is evidence for migration from FCMs into food or food simulant (Zimmermann et al., 2022).

Currently, in the United States (US), Canada, the European Union (EU), China and other countries, chemical risk assessment is required for all migrating substances (Fig. 2). In practice, however, it is predominantly the intentionally used substances that are assessed for their risk to human health (Muncke et al., 2017). Humans are exposed to many more FCCs that are non-intentionally added to the finished food contact material or foodstuff. These NIAS include impurities of the starting substances, reaction by-products, or degradation products of starting substances (like additives) (Horodytska et al., 2020; Qian et al., 2018; Bradley and Coulier, 2007; Bauer et al., 2019), and also contaminants that accumulate in reused or recycled FCMs (Geueke et al., 2023; Geueke et al., 2018; Biedermann et al., 2013; Oldring et al., 2023). NIAS most often are unidentified, they are common in FCMs with high chemical complexity, and they are likely to be biologically active (Geueke, 2018). Under the current chemicals risk assessment paradigm for FCMs, where a chemical's identity must be known, unidentified FCCs cannot be assessed, although, for example, the EU plastic food contact



**Fig. 2.** Chemical risk assessment for food contact chemicals (FCCs): current practice. The current approach for assessing the safety of FCCs focuses on testing single substances that are intentionally used to make FCMs. Only genotoxic carcinogenicity is currently determined as a human-health relevant endpoint. However, many more chemicals can migrate simultaneously from the finished FCM, including unidentified compounds that are non-intentionally added substances (NIAS). The migrating mixture is known as the overall migrate, and it can exert adverse effects (mixture toxicity). Currently, the assessment of the overall migrate's mixture toxicity is not legally required.

**Table 1**

Food contact chemicals (FCCs) associated with non-communicable diseases (NCDs) from each of the Six Clusters of Disease (SCOD) (non-exhaustive and non-systematic overview of epidemiological studies). Identification of FCCs was based on the Food Contact Chemicals database (FCCdb) (Groh et al., 2021) and the database on migrating and extractable food contact chemicals (FCCmigex) (Geueke et al., 2022). This overview is not a complete list of FCCs that are associated with adverse health outcomes. Cancer agents are classified by cancer site (IARC, 2022).

Disease Cluster	Example disease	Associated FCC exposure	References
Cancers	Testicular cancer	PFOA	IARC (2016); ATSDR (2021)
	Kidney cancer	PFOA	IARC (2016); Melnick (2001)
	Breast cancer	PFOA Ortho-phthalates	Wan et al. (2022) Wan et al. (2022)
Cardiovascular diseases	Cardiovascular diseases: including myocardial infarction, arrhythmias, dilated cardiomyopathy, atherosclerosis, and hypertension	BPA	Moon et al. (2021); Zhang et al. (2020b); Wehbe et al. (2020); Ramadan et al. (2020) Fu et al. (2020)
		Ortho-phthalates	
Brain-related disorders	Hypothyroid	BPA	Rebolledo-Solleiro et al. (2021)
		Ortho-phthalates	Radke et al. (2020)
		Perchlorate	Radke et al. (2020)
	Abnormal neurodevelopment	PFAS	Piekarski et al. (2020)
		Ortho-phthalates: DEHP, DBP, BBP and DEP	Eales et al. (2022)
	Attention Deficit Hyperactivity Disorder/behavior	Lead, BPA, ortho-phthalates	Moore et al. (2022); Li et al. (2020); Park et al. (2015)
	Lower Intelligence Quotient	Endocrine disrupting chemical (EDC) mixture (Ortho-phthalates)	Tanner et al. (2020); van den Dries et al. (2020)
Language delay	EDC mixture	Caporale et al. (2022)	
Metabolic and endocrine diseases	Type-1 diabetes	BPA, Ortho-phthalates, PFAS	Predieri et al. (2020)
	Type-2 diabetes	BPA	Wang et al. (2019); Rancière et al. (2015); Akash et al. (2020)
	Pre-diabetes and diabetes	PFOA	He et al. (2018)
		Ortho-phthalates	Eales et al. (2022); Radke et al. (2018); Dales et al. (2018)
	Obesity (BMI, waist circumference)	BPA	Fu et al. (2020); Pérez-Bermejo et al. (2021); Wu et al. (2020)
PFAS		Liu et al. (2018); Geiger et al. (2021)	

**Table 1 (continued)**

Disease Cluster	Example disease	Associated FCC exposure	References
	Childhood Obesity	BPA	Ribeiro et al. (2020)
		Ortho-phthalates	Buckley et al. (2016)
	Gestational diabetes	Antimony	Zhang et al. (2020a)
		Ortho-phthalates	Shaffer et al. (2019)
Non-alcoholic fatty liver disease		EDC mixture	Midya et al. (2022)
		PFAS	Stratakis et al. (2020)
Immunological disorders	Immunosuppression	PFAS: PFOS and PFOA	DeWitt et al. (2019)
	Childhood asthma	Ortho-phthalates: DEHP and BBzP	Eales et al. (2022)
	Kidney damage	Melamine	Hsieh et al. (2012)
Reproductive disorders	Male infertility	BPA	Sharma et al. (2020)
		Dibutyl phthalate	Estill et al. (2019)
	Semen quality	Ortho-phthalates: DBP, BBP, DEHP, and DINP	Radke et al. (2018); Eales et al. (2022); Thurston et al. (2016)
Female infertility (reduced follicular count)	DEHP	Messerlian et al. (2016)	

regulation requires the risk assessment of NIAS (EU, 2011), and also US FDA’s Food Contact Notification has information requirements on impurities and reaction by-products (FDA, 2007).


A second problem is the lack of testing of substances present in the finished food contact material. Several approaches have been developed to approximate the health risks of unknown NIAS (Koster et al., 2015; Koster et al., 2013; Piek et al., 2017; Taylor and Sapozhnikova, 2022; Leeman and Krul, 2015; Omer et al., 2019; Sapozhnikova et al., 2021), but these approaches contain substantial uncertainties related to hazard estimation, chemical identification, and quantification (Bschor, accepted; Van Bossuyt et al., 2017) because they are based on assumptions that cannot be entirely supported by empirical evidence. For example, generic thresholds for chronic exposures to nongenotoxic carcinogens were derived from testing chemicals at maximum tolerable doses (MTD) and at 1/2 MTD, but it depends on the exact mechanism by which a chemical exerts its toxicity whether a low-dose extrapolation from MTD dosing is appropriate or not (Bailey et al., 2009; Williams et al., 2009).

Finally, because some laws prohibit the use of chemicals that cause cancer in humans or animals, testing methods currently focus on genotoxicity as a proxy for predicting cancer risk (Muncke et al., 2017; Neltner et al., 2013b). But other hazards that are not related to genotoxic effects are currently not systematically assessed, including outcomes relevant to other chronic NCDs. Thus, there is a need for novel and more comprehensive approaches to more fully evaluate all the relevant hazards to human health associated with FCCs.

### 3. Our vision: To make safer food contact materials

#### 3.1. Assessing toxicological effects relevant to the Six Clusters Of Disease

Chronic exposure to hazardous chemicals is a known modifiable risk factor for cancer and reducing exposure to hazardous or untested



**Fig. 3.** The Six Clusters of Disease (SCOD) concept comprises non-communicable diseases (NCDs) that are highly prevalent in the global human population, of increasing concern and associated with hazardous chemical exposures that can be clustered by disease type. The SCOD are of major concern for public health and require novel approaches for prevention, namely the identification of chemical contributors.

chemicals from consumer products, including FCMs, is a recommended preventive measure (Madia et al., 2019). It is reasonable to assume that the same holds true for other NCDs that are associated with chemical exposures, especially for endocrine disrupting chemicals (Table 1). Indeed, exposure reductions can lower the incidence of disease (Scholz et al., 2022), for example for neurodevelopmental disorders (Bennett et al., 2016), obesity (Mohanto et al., 2021) or male reproductive disorders (Foresta et al., 2018).

NCDs that are increasingly prevalent in the human population and that are associated with hazardous chemical exposures can be grouped into disease clusters. On this basis, we have developed the novel concept of SCOD (Fig. 3). The SCOD concept emerged from discussions with the Food Packaging Forum's Scientific Advisory Board (SAB) during several meetings between 2016 and 2022. The SCOD concept provides for the first time a framework for systematically assessing the safety of chemicals in FCMs, with a focus on the prevention of chemical-associated, highly prevalent and severe NCDs. As such, the SCOD concept expands current efforts for chemical risk assessment of FCCs beyond cancers induced via a genotoxic mechanism (Beneventi et al., 2020; Muncke et al., 2017). For each disease cluster within the SCOD, many widely used FCCs have been associated with relevant diseases in both epidemiology and animal studies (Tables 1 and 2). For some, mechanistic evidence strengthens these associations (Table 2). It is also this mechanistic evidence that provides opportunities to use *in-silico* and *in-vitro* assays to better map toxicity profiles of individual FCCs in finished FCMs, before they are placed on the market, as well as mixtures, extracts and migrates from FCMs and food contact articles. The SCOD provides organizing principles for such an approach.

### 3.2. Assessing real-life chemical exposures: Testing overall migrate from food contact materials

All FCCs that are relevant for human exposure should be tested, in other words, FCCs used in the manufacturing of FCMs should be tested as single substances, and the real-life mixture of all migrating FCCs, the overall migrate, should also be tested. If the overall migrate displays positive findings in the *in-vitro* assays, it should be subjected to non-targeted chemical analyses in order to elucidate its chemical composition, including NIAS, and to identify the substances driving the overall migrate's toxicity (Nerín et al., 2022). This combined testing and chemical identification approach could inform the development of safer FCMs by selecting less hazardous ingredients and developing manufacturing processes that generate fewer and less biologically active NIAS. Such an approach would be aligned with the proposed Safe and Sustainable by Design framework included in the EU's Chemicals Strategy for Sustainability (EU, 2020a).

The already available as well as emerging *in-vitro* assays provide an opportunity to identify hazardous properties of single substances and of the overall migrate. *In-vitro* test systems are small-scale, often single-cell or small organism systems, for example human cancer cell lines, bacteria, and fungi (e.g. yeast). Other high-throughput screening assays utilize embryos and larvae from vertebrates such as zebrafish (*Danio*

*rerio*) or African clawed frog (*Xenopus laevis*). These assays can be performed efficiently both in terms of time and cost and are usually based on mechanistic pathways (Groh and Muncke, 2017; Severin et al., 2017; Akoueson et al., 2023).

Test batteries, where several relevant assays are combined simultaneously, can also be operated as high-throughput screening methods such as those developed in Tox21 and ToxCast (Richard et al., 2016; Tice et al., 2013; Filer et al., 2022), which demonstrate the feasibility of this approach. In this way, diverse information about the interaction properties of a single chemical with different biological systems can be generated efficiently, and with lower cost, compared to whole-animal testing used in traditional toxicology. Further, overall migrate, that is, mixtures of chemicals migrating from FCMs, can be tested in such assays, too (Akoueson et al., 2023; Zimmermann et al., 2019; Bengtstrom et al., 2016).

These assessments should be guided by the SCOD concept. However, gaps exist in the current understanding of molecular pathways related to the SCOD, and these *in-vitro* assays remain insufficient to identify the full panoply of potential hazards, especially those mediated by endocrine mechanisms. *In-vitro* assays included in high-throughput test batteries need to be appropriate for predicting relevant human health outcomes; should be demonstrated to be reproducible, sufficiently specific and sensitive; and must be executed transparently (Janesick et al., 2016; Schug et al., 2013). Because of the limited *in-vitro* assays for known pathways and mechanisms of action associated with endocrine disruption and other complex biological cascades, animal testing needs to continue, but actual *in-vivo* tests will be fewer than in the past. For example, no current *in-vitro* approaches would have revealed what is now known to be a feature of some chemical exposures, e.g., trans-generational epigenetic inheritance (Fitz-James and Cavalli, 2022). Acknowledging these and other gaps, the European Commission is funding EURION, a program to develop new testing and screening methods (including many *in-vitro* approaches) for identifying endocrine disrupting chemicals (Street et al., 2021).

### 3.3. Shifting from the status quo to a more comprehensive approach to testing

Within the SCOD, increasingly available mechanistic information enables an understanding of how chemicals contribute to highly prevalent NCDs. Two emerging frameworks are being implemented to describe how chemicals affect complex diseases and to provide a more uniform approach to evaluating mechanistic evidence: the key characteristics concept, and adverse outcome pathways (AOPs). Both offer opportunities to shift from the status quo, modernize hazard assessments, and develop suitable *in-vitro* assays.

#### 3.3.1. The key characteristics concept: Modernizing chemical hazard assessments

The key characteristics concept makes use of information about the properties of hazardous chemicals that have empirical evidence linking them causally to relevant apical (disease) endpoints (Smith et al., 2016).

**Table 2**

Examples of food contact chemicals (FCCs) that are associated with diseases from the Six Clusters of Disease (SCOD) by mechanisms from *in vitro* and/or *in vivo* studies (not including epidemiological studies). Not a complete list: select references only.

Disease Cluster	Food Contact Chemical	Reference	
Cancers	<b>Melamine</b> (CAS 108-78-1)	IARC (2019)	
	<b>Formaldehyde</b> (CAS 50-00-0)	IARC (2012a)	
	<b>Benzidine</b> (CAS 92-87-5)	IARC (2010)	
	<b>4,4'-Diamino-3,3'-Dichlorodiphenylmethane (MOCA)</b> (CAS 101-14-4)	IARC (2012a)	
	<b>Antimony trioxide</b> (CAS 1309-64-4)	NTP (2018)	
	<b>Perfluorooctanoic acid (PFOA)</b> (CAS 335-67-1)	Temkin et al. (2020); Pierozan et al. (2018); Charazac et al. (2022)	
	<b>Di (2-ethylhexyl) phthalate (DEHP)</b> (CAS 117-81-7)	Hager et al. (2022); IARC (2012b)	
	<b>Bisphenol A (BPA)</b> (CAS 80-05-7)	Sang et al. (2021); Jun et al. (2021); Dhimolea et al. (2014); Prins et al. (2014)	
	Cardiovascular diseases	<b>Bisphenol A (BPA)</b> (CAS 80-05-7)	Pant et al. (2011); Gao and Wang (2014); Kofron et al. (2021); Hyun et al. (2021); Krishna et al. (2021); Cooper and Posnack (2022)
		<b>Triclosan</b> (CAS 3380-34-5)	Krishna et al. (2021)
<b>Tributyltin chloride</b> (CAS 1461-22-9)		Krishna et al. (2021)	
<b>Diethanolamine</b> (CAS 111-42-2)		Jokinen et al. (2005)	
<b>DEHP</b>		Mariana et al. (2016)	
Brain-related disorders	<b>Perchlorate</b> (CAS 14797-73-0)	Kirk (2006)	
	<b>Ortho-phthalates</b>	Hlisenfková et al. (2021)	
	<b>BPA</b>	McDonough et al. (2021); Wang et al. (2020)	
Metabolic diseases	<b>Bisphenol S (BPS)</b> (CAS 80-09-1)	Naderi and Kwong (2020); McDonough et al. (2021)	
	<b>BPA</b>	Villar-Pazos et al. (2017); Martinez-Pinna et al. (2019); Wassenaar et al. (2017); Desai et al. (2018a); Manikkam et al. (2013)	
	<b>Bisphenol A diglycidyl ether (BADGE)</b> (CAS 1675-54-3)	Wang et al. (2021)	
	<b>Organotins</b>	Rotenberg et al. (1978)	
	<b>Perchlorate</b>	Larsson-Nyrén et al. (2001)	
	<b>Perfluorooctanesulfonic acid (PFOS)</b> (CAS 1763-23-1)	Qin et al. (2020); Sant et al. (2017)	
	<b>Bisphenol F (BPF)</b> (CAS 620-92-8)	Marroqui et al. (2021)	
	<b>BPS</b>	Marroqui et al. (2021)	
	<b>2,4,7,9-tetramethyl-5-decyne-4,7-diol (TMDD; Surfynol)</b> (CAS 126-86-3)	Garcia-Calvo et al. (2020b); Nerin et al. (2018); Nerin et al. (2014)	
	<b>DEHP</b>	Li et al. (2019); Manikkam et al. (2013)	
Immunological disorders	<b>Melamine</b>	IARC (2019)	
	<b>BPA</b>	McDonough et al. (2021)	
	<b>BPF</b>	McDonough et al. (2021)	
	<b>BPS</b>	McDonough et al. (2021); Nowak et al. (2019)	

**Table 2 (continued)**

Disease Cluster	Food Contact Chemical	Reference
Reproductive disorders	<b>2,4-di-tert-butylphenol</b> (CAS 96-76-4)	Liu et al. (2022)
	<b>DEHP</b>	Nowak et al. (2019); Hessel et al. (2015)
	<b>BPA</b>	Liu (2021); Vessa et al. (2022b); Wang et al. (2014); Mahalingam et al. (2017)
	<b>BADGE</b>	Wang et al. (2021); Nerin et al. (2014)
	<b>BPS</b>	Desmarchais et al. (2020)
	<b>DEHP</b>	Vessa et al. (2022b); Mariana et al. (2016)

The underlying premise is that chemicals that cause the same disease outcomes in whole organisms share molecular properties (i.e., key characteristics) that are relevant for their hazardous properties. The key characteristics for different disease outcomes are hence defined using empirical evidence for well-characterized chemicals, combined from epidemiological, *in vivo* and mechanistic studies. These disease-specific key characteristics can then be used to develop mechanistic *in vitro* assays to screen chemicals for their propensity to contribute to different disease clusters and thereby reduce the need for *in vivo* experiments while still decreasing scientific uncertainty normally associated with *in vitro* data.

The key characteristics were first developed for carcinogens, drawing from existing mechanistic information from thoroughly assessed chemicals that are known to be carcinogenic in humans (Smith et al., 2016; Guyton et al., 2018; Krewski et al., 2019; Al-Zoughool et al., 2019; Guyton and Schubauer-Berigan, 2021). Additional key characteristics of other disease-causing chemicals have also been described, such as for hepatotoxicants (Rusyn et al., 2021), endocrine disrupting chemicals (Merrill et al., 2020), female reproductive toxicants (Luderer et al., 2019), male reproductive toxicants (Arzuaga et al., 2019), cardiovascular toxicants (Lind et al., 2021), and immunotoxicants (Germolec et al., 2022). For metabolic toxicants and neurotoxicants, work to describe key characteristics is ongoing. Taken together, the key characteristics approach provides an excellent starting point for the mechanistic understanding of how certain chemicals are associated with NCDs, such as those covered in the SCOD.

### 3.3.2. Using other mechanistic information to develop suitable *in vitro* assays

In addition to the key characteristics, further important mechanistic understanding is becoming available and can be useful to inform the development of dedicated *in vitro* screening assays for hazard assessments of FCCs. Chemicals exert toxic effects by combinations of many different molecular-level events. These mechanistic events leading to apical endpoints of toxicity can be organized in an AOP (Ankley et al., 2010). Several AOPs relevant to NCDs in the SCOD have been proposed, such as estrogen receptor activation leading to breast cancer (Coumoul et al., 2022) and the upregulation of thyroid hormone catabolism (via activation of hepatic nuclear receptors) leading to subsequent adverse neurodevelopmental outcomes in mammals, specifically the loss of cochlear function (Friedman et al., 2022). Thus, AOPs are an emerging approach to organize mechanistic information so that molecular or cellular-level targets can be identified for developing *in vitro* assays that are relevant to the SCOD.

### 3.3.3. The novel approach: A vision for safer food contact materials

Based on the presumption that mechanistic *in vitro* testing of chemicals supports the prevention of NCDs within the SCOD, we propose a novel approach for testing FCCs that

- (1) covers individual FCCs as well as real-life mixtures, migrating (or extractable) from finished FCMs, including all known and unknown NIAS,
- (2) assesses the health impacts of FCCs and real-life mixtures with respect to the most prevalent NCDs in the human population, and
- (3) evaluates effects that are upstream from the disease, relying on mechanistic information and *in-vitro* screening approaches (wherever possible) to predict health effects induced by FCCs and migrates.

This shift from current practice to the proposed approach is summarized in Fig. 4, and a detailed overview is provided in Fig. 5. Our approach overcomes the most challenging shortcomings of the current testing paradigm of chemical hazard assessment of FCMs, fully recognizing that to assess all adverse effects of chemicals on biological systems, adequate *in-vivo* testing is required, where additional aspects would be addressed such as metabolic activation, unknown modes of action leading to apical endpoints, and transgenerational effects. However, we also realize that such extensive, multigeneration *in-vivo* testing may not always be feasible for various reasons, including ethical and practical ones. Therefore, we propose this vision to improve FCC testing from the currently too limited scope towards a much more comprehensive yet feasible approach that holds promise for better protection of public health.

#### 4. Implementing the vision: Assessing impacts of FCCs and relevant mixtures on human health outcomes in the SCOD using mechanistic approaches

Here we review the mechanistic basis for each of the disease clusters included in the SCOD, and selectively highlight available *in-vitro* testing

methods. Importantly, some available assays cover key characteristics that are relevant for several disease clusters.

This vision for expanded hazard assessment of FCMs is based on the finding that for each of the disease clusters included in the SCOD, some mechanistic understanding is available for the way that chemicals cause disease (Table 2).

##### 4.1. Cancer

As defined by Willis,

A neoplasm is an abnormal mass of tissue, the growth of which exceeds and is uncoordinated with that of the normal tissues and persists in the same excessive manner after cessation of the stimulus which evoked the change (Willis, 1948).

Regarding cancer causation, the somatic mutation theory posits that cancer is a cellular disease caused by mutations of genes that disrupt the control of cell proliferation. Yet, substantive contradictions exist between this theory and empirical evidence (Naxerova, 2021), which inspired competing theories to consider cancer as a problem of tissue organization akin to organogenesis (Sonnenschein and Soto, 2020; Maffini et al., 2004; Rønnov-Jessen and Bissell, 2009). Importantly, not all carcinogens are mutagens (Keri et al., 2007) and, thus, carcinogenicity cannot be equated with genotoxicity. Yet, because legal requirements restrict the use of cancer-causing agents in FCMs, testing of FCCs has focused on genotoxicity as a proxy to identify carcinogenic substances.

Both carcinogens and mutagens are found in FCMs including 1) formaldehyde, a known human carcinogen (IARC Group 1) (IARC, 2012a), which migrates from various plastics including melamine-formaldehyde plastics used as tableware for children, and





Fig. 4. Overview of the current vs. proposed approach to food contact chemical (FCC) testing. The proposed new approach focuses on testing the overall migrate (i. e., the human exposure-relevant mixture of all migrating FCCs) for its potential to contribute to the Six Clusters of Disease (SCOD).



**Fig. 5.** The vision for a novel approach to safety assessment of FCMs and food contact articles. Finished food contact articles are tested for their real-life mixture of all migrating chemicals (the *overall migrate*, i.e. the mixture of all migrating chemicals), using *in-vitro* screening assays. The screening assays are mechanism-based and identify the key characteristics, key initiating events, or other mechanisms of action of the overall migrate. Screening assays are selected around the Six Clusters of Disease (SCOD). For overall migrate displaying positive findings in the *in-vitro* assays, non-targeted chemical analyses are carried out to identify the substances driving the overall migrate's toxicity.

polyethylene terephthalate plastic (PET) (Kim et al., 2021; Bach et al., 2013); 2) antimony trioxide, which “is reasonably anticipated to be a human carcinogen” (NTP, 2021) and “probably carcinogenic to humans” (IARC Group 2A) (IARC, 2022), and is used in the manufacture of PET, where antimony is found to migrate into soft drinks (Westerhoff et al., 2008; Bach et al., 2013); and 3) per- and polyfluoroalkyl substances (PFAS), which are widely used in the manufacture of FCMs as processing aids in plastic and paper food contact material production (Trier et al., 2011; Minet et al., 2022), and perfluorooctanoic acid has limited evidence for testicular and kidney cancers in humans and is “possibly carcinogenic to humans” (IARC Group 2B) (Benbrahim-Tallaa et al., 2014).

The key characteristics for carcinogens reveal that these chemicals can be mutagens, but that there are numerous other common features for these agents as well, such as inducing oxidative stress, modulating receptor-mediated effects and inducing chronic inflammation (Smith et al., 2016; Guyton et al., 2018; Krewski et al., 2019; Al-Zoughool et al., 2019; Guyton and Schubauer-Berigan, 2021). Guyton and Schubauer-Berigan (2021) recommended the use of *in-vitro* assays based on the key characteristics to identify carcinogens in high-throughput screening (Guyton and Schubauer-Berigan, 2021). Further, Rider et al. (2021) proposed methods to use the key characteristics to test chemical mixtures and their propensity to affect cancer development including in mixtures of chemicals with different key characteristics of carcinogens (Rider et al., 2021). Approaches such as these will provide important information for testing mixtures such as the overall migrate from finished FCMs. Methods for evaluating genotoxicity are readily available, validated, and trusted. Chemicals are considered genotoxic if they damage the structure, information content, or segregation of DNA, with mutagenicity (i.e. changes to the nucleotide sequence) being a sub-type of genotoxicity (OECD, 2015).

These methods include (Muncke, 2009):

- **Mutagenicity:** The Ames test, based on bacterial reverse mutagenicity, is the most employed test for mutagenicity (Organisation for Economic Co-operation and Development (OECD) test guideline (TG) 471). A mammalian cell (mouse lymphoma) gene mutation test (OECD TG 490) is also available (OECD, 2022)
- **Chromosomal aberration:** Cultured mammalian cells are assessed for the presence of chromatid-type and chromosome-type aberrations during metaphase (OECD TG 473)
- **Micronucleus:** Micronuclei represent chromosomal damage (chromosome fragments or whole chromosomes) that have been transmitted to daughter cells. Micronuclei can be assessed *in-vitro* by using

mammalian cells (OECD TG 487) or *in-vivo* with erythrocytes collected from bone marrow or peripheral blood (OECD TG 874)

These methods are recommended or required for assessing intentionally used FCCs (EFSA, 2008; FDA, 2007). Several other *in-vitro* assays for assessing the genotoxic potential of FCCs are also available (Pinter et al., 2020). However, these strategies have not kept pace with discoveries in cancer biology (Chiara et al., 2020). Currently, no *in-vitro* assays are available that capture features of carcinogenicity beyond genotoxicity, but research is underway to address this technical gap (Hwang et al., 2020). On the other hand, the causal role of the micro-environment in carcinogenicity, as put forward by tissue-based theories on carcinogenicity (Maffini et al., 2004), is not captured by such *in-vitro* assays, because the reciprocal interactions between stroma and parenchyma during development, regeneration, and remodeling are not considered (Soto et al., 2013). Although *in-vivo* assays involving mammals are available, traditional 2-year rodent carcinogenicity studies (OECD TG 451), either alone or in combination with chronic toxicity studies, are rarely performed for FCCs.

#### 4.2. Cardiovascular diseases

Cardiovascular diseases (CVDs) are a group of disorders arising due to dysfunction of the heart and blood vessels. The most recognized forms of CVD, coronary heart disease and cerebrovascular disease, result in damage to tissues caused by limited or complete loss of blood supply (WHO, 2021).

FCCs including several phthalates and bisphenols contribute to the causation of CVDs, independent of obesity and diabetes (Lind et al., 2021). Bisphenols can disrupt calcium signalling in myocardium and vasculature; and phthalates and bisphenols are oxidant stressors that accelerate coronary and other arterial inflammation (Lind et al., 2021). In the US alone, 100,000 premature deaths from CVD among 55–64-year-olds each year are attributed to exposure to one phthalate, DEHP (Wen et al., 2022). Other FCCs, such as antimony, may also impair cardiovascular function and accelerate CVDs (El-Kersh et al., 2022).

Lind et al. (2021) compiled the key characteristics of cardiovascular toxicants and provided a comprehensive overview of robust and sensitive *in-vitro*, *ex-vivo* and *in-vivo* assays that are available for measuring dysregulation of  $Ca^{2+}$  ion homeostasis and resulting arrhythmogenic activities of chemicals. For example, the increased risk for CVDs associated with higher exposures to BPA is mechanistically associated with  $Ca^{2+}$  release and reuptake, resulting in proarrhythmic delays after depolarizations in isolated cardiomyocytes. BPA promotes  $Ca^{2+}$ -mediated



arrhythmias *ex vivo* in the whole heart of rats and mice (Yan et al., 2011). However, this is only one of many possible mechanisms for inducing CVDs, and further assay development is required.

Although several FCCs have been associated with CVDs, cardiovascular toxicity is generally not evaluated for FCCs. This is in part due to a reliance on *in-vivo* guideline testing of general toxicity for chemicals migrating at very high levels and limited to assessment of neoplastic and non-neoplastic cardiac lesions in rodent models, which can be confounded by a high incidence of background pathology in many of the rodent strains used for toxicity testing (Gear et al., 2017). However, these are insensitive apical endpoints that only identify highly cardiotoxic chemicals that result in robust pathology but miss subtle molecular effects (Gao and Wang, 2014; Jokinen et al., 2011).

We recommend that comprehensive testing for all new chemicals include *in-vitro* and *in-silico* testing harmonized with the Comprehensive *in-vitro* Proarrhythmia Assay approach (CiPA) (FDA, 2022; CIPA, 2019). The CiPA initiative was launched to address limitations in the current cardiac safety testing methods used to assess the risk for adverse cardiac events of new drugs. The CiPA aims to develop a new approach for evaluating the potential of drugs to cause cardiac arrhythmias, particularly a specific type known as Torsades de Pointes (TdP). It is a multi-step approach that combines *in-vitro* assays and computational modeling to assess proarrhythmic risk and predict the risk of TdP by considering the complex interactions of multiple ion channels and cellular components involved in cardiac electrophysiology. The CiPA could be used as an important approach for identifying cardiotoxic hazards of FCCs.

#### 4.3. Brain-based disorders

Disrupted neurodevelopment can have numerous consequences including a lower intelligence quotient, delayed language acquisition, attention deficit hyperactivity disorder (ADHD), and autism (Caporale et al., 2022; Bornehag et al., 2021; Kim et al., 2022). Because the role of thyroid hormone in brain development is well established, hypothyroidism, especially during early development, is also a condition of concern upstream of neurodevelopmental disorders. Neurotoxicity can also result from impaired neuronal function due to a variety of factors, such as neuronal misplacement during development, altered synapses, hypomyelin, or degeneration. Other neurodegenerative conditions that typically arise later in life include Parkinson's disease, Alzheimer's disease, and other forms of dementia.

The role of FCCs in the causation of many brain-based disorders is well established, with substantial contribution to the burden of disease for both neurodevelopmental and neurodegenerative disorders (Attina et al., 2016). For example, FCCs that interfere with thyroid hormone systems or sex steroids (e.g., phthalates and perchlorate) can affect brain development as well as cognitive function in adults (Bennett et al., 2016; Grandjean and Landrigan, 2006). The vulnerability of the developing brain and the lack of systematic assessment of neurodevelopmental toxicity for FCCs raises serious concerns (Maffini et al., 2016; Mustieles and Fernández, 2020). At present, the key characteristics of neurotoxicants remain undescribed, but relevant work is ongoing.

In addition to assays covering interference with the thyroid and sex steroid axes, *in-vitro* testing of neurotoxicants requires sophisticated and reliable models due to the complexity of the brain (Marty et al., 2021). Neuronal cell lines, primary central nervous system cells, transformed neuronal precursors and stem cell derived progenitor cells are used in neurotoxicity assays (Arshajyothirmayi and Gulia, 2022) to evaluate endpoints including migration, synapsis formation, network activity and differentiation. Although single-cell cultures are informative, multi-cell type and three-dimensional models utilizing microfluidics more adequately represent the diversity and spatial properties of the brain (Caffrey et al., 2021; Kilic et al., 2016; Maoz, 2021; Park et al., 2021), but high-throughput versions of these methods are not yet available, and thus their use in evaluating FCCs has been limited. Additional *in-vitro* assays for chemical screening of neurotoxicants are under development

in EU-funded research programs (Cediell-Ulloa et al., 2022) and research is ongoing to develop further *in-vitro* assays targeting the thyroid system (Kortenkamp et al., 2020). Recently, the establishment of a human cell-based *in-vitro* battery has been reported; it combines 10 assays selected to cover major key events in the relevant AOPs (Sachana et al., 2021) and was shown to provide 82 % sensitivity in that it was able to identify 24 out of 28 known neurotoxicants (Blum et al., 2022).

New low- and medium-throughput screening assays have been developed. For example, the nematode is a promising model for evaluating known neurodevelopmental toxicants, and this could be expanded to profiling chemicals with unknown neurotoxicity (Ruszkiewicz et al., 2018; Hunt et al., 2018). Spontaneous movements (Parg et al., 2007), number and location of neurons (Rericha et al., 2022), and behavioral effects (Fitzgerald et al., 2021) are some of the neurological endpoints measured in zebrafish. Validated high-throughput screening assays using African clawed frog tadpoles are also available (OECD TG 248).

*In-vivo* testing in rodents can be used to assess different functional aspects of neurotoxicity including impacts on cognition, learning and memory, and anxiety-like, depressive-like and reproductive behaviors. OECD developmental neurotoxicity (OECD TG 426) and extended one-generation reproductive toxicity assays (OECD TG 443) include optional measurements of learning and memory, motor and sensory function, motor activity, and auditory startle. Neurodegeneration is not covered because this endpoint can only be studied in animals which age, and animals used in assays are typically not kept until the end of their natural lifetime where neurodegeneration would manifest itself (Huff et al., 2008).

#### 4.4. Obesity and metabolic diseases

Metabolic diseases, including obesity, involve the many tissues that comprise the metabolic system (Mohajer et al., 2021). These include adipose tissue, skeletal muscle, pancreas, liver, gastrointestinal tract, bone, and brain. Type-2 diabetes, an important metabolic disease with increasing prevalence in human populations, occurs due to systemic insulin resistance, often with an increasing production of insulin by the pancreas. Type-1 diabetes occurs due to a progressive loss of  $\beta$ -cell insulin secretion. Non-alcoholic fatty liver disease is another metabolic disease with increasing prevalence in human populations.

While poor diet and insufficient physical activity are considered the chief drivers of the obesity and diabetes twin pandemics, chemical exposures (for example, to phthalates, bisphenols, parabens, PFAS, etc.) can disrupt the balance between energy expenditure and energy intake (Heindel et al., 2022). A large comprehensive review of metabolic disrupting chemicals, including those that can induce obesity (obesogens), provides strong evidence that numerous FCCs are associated with type-2 diabetes, obesity, and fatty liver disease (Heindel, 2019). The key characteristics of metabolic disruptors and obesogens are being compiled. Rusyn et al. (2021) have described the key characteristics of acute and chronic human hepatotoxicants and note that only one of 12 key characteristics are specific to liver tissue (KC9: causing cholestasis) (Rusyn et al., 2021), indicating that there are overlaps with the key characteristics of other toxicants, i.e. carcinogens, cardiovascular toxicants, endocrine disrupting chemicals, and male and female reproductive toxicants.

The simplest assays to identify an obesity hazard are those that measure the effect of chemical exposures on the development of adipocytes (Kassotis et al., 2022; Kassotis and Stapleton, 2019; Seo et al., 2019). Primary preadipocyte cultures, or mesenchymal stem cell assays, use animal or human cells to assess proliferation and differentiation into adipocytes (Desai et al., 2018b; Shoucri et al., 2018; Kassotis and Stapleton, 2019; Chamorro-Garcia and Blumberg, 2019; Lane et al., 2014; Tang et al., 2004; Pillai et al., 2014). Using this *in-vitro* assay, a recent study found that around one third of tested plastic food contact articles contained metabolic disrupting chemicals (Völker et al., 2022).

Recently, spheroid adipocyte models have been developed that

improve the efficiency and speed of differentiation (Turner et al., 2017) and can be used for a more comprehensive understanding of adipocyte physiology than monolayer cultures. The zebrafish obesogenic test offers an *in-vivo* approach to screening chemicals that target adiposity; it measures adipocyte lipid droplet size and normalized triacylglycerol content as an assessment of adiposity in a whole-organism assay of larvae to test for obesogenic and anti-obesogenic chemicals and mixtures (Tingaud-Sequeira et al., 2011).

Other non-adipocyte cell lines, when well characterized such as the mouse bone marrow-derived mesenchymal stem cells (mBMSCs), are also useful for mechanistic studies (Auerbach et al., 2016; Janesick et al., 2016). In addition to adipocyte differentiation, several other mechanisms are implicated with metabolic disease causations, for example the disruption of energy homeostasis at the level of the hypothalamus and brain. Therefore, *in-vitro* assays that examine effects on hypothalamic neurons are useful (Ye et al., 2016; Loganathan et al., 2018).

No assays have been developed to identify metabolic disruptors acting as diabetogens. Ongoing projects are developing assays to measure  $\beta$ -cell function and survival (Audouze et al., 2020; Legler et al., 2020; Küblbeck et al., 2020) using rodent  $\beta$ -cell lines (INS-1E and MIN-6) and a human  $\beta$ -cell line (ENDOC- $\beta$ H1). Assays of insulin function on the human liver cell line HepaRG, the skeletal muscle cell line C2C12, and adipocytes are also under investigation (Legler et al., 2020). One well established system of assays employing both *in-vitro* and *in-vivo* methods has been used to explore the relationship between BPA and type-2 diabetes (Dos Santos et al., 2022).

The most used assays to screen chemicals for effects on the liver use the HepaRG and HepG2 cell lines. The HepG2 cell line can be customized with different expression levels of various drug metabolizing enzymes (Tolosa et al., 2018). Other 2D and 3D *in-vitro* approaches use primary hepatocytes, immortalized liver cell lines, and hepatocytes derived from stem cells that are grown in monolayers, as spheroids or organoids, or used in emerging technologies (like organ-on-a-chip) to identify liver toxicants (Yang et al., 2023). Each of the approaches available has strengths and weaknesses; for example, the use of human primary hepatocytes in 2D culture can produce patient-specific evaluations that account for differences in metabolism and sensitivity, but such an approach comes at high cost. Several of these methods are currently being used to evaluate liver toxicity in the screening of pharmaceuticals (Serras et al., 2021), making them similarly well suited to evaluate FCCs for potential effects on liver outcomes.

#### 4.5. Immunological disorders

The immune system is an intricate network of many different, highly specialized cells interacting with each other and with the nervous and endocrine systems (Nicholson, 2016). Disorders of the immune system include autoimmune disorders such as multiple sclerosis, Graves' and Hashimoto's diseases, lupus, Celiac's, Addison's, and rheumatoid arthritis, among others. Other diseases including type-1 diabetes and asthma have an important immune component. Therefore, assays for immunotoxicity need to capture a multitude of potential effects, including immunosuppression, immunostimulation, hypersensitivity reactions, mechanisms of autoimmunity, and developmental immunotoxicity, e.g., delayed immunotoxic responses to toxic influences (Germolec et al., 2017).

The human immune system is highly effective, but also sensitive to synthetic chemical insults during development and adult life. Effects of chemicals on the immune system are less well understood in humans than other disease endpoints, but emerging evidence implicates PFAS exposure with reduced immune response to vaccines and increased susceptibility to infections in early life (Grandjean et al., 2017). Other FCCs including bisphenols and phthalates increase the risk of atopy and asthma (Xie et al., 2016; Wang et al., 2016; Kim et al., 2017), and infections in early life (Gascon et al., 2015).

The key characteristics of immunotoxicants have been described

(Germolec et al., 2022). This offers a starting point for the development of suitable *in-vitro* assays for testing FCCs for immunotoxicity. Due to the complexity of the immune system components and responses, a comprehensive battery of *in-vitro* assays covering all relevant aspects of immunotoxicity has not been established. However, several *in-vitro* assays, dealing for example with direct immunosuppression, allergic hypersensitivity, or autoimmunity, are being developed to detect a range of immunotoxicants (Corsini and Roggen, 2017; Luebke, 2012; Deproux et al., 2022; Naidenko et al., 2021) and these assays could be used to screen FCCs (Ogungbesan et al., 2019).

#### 4.6. Reproductive disorders

In industrialized countries, male reproductive health has declined over the past decades, including a 50–60 % decrease in sperm counts since 1973 (Levine et al., 2017; Skakkebaek et al., 2022; Levine et al., 2022) and an increase in testicular cancer (Znaor et al., 2022). Female fertility is also affected, as are maternal health and pregnancy outcomes, and conditions such as polycystic ovary syndrome (PCOS), endometriosis, and premature ovarian failure (Boomsma et al., 2006).

The sperm count decrease is associated with chemical exposures (to, e.g. phthalates), especially during fetal development (Skakkebaek et al., 2001). Strong evidence from animal experiments support this interpretation (Gore et al., 2015; Skakkebaek et al., 2016; Jorgensen et al., 2021; Kortenkamp, 2020). FCC exposures are also associated with PCOS (Soave et al., 2020), and other aspects of reproductive toxicity (Nerfin et al., 2020; Garcia-Calvo et al., 2020a). These adverse outcomes have also been found for FCCs promoted as safer alternatives to hazardous chemicals such as the plasticizer 1,2-cyclohexane dicarboxylic acid diisononyl ester (tradename Hexamoll DINCH) (Boisvert et al., 2016), which is used as a replacement for DEHP and other phthalates. Several FCCs such as BPA have been studied for mechanistic-level impacts on female fertility, including oogenesis, folliculogenesis, and altered expression of gonadotropin and gonadotropin hormone-releasing hormone receptors (Vessa et al., 2022a). The key characteristics of male (Arzuaga et al., 2019) and female reproductive toxicants (Luderer et al., 2019) have been described. Development and function of the reproductive system are fundamentally dependent on sex hormone action. Thus, the key characteristics of endocrine disrupting chemicals (Merrill et al., 2020) are also relevant to the study of chemicals that affect reproductive outcomes. However, a systematic overview of available *in-vitro* assays for hazard identification of endocrine disrupting chemicals that affect male and female fertility is currently not available.

*In-vitro* assays that identify chemical interference with sex hormone production and signalling have been validated (OECD TG 493, 455, 458, 456). These include assays based on nuclear receptor activation and steroid hormone synthesis. The bovine oocyte maturation assay (ECVAM TM 2010–05) is also a reproduction-relevant *in-vitro* assay. A good correlation between *in-vitro* results and *in-vivo* observations has been established for female fertility endpoints (Corton et al., 2022; Pinto et al., 2018). Validated *in-vivo* assays exist to evaluate reproductive toxicity for impacts on both male and female fertility (OECD TG 443), but these may not be sufficiently sensitive or comprehensive, as they fail to evaluate numerous key characteristics of male and female reproductive toxicants (Luderer et al., 2019; Arzuaga et al., 2019).

### 5. What is needed to implement the vision for safer food contact materials?

To achieve our vision, we propose a multi-pronged approach that is grounded in the SCOD concept, which includes many of the most prevalent NCDs of high relevance to human health. We identified three components needed to realize this vision: testing strategies and analytical methods, data interpretation and integration, and science to inform decision making.

### 5.1. Testing strategies and analytical methods

In Section 4 we list several available and emerging assays used in the identification of hazard for each of the SCOD. However, much more is needed, especially high-throughput non-animal and low-medium throughput assays with non-mammalian models that are specific, sensitive, reliable and robust, and adequate for predicting effects relevant to humans. These assays would overcome challenges with cost, time, and scientific relevance as the selection of suitable *in-vitro* assays would be based on robust mechanistic evidence from key characteristics and AOPs. Identification of the key characteristics for brain disorders and metabolic diseases will form the basis for identification and/or development of relevant *in-vitro* assays to identify hazardous chemicals related to these clusters. For *in-vitro* testing based on mechanistic pathways to succeed, additional dedicated expertise and financial support are needed to identify and/or develop assays that would address relevant key characteristics. This work is ongoing and the website [keycharacteristics.org](https://keycharacteristics.org) collates all available information and publications in this area (Key Characteristics, 2022). It also remains to be shown if *in-vitro* assays based on the key characteristics of hazardous chemicals will be sufficiently predictive of chemical hazards when used in pre-market assessments, rather than in *ex-ante* evaluations (where the key characteristics are currently used).

Another important aspect of testing is the development and validation of methods that reflect real-world chemical exposures from FCMs, including the effects of metabolites formed from FCCs in the human body. Migration testing protocols exist but ongoing research efforts need to be expanded and validated to ensure minimal loss of potentially hazardous chemicals during sample preparation (e.g. by using polar and apolar food simulants and by capturing not only non-volatile compounds, but also those that are semi-volatile and volatile) (Nerín et al., 2022; Oldring et al., 2023).

Lastly, a battery of screening assays addressing the SCOD needs to be defined and validated. This step will need the contribution of experts in each field to ensure that the selected endpoints are reliable and result in high confidence.

### 5.2. Data interpretation and integration

Methods must be developed to interpret and corroborate *in-vitro* test results. Individual assays should be integrated into an overall high-level/aggregated scheme (e.g. using visualization approaches such as ToxPi (Rajkumar et al., 2022; Marvel et al., 2018)). Also, non-targeted chemical analyses are challenging, as confirmation of identified compounds is very time- and labour intensive, and at times not possible at all. Also, reliable quantification of chemicals that lack analytical standards is not possible. Therefore, non-targeted approaches need to be advanced to allow for better identification of currently unknown compounds, especially when present at low concentrations. One way to improve the latter is to create comprehensive and open mass spectrometry libraries of FCCs, including NIAS. Ideally, an open-access repository of information about food contact material manufacturing processes and the major FCCs associated with specific materials should be generated. Confidential business information poses a critical obstacle, as the full disclosure of the chemical composition of FCMs is commonly not available. Accordingly, a mechanism needs to be developed that enables such an FCC library without infringing on intellectual property rights.

### 5.3. Science to inform decision making

The results of testing single chemicals or overall migrate (or, for a worst-case assessment, the extract) from a finished FCA using a battery of assays for each of the SCOD would need to be interpreted and integrated with available evidence to reach a conclusion within a regulatory context. A framework, similar to that available for read-across

(Lizarraga et al., 2023; European Chemicals Agency, 2017), should be developed to effectively utilize results and support conclusions that are actionable for policy makers and regulatory enforcement. The experience gained from the development of effect-based trigger values for water quality assessment in Europe could be highly informative (Escher et al., 2018; Neale et al., 2023). Here, effect-based trigger values have been developed as a means to interpret the results of *in-vitro* assays through linking the existing water quality guideline values to observed levels of bioactivity elicited by a reference chemical. Then, if a test chemical or mixture causes an activity above the trigger value set for a specific assay, it is highlighted for a follow-up assessment, such as calculation of concentration factors and *in-vitro* to *in-vivo* extrapolation (Escher and Neale, 2021; Robitaille et al., 2022; Escher et al., 2015). In theory, effect-based trigger values for FCMs could be developed following the same principle, e.g. by matching effect concentrations in relevant bioassays with existing specific migration limits for FCCs of concern, and possibly factoring in additional exposure-related parameters. This approach appears highly promising, since it has been demonstrated that derivation of effect-based trigger values greatly facilitates regulatory and practical uptake of *in-vitro* methods into specific assessment pipelines (Neale et al., 2023), but it is evident that further dedicated efforts are required for an effective implementation of such an approach to the safety assessment of FCMs.

## 6. Conclusion

The novel approach we present here is in line with the goals laid out in the EU's Chemicals Strategy for Sustainability (EU, 2020a), the EU Farm to Fork Strategy (EU, 2020b), and the European Parliament's report on FCMs (European Parliament, 2016), which emphasize the need for revising the food contact material regulation in Europe to adequately reflect recent scientific understanding and improve compliance. Further, this work adds to previous publications on policies and methods related to the risk assessment of food contact chemicals and materials (Zimmermann et al., 2022; Muncke et al., 2020; Muncke et al., 2017), and to the use of new approach methodologies for assessing the health impacts of industrial chemicals (Stucki et al., 2022).

We think that our vision to create safer FCMs by linking hazard identification more directly to human health has the potential to spur innovation in assay development and testing, and ultimately, for safer materials as such. Additionally, new findings on the key characteristics for the NCDs included in the SCOD, as well as mechanistic understanding derived from AOP research, will support the development of new assays.

Awareness of adverse health effects of synthetic chemicals is increasing globally, and the need is obvious for significant and urgent improvements in the ways that risks are assessed and managed for FCCs (Fenner and Scheringer, 2021).

## Declaration of Competing Interest

The authors have no competing interests to declare. For the sake of transparency, the authors list their relationships with various research funders and other organizations in the following. As researchers employed by the Food Packaging Forum Foundation (FPF) (JMB, BG, JM, LZ) or working *pro bono* as members of the Foundation's board (TB, JPM, MS) and its Scientific Advisory Board (SAB) (AMA, TJC, KJG, JJH, MVM, OVM, AN, CN, AMS, LT, MW, RTZ), we are reporting that the FPF receives donations from diverse companies that may be affected by the research reported in the enclosed paper. FPF funders have no influence on any of the work at FPF and were not involved in any way in the preparation of this manuscript. TB declares that he serves as the board member of the International Panel on Chemical Pollution (IPCP), the Swedish Toxicological Council and the EU Commission's Committee on Health, Environmental and Emerging Risks (SCHEER). All those activities are *pro bono* and have no bearing on the content of the manuscript.

None of the aforementioned organizations have had any impact on the content of the manuscript. TJC declares that he is the creator-founder of Sudoc, LLC, which deploys TAML catalysts for many applications and has potential for remediating FCCs in water. JL reports that she receives funding for another research project (ZonMw/Health-Holland Microplastics and Health project MOMENTUM 458001101) of which some partners may be affected by the research reported here. MVM is a paid consultant to the FPF. OVM is one of the representatives of the European Parliament on the European Chemical Agency's Management Board. JPM is co-founder and board member of Sudoc and he declares to have given all his shares to an irrevocable grantor trust so that he will not benefit financially if the company is successful. AN declares to have received travel reimbursement from universities, NGOs and scientific societies, to speak about endocrine-disrupting chemicals. LNV has received travel reimbursements from universities, governments, NGOs, and industry. She has received research funding from the US National Institutes of Health, the University of Massachusetts Amherst, and NGOs including the Cornell Douglas Foundation, the Allen Family Foundation, and the Great Neck Breast Cancer Coalition. She is a scientific advisor to Sudoc LLC. The FPF foundation board, whose members have no connection with any of the FPF's funders and receive no remuneration for their work, is legally obliged to guarantee that the work of the FPF is in no way influenced by the interests or views of the funders.

#### Data availability

No data were used for the research described in the article.

#### Acknowledgements

We are grateful to Michele La Merrill for constructive comments on this manuscript.

#### Authors' contributions

This manuscript was initiated by the FPF's SAB and guests participating in SAB meetings in 2017, 2018, 2019, 2020, 2021; AMA, MVM, JM, JPM, RTZ and MS were responsible for preparing an outline and a first version; JM, LVM, MVM and MS edited the final draft, and all authors contributed to the various intermediate versions, wrote separate sections of the manuscript, and approved the final version.

#### Funding

This work was supported by the FPF's own resources. FPF is a charitable foundation and it funded four meetings of its SAB with external scientific experts as guests (SMB, BCA, FAvH, JL, LNV) during which this manuscript was prepared. Funding for travel and accommodation was provided for three meetings and two additional meetings were held online. Neither FPF's SAB members nor guest participants were reimbursed for their contributions to this manuscript. FPF's funding sources are declared on its website (<https://www.foodpackagingforum.org/about-us/funding>). MS acknowledges funding by the CETOCOEN PLUS project (CZ.02.1.01/0.0/0.0/15\_003/0000469), the project CETOCOEN EXCELLENCE (CZ.02.1.01/0.0/0.0/17\_043/0009632), and RECETOX RI (LM2018121) financed by the Czech Ministry of Education, Youth and Sports.

#### Availability of data and materials

Not applicable.

#### Appendix A. Supplementary material

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.envint.2023.108161>.

#### References

- Akash, M.S., Hamid, S.S., Rehman, K., 2020. Bisphenol A-induced metabolic disorders: From exposure to mechanism of action. *Environ. Toxicol. Pharmacol.* 77, 103373.
- Akousson, F., Paul-Pont, I., Tallec, K., Huvet, A., Doyen, P., Dehaut, A., Duflos, G., 2023. Additives in polypropylene and polylactic acid food packaging: Chemical analysis and bioassays provide complementary tools for risk assessment. *Sci. Total Environ.* 857, 159318.
- Alger, H.M., Maffini, M.V., Kulkarni, N.R., Bongard, E.D., Neltner, T., 2013. Perspectives on how FDA assesses exposure to food additives when evaluating their safety: workshop proceedings. *Compr. Rev. Food Sci. Food Saf.* 12, 90–119.
- Al-Zoughool, M., Bird, M., Rice, J., Baan, R.A., Billard, M., Birkett, N., Krewski, D., Zielinski, J.M., 2019. Development of a database on key characteristics of human carcinogens. *J. Toxicol. Environ. Health. Part B* 22, 264–287.
- Ankley, G.T., Bennett, R.S., Erickson, R.J., Hoff, D.J., Hornung, M.W., Johnson, R.D., Mount, D.R., Nichols, J.W., Russom, C.L., Schmieder, P.K., Serrano, J.A., Tietge, J.E., Villeneuve, D.L., 2010. Adverse outcome pathways: a conceptual framework to support ecotoxicology research and risk assessment. *Environ. Toxicol. Chem.* 29, 730–741.
- Arshajothirmayi, V.A., Gulia, K.K., 2022. 26 - Neurotoxicity assays. In: Mohanan, P.V. (Ed.), *Biomedical Product and Materials Evaluation*. Woodhead Publishing.
- Arzuaga, X., Smith, M.T., Gibbons, C.F., Skakkebaek, N.E., Yost, E.E., Beverly, B.E.J., Hotchkiss, A.K., Hauser, R., Pagani, R.L., Schrader, S.M., Zeise, L., Prins, G.S., 2019. Proposed key characteristics of male reproductive toxicants as an approach for organizing and evaluating mechanistic evidence in human health hazard assessments. *Environ. Health Perspect.* 127, 65001.
- ATSDR, 2021. *Toxicological Profile for Perfluoroalkyls*, Accessed 23 Dec 2022. <https://www.cdc.gov/TSP/ToxProfiles/ToxProfiles.aspx?id=1117&tid=237>.
- Attina, T.M., Hauser, R., Sathyanarayana, S., Hunt, P.A., Bourguignon, J.-P., Myers, J.P., Joseph DiGangi, R., Zoeller, T., Trasande, L., 2016. Exposure to endocrine-disrupting chemicals in the USA: a population-based disease burden and cost analysis. *Lancet Diabetes Endo* 4, 996–1003.
- Audouze, K., Sarigiannis, D., Alonso-Magdalena, P., Brochot, C., Casas, M., Vrijheid, M., Babin, P.J., Karakitsios, S., Coumoul, X., Barouki, R., 2020. Integrative strategy of testing systems for identification of endocrine disruptors inducing metabolic disorders—an introduction to the OBERON project. *Int. J. Mol. Sci.* 21, 2988.
- Auerbach, S., Filer, D., Reif, D., Walker, V., Holloway, A.C., Schlezinger, J., Srinivasan, S., Svoboda, D., Judson, R., Bucher, J.R., Thayer, K.A., 2016. Prioritizing environmental chemicals for obesity and diabetes outcomes research: a screening approach using ToxCast™ high-throughput data. *Environ. Health Perspect.* 124, 1141–1154.
- Bach, C., Dauchy, X., Severin, I., Munoz, J.F., Etienne, S., Chagnon, M.C., 2013. Effect of temperature on the release of intentionally and non-intentionally added substances from polyethylene terephthalate (PET) bottles into water: chemical analysis and potential toxicity. *Food Chem.* 139, 672–680.
- Bailey, G.S., Reddy, A.P., Pereira, C.B., Harttig, U., Baird, W., Spitsbergen, J.M., Hendricks, J.D., Orner, G.A., Williams, D.E., Swenberg, J.A., 2009. Nonlinear cancer response at ultralow dose: a 40800-animal ED001 tumor and biomarker study. *Chem. Res. Toxicol.* 22, 1264–1276.
- Barr, D.B., Silva, M.J., Kato, K., Reidy, J.A., Malek, N.A., Hurtz, D., Sadowski, M., Needham, L.L., Calafat, A.M., 2003. Assessing human exposure to phthalates using monoesters and their oxidized metabolites as biomarkers. *Environ. Health Perspect.* 111, 1148–1151.
- Bauer, A., Jesús, F., Ramos, M.J.G., Lozano, A., Fernández-Alba, A.R., 2019. Identification of unexpected chemical contaminants in baby food coming from plastic packaging migration by high resolution accurate mass spectrometry. *Food Chem.* 295, 274–288.
- Benbrahim-Tallaa, L., Lauby-Secretan, B., Loomis, D., Guyton, K.Z., Grosse, Y., El Ghissassi, F., Bouvard, V., Guha, N., Mattock, H., Straif, K., 2014. Carcinogenicity of perfluorooctanoic acid, tetrafluoroethylene, dichloromethane, 1,2-dichloropropane, and 1,3-propane sultone. *Lancet Oncol.* 15, 924–925.
- Beneventi, E., Tietz, T., Merkel, S., 2020. Risk assessment of food contact materials. *EFSA J.* 18, e181109.
- Bengtstrom, L., Rosenmai, A.K., Trier, X., Jensen, L.K., Granby, K., Vinggaard, A.M., Driffield, M., Hojlslev Petersen, J., 2016. Non-targeted screening for contaminants in paper and board food-contact materials using effect-directed analysis and accurate mass spectrometry. *Food Addit. Contam. Part A: Chem. Anal. Control Expo. Risk Assess.* 33, 1080–1093.
- Bennett, D., Bellinger, D.C., Birnbaum, L.S., Bradman, A., Chen, A., Cory-Slechta, D.A., Engel, S.M., Daniele Fallin, M., Halladay, A., Hauser, R., Hertz-Picciotto, L., Kwiatkowski, C.F., Lanphear, B.P., Marquez, E., Marty, M., McPartland, J., Newschaffer, C.J., Payne-Sturges, D., Patisaul, H.B., Perera, F.P., Ritz, B., Sassi, J., Schantz, S.L., Webster, T.F., Whyatt, R.M., Woodruff, T.J., Thomas Zoeller, R., Anderko, L., Campbell, C., Conry, J.A., DeNicola, N., Gould, R.M., Hirtz, D., Huffling, K., Landrigan, P.J., Lavin, A., Miller, M., Mitchell, M.A., Rubin, L., Schettler, T., Tran, H.L., Acosta, A., Brody, C., Miller, E., Miller, P., Swanson, M., Witherspoon, N.O., 2016. Project TENDR: targeting environmental neurodevelopmental risks the TENDR Consensus Statement. *Environ. Health Perspect.* 124, A118–A122.
- Bergman, A., Andersson, A.-M., Becher, G., van den Berg, M., Blumberg, B., Bjerregaard, P., Bornehag, C.-G., Bornman, R., Brandt, I., Brian, J., Casey, S., Fowler, P., Frouin, H., Giudice, L., Iguchi, T., Hass, U., Jobling, S., Juul, A., Kidd, K., Kortenkamp, A., Lind, M., Martin, O., Muir, D., Ochieng, R., Olea, N., Norrgren, L., Ropstad, E., Ross, P., Ruden, C., Scheringer, M., 2013. Science and policy on endocrine disruptors must not be mixed: a reply to a “common sense” intervention by toxicology journal editors. *Environ. Health* 12, 69.

- Biedermann, M., Ingenhoff, J.-E., Zurfuh, M., Richter, L., Simat, T., Harling, A., Altkofer, W., Helling, R., Grob, K., 2013. Migration of mineral oil, photoinitiators and plasticizers from recycled paperboard into dry foods: a study under controlled conditions. *Food Addit. Contam. Part A: null-null*.
- Bil, W., Govarts, E., Zeilmaker, M.J., Woutersen, M., Bessems, J., Ma, Y., Thomsen, C., Haug, L.S., Lignell, S., Gyllenhammar, L., Palkovicova Murinova, L., Fabelova, L., Snoj Tratnik, J., Kosjek, T., Gabriel, C., Sarigiannis, D., Pedraza-Diaz, S., Esteban-López, M., Castaño, A., Rambaud, L., Riou, M., Franken, C., Colles, A., Vogel, N., Kolossa-Gehring, M., Halldorsson, T.I., Uhl, M., Schoeters, G., Santonen, T., Vinggaard, A.M., 2023. Approaches to mixture risk assessment of PFASs in the European population based on human hazard and biomonitoring data. *Int. J. Hyg. Environ. Health* 247, 114071.
- Biryol, D., Nicolas, C.L., Wambaugh, J., Phillips, K., Isaacs, K., 2017. High-throughput dietary exposure predictions for chemical migrants from food contact substances for use in chemical prioritization. *Environ. Int.* 108, 185–194.
- Blum, J., Masjosthusmann, S., Bartmann, K., Bendt, F., Dolde, X., Dönmez, A., Förster, N., Holzer, A.K., Hübenal, U., Keßel, H.E., Kilic, S., Klose, J., Pahl, M., Stürzl, L.C., Mangas, I., Terron, A., Crofton, K.M., Scholze, M., Mosig, A., Leist, M., Fritsche, E., 2022. Establishment of a human cell-based in vitro battery to assess developmental neurotoxicity hazard of chemicals. *Chemosphere* 311, 137035.
- Boisvert, A., Jones, S., Issop, L., Erythropel, H.C., Papadopoulos, V., Culty, M., 2016. In vitro functional screening as a means to identify new plasticizers devoid of reproductive toxicity. *Environ. Res.* 150, 496–512.
- Boomsma, C.M., Eijkemans, M.J., Hughes, E.G., Visser, G.H., Fauser, B.C., Macklon, N.S., 2006. A meta-analysis of pregnancy outcomes in women with polycystic ovary syndrome. *Hum. Reprod. Update* 12, 673–683.
- Bornehag, C.-G., Kitraki, E., Stamatakis, A., Panagiotidou, E., Rudén, C., Shu, H., Lindh, C., Ruegg, J., Gennings, C., 2019. A novel approach to chemical mixture risk assessment—linking data from population-based epidemiology and experimental animal tests. *Risk Anal.* 39, 2259–2271.
- Bornehag, C.-G., Engdahl, E., Hallerback, M.U., Wikström, S., Lindh, C., Ruegg, J., Tanner, E., Gennings, C., 2021. Prenatal exposure to bisphenols and cognitive function in children at 7 years of age in the Swedish SELMA study. *Environ. Int.* 150, 106433.
- Bradley, E., Coulier, L., 2007. An investigation into the reaction and breakdown products from starting substances used to produce food contact plastics. In: *Food Standards Agency (Ed.). Central Science Laboratory, London*.
- Bradley, E.L., Driffield, M., Harmer, N., Oldring, P.K.T., Castle, L., 2008. Identification of potential migrants in epoxy phenolic can coatings. *Int. J. Polym. Anal. Charact.* 13, 200–223.
- Bschir, K. (accepted). Risk, uncertainty and precaution in science: the case of the threshold of toxicological concern approach in food toxicology. *J. Sci. Eng. Ethics*.
- Buckley, J.P., Engel, S.M., Braun, J.M., Whyatt, R.M., Daniels, J.L., Mendez, M.A., Richardson, D.B., Xu, Y., Calafat, A.M., Wolff, M.S., Lanphear, B.P., Herring, A.H., Rundle, A.G., 2016. Prenatal phthalate exposures and body mass index among 4- to 7-year-old children: a pooled analysis. *Epidemiology* 27, 449–458.
- Caffrey, T.M., Button, E.B., Robert, J., 2021. Toward three-dimensional in vitro models to study neurovascular unit functions in health and disease. *Neural Regen. Res.* 16, 2132–2140.
- Calafat, A., Kuklennyik, Z., Reidy, J.A., Caudill, S.P., Ekong, J., Needham, L.L., 2005. Urinary concentrations of bisphenol A and 4-nonylphenol in a human reference population. *Environ. Health Perspect.* 113, 391–395.
- Caporale, N., Leemans, M., Birgersson, L., Germain, P.-L., Cheroni, C., Borbély, G., Engdahl, E., Lindh, C., Bressan, R.B., Cavallo, F., Choren, N.E., D'Agostino, G.A., Pollard, S.M., Rigoli, M.T., Tenderini, E., Tobon, A.L., Trattaro, S., Troglia, F., Zanella, M., Bergman, Å., Ddamimopoulou, P., Jönsson, M., Kiess, W., Kitraki, E., Kiviranta, H., Nånberg, E., Öberg, M., Rantakokko, P., Rudén, C., Söder, O., Bornehag, C.-G., Demeneix, B., Fini, J.-B., Gennings, C., Ruegg, J., Sturve, J., Testa, G., 2022. From cohorts to molecules: adverse impacts of endocrine disrupting mixtures. *Science* 375, eabe8244.
- Castle, L., Mayo, A., Crews, C., Gilbert, J., 1989. Migration of Poly(ethylene terephthalate) (PET) Oligomers from PET Plastics into Foods during Microwave and Conventional Cooking and into Bottled Beverages. *J. Food Prot.* 52, 337–342.
- Cediell-Ulloa, A., Lupu, D.L., Johansson, Y., Hinojosa, M., Özel, F., Ruegg, J., 2022. Impact of endocrine disrupting chemicals on neurodevelopment: the need for better testing strategies for endocrine disruption-induced developmental neurotoxicity. *Expert. Rev. Endocrinol. Metab.* 17, 131–141.
- Chakori, S., Aziz, A.A., Smith, C., Dargusch, P., 2021. Untangling the underlying drivers of the use of single-use food packaging. *Ecol. Econ.* 185, 107063.
- Chamorro-García, R., Blumberg, B., 2019. Current research approaches and challenges in the obesogen field. *Front. Endocrinol. (Lausanne)* 10, 167.
- Chamorro-García, R., Diaz-Castillo, C., Shoucri, B.M., Käch, H., Leavitt, R., Shioda, T., Blumberg, B., 2017. Ancestral perinatal obesogen exposure results in a transgenerational thrifty phenotype in mice. *Nat. Commun.* 8, 2012.
- Charazac, A., Hinault, C., Dolfi, B., Hautier, S., Decondé Le Butor, C., Bost, F., Chevalier, N., 2022. Low doses of PFOA promote prostate and breast cancer cells growth through different pathways. *Int. J. Mol. Sci.* 23.
- Chiara, F., Indraccolo, S., Trevisan, A., 2020. Filling the gap between risk assessment and molecular determinants of tumor onset. *Carcinogenesis* 42, 507–516.
- CIPA, 2019. CIPA Initiative. Accessed 23 Dec 2022. <https://cipaproject.org/>.
- Cooper, B.L., Posnack, N.G., 2022. Characteristics of bisphenol cardiotoxicity: impaired excitability, contractility, and relaxation. *Cardiovasc. Toxicol.* 22, 273–280.
- Correia-Sá, L., Schütze, A., Norberto, S., Calhau, C., Domingues, V.F., Koch, H.M., 2017. Exposure of Portuguese children to the novel non-phthalate plasticizer di-(isononyl)-cyclohexane-1,2-dicarboxylate (DINCH). *Environ. Int.* 102, 79–86.
- Corsini, E., Roggen, E.L., 2017. Overview of in vitro assessment of immunotoxicity. *Curr. Opin. Toxicol.* 5, 13–18.
- Cortéjade, A., Buleté, A., Prouteau, L., Chatti, S., Cren, C., Vulliet, E., 2017. Development and optimisation of home-made stir bar sorptive extraction for analysis of plastic additives: application in human urine. *Anal. Methods* 9, 3549–3560.
- Corton, J.C., Liu, J., Kleinstreuer, N., Gwinn, M.R., Ryan, N., 2022. Towards replacement of animal tests with in vitro assays: a gene expression biomarker predicts in vitro and in vivo estrogen receptor activity. *Chem. Biol. Interact.* 363, 109995.
- Coumoul, X., Barouki, R., Koulal, M., Audouze, K., Tomkiewicz, C., 2022. Activation of the AhR leading to breast cancer. *AOP* 439. Accessed 23 Dec 2022. <https://aopwiki.org/aops/439>.
- Dales, R.E., Kauri, L.M., Cakmak, S., 2018. The associations between phthalate exposure and insulin resistance,  $\beta$ -cell function and blood glucose control in a population-based sample. *Sci. Total Environ.* 612, 1287–1292.
- Demeneix, B.A., Slama, R., 2019. Endocrine Disruptors: From the scientific evidence to human health protection. In: *Report to the European Parliament. European Parliament*.
- Deprouw, C., Courties, A., Fini, J.B., Clerget-Froidevaux, M.S., Demeneix, B., Berenbaum, F., Sellam, J., Louati, K., 2022. Pollutants: a candidate as a new risk factor for osteoarthritis—results from a systematic literature review. *RMD Open* 8.
- Desai, M., Ferrini, M.G., Han, G., Jellyman, J.K., Ross, M.G., 2018a. In vivo maternal and in vitro BPA exposure effects on hypothalamic neurogenesis and appetite regulators. *Environ. Res.* 164, 45–52.
- Desai, M., Ferrini, M.G., Jellyman, J.K., Han, G., Ross, M.G., 2018b. In vivo and in vitro bisphenol A exposure effects on adiposity. *J. Dev. Health Dis.* 9, 678–687.
- Desmarchais, A., Têteau, O., Papillier, P., Jaubert, M., Duart, X., Binet, A., Maillard, V., Elis, S., 2020. Bisphenol S impaired in vitro ovine early developmental oocyte competence. *Int. J. Mol. Sci.* 21, 1238.
- DeWitt, J.C., Blossom, S.J., Schaidler, L.A., 2019. Exposure to per-fluoroalkyl and polyfluoroalkyl substances leads to immunotoxicity: epidemiological and toxicological evidence. *J. Exposure Sci. Environ. Epidemiol.* 29, 148–156.
- Dhimolea, E., Wadia, P.R., Murray, T.J., Settles, M.L., Treitman, J.D., Sonnenschein, C., Shioda, T., Soto, A.M., 2014. Prenatal exposure to BPA alters the epigenome of the rat mammary gland and increases the propensity to neoplastic development. *PLoS One* 9, e99800.
- Dionisi, G., Oldring, P.K., 2002. Estimates of per capita exposure to substances migrating from canned foods and beverages. *Food Addit. Contam.* 19, 891–903.
- Domínguez-Romero, E., Komprdová, K., Kalina, J., Bessems, J., Karakitsios, S., Sarigiannis, D.A., Scheringer, M., 2022. Time-trends in human urinary concentrations of phthalates and substitutes DEHT and DINCH in Asian and North American countries (2009–2019). *J. Exposure Sci. Environ. Epidemiol.*
- Eales, J., Bethel, A., Galloway, T., Hopkins, P., Morrissey, K., Short, R.E., Garside, R., 2022. Human health impacts of exposure to phthalate plasticizers: an overview of reviews. *Environ. Int.* 158, 106903.
- EFSA, 2008. Guidance document on the submission of a dossier on a substance to be used in Food Contact Materials for evaluation by EFSA by the Panel on additives, flavourings, processing aids and materials in contact with food (AFC). *EFSA J.*
- El-Kersh, K., Danielle Hopkins, C., Wu, X., Rai, S.N., Cai, L., Huang, J., 2022. Plasma level of antimony correlates with pulmonary arterial hypertension severity. *Curr. Res. Toxicol.* 3, 100080.
- Escher, B.I., Neale, P.A., Leusch, F.D., 2015. Effect-based trigger values for in vitro bioassays: reading across from existing water quality guideline values. *Water Res.* 81, 137–148.
- Escher, B.I., Ait-Aïssa, S., Behnisch, P.A., Brack, W., Brion, F., Brouwer, A., Buchinger, S., Crawford, S.E., Pasquier, D.D., Hamers, T., Hettwer, K., Hilscherová, K., Hollert, H., Kase, R., Kienle, C., Tindall, A.J., Tuerk, J., van der Oost, R., Vermeirssen, E., Neale, P.A., 2018. Effect-based trigger values for in vitro and in vivo bioassays performed on surface water extracts supporting the environmental quality standards (EQS) of the European Water Framework Directive. *Sci. Total Environ.* 628–629, 748–765.
- Escher, B.I., Neale, P.A., 2021. Effect-based trigger values for mixtures of chemicals in surface water detected with in vitro bioassays. *Environ. Toxicol. Chem.* 40, 487–499.
- Estill, M., Hauser, R., Nassan, F.L., Moss, A., Krawetz, S.A., 2019. The effects of di-butyl phthalate exposure from medications on human sperm RNA among men. *Sci. Rep.* 9, 12397.
- EU, 2011. COMMISSION REGULATION (EU) No 10/2011 of 14 January 2011 on plastic materials and articles intended to come into contact with food. In: *EU 10/2011, edited by European Union. European Union, Brussels*.
- EU, 2020a. Chemicals Strategy for Sustainability. Towards a Toxic-Free Environment. *European Commission*.
- EU, 2020b. A Farm to Fork Strategy for a fair, healthy and environmentally-friendly food system. *European Commission*.
- European Chemicals Agency, 2017. Read-Across Assessment Framework (RAAF). *European Chemicals Agency*.
- European Parliament, 2016. Implementation of the Food Contact Materials Regulation. *European Parliament resolution of 6 October 2016 on the implementation of the Food Contact Materials Regulation (EC) No 1935/2004 (2015/2259(INI))*.
- FDA, 2007. Guidance for Industry: Preparation of Premarket Submissions for Food Contact Substances: Chemistry Recommendations. Accessed 23 December 2022. <https://www.fda.gov/regulatory-information/search-fda-guidance-documents/guidance-industry-preparation-premarket-submissions-food-contact-substances-chemistry>.
- FDA, 2022. Impact Story: Improved Assessment of Cardiotoxic Risk in Drug Candidates: The Comprehensive in vitro Proarrhythmia Assay. Accessed 23 Dec 2022. <https://www.fda.gov/drugs/regulatory-science-action/impact-story-improved-assessment-cardiotoxic-risk-drug-candidates-comprehensive-vitro-proarrhythmia>.

- Feil, R., Fraga, M.F., 2012. Epigenetics and the environment: emerging patterns and implications. *Nat. Rev. Genet.* 13, 97–109.
- Fenner, K., Scherlinger, M., 2021. The need for chemical simplification as a logical consequence of ever-increasing chemical pollution. *Environ. Sci. Technol.* 55, 14470–14472.
- Filer, D.L., Hoffman, K., Sargis, R.M., Trasande, L., Kassotis, C.D., 2022. On the utility of ToxCast-based predictive models to evaluate potential metabolic disruption by environmental chemicals. *Environ. Health Perspect.* 130, 57005.
- Fitzgerald, J.A., Könemann, S., Krümpelmann, L., Županič, A., vom Berg, C., 2021. Approaches to test the neurotoxicity of environmental contaminants in the zebrafish model: from behavior to molecular mechanisms. *Environ. Toxicol. Chem.* 40, 989–1006.
- Fitz-James, M.H., Cavalli, G., 2022. Molecular mechanisms of transgenerational epigenetic inheritance. *Nat. Rev. Genet.* 23, 325–341.
- Foresta, C., Tescari, S., Di Nisio, A., 2018. Impact of perfluorochemicals on human health and reproduction: a male's perspective. *J. Endocrinol. Invest.* 41, 639–645.
- Friedman, K.P., Crofton, K., Gilbert, M., 2022. Upregulation of Thyroid Hormone Catabolism via Activation of Hepatic Nuclear Receptors, and Subsequent Adverse Neurodevelopmental Outcomes in Mammals. *AOP 8*. Accessed 23 Dec 2022. <https://aopwiki.org/aops/8>.
- Fu, X., Xu, J., Zhang, R., Yu, J., 2020. The association between environmental endocrine disruptors and cardiovascular diseases: a systematic review and meta-analysis. *Environ. Res.* 187, 109464.
- Gao, X., Wang, H.S., 2014. Impact of bisphenol a on the cardiovascular system - epidemiological and experimental evidence and molecular mechanisms. *Int. J. Environ. Res. Public Health* 11, 8399–8413.
- García-Calvo, E., Machuca, A., Nerín, C., Rosales-Conrado, N., Anunciação, D.S., Luque-García, J.L., 2020. Integration of untargeted and targeted mass spectrometry-based metabolomics provides novel insights into the potential toxicity associated to surfynol. *Food Chem. Toxicol.* 146, 111849.
- Gascon, M., Casas, M., Morales, E., Valvi, D., Ballesteros-Gómez, A., Luque, N., Rubio, S., Monfort, N., Ventura, R., Martínez, D., Sunyer, J., Vrijheid, M., 2015. Prenatal exposure to bisphenol A and phthalates and childhood respiratory tract infections and allergy. *J. Allergy Clin. Immunol.* 135, 370–378.e7.
- Gear, R., Kendzioriski, J.A., Belcher, S.M., 2017. Effects of bisphenol A on incidence and severity of cardiac lesions in the NCTR-Sprague-Dawley rat: a CLARITY-BPA study. *Toxicol. Lett.* 275, 123–135.
- Geiger, S.D., Yao, P., Vaughn, M.G., Qian, Z., 2021. PFAS exposure and overweight/obesity among children in a nationally representative sample. *Chemosphere* 268, 128852.
- Germolec, D.R., Lebec, H., Anderson, S.E., Burleson, G.R., Cardenas, A., Corsini, E., Elmore, S.E., Kaplan, B.L.F., Lawrence, B.P., Lehmann, G.M., Maier, C.C., McHale, C.M., Myers, L.P., Pallardy, M., Rooney, A.A., Zeise, L., Zhang, L., Smith, M.T., 2022. Consensus on the key characteristics of immunotoxic agents as a basis for hazard identification. *Environ. Health Perspect.* 130, 105001.
- Germolec, D., Luebke, R., Rooney, A., Shipkowski, K., Vandebriel, R., van Loveren, H., 2017. Immunotoxicology: A brief history, current status and strategies for future immunotoxicity assessment. *Curr. Opin. Toxicol.* 5, 55–59.
- Geueke, B., Groh, K., Muncke, J., 2018. Food packaging in the circular economy: overview of chemical safety aspects for commonly used materials. *J. Clean. Prod.* 193, 491–505.
- Geueke, B., Groh, K.J., Maffini, M.V., Martin, O.V., Boucher, J.M., Chiang, Y.-T., Gwosdz, F., Jieh, P., Kassotis, C.D., Łańska, P., Myers, J.P., Odermatt, A., Parkinson, L.V., Schreier, V.N., Srebny, V., Zimmermann, L., Scherlinger, M., Muncke, J., 2022. Systematic evidence on migrating and extractable food contact chemicals: Most chemicals detected in food contact materials are not listed for use. *Crit. Rev. Food Sci. Nutr.* 1–11.
- Geueke, B., Phelps, D.W., Parkinson, L.V., Muncke, J., 2023. Hazardous chemicals in recycled and reusable plastic food packaging. *Cambridge Prisms: Plastics* 1, e7.
- Geueke, B., 2018. Non-intentionally added substances (NIAS). In: Dossier, P.F.F. (Ed.), *Food Packaging Forum*. Food Packaging Forum Foundation.
- Goralczyk, K., 2021. A review of the impact of selected anthropogenic chemicals from the group of endocrine disruptors on human health. *Toxics*.
- Gore, A.C., Chappell, V.A., Fenton, S.E., Flaws, J.A., Nadal, A., Prins, G.S., Toppari, J., Zoeller, R.T., 2015. EDC-2: the endocrine society's second scientific statement on endocrine-disrupting chemicals. *Endocr. Rev.* 36, E1–E150.
- Grandjean, P., Landrigan, P.J., 2006. Developmental neurotoxicity of industrial chemicals. *Lancet* 368, 2167–2178.
- Grandjean, P., Heilmann, C., Weihe, P., Nielsen, F., Mogensén, U.B., Timmermann, A., Budtz-Jørgensen, E., 2017. Estimated exposures to perfluorinated compounds in infancy predict attenuated vaccine antibody concentrations at age 5-years. *J. Immunotoxicol.* 14, 188–195.
- Grob, K., Biedermann, M., Scherbaum, E., Roth, M., Rieger, K., 2006. Food contamination with organic materials in perspective: packaging materials as the largest and least controlled source? A view focusing on the European situation. *Crit. Rev. Food Sci. Nutr.* 46, 529–535.
- Groh, K.J., Geueke, B., Martin, O., Maffini, M., Muncke, J., 2021. Overview of intentionally used food contact chemicals and their hazards. *Environ. Int.* 150, 106225.
- Groh, K.J., Muncke, J., 2017. In vitro toxicity testing of food contact materials: state-of-the-art and future challenges. *Compr. Rev. Food Sci. Food Saf.* 16, 1123–1150.
- Guyton, K.Z., Schubauer-Berigan, M.K., 2021. Invited perspective: prioritizing chemical testing and evaluation using validated in vitro assays relevant to key characteristics. *Environ. Health Perspect.* 129, 071303.
- Guyton, K.Z., Rusyn, I., Chiu, W.A., Corpet, D.E., van den Berg, M., Ross, M.K., Christiani, D.C., Beland, F.A., Smith, M.T., 2018. Application of the key characteristics of carcinogens in cancer hazard identification. *Carcinogenesis* 39, 614–622.
- Hager, E., Chen, J., Zhao, L., 2022. Minireview: parabens exposure and breast cancer. *Int. J. Environ. Res. Public Health* 19.
- He, X., Liu, Y., Xu, B., Gu, L., Tang, W., 2018. PFOA is associated with diabetes and metabolic alteration in US men: National Health and Nutrition Examination Survey 2003–2012. *Sci. Total Environ.* 625, 566–574.
- Heindel, J.J., 2019. History of the obesogen field: looking back to look forward. *Front. Endocrinol. (Lausanne)* 10, 14.
- Heindel, J.J., Howard, S., Agay-Shay, K., Arrebola, J.P., Audouze, K., Babin, P.J., Barouki, R., Bansal, A., Blanc, E., Cave, M.C., Chatterjee, S., Chevalier, N., Choudhury, M., Collier, D., Connolly, L., Coumoul, X., Garruti, G., Gilbertson, M., Hoepner, L.A., Holloway, A.C., Howell 3rd, G., Kassotis, C.D., Kay, M.K., Kim, M.J., Lagadic-Gossman, D., Langouet, S., Legrand, A., Li, Z., Le Mentec, H., Lind, L., Monica Lind, P., Lustig, R.H., Martin-Chouly, C., Munic Kos, V., Podgehard, N., Roepeke, T.A., Sargis, R.M., Starling, A., Tomlinson, C.R., Touma, C., Vondracek, J., Vom Saal, F., Blumberg, B., 2022. Obesity II: Establishing causal links between chemical exposures and obesity. *Biochem. Pharmacol.* 199, 115015.
- Hessel, E.V.S., Tonk, E.C.M., Bos, P.M.J., van Loveren, H., Piersma, A.H., 2015. Developmental immunotoxicity of chemicals in rodents and its possible regulatory impact. *Crit. Rev. Toxicol.* 45, 68–82.
- Hliseníková, H., Petrovičová, I., Kolena, B., Šidlovská, M., Sirotkin, A., 2021. Effects and mechanisms of phthalates' action on neurological processes and neural health: a literature review. *Pharmacol. Rep.* 73, 386–404.
- Horodytska, O., Cabanes, A., Fullana, A., 2020. Non-intentionally added substances (NIAS) in recycled plastics. *Chemosphere* 251, 126373.
- Hsieh, T.J., Hsieh, P.C., Tsai, Y.H., Wu, C.F., Liu, C.C., Lin, M.Y., Wu, M.T., 2012. Melamine induces human renal proximal tubular cell injury via transforming growth factor- $\beta$  and oxidative stress. *Toxicol. Sci.* 130, 17–32.
- Huff, J., Jacobson, M.F., Davis, D.L., 2008. The limits of two-year bioassay exposure regimens for identifying chemical carcinogens. *Environ. Health Perspect.* 116, 1439–1442.
- Hunt, P.R., Olejnik, N., Bailey, K.D., Vaught, C.A., Sprando, R.L., 2018. *C. elegans* development and activity test detects mammalian developmental neurotoxins. *Food Chem. Toxicol.* 121, 583–592.
- Hwang, S.H., Yeom, H., Han, B.I., Ham, B.J., Lee, Y.M., Han, M.R., Lee, M., 2020. Predicting carcinogenic mechanisms of non-genotoxic carcinogens via combined analysis of global DNA methylation and in vitro cell transformation. *Int. J. Mol. Sci.* 21.
- Hyun, S.A., Lee, C.Y., Ko, M.Y., Chon, S.H., Kim, Y.J., Seo, J.W., Kim, K.K., Ka, M., 2021. Cardiac toxicity from bisphenol A exposure in human-induced pluripotent stem cell-derived cardiomyocytes. *Toxicol. Appl. Pharmacol.* 428, 115696.
- IARC, 2010. Some Aromatic Amines, Organic Dyes, and Related Exposures. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Volume 99. In: IARC Monographs, edited by International Agency for Research on Cancer. IARC, Lyon, France.
- IARC, 2012a. Chemical Agents and Related Occupations. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Volume 100F. In: IARC Monographs, International Agency for Research on Cancer. International Agency for Research on Cancer, Lyon, France.
- IARC, 2012b. Some chemicals present in industrial and consumer products, food and drinking-water. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Volume 101. In: IARC Monographs, edited by International Agency for Research on Cancer. IARC, Lyon, France.
- IARC, 2016. Some chemicals used as solvents and in polymer manufacture. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Volume 110. In: IARC Monographs, International Agency for Research on Cancer. Lyon, France.
- IARC, 2019. Some chemicals that cause tumours of the urinary tract in rodents. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans Volume 119. In: IARC Monographs, edited by International Agency for Research on Cancer. Lyon, France.
- IARC, 2022. Agents classified by the IARC monographs, Volumes 1–129. International Agency for Research on Cancer. <https://monographs.iarc.who.int/agents-classified-by-the-iarc/>.
- Isaacs, K.K., Wall, J.T., Williams, A.R., Hobbie, K.A., Sobus, J.R., Ulrich, E., Lyons, D., Dionisio, K.L., Williams, A.J., Grulke, C., Foster, C.A., McCoy, J., Bevington, C., 2022. A harmonized chemical monitoring database for support of exposure assessments. *Sci. Data* 9, 314.
- Janesick, A.S., Dimastrogiovanni, G., Vanek, L., Boulos, C., Chamorro-García, R., Tang, W., Blumberg, B., 2016. On the Utility of ToxCast™ and ToxPi as Methods for Identifying New Obesogens. *Environ. Health Perspect.* 124, 1214–1226.
- Jeddi, Z., Maryam, N.B., Hopf, S.V., Price, A.B., Paini, A., van Thriel, C., Benfenati, E., Ndaw, S., Bessems, J., Behnisch, P.A., Leng, G., Duca, R.-C., Verhagen, H., Cubadda, F., Brennan, L., Ali, I., David, A., Mustieles, V., Fernandez, M.F., Louro, H., Pasanen-Kase, R., 2021. Towards a systematic use of effect biomarkers in population and occupational biomonitoring. *Environ. Int.* 146, 106257.
- Jickells, S.M., Gancedo, P., Nerin, C., Castle, L., Gilbert, J., 1993. Migration of styrene monomer from thermoset polyester cookware into foods during high temperature applications. *Food Addit. Contam.* 10, 567–573.
- Jokinen, M.P., Lieuallen, W.G., Johnson, C.L., Dunnick, J., Nyska, A., 2005. Characterization of spontaneous and chemically induced cardiac lesions in rodent model systems: the national toxicology program experience. *Cardiovasc. Toxicol.* 5, 227–244.
- Jokinen, M.P., Lieuallen, W.G., Boyle, M.C., Johnson, C.L., Malarkey, D.E., Nyska, A., 2011. Morphologic aspects of rodent cardiotoxicity in a retrospective evaluation of national toxicology program studies. *Toxicol. Pathol.* 39, 850–860.

- Jorgensen, A., Svingen, T., Miles, H., Chetty, T., Stukenborg, J.B., Mitchell, R.T., 2021. Environmental Impacts on Male Reproductive Development: Lessons from Experimental Models. *Hormone Research in Paediatrics*.
- Jun, J.H., Oh, J.E., Shim, J.K., Kwak, Y.L., Cho, J.S., 2021. Effects of bisphenol A on the proliferation, migration, and tumor growth of colon cancer cells: In vitro and in vivo evaluation with mechanistic insights related to ERK and 5-HT3. *Food Chem. Toxicol.* 158, 112662.
- Jung, S.K., Choi, W., Kim, S.Y., Hong, S., Jeon, H.L., Joo, Y., Lee, C., Choi, K., Kim, S., Lee, K.J., Yoo, J., 2022. Profile of environmental chemicals in the Korean population—results of the Korean National Environmental Health Survey (KoNEHS) Cycle 3, 2015–2017. *Int. J. Environ. Res. Public Health* 19.
- Kassotis, C.D., Stapleton, H.M., 2019. Endocrine-mediated mechanisms of metabolic disruption and new approaches to examine the public health threat. *Front. Endocrinol. (Lausanne)* 10, 39.
- Kassotis, C.D., Vandenberg, L.N., Demeneix, B.A., Porta, M., Slama, R., Trasande, L., 2020. Endocrine-disrupting chemicals: economic, regulatory, and policy implications. *Lancet Diabetes Endocrinol.* 8, 719–730.
- Kassotis, C.D., Vom Saal, F.S., Babin, P.J., Lagadic-Gossman, D., Le Mentec, H., Blumberg, B., Mohajer, N., Legrand, A., Munic Kos, V., Martin-Chouly, C., Podechard, N., Langouët, S., Touma, C., Barouki, R., Kim, M.J., Audouze, K., Choudhury, M., Shree, N., Bansal, A., Howard, S., Heindel, J.J., 2022. Obesity III: Obesogen assays: limitations, strengths, and new directions. *Biochem. Pharmacol.* 199, 115014.
- Keri, R.A., Ho, S.-M., Hunt, P.A., Knudsen, K.E., Soto, A.M., Prins, G.S., 2007. An evaluation of evidence for the carcinogenic activity of bisphenol A. *Reprod. Toxicol.* 24, 240–252.
- Key Characteristics, 2022. Key Characteristics. Identifying the Key Characteristics of Hazardous Chemicals and Other Exposures: A Collaborative Approach.** Accessed 23 Dec 2022. <https://keycharacteristics.org/>.
- Kilic, O., Pamiés, D., Lavell, E., Schiapparelli, P., Feng, Y., Hartung, T., Bal-Price, A., Hogberg, H.T., Quinones-Hinojosa, A., Guerrero-Cazares, H., Levchenko, A., 2016. Brain-on-a-chip model enables analysis of human neuronal differentiation and chemotaxis. *Lab Chip* 16, 4152–4162.
- Kim, E.H., Jeon, B.H., Kim, J., Kim, Y.M., Han, Y., Ahn, K., Cheong, H.K., 2017. Exposure to phthalates and bisphenol A are associated with atopic dermatitis symptoms in children: a time-series analysis. *Environ. Health* 16, 24.
- Kim, H.S., Lee, Y.J., Koo, Y.J., Pack, E.C., Lim, K.M., Choi, D.W., 2021. Migration of monomers, plastic additives, and non-intentionally added substances from food utensils made of melamine–formaldehyde resin following ultraviolet sterilization. *Food Control* 125, 107981.
- Kim, J.I., Lee, Y.A., Shin, C.H., Hong, Y.-C., Kim, B.-N., Lim, Y.-H., 2022. Association of bisphenol A, bisphenol F, and bisphenol S with ADHD symptoms in children. *Environ. Int.* 161, 107093.
- Kirk, A.B., 2006. Environmental perchlorate: why it matters. *Anal. Chim. Acta* 567, 4–12.
- Koch, H.M., Calafat, A.M., 2009. Human body burdens of chemicals used in plastic manufacture. *Philos. Trans. R. Soc. Lond. B: Biol. Sci.* 364, 2063–2078.
- Kofron, C.M., Kim, T.Y., Munarin, F., Soepriatna, A.H., Kant, R.J., Mende, U., Choi, B.-R., Coulombe, K.L.K., 2021. A predictive in vitro risk assessment platform for pro-arrhythmic toxicity using human 3D cardiac microtissues. *Sci. Rep.* 11, 10228.
- Kortenkamp, A., 2020. Which chemicals should be grouped together for mixture risk assessments of male reproductive disorders? *Mol. Cell. Endocrinol.* 499, 110581.
- Kortenkamp, A., Faust, M., 2018. Regulate to reduce chemical mixture risk. *Science* 361, 224–226.
- Kortenkamp, A., Axelstad, M., Baig, A.H., Bergman, Å., Bornehag, C.-G., Ceniñ, P., Christiansen, S., Demeneix, B., Derakhshan, A., Fini, J.-B., Frädrich, C., Hamers, T., Hellwig, L., Köhrle, J., Korevaar, T.I.M., Lindberg, J., Martin, O., Meima, M.E., Mergenthaler, P., Nikolov, N., Pasquier, D.D., Peeters, R.P., Platzack, B., Ramhøj, L., Rемаud, S., Renko, K., Scholze, M., Stachelscheid, H., Svingen, T., Wagenaar, F., Wedebye, E.B., Thomas Zoeller, R., 2020. Removing critical gaps in chemical test methods by developing new assays for the identification of thyroid hormone system-disrupting chemicals—the ATHENA project. *Int. J. Mol. Sci.* 21, 3123.
- Koster, S., Rennen, M., Leeman, W., Houben, G., Muilwijk, B., van Acker, F., Krul, L., 2013. A novel safety assessment strategy for non-intentionally added substances (NIAS) in carton food contact materials. *Food Additi. Contaminants: Part A* 31, 422–443.
- Koster, S., Bani-Estivals, M.H., Bonuomo, M., Bradley, E., Chagnon, M.C., Garcia, M.L., Godts, F., Gude, T., Helling, R., Paseiro-Losada, P., Pieper, G., Rennen, M., Simat, T., Spack, L., 2015. Guidance on best practices on the risk assessment of non-intentionally added substances (NIAS) in food contact materials and articles. *ILSI Europe*.
- Krewski, D., Bird, M., Al-Zoughool, M., Birkett, N., Billard, M., Milton, B., Rice, J.M., Grosse, Y., Coglian, V.J., Hill, M.A., Baan, R.A., Little, J., Zielinski, J.M., 2019. Key characteristics of 86 agents known to cause cancer in humans. *J. Toxicol. Environ. Health, Part B* 22, 244–263.
- Krishna, S., Berridge, B., Kleinstreuer, N., 2021. High-Throughput Screening to Identify Chemical Cardiotoxic Potential. *Chem. Res. Toxicol.* 34, 566–583.
- Küblbeck, J., Vuorio, T., Niskanen, J., Fortino, V., Braeuning, A., Abass, K., Rautio, A., Hakola, J., Honkakoski, P., Levenon, A.-L., 2020. The EDCMET Project: Metabolic Effects of Endocrine Disruptors. *Int. J. Mol. Sci.* 21, 3021.
- Lane, J.M., Doyle, J.R., Fortin, J.P., Kopin, A.S., Ordovás, J.M., 2014. Development of an OP9 derived cell line as a robust model to rapidly study adipocyte differentiation. *PLoS One* 9, e112123.
- Larsson-Nyrén, G., Sehlin, J., Rorsman, P., Renström, E., 2001. Perchlorate stimulates insulin secretion by shifting the gating of L-type Ca<sup>2+</sup> currents in mouse pancreatic B-cells towards negative potentials. *Pflugers Arch.* 441, 587–595.
- Leeman, W., Krul, L., 2015. Non-intentionally added substances in food contact materials: how to ensure consumer safety. *Curr. Opin. Food Sci.* 6, 33–37.
- Legler, J., Zalko, D., Jourdan, F., Jacobs, M., Fromenty, B., Balaguer, P., Bourguet, W., Kos, V.M., Nadal, A., Beausoleil, C., Cristobal, S., Remy, S., Ermler, S., Margiotta-Casaluci, L., Griffin, J.L., Blumberg, B., Chesné, C., Hoffmann, S., Andersson, P.L., Kamstra, J.H., 2020. The GOLIATH Project: Towards an Internationally Harmonised Approach for Testing Metabolism Disrupting Compounds. *Int. J. Mol. Sci.* 21, 3480.
- Levine, H., Jørgensen, N., Martino-Andrade, A., Mendiola, J., Weksler-Derri, D., Mindlis, I., Pinotti, R., Swan, S.H., 2017. Temporal trends in sperm count: a systematic review and meta-regression analysis. *Hum. Reprod. Update* 23, 646–659.
- Levine, H., Jørgensen, N., Martino-Andrade, A., Mendiola, J., Weksler-Derri, D., Jolles, M., Pinotti, R., Swan, S.H., 2022. Temporal trends in sperm count: a systematic review and meta-regression analysis of samples collected globally in the 20th and 21st centuries. *Hum. Reprod. Update* 29, 157–176.
- Li, N., Papandonatos, G.D., Calafat, A.M., Yolton, K., Lanphear, B.P., Chen, A., Braun, J.M., 2020. Gestational and childhood exposure to phthalates and child behavior. *Environ. Int.* 144, 106036.
- Li, J., Zheng, L., Wang, X., Yao, K., Shi, L., Sun, X., Yang, G., Jiang, L., Zhang, C., Wang, Y., Jiang, L., Liu, X., 2019. Taurine protects INS-1 cells from apoptosis induced by Di(2-ethylhexyl) phthalate via reducing oxidative stress and autophagy. *Toxicol. Mech. Methods* 29, 445–456.
- Lind, L., Araujo, J.A., Barchowsky, A., Belcher, S., Berridge, B.R., Chiamvimonvat, N., Chiu, W.A., Coglian, V.J., Elmore, S., Farraj, A.K., Gomes, A.V., McHale, C.M., Meyer-Tamaki, K.B., Posnack, N.G., Vargas, H.M., Yang, X., Zeise, L., Zhou, C., Smith, M.T., 2021. Key Characteristics of Cardiovascular Toxicants. *Environ. Health Perspect.* 129, 095001.
- Liu, Q., 2021. Effects of Environmental Endocrine-Disrupting Chemicals on Female Reproductive Health. In: Zhang, H., Yan, J. (Eds.), *Environment and Female Reproductive Health*. Springer Singapore, Singapore.
- Liu, P., Yang, F., Wang, Y., Yuan, Z., 2018. Perfluorooctanoic acid (PFOA) exposure in early life increases risk of childhood adiposity: a meta-analysis of prospective cohort studies. *Int. J. Environ. Res. Public Health* 15.
- Liu, W., Zhang, J., Liang, X., Wang, Y., Liu, R., Zhang, R., Zha, J., Martyniuk, C.J., 2022. Environmental concentrations of 2, 4-DTBP cause immunotoxicity in zebrafish (*Danio rerio*) and may elicit ecological risk to wildlife. *Chemosphere* 308, 136465.
- Lizarraga, L.E., Suter, G.W., Lambert, J.C., Patlewicz, G., Zhao, J.Q., Dean, J.L., Kaiser, P., 2021. Advancing the science of a read-across framework for evaluation of data-poor chemicals incorporating systematic and new approach methods. *Regul. Toxicol. Pharm.* 137, 105293.
- Loganathan, N., Salehi, A., Chalmers, J.A., Belsham, D.D., 2018. Bisphenol A Alters Bmal1, Per2, and Rev-Erba mRNA and Requires Bmal1 to Increase Neuropeptide Y Expression in Hypothalamic Neurons. *Endocrinology* 160, 181–192.
- Luderer, U., Eskenazi, B., Hauser, R., Korach, K.S., McHale, C.M., Moran, F., Rieswijk, L., Solomon, G., Udagawa, O., Zhang, L., Zlatnik, M., Zeise, L., Smith, M.T., 2019. Proposed Key Characteristics of Female Reproductive Toxicants as an Approach for Organizing and Evaluating Mechanistic Data in Hazard Assessment. *Environ. Health Perspect.* 127, 075001.
- Luebke, R., 2012. Immunotoxicant screening and prioritization in the twenty-first century. *Toxicol. Pathol.* 40, 294–299.
- Madia, F., Worth, A., Whelan, M., Corvi, R., 2019. Carcinogenicity assessment: Addressing the challenges of cancer and chemicals in the environment. *Environ. Int.* 128, 417–429.
- Maffini, M.V., Soto, A.M., Calabro, J.M., Ucci, A.A., Sonnenschein, C., 2004. The stroma as a crucial target in rat mammary gland carcinogenesis. *J. Cell Sci.* 117, 1495–1502.
- Maffini, M.V., Alger, H.M., Olson, E.D., Neltner, T.G., 2013. Looking back to look forward: a review of FDA's food additives safety assessment. *Compr. Rev. Food Sci. Food Saf.* 12.
- Maffini, M.V., Trasande, L., Neltner, T.G., 2016. Perchlorate and diet: human exposures, risks, and mitigation strategies. *Curr. Environ. Health Rep.* 3, 107–117.
- Mahalingam, S., Ther, L., Gao, L., Wang, W., Ziv-Gal, A., Flaws, J.A., 2017. The effects of in utero bisphenol A exposure on ovarian follicle numbers and steroidogenesis in the F1 and F2 generations of mice. *Reprod. Toxicol.* 74, 150–157.
- Manikkam, M., Tracey, R., Guerrero-Bosagna, C., Skinner, M.K., 2013. Plastics derived endocrine disruptors (BPA, DEHP and DBP) induce epigenetic transgenerational inheritance of obesity, reproductive disease and sperm epimutations. *PLoS One* 8, e55387.
- Maoz, B.M., 2021. Brain-on-a-Chip: Characterizing the next generation of advanced in vitro platforms for modeling the central nervous system. *APL Bioeng.* 5, 030902.
- Mariana, M., Feiteiro, J., Verde, I., Cairrao, E., 2016. The effects of phthalates in the cardiovascular and reproductive systems: a review. *Environ. Int.* 94, 758–776.
- Marroqui, L., Martínez-Pinna, J., Castellano-Muñoz, M., Dos Santos, R.S., Medina-Gali, R.M., Soriano, S., Quesada, I., Gustafsson, J.A., Encinar, J.A., Nadal, A., 2021. Bisphenol-S and Bisphenol-F alter mouse pancreatic  $\beta$ -cell ion channel expression and activity and insulin release through an estrogen receptor ER $\beta$  mediated pathway. *Chemosphere* 265, 129051.
- Martínez-Ibarra, A., Martínez-Razo, L.D., MacDonald-Ramos, K., Morales-Pacheco, M., Vázquez-Martínez, E.R., López-López, M., Rodríguez Dorantes, M., Cerbón, M., 2021. Multisystemic alterations in humans induced by bisphenol A and phthalates: Experimental, epidemiological and clinical studies reveal the need to change health policies. *Environ. Pollut.* 271, 116380.
- Martínez-Pinna, J., Marroqui, L., Hmadcha, A., Lopez-Beas, J., Soriano, S., Villar-Pazos, S., Alonso-Magdalena, P., Dos Santos, R.S., Quesada, I., Martín, F., Soria, B., Gustafsson, J.A., Nadal, A., 2019. Oestrogen receptor  $\beta$  mediates the actions of bisphenol-A on ion channel expression in mouse pancreatic beta cells. *Diabetologia* 62, 1667–1680.

- Marty, S., Beekhuijzen, M., Charlton, A., Hallmark, N., Hannas, B.R., Jacobi, S., Melching-Kollmuss, S., Sauer, U.G., Sheets, L.P., Strauss, V., Urbisch, D., Botham, P. A., van Ravenzwaay, B., 2021. Towards a science-based testing strategy to identify maternal thyroid hormone imbalance and neurodevelopmental effects in the progeny – part II: how can key events of relevant adverse outcome pathways be addressed in toxicological assessments? *Crit. Rev. Toxicol.* 51, 328–358.
- Marvel, S.W., To, K., Grimm, F.A., Wright, F.A., Rusyn, I., Reif, D.M., 2018. ToxPi Graphical User Interface 2.0: Dynamic exploration, visualization, and sharing of integrated data models. *BMC Bioinf.* 19, 80.
- McDonough, C.M., Hannah Shibo, X.u., Guo, T.L., 2021. Toxicity of bisphenol analogues on the reproductive, nervous, and immune systems, and their relationships to gut microbiome and metabolism: insights from a multi-species comparison. *Crit. Rev. Toxicol.* 51, 283–300.
- Melnick, R.L., 2001. Is peroxisome proliferation an obligatory precursor step in the carcinogenicity of di(2-ethylhexyl)phthalate (DEHP)? *Environ. Health Perspect.* 109, 437–442.
- Merrill, L., Michele, A., Vandenberg, L.N., Smith, M.T., Goodson, W., Browne, P., Patisaul, H.B., Guyton, K.Z., Kortenkamp, A., Cogliano, V.J., Woodruff, T.J., Rieswijk, L., Sone, H., Korach, K.S., Gore, A.C., Zeise, L., Thomas Zoeller, R., 2020. Consensus on the key characteristics of endocrine-disrupting chemicals as a basis for hazard identification. *Nat. Rev. Endocrinol.* 16, 45–57.
- Messlerian, C., Souter, I., Gaskins, A.J., Williams, P.L., Ford, J.B., Chiu, Y.H., Calafat, A. M., Hauser, R., 2016. Urinary phthalate metabolites and ovarian reserve among women seeking infertility care. *Hum. Reprod.* 31, 75–83.
- Midya, V., Colicino, E., Conti, D.V., Berhane, K., Garcia, E., Stratakis, N., Andrusaityte, S., Basagaña, X., Casas, M., Fossati, S., Gražulevičienė, R., Haug, L.S., Heude, B., Maitre, R., McEachan, R., Papadopoulou, E., Roumeliotaki, T., Philippat, C., Thomsen, C., Qurquiza, J., Vafeiadi, M., Varo, N., Vos, M.B., Wright, J., McConnell, R., Vrijheid, M., Chatzi, L., Valvi, D., 2022. Association of prenatal exposure to endocrine-disrupting chemicals with liver injury in children. *JAMA Netw. Open* 5, e2220176–e2220276.
- Minet, L., Wang, Z., Shalin, A., Bruton, T.A., Blum, A., Peaslee, G.F., Schwartz-Narbonne, H., Venier, M., Whitehead, H., Yan, W.u., Diamond, M.L., 2022. Use and release of per- and polyfluoroalkyl substances (PFASs) in consumer food packaging in U.S. and Canada. *Environ. Sci. Processes Impacts* 24, 2032–2042.
- Mohajer, N., Du, C.Y., Checkinco, C., Blumberg, B., 2021. Obesogens: How They Are Identified and Molecular Mechanisms Underlying Their Action. *Front. Endocrinol. (Lausanne)* 12, 780888.
- Mohanto, N.C., Ito, Y., Kato, S., Kamijima, M., 2021. Life-Time Environmental Chemical Exposure and Obesity: Review of Epidemiological Studies Using Human Biomonitoring Methods. *Front. Endocrinol.* 12, 100.
- Moon, S., Sung Hoon, Y.u., Lee, C.B., Park, Y.J., Yoo, H.J., Kim, D.S., 2021. Effects of bisphenol A on cardiovascular disease: An epidemiological study using National Health and Nutrition Examination Survey 2003–2016 and meta-analysis. *Sci. Total Environ.* 763, 142941.
- Moore, S., Paalanen, L., Melymuk, L., Katsonouri, A., Kolossa-Gehring, M., Tolonen, H., 2022. The Association between ADHD and Environmental Chemicals — A Scoping Review. *Int. J. Environ. Res. Public Health* 19, 2849.
- Muncke, J., 2009. Exposure to endocrine disrupting compounds via the food chain: Is packaging a relevant source? *Sci. Total Environ.* 407, 4549–4559.
- Muncke, J., 2021. Tackling the toxics in plastics packaging. *PLoS Biol.* 19, e3000961.
- Muncke, J., Backhaus, T., Geueke, B., Maffini, M.V., Martin, O.V., Myers, J.P., Soto, A.M., Trasande, L., Trier, X., Scheringer, M., 2017. Scientific challenges in the risk assessment of food contact materials. *Environ. Health Perspect.* 125, 095001.
- Muncke, J., Andersson, A.-M., Backhaus, T., Boucher, J.M., Almroth, B.C., Castillo, A.C., Chevrier, J., Demeneix, B.A., Emmanuel, J.A., Fini, J.-B., Gee, D., Geueke, B., Groh, K., Heindel, J.J., Houlihan, J., Kossatos, C.D., Kwiatkowski, C.F., Lefferts, L.Y., Maffini, M.V., Martin, O.V., Myers, J.P., Nadal, A., Nerin, C., Pelch, K.E., Fernández, S.R., Sargis, R.M., Soto, A.M., Trasande, L., Vandenberg, L.N., Wagner, M., Changqing, W.u., Thomas Zoeller, R., Scheringer, M., 2020. Impacts of food contact chemicals on human health: a consensus statement. *Environ. Health* 19, 25.
- Mustieles, V., Fernández, M.F., 2020. Bisphenol A shapes children's brain and behavior: towards an integrated neurotoxicity assessment including human data. *Environ. Health* 19, 66.
- Naderi, M., Kwong, R.W.M., 2020. A comprehensive review of the neurobehavioral effects of bisphenol S and the mechanisms of action: New insights from in vitro and in vivo models. *Environ. Int.* 145, 106078.
- Naidenko, O.V., Andrews, D.Q., Temkin, A.M., Stoiber, T., Uche, U.I., Evans, S., Perrone-Gray, S., 2021. Investigating Molecular Mechanisms of Immunotoxicity and the Utility of ToxCast for Immunotoxicity Screening of Chemicals Added to Food. *Int. J. Environ. Res. Public Health* 18, 3332.
- Naxerova, K., 2021. Mutation fingerprints encode cellular histories. *Nature* 597, 334–36.
- Neale, P.A., Escher, B.I., de Baat, M.L., Enault, J., Leusch, F.D.L., 2023. Effect-Based Trigger Values Are Essential for the Uptake of Effect-Based Methods in Water Safety Planning. *Environ. Toxicol. Chem.* 42, 714–726.
- Neltner, T.G., Alger, H.M., Leonard, J.E., Maffini, M.V., 2013a. Data Gaps in Toxicity Testing of Chemicals Allowed in Food in the United States. *Reprod. Toxicol.*
- Neltner, T.G., Alger, H.M., Leonard, J.E., Maffini, M.V., 2013b. Data gaps in toxicity testing of chemicals allowed in food in the United States. *Reprod. Toxicol.* 42, 85–94.
- Nerin, C., Alfaro, P., Aznar, M., Domeño, C., 2013. The challenge of identifying non-intentionally added substances from food packaging materials: A review. *Anal. Chim. Acta* 775, 14–24.
- Nerin, C., Asensio, E., 2007. Migration of organic compounds from a multilayer plastic-paper material intended for food packaging. *Anal. Bioanal. Chem.* 389, 589–596.
- Nerín, C., Bourdoux, S., Faust, B., Gude, T., Lesueur, C., Simat, T., Stoermer, A., Van Hoek, E., Oldring, P., 2022. Guidance in selecting analytical techniques for identification and quantification of non-intentionally added substances (NIAS) in food contact materials (FCMS). *Food Addit. Contaminants: Part A* 39, 620–643.
- Nerín, C., Ubeda, J.L., Alfaro, P., Dahmani, Y., Aznar, M., Canellas, E., Aulsejo, R., 2014. Compounds from multilayer plastic bags cause reproductive failures in artificial insemination. *Sci. Rep.* 4.
- Nerín, C., Canellas, E., Vera, P., Garcia-Calvo, E., Luque-Garcia, J.L., Cámara, C., Aulsejo, R., Miguel, J., Mendoza, N., 2018. A common surfactant used in food packaging found to be toxic for reproduction in mammals. *Food Chem. Toxicol.* 113, 115–124.
- Nerín, C., Su, Q.Z., Vera, P., Mendoza, N., Aulsejo, R., 2020. Influence of nonylphenol from multilayer plastic films on artificial insemination of sows. *Anal. Bioanal. Chem.* 412, 6519–6528.
- Nicholson, L.B., 2016. The immune system. *Essays Biochem.* 60, 275–301.
- Nowak, K., Jabłońska, E., Ratajczak-Wrona, W., 2019. Immunomodulatory effects of synthetic endocrine disrupting chemicals on the development and functions of human immune cells. *Environ. Int.* 125, 350–364.
- NTP, 2018. Report on Carcinogens. Monograph on Antimony Trioxide. In: RoC Monograph, edited by National Toxicology Program. U.S. Department of Health and Human Services.
- NTP, 2021. RoC Review of Antimony Trioxide. U.S. Department of Health and Human Services, Accessed 23 Dec 2022. <https://ntp.niehs.nih.gov/whatwestudy/assessment/s/cancer/completed/antimony/index.html>.
- Obsekov, V., Kahn, L.G., Trasande, L., 2022a. Leveraging Systematic Reviews to Explore Disease Burden and Costs of Per- and Polyfluoroalkyl Substance Exposures in the United States. *Exposure and Health.*
- OECD, 2015. Guidance Document on Revisions to OECD Genetic Toxicology Test Guidelines. Edited by Organisation for Economic Co-operation and Development.
- OECD, 2022. OECD Test Guidelines for Chemicals. Accessed 23 Dec 2022. <https://www.oecd.org/chemicalsafety/testing/oecdguidelinesforthetestingofchemicals.htm>.
- Ogungbesan, A., Neal-Kluever, A., Rice, P., 2019. Exploring the use of current immunological assays for the developmental immunotoxicity assessment of food contact materials. *Food Chem. Toxicol.* 133, 110801.
- Oldring, P.K.T., Castle, L., O'Mahony, C., Dixon, J., 2014. Estimates of dietary exposure to bisphenol A (BPA) from light metal packaging using food consumption and packaging usage data: a refined deterministic approach and a fully probabilistic (FACET) approach. *Food Addit. Contaminants: Part A* 31, 466–489.
- Oldring, P., Faust, B., Gude, T., Lesueur, C., Simat, T., Stoermer, A., Van Hoek, E., Nerin, C., 2023. An Overview of Approaches for Analysing NIAS from different FCMS. *ILSI Europe. Zenodo: ILSI Europe.*
- Omer, E., Bichon, E., Hutinet, S., Royer, A.-L., Monteau, F., Germon, H., Hill, P., Rемаud, G., Dervilly-Pinel, G.u., Cariou, R., Le Bizec, B., 2019. Toward the characterisation of non-intentionally added substances migrating from polyester-polyurethane lacquers by comprehensive gas chromatography-mass spectrometry technologies. *J. Chromatogr. A* 1601, 327–334.
- Pant, J., Ranjan, P., Deshpande, S.B., 2011. Bisphenol A decreases atrial contractility involving NO-dependent G-cyclase signaling pathway. *J. Appl. Toxicol.* 31, 698–702.
- Park, S.E., Ahn, J., Jeong, H.-E., Youn, I., Huh, D., Chung, S., 2021. A three-dimensional in vitro model of the peripheral nervous system. *NPJ Asia Mater.* 13, 2.
- Park, S., Lee, J.M., Kim, J.W., Cheong, J.H., Yun, H.J., Hong, Y.C., Kim, Y., Han, D.H., Yoo, H.J., Shin, M.S., Cho, S.C., Kim, B.N., 2015. Association between phthalates and externalizing behaviors and cortical thickness in children with attention deficit hyperactivity disorder. *Psychol. Med.* 45, 1601–1612.
- Parg, C., Roy, N.M., Ton, C., Lin, Y., McGrath, P., 2007. Neurotoxicity assessment using zebrafish. *J. Pharmacol. Toxicol. Methods* 55, 103–112.
- Pérez-Bermejo, M., Mas-Pérez, I., Murillo-Llorente, M.T., 2021. The role of the bisphenol A in diabetes and obesity. *Biomedicines* 9.
- Piekarski, D.J., Diaz, K.R., McNeerney, M.W., 2020. Perfluoroalkyl chemicals in neurological health and disease: Human concerns and animal models. *Neurotoxicology* 77, 155–168.
- Pieke, E.N., Granby, K., Trier, X., Smedsgaard, J., 2017. A framework to estimate concentrations of potentially unknown substances by semi-quantification in liquid chromatography electrospray ionization mass spectrometry. *Anal. Chim. Acta* 975, 30–41.
- Pierozan, P., Jermeren, F., Karlsson, O., 2018. Perfluorooctanoic acid (PFOA) exposure promotes proliferation, migration and invasion potential in human breast epithelial cells. *Arch. Toxicol.* 92, 1729–1739.
- Pillai, H.K., Fang, M., Beglov, D., Kozakov, D., Vajda, S., Stapleton, H.M., Webster, T.F., Schlezinger, J.J., 2014. Ligand binding and activation of PPAR $\gamma$  by Firemaster® 550: effects on adipogenesis and osteogenesis <i>in vitro</i>. *Environ. Health Perspect.* 122, 1225–1232.
- Pinter, E., Rainer, B., Czerny, T., Riegel, E., Schilter, B., Marin-Kuan, M., Tacker, M., 2020. Evaluation of the suitability of mammalian in vitro assays to assess the genotoxic potential of food contact materials. *Foods*, 9.
- Pinto, C.L., Markey, K., Dix, D., Browne, P., 2018. Identification of candidate reference chemicals for in vitro steroidogenesis assays. *Toxicol. In Vitro* 47, 103–119.
- Poças, M.F.F., Oliveira, J.C., Pinto, H.J., Zacarias, M.E., Hogg, T., 2009. Characterization of patterns of food packaging usage in Portuguese homes. *Food Addit. Contam. Part A: Chem. Anal. Control Expo. Risk Assess.* 26, 1314–1324.
- Pouech, C., Kiss, A., Lafay, F., Léonard, D., Wiest, L., Cren-Olivé, C., Vulliet, E., 2015. Human exposure assessment to a large set of polymer additives through the analysis of urine by solid phase extraction followed by ultra high performance liquid chromatography coupled to tandem mass spectrometry. *J. Chromatogr. A* 1423, 111–123.



- Predieri, B., Bruzzi, P., Bigi, E., Ciancia, S., Madeo, S.F., Lucaccioni, L., Iughetti, L., 2020. Endocrine disrupting chemicals and type 1 diabetes. *Int. J. Mol. Sci.* 21, 2937.
- Prins, G.S., Hu, W.Y., Shi, G.B., Hu, D.P., Majumdar, S., Li, G., Huang, K., Nelles, J.L., Ho, S.M., Walker, C.L., Kajdacsy-Balla, A., van Breemen, R.B., 2014. Bisphenol A promotes human prostate stem-progenitor cell self-renewal and increases in vivo carcinogenesis in human prostate epithelium. *Endocrinology* 155, 805–817.
- Qian, S., Ji, H., XiaoXiao, W., Li, N., Yang, Y., Jiangtao, B., Zhang, X., Qiao, L., Henglin, Y., Ning, X., Zhang, C., 2018. Detection and quantification analysis of chemical migrants in plastic food contact products. *PLoS One* 13, e0208467.
- Qin, W.-P., Cao, L.-Y., Li, C.-H., Guo, L.-H., Colbourne, J., Ren, X.-M., 2020. Perfluoroalkyl substances stimulate insulin secretion by islet  $\beta$  cells via G protein-coupled receptor 40. *Environ. Sci. Technol.* 54, 3428–3436.
- Radke, E.G., Braun, J.M., Meeker, J.D., Cooper, G.S., 2018. Phthalate exposure and male reproductive outcomes: a systematic review of the human epidemiological evidence. *Environ. Int.* 121, 764–793.
- Radke, E.G., Braun, J.M., Nachman, R.M., Cooper, G.S., 2020. Phthalate exposure and neurodevelopment: a systematic review and meta-analysis of human epidemiological evidence. *Environ. Int.* 137, 105408.
- Rajkumar, A., Luu, T., Beal, M.A., Barton-Maclaren, T.S., Hales, B.F., Robaire, B., 2022. Phthalates and alternative plasticizers differentially affect phenotypic parameters in gonadal somatic and germ cell lines. *Biol. Reprod.* 106, 613–627.
- Ramadan, M., Cooper, B., Posnack, N.G., 2020. Bisphenols and phthalates: plastic chemical exposures can contribute to adverse cardiovascular health outcomes. *Birth Defects Res.* 112, 1362–1385.
- Rancière, F., Lyons, J.G., Loh, V.H.Y., Botton, J., Galloway, T., Wang, T., Shaw, J.E., Magliano, D.J., 2015. Bisphenol A and the risk of cardiometabolic disorders: a systematic review with meta-analysis of the epidemiological evidence. *Environ. Health* 14, 1–23.
- Rebollo-Solheiro, D., Castillo Flores, L.Y., Solheiro-Villavicencio, H., 2021. Impact of BPA on behavior, neurodevelopment and neurodegeneration. *FBL*, 26, 363–400.
- Reicha, Y., Truong, L., Leong, C., Cao, D., Field, J.A., Tanguay, R.L., 2022. Dietary Perfluorohexanoic Acid (PFHxA) Exposures in Juvenile Zebrafish Produce Subtle Behavioral Effects across Generations. *Toxics* 10, 372.
- Ribeiro, C.M., Beserra, B.T.S., Silva, N.G., Lima, C.L., Rocha, P.R.S., Coelho, M.S., Neves, F.A.R., Amato, A.A., 2020. Exposure to endocrine-disrupting chemicals and anthropometric measures of obesity: a systematic review and meta-analysis. *BMJ Open* 10, e033509.
- Richard, A.M., Judson, R.S., Houck, K.A., Grulke, C.M., Volarath, P., Thillainadarajah, I., Yang, C., Rathman, J., Martin, M.T., Wambaugh, J.F., Knudsen, T.B., Kancherla, J., Mansouri, K., Patlewicz, G., Williams, A.J., Little, S.B., Crofton, K.M., Thomas, R.S., 2016. ToxCast Chemical Landscape: Paving the Road to 21st Century Toxicology. *Chem. Res. Toxicol.* 29, 1225–1251.
- Rider, C.V., McHale, C.M., Webster, T.F., Lowe, L., Goodson, W.H., La Merrill, M.A., Rice, G., Zeise, L., Zhang, L., Smith, M.T., 2021. Using the Key Characteristics of Carcinogens to Develop Research on Chemical Mixtures and Cancer. *Environ. Health Perspect.* 129, 035003.
- Robitaille, J., Denslow, N.D., Escher, B.I., Kurita-Oyamada, H.G., Marlatt, V., Martyniuk, C.J., Navarro-Martín, L., Prosser, R., Sanderson, T., Yargeau, V., Langlois, V.S., 2022. Towards regulation of Endocrine Disrupting chemicals (EDCs) in water resources using bioassays - A guide to developing a testing strategy. *Environ. Res.* 205, 112483.
- Rojas-Rueda, D., Morales-Zamora, E., Alsufyani, W.A., Herbst, C.H., AlBalawi, S.M., Alskait, R., Alomran, M., 2021. Environmental Risk Factors and Health: An Umbrella Review of Meta-Analyses. *Int. J. Environ. Res. Public Health* 18, 704.
- Rønnov-Jessen, L., Bissell, M.J., 2009. Breast cancer by proxy: can the microenvironment be both the cause and consequence? *Trends Mol. Med.* 15, 5–13.
- Roser, M., Ritchie, H., Spooner, F., 2021. Burden of disease. [OurWorldInData.org](https://ourworldindata.org/burden-of-disease). Accessed 23 December 2022. <https://ourworldindata.org/burden-of-disease>.
- Rotenberg Iu, S., Mazaev, V.T., Shlepina, T.G., 1978. Peculiarities of alkyl tin effects on respiration and oxidative phosphorylation of rat liver mitochondria. *Ukr Biokhim Zh.* 50, 695–700.
- Ruan, Y., Lalwani, D., Kwok, K.Y., Yamazaki, E., Taniyasu, S., Kumar, N.J.I., Lam, P.K.S., Yamashita, N., 2019. Assessing exposure to legacy and emerging per- and polyfluoroalkyl substances via hair – The first nationwide survey in India. *Chemosphere* 229, 366–373.
- Rudel, R.A., Gray, J.M., Engel, C.L., Rawsthorne, T.W., Dodson, R.E., Ackerman, J.M., Rizzo, J., Nudelman, J.L., Brody, J.G., 2011. Food Packaging and Bisphenol A and Bis(2-Ethylhexyl) Phthalate Exposure: Findings from a Dietary Intervention. *Environ. Health Perspect.* 119, 914–920.
- Rusyn, I., Arzuaga, X., Cattley, R.C., Corton, J.C., Ferguson, S.S., Godoy, P., Guyton, K.Z., Kaplowitz, N., Khetani, S.R., Roberts, R.A., Roth, R.A., Smith, M.T., 2021. Key Characteristics of Human Hepatotoxicants as a Basis for Identification and Characterization of the Causes of Liver Toxicity. *Hepatology* 74, 3486–3496.
- Ruszkiewicz, J.A., Pinkas, A., Miah, M.R., Weitz, R.L., Lawes, M.J.A., Akinyemi, A.J., Ijomone, O.M., Aschner, M., 2018. *C. elegans* as a model in developmental neurotoxicology. *Toxicol. Appl. Pharmacol.* 354, 126–135.
- Sachana, M., Willett, C., Pistollato, F., Bal-Price, A., 2021. The potential of mechanistic information organised within the AOP framework to increase regulatory uptake of the developmental neurotoxicity (DNT) in vitro battery of assays. *Reprod. Toxicol.* 103, 159–170.
- Sanchis, Y., Yusà, V., Coscollà, C., 2017. Analytical strategies for organic food packaging contaminants. *J. Chromatogr. A* 1490, 22–46.
- Sang, C., Song, Y., Jin, T.W., Zhang, S., Fu, L., Zhao, Y., Zou, X., Wang, Z., Gao, H., Liu, S., 2021. Bisphenol A induces ovarian cancer cell proliferation and metastasis through estrogen receptor- $\alpha$  pathways. *Environ. Sci. Pollut. Res. Int.* 28, 36060–36068.
- Sant, K.E., Jacobs, H.M., Borofski, K.A., Moss, J.B., Timme-Laragy, A.R., 2017. Embryonic exposures to perfluorooctanesulfonic acid (PFOS) disrupt pancreatic organogenesis in the zebrafish, *Danio rerio*. *Environ. Pollut.* 220, 807–817.
- Santos, D., Sousa, R., Medina-Gali, R.M., Babiloni-Chust, I., Marroqui, L., Nadal, A., 2022. In Vitro Assays to Identify Metabolism-Disrupting Chemicals with Diabetogenic Activity in a Human Pancreatic  $\beta$ -Cell Model. *Int. J. Mol. Sci.* 23, 5040.
- Sapozhnikova, Y., Nuñez, A., Johnston, J., 2021. Screening of chemicals migrating from plastic food contact materials for oven and microwave applications by liquid and gas chromatography - Orbitrap mass spectrometry. *J. Chromatogr. A* 1651, 462261.
- Scholz, S., Brack, W., Escher, B.I., Hackermüller, J., Liess, M., von Bergen, M., Wick, L.Y., Zenclussen, A.C., Altenburger, R., 2022. The EU chemicals strategy for sustainability: an opportunity to develop new approaches for hazard and risk assessment. *Arch. Toxicol.* 96, 2381–2386.
- Schug, T.T., Abagyan, R., Blumberg, B., Collins, T.J., Crews, D., DeFur, P.L., Dickerson, S.M., Edwards, T.M., Gore, A.C., Guillette, L.J., Hayes, T., Heindel, J.J., Moores, A., Patisaul, H.B., Tal, T.L., Thayer, K.A., Vandenberg, L.N., Warner, J.C., Watson, C.S., vom Saal, F.S., Zoeller, R.T., O'Brien, K.P., Myers, J.P., 2013. Designing endocrine disruption out of the next generation of chemicals. *Green Chem.*
- Seo, Y., Shin, T.H., Kim, H.S., 2019. Current strategies to enhance adipose stem cell function: an update. *Int. J. Mol. Sci.* 20.
- Serras, A.S., Rodrigues, J.S., Cipriano, M., Rodrigues, A.V., Oliveira, N.G., Miranda, J.P., 2021. A critical perspective on 3D liver models for drug metabolism and toxicology studies. *Front. Cell Dev. Biol.* 9.
- Severin, I., Soutou, E., Dahbi, L., Chagnon, M.C., 2017. Use of bioassays to assess hazard of food contact material extracts: State of the art. *Food Chem. Toxicol.* 105, 429–447.
- Shaffer, R.M., Ferguson, K.K., Sheppard, L., James-Todd, T., Butts, S., Chandrasekaran, S., Swan, S.H., Barrett, E.S., Nguyen, R., Bush, N., McElrath, T.F., Sathyanarayana, S., 2019. Maternal urinary phthalate metabolites in relation to gestational diabetes and glucose intolerance during pregnancy. *Environ. Int.* 123, 588–596.
- Sharma, A., Mollier, J., Brocklesby, R.W.K., Caves, C., Jayasena, C.N., Minhas, S., 2020. Endocrine-disrupting chemicals and male reproductive health. *Reprod. Med. Biol.* 19, 243–253.
- Shoucri, B.M., Hung, V.T., Chamorro-García, R., Shioda, T., Blumberg, B., 2018. Retinoid X Receptor Activation During Adipogenesis of Female Mesenchymal Stem Cells Programs a Dysfunctional Adipocyte. *Endocrinology* 159, 2863–2883.
- Silva, M.J., Slakman, A.R., Reidy, J.A., Preau Jr., J.L., Herbert, A.R., Samandar, E., Needham, L.L., Calafat, A.M., 2004. Analysis of human urine for fifteen phthalate metabolites using automated solid-phase extraction. *J. Chromatogr. B Anal. Technol. Biomed. Life Sci.* 805, 161–167.
- Skakkebaek, N.E., Rajpert-De Meyts, E., Main, K.M., 2001. Testicular dysgenesis syndrome: an increasingly common developmental disorder with environmental aspects. *Opinion. Hum. Reprod.* 16, 972–998.
- Skakkebaek, N.E., Lindahl-Jacobsen, R., Levine, H., Andersson, A.-M., Jørgensen, N., Main, K.M., Lidgaard, Ø., Priskorn, L., Holmboe, S.A., Bråuner, E.V., Almstrup, K., Franca, L.R., Znaor, A., Kortenkamp, A., Hart, R.J., Juul, A., 2022. Environmental factors in declining human fertility. *Nat. Rev. Endocrinol.* 18, 139–157.
- Skakkebaek, N.E., Rajpert-De Meyts, E., Buck Louis, G.M., Toppari, J., Andersson, A.M., Eisenberg, M.L., Jensen, T.K., Jørgensen, N., Swan, S.H., Sapra, K.J., Ziebe, S., Priskorn, L., Juul, A., 2016. Male Reproductive Disorders and Fertility Trends: Influences of Environment and Genetic Susceptibility. *Physiol. Rev.* 96, 55–97.
- Smith, M.T., Guyton, K.Z., Gibbons, C.F., Fritz, J.M., Portier, C.J., Rusyn, I., DeMarini, D.M., Caldwell, J.C., Kavlock, R.J., Lambert, P.F., Hecht, S.S., Bucher, J.R., Stewart, B.W., Baan, R.A., Coglian, V.J., Straif, K., 2016. Key Characteristics of Carcinogens as a Basis for Organizing Data on Mechanisms of Carcinogenesis. *Environ. Health Perspect.* 124, 713–721.
- Soave, I., Occhiali, T., Assorgi, C., Marci, R., Caserta, D., 2020. Environmental toxin exposure in polycystic ovary syndrome women and possible ovarian neoplastic repercussion. *Curr. Med. Res. Opin.* 36, 693–703.
- Sonnenschein, C., Soto, A.M., 2020. Over a century of cancer research: Inconvenient truths and promising leads. *PLoS Biol.* 18, e3000670.
- Soto, A.M., Briskin, C., Schaeberle, C.M., Sonnenschein, C., 2013. Does cancer start in the womb? Altered mammary gland development and predisposition to breast cancer due to in utero exposure to endocrine disruptors. *J. Mammary Gland Biol. Neoplasia* 18, 199–208.
- Soutou, E., Severin, I., Le Hegarat, L., Hogeveen, K., Aljawish, A., Fessard, V., Marie-Christine, C., 2017. Genotoxic effects of food contact recycled paperboard extracts on two human hepatic cell lines. *Food Addit. Contaminants Part A* 1–12.
- Stratakis, N., Conti, V., Jin, R., Margetaki, K., Valvi, D., Siskos, A.P., Maitre, L., Garcia, E., Varo, N., Zhao, Y., Roumeliotaki, T., Vafeiadi, M., Urquiza, J., Fernández-Barrés, S., Heude, B., Basagana, X., Casas, M., Fossati, S., Gražulevičienė, R., Andrušaitytė, S., Uppal, K., McEachan, R.R.C., Papadopoulou, E., Robinson, O., Haug, L.S., Wright, J., Vos, M.B., Keun, H.C., Vrijheid, M., Berhane, K.T., McConnell, R., Chatzi, L., 2020. Prenatal exposure to perfluoroalkyl substances associated with increased susceptibility to liver injury in children. *Hepatology* 72, 1758–1770.
- Street, M.E., Audouze, K., Legler, J., Sone, H., Palanza, P., 2021. Endocrine Disrupting Chemicals: Current Understanding, New Testing Strategies and Future Research Needs. *Int. J. Mol. Sci.* 22, 933.
- Stucki, A.O., Barton-Maclaren, T.S., Bhuller, Y., Henriquez, J.E., Henry, T.R., Hirn, C., Miller-Holt, J., Nagy, E.G., Perron, M.M., Ratzlaff, D.E., Stedeford, T.J., Clippinger, A.J., 2022. Use of new approach methodologies (NAMs) to meet regulatory requirements for the assessment of industrial chemicals and pesticides for effects on human health. *Front. Toxicol.* 4, 964553.

- Susmann, H.P., Schaidler, L.A., Rodgers, K.M., Rudel, R., 2019. Dietary Habits Related to Food Packaging and Population Exposure to PFASs. *Environ. Health Perspect.* 127, 10.
- Svensson, K., Tanner, E., Gennings, C., Lindh, C., Kiviranta, H., Wikström, S., Bornehag, C.-G., 2021. Prenatal exposures to mixtures of endocrine disrupting chemicals and children's weight trajectory up to age 5.5 in the SELMA study. *Sci. Rep.* 11, 11036.
- Symeonides, C., Brunner, M., Mulders, Y., Toshiwal, P., Cantrell, M., Mofflin, L., Dunlop, S., 2021. Buy-now-pay-later: Hazards to human and planetary health from plastics production, use and waste. *J. Paediatr. Child Health* 57, 1795–1804.
- Tang, Q.Q., Otto, T.C., Lane, M.D., 2004. Commitment of C3H10T1/2 pluripotent stem cells to the adipocyte lineage. *PNAS* 101, 9607–9611.
- Tanner, E.M., Hallerback, M.U., Wikström, S., Lindh, C., Kiviranta, H., Gennings, C., Bornehag, C.-G., 2020. Early prenatal exposure to suspected endocrine disruptor mixtures is associated with lower IQ at age seven. *Environ. Int.* 134, 105185.
- Taylor, R.B., Sapozhnikova, Y., 2022. Assessing Chemical Migration from Plastic Food Packaging into Food Simulant by Gas and Liquid Chromatography with High-Resolution Mass Spectrometry. *J. Agric. Food Chem.* 70, 4805–4816.
- Temkin, A.M., Hocevar, B.A., Andrews, D.Q., Naidenko, O.V., Kamendulis, L.M., 2020. Application of the Key Characteristics of Carcinogens to Per and Polyfluoroalkyl Substances. *Int. J. Environ. Res. Public Health* 17, 1668.
- Thurston, S.W., Mendiola, J., Bellamy, A.R., Levine, H., Wang, C., Sparks, A., Redmon, J. B., Drobnis, E.Z., Swan, S.H., 2016. Phthalate exposure and semen quality in fertile US men. *Andrology* 4, 632–638.
- Tice, R.R., Austin, C.P., Kavlock, R.J., Bucher, J.R., 2013. Improving the human hazard characterization of chemicals: a Tox21 update. *Environ. Health Perspect.* 121, 756–765.
- Tingaud-Sequeira, A., Ouadah, N., Babin, P.J., 2011. Zebrafish obesogenic test: a tool for screening molecules that target adiposity. *J. Lipid Res.* 52, 1765–1772.
- Tisler, S., Christensen, J.H., 2022. Non-target screening for the identification of migrating compounds from reusable plastic bottles into drinking water. *J. Hazard. Mater.* 429, 128331.
- Tolosa, L., Jiménez, N., Pérez, G., Castell, J.V., José Gómez-Lechón, M., Teresa Donato, M., 2018. Customised in vitro model to detect human metabolism-dependent idiosyncratic drug-induced liver injury. *Arch. Toxicol.* 92, 383–399.
- Trasande, L., Zoeller, R.T., Hass, U., Kortenkamp, A., Grandjean, P., Myers, J.P., DiGangi, J., Bellanger, M., Hauser, R., Legler, J., Skakkebaek, N.E., Heindel, J.J., 2015. Estimating burden and disease costs of exposure to endocrine-disrupting chemicals in the European union. *J. Clin. Endocrinol. Metab.* 100, 1245–1255.
- Trasande, L., Zoeller, R.T., Hass, U., Kortenkamp, A., Grandjean, P., Myers, J.P., DiGangi, J., Hunt, P.M., Rudel, R., Sathyanarayana, S., Bellanger, M., Hauser, R., Legler, J., Skakkebaek, N.E., Heindel, J.J., 2016. Burden of disease and costs of exposure to endocrine disrupting chemicals in the European Union: an updated analysis. *Andrology* 4, 565–572.
- Trier, X., Granby, K., Christensen, J., 2011. Polyfluorinated surfactants (PFS) in paper and board coatings for food packaging. *Environ. Sci. Pollut. Res.* 1–13.
- Tsochatzis, E.D., Lopes, J.A., Gika, H., Dalsgaard, T.K., Theodoridis, G., 2021. A fast SALLE GC-MS/MS multi-analyte method for the determination of 75 food packaging substances in food simulants. *Food Chem.* 361, 129998.
- Turner, P.A., Gurumurthy, B., Bailey, J.L., Elks, C.M., Janorkar, A.V., 2017. Adipogenic Differentiation of Human Adipose-Derived Stem Cells Grown as Spheroids. *Process Biochem.* 59, 312–320.
- Van Bossuyt, M., Van Hoeck, E., Vanhaecke, T., Rogiers, V., Mertens, B., 2016. Printed paper and board food contact materials as a potential source of food contamination. *Regul. Toxicol. Pharm.* 81, 10–19.
- Van Bossuyt, M., Van Hoeck, E., Vanhaecke, T., Rogiers, V., Mertens, B., 2017. Safeguarding human health using in silico tools? *Arch. Toxicol.* 91, 2705–2706.
- Van Bossuyt, M., Van Hoeck, E., Vanhaecke, T., Rogiers, V., Mertens, B., 2019. Prioritizing Substances of Genotoxic Concern for In-Depth Safety Evaluation Using Non-Animal Approaches: The Example of Food Contact Materials. *Altox-Alternat. Animal Exp.* 36, 215–230.
- van den Dries, M.A., Guxens, M., Spaan, S., Ferguson, K.K., Philips, E., Santos, S., Jaddoe, V.W.V., Ghassabian, A., Trasande, L., Tiemeier, H., Pronk, A., 2020. Phthalate and Bisphenol Exposure during Pregnancy and Offspring Nonverbal IQ. *Environ. Health Perspect.* 128, 77009.
- Vessa, B., Perlman, B., McGovern, P.G., Morelli, S.S., 2022. Endocrine disruptors and female fertility: a review of pesticide and plasticizer effects. *F S Rep.* 3, 86–90.
- Villar-Pazos, S., Martínez-Pinna, J., Castellano-Muñoz, M., Alonso-Magdalena, P., Marroqui, L., Quesada, I., Gustafsson, J.A., Nadal, A., 2017. Molecular mechanisms involved in the non-monotonic effect of bisphenol-a on ca2+ entry in mouse pancreatic  $\beta$ -cells. *Sci. Rep.* 7, 11770.
- Völker, J., Ashcroft, F., Vedøy, Å., Zimmermann, L., Wagner, M., 2022. Adipogenic Activity of Chemicals Used in Plastic Consumer Products. *Environ. Sci. Technol.* 56, 2487–2496.
- Walker, V.R., Boyles, A.L., Pelch, K.E., Holmgren, S.D., Shapiro, A.J., Blystone, C.R., Devito, M.J., Newbold, R.R., Blain, R., Hartman, P., Thayer, K.A., Rooney, A.A., 2018. Human and animal evidence of potential transgenerational inheritance of health effects: An evidence map and state-of-the-science evaluation. *Environ. Int.* 115, 48–69.
- Wan, M.L.Y., Co, V.A., El-Nezami, H., 2022. Endocrine disrupting chemicals and breast cancer: a systematic review of epidemiological studies. *Crit. Rev. Food Sci. Nutr.* 62, 6549–6576.
- Wang, Z., Alderman, M.H., Asgari, C., Taylor, H.S., 2020. Fetal Bisphenol-A Induced Changes in Murine Behavior and Brain Gene Expression Persisted in Adult-aged Offspring. *Endocrinology* 161.
- Wang, L.J., Chen, C.-Y., Bornehag, C.-G., 2016. Bisphenol A exposure may increase the risk of development of atopic disorders in children. *Int. J. Hyg. Environ. Health* 219, 311–316.
- Wang, W., Hafner, K.S., Flaws, J.A., 2014. In utero bisphenol A exposure disrupts germ cell nest breakdown and reduces fertility with age in the mouse. *Toxicol. Appl. Pharmacol.* 276, 157–164.
- Wang, B., Li, M., Zhao, Z., Lu, J., Chen, Y., Xu, Y., Xu, M., Wang, W., Wang, T., Bi, Y., Ning, G., 2019. Urinary bisphenol A concentration and glucose homeostasis in non-diabetic adults: a repeated-measures, longitudinal study. *Diabetologia* 62, 1591–1600.
- Wang, D., Zhao, H., Fei, X., Synder, S.A., Fang, M., Liu, M., 2021. A comprehensive review on the analytical method, occurrence, transformation and toxicity of a reactive pollutant: BADGE. *Environ. Int.* 155, 106701.
- Wassenaar, P.N.H., Trasande, L., Legler, J., 2017. Systematic Review and Meta-Analysis of Early-Life Exposure to Bisphenol A and Obesity-Related Outcomes in Rodents. *Environ. Health Perspect.* 125, 106001.
- Wehbe, Z., Nasser, S.A., El-Yazbi, A., Nasreddine, S., Eid, A.H., 2020. Estrogen and Bisphenol A in Hypertension. *Curr. Hypertens. Rep.* 22, 23.
- Wen, Z.-J., Wang, Z.-Y., Zhang, Y.-F., 2022. Adverse cardiovascular effects and potential molecular mechanisms of DEHP and its metabolites—a review. *Sci. Total Environ.* 847, 157443.
- Westerhoff, P., Prapaipong, P., Shock, E., Hillaireau, A., 2008. Antimony leaching from polyethylene terephthalate (PET) plastic used for bottled drinking water. *Water Res.* 42, 551–556.
- WHO, 2018. Noncommunicable diseases. Fact sheet. <http://www.who.int/mediacentre/factsheets/fs355/en/>.
- WHO, 2020. Infertility, WHO, Accessed 23 Dec 2022. <https://www.who.int/news-room/fact-sheets/detail/infertility>.
- WHO, 2021. Cardiovascular diseases (CVDs), Accessed 23 Dec 2022. <https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-cvds>.
- Williams, D.E., Orner, G., Willard, K.D., Tilton, S., Hendricks, J.D., Pereira, C., Benninghoff, A.D., Bailey, G.S., 2009. Rainbow Trout (*Oncorhynchus mykiss*) and Ultra-Low Dose Cancer Studies. *Comp. Biochem. Physiol.* 149, 175–181.
- Willis, R.A., 1948. Pathology of tumors. Butterworth & Co Ltd, London, England.
- Wu, W., Li, M., Liu, A., Wu, C., Li, D., Deng, Q., Zhang, B., Du, J., Gao, X., Hong, Y., 2020. Bisphenol A and the Risk of Obesity a Systematic Review With Meta-Analysis of the Epidemiological Evidence. *Dose Response* 18, 1559325820916949.
- Xie, M.-Y., Ni, H., Zhao, D.-S., Wen, L.-Y., Li, K.-S., Yang, H.-H., Wang, S.-S., Zhang, H., Hong, S., 2016. Exposure to bisphenol A and the development of asthma: A systematic review of cohort studies. *Reprod. Toxicol.* 65, 224–229.
- Yan, S., Chen, Y., Dong, M., Song, W., Belcher, S.M., Wang, H.-S., 2011. Bisphenol A and 17 $\beta$ -Estradiol Promote Arrhythmia in the Female Heart via Alteration of Calcium Handling. *PLoS One* 6, e25455.
- Yang, S., Ooka, M., Margolis, R.J., Xia, M., 2023. Liver three-dimensional cellular models for high-throughput chemical testing. *Cell Reports Methods* 3, 100432.
- Ye, W., Ramos, E.H., Wong, B.C., Belsham, D.D., 2016. Beneficial Effects of Metformin and/or Salicylate on Palmitate- or TNF $\alpha$ -Induced Neuroinflammatory Marker and Neuropeptide Gene Regulation in Immortalized NPY/AgRP Neurons. *PLoS One* 11, e0166973.
- Zhang, Q., Li, X., Liu, X., Dong, M., Xiao, J., Wang, J., Zhou, M., Wang, Y., Ning, D., Ma, W., Zhu, W., Liu, T., Zhang, B.o., 2020a. Association between maternal antimony exposure and risk of gestational diabetes mellitus: a birth cohort study. *Chemosphere* 246, 125732.
- Zhang, Y.-F., Shan, C., Wang, Y., Qian, L.-L., Jia, D.-D., Zhang, Y.-F., Hao, X.-D., Xu, H.-M., 2020b. Cardiovascular toxicity and mechanism of bisphenol A and emerging risk of bisphenol S. *Sci. Total Environ.* 723, 137952.
- Zimmermann, L., Dierkes, G., Ternes, T.A., Völker, C., Wagner, M., 2019. Benchmarking the in vitro toxicity and chemical composition of plastic consumer products. *Environ. Sci. Technol.*
- Zimmermann, L., Scheringer, M., Geueke, B., Boucher, J.M., Parkinson, L.V., Groh, K.J., Muncke, J., 2022. Implementing the EU chemicals strategy for sustainability: the case of food contact chemicals of concern. *J. Hazard. Mater.*, 129167.
- Znaor, A., Skakkebaek, N.E., Meys, E.-D., Kuliš, T., Laversanne, M., Gurney, J., Sarfati, D., McGlynn, K.A., Bray, F., 2022. Global patterns in testicular cancer incidence and mortality in 2020. *Int. J. Cancer* 151, 692–698.