

Adult Cerebral Malaria: Acute and Subacute Imaging Findings, Long-term Clinical Consequences

Charles Coughlan,¹ Hans Rolf Jäger,^{2,3,4} David Brealey,⁵ Francesco Carletti,^{2,3} Harpreet Hyare,² Rajyabardhan Pattnaik,⁶ Praveen K. Sahu,⁷ Sanjib Mohanty,⁷ Sarah Logan,¹ Angelika Hoffmann,⁸ Samuel C. Wassmer,⁹ and Anna M. Checkley^{1,9}

¹Department of Infectious Diseases, University College London Hospitals NHS Foundation Trust, London, United Kingdom; ²Department of Radiology, University College London Hospitals NHS Foundation Trust, London, United Kingdom; ³Lysholm Department of Neuroradiology, National Hospital for Neurology and Neurosurgery, London, United Kingdom; ⁴Neuroradiological Academic Unit, Queen Square Institute of Neurology, London, United Kingdom; ⁵Department of Intensive Care Medicine, University College London Hospitals NHS Foundation Trust, London, United Kingdom; ⁶Ispat General Hospital, Rourkela, India; ⁷Center for the Study of Complex Malaria in India, Community Welfare Society Hospital, Rourkela, India; ⁸University Institute of Diagnostic and Interventional Neuroradiology, University Hospital Bern, Inselspital, University of Bern, Bern, Switzerland; and ⁹Department of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London, United Kingdom

Cerebral malaria is an important cause of mortality and neurodisability in endemic regions. We show magnetic resonance imaging (MRI) features suggestive of cytotoxic and vasogenic cerebral edema followed by microhemorrhages in 2 adult UK cases, comparing them with an Indian cohort. Long-term follow-up images correlate ongoing changes with residual functional impairment.

Keywords. malaria; cerebral malaria; neuro-imaging; MRI; neuro-disability.

CASE SUMMARIES

Case 1

A 28-year-old female presented to her local hospital 2 weeks after returning from Ghana with 5 days of fever, diarrhea, vomiting, and confusion. She had not taken antimalarial prophylaxis. She was febrile, agitated, and confused, with tachycardia and hypotension. Blood film showed 22.1% *Plasmodium falciparum* trophozoites. There was lactic acidosis (pH 7.2, pCO₂ 33 mmHg, lactate 8 mmol/L), hemolytic anemia (Hb 105 g/L, unconjugated bilirubin 1.30 mg/dL), severe thrombocytopenia (6×10^9 cells/L), raised

inflammatory markers (white cell count 20×10^9 cells/L, C-reactive protein 150 mg/L), acute kidney injury (serum creatinine 2.85 mg/dL), and disseminated intravascular coagulation (INR 1.5, fibrinogen 0.6 g/L). She was intubated and ventilated and treated with intravenous artesunate, broad-spectrum intravenous antibiotics, inotropes, continuous veno-venous hemodiafiltration and multiple red cell, platelet and fresh frozen plasma transfusions. Blood cultures were negative, and there was no other evidence of bacterial sepsis.

On transfer to our center (day 4), cumulative fluid balance was 10L positive. She received 9 days of IV artesunate and 5 days of oral artemether-lumefantrine. Malaria films were negative from day 5. A lumbar puncture was not done due to the patient's low platelet count, so the possibility of concurrent bacterial meningitis was not completely excluded. On day 7 Glasgow coma scale (GCS) was 4. Magnetic resonance imaging (MRI) demonstrated a right-sided subacute striatocapsular infarct, restricted diffusion in the corpus callosum, basal ganglia, and the hippocampus, and diffuse severe vasogenic edema, particularly affecting the posterior subcortical and deeper white matter tracts, with brain swelling and effacement of the posterior sulci (Figures 1A and 1B). By day 14, GCS was 7. Day 15 MRI showed significant reduction of brain swelling with diminishing vasogenic edema, resolved restricted diffusion (Figures 1A and 1B), and striatocapsular infarct remodeling. Signal alteration in the hippocampus and adjacent limbic structures persisted. Susceptibility weighted imaging (SWI) showed numerous new susceptibility artefacts in the subcortical white matter, corpus callosum, basal ganglia and cerebellum (Figure 1A). On day 23, the patient was extubated, was GCS 10, and was able to follow 1-step commands. By day 30, she could speak in short sentences, with pseudoathetic movements and clumsiness in her left arm. Day 50 MRI showed complete edema resolution and maturation of the right striatocapsular infarct. The hippocampus, parahippocampal gyrus, and insula continued to show T2-hyperintensity, suggestive of gliosis, and persistent microhemorrhages (Figure 1A).

Upon discharge (day 62) she could walk with assistance and communicate in full sentences, with subtle word-finding difficulties. Her Montreal Cognitive Assessment score was 21/30. Days 70 and 253 neurocognitive assessments (Supplementary Data) showed reduced mental processing speed and semantic fluency, and fatigability on prolonged concentration. All indices except language function improved between the 2 time-points; she remained aware of word-finding difficulties and reduced short-term memory a year later.

Case 2

A 17-year-old female travelled to Sierra Leone and Morocco for 3 weeks, taking malaria prophylaxis intermittently, and

Received 14 March 2023; editorial decision 03 October 2023; accepted 27 October 2023; published online 28 October 2023

Correspondence: A. Checkley, Department of Infectious Diseases, UCLH Hospitals NHS Foundation Trust, 250 Euston Rd, London NW1 2PG, United Kingdom (anna.checkley@nhs.net); S. Wassmer, Department of Infection Biology, London School of Hygiene and Tropical Medicine, Keppel St, London WC1E 7HT (sam.wassmer@lshtm.ac.uk).

Clinical Infectious Diseases®

© The Author(s) 2023. Published by Oxford University Press on behalf of Infectious Diseases Society of America.

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs licence (<https://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial reproduction and distribution of the work, in any medium, provided the original work is not altered or transformed in any way, and that the work is properly cited. For commercial re-use, please contact journals.permissions@oup.com <https://doi.org/10.1093/cid/ciad651>




Figure 1. Comparative MRI findings between case 1 and a patient from an Indian cerebral malaria series [1]. The acute stage of the disease is seen in day 1 images from Rourkela, India. The early subacute stage is seen in day 7 images from London and day 4 images from Rourkela, and residual changes are seen in day 50 images from London. **A**, Representative T2-weighted images, ADC maps, and susceptibility-weighted images (SWI) of case 1 are displayed at the level of the basal ganglia. **Day 7 (GCS 4)**: The T2-weighted image shows a right T2 hyperintense striatocapsular infarct (arrow, first row). In addition, there is more ill-defined T2 hyperintense signal in the subcortical white matter bilaterally (arrowhead) and corona radiata representing vasogenic edema with corresponding increase in ADC. There is sulcal effacement posteriorly (and to a lesser degree, anteriorly) indicating brain swelling. In addition to the striatocapsular infarct, there is symmetrical ADC decrease in the putamen bilaterally (arrows, second row) and in the corpus callosum, consistent with mild cytotoxic edema. The SWI shows subtle areas of punctate susceptibility artefacts in the internal capsules and subcortical white matter. **Day 15 (GCS 7)**: T2 hyperintensities in the right striatum and posterior internal capsule and corona radiata remain, while sulci are now clearly visible, indicating brain swelling reversal. There is no longer evidence of cytotoxic edema with reduced ADC. The areas of increased T2 signal show increased ADC corresponding to residual vasogenic edema in the subcortical white matter and corona radiata. Numerous areas of susceptibility artefacts are now much more clearly visible in the subcortical white matter (arrowhead points at susceptibility artefacts in the occipital subcortical white matter), corpus callosum, corona radiata and internal capsules, corresponding to microthrombi or microhemorrhages. **Day 50 (GCS 15)**: A well-defined T2 hyperintensity indicates maturation of the right capsular infarct (arrow). The ADC signal has normalised. Extensive areas of susceptibility artefact persist within the same distribution as at day 15, appearing more prominent as imaging has been performed at 3 T compared to 1.5 T previously (arrowhead points at susceptibility artefacts in the occipital subcortical white matter). Thus, at day 50 brain swelling has completely reversed, with a mature right striatocapsular infarct and microthrombi/microhemorrhages in the subcortical white matter, corpus callosum, corona radiata and internal capsules. **B**, ADC maps at the level of the superior corona radiata/centrum semiovale of case 1 show an ADC decrease at day 7 (arrows), a mildly increased ADC on day 15 (arrow) and a fully normalised ADC at day 50. **C**, representative T2-weighted images, ADC maps and SWI of a 32-year-old Indian male with CM are displayed at the level of the basal ganglia. **Day 1 (GCS 5)**: An increased T2 signal is evident in the cortex and the slightly swollen lateral basal ganglia (putamen, arrows). Sulci are narrowed. ADC decrease is seen in the subcortical white matter (arrowhead) and basal ganglia (putamen, arrows), as in 1A. On SWI no clear pathological signal is apparent. Brain swelling is evident with mild cytotoxic edema in the subcortical white matter and basal ganglia. **Day 4 (GCS 15)**: The T2 signal in the basal ganglia (arrows) and the cortex has normalised. Residual ADC decrease is seen in the occipital white matter (arrowhead). On the SWI image multiple susceptibility artefacts are seen in the subcortical white matter, corpus callosum and basal ganglia. Thus, brain swelling has reversed with remaining slight cytotoxic edema in the occipital white matter and newly apparent microthrombi/microhemorrhages in the subcortical white matter (arrowhead), corpus callosum and basal ganglia. Microthrombi/microhemorrhages have the same distribution as in 1A. **D**, ADC map of a 22-year-old Indian male with CM at the level of the corona radiata. **Day 1 (GCS 5)**: Cytotoxic edema, evidenced as ADC decrease is evident in the subcortical white matter (arrowhead) with sparing of the corona radiata. **Day 4 (GCS 15)**: Cytotoxic edema, evidenced as ADC decrease is now apparent in the corona radiata (arrows), whereas most of the previously affected subcortical white matter does not show an ADC decrease anymore, similar to the appearance in 1B. Diffusion restriction was also present in the basal ganglia on day 1, resolved by day 4 (not shown). Abbreviations: ADC, apparent diffusion coefficient; CM, cerebral malaria; GCS, Glasgow coma scale; MRI, magnetic resonance imaging; SWI, susceptibility weighted imaging.

becoming unwell 12 days after her return to the United Kingdom, with 5 days of fever, vomiting, confusion, and unsteadiness. Blood film showed *Plasmodium falciparum* parasitemia 1.1%. Hemoglobin was 7.3 g/L, white cell count normal ($8.74 \times 10^9/L$), platelets $26 \times 10^9/L$. There was lactic acidosis, with pH 7.1 and lactate 7.7 mmol/L, normal renal function, and no evidence of respiratory or renal compromise. GCS was 14 on admission and subsequently dropped to 11. She was treated with IV artesunate (10 days total) and ceftriaxone (5 days total) and transferred to our unit. Malaria treatment was completed with 3 days of oral artemether-lumefantrine.

Lumbar puncture showed <1 white blood cell/cu mm, 6 red blood cells/cu mm, cerebrospinal fluid (CSF) protein 0.84 g/L, CSF glucose 3.8 mmol/L (plasma glucose 6.2), CSF lactate 2.54 mmol/L. Blood cultures were negative, and there was no evidence of bacterial sepsis. By day 4, GCS was 6, and MRI on day 5 showed restricted diffusion in the corpus callosum and subcortical white matter, vasogenic edema in the basal ganglia and susceptibility artefacts in the subcortical white matter, corpus callosum, and internal capsule (Supplementary Figure 1A). She was extubated on day 8, and by day 15 could walk with assistance, was orientated to time but not place, with limited attention and slow processing, and able to follow 1-stage commands. MRI on day 21 showed nearly complete resolution of the restricted diffusion as well as the vasogenic edema, with more prominent susceptibility artefacts (Supplementary Figure 1B). Upon discharge on day 27, she was fully orientated and ready to return to school. Neurocognitive testing (WAIS III) on day 117 showed mild to moderate under-functioning in verbal and non-verbal domains from high average premorbid estimates. Poor visual recall memory and executive functioning tests suggested medial temporal and frontal weakness.

DISCUSSION

We present 2 cerebral malaria (CM) cases in non-immune patients returning from West Africa, comparing MRI changes with our cohort of 85 cases in Rourkela, India [1], a malaria-endemic setting with seasonal transmission. Follow-up imaging and detailed psychometric assessments provide unique insights into the evolution of neurologic changes in adult patients.

Both UK and Indian cases had diffusion restriction and vasogenic edema in hypoxia-sensitive areas of the brain (Figure 1, Supplementary Figure 1) [2, 3]. Restricted diffusion with decreased apparent diffusion coefficient (ADC) values likely represents cytotoxic edema secondary to ischaemia caused by *P. falciparum* sequestration in the central nervous system (CNS) microvasculature [4]. Although diffusion restriction can also be caused by intramyelinic edema (more common in toxic and metabolic conditions), it appears less likely to be the cause for diffusion restriction observed here.

In case 1, MRI after 1 week showed generalized brain swelling and significant vasogenic edema with mild cytotoxic edema

in the subcortical and deep white matter [2]. There was also a basal ganglia infarct likely resulting from extensive sequestration [5]. Case 1 might not have survived outside of this well-resourced setting. Case 2 was less severely affected, with cytotoxic edema in the subcortical white matter and corpus callosum and milder generalized brain swelling and vasogenic edema in the basal ganglia. The vasogenic edema in case 1 was unusually marked, possibly due to the initial aggressive fluid resuscitation [6]. The same brain areas were involved in our cohort of Indian CM patients imaged at admission, but cytotoxic edema was the dominant feature with a lesser degree of vasogenic edema. (Figure 1, Supplementary Figure 1) [1].

We suggest that cytotoxic edema develops first in hypoxia-sensitive regions during the acute stage of CM, followed by vasogenic edema in the subacute stage (Figure 1A–D), as described in other forms of hypoxic brain injury. Depending on the degree of cytotoxic edema, it can reverse completely, for example, in rapidly recanalized stroke patients, or turn into vasogenic edema in more severely affected brain regions. This mechanism of initial cytotoxic edema development may explain why adjunctive steroids, hypertonic saline and mannitol are not efficacious in adult CM [7].

In both settings, reversal of brain swelling was accompanied by the development of numerous susceptibility artefacts in the subcortical gray matter, corpus callosum, internal capsule and basal ganglia, representing microhemorrhages or intravascular microthrombi (Figure 1C, Supplementary Figure 1). Both microthrombi at autopsy and SWI abnormalities consistent with microhemorrhages are associated with pediatric CM [8]. Damaged or previously thrombosed microvessels cannot withstand the pressure of normalised blood flow following parasite clearance, leading to rupture and microhemorrhage. Similar changes have been reported in coronavirus disease 2019 (COVID-19), severe respiratory failure, and disseminated intravascular coagulation [9, 10]. There was no clinical evidence of bacterial sepsis or significant hypoxia in the UK cases although disseminated intravascular coagulation (DIC) could have contributed to the microhemorrhagic pattern in case 1.

Although brain swelling and vasogenic edema persisted for weeks in case 1, repeat neuroimaging within 48–72 hours of treatment in the adult Indian cohort showed a much faster resolution of these changes [1]. The UK cases had severe disease with longer intensive care unit (ICU) stays and slower neurologic recoveries compared to the Indian cases, all of whom showed a GCS improvement to 14 or 15 within 72 hours. Long-term neurocognitive follow-up testing showed ongoing functional deficits. In case 1 these were in memory and processing, correlating with persistent imaging changes in the hippocampus.

CONCLUSION

We describe imaging findings consistent with acute cytotoxic edema and subacute vasogenic edema and likely microhemorrhage in

2 UK adults with CM. These findings are similar to those described in a large Indian cohort, despite different genetic backgrounds, degrees of pre-existing immunity, and geographical settings. Our findings support the hypothesis that CM is caused by hypoxia-induced edema through sequestration of *Plasmodium falciparum*-parasitized erythrocytes within the cerebral microvasculature [1], followed by vasogenic edema. These are the first cases we are aware of where long-term neurocognitive follow-up testing shows ongoing functional deficits and, in case 1, sequential imaging showing persistent brain alterations correlating with these functional impairments.

Historically, long-term neurologic sequelae of CM have been considered rare in adults [7]. CM is a major cause of childhood epilepsy and neurodisability in endemic regions [11], and there is mounting concern that it may cause unrecognized lifelong neurocognitive deficits in pediatric and adult survivors [12, 13]. With emerging evidence of neuroimaging changes in adults with both uncomplicated and severe non-cerebral malaria [14], there is an urgent need to investigate long-term neurocognitive outcomes following falciparum malaria.

Supplementary Data

Supplementary materials are available at *Clinical Infectious Diseases* online. Consisting of data provided by the authors to benefit the reader, the posted materials are not copyedited and are the sole responsibility of the authors, so questions or comments should be addressed to the corresponding author.

Notes

Author Contributions. C. C., A. C., S. W., and A. H. wrote the article. A. H., R. J., and F. C. analyzed the radiological images. All authors commented on the article.

Acknowledgments. The authors would like to thank the patients and their families for their permission to publish their clinical details.

Financial support. S. C. W. is supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health (NIH) under award number U19AI089676, and the Medical Research Council UK under award number MR/S009450/1. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

Potential conflicts of interest. H. R. J. reports royalties from Springer Nature Publishers for editing textbook of Clinical Neuroradiology; participation on Parexel MS Study Merck KGaA MS700568_0022 (Evaluation of Potential Progressive Multifocal Leukoencephalopathy [PML] Lesions on Imaging). S. C. W. reports grants unrelated to this work (NIH, USA (ref: R21AI142472) and Wellcome, UK (ref: 214227/Z/18/Z)). All other authors report no potential conflicts.

All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Sahu PK, Hoffmann A, Majhi M, et al. Brain magnetic resonance imaging reveals different courses of disease in pediatric and adult cerebral malaria. *Clin Infect Dis* **2021**; 73:e2387–96.
2. Mohanty S, Benjamin LA, Majhi M, et al. Magnetic resonance imaging of cerebral malaria patients reveals distinct pathogenetic processes in different parts of the brain. *mSphere* **2017**; 2:e00193-17.
3. Seydel KB, Kampondeni SD, Valim C, et al. Brain swelling and death in children with cerebral malaria. *N Engl J Med* **2015**; 372:1126–37.
4. Ponsford MJ, Medana IM, Prapansilp P, et al. Sequestration and microvascular congestion are associated with coma in human cerebral malaria. *J Infect Dis* **2012**; 205:663–71.
5. Rasalkar DD, Paunipagar BK, Sanghvi D, Sonawane BD, Loniker P. Magnetic resonance imaging in cerebral malaria: a report of four cases. *Br J Radiol* **2011**; 84: 380–5.
6. van der Jagt M. Fluid management of the neurological patient: a concise review. *Crit Care* **2016**; 20:126.
7. Mishra SK, Newton CR. Diagnosis and management of the neurological complications of falciparum malaria. *Nat Rev Neurol* **2009**; 5:189–98.
8. Dorovini-Zis K, Schmidt K, Huynh H, et al. The neuropathology of fatal cerebral malaria in Malawian children. *Am J Pathol* **2011**; 178:2146–58.
9. Riech S, Kallenberg K, Moerer O, et al. The pattern of brain microhemorrhages after severe lung failure resembles the one seen in high-altitude cerebral edema. *Crit Care Med* **2015**; 43:e386–9.
10. Neligan A, Rajakulendran S, Nortley R, Manji H. Extensive cerebral microhemorrhages caused by acute disseminated intravascular coagulation secondary to sepsis. *JAMA Neurol* **2014**; 71:510–1.
11. Birbeck GL, Molyneux ME, Kaplan PW, et al. Blantyre malaria project epilepsy study (BMPES) of neurological outcomes in retinopathy-positive paediatric cerebral malaria survivors: a prospective cohort study. *Lancet Neurol* **2010**; 9: 1173–81.
12. Oluwayemi IO, Brown BJ, Oyedeji OA, Oluwayemi MA. Neurological sequelae in survivors of cerebral malaria. *Pan Afr Med J* **2013**; 15:88.
13. Peixoto B, Kalei I. Neurocognitive sequelae of cerebral malaria in adults: a pilot study in Benguela Central Hospital, Angola. *Asian Pac J Trop Biomed* **2013**; 3: 532–5.
14. Mohanty S, Sahu PK, Pattnaik R, et al. Evidence of brain alterations in noncerebral falciparum malaria. *Clin Infect Dis* **2022**; 75:11–8.