

1 **Insulin-like growth factor binding protein-2 in at-risk adults** 2 **and autopsy-confirmed Alzheimer brains**

3 Marc James Quesnel,^{1,2} Anne Labonté,^{2,3} Cynthia Picard,^{2,3} Henrik Zetterberg,^{4,5,6,7,8,9} Kaj
4 Blennow,^{4,5,10,11} Ann Brinkmalm,^{4,5} Sylvia Villeneuve^{1,2,3} and Judes Poirier^{1,2,3} for the
5 Alzheimer's Disease Neuroimaging Initiative and the PREVENT-AD Research Group

6 **Abstract**

7 Insulin, insulin-like growth factors (IGF) and their receptors are highly expressed in the adult
8 hippocampus. Thus, disturbances in the insulin-IGF signaling pathway may account for the
9 selective vulnerability of the hippocampus to nascent Alzheimer's disease (AD) pathology. In the
10 present study, we examined the predominant IGF-binding protein (IGFBP) in the cerebrospinal
11 fluid (CSF) – IGFBP2.

12 CSF was collected from 109 asymptomatic members of the parental history-positive PREVENT-
13 AD cohort. CSF levels of IGFBP2, core AD biomarkers and synaptic biomarkers were measured
14 using proximity extension assay, ELISA and mass spectrometry. Cortical amyloid-beta (A β) and
15 tau deposition were examined using ¹⁸F-NAV4694 and flortaucipir. Cognitive assessments were
16 performed up to 8 years of follow-up, using the Repeatable Battery for the Assessment of
17 Neuropsychological Status. T1-weighted structural MRI scans were acquired, and neuroimaging
18 analyses were performed on pre-specified temporal and parietal brain regions. Next, in an
19 independent cohort, we allocated 241 dementia-free ADNI-1 participants into four stages of AD
20 progression based on the biomarkers CSF A β ₄₂ and total-tau (t-tau). In this analysis, differences in
21 CSF and plasma IGFBP2 levels were examined across the pathological stages. Finally, IGFBP2
22 mRNA and protein levels were examined in the frontal cortex of 55 autopsy-confirmed AD and
23 31 control brains from the QFP cohort, a unique population isolate from Eastern Canada.

24 CSF IGFBP2 progressively increased over 5 years in asymptomatic PREVENT-AD participants.
25 Baseline CSF IGFBP2 was positively correlated with CSF AD biomarkers and synaptic
26 biomarkers, and was negatively correlated with longitudinal changes in delayed memory ($P =$
27 0.024) and visuospatial abilities ($P = 0.019$). CSF IGFBP2 was negatively correlated at a trend-

1 level with entorhinal cortex volume ($P = 0.082$) and cortical thickness in the piriform ($P = 0.039$),
2 inferior temporal ($P = 0.008$), middle temporal ($P = 0.014$) and precuneus ($P = 0.033$) regions. In
3 ADNI-1, CSF ($P = 0.009$) and plasma ($P = 0.001$) IGFBP2 were significantly elevated in Stage 2
4 (CSF A β (+)/t-tau(+)). In survival analyses in ADNI-1, elevated plasma IGFBP2 was associated
5 with a greater rate of AD conversion (HR = 1.62, $P = 0.021$). In the QFP cohort, IGFBP2 mRNA
6 was reduced ($P = 0.049$), however IGFBP2 protein levels did not differ in the frontal cortex of
7 autopsy-confirmed AD brains ($P = 0.462$).

8 Nascent AD pathology may induce an upregulation in IGFBP2, in asymptomatic individuals. CSF
9 and plasma IGFBP2 may be valuable markers for identifying CSF A β (+)/t-tau(+) individuals and
10 those with a greater risk of AD conversion.

11

12 **Author affiliations:**

13 1 McGill University, Department of Psychiatry, Montréal, Québec, Canada, H3A 1A1

14 2 Douglas Mental Health University Institute, Research Centre, Montréal, Québec, Canada, H4H
15 1R3

16 3 Centre for the Studies in the Prevention of Alzheimer's Disease, Douglas Mental Health
17 University Institute, Montréal, Québec, Canada, H4H 1R3

18 4 Department of Psychiatry and Neurochemistry, Institute of Neuroscience and Physiology, The
19 Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

20 5 Clinical Neurochemistry Department, Sahlgrenska University Hospital, Mölndal, Sweden

21 6 Department of Neurodegenerative Disease, UCL Institute of Neurology, London, WC1N 3BG,
22 UK

23 7 UK Dementia Research Institute at UCL, London, WC1E 6BT, UK

24 8 Hong Kong Center for Neurodegenerative Diseases, Hong Kong, China

25 9 Wisconsin Alzheimer's Disease Research Center, University of Wisconsin School of Medicine
26 and Public Health, University of Wisconsin-Madison, Madison, WI 53792-2420 USA

1 10 Paris Brain Institute, ICM, Pitié-Salpêtrière Hospital, Sorbonne University, 21 414 – 75646
2 Paris Cedex 13, Paris, France

3 11 Neurodegenerative Disorder Research Center, Division of Life Sciences and Medicine, and
4 Department of Neurology, Institute on Aging and Brain Disorders, University of Science and
5 Technology of China and First Affiliated Hospital of USTC, Hefei, P.R. China

6
7 Correspondence to: Dr Judes Poirier

8 Centre for the Studies in the Prevention of Alzheimer's Disease, Douglas Mental Health University
9 Institute, 6875, Lasalle, Montréal, Quebec H4H 1R3, Canada

10 E-mail: judes.poirier@mcgill.ca

11 **Running title:** Role of IGFBP2 across the AD spectrum

12
13 **Keywords:** insulin-like growth factor-binding protein-2; insulin-like growth factor; Alzheimer's
14 disease; cerebrospinal fluid; post-mortem brain tissue; RBANS

15 **Abbreviations:** A β , amyloid-beta; A β ₄₂, amyloid-beta 42; AD, Alzheimer's disease; ADNI,
16 Alzheimer's Disease Neuroimaging Initiative; ANCOVA, analysis of covariance; APOE,
17 Apolipoprotein E; BBB, blood-brain barrier; BIOMARKAPD, Biomarkers for Alzheimer's and
18 Parkinson's Disease; BMI, body mass index; BP, blood pressure; CAIDE, Cardiovascular Risk
19 Factors, Aging, and Incidence of Dementia; CI, confidence interval; CTL, control; CU, cognitively
20 unaffected; GAP43, growth-associated protein 43; GSK3, glycogen synthase kinase-3; HbA1c,
21 hemoglobin A1C; HR, hazard ratio; ICV, intracranial volume; IGF: insulin-like growth factor;
22 IGF-1/IGF-I, insulin-like growth factor 1; IGF-2/IGF-II, insulin-like growth factor 2; IGFBP,
23 insulin-like growth factor binding protein; IGFBP2, insulin-like growth factor binding protein-2;
24 MCI, mild cognitive impairment; MoCA, Montréal cognitive assessment; NINCDS-ADRDA,
25 National Institute of Neurological and Communicative Disorders and Stroke and the Alzheimer's
26 Disease and Related Disorders Association; NPX, Normalized Protein eXpression; NRG1,
27 neurogranin; PET, positron emission tomography; PI3K-AKT, phosphoinositide 3-kinase-protein
28 kinase B; PNS, peripheral nervous system; PREVENT-AD, PRE-symptomatic EValuation of
29 Experimental or Novel Treatments for Alzheimer's Disease; p₁₈₁-tau, phosphorylated tau 181;

1 RBANS, Repeatable Battery for the Assessment of Neuropsychological Status; ROI, region of
2 interest; Rpm, revolutions per minute; SHANK3, SH3 And Multiple Ankyrin Repeat Domains 3;
3 SNAP, suspected non-Alzheimer pathology; SNAP25, synaptosomal-associated protein 25;
4 SUVR, standardized uptake value ratio; SYT1, synaptotagmin-1; t-tau: total tau
5
6

7 **Introduction**

8 By 2050, is it estimated that 152.8 million individuals worldwide will be affected by dementia.¹
9 Furthermore, the cost of dementia care is projected to reach \$16.9 trillion in 2050.² Alzheimer's
10 disease (AD) remains one of the most challenging medical mysteries, as it is the most common
11 cause of dementia - accounting for 60-80% of all cases.³ The neuropathological hallmarks of AD
12 include the accumulation of extracellular amyloid-beta ($A\beta$) plaques, intracellular neurofibrillary
13 tangles, as well as the loss of synapses and neurons.⁴ These brain changes are believed to begin up
14 to 20 years or more, before the onset of symptoms.⁵

15 It has been well established that insulin resistance and diabetes are risk factors for developing
16 AD.^{6,7} Indeed, impaired insulin and insulin-like growth factor (IGF) signalling plays a critical role
17 in the pathogenesis of AD.⁸⁻¹⁰ Post-mortem studies have demonstrated that insulin, IGFs as well
18 as their receptors and downstream signalling molecules, are decreased in the AD brain.⁸⁻¹¹
19 Furthermore, the insulin-IGF system has been shown to directly modulate $A\beta$ degradation¹² and
20 clearance,¹³ phospho-tau production,^{14,15} synaptic integrity¹⁶ and neuronal survival.¹⁷ It is also
21 known that insulin, IGFs and their receptors are highly expressed in the hippocampus, relative to
22 the frontal cortex in the human brain.⁸ Overall, these findings suggest that impairments in insulin-
23 IGF signaling may account for the selective vulnerability of the hippocampus to nascent AD
24 pathology, and therefore, account for early impairments in episodic memory. Similarly,
25 deficiencies in insulin-IGF signaling may contribute to the reductions in glucose metabolism that
26 are seen in patients with AD and individuals at risk for AD.¹⁸ The importance of insulin and IGFs
27 has been emphasized in pilot clinical trials in which the administration of intranasal insulin
28 improved memory, caregiver-rated functional abilities, and glucose metabolism in individuals with
29 mild cognitive impairment (MCI) or mild AD.^{19,20}

1 The insulin-IGF system encompasses a complex collection of proteins that play pivotal roles in
2 glucose metabolism, neurogenesis, synaptogenesis and cell survival.^{17,21,22} Insulin, IGF-I and IGF-
3 II are the key proteins, which bind to their cell surface receptors.²² The actions of IGF-I and IGF-
4 II are modulated by six IGF-binding proteins (IGFBP), which can bind to IGFs with an equal or
5 greater affinity than the IGF receptors.^{23,24} Indeed, in the circulation, CSF and local tissues, most
6 extracellular IGFs are bound to so-called IGFBPs, which prolong the half-life of IGFs.²²⁻²⁴ For
7 instance, it has been proposed that IGFBPs may prevent the degradation of IGFs during transport
8 and mobilisation, and target IGFs to their receptors.²²⁻²⁴ The latter may be achieved through
9 IGFBPs binding to cell-surface proteoglycans²⁵ and integrins,²⁶ or proteolytic cleavage,²⁷ both of
10 which reduce the binding affinity of IGFBPs for IGFs and promote IGF release.²²

11 The main objective of the current study is to examine a less studied member of the IGF molecular
12 cascade, insulin-like growth factor binding protein-2 (IGFBP2), in both the pre-symptomatic and
13 symptomatic stages of AD. Since IGFBP2 is the most abundant IGF-binding protein in the
14 CSF,^{28,29} we hypothesize IGFBP2 plays a critical role in the neurodegenerative process, most likely
15 at the level of neuroprotection and resilience.²² IGFBP2 is increased in the CSF and plasma of
16 patients with a clinical diagnosis of AD,³⁰⁻³⁵ and is associated with longitudinal atrophy in
17 entorhinal, parahippocampal and inferior temporal regions.³⁶ Moreover, elevated circulating levels
18 of IGFBP2 have been associated with an increased risk of developing AD.^{35,37,38} Finally, in a pilot
19 study, IGFBP2 has been found to be decreased in the temporal cortex of AD patients.⁹ These and
20 other findings will be verified in the pre-symptomatic and symptomatic stages of the disease, as
21 well as in autopsy-confirmed AD brains.

22

23 **Materials and methods**

24 **PREVENT-AD cohort**

25 **Study participants**

26 The PRE-symptomatic EVALuation of Experimental or Novel Treatments for Alzheimer's Disease
27 (PREVENT-AD) cohort consists of asymptomatic, "at-risk", individuals with a parental or multi-
28 sibling history of sporadic Alzheimer's disease.³⁹ The majority of participants were over the age

1 of 60, however individuals aged 55-59 years were included if they were within 15 years of the
2 onset of their youngest-affected relative's symptoms. In order to confirm normal cognition, the
3 Clinical Dementia Rating and Montreal Cognitive Assessment (MoCA) were used at the study
4 eligibility visit. 386 active PREVENT-AD participants have been followed longitudinally, as
5 annual visits include cognitive assessments, neurosensory tests, blood and (for a subset of
6 individuals) CSF collections, structural and functional MRI scans, as well as PET scans. Each
7 participant and their study partner provided written informed consent. All procedures were
8 approved by the McGill University Faculty of Medicine Institutional Review Board and complied
9 with the ethical principles of the Declaration of Helsinki. A detailed description of the PREVENT-
10 AD cohort is available elsewhere.³⁹

11

12 **Cerebrospinal fluid measurements**

13 A Sprotte 24-gauge atraumatic needle was used to perform lumbar punctures (LP) in PREVENT-
14 AD participants, following an overnight fast. In order to exclude cells and insoluble material, CSF
15 samples were centrifuged (~2000g) within 4 hours, for 10 minutes at room temperature. Finally,
16 the CSF samples were aliquoted (0.5 mL) into polypropylene cryotubes and stored at -80°C.

17 CSF IGFBP2 levels were measured in a subset of PREVENT-AD participants ($n = 109$) using the
18 Olink Cardiovascular III panel (Uppsala, Sweden), which employs proximity extension assay
19 technology. Olink measurements are expressed in arbitrary Normalized Protein eXpression units
20 (NPX), which are on a \log_2 scale.

21 CSF AD biomarkers amyloid-beta 42 ($A\beta_{42}$), phosphorylated tau (p_{181} -tau) and total tau (t-tau)
22 were measured in a subset ($n = 101$) of PREVENT-AD participants, using the validated Innotech
23 ELISA kit (Fujirebio, Ghent, Belgium) following the standardized protocols established by the
24 BIOMARKAPD consortium ($A\beta_{42}$ Cat.# 81583, p_{181} -tau Cat.# 81581 and t-tau Cat.# 81579).

25 Of the 109 PREVENT-AD participants that had CSF IGFBP2 measurements, 106 individuals had
26 the synaptic proteins synaptosomal-associated protein 25 (SNAP25) and synaptotagmin-1 (SYT1)
27 assayed. CSF SNAP25 and SYT1 were immunoprecipitated and their concentrations were
28 determined by mass spectrometry, as previously described.^{40,41,42} Mass spectrometry results are
29 expressed in arbitrary units.⁴³ As previously reported, CSF levels of growth-associated protein 43

1 (GAP43) and neurogranin (NRGN) were assessed by validated ELISAs in a subset of PREVENT-
2 AD individuals ($n = 46$).^{44,45}

3

4 **Neuroimaging acquisition and processing**

5 *In vivo* cortical A β and phosphorylated tau pathologies were determined using PET tracers ¹⁸F-
6 NAV4694 (Navidea Biopharmaceuticals, Dublin, OH, USA) and flortaucipir (¹⁸F-AV1451; Eli
7 Lilly & Company, Indianapolis, IN, USA), in a subset of PREVENT-AD participants that also had
8 CSF IGFBP2 measurements ($n = 46$, $n = 49$ respectively). A β and tau PET scans were performed
9 40 to 70 minutes and 80 to 100 minutes post-injection, respectively. A 3T Siemens Trio scanner
10 was used to acquire T1-weighted structural MRI scans at the Douglas Mental Health University
11 Institute (Montreal). A Siemens standard 12 or 32-channel coil was used (Siemens Medical
12 Solutions, Erlangen). FreeSurfer 5.3 was used to process the MRI scans, and the Desikan-Killiany
13 atlas was used for parcellation. The preprocessing pipeline for PET images has previously been
14 described.⁴⁶ Briefly, standardized uptake value ratios (SUVRs) were generated by dividing the
15 signal in the regions of interest (ROI) by the signal in the reference region. Thus, cerebellar grey
16 matter was used as a reference region for ¹⁸F-NAV4694, whilst the inferior cerebellar grey matter
17 was used for flortaucipir. A global cortical ROI was computed to evaluate A β deposition, whilst
18 tau deposition was assessed by averaging flortaucipir SUVRs in the entorhinal cortex and lingual
19 gyri. The imaging processing pipeline CIVET 1.1.12 was used to estimate cortical thickness from
20 T1-weighted images ($n = 104$).⁴⁷ Brain volumes were computed using a volumetric pipeline that
21 has been previously described.⁴⁸

22

23 **Apolipoprotein E genotyping**

24 The QIASymphony apparatus and DNA Blood Mini QIA Kit were used to isolate DNA from 200
25 μ l whole blood (Qiagen, Valencia, CA, USA). The standard QIASymphony isolation program was
26 used following the manufacturer's instructions. The PyroMark Q96 pyrosequencer (Qiagen,
27 Toronto, Ontario, Canada) was used to determine apolipoprotein E (APOE) genotype in
28 PREVENT-AD. qPCR was used to amplify DNA, with primers rs429358 amplification forward
29 5'-ACGGCTGTCCAAGGAGCTG-3', rs429358 amplification reverse biotinylated 5'-

1 CACCTCGCCGCGGTACTG-3', rs429358 sequencing 5'CGGACATGGAGGACG-3', rs7412
2 amplification forward 5'-CTCCGCGATGCCGATGAC-3', rs7412 amplification reverse
3 biotinylated 5'-CCCCGGCCTGGTACACTG-3' and rs7412 sequencing 5'-CGA
4 TGACCTGCAGAAG-3'.

6 **Cognitive testing**

7 At annual visits, the Repeatable Battery for the Assessment of Neuropsychological Status
8 (RBANS) was used to assess the cognitive performance of PREVENT-AD participants. The
9 RBANS possesses an excellent sensitivity in differentiating normal cognition from MCI.⁴⁹ Five
10 cognitive domains are evaluated, which include immediate memory, delayed memory, attention,
11 language and visuospatial abilities.⁴⁹ A total summary score is included as well. Each participant's
12 score is standardized by their age, such that a score of 100 represents the expected cognitive
13 performance for a given age.⁴⁹ In order to reduce practice effects in longitudinal assessment, the
14 RBANS was available in four equivalent versions. Furthermore, the battery was administered in
15 English or French depending on the participants' preferred language. Cognitive measurements are
16 available for up to 8 years of follow-up.

18 **ADNI-1 cohort**

19 **Study participants**

20 Led by Principal Investigator Michael W. Weiner, MD, the primary objective of the Alzheimer's
21 Disease Neuroimaging Initiative (ADNI) has been to detect the earliest changes associated with
22 AD, and to track the progression of AD pathology. Given our interest in the earliest possible stages
23 of AD, we restricted our primary analyses to 241 ADNI-1 participants with CSF data available
24 from 92 cognitively unaffected (CU) individuals and 149 individuals with MCI. For analyses
25 involving plasma samples, we restricted our analyses to 58 CU individuals and 396 individuals
26 with MCI that had available data. Two individuals with ambiguous diagnoses were excluded from
27 analyses.

28

1 **Cerebrospinal fluid measurements**

2 LPs were performed with a 20- or 24- gauge spinal needle, following an overnight fast. CSF
3 samples were frozen within 1 hour after collection and shipped on dry ice to the ADNI Biomarker
4 Core laboratory. Following thawing (1h) with gentle mixing at room temperature, the samples
5 were aliquoted (0.5 mL) into polypropylene vials and stored at -80°C. CSF AD biomarkers A β ₄₂,
6 p181-tau and t-tau were measured in ADNI-1 samples using the INNO-BIA AlzBio3 immunoassay
7 kits (Fujirebio, Ghent, Belgium) and the xMap Luminex platform (Austin, Texas, USA).

8 CSF levels of 159 inflammatory, metabolic and lipid analytes, including IGFBP2 had been
9 assessed with the Human Discovery Map panel, a multiplex immunoassay panel developed by
10 Rules Based Medicine. In the case of IGFBP2, 8 (imputed) samples exceeded the detectable
11 analyte concentration range and were omitted from subsequent analyses. Finally, in supplementary
12 analyses, we further analyzed multiple reaction monitoring mass spectrometry measurements of
13 CSF IGFBP2.

14 **Pathological staging of participants**

15 Following the recent emergence of biological frameworks for defining AD,⁵⁰ we used baseline
16 CSF A β ₄₂ and CSF t-tau measurements to stage 90 CU individuals and 145 individuals with MCI.
17 We applied the recommended CSF A β ₄₂ and CSF t-tau thresholds of <192 pg/mL and >93 pg/mL,
18 respectively.⁵¹ These cutoff values have been generated from autopsy-based AD CSF samples and
19 have been reported to detect mild AD and predict the conversion from MCI to AD.⁵¹ Two
20 individuals with biomarker measurements equivalent to the threshold values were removed.

21 ADNI-1 participants were assigned to Stage 0, A β (-)/t-tau(-), if they had normal levels of CSF
22 A β ₄₂ and CSF t-tau. Participants in Stage 1, A β (+)/t-tau(-), exhibited early amyloid pathology, as
23 reflected by reduced levels of CSF A β ₄₂. However, individuals in Stage 1 did not display
24 significant levels of neuronal loss, as reflected by low levels of CSF t-tau. In Stage 2, A β (+)/t-
25 tau(+), participants exhibited low levels of CSF A β ₄₂ and elevated levels of CSF t-tau. Finally, the
26 Suspected Non-Alzheimer Pathology (SNAP) group, A β (-)/t-tau(+), exhibited normal levels of
27 CSF A β ₄₂ and elevated levels of CSF t-tau, thus suggesting other causes of neurodegeneration
28 and/or dementia.

29

1 **Plasma measurements**

2 At the baseline visit, plasma samples were drawn following an overnight fast. 192 analytes that
3 have been reported to be altered in cancer, cardiovascular disease, metabolic disorders,
4 inflammation and AD were analyzed with the Human Discovery Map panel, a multiplex
5 immunoassay panel developed on the Luminex xMAP platform by Rules Based Medicine. In order
6 to meet model assumptions, plasma IGFBP2 levels were log₁₀ transformed by the ADNI
7 investigators.

8 9 **Apolipoprotein E genotyping**

10 The ABI 7900 real-time thermo-cycler (Applied Biosystems, Foster City, CA) was used to
11 determine the APOE genotype of ADNI participants. TaqMan qPCR was applied to DNA prepared
12 from EDTA whole blood.

13 14 **QFP cohort**

15 **Study participants**

16 The Quebec Founder Population (QFP) is composed of the descendants of a few thousand French
17 settlers that colonized Nouvelle France in the 17th and 18th centuries.⁵² The migration and the
18 isolated nature of settlements created a founder effect, which resulted in a population with less
19 genetic heterogeneity.⁵² Genealogical information for this population, for almost four centuries, is
20 available in the BALSAC database. In the present study, we analyzed the brains of 55 autopsy-
21 confirmed AD cases and 31 autopsy-confirmed elderly controls, which were obtained from the
22 Douglas-Bell Canada Brain Bank. According to medical record reviews, neuropsychological
23 examinations and caregiver interviews, there was no evidence of memory problems, neurological
24 or neuropsychiatric diseases in the elderly control group. Furthermore, controls only exhibited
25 neuropathology that is associated with healthy aging (plaque and tangle densities <10/mm³ and <
26 20/mm³ in at least one hippocampal and neocortical section). AD cases had to fulfill the
27 histopathological NINCDS-ADRDA criteria for definite AD.⁵³ This study was conformed to the
28 Code of Ethics of the World Medical Association and was approved by the Ethics Board of the

1 Douglas Mental Health University Institute. This study complied with the ethical principles of the
2 Declaration of Helsinki. Each participant provided written informed consent.

3

4 **IGFBP2 gene expression in the frontal cortex**

5 In the frontal cortex of 31 autopsy-confirmed controls and 55 autopsy-confirmed AD cases,
6 transcriptome-wide gene expression was measured using the human Clariom D Assay, by Génome
7 Québec. Briefly, the Nanodrop Spectrophotometer ND-1000 was used to measure total RNA
8 (NanoDrop Technologies, Inc.). RNA integrity was evaluated with the Agilent 2100 Bioanalyzer.
9 10 ng of total RNA was used to synthesize sense-strand cDNA. The GeneChip WT Terminal
10 Labeling Kit was used to fragment and label single-stranded cDNA, following the manufacture's
11 instructions. 5 µg cDNA was hybridized on the GeneChip cartridge array and incubated at 45°C
12 for 17 hours in the GeneChip Hybridization Oven 640, at 60 rpm. The microarrays were washed
13 in the GeneChip Fluidics Station 450, using the GeneChip Hybridization, Wash, and Stain Kit,
14 according to the manufacture's instructions. Finally, microarrays were scanned in a GeneChip
15 Scanner 3000. IGFBP2 mRNA levels are presented on a log₂ scale.

16

17 **IGFBP2 protein levels in the frontal cortex**

18 Of the 86 brains that had IGFBP2 mRNA levels measured, 78 brains ($n = 25$ Controls, $n = 53$ AD)
19 had IGFBP2 protein levels measured. Frontal cortex brain samples were placed in pre-filled tubes
20 containing 2.8 mm ceramic beads (Omni International, GA, USA). One tablet of protease inhibitor
21 was dissolved in 50 mL of cold phosphate-buffered saline. 1 mL of protease inhibitor solution was
22 added to each tube. The Bead Ruptor 24 (Omni International, Kennesaw, GA, USA) was used to
23 mechanically homogenize brain samples. The Bead Ruptor 24 was set twice at 5.65m/s for 30
24 seconds, with a 15 second pause between runs. Following homogenization, the samples were
25 stored overnight at -20°C. In order to break the cell membranes, two freeze-thaw cycles were
26 performed. Finally, the homogenates were centrifuged for 5 minutes at 5000 rpm and 4°C. The
27 supernatant was collected and stored for future use at -80°C.

28 Frontal cortex IGFBP2 protein levels were measured using a commercially available ELISA kit
29 (Cat.# OKEH00084, Aviva Systems Biology, CA, USA). Protocols were performed according to

1 the manufacturers' instructions and results were obtained using the BioTek Synergy H1 microplate
2 reader (Winooski, Vermont, USA). Sample replicates had a coefficient of variability of less than
3 20%. Finally, in order to normalize IGFBP2 protein levels, total protein concentration was
4 measured using a commercially available bicinchoninic acid assay developed by Pierce (Cat.#
5 23225). Finally, normalized IGFBP2 protein ratios were \log_2 transformed in order to meet model
6 assumptions.

7

8 **Apolipoprotein E genotyping**

9 DNA was extracted from brain tissue with the DNeasy Tissue Kit (Qiagen). As previously
10 described in PREVENT-AD,³⁹ the PyroMark Q96 pyrosequencer was used to determine *APOE*
11 genotype.

12

13 **Statistical analyses**

14 Annual changes in CSF IGFBP2 levels were assessed in a subset of PREVENT-AD participants
15 ($n = 27$). Each participant's trajectory was analyzed using a linear mixed model with a random
16 intercept and slope. Age, sex and *APOE* $\epsilon 4$ carrier status – adjusted linear regression models were
17 used to examine the associations between baseline CSF IGFBP2 and baseline measurements of
18 CSF AD biomarkers ($A\beta_{42}$, p_{181} -tau, t-tau), CSF synaptic proteins (SYT1, SNAP25, GAP43,
19 NRGN) as well as PET and structural (volumetric, cortical thickness) neuroimaging data. For each
20 PREVENT-AD participant that was followed for 5-8 years ($n = 89$), an estimated cognitive
21 performance trajectory slope was calculated for each of the five cognitive domains of the RBANS.
22 Thus, linear regression models adjusted for age, sex, *APOE* $\epsilon 4$ carrier status and years of education
23 were used to examine the relationship between rate of change in cognition and baseline CSF
24 IGFBP2 levels. Across all analyses, IGFBP2 was assigned as a dependent variable, except for the
25 association with CSF $A\beta_{42}$. Finally, given the critical role of the insulin-IGF system in diverse
26 metabolic processes, we further controlled for clinical covariates such as body mass index (BMI),
27 systolic blood pressure and hemoglobin A1C levels (HbA1c), in supplementary analyses.
28 However, the results of these analyses were similar to those of the original model. Therefore, the
29 following results are presented according to the original model.

1 In the ADNI-1 cohort, analysis of covariance (ANCOVA) was used to assess the relationship
2 between baseline CSF and baseline plasma IGFBP2 with pathological stage as determined by CSF
3 $A\beta_{42}$ and CSF t-tau positivity.^{50,51} To correct for multiple planned comparisons between
4 pathological stages, statistical significance was considered at $P \leq 0.01$. Finally, Cox proportional
5 hazards models examined the association between baseline plasma IGFBP2 levels and rate of
6 conversion to AD. CU participants and individuals with MCI were followed from the baseline visit
7 to the time of diagnosis (of AD), or to the time the participant was last confirmed to be free of AD.
8 Cox models were adjusted for age, gender and *APOE* $\epsilon 4$ carrier status.

9 In the QFP cohort, the relationship between frontal cortex mRNA, protein levels, and diagnosis
10 was evaluated with ANCOVA adjusted for age at death, sex, *APOE* $\epsilon 4$ carrier status and post-
11 mortem delay. Statistical significance was considered at $P \leq 0.05$. R^2 values are presented as
12 adjusted R^2 . All Analyses were two-tailed and performed in SPSS 23 (IBM) and JMP Pro 16
13 (SAS).

14

15 **Results**

16 **Demographics**

17 Table 1 summarizes the demographic characteristics of the three cohorts that were used to analyze
18 the role of IGFBP2 in the CSF of asymptomatic (PREVENT-AD) and symptomatic individuals
19 (ADNI-1), as well as in the frontal cortex of autopsy-confirmed, age-matched control and AD
20 cases (QFP).

21

22 **PREVENT-AD cohort**

23 **CSF IGFBP2 increases annually and is associated with CSF and PET** 24 **biomarkers in asymptomatic AD**

25 In a subset of CU PREVENT-AD participants that had longitudinal CSF IGFBP2 measurements
26 available ($n = 27$), a random intercept and random slope linear mixed model revealed that CSF
27 IGFBP2 levels progressively increase over the course of five years ($\beta = 0.132$, $P = 0.005$, Fig. 1).

1 In PREVENT-AD participants ($n = 101$) that had CSF AD pathological biomarker measurements,
2 baseline CSF IGFBP2 levels were positively correlated with CSF $A\beta_{42}$ ($R^2 = 0.055$, $\beta = 91.232$, P
3 $= 0.023$, Fig. 2A), CSF p_{181} -tau ($R^2 = 0.162$, $\beta = 0.014$, $P = 4.50 \times 10^{-5}$, Fig. 2B) and CSF t-tau
4 ($R^2 = 0.121$, $\beta = 0.001$, $P = 0.001$, Fig. 2C). In a subset of PREVENT-AD participants that
5 underwent $A\beta$ and tau PET scans ($n = 46$ and $n = 49$), CSF IGFBP2 was not associated with global
6 cortical $A\beta$ deposition ($P = 0.540$, Fig. 2D). However, CSF IGFBP2 was positively correlated
7 (trend-level) with tau deposition in the entorhinal cortex ($R^2 = 0.027$, $\beta = 1.498$, $P = 0.082$, Fig.
8 2E) and lingual gyrus ($R^2 = 0.034$, $\beta = 2.226$, $P = 0.067$, Supplementary Fig. 1).

9 Finally, CSF IGFBP2 was positively associated with synaptic proteins in the CSF, including
10 SNAP25 (trend-level, $R^2 = 0.051$, $\beta = 0.014$, $P = 0.076$, Fig. 2F), SYT1 ($R^2 = 0.060$, $\beta = 0.008$, P
11 $= 0.043$, Fig. 2G), GAP43 ($R^2 = 0.243$, $\beta = 3.06 \times 10^{-4}$, $P = 2.52 \times 10^{-4}$, Fig. 2H), and NRGN
12 (trend-level, $R^2 = 0.033$, $\beta = 0.002$, $P = 0.063$, Fig. 2I).

13

14 **CSF IGFBP2 is linked to changes in delayed memory and visuospatial abilities** 15 **in asymptomatic AD**

16 Upon computing RBANS cognitive performance trajectory slopes estimated over the course of 5
17 to 8 years in a subset of PREVENT-AD participants ($n = 89$), baseline CSF IGFBP2 levels were
18 negatively correlated with estimated rates of change in delayed memory scores ($R^2 = 0.072$, $\beta = -$
19 0.098 , $P = 0.024$, Fig. 3). Furthermore, baseline CSF IGFBP2 levels were negatively correlated
20 with rates of change in visuospatial constructional abilities ($R^2 = 0.077$, $\beta = -0.082$, $P = 0.019$, Fig.
21 3). However, baseline IGFBP2 levels were not associated with changes in immediate memory (P
22 $= 0.191$), language ($P = 0.332$) or attention ($P = 0.679$) (data not shown).

23

24 **CSF IGFBP2 is associated with cortical atrophy in AD-specific brain regions in** 25 **asymptomatic AD**

26 We analyzed baseline structural neuroimaging data collected from a subset of PREVENT-AD
27 individuals ($n = 104$) in a cross-sectional fashion. Four individuals were omitted from analyses
28 due to failed quality control regarding subject-specific stereotaxic registration and/or brain

1 masking. After adjusting for total intracranial volume (ICV), baseline CSF IGFBP2 was negatively
2 correlated with entorhinal cortex volumes in the left hemisphere at a trend-level ($R^2 = 0.117$, $\beta = -$
3 39.720 , $P = 0.082$, Fig. 4A). However, CSF IGFBP2 was not associated with entorhinal cortex
4 volume in the right hemisphere ($P = 0.962$).

5 Next, we analyzed baseline cortical thickness in pre-specified temporal and parietal brain regions
6 that are vulnerable to early AD pathology. Baseline CSF IGFBP2 was found to be negatively
7 correlated with cortical thickness in the piriform cortex (trend, left hemisphere: $R^2 = 0.121$, $\beta = -$
8 0.628 , $P = 0.064$; right hemisphere: $R^2 = 0.128$, $\beta = -0.709$, $P = 0.039$, Fig. 4B), inferior temporal
9 gyrus (left hemisphere: $R^2 = 0.153$, $\beta = -1.253$, $P = 0.008$; right hemisphere: $P = 0.315$, Fig. 4C),
10 middle temporal gyrus (left hemisphere: $R^2 = 0.144$, $\beta = -1.369$, $P = 0.014$; right hemisphere: $R^2 =$
11 0.116 , $\beta = -0.959$, $P = 0.090$, Fig. 4D), and precuneus (left hemisphere: $P = 0.123$; right
12 hemisphere: $R^2 = 0.131$, $\beta = -1.353$, $P = 0.033$, Supplementary Fig. 2).

13

14 **Changes in CSF IGFBP2 are not caused by alterations of the blood brain** 15 **barrier integrity in asymptomatic subjects**

16 To examine possible blood-brain barrier (BBB) dysfunction and possible peripheral vascular
17 contributions to CSF IGFBP2 levels, we measured microprotein levels, red blood cell count and
18 white blood cell count in the CSF. We did not find any associations between these vascular factors
19 and CSF IGFBP2 (Supplementary Fig. 3); consistent with a relatively intact BBB in asymptomatic
20 PREVENT-AD participants.⁵⁴

21

22 **ADNI-1 cohort**

23 **CSF and plasma IGFBP2 concentrations are elevated in CSF A β (+)/t-tau(+)** 24 **individuals**

25 89 CU individuals and 144 individuals with MCI from ADNI were staged as amyloid and/or tau
26 positive according to recommended CSF A β_{42} and CSF t-tau thresholds of 192 pg/mL and 93
27 pg/mL, respectively (Fig. 5A).^{50,51} The results from the CSF multiplex immunoassay (Fig. 5B)
28 revealed that baseline CSF IGFBP2 levels did not differ between Stages 0 ($n = 80$) A β (-)/t-tau(-)

1 and Stage 1 ($n = 68$) $A\beta(+)/t\text{-tau}(-)$, $P = 0.541$. However, CSF IGFBP2 was significantly elevated
2 at Stage 2 ($n = 77$) $A\beta(+)/t\text{-tau}(+)$ relative to Stage 0 ($P = 0.009$) and Stage 1 ($P = 0.001$). Finally,
3 CSF IGFBP2 was significantly increased in SNAP ($n = 8$) $A\beta(-)/t\text{-tau}(+)$, relative to Stage 0 ($P =$
4 0.010).

5 To reproduce our findings, we performed replication analyses using CSF mass spectrometry data
6 acquired from a subset of the same ADNI participants. These supplementary analyses revealed
7 that both CSF IGFBP2 peptides, HGLYNLK (Supplementary Fig. 4) and LIQGAPTIR
8 (Supplementary Fig. 5) were significantly reduced at Stage 1 ($n = 62$) relative to Stage 0 ($n = 75$),
9 $P = 0.005$ and $P = 0.051$ (trend). Although a similar reduction at Stage 1 was observed with the
10 multiplex immunoassay data, it was not statistically significant. However, similar to the multiplex
11 immunoassay data, both IGFBP2 peptides were significantly elevated at Stage 2 ($n = 72$) relative
12 to Stage 1, $P = 1.50 \times 10^{-5}$ and $P = 2.21 \times 10^{-4}$. Likewise, both IGFBP2 peptides were markedly
13 increased in SNAP ($n = 9$) relative to Stage 0, $P = 4.90 \times 10^{-5}$ and $P = 0.002$, consistent with the
14 immunoassay results.

15 Finally, we staged 58 CU individuals and 196 individuals with MCI from ADNI, that had both
16 baseline plasma IGFBP2 measurements and CSF $A\beta_{42}$ and CSF t-tau measurements (Fig. 5C).
17 Baseline plasma IGFBP2 levels were significantly elevated in Stage 2 ($n = 84$) relative to Stage 0
18 ($n = 98$), $P = 0.001$, and to Stage 1 ($n = 60$), $P = 0.041$ (trend). However, plasma IGFBP2 did not
19 differ between Stage 0 and Stage 1 ($P = 0.208$), or between Stage 0 and SNAP ($n = 12$), $P = 0.874$.

21 **Elevated plasma IGFBP2 is associated with a faster rate of conversion to AD**

22 In the primary analysis for conversion to AD in ADNI, we established baseline plasma IGFBP2
23 threshold values at the 25th percentile (≤ 1.83251 , first quartile, \log_{10} transformed) and above the
24 75th percentile (≥ 2.09342 , fourth quartile, \log_{10} transformed). A total of 226 individuals that were
25 either CU or had MCI were included in these analyses. Of these dementia-free participants, 107
26 individuals eventually met the clinical criteria for a diagnosis of AD (mean follow-up, 3.8 years;
27 range, 0.5-16.5 years). Cox proportional hazards models revealed that individuals with plasma
28 IGFBP2 values greater than the 75th percentile exhibited a faster rate of conversion to AD, than

1 individuals with plasma IGFBP2 values less than the 25th percentile (hazard ratio (HR) 1.616, 95%
2 confidence interval (CI) 1.074-2.430, $P = 0.021$, Fig. 6).

3 In the secondary analysis, baseline IGFBP2 plasma levels were kept as continuous, and 439 ADNI-
4 1 dementia-free participants with plasma IGFBP2 measurements were included. Of these
5 individuals, 214 were eventually diagnosed with AD (mean follow-up, 3.6 years; range, 0.5-16.5
6 years). Similar to the first model, elevated plasma IGFBP2 was associated with a greater rate of
7 conversion to AD (HR 1.857, 95% CI 1.054-3.270, $P = 0.032$).

8

9 **QFP cohort**

10 **Despite reductions in IGFBP2 mRNA, protein levels do not differ in the frontal** 11 **cortex of AD brains**

12 IGFBP2 gene expression was assessed by DNA microarray in the QFP cohort, and demonstrated
13 to be significantly reduced in the frontal cortex of autopsy-confirmed AD brains ($n = 55$),
14 compared to elderly controls ($n = 31$) ($P = 0.049$, Fig. 7A). However, as demonstrated through
15 ELISA, IGFBP2 protein levels did not differ in the frontal cortex of AD cases ($n = 53$) after
16 controlling for total protein levels, relative to controls ($n = 25$, $P = 0.462$, Fig. 7B).

17

18 **Discussion**

19 Our results suggest that nascent AD pathology induces a marked upregulation in IGFBP2, in
20 asymptomatic individuals. CSF and plasma IGFBP2 behave as valuable biomarkers for identifying
21 pre-clinical CSF A β (+)/t-tau(+) individuals, and those with a greater risk of AD conversion.

22 It has been well established that impaired insulin-IGF signalling plays a critical role in AD.⁸⁻¹⁰
23 Furthermore, insulin and IGF proteins are highly expressed in the hippocampus.⁸ Therefore, it is
24 possible that impairments in insulin-IGF signaling may account for the selective vulnerability of
25 the hippocampal formation to nascent AD pathology. Hence, targeting the insulin-IGF system may
26 offer a promising solution to delay, slow down and/or prevent AD, either alone or in combination
27 therapies. However, to develop these therapies, unraveling the molecular intricacies of the insulin-

1 IGF system is necessary. To this end, we investigated the role of the most abundant IGF-binding
2 protein in the CSF, IGFBP2,^{28,29} during the earliest possible asymptomatic stage of AD, in “at-
3 risk”, parental history-positive PREVENT-AD participants.

4 We observed a positive relationship between IGFBP2 and CSF A β ₄₂ (Fig. 2A), which is consistent
5 with the proposed IGF-mediated clearance of A β .^{12,13} This notion is in agreement with the finding
6 that current A β -lowering therapies induce an increase in CSF A β ₄₂ in participants with MCI or
7 mild AD.⁵⁵ Furthermore, our data suggest IGFBP2 may be upregulated as a result of early neuronal
8 loss in the asymptomatic stage of the disease (Fig. 2C); which is consistent with the extensive
9 literature regarding the upregulation of IGFs and IGFBPs following several rodent models of brain
10 damage and recovery.⁵⁶⁻⁶³ Furthermore, the administration of des-IGF-I, an analogue of IGF-I
11 with a low affinity for IGFBPs, failed to attenuate neuronal cell death in mice with experimental
12 hypoxic ischemic injuries.⁶⁴ However, the administration of IGFBP-compatible IGF-I
13 significantly reduced the observed neuronal loss.⁶⁴

14 Our findings also suggest that IGFBP2 may be modulated by early CSF p₁₈₁-tau production (Fig.
15 2B) and early deposition in asymptomatic individuals. PET imaging analyses revealed that
16 IGFBP2 is positively associated (at a trend-level) with significant tau deposition in the entorhinal
17 cortex (Fig. 2E) and lingual gyrus (Supplementary Fig. 1), features that are typically associated
18 with early Braak Stages 2-3.⁶⁵ These findings are certainly consistent with the fact that insulin and
19 IGFs normally inhibit glycogen synthase kinase-3 (GSK3) activity and phospho-tau production in
20 human neurons, through the phosphoinositide 3-kinase-protein kinase B (PI3K-AKT) signaling
21 pathway.^{14,15} For instance, the pharmacological inhibition of GSK3 has been linked to increases in
22 IGF-I in the rodent brain,⁶⁶ whereas conditional transgenic mice overexpressing GSK3 in the
23 cortex and hippocampus display increased tau phosphorylation in AD relevant epitopes,⁶⁷
24 degeneration of the dentate gyrus⁶⁸ and spatial memory impairments.⁶⁹ Finally, mice
25 overexpressing IGFBP2 display an increase in AKT activity, a GSK3 inhibitor, in the brain.⁷⁰
26 Thus, it is tempting to postulate that IGF regulation of GSK3 activity in turn modulates IGFBP2
27 production via a phospho-tau mediated process that ensures some form of local autoregulation
28 and/or resilience.

29 Given the prominent loss of synapses in AD, we examined the relationship between IGFBP2 and
30 synaptic proteins in the CSF, namely SNAP25 (Fig. 2F), SYT1 (Fig. 2G), GAP43 (Fig. 2H) and

1 NRGN (Fig. 2I).⁴⁰⁻⁴⁵ The present study's results suggest that synaptic dysfunction and loss may
2 trigger an increase in IGFBP2 synthesis and secretion. This finding is in agreement with the
3 upregulation and regenerative abilities of IGFs to grow axons during the development of the CNS
4 and regrow axons during repair following injury to the CNS and/or PNS.^{21,71-76}

5 Considering the relationship between CSF synaptic markers and CSF IGFBP2, we were interested
6 in examining the relationship between changes in cognition and baseline CSF IGFBP2 in
7 PREVENT-AD. We have demonstrated that CSF IGFBP2 levels are associated with cognitive
8 decline in RBANS delayed memory and visuospatial abilities over a 5-8-year period, in a subset
9 of PREVENT-AD participants (Fig. 3). This finding is consistent with a report that plasma levels
10 of IGFBP2 were negatively correlated with episodic memory performances in participants from
11 the ADNI cohort.⁷⁷ Thus, given the critical role of the hippocampus in the consolidation of
12 declarative memory and in spatial processing, our data provide compelling evidence that IGFBP2
13 plays a pivotal role in the integrity of the hippocampal formation, which displays elevated
14 expression levels of insulin, IGFs and their receptors – compared to other brain regions, such as
15 the frontal cortex.⁸ Indeed, plasma IGFBP2 levels have been demonstrated to be negatively
16 correlated with hippocampal volumes in amyloid-negative individuals from the ADNI cohort.⁷⁷
17 Consistent with this view, IGFBP2 has been demonstrated to be expressed by neurons and
18 astrocytes in the hippocampus during development and following CNS injury.^{57-59,62,78,79}
19 Furthermore, the relationship between IGFBP2, delayed memory and hippocampal structure is
20 consistent with the finding that the administration of IGFBP2 has been shown to increase the
21 number of dendritic spines in the dentate gyrus of rodent models of post-traumatic stress disorder.⁸⁰
22 In a similar fashion, in cell culture experiments, antibodies targeted against IGFBP2 have been
23 demonstrated to inhibit neurogenesis – a phenomenon that occurs in the dentate gyrus.⁸¹ Moreover,
24 *IGFBP2* knockout mice exhibit deficits in long-term potentiation as well as impaired performances
25 on the Morris water maze, which heavily relies on the hippocampus.⁷⁹ Finally, the administration
26 of an IGFBP2-derived peptide has been demonstrated to rescue deficits in synaptic plasticity,
27 memory and learning in a mouse model of *SHANK3*-mediated post-synaptic deficits.⁸² Overall,
28 these findings suggest that IGFBP2 in certain circumstances is linked to neuroprotection and
29 hippocampal-mediated cognitive abilities.

30 Indeed, it is important to take into consideration the early involvement of IGFBP2 in the
31 presymptomatic phase of AD, when presumably resilience is a major player in the brain response

1 to early neurodegeneration. For instance, several neuroprotective genes have been demonstrated
2 to be upregulated by the (soluble) α -secretase cleaved fragment of amyloid precursor protein
3 (sAPP α), amongst them, *IGFBP2* and *IGF2*.⁸³ Thus, it is possible that an increase in binding of
4 IGFBP2 to IGFs may attenuate IGF degradation and/or promote the targeting of IGFs to their
5 receptors, thus enhancing synaptic and terminal resilience in face of the emerging
6 neurodegenerative process. This receptor targeting hypothesis is notably supported by evidence
7 that transgenic mice overexpressing IGFBP2 lacking a proteoglycan-binding domain exhibit
8 severe reductions in synaptic markers and hippocampal weight.⁸⁴ Finally, in a pilot study
9 conducted on AD brains, temporal cortex IGFBP2 levels were negatively correlated with senile
10 plaque levels – suggesting a role for IGFBP2 in neuroprotection and resilience.⁹

11 Given the negative relationship between IGFBP2 and cognitive abilities, we were prompted to
12 examine whether baseline levels of CSF IGFBP2 were associated with anatomical changes in the
13 brain. Interestingly, in asymptomatic PREVENT-AD participants, IGFBP2 was associated with
14 atrophy in the left entorhinal cortex, at a trend-level (Fig. 4A). This is a critical finding, as the
15 entorhinal cortex is the first region that is affected in AD.^{65,85,86} Furthermore, IGFBP2 was
16 associated with cortical thinning in several pre-specified temporal and parietal brain regions that
17 are known to be affected early on in AD, such as the piriform cortex (Fig. 4B), inferior (Fig. 4C)
18 and middle temporal gyri (Fig. 4D), and precuneus (Supplementary Fig. 2).^{87,88} Overall, the
19 structural neuroimaging results are in agreement with previous reports of IGFBP2 being associated
20 with atrophy in AD-associated brain regions.^{36,89,90}

21 Finally, in the PREVENT-AD cohort, changes in CSF IGFBP2 appear to be mostly specific to
22 changes in the CNS, as we did not detect any significant peripheral vascular contributions or blood
23 brain barrier dysfunction in these asymptomatic individuals (Supplementary Fig. 3).⁵⁴

24 To independently validate our observations, we further analyzed data from a well-characterized
25 cohort of CU individuals and individuals with MCI from ADNI-1 (Fig. 5A). Our results suggest
26 that elevated CSF (Fig. 5B, Supplementary Fig. 4, Supplementary Fig. 5) and plasma IGFBP2
27 (Fig. 5C) may be valuable biomarkers of CSF A β (+)/t-tau(+) individuals and thus, facilitate
28 screening for suitable patients for clinical trials.^{50,51} This is not the first time that we identified a
29 strong biphasic response as individuals progress from the CSF A β (+)/t-tau(-) stage to the CSF
30 A β +/t-tau(+) stage (Supplementary Fig. 4, Supplementary Fig. 5). Certain inflammatory proteins

1 such as IL-8, IL-12 and IL-15 display the same initial reduction in the CSF A β (+)/t-tau(-) stage
2 followed by marked increases in the CSF A β (+)/t-tau(+) stage.⁹¹ These results support recent
3 observations that immune activation may become apparent only after the onset of both amyloid
4 and tau pathologies. Unexpectedly, these results also suggest that immune marker activity and
5 IGFBP2 may diminish upon the earliest appearance of amyloid plaque pathology. However, as
6 IGFBP2 regulates IGF signaling in neurons, one might speculate that an increase in binding of
7 IGFBP2 to a limited amount of IGFs may block IGF-mediated suppression of tau phosphorylation,
8 leading to increased levels of phospho-tau and ultimately, promoting tau deposition, neuronal
9 damage and death.

10 Consistent with this working model and upon conducting survival analyses, we found that elevated
11 circulating IGFBP2 levels were associated with a pronounced rate of conversion to AD in the
12 ADNI cohort (Fig. 6). Overall, our results are consistent with the existing literature that plasma
13 IGFBP2 has been associated with a greater risk of developing AD.^{35,37,38} Finally, in contrast to the
14 PREVENT-AD cohort, CSF IGFBP2 immunoassay measurements have been previously found to
15 correlate with plasma IGFBP2 levels in the ADNI cohort – which incorporates individuals with a
16 disrupted BBB.⁷⁷

17 Finally, given the regional-specificity of the insulin-IGF system, we further examined IGFBP2
18 levels in the human brain.⁸ IGFBP2 mRNA levels were reduced in the frontal cortex of autopsy-
19 confirmed AD brains from the QFP cohort (Fig. 7A). However, cortical IGFBP2 protein levels did
20 not differ between AD brains and elderly controls (Fig. 7B). Thus, our results provide further
21 evidence that support the (hippocampal) regional specificity of the insulin-IGF system, potentially
22 including the IGFBP2 protein.⁸ However, due to the scarcity of medial temporal lobe tissues, we
23 were unable to confirm the regional specificity of the IGFBP2 protein in the QFP cohort.
24 Nevertheless, our results differ from those of a pilot study, that found IGFBP2 was decreased in
25 the temporal cortex of AD brains.⁹ It is possible our results differ since we measured cytoplasmic
26 and extracellular IGFBP2, whilst the pilot study measured membrane-bound IGFBP2.⁹ Finally,
27 although IGFBP2 has been demonstrated to contain a nuclear localization signal and regulate the
28 expression of several genes such as vascular endothelial growth factor, we were unable to
29 differentiate between IGF-dependent and IGF-independent IGFBP2.⁹²

1 Overall, our data suggest that IGFBP2 may play a critical role in neuroprotection during the
2 asymptomatic stage of AD. However, as amyloid and tau pathology accumulate, the involvement
3 of IGFBP2 appears to notably change. At that stage, elevated concentrations of IGFBP2 are
4 associated with an accelerated rate of MCI to AD conversion in the ADNI cohort, and with subtle
5 declines in delayed memory and visuospatial abilities in asymptomatic PREVENT-AD
6 participants.

7 In the larger context of the insulin-IGF system, it will be exciting to follow the results of ongoing
8 phase 3 clinical trials involving the administration of semaglutide (Ozempic), a glucagon-like
9 peptide-1 receptor agonist that is clinically approved for the treatment of type 2 diabetes.⁹³
10 Considering the role of semaglutide in stimulating insulin secretion and signalling,⁹³ exploring
11 therapeutic strategies that enhance IGF signalling in the brain,⁹⁴ perhaps through IGFBP2, may
12 also warrant further investigation in presymptomatic AD. At the moment, we favor the hypothesis
13 that an anti-IGFBP2 therapy may be effective in amyloid positive individuals whose tau deposition
14 is minimal. As tau pathology emerges, it may prove difficult to change this complex metabolic
15 process as IGFBP2 activity appears less protective and more reactive, similar to many other
16 immune regulators.

17

18 **Data availability**

19 Data pertaining to the PREVENT-AD cohort can be downloaded from data release 6.0 at
20 <https://openpreventad.loris.ca/>. CSF, plasma, genetic and clinical data from the ADNI-1 cohort
21 were downloaded from the ADNI website (<http://adni.loni.usc.edu/>). Data collected from the QFP
22 cohort are not publicly available, however the data are available from the corresponding author
23 upon reasonable request.

24

25 **Acknowledgements**

26 The authors would like to thank Dr. Naguib Mechawar at the Douglas Institute/Bell Canada Brain
27 Bank for providing human brain tissues from the Quebec Founder Population. We also wish to
28 thank Mrs. Jennifer Tremblay-Mercier, Marie-Elyse Lafaille-Magnan and Melissa Savard as well

1 as Drs. Pedro Rosa-Neto, Daniel Auld and David Lafontaine for their technical expertise. Data
2 used in preparation of this article were obtained from the Alzheimer's Disease Neuroimaging
3 Initiative (ADNI) database (adni.loni.usc.edu). As such, the investigators within the ADNI
4 contributed to the design and implementation of ADNI and/or provided data but did not participate
5 in analysis or writing of this report. A complete listing of ADNI investigators can be found
6 at:http://adni.loni.usc.edu/wpcontent/uploads/how_to_apply/ADNI_Acknowledgement_List.pdf.
7 Data used in preparation of this article were obtained from the PRE-symptomatic EVALuation of
8 Experimental or Novel Treatments for Alzheimer's Disease (PREVENT-AD) program at the
9 Centre for Studies on Prevention of Alzheimer's Disease (StoP-AD), Douglas Mental Health
10 University Institute Research Centre (<http://douglas.research.mcgill.ca/stop-ad-centre>). A
11 complete listing of the PREVENT-AD Research Group can be found at:
12 <https://preventad.loris.ca/acknowledgements/acknowledgements.php?date=2023-05-01>.

14 **Funding**

15 JP is supported by the Fonds de recherche du Québec-Santé (FRQS), the Canadian Institutes of
16 Health Research (CIHR #PJT 153287) and the J.L. Levesque Foundation. SV is supported by a
17 Canada Research Chair and a Canada Fund for Innovation grant, the FRQS, the CIHR, Brain
18 Canada, McGill University and the Alzheimer's Association. HZ is a Wallenberg Scholar
19 supported by grants from the Swedish Research Council (#2022-01018 and #2019-02397), the
20 European Union's Horizon Europe research and innovation programme under grant agreement No
21 101053962, Swedish State Support for Clinical Research (#ALFGBG-71320), the Alzheimer Drug
22 Discovery Foundation (ADDF), USA (#201809-2016862), the AD Strategic Fund and the
23 Alzheimer's Association (#ADSF-21-831376-C, #ADSF-21-831381-C, and #ADSF-21-831377-
24 C), the Bluefield Project, the Olav Thon Foundation, the Erling-Persson Family Foundation,
25 Stiftelsen för Gamla Tjänarinnor, Hjärfonden, Sweden (#FO2022-0270), the European Union's
26 Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant
27 agreement No 860197 (MIRIADE), the European Union Joint Programme – Neurodegenerative
28 Disease Research (JPND2021-00694), the National Institute for Health and Care Research
29 University College London Hospitals Biomedical Research Centre, and the UK Dementia
30 Research Institute at UCL (UKDRI-1003). KB is supported by the Swedish Research Council

1 (#2017-00915 and #2022-00732), the Swedish Alzheimer Foundation (#AF-930351, #AF-939721
2 and #AF-968270), Hjärnfonden, Sweden (#FO2017-0243 and #ALZ2022-0006), the Swedish state
3 under the agreement between the Swedish government and the County Councils, the ALF-
4 agreement (#ALFGBG-715986 and #ALFGBG-965240), the European Union Joint Program for
5 Neurodegenerative Disorders (JPND2019-466-236), the Alzheimer's Association 2021 Zenith
6 Award (ZEN-21-848495), and the Alzheimer's Association 2022-2025 Grant (SG-23-1038904
7 QC). MJQ is supported by the FRQS.

8 PREVENT-AD was launched in 2011 as a \$13.5 million, 7-year public-private partnership using
9 funds provided by McGill University, FRQS, an unrestricted research grant from Pfizer Canada,
10 the Levesque Foundation, the Douglas Hospital Research Centre and Foundation, the Government
11 of Canada, and the Canada Fund for Innovation. Private sector contributions are facilitated by the
12 Development Office of the McGill University Faculty of Medicine and by the Douglas Hospital
13 Research Centre Foundation (<http://www.douglas.qc.ca/>).

14 Data collection and sharing for this project was funded by the ADNI (National Institutes of Health
15 Grant U01 AG024904) and DOD ADNI (Department of Defense award number W81XWH-12-2-
16 0012). ADNI is funded by the National Institute on Aging, the National Institute of Biomedical
17 Imaging and Bioengineering, and through generous contributions from the following: AbbVie,
18 Alzheimer's Association; Alzheimer's Drug Discovery Foundation; Araclon Biotech; BioClinica,
19 Inc.; Biogen; Bristol-Myers Squibb Company; CereSpir, Inc.; Cogstate; Eisai Inc.; Elan
20 Pharmaceuticals, Inc.; Eli Lilly and Company; EuroImmun; F. Hoffmann-La Roche Ltd and its
21 affiliated company Genentech, Inc.; Fujirebio; GE Healthcare; IXICO Ltd.; Janssen Alzheimer
22 Immunotherapy Research & Development, LLC.; Johnson & Johnson Pharmaceutical Research &
23 Development LLC.; Lumosity; Lundbeck; Merck & Co., Inc.; Meso Scale Diagnostics, LLC.;
24 NeuroRx Research; Neurotrack Technologies; Novartis Pharmaceuticals Corporation; Pfizer Inc.;
25 Piramal Imaging; Servier; Takeda Pharmaceutical Company; and Transition Therapeutics. The
26 CIHR is providing funds to support ADNI clinical sites in Canada. Private sector contributions are
27 facilitated by the Foundation for the National Institutes of Health (www.fnih.org). The grantee
28 organization is the Northern California Institute for Research and Education, and the study is
29 coordinated by the Alzheimer's Therapeutic Research Institute at the University of Southern
30 California. ADNI data are disseminated by the Laboratory for Neuro Imaging at the University of
31 Southern California.

1

2 **Competing interests**

3 JP serves as a scientific advisor to the Alzheimer Society of France. HZ has served at scientific
4 advisory boards and/or as a consultant for Abbvie, Acumen, Alector, Alzinova, ALZPath,
5 Annexon, Apellis, Artery Therapeutics, AZTherapies, CogRx, Denali, Eisai, Nervgen, Novo
6 Nordisk, Optoceutics, Passage Bio, Pinteon Therapeutics, Prothena, Red Abbey Labs, reMYND,
7 Roche, Samumed, Siemens Healthineers, Triplet Therapeutics, and Wave, has given lectures in
8 symposia sponsored by Celectricon, Fujirebio, Alzecure, Biogen, and Roche, and is a co-founder
9 of Brain Biomarker Solutions in Gothenburg AB (BBS), which is a part of the GU Ventures
10 Incubator Program (outside submitted work). KB has served as a consultant and at advisory boards
11 for Acumen, ALZPath, BioArctic, Biogen, Eisai, Lilly, Moleac Pte. Ltd, Novartis, Ono Pharma,
12 Prothena, Roche Diagnostics, and Siemens Healthineers; has served at data monitoring committees
13 for Julius Clinical and Novartis; has given lectures, produced educational materials and
14 participated in educational programs for AC Immune, Biogen, Celdara Medical, Eisai and Roche
15 Diagnostics; and is a co-founder of Brain Biomarker Solutions in Gothenburg AB (BBS), which
16 is a part of the GU Ventures Incubator Program, outside the work presented in this paper.

17

18 **Supplementary material**

19 Supplementary material is available at *Brain* online.

20

21 **References**

- 22 1. GBD 2019 Dementia Forecasting Collaborators. Estimation of the global prevalence of
23 dementia in 2019 and forecasted prevalence in 2050: an analysis for the Global Burden of
24 Disease Study 2019. *Lancet Public Health*. 2022;7(2):e105-e125. doi:10.1016/S2468-
25 2667(21)00249-8

- 1 2. Nandi A, Counts N, Chen S, *et al.* Global and regional projections of the economic burden
2 of Alzheimer's disease and related dementias from 2019 to 2050: A value of statistical life
3 approach. *EClinicalMedicine*. 2022;51:101580. doi:10.1016/j.eclinm.2022.101580
- 4 3. Alzheimer's Association. 2022 Alzheimer's disease facts and figures. *Alzheimers Dement*.
5 2022;18(4):700-789. doi:10.1002/alz.12638
- 6 4. Serrano-Pozo A, Frosch MP, Masliah E, Hyman BT. Neuropathological alterations in
7 Alzheimer disease. *Cold Spring Harbor Perspect Med*. 2011;1(1):a006189.
8 doi:10.1101/cshperspect.a006189
- 9 5. Jack CR Jr, Knopman DS, Jagust WJ, *et al.* Hypothetical model of dynamic biomarkers of
10 the Alzheimer's pathological cascade. *Lancet Neurol*. 2010;9(1):119-128.
11 doi:10.1016/S1474-4422(09)70299-6
- 12 6. Arvanitakis Z, Wilson RS, Bienias JL, Evans DA, Bennett DA. Diabetes mellitus and risk
13 of Alzheimer disease and decline in cognitive function. *Arch Neurol*. 2004;61(5):661-666.
14 doi:10.1001/archneur.61.5.661
- 15 7. Arnold SE, Arvanitakis Z, Macauley-Rambach SL, *et al.* Brain insulin resistance in type 2
16 diabetes and Alzheimer disease: concepts and conundrums. *Nat Rev Neurol*.
17 2018;14(3):168-181. doi:10.1038/nrneurol.2017.185
- 18 8. Steen E, Terry BM, J Rivera E, *et al.* Impaired insulin and insulin-like growth factor
19 expression and signaling mechanisms in Alzheimer's disease—is this type 3 diabetes?. *J*
20 *Alzheimers Dis*. 2005;7(1):63-80. doi:10.3233/jad-2005-7107
- 21 9. Moloney AM, Griffin RJ, Timmons S, O'Connor R, Ravid R, O'Neill C. Defects in IGF-
22 1 receptor, insulin receptor and IRS-1/2 in Alzheimer's disease indicate possible resistance
23 to IGF-1 and insulin signalling. *Neurobiol Aging*. 2010;31(2):224-243.
24 doi:10.1016/j.neurobiolaging.2008.04.002
- 25 10. Talbot K, Wang HY, Kazi H, *et al.* Demonstrated brain insulin resistance in Alzheimer's
26 disease patients is associated with IGF-1 resistance, IRS-1 dysregulation, and cognitive
27 decline. *J Clin Invest*. 2012;122(4):1316-1338. doi:10.1172/JCI59903
- 28 11. Kar S, Poirier J, Guevara J, *et al.* Cellular distribution of insulin-like growth factor-
29 II/mannose-6-phosphate receptor in normal human brain and its alteration in Alzheimer's

- 1 disease pathology. *Neurobiol Aging*. 2006;27(2):199-210.
2 doi:10.1016/j.neurobiolaging.2005.03.005
- 3 12. Farris W, Mansourian S, Chang Y, *et al*. Insulin-degrading enzyme regulates the levels of
4 insulin, amyloid β -protein, and the β -amyloid precursor protein intracellular domain in
5 vivo. *Proc Natl Acad Sci U S A*. 2003;100(7):4162-4167. doi:10.1073/pnas.0230450100
- 6 13. Carro E, Trejo JL, Gomez-Isla T, LeRoith D, Torres-Aleman I. Serum insulin-like growth
7 factor I regulates brain amyloid- β levels. *Nat Med*. 2002;8(12):1390-1397.
8 doi:10.1038/nm1202-793
- 9 14. Hong M, Lee VM. Insulin and insulin-like growth factor-1 regulate tau phosphorylation in
10 cultured human neurons. *J Biol Chem*. 1997;272(31):19547-19553.
11 doi:10.1074/jbc.272.31.19547
- 12 15. Schubert M, Brazil DP, Burks DJ, *et al*. Insulin receptor substrate-2 deficiency impairs
13 brain growth and promotes tau phosphorylation. *J Neurosci*. 2003;23(18):7084-7092.
14 doi:10.1523/JNEUROSCI.23-18-07084.2003
- 15 16. De Felice FG, Vieira MN, Bomfim TR, *et al*. Protection of synapses against Alzheimer's-
16 linked toxins: insulin signaling prevents the pathogenic binding of A β oligomers. *Proc Natl*
17 *Acad Sci U S A*. 2009;106(6):1971-1976. doi:10.1073/pnas.0809158106
- 18 17. Doré S, Kar S, Quirion R. Insulin-like growth factor I protects and rescues hippocampal
19 neurons against beta-amyloid- and human amylin-induced toxicity. *Proc Natl Acad Sci U*
20 *S A*. 1997;94(9):4772-4777. doi:10.1073/pnas.94.9.4772
- 21 18. Willette AA, Bendlin BB, Starks EJ, *et al*. Association of Insulin Resistance With Cerebral
22 Glucose Uptake in Late Middle-Aged Adults at Risk for Alzheimer Disease. *JAMA Neurol*.
23 2015;72(9):1013-1020. doi:10.1001/jamaneurol.2015.0613
- 24 19. Reger MA, Watson GS, Green PS, *et al*. Intranasal insulin improves cognition and
25 modulates beta-amyloid in early AD. *Neurology*. 2008;70(6):440-448.
26 doi:10.1212/01.WNL.0000265401.62434.36
- 27 20. Craft S, Baker LD, Montine TJ, *et al*. Intranasal insulin therapy for Alzheimer disease and
28 amnesic mild cognitive impairment: a pilot clinical trial. *Arch Neurol*. 2012;69(1):29-38.
29 doi:10.1001/archneurol.2011.233

- 1 21. O'Kusky JR, Ye P, D'Ercole AJ. Insulin-like growth factor-I promotes neurogenesis and
2 synaptogenesis in the hippocampal dentate gyrus during postnatal development. *J*
3 *Neurosci.* 2000;20(22):8435-8442. doi:10.1523/JNEUROSCI.20-22-08435.2000
- 4 22. Russo VC, Gluckman PD, Feldman EL, Werther GA. The insulin-like growth factor
5 system and its pleiotropic functions in brain. *Endocr Rev.* 2005;26(7):916-43.
6 doi:10.1210/er.2004-0024
- 7 23. Allard JB, Duan C. IGF-binding proteins: why do they exist and why are there so many?.
8 *Front Endocrinol (Lausanne).* 2018;9:117. doi:10.3389/fendo.2018.00117
- 9 24. Hwa V, Oh Y, Rosenfeld RG. The insulin-like growth factor-binding protein (IGFBP)
10 superfamily. *Endocr Rev.* 1999;20(6):761-787. doi:10.1210/edrv.20.6.0382
- 11 25. Russo VC, Bach LA, Fosang AJ, Baker NL, Werther GA. Insulin-like growth factor
12 binding protein-2 binds to cell surface proteoglycans in the rat brain olfactory bulb.
13 *Endocrinology.* 1997;138(11):4858-4867. doi:10.1210/endo.138.11.5472
- 14 26. Schütt BS, Langkamp M, Rauschnabel U, Ranke MB, Elmlinger MW. Integrin-mediated
15 action of insulin-like growth factor binding protein-2 in tumor cells. *J Mol Endocrinol.*
16 2004;32(3):859-868. doi:10.1677/jme.0.0320859
- 17 27. Russo VC, Rekaris G, Baker NL, Bach LA, Werther GA. Basic fibroblast growth factor
18 induces proteolysis of secreted and cell membrane-associated insulin-like growth factor
19 binding protein-2 in human neuroblastoma cells. *Endocrinology.* 1999;140(7):3082-3090.
20 doi:10.1210/endo.140.7.6771
- 21 28. Roghani M, Lassarre C, Zapf J, Pova G, Binoux M. Two insulin-like growth factor (IGF)-
22 binding proteins are responsible for the selective affinity for IGF-II of cerebrospinal fluid
23 binding proteins. *J Clin Endocrinol Metab.* 1991;73(3):658-666. doi:10.1210/jcem-73-3-
24 658
- 25 29. Ocrant I, Fay CT, Parmelee JT. Characterization of insulin-like growth factor binding
26 proteins produced in the rat central nervous system. *Endocrinology.* 1990;127(3):1260-
27 1267. doi:10.1210/endo-127-3-1260
- 28 30. Tham A, Nordberg A, Grissom FE, Carlsson-Skwirut C, Viitanen M, Sara VR. Insulin-like
29 growth factors and insulin-like growth factor binding proteins in cerebrospinal fluid and

- 1 serum of patients with dementia of the Alzheimer type. *J Neural Transm Park Dis Dement*
2 *Sect.*1993;5(3):165-176. doi:10.1007/BF02257671
- 3 31. Salehi Z, Mashayekhi F, Naji M. Insulin like growth factor-1 and insulin like growth factor
4 binding proteins in the cerebrospinal fluid and serum from patients with Alzheimer's
5 disease. *Biofactors.* 2008;33(2):99-106. doi:10.1002/biof.5520330202
- 6 32. Hertze J, Nägga K, Minthon L, Hansson O. Changes in cerebrospinal fluid and blood
7 plasma levels of IGF-II and its binding proteins in Alzheimer's disease: an observational
8 study. *BMC Neurol.* 2014;14:64. doi:10.1186/1471-2377-14-64
- 9 33. Åberg D, Johansson P, Isgaard J, *et al.* Increased cerebrospinal fluid level of insulin-like
10 growth factor-II in male patients with Alzheimer's disease. *J Alzheimers Dis.*
11 2015;48(3):637-646. doi:10.3233/JAD-150351
- 12 34. O'Bryant SE, Xiao G, Barber R, *et al.* A serum protein-based algorithm for the detection
13 of Alzheimer disease. *Arch Neurol.* 2010;67(9):1077-1081.
14 doi:10.1001/archneurol.2010.215
- 15 35. Doecke JD, Laws SM, Faux NG, *et al.* Blood-based protein biomarkers for diagnosis of
16 Alzheimer disease. *Arch Neurol.* 2012;69(10):1318-1325.
17 doi:10.1001/archneurol.2012.1282
- 18 36. Bonham LW, Geier EG, Steele NZ, *et al.* Insulin-like growth factor binding protein 2 is
19 associated with biomarkers of Alzheimer's disease pathology and shows differential
20 expression in transgenic mice. *Front Neurosci.* 2018;12:476.
21 doi:10.3389/fnins.2018.00476
- 22 37. McGrath ER, Himali JJ, Levy D, *et al.* Circulating IGFBP-2: a novel biomarker for incident
23 dementia. *Ann Clin Transl Neurol.* 2019;6(9):1659-1670. doi:10.1002/acn3.50854
- 24 38. Rocha de Paula M, Gómez Ravetti M, Berretta R, Moscato P. Differences in abundances
25 of cell-signalling proteins in blood reveal novel biomarkers for early detection of clinical
26 Alzheimer's disease. *PLoS One.* 2011;6(3):e17481. doi:10.1371/journal.pone.0017481
- 27 39. Tremblay-Mercier J, Madjar C, Das S, *et al.* Open science datasets from PREVENT-AD,
28 a longitudinal cohort of pre-symptomatic Alzheimer's disease. *Neuroimage Clin.*
29 2021;31:102733. doi:10.1016/j.nicl.2021.102733

- 1 40. Brinkmalm A, Brinkmalm G, Honer WG, *et al.* SNAP-25 is a promising novel
2 cerebrospinal fluid biomarker for synapse degeneration in Alzheimer's disease. *Mol*
3 *Neurodegener.* 2014;9:53. doi:10.1186/1750-1326-9-53
- 4 41. Öhrfelt A, Brinkmalm A, Dumurgier J, *et al.* The pre-synaptic vesicle protein
5 synaptotagmin is a novel biomarker for Alzheimer's disease. *Alzheimers Res Ther.*
6 2016;8(1):41. doi:10.1186/s13195-016-0208-8
- 7 42. Tible M, Sandelius Å, Höglund K, *et al.* Dissection of synaptic pathways through the CSF
8 biomarkers for predicting Alzheimer disease. *Neurology.* 2020;95(8):e953-e961.
9 doi:10.1212/WNL.0000000000010131
- 10 43. Xu C, Sellgren CM, Fatouros-Bergman H, *et al.* CSF levels of synaptosomal-associated
11 protein 25 and synaptotagmin-1 in first-episode psychosis subjects. *IBRO Rep.*
12 2020;8:136-142. doi:10.1016/j.ibror.2020.04.001
- 13 44. Sandelius Å, Portelius E, Källén Å, *et al.* Elevated CSF GAP-43 is Alzheimer's disease
14 specific and associated with tau and amyloid pathology. *Alzheimers Dement.*
15 2019;15(1):55-64. doi:10.1016/j.jalz.2018.08.006
- 16 45. Portelius E, Olsson B, Höglund K, *et al.* Cerebrospinal fluid neurogranin concentration in
17 neurodegeneration: relation to clinical phenotypes and neuropathology. *Acta Neuropathol.*
18 2018;136(3):363-376. doi:10.1007/s00401-018-1851-x
- 19 46. McSweeney M, Binette AP, Meyer PF, *et al.* Intermediate flortaucipir uptake is associated
20 with A β -PET and CSF tau in asymptomatic adults. *Neurology.* 2020;94(11):e1190-1200.
21 doi:10.1212/WNL.0000000000008905
- 22 47. Zijdenbos AP, Forghani R, Evans AC. Automatic "pipeline" analysis of 3-D MRI data for
23 clinical trials: application to multiple sclerosis. *IEEE Trans Med Imaging.*
24 2002;21(10):1280–1291. doi:10.1109/TMI.2002.806283
- 25 48. Aubert-Broche B, Fonov VS, García-Lorenzo D, *et al.* A new method for structural volume
26 analysis of longitudinal brain MRI data and its application in studying the growth
27 trajectories of anatomical brain structures in childhood. *Neuroimage.* 2013;82:393-402.
28 doi:10.1016/j.neuroimage.2013.05.065

- 1 49. Randolph C, Tierney MC, Mohr E, Chase TN. The Repeatable Battery for the Assessment
2 of Neuropsychological Status (RBANS): preliminary clinical validity. *J Clin Exp*
3 *Neuropsychol.* 1998;20(3):310-319. doi:10.1076/jcen.20.3.310.823
- 4 50. Jack CR Jr, Bennett DA, Blennow K, *et al.* NIA-AA research framework: toward a
5 biological definition of Alzheimer's disease. *Alzheimers Dement.* 2018;14(4):535-562.
6 doi:10.1016/j.jalz.2018.02.018
- 7 51. Shaw LM, Vanderstichele H, Knapik-Czajka M, *et al.* Cerebrospinal fluid biomarker
8 signature in Alzheimer's disease neuroimaging initiative subjects. *Ann Neurol.*
9 2009;65(4):403-413. doi:10.1002/ana.21610
- 10 52. Scriver CR. Human genetics: lessons from Quebec populations. *Annu Rev Genomics Hum*
11 *Genet.* 2001;2:69-101. doi:10.1146/annurev.genom.2.1.69
- 12 53. McKhann G, Drachman D, Folstein M, Katzman R, Price D, Stadlan EM. Clinical
13 diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA Work Group* under
14 the auspices of Department of Health and Human Services Task Force on Alzheimer's
15 Disease. *Neurology.* 1984;34(7):939-944. doi:10.1212/wnl.34.7.939
- 16 54. Picard C, Nilsson N, Labonté A, *et al.* Apolipoprotein B is a novel marker for early tau
17 pathology in Alzheimer's disease. *Alzheimers Dement.* 2022;18(5):875-887.
18 doi:10.1002/alz.12442
- 19 55. van Dyck CH, Swanson CJ, Aisen P, *et al.* Lecanemab in Early Alzheimer's Disease. *N*
20 *Engl J Med.* 2023;388(1):9-21. doi:10.1056/NEJMoa2212948
- 21 56. Gluckman P, Klempt N, Guan J, *et al.* A role for IGF-1 in the rescue of CNS neurons
22 following hypoxic-ischemic injury. *Biochem Biophys Res Commun.* 1992;182(2):593-599.
23 doi:10.1016/0006-291x(92)91774-k
- 24 57. Klempt ND, Klempt M, Gunn AJ, Singh K, Gluckman PD. Expression of insulin-like
25 growth factor-binding protein 2 (IGF-BP 2) following transient hypoxia-ischemia in the
26 infant rat brain. *Brain Res Mol Brain Res.* 1992;15(1-2):55-61. doi:10.1016/0169-
27 328x(92)90151-z

- 1 58. Beilharz EJ, Russo VC, Butler G, *et al.* Co-ordinated and cellular specific induction of the
2 components of the IGF/IGFBP axis in the rat brain following hypoxic–ischemic injury.
3 *Brain Res Mol Brain Res.* 1998;59(2):119-134. doi:10.1016/s0169-328x(98)00122-3
- 4 59. Sandberg Nordqvist AC, Von Holst H, Holmin S, Sara VR, Bellander BM, Schalling M.
5 Increase of insulin-like growth factor (IGF)-1, IGF binding protein-2 and– 4 mRNAs
6 following cerebral contusion. *Brain Res Mol Brain Res.* 1996;38(2):285-293.
7 doi:10.1016/0169-328x(95)00346-t
- 8 60. Walter HJ, Berry M, Hill DJ, Logan A. Spatial and temporal changes in the insulin-like
9 growth factor (IGF) axis indicate autocrine/paracrine actions of IGF-I within wounds of
10 the rat brain. *Endocrinology.* 1997;138(7):3024-3034. doi:10.1210/endo.138.7.5284
- 11 61. Fletcher L, Isgor E, Sprague S, *et al.* Spatial distribution of insulin-like growth factor
12 binding protein-2 following hypoxic-ischemic injury. *BMC Neurosci.* 2013;14:158.
13 doi:10.1186/1471-2202-14-158
- 14 62. Breese CR, D'Costa A, Rollins YD, *et al.* Expression of insulin-like growth factor-1 (IGF-
15 1) and IGF-binding protein 2 (IGF-BP2) in the hippocampus following cytotoxic lesion of
16 the dentate gyrus. *J Comp Neurol.* 1996;369(3):388-404. doi:10.1002/(SICI)1096-
17 9861(19960603)369:3<388::AID-CNE5>3.0.CO;2-1
- 18 63. Kar S, Baccichet A, Quirion R, Poirier J. Entorhinal cortex lesion induces differential
19 responses in [125I]insulin-like growth factor I, [125I]insulin-like growth factor II and
20 [125I]insulin receptor binding sites in the rat hippocampal formation. *Neuroscience.*
21 1993;55(1):69-80. doi:10.1016/0306-4522(93)90455-o
- 22 64. Guan J, Williams CE, Skinner SJ, Mallard EC, Gluckman PD. The effects of insulin-like
23 growth factor (IGF)-1, IGF-2, and des-IGF-1 on neuronal loss after hypoxic-ischemic brain
24 injury in adult rats: evidence for a role for IGF binding proteins. *Endocrinology.*
25 1996;137(3):893-898. doi:10.1210/endo.137.3.8603600
- 26 65. Braak H, Braak E. Staging of Alzheimer's disease-related neurofibrillary changes.
27 *Neurobiol Aging.* 1995;16(3):271-284. doi:10.1016/0197-4580(95)00021-6

- 1 66. Bolós M, Fernandez S, Torres-Aleman I. Oral administration of a GSK3 inhibitor increases
2 brain insulin-like growth factor I levels. *J Biol Chem.* 2010;285(23):17693-17700.
3 doi:10.1074/jbc.M109.096594
- 4 67. Lucas JJ, Hernández F, Gómez-Ramos P, Morán MA, Hen R, Avila J. Decreased nuclear
5 beta-catenin, tau hyperphosphorylation and neurodegeneration in GSK-3beta conditional
6 transgenic mice. *EMBO J.* 2001;20(1-2):27-39. doi:10.1093/emboj/20.1.27
- 7 68. Engel T, Lucas JJ, Gómez-Ramos P, Moran MA, Avila J, Hernández F. Coexpression of
8 FTDP-17 tau and GSK-3beta in transgenic mice induce tau polymerization and
9 neurodegeneration. *Neurobiol Aging.* 2006;27(9):1258-1268.
10 doi:10.1016/j.neurobiolaging.2005.06.010
- 11 69. Hernández F, Borrell J, Guaza C, Avila J, Lucas JJ. Spatial learning deficit in transgenic
12 mice that conditionally over-express GSK-3beta in the brain but do not form tau filaments.
13 *J Neurochem.* 2002;83(6):1529-1533. doi:10.1046/j.1471-4159.2002.01269.x
- 14 70. Hoeflich A, Reyer A, Ohde D, *et al.* Dissociation of somatic growth, time of sexual
15 maturity, and life expectancy by overexpression of an RGD-deficient IGFBP-2 variant in
16 female transgenic mice. *Aging Cell.* 2016;15(1):111-117. doi:10.1111/acel.12413
- 17 71. Kanje M, Skottner A, Sjöberg J, Lundborg G. Insulin-like growth factor I (IGF-I)
18 stimulates regeneration of the rat sciatic nerve. *Brain Res.* 1989;486(2):396-398.
19 doi:10.1016/0006-8993(89)90531-3
- 20 72. Near SL, Whalen LR, Miller JA, Ishii DN. Insulin-like growth factor II stimulates motor
21 nerve regeneration. *Proc Natl Acad Sci U S A.* 1992;89(24):11716-11720.
22 doi:10.1073/pnas.89.24.11716
- 23 73. Guthrie KM, Nguyen T, Gall CM. Insulin-like growth factor-1 mRNA is increased in
24 deafferented hippocampus: spatiotemporal correspondence of a trophic event with axon
25 sprouting. *J Comp Neurol.* 1995;352(1):147-160. doi:10.1002/cne.903520111
- 26 74. Duan X, Qiao M, Bei F, Kim IJ, He Z, Sanes JR. Subtype-specific regeneration of retinal
27 ganglion cells following axotomy: effects of osteopontin and mTOR signaling. *Neuron.*
28 2015;85(6):1244-1256. doi:10.1016/j.neuron.2015.02.017

- 1 75. Anderson MA, O'Shea TM, Burda JE, *et al.* Required growth facilitators propel axon
2 regeneration across complete spinal cord injury. *Nature*. 2018;561(7723):396-400.
3 doi:10.1038/s41586-018-0467-6
- 4 76. Liu Y, Wang X, Li W, *et al.* A Sensitized IGF1 Treatment Restores Corticospinal Axon-
5 Dependent Functions. *Neuron*. 2017;95(4):817-833.e4. doi:10.1016/j.neuron.2017.07.037
- 6 77. Lane EM, Hohman TJ, Jefferson AL, Alzheimer's Disease Neuroimaging Initiative.
7 Insulin-like growth factor binding protein-2 interactions with Alzheimer's disease
8 biomarkers. *Brain Imaging Behav*. 2017;11(6):1779-1786. doi:10.1007/s11682-016-9636-
9 0
- 10 78. Lee WH, Michels KM, Bondy CA. Localization of insulin-like growth factor binding
11 protein-2 messenger RNA during postnatal brain development: correlation with insulin-
12 like growth factors I and II. *Neuroscience*. 1993;53(1):251-265. doi:10.1016/0306-
13 4522(93)90303-w
- 14 79. Khan S, Lu X, Huang Q, *et al.* IGFBP2 plays an essential role in cognitive development
15 during early life. *Adv Sci (Weinh)*. 2019;6(23):1901152. doi:10.1002/advs.201901152
- 16 80. Burgdorf J, Colechio EM, Ghoreishi-Haack N, *et al.* IGFBP2 produces rapid-acting and
17 long-lasting effects in rat models of posttraumatic stress disorder via a novel mechanism
18 associated with structural plasticity. *Int J Neuropsychopharmacol*. 2017;20(6):476-484.
19 doi:10.1093/ijnp/pyx007
- 20 81. Brooker GJ, Kalloniatis M, Russo VC, Murphy M, Werther GA, Bartlett PF. Endogenous
21 IGF-1 regulates the neuronal differentiation of adult stem cells. *J Neurosci Res*.
22 2000;59(3):332-341. doi:10.1002/(sici)1097-4547(20000201)59:3<332::aid-
23 jnr6>3.0.co;2-2
- 24 82. Burgdorf JS, Yoon S, Dos Santos M, Lammert CR, Moskal JR, Penzes P. An IGFBP2-
25 derived peptide promotes neuroplasticity and rescues deficits in a mouse model of Phelan-
26 McDermid syndrome. *Mol Psychiatry*. 2023;28(3):1101-1111. doi:10.1038/s41380-022-
27 01904-0
- 28 83. Stein TD, Anders NJ, DeCarli C, Chan SL, Mattson MP, Johnson JA. Neutralization of
29 transthyretin reverses the neuroprotective effects of secreted amyloid precursor protein

- 1 (APP) in APPSW mice resulting in tau phosphorylation and loss of hippocampal neurons:
2 support for the amyloid hypothesis. *J Neurosci.* 2004;24(35):7707-7717.
3 doi:10.1523/JNEUROSCI.2211-04.2004
- 4 84. Schindler N, Mayer J, Saenger S, *et al.* Phenotype analysis of male transgenic mice
5 overexpressing mutant IGFBP-2 lacking the Cardin-Weintraub sequence motif: Reduced
6 expression of synaptic markers and myelin basic protein in the brain and a lower degree of
7 anxiety-like behaviour. *Growth Horm IGF Res.* 2017;33:1-8.
8 doi:10.1016/j.ghir.2016.11.003
- 9 85. Gómez-Isla T, Price JL, McKeel DW Jr, Morris JC, Growdon JH, Hyman BT. Profound
10 loss of layer II entorhinal cortex neurons occurs in very mild Alzheimer's disease. *J*
11 *Neurosci.* 1996;16(14):4491-4500. doi:10.1523/JNEUROSCI.16-14-04491.1996
- 12 86. Dickerson BC, Goncharova I, Sullivan MP, *et al.* MRI-derived entorhinal and hippocampal
13 atrophy in incipient and very mild Alzheimer's disease. *Neurobiol Aging.* 2001;22(5):747-
14 754. doi:10.1016/s0197-4580(01)00271-8
- 15 87. Dickerson BC, Bakkour A, Salat DH, *et al.* The cortical signature of Alzheimer's disease:
16 regionally specific cortical thinning relates to symptom severity in very mild to mild AD
17 dementia and is detectable in asymptomatic amyloid-positive individuals. *Cereb Cortex.*
18 2009;19(3):497-510. doi:10.1093/cercor/bhn113
- 19 88. Schwarz CG, Gunter JL, Wiste HJ, *et al.* A large-scale comparison of cortical thickness
20 and volume methods for measuring Alzheimer's disease severity. *Neuroimage Clin.*
21 2016;11:802-812. doi:10.1016/j.nicl.2016.05.017
- 22 89. Toledo JB, Da X, Bhatt P, *et al.* Relationship between plasma analytes and SPARE-AD
23 defined brain atrophy patterns in ADNI. *PLoS One.* 2013;8(2):e55531.
24 doi:10.1371/journal.pone.0055531
- 25 90. McLimans KE, Webb JL, Anantharam V, Kanthasamy A, Willette AA, Alzheimer's
26 Disease Neuroimaging Initiative. Peripheral versus Central Index of Metabolic
27 Dysfunction and Associations with Clinical and Pathological Outcomes in Alzheimer's
28 Disease. *J Alzheimers Dis.* 2017;60(4):1313-1324. doi:10.3233/JAD-170263

- 1 91. Meyer PF, Savard M, Poirier J, *et al.* Bi-directional Association of Cerebrospinal Fluid
2 Immune Markers with Stage of Alzheimer's Disease Pathogenesis. *J Alzheimers Dis.*
3 2018;63(2):577-590. doi:10.3233/JAD-170887
- 4 92. Azar WJ, Zivkovic S, Werther GA, Russo VC. IGFBP-2 nuclear translocation is mediated
5 by a functional NLS sequence and is essential for its pro-tumorigenic actions in cancer
6 cells. *Oncogene.* 2014;33(5):578-588. doi:10.1038/onc.2012.630
- 7 93. Mahapatra MK, Karuppasamy M, Sahoo BM. Therapeutic Potential of Semaglutide, a
8 Newer GLP-1 Receptor Agonist, in Abating Obesity, Non-Alcoholic Steatohepatitis and
9 Neurodegenerative diseases: A Narrative Review. *Pharm Res.* 2022;39(6):1233-1248.
10 doi:10.1007/s11095-022-03302-1
- 11 94. Doré S, Kar S, Quirion R. Rediscovering an old friend, IGF-I: potential use in the treatment
12 of neurodegenerative diseases. *Trends Neurosci.* 1997;20(8):326-331. doi:10.1016/s0166-
13 2236(96)01036-3

16 Figure legends

17

18 **Figure 1 CSF IGFBP2 levels progressively increase over 5 years in asymptomatic**
19 **PREVENT-AD participants.** IGFBP2 was measured in the CSF of a subset of PREVENT-AD
20 participants ($n = 27$) at baseline and at follow-up visits, using the Olink Proximity Extension
21 Assay. Linear mixed models accounting for participant-specific trajectories demonstrate CSF
22 IGFBP2 levels increase in a subset of at-risk individuals that have been followed for 5 years. β and
23 P values are located in the top left corner. PREVENT-AD, Pre-symptomatic EVAluation of
24 Experimental or Novel Treatments for Alzheimer's Disease; CSF, cerebrospinal fluid; IGFBP2,
25 insulin-like growth factor binding protein-2; NPX, Normalized Protein eXpression.

26

1 **Figure 2 CSF IGFBP2 is associated with CSF and PET AD biomarkers in the asymptomatic**
 2 **PREVENT-AD cohort.** CSF IGFBP2 levels were measured using the Olink Proximity Extension
 3 Assay ($n = 109$). (A) CSF AD biomarkers $A\beta_{42}$, (B) p_{181} -tau and (C) t-tau were measured using
 4 validated Innotech ELISA kits, following the standardized protocols established by the
 5 BIOMARKAPD consortium ($n = 101$). (D) Global cortical amyloid SUVR was measured using
 6 ^{18}F -NAV4694 ($n = 46$). (E) Tau deposition in the entorhinal cortex was measured with flortaucipir
 7 ($n = 49$). The synaptic markers (F) SNAP-25 ($n = 106$), (G) SYT-1 ($n = 106$), (H) GAP43 ($n =$
 8 46) and (I) NRGN ($n = 46$) were quantified using immunoprecipitation followed by mass
 9 spectrometry. Significant or trend-level linear regressions are represented with a blue confidence
 10 region of the fitted line. R^2 and P values are located in the top left corners of each panel. Analyses
 11 were adjusted for age, sex and *APOE* $\epsilon 4$ carrier status. PREVENT-AD, PRe-symptomatic
 12 EValuation of Experimental or Novel Treatments for Alzheimer's Disease; CSF, cerebrospinal
 13 fluid; IGFBP2, insulin-like growth factor binding protein-2; NPX, Normalized Protein eXpression;
 14 AD, Alzheimer's disease; $A\beta_{42}$, amyloid-beta 42; p_{181} -tau, phosphorylated tau at residue 181; t-
 15 tau, total tau; ELISA, enzyme-linked immunosorbent assay; BIOMARKAPD, Biomarker's for
 16 Alzheimer's and Parkinson's disease; PET, positron emission tomography; SUVR, standardized
 17 uptake value ratio; SNAP-25, synaptosomal-associated protein 23kDa; a.u., arbitrary units; SYT1,
 18 synaptotagmin-1; GAP43, growth-associated protein 43; pg/mL, picogram per millilitre NRGN,
 19 neurogranin; APOE, apolipoprotein E.

20
 21 **Figure 3 CSF IGFBP2 is associated with longitudinal changes in delayed memory and**
 22 **visuospatial abilities over 5-8 years in PREVENT-AD.** CSF IGFBP2 levels were measured
 23 using the Olink Proximity Extension Assay ($n = 109$). Cognitive performance trajectory slopes
 24 were computed for each cognitive domain of the RBANS (delayed memory, visuospatial abilities,
 25 language, immediate memory and attention) in a subset of PREVENT-AD participants that were
 26 followed for 5-8 years ($n = 89$). Significant linear regressions are represented with a confidence
 27 region of the fitted line (blue for delayed memory and red for visuospatial abilities). R^2 , β and P
 28 values are located in the top right corner. Analyses were adjusted for age, sex, *APOE* $\epsilon 4$ carrier
 29 status and years of education. PREVENT-AD, PRe-symptomatic EValuation of Experimental or
 30 Novel Treatments for Alzheimer's Disease; CSF, cerebrospinal fluid; IGFBP2, insulin-like growth

1 factor binding protein-2; NPX, Normalized Protein eXpression; RBANS, Repeatable Battery for
2 the Assessment of Neuropsychological Status; APOE, apolipoprotein E.

3

4 **Figure 4 CSF IGFBP2 is associated with atrophy in AD-related brain regions in PREVENT-**
5 **AD.** CSF IGFBP2 levels were measured using the Olink Proximity Extension Assay ($n = 109$).
6 T1-weighted structural MRI scans were performed on a subset of PREVENT-AD participants (n
7 = 104). The imaging processing pipeline CIVET 1.1.12 was used to analyze neuroimaging data.
8 (A) Entorhinal cortex volumes were normalized by total intracranial volumes. Cortical thickness
9 measurements were acquired from AD-related brain regions, such as the (B) piriform cortex, (C)
10 inferior temporal gyrus and (D) middle temporal gyrus. Significant or trend-level linear regressions
11 are represented with a confidence region of the fitted line (blue for left hemisphere and red for
12 right hemisphere). R^2 and P values are located in the top right corner of each panel. Analyses were
13 adjusted for age, sex and *APOE* $\epsilon 4$ carrier status. PREVENT-AD, PRE-symptomatic EVAluation
14 of Experimental or Novel Treatments for Alzheimer's Disease; AD, Alzheimer's disease; CSF,
15 cerebrospinal fluid; IGFBP2, insulin-like growth factor binding protein-2; NPX, Normalized
16 Protein eXpression; ICV, intracranial volume; mm, millimetres; MRI, magnetic resonance
17 imaging; APOE, apolipoprotein E.

18

19 **Figure 5 CSF and plasma IGFBP2 is elevated in CSF A β (+)/t-tau(+) individuals from the**
20 **ADNI-1 cohort.** (A) Cognitively unaffected participants (red, $n = 92$) and participants with MCI
21 (blue, $n = 149$) from the ADNI-1 cohort were staged as CSF amyloid and/or CSF total tau positive
22 according to the recommended thresholds of 192 pg/mL and 93 pg/mL. Linear models, adjusted
23 for age, sex and *APOE* $\epsilon 4$ carrier status were used to examine mean differences in IGFBP2 protein
24 levels across stages. (B) CSF IGFBP2 was elevated at Stage 2 ($n = 77$) relative to Stage 0 ($n = 80$)
25 and Stage 1 ($n = 68$). Furthermore, CSF IGFBP2 was elevated in SNAP ($n = 8$) compared to Stage
26 0. (C) Plasma IGFBP2 was elevated at Stage 2 ($n = 84$) relative to Stage 0 ($n = 98$) and Stage 1 (n
27 = 60). However, plasma IGFBP2 did not significantly differ between SNAP ($n = 12$) and Stage 0.
28 The data are represented as mean \pm SEM. * $P < 0.05$; ** $P < 0.01$. ADNI, Alzheimer's disease
29 neuroimaging initiative; CSF, cerebrospinal fluid; IGFBP2, insulin-like growth factor binding
30 protein-2; NPX, Normalized Protein eXpression; A β , amyloid-beta; A β_{42} , amyloid-beta 42; t-tau,

1 total tau; pg/mL, picogram per millilitre; SNAP, suspected non-Alzheimer pathology; MCI, mild
2 cognitive impairment; ng/mL, nanogram per millilitre; log, logarithm base 10; APOE,
3 apolipoprotein E; SEM, standard error of the mean.

4

5 **Figure 6 Elevated plasma IGFBP2 is associated with a greater rate of conversion to AD in**
6 **individuals from the ADNI-1 cohort.** Cox proportional hazards models examined the association
7 between baseline plasma IGFBP2 levels and rate of conversion to AD. The first quartile (blue) and
8 fourth (red) values of plasma IGFBP2 were contrasted. Participants were followed from the
9 baseline visit to the time of diagnosis (of AD), or to the time the participant was last confirmed to
10 be free of AD (mean follow-up, 3.8 years; range, 0.5-16.5 years). Of the 226 individuals that were
11 followed longitudinally, 107 individuals progressed to AD. Individuals with plasma IGFBP2
12 values in the fourth quartile exhibited a greater rate of conversion to AD, compared to the first
13 quartile. HR and *P* values are located in the top right corner. Cox models were adjusted for age,
14 gender and *APOE* ϵ 4 carrier status. ADNI, Alzheimer's disease neuroimaging initiative; IGFBP2,
15 insulin-like growth factor binding protein-2; NPX, Normalized Protein eXpression; AD,
16 Alzheimer's disease; HR, hazard ratio; APOE, apolipoprotein E.

17

18 **Figure 7 Frontal cortex IGFBP2 gene expression is reduced in autopsy-confirmed AD brains,**
19 **however protein levels do not differ from elderly controls.** (A) Microarray technology was used
20 to measure IGFBP2 mRNA levels in the frontal cortex of autopsy-confirmed AD brains ($n = 55$)
21 and elderly controls ($n = 31$) from the QFP cohort. (B) IGFBP2 protein levels in the frontal cortex
22 were measured in AD brains ($n = 53$) and control brains ($n = 25$) using a commercially available
23 ELISA kit. Analyses were adjusted for age, sex, *APOE* ϵ 4 carrier status and post-mortem interval.
24 The data are represented as mean \pm SEM. * $P < 0.05$. QFP, Quebec Founder Population; IGFBP2,
25 insulin-like growth factor binding protein-2; mRNA, messenger RNA; log₂, logarithm base 2;
26 CTL, control; AD, Alzheimer's disease; ELISA, enzyme-linked immunosorbent assay; SEM,
27 standard error of the mean; APOE, apolipoprotein E, SEM, standard error of the mean.

28

29




Figure 1
159x100 mm (x DPI)




Figure 2
150x198 mm (x DPI)

1
2
3
4




Figure 3
159x124 mm (x DPI)

1
2
3
4




Figure 4
159x136 mm (x DPI)

1
2
3
4




Figure 5
159x180 mm (x DPI)

1
2
3
4




Figure 6
159x142 mm (x DPI)

1
2
3
4




Figure 7
159x71 mm (x DPI)

1
2
3
4

ACCEPTED MANUSCRIPT

1 **Table 1 Baseline Participant Demographics**

	PREVENT-AD	ADNI-I			QFP	
	CU	CU	MCI	AD	CU	AD
N Sample	109	58	395	111	31	55
Mean Age, Years (SD)	62.60 (5.43)	75.11 (5.77)	74.73 (7.40)	74.73 (8.08)	77.39 (11.37)	80.71 (6.39)
N Female (%)	76 (69.72)	28 (48.28)	140 (35.44)	47 (42.34)	11 (35.48)	23 (41.81)
N APOE ε4+ (%)	43 (39.44)	5 (8.62)	210 (53.16)	75 (67.57)	9 (29.03)	32 (58.18)
Mean BMI, kg/m ² (SD)	27.11 (4.47)	27.02 (4.12)	26.09 (3.97)	25.59 (3.82)	-	-
Mean HbA1c, % (SD)	5.40 (0.40) ^a	-	-	-	-	-
Mean Systolic BP, mmHg (SD)	120.20 (13.85)	131.41 (17.65)	132.79 (18.14)	135.05 (17.11)	-	-
Mean Education, Years (SD)	14.88 (2.94) ^b	15.67 (2.78)	15.64 (3.04)	15.09 (3.21)	-	-
N Amyloid Positive (%)	37 (33.94)	21 (36.21)	205 (51.90)	91 (81.98)	0(0)	55 (100)
Mean CSF Aβ ₄₂ , pg/mL (SD)	1145.73 (277.62) ^{c,d}	250.85 (21.08)	163.48 (52.90)	142.56 (39.32)	-	-
Mean CSF p ₁₈₁ -tau, pg/mL (SD)	46.83 (18.00) ^{c,d}	21.07 (8.43)	36.15 (19.32)	42.05 (19.96)	-	-
Mean CSF t-tau, pg/mL (SD)	273.09 (129.97) ^{c,d}	63.62 (21.76)	102.33 (59.78)	120.47 (56.58)	-	-
Mean CSF IGFBP2, NPX (SD)	6.89 (0.67)	-	-	-	-	-
Mean CSF IGFBP2, ng/mL (SD)	-	100.85 (15.85)	104.93 (18.87)	103.02 (18.76)	-	-
Mean Plasma IGFBP2, log ₁₀ (SD)	-	1.88 (0.20)	1.99 (0.23)	1.91 (0.12)	-	-
Mean Global Aβ, SUVR (SD)	1.30 (0.27) ^e	-	-	-	-	-
Mean Tau metaROI, SUVR (SD)	1.17 (0.07) ^f	-	-	-	-	-
Mean Cortical IGFBP2, log ₂ (SD)	-	-	-	-	2.43 (0.56) ^g	2.50 (0.83) ^g
Mean Postmortem interval, h (SD)	-	-	-	-	30.03 (19.85)	21.07 (10.36)

2 PREVENT-AD, Presymptomatic Evaluation of Experiment or Novel Treatments for Alzheimer's disease; ADNI-I, Alzheimer's Disease
3 Neuroimaging Initiative; QFP, Quebec Founder Population (autopsy confirmed cases only, amyloid positivity dependent on plaque density); CU,
4 cognitively unaffected; MCI, mild cognitive impairment; AD, Alzheimer's disease; APOE ε4+, Apolipoprotein ε4 carriers; BMI, body mass index;
5 kg/m², kilograms per square meter; HbA1c, hemoglobin A1c; BP, blood pressure; mmHg, millimetres of mercury; CSF, cerebrospinal fluid; Aβ₄₂,
6 amyloid beta 42; p₁₈₁-tau, phosphorylated tau 181; t-tau, total tau; IGFBP2, insulin-like growth factor binding protein-2; NPX, Normalized Protein
7 eXpression; SUVR, standardized uptake value ratio; ROI, region of interest; h, hours; SD, standard deviation; N, sample size; pg/mL, picograms
8 per millilitre; ng/mL, nanograms per millilitre.

9 ^a107 participants had HbA1c values available.

10 ^b106 participants had RBANS (Total Score) values available.

11 ^c101 PREVENT-AD participants had CSF Aβ₄₂, p₁₈₁-tau and t-tau (pg/mL) values available.

12 ^dPREVENT-AD (Fujirebio Innotest ELISA) and ADNI (INNO-BIA AlzBio3 Immunoassay) used different assays to measure the core CSF AD
13 biomarkers, which explains the differences.

14 ^e46 PREVENT-AD participants had Global Aβ SUVR values available.

15 ^f49 PREVENT-AD participants had Tau metaROI SUVR values available.

16 ^g78 QFP participants had cortical IGFBP2 protein values available.

17