



# User feedback on the NHS test & Trace Service during COVID-19: The use of machine learning to analyse free-text data from 37,914 England adults

P. Bondaronek<sup>a,b,\*</sup>, T. Papakonstantinou<sup>a</sup>, C. Stefanidou<sup>a</sup>, T. Chadborn<sup>a</sup>

<sup>a</sup> Office for Health Improvement & Disparities, Department of Health and Social Care, London, SW1H 0EU, United Kingdom

<sup>b</sup> Institute of Health Informatics, University College London, London, NW1 2DA, United Kingdom

## ARTICLE INFO

### Keywords:

Public health  
Machine learning  
Qualitative data  
Contact tracing  
COVID-19

## ABSTRACT

**Objectives:** The UK government's approach to the pandemic relies on a test, trace and isolate strategy, mainly implemented via the digital NHS Test & Trace Service. Feedback on user experience is central to the successful development of public-facing Services. As the situation dynamically changes and data accumulate, interpretation of feedback by humans becomes time-consuming and unreliable. The specific objectives were to 1) evaluate a human-in-the-loop machine learning technique based on structural topic modelling in terms of its Service ability in the analysis of vast volumes of free-text data, 2) generate actionable themes that can be used to increase user satisfaction of the Service.

**Methods:** We evaluated an unsupervised Topic Modelling approach, testing models with 5–40 topics and differing covariates. Two human coders conducted thematic analysis to interpret the topics. We identified a Structural Topic Model with 25 topics and metadata as covariates as the most appropriate for acquiring insights.

**Results:** Results from analysis of feedback by 37,914 users from May 2020 to March 2021 highlighted issues with the Service falling within three major themes: multiple contacts and incompatible contact method and incompatible contact method, confusion around isolation dates and tracing delays, complex and rigid system.

**Conclusions:** Structural Topic Modelling coupled with thematic analysis was found to be an effective technique to rapidly acquire user insights. Topic modelling can be a quick and cost-effective method to provide high quality, actionable insights from free-text feedback to optimize public health Services.

## 1. Introduction

NHS Test & Trace is a crucial component of the UK government's COVID-19 recovery strategy [1]. The Test & Trace Service traces the contacts of COVID-19-positive cases, and provides guidance on the next steps for both cases and contacts. The Service comprises of web-based, phone contacts, emails and text messaging. However, adherence to all parts of the process has been poor [2]. Improving how the NHS Test & Trace Service works for users is a necessary part of solving this problem.

Feedback on user experience is central to the successful development of public-facing Services, no less so when systems have been developed rapidly in a pandemic [3]. If individuals have a negative experience with the Service, they may not complete data input, be less likely to use it in future, and more likely to report negative experiences to others. In addition, not believing that contact tracing systems are effective predicts not engaging with them [4].

Users of the Test & Trace Service are asked to rate their experience

and leave feedback on how the system can be improved. A large volume of free-text data has been gathered with >1,800,000 words produced up to March 2021 just by those who said they were dissatisfied/very dissatisfied. Human coders have only been able to examine a portion of the data (0.04% of the data collected from May to October 2020).

Such data requires an automated approach that is enabled by Artificial Intelligence (AI) and Machine Learning (ML) approaches. Topic modelling is an unsupervised Natural Language Processing (NLP) approach that has gained popularity in the analysis of free text data. Topic models are generative models of word counts, that are able to automatically infer latent topics from text. They are particularly useful as tools to analyse large volumes of open-ended free-text responses in an exploratory bottom-up manner. Our approach is based on an application of the Structural Topic Model (STM) [5] in particular. The STM is a general framework for topic modelling that enables the researchers to include document-level metadata as covariates in a topic model and estimate their relationships to specific topics. We employed STM as the

\* Corresponding author. UCL Institute Of Health Informatics, 222 Euston Road, NW1 2DA, London, United Kingdom.

E-mail address: [p.bondaronek@ucl.ac.uk](mailto:p.bondaronek@ucl.ac.uk) (P. Bondaronek).

<https://doi.org/10.1016/j.puhip.2023.100401>

Received 1 October 2022; Received in revised form 19 June 2023; Accepted 19 June 2023

Available online 29 June 2023

2666-5352/Crown Copyright © 2023 Published by Elsevier Ltd on behalf of The Royal Society for Public Health. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

primary component in our hybrid approach, which combines ML with qualitative analysis. Our objective was to create and test a framework aimed at efficiently analysing large volumes of unstructured text data in a swift manner.

During the course of the pandemic topic modelling has been used to help provide insights into the public's behaviours and attitudes toward COVID-19 and the various measures put in place. Some examples of relevant applications are: understanding COVID-19 concerns and attitudes as expressed in social media and online forums [6–8]; categorising disinformation and mistrust [9]; surveillance of non-pharmacological interventions across the globe [10]; monitoring the content of news articles about COVID-19 [11]; and analysis of attitudes on mask-wearing [12]. In all instances, topic modelling has proved to be an excellent tool to acquire rapid, high-level insights on the various issues explored.

A study by Guetterman and colleagues [13] explored the use of NLP as a tool to augment and validate qualitative analysis. They found an NLP clustering technique based on the Wu-Palmer similarity measure [14] to be suitable both for providing a foundation for speeding up human qualitative analysis and for validating the human analysis. Similarly, another recent study [15] found topic modelling and graphic clustering of free-text data to be an effective assistive technique for human qualitative coding.

There is a need to develop a method to accelerate the understanding of the complexity of what influences human behaviours. Indeed, human behaviour is central to the transmission of SARS-Cov-2, and changing behaviour is crucial to preventing transmission. Automated approaches to data analysis of instructed data is needed to respond quickly to the changing situation. Moreover, other pandemic-related digital Services face similar problems of rapid rollout and need rapid analysis of user feedback.

## 2. Aims and objectives

The aim of this study was to evaluate the use of ML as a tool to help understand the issues with the NHS Test & Trace Service as expressed in the free-text user feedback responses. The specific objectives were to 1) provide proof of concept for the application of a qualitative analysis framework based on unsupervised machine learning, 2) create actionable themes that can be used to increase user satisfaction of the Test & Trace Service.

## 3. Methods

### 3.1. Data

This study was conducted as a service evaluation of the contact tracing service. The permission to publish the study was granted by the Publication Standards Board at the UK Health Security Agency.

User feedback has been collected since the start of the Service, on 28 May 2020. We are using data from that date until 7 March 2021. Because of time constraints and the pragmatic nature of this study as a methodological proof of concept, we did not include more recent data in the reported analysis. As part of the contact tracing process in the UK, data was gathered through the NHS Test & Trace Service. Upon completion of the process, whether via an online/in-app form or a phone call, users are prompted to rate their satisfaction with the Service received. The available options for rating range from "Very satisfied" to "Very dissatisfied." Users are finally asked "How could we improve this Service?", to which they can respond in free text. The responses to this question provide a rich dataset of recommendations that can be used to improve the Service to better cater to its users' needs and respond to a dynamic situation.

On 7 March 2021, a total of 153,128 responses were collected. In this study, we are only analysing the free-text responses from users who indicated they were "dissatisfied" or "very dissatisfied" with the Service

(64,492 responses, 42%). We made the decision to only analyse this subset of responses in order to more easily and quickly identify those aspects of the Service users were most often dissatisfied with and minimise the amount of noise in the dataset. This approach allows us to swiftly generate recommendations for enhancing the Service by identifying the barriers users encountered to engaging with it or adhering to guidance in general. By identifying these barriers, we can effectively pinpoint areas that require improvement and suggest actionable steps for enhancing user engagement and compliance.

In addition to the free-text data, more structured data were collected, such as whether the user is a case (person testing positive for SARS-CoV-2) or contact (person named by a case as potentially exposed), the region where they are based, demographic information and "tier". The tier variable refers to the user journey through the contact tracing system. Tier 1 is the first phase of the journey, where most cases are managed through the automated system, and complex cases are contacted via phone. If individuals fail to complete the online contact tracing process within 24 h, they are escalated to Tier 2 and contacted by call agents. Additionally, all identified contacts of cases are assigned to Tier 2 and contacted by call agents. Cases and contacts requiring self-isolation are escalated to Tier 3, where they receive follow-up calls from the Service.

### 3.2. Text analysis procedure

#### 3.2.1. Preprocessing

The original dataset includes data collected since the Service was first implemented in May 2020, up until March 2021, from users in the England. We preprocessed the data using R (version 3.5.2), and cleaned the free text responses using base R functions, the *quanteda* [16] and *stm* [5] packages. We deleted observations with missing values and duplicate data and prepared the covariates for analysis. After preprocessing, the remaining data available for analysis included 37,914 responses. These data were converted into token units using the *quanteda* package after punctuation and numbers were removed. Data preprocessing was completed by deleting stop words and stemming the tokens.

#### 3.2.2. Structural topic model

Prior to running the models, we produced some diagnostic plots to identify the optimal number of topics, according to both the relevant metrics and this analysis' aims, focusing on the trade-off between semantic coherence and exclusivity [17]. We evaluated an unsupervised Topic Modelling approach, testing models with 5–40 topics and differing covariates in terms of coherence, residuals and interpretability by human coders (see <https://osf.io/cvzd3/>). We identified a Structural Topic Model with 25 topics and metadata as covariates as the most appropriate for acquiring insights.

#### 3.2.3. Output

The output analysed consisted of lists of 8 stemmed keywords with different weightings which is what the model interprets as the 'topic' and 10 representative quotes for each topic. Different types of word weightings were generated with each topic where the following two types were analysed in subsequent qualitative analysis: 1) Highest Prob (words within each topic with the highest probability) and 2) FREX (words that are both frequent and exclusive, identifying words that distinguish topics). In addition to that, a list of the most representative quotes for a given topic was generated by the algorithm. See Fig. 1 for an example of the output.

### 3.3. Qualitative analysis process

The model's output was analysed systematically in two stages. In Stage 1, the descriptive titles for the topics were generated and agreed. In Stage 2, the six stages process of conducting a thematic analysis was adapted to guide to analyse the topics generated by the text analysis

## Topic 14

### Keywords

Highest Prob: call, time, happi, keep, said, unwell, feel  
 FREX: caller, amount, hes, call, satisfi, unwell, rung

### Quotes

Not dissatisfied with my call but frustrated by the number of calls. They are all feeling unwell and [...] has anxiety

They have been called multiple times, 25 times I think he said. They feel unwell and are very unhappy about how many times they have been called.

She's not happy with the amount of calls.

Not me personally it's just system as she's had three calls with all her children when it could have been done in one call.

Called lots of times - fed up - feel harassed

Too meny nenessery calls

She's received numerous calls so would be happier if it was one call per household

One call per household , very frustrated with the amount of calls he has received.

No happy that he's had to provide information for the household several times on several calls. So not very happy to hear the information over and over again and lots of times on calls.

Client said he had been called before twice and there was no record he had been called before which he was not satisfied about

Fig. 1. Example of the Structural Topic Model output.

[18]. Two researchers (PB, TP) conducted the initial coding of the topics generated through test analysis. The process is summarised below.

First, the researchers familiarised themselves with the data by repeatedly reading the themes with associated quotes whilst searching for patterns. For this analysis, the emphasis was on identifying the issues that users of the Test & Trace Service were unsatisfied with. Secondly, the initial descriptions of the topics were generated. Up until this stage, the two reviewers were blinded (worked independently on their codes). Next, the researchers identified the initial themes and discussed the similarities and discrepancies between them. We use the term “topics” to refer to our labels of the topics generated by the STM model output and the term “themes” to refer to human-generated themes created through analysing the topics and identifying common patterns. In the thematic analysis context, these terms would be analogous to the terms “codes” and “themes” respectively. Fourth, the researchers reviewed the themes whilst looking for any user feedback that might have been present in the topics identified but not represented in the major themes. This was a continuous process that was reviewed and re-defined (step five). Lastly, the report summarising the themes with illustrative quotes was generated (step six).

## 4. Results

### 4.1. Descriptive statistics

Descriptive statistics for respondents are displayed in Table 1. The sample of dissatisfied users is majority female, with 57% falling within the 30–59 age group, evenly split between contacts and cases. Finally, the majority completed the procedure entirely online.

Table 1  
 Sample description.

	Variable	n	%
	n	37914	100%
Age group	0–9	1096	3%
	10–19	2674	7%
	20–29	6960	18%
	30–39	7201	19%
	40–49	7203	19%
	50–59	7264	19%
	60–69	3565	9%
	70–79	1259	3%
	80–89	503	1%
	90+	189	0%
Sex	Female	20817	55%
	Male	16973	45%
	Not specified	124	0%
Region	East Midlands	3282	9%
	East of England	3690	10%
	London	5897	16%
	North East	2114	6%
	North West	6215	16%
	South East	5469	14%
	South West	3026	8%
	West Midlands	3705	10%
Category	Yorkshire and Humber	4516	12%
	Case	19299	51%
Tier	Contact	18615	49%
	Automated	25681	68%
	Tier 1	20	0%
	Tier 2	5399	14%
	Tier 3	6442	17%
	Other	372	1%

#### 4.2. Inclusion of the topics in the qualitative analysis

Of 25 topics analysed qualitatively, 19 topics were included in the analysis as they provided substantial insights on various issues as expressed by the users' feedback. The full output of the topic analysis is available in the online repository (<https://osf.io/cvzd3/>).

The rationale for the exclusion of 6 topics from the analysis is provided below:

- Topic 22 related to the feedback on the testing Service, specifically comments on inconclusive and inconsistent results, delay/not receiving tests, comments on perceived lack of the need to self-isolate due to negative results following positive ones. Although this topic was important, it related to a separate issue than the Test & Trace Service per se
- One topic (topic 1) was deemed as incoherent due to the text analysis based on the Service name which was used repeatedly in the free text, i.e., "track" and "trace" and was excluded from further analysis
- Four topics included mixed issues that were represented in other themes (topics 11,5,3,6).

These remaining 19 topics were grouped into three major themes representing the family of issues as expressed by the users of contact tracing (Fig. 2):

##### 4.2.1. Major theme 1: multiple contacts and incompatible contact method

The feedback expressed in this major theme related to the distress caused by receiving too many calls, especially in the context of experiencing Covid-19 symptoms. Users complained about multiple contacts per household reiterating the same process and expressed their views that the households should be 'linked' together. Users also perceived that too much information was collected, and there was some indication of data privacy concerns. In addition, participants expressed that multiple accounts were generated for one person, and hence they experienced multiple contact attempts through different channels, i.e., calls, emails and messages.

One topic related to the issue where the contact method did not match users' digital skills and preferences. For example, users who did not own a smartphone were unable to access the links in the text message; those who did not have internet access; or were unable to fill in the online form due to lower digital skills.




Fig. 2. Visualisation of the qualitative analysis of the 19 topics.

#### 4.2.2. Major theme 2: confusion around isolation dates and tracing

The feedback expressed in this major theme related to receiving numerous and often conflicting information about isolation dates from different sources of communication. In many cases, users were already contacted by the case and started their isolation period. They expressed their concern about their perceived delay of the contact attempts by the Service because they “could have been carrying on as normal & spreading the disease”.

#### 4.2.3. Major theme 3: complex and rigid operational system

The feedback expressed in this major theme related to the complexity of the online form, especially around the difficulties with setting appropriate passwords and issues with completing the form in the context of experiencing Covid-19 symptoms. The form was seen as repetitive and rigid, with an inability to go back and adjust responses. Some questions were seen as repetitive, irrelevant and phrased in a complex manner.

There were various issues with the logic flow of the form, and the inability to input information that was perceived as important. For example, the list of symptoms was perceived as not exhaustive, there were issues with entering postcodes, the ambiguity between home and current address, and error messages when inputting addresses outside the UK, as well the lack of representativeness of certain ethnic backgrounds. The users perceived the options for information input as limited and, at the same time, some input requests were seen as repetitive.

## 5. Discussion

### 5.1. A statement of overall findings

This study provides some important insights into the user experience of the Test & Trace Service as expressed through the free-text feedback survey based on over 37 thousand responses. The main issues related to multiple contact attempts, delays in contact tracing, and technical issues with the form, specifically the complexity and rigidity of the survey. In line with previous research, technical issues were found to be a major cause of distress for users of the Services, making their engagement with the Service challenging and, as a consequence, affecting their ability to adhere to guidance [19]. Users expressed that the contact method used was incompatible with their level of digital skills, which was a substantial issue. This finding underscores existing concerns regarding equity and accessibility of digital public-facing Services [20]. Another important finding showed that users reported feeling overwhelmed and harassed by the number of contact attempts from contact tracers. This is in line with previous findings by Wright and colleagues [21].

Overall, Structural Topic Modelling was found to be an effective technique to rapidly acquiring user insights. The 25 topics provided highly specific insights into issues that can be utilized towards improving the Test & Trace Service, with minimal human involvement and low maintenance requirements, making them ideally suited as tools for the evaluation of pandemic response Services.

This study provides a proof of concept of a method to analyse large quantities of free text data in Service improvement, combining an AI-enabled text analytics followed by a qualitative analysis conducted by a human. As such, this method has the potential to be to contribute to the improvement of the Test & Trace Service, and other digital systems for

public health safety or health improvement. This framework is represented in the diagram below (Fig. 3).

### 5.2. Strengths and limitations

First, the strength of this study relates to the use of the advancement in AI and applying the methods to analyse large volumes of unstructured data in a public health emergency. Second, the systematically developed method which includes text analytics and qualitative data analysis demonstrates a proof of concept and paves the way for further application of this method to various public health domains.

This study has several limitations. First, the Service included users in England and we only analysed those that indicated they were dissatisfied. Hence, the results might not be generalisable to other contexts. In addition, these results might not be generalisable to those that disengaged with or who completed the Service but did not provide any feedback. This study does not provide an in-depth analysis and an exhaustive issue expressed by the users. However, the results are congruent with the qualitative analysis of the interviews conducted with the cases and contacts [22]. Second, we have not assessed the differences between different periods throughout the functioning of the Service.

With regards to methodology, an issue with applying STM relates to the lack of control over the variables explored. Although this is one of the strengths of the method, it also limits the researchers' ability to make strong inferences about hypothesised results, making the method less robust to interpretation biases. The quality of the STM output was generally high, as the majority of the 25 topics generated exhibited coherence and allowed for easy interpretation, highlighting the reliability of the results. Only one topic out of the 25 was found to be incoherent, indicating a high level of overall coherence and meaningful thematic representation in the output. A limitation of interpreting this output relates to missing data. After removing responses with missing data, the sample of responses analysed was reduced. By removing these records, there is a potential loss of valuable information and insights that could have been derived from the excluded data points. This reduction in the dataset could affect the representativeness and generalisability of the findings, and could have introduced bias. Research has shown that excluding incomplete datapoints might often result in disproportionate bias toward marginalised groups [23].

### 5.3. Implications for research, practice, policy

#### 5.3.1. Research

This study has several implications for research and further development of the model for automatic and pragmatic free-text data analysis. Specifically:

1. Further research is needed to develop a more sophisticated method that can be applied to monitor the changes in the Service over time.
2. As individuals have different experiences of the Service highlighting social inequalities, findings need to be stratified by sociodemographics to investigate the effects of this Service (and other such Services) based on sociodemographics, especially to emphasize the underrepresented and marginalised communities.
3. Need to develop a scalable model and apply it to accelerate understanding of different behaviours beyond the pandemic and into the




Fig. 3. Diagram representing a framework for analysis of qualitative data for Service improvement.



post-pandemic recovery, namely the behaviours associated with poorer outcomes and higher mortality rates in those infected with SARS-Cov-2, such as unhealthy diet, physical inactivity, alcohol consumption and smoking.

4. The current model needs to be applied to different datasets in order to increase its generalisability and usability of the model for public health professionals.
5. The present model and analysis framework should be validated by comparing it to human qualitative analysis in order to identify discrepancies and assess its reliability compared to conventional qualitative methods. There is evidence that the quality of the analysis produced by this framework is equal to that of human-only qualitative analysis, with substantial time and resource savings [24].

### 5.3.2. Practice and policy

There is a need to systematically reflect and gather the learnings from the pandemic in order to develop a more efficient response in the future. Some of the issues with the Service related to usability may be addressed in a time-efficient manner by simple technological changes. Some issues might require a more complex approach, for example, beliefs about the inefficiency of the Service. It is important that these issues are recognised and that various interventions addressing different problems are developed in order to develop a more efficient method of contact tracing in the context of the pandemic. Future research that incorporates data from different phases of the pandemic and subsequent iterations of the Service would offer valuable insights. By examining the evolution of themes over time, researchers can determine if similar concerns persist or if new issues emerge as the Service matures and adapts. Additionally, by stratifying the analysis based on demographic variables, a more nuanced understanding of the experiences and needs of specific populations can be obtained. This approach would enable targeted support and interventions tailored to address the unique challenges faced by different demographic groups.

## 6. Conclusion

This study provides a proof of concept of a method to analyse large quantities of free text data in Service improvement. It includes an AI-enabled text analytics followed by a qualitative analysis conducted by a human. As such, this method has the potential to contribute to the improvement of the Test & Trace Service, and other digital systems for public health safety or health improvement. More specifically, this method is best suited for cases where qualitative analysis is either needed rapidly or applied to large volumes of narrow-in-scope, short free-text responses. Importantly, this method is not suitable in cases where a nuanced qualitative analysis is the goal but can be a tool to augment and speed up existing qualitative approaches.

### Ethics approval and consent to participate

This study was conducted as a service evaluation of the contact tracing service using secondary data. The permission to publish the study was granted by the Publication Standards Board at the UK Health Security Agency.

### Consent for publication

Not applicable; no individual person's data is included.

### Availability of data and materials

The dataset used in this research cannot be made available as it includes free-text data that potentially compromise individual privacy. There was no consent obtained for the data to be made publicly available. The code used to analyse the data has been made available in an online repository (<https://doi.org/10.17605/OSF.IO/CVZD3>).

## Declaration of competing interests

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: During the conduct of the study (data collection, analysis and interpretation), all authors were employed at Public Health England Behavioural Insights. From 1st October, PHE was dissolved, and all authors have been transferred to the newly created Office for Health Improvement and Disparities at the Department of Health and Social Care. Neither PHE nor DHSC had any influence over the project or dissemination of the findings.

## Funding

This work was supported by Public Health England, United Kingdom.

## Contributions

PB conceptualised the study, facilitated access to data and resources, identified and developed the methodology, conducted qualitative analysis and contributed to writing the original draft of the manuscript and editing later versions. TP developed the methodology, conducted quantitative and qualitative analysis of the data, contributed to writing the original draft of the manuscript and editing later versions. CS conceptualised the study, conducted preliminary data analysis and contributed to editing later versions of the manuscript. TC supervised all aspects of the study.

## Acknowledgements

We thank Charlotte Anderson and the team at Contact Tracing Cell: data and surveillance (Public Health England, now transitioned to UK Health Security Agency) for access to the data; we thank Julian Flowers (Public Health Data Science, Public Health England) for providing guidance on statistical methods; we thank Will Nicholson from Behavioural Science Team, Public Health England for helping us explore the various supervised methods for data analysis.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.puhip.2023.100401>.

## References

- [1] A. Briggs, D. Jenkins, C. Fraser, NHS Test and Trace: the Journey So Far, Health Foundation, 2020. <https://www.health.org.uk/publications/long-reads/nhs-test-and-trace-the-journey-so-far>. (Accessed 13 April 2022).
- [2] L.E. Smith, H.W.W. Potts, R. Amlôt, et al., Adherence to the test, trace, and isolate system in the UK: results from 37 nationally representative surveys, *BMJ* (2021) 372, <https://doi.org/10.1136/bmj.n608>.
- [3] H. Potts, F. Death, C. Garattini, et al., Rapid evaluation of digital health products during the COVID-19 pandemic. <https://www.gov.uk/guidance/rapid-evaluation-of-digital-health-products-during-the-covid-19-pandemic>, 2020. (Accessed 13 April 2022).
- [4] L.E. Smith, R. Amlôt, H. Lambert, et al., Factors associated with adherence to self-isolation and lockdown measures in the UK: a cross-sectional survey, *Publ. Health* 187 (2020) 41–52, <https://doi.org/10.1016/j.puhe.2020.07.024>.
- [5] M.E. Roberts, B.M. Stewart, D. Tingley, *Stm: an R package for structural topic models*, *J. Stat. Software* (2019) 91, <https://doi.org/10.18637/jss.v091.i02>.
- [6] W. Jo, J. Lee, J. Park, et al., Online information exchange and anxiety spread in the early stage of the novel coronavirus (COVID-19) outbreak in South Korea: structural topic model and network analysis, *J. Med. Internet Res.* 22 (6) (2020), <https://doi.org/10.2196/19455>.
- [7] H. Yin, X. Song, S. Yang, et al., Sentiment analysis and topic modeling for COVID-19 vaccine discussions, *World Wide Web* (2022) 1–17, <https://doi.org/10.1007/s11280-022-01029-y>.
- [8] M. Zamani, H.A. Schwartz, J. Eichstaedt, et al., Understanding weekly COVID-19 concerns through dynamic content-specific LDA topic modeling, 2020, *Proc Conf Empir Methods Nat Lang Process* (2020) 193–198, <https://doi.org/10.18653/v1/2020.nlpccs-1.21>.

- [9] X. Song, J. Petrak, Y. Jiang, et al., Classification aware neural topic model for COVID-19 disinformation categorisation, *PLoS One* (2 February) (2021) 16, <https://doi.org/10.1371/journal.pone.0247086>.
- [10] Y. Li, P. Nair, Z. Wen, et al., Global surveillance of COVID-19 by mining news media using a multi-source dynamic embedded topic model, in: *BCB '20: Proceedings of the 11th ACM International Conference on Bioinformatics, Computational Biology and Health Informatics, 2020*, <https://doi.org/10.1145/3388440.341241815>.
- [11] Y. Bai, S. Jia, L. Chen, Topic evolution analysis of COVID-19 news articles, *J. Phys. Conf. Ser.* 1601 (2020), 052009, <https://doi.org/10.1088/1742-6596/1601/5/052009>.
- [12] K.R. Lee, B. Kim, D. Nan, et al., Structural topic model analysis of mask-wearing issue using international news big data, *Int. J. Environ. Res. Publ. Health* 18 (12) (2021), <https://doi.org/10.3390/ijerph18126432>.
- [13] T.C. Guetterman, T. Chang, M. DeJonckheere, et al., Augmenting qualitative text analysis with natural language processing: methodological study, *J. Med. Internet Res.* 20 (6) (2018), <https://doi.org/10.2196/JMIR.9702>.
- [14] Z. Wu, M. Palmer, *Verb semantics and lexical selection, 32nd Annual Meeting of the Association for Computational Linguistics* (1994) 133–138.
- [15] R.P. Lennon, R. Fraleigh, L.J. van Scoy, et al., Developing and testing an automated qualitative assistant (AQUA) to support qualitative analysis, *Fam Med Community Health* (2021) 9, <https://doi.org/10.1136/fmch-2021-001287>.
- [16] K. Benoit, K. Watanabe, H. Wang, et al., quanteda: an R package for the quantitative analysis of textual data, *J. Open Source Softw.* 3 (30) (2018), <https://doi.org/10.21105/joss.00774>.
- [17] M.E. Roberts, B.M. Stewart, D. Tingley, et al., Structural topic models for open-ended survey responses, *Am. J. Polit. Sci.* 58 (4) (2014), <https://doi.org/10.1111/ajps.12103>.
- [18] V. Braun, V. Clarke, Using thematic analysis in psychology, *Qual. Res. Psychol.* 3 (2) (2006), <https://doi.org/10.1191/1478088706qp0630a>.
- [19] O. Megnin-Viggars, P. Carter, G.J. Melendez-Torres, et al., Facilitators and barriers to engagement with contact tracing during infectious disease outbreaks: a rapid review of the evidence, *PLoS One.* 15 (10) (2020), e024147.
- [20] A.M. Kaihlanen, L. Virtanen, U. Buchert, N. Safarov, P. Valkonen, L. Hietapakka, I. Hörhammer, S. Kujala, A. Kouvonon, T. Heponiemi, Towards digital health equity—a qualitative study of the challenges experienced by vulnerable groups in using digital health services in the COVID-19 era, *BMC Health Serv. Res.* 22 (1) (2022 Feb 12) 188, <https://doi.org/10.1186/s12913-022-07584-4>.
- [21] L. Wright, A. Burton, A. McKinlay, et al., Public opinion about the UK government during COVID-19 and implications for public health: a topic modelling analysis of open-ended survey response data, March, medRxiv 21254094 [Preprint] 24 (2021), <https://doi.org/10.1101/2021.03.24.21254094>.
- [22] T. Shelby, C. Arechiga, A.J. Gupta, R. Hennein, C. Schenck, B. Weeks, M. Bond, L. Niccolai, J.L. Davis, L.E. Grau, I can't do it": A qualitative study exploring case and contact experiences with COVID-19 contact tracing, *BMC Public Health* 22 (1) (2022) 1963.
- [23] Y. Zhang, Long Q assessing fairness in the presence of missing data, *Adv. Neural Inf. Process. Syst.* 34 (2021) 16007–16019.
- [24] L.B. Towler, P. Bondaronek, T. Papakonstantinou, R. Amlôt, T. Chadborn, B. Ainsworth, L. Yardley, Applying machine-learning to rapidly analyse large qualitative text datasets to inform the COVID-19 pandemic response: comparing human and machine-assisted topic analysis techniques, medRxiv (2022).