

Adjuvant chemotherapy for adenocarcinoma arising from intraductal papillary mucinous neoplasia: multicentre ADENO-IPMN study

James Lucocq¹ , Jake Hawkyard², Beate Haugk², Omar Mownah³, Krishna Menon³, Takaki Furukawa⁴, Yosuke Inoue⁴, Yuki Hirose⁴, Naoki Sasahira⁴, Michael Feretis⁵, Anita Balakrishnan⁵, Carlo Ceresa⁶, Brian Davidson⁶, Rupaly Pande⁷, Bobby Dasari⁷, Lulu Tanno⁸, Dimitrios Karavias⁸ , Jack Helliwell⁹, Alistair Young⁹, Quentin Nunes¹⁰ , Tomas Urbonas¹¹, Michael Silva¹¹, Alex Gordon-Weeks¹¹, Jenifer Barrie¹², Dhanny Gomez¹², Stijn Van Laarhoven¹³, Francis Robertson², Hossain Nawara¹³, Joseph Doyle¹⁴, Ricky Bhogal¹⁴, Ewen Harrison¹⁵, Marcus Roalso¹⁶, Debora Ciprani¹⁷, Somaiah Aroori¹⁷, Bathiya Ratnayake¹⁸ , Jonathan Koea¹⁸, Gabriele Capurso¹⁹, Ruben Bellotti²⁰, Stefan Stättner²⁰, Tareq Alsaoudi²¹, Neil Bhardwaj²¹, Srujan Rajesh²¹, Fraser Jeffery²², Saxon Connor²² , Andrew Cameron²³, Nigel Jamieson²³ , Amy Sheen²⁴, Anubhav Mittal²⁵, Jas Samra²⁵, Anthony Gill^{24,25,26}, Keith Roberts¹, Kjetil Søreide¹⁶ and Sanjay Pandanaboyana^{2,27,*} 

¹Department of General Surgery, NHS Lothian, Edinburgh, UK

²Hepatopancreatobiliary and Transplant Unit, Freeman Hospital, Newcastle upon Tyne, UK

³Department of Hepatobiliary and Pancreatic Surgery, King's College Hospital, London, UK

⁴Hepato-Biliary-Pancreatic Medicine Department, Cancer Institute Hospital of Japanese Foundation for Cancer Research, Tokyo, Japan

⁵Cambridge Hepatobiliary and Pancreatic Surgery Unit, Addenbrooke's Hospital, Cambridge, UK

⁶Hepatobiliary and Pancreatic Surgery Unit, Royal Free Hospital, London, UK

⁷Hepatobiliary and Pancreatic Surgery Unit, University Hospitals Birmingham NHS Foundation Trust, Queen Elizabeth Hospital, Birmingham, UK

⁸Hepatobiliary and Pancreatic Surgery Unit, University Hospital Southampton, Southampton, UK

⁹Hepatobiliary and Pancreatic Surgery Unit, Leeds Teaching Hospitals NHS Trust, Leeds, UK

¹⁰Department of Hepatopancreatobiliary Surgery, East Lancashire Teaching Hospitals NHS Trust, Blackburn, UK

¹¹Oxford Hepato-Pancreato-Biliary Surgical Unit, Oxford University Hospitals NHS Foundation Trust, Oxford, UK

¹²Nottingham Hepato-Pancreato-Biliary Service, Nottingham University Hospitals NHS Foundation Trust, Nottingham, UK

¹³Department of General Surgery, University Hospitals Bristol and Weston NHS Foundation Trust, Bristol, UK

¹⁴Gastrointestinal Unit, Royal Marsden NHS Foundation Trust, London, UK

¹⁵Department of Clinical Surgery, University of Edinburgh, Edinburgh, UK

¹⁶Department of Gastrointestinal Surgery, Stavanger University Hospital, Stavanger, Norway

¹⁷Hepatopancreatobiliary Unit, University Hospitals Plymouth NHS Trust, Plymouth, UK

¹⁸Hepato-pancreato-biliary/Upper Gastrointestinal Unit, North Shore Hospital, Auckland, New Zealand

¹⁹Pancreatico-Biliary Endoscopy and Endosonography Division, Pancreas Translational and Clinical Research Centre, San Raffaele Scientific Institute IRCCS, Vita-Salute San Raffaele University, Milan, Italy

²⁰Department of Visceral, Transplant and Thoracic Surgery, Innsbruck Medical University, Innsbruck, Austria

²¹Leicester Hepatopancreatobiliary Unit, University Hospitals of Leicester NHS Trust, Leicester, UK

²²Department of General and Vascular Surgery, Christchurch Hospital, Canterbury District Health Board, Christchurch, New Zealand

²³Wolfson Wohl Cancer Research Centre, Research Institute of Cancer Sciences, University of Glasgow, Glasgow, UK

²⁴New South Wales Health Pathology, Department of Anatomical Pathology, Royal North Shore Hospital, Sydney, New South Wales, Australia

²⁵Department of Hepatopancreatobiliary Surgery, Royal North Shore Hospital, Sydney, New South Wales, Australia

²⁶Sydney Medical School, University of Sydney, Sydney, New South Wales, Australia

²⁷Population Health Sciences Institute, Newcastle University, Newcastle upon Tyne, UK

*Correspondence to: Sanjay Pandanaboyana, Hepatopancreatobiliary and Transplant Unit, Freeman Hospital, Freeman Road, Newcastle upon Tyne, NE7 7DN, UK (e-mail: sanjay.pandanaboyana@ncl.ac.uk)

Abstract

Background: The clinical impact of adjuvant chemotherapy after resection for adenocarcinoma arising from intraductal papillary mucinous neoplasia is unclear. The aim of this study was to identify factors related to receipt of adjuvant chemotherapy and its impact on recurrence and survival.

Methods: This was a multicentre retrospective study of patients undergoing pancreatic resection for adenocarcinoma arising from intraductal papillary mucinous neoplasia between January 2010 and December 2020 at 18 centres. Recurrence and survival outcomes for patients who did and did not receive adjuvant chemotherapy were compared using propensity score matching.

Results: Of 459 patients who underwent pancreatic resection, 275 (59.9%) received adjuvant chemotherapy (gemcitabine 51.3%, gemcitabine–capecitabine 21.8%, FOLFIRINOX 8.0%, other 18.9%). Median follow-up was 78 months. The overall recurrence rate was 45.5% and the median time to recurrence was 33 months. In univariable analysis in the matched cohort, adjuvant chemotherapy was not associated with reduced overall ($P = 0.713$), locoregional ($P = 0.283$) or systemic ($P = 0.592$) recurrence, disease-free survival ($P = 0.284$) or overall survival ($P = 0.455$). Adjuvant chemotherapy was not associated with reduced site-specific recurrence. In multivariable analysis, there was no association between adjuvant chemotherapy and overall recurrence (HR 0.89, 95% c.i. 0.57 to

Received: August 23, 2023. Revised: March 23, 2024. Accepted: March 26, 2024

© The Author(s) 2024. Published by Oxford University Press on behalf of BJS Foundation Ltd.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<https://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

1.40), disease-free survival (HR 0.86, 0.59 to 1.30) or overall survival (HR 0.77, 0.50 to 1.20). Adjuvant chemotherapy was not associated with reduced recurrence in any high-risk subgroup (for example, lymph node-positive, higher AJCC stage, poor differentiation). No particular chemotherapy regimen resulted in superior outcomes.

Conclusion: Chemotherapy following resection of adenocarcinoma arising from intraductal papillary mucinous neoplasia does not appear to influence recurrence rates, recurrence patterns or survival.

Introduction

Intraductal papillary mucinous neoplasms (IPMNs) arise from mucin-producing cells in the main pancreatic duct and/or its branches. Adenocarcinoma may complicate IPMNs in 20% of cases, but make up approximately 5% of all pancreatic adenocarcinomas¹. Following resection of adenocarcinoma arising from IPMN, patients are at significant risk of recurrence, which is estimated to occur in 32–43% of patients. Hence, adjuvant chemotherapy has been proposed along the lines of multimodal treatment for pancreatic cancer^{2–6}.

In patients with pancreatic ductal adenocarcinoma (PDAC), not associated with an underlying IPMN, the standard of care is surgical resection and adjuvant chemotherapy. More recently, the PREOPANC-1⁷ and Prep-02JSAPO5⁸ trials have demonstrated the benefit of a neoadjuvant treatment strategy in PDAC. Trials that demonstrated the benefit of adjuvant therapy in pancreatic cancer included patients with primary PDAC without associated IPMN. As such, the benefit of adjuvant therapy is unclear in patients with adenocarcinoma arising from IPMN^{6,9–14}. In particular, the effect of adjuvant chemotherapy on recurrence, its impact on site-specific recurrence, and the outcomes of different adjuvant chemotherapy regimens are largely unknown^{3–6}.

The aim of this study was to investigate the impact of adjuvant chemotherapy on recurrence and survival, its effect on site-specific recurrence, and to compare different chemotherapy regimens in a large, international, multicentre cohort of patients undergoing pancreatic resection for adenocarcinoma arising from IPMN.

Methods

This was a retrospective multicentre study of consecutive patients undergoing pancreatic resection for adenocarcinoma arising from IPMN between January 2010 and December 2020 at 18 academic pancreatic cancer centres in Europe, Asia, Australia, and New Zealand. The methods used to gather data on these patients were published previously in a study investigating the impact of treatment of recurrence on survival¹⁵. The presence of adenocarcinoma arising from IPMN was identified retrospectively based on histopathological specimens. Patients with PDAC with concomitant IPMN elsewhere in the specimen were excluded. A primary investigator was appointed for each centre and was responsible for leading data collection. The Research Electronic Data Capture (REDCap) system was used to store anonymized information (anonymized at the source), which was then maintained by Newcastle Joint Research Office. The institutional review board of each participating institution approved the study before initiation. The study was conducted according to the Declaration of Helsinki. Informed patient consent was not required given the retrospective nature of the study. STROBE recommendations were followed¹⁶.

Data collection, outcomes and definitions

Adenocarcinoma arising from IPMN was defined as an invasive carcinoma derived from IPMN, in accordance with the WHO

classification¹⁷. The type of adenocarcinoma was classified as either tubular or colloid, and the invasive component of the IPMN tumours were staged according to the TNM Classification of Malignant Tumours, 8th edition, published by the AJCC. Determination of R status was based on a cut-off distance from the tumour to the resection margin of less than 1 mm¹⁸. Where the underlying precursor epithelial lesion or subtype was not documented in the original reporting, the pathology slides were reviewed again by a pathologist with expertise in pancreatic cancer.

Duct type was classified as main duct type, branch duct type, or mixed type, according to consensus guidelines¹⁹. The type of surgical procedure and decision to administer adjuvant chemotherapy were at the discretion of each participating institution based on the location, degree, and extent of the tumour. The type and number of cycles of adjuvant chemotherapy were recorded.

In patients with resectable pancreatic cancer, upfront resection was performed and adjuvant chemotherapy was considered¹⁹. Patients underwent either Whipple's resection, pylorus-preserving pancreatoduodenectomy, distal pancreatectomy with splenectomy, distal pancreatectomy without splenectomy, or total pancreatectomy. In patients with borderline-resectable tumours portal or superior mesenteric vein resection was undertaken if infiltration was suspected.

Recurrence was diagnosed radiologically or histologically when available. Given the lack of current guidelines, follow-up was at the discretion of each centre. Overall recurrence was categorized as locoregional or systemic, and recurrence sites were reported. The overall rate and median time of overall (OS) and disease-free (DFS) survival were determined for the entire cohort. DFS was defined as survival in the absence of recurrence.

Statistical analysis

Clinicopathological variables associated with recurrence were identified using Kaplan–Meier (KM) analysis and the log rank test. In all statistical tests, $P < 0.050$ was considered significant.

The type of adjuvant chemotherapy used was compared between the pre-2017 and 2017-onwards groups using Pearson's χ^2 test to determine whether there had been a change in practice since the ESPAC-4 trial¹⁴. The administration rate of adjuvant chemotherapy between centres was compared using Pearson's χ^2 test. Clinicopathological variables associated with adjuvant chemotherapy use were identified by univariable analysis. Variables demonstrating statistical significance with a cut-off of $P < 0.100$ were included in a multivariable logistic regression model, with adjuvant chemotherapy as the dependent variable.

To allow for immortal time bias relating to administration of adjuvant chemotherapy, landmark analysis was performed at 6 months, whereby all patients who died before this time point were excluded. Postlandmark analysis propensity score matching (PSM) was performed to determine the impact of adjuvant chemotherapy on outcome; treatment (adjuvant chemotherapy) and control (no adjuvant chemotherapy) groups were matched for clinicopathological variables to decrease the effects of confounding. Propensity scores were calculated using

Table 1 Clinicopathological features of the overall cohort and predictors of recurrence

	Recurrence (n = 209)	No recurrence (n = 250)	P*
Age (years), median (range)	70 (27–87)	70 (36–92)	0.653†
Sex			0.533
Male	110 (52.6)	140 (56.0)	
Female	99 (47.4)	110 (44.0)	
Charlson Co-morbidity Index score, median (range)	4 (0–9)	4 (0–9)	0.453†
Borderline resectable	15 (3.3)	6 (2.4)	0.651
Operation			0.190
Whipple's	78 (19.1)	94 (37.6)	
PPPD	45 (21.5)	45 (18.0)	
DPS	38 (9.3)	34 (13.6)	
DPNS	12 (2.9)	18 (7.2)	
TP	36 (8.8)	59 (23.6)	
Multivisceral resection	38 (9.3)	31 (12.4)	0.001
Duct location			0.006
Main duct	114 (27.9)	152 (60.8)	
Side	33 (8.1)	30 (12.0)	
Mixed	43 (10.5)	56 (22.4)	
Tumour location			0.390
Head	138 (33.7)	154 (61.6)	
Body	21 (5.1)	31 (12.4)	
Tail	32 (7.8)	37 (14.8)	
Tumour size (mm), median (i.q.r.)	28 (20–35)	20 (10–30)	< 0.001†
Cyst size (mm), median (i.q.r.)	30 (24–45)	33 (20–47.5)	0.817†
Precursor epithelium			0.256
Gastric	42 (10.3)	44 (17.6)	
Intestinal	35 (8.6)	62 (24.8)	
Pancreatobiliary	70 (17.1)	75 (30.0)	
IOPN	9 (4.3)	11 (4.2)	
Differentiation grade			< 0.001
Well	24 (11.5)	58 (23.2)	
Moderately	102 (48.8)	129 (51.6)	
Poor	76 (36.4)	42 (16.8)	
Invasive component			< 0.001
Ductal	170 (81.3)	173 (69.2)	
Colloid	19 (9.1)	50 (20.0)	
Lymphovascular invasion	129 (61.7)	107 (42.8)	< 0.001
Perineural invasion	143 (68.4)	113 (45.2)	< 0.001
R1 resection	102 (48.8)	75 (30.0)	< 0.001
Adjuvant chemotherapy	143 (68.4)	132 (52.8)	< 0.001
AJCC stage			< 0.001
Ia	16 (7.7)	75 (30.0)	
Ib	13 (6.2)	39 (15.6)	
IIa	29 (13.9)	40 (16.0)	
IIb	125 (59.8)	75 (30.0)	
III	26 (12.4)	21 (8.4)	

Values are n (%) unless otherwise indicated. PPPD, pylorus-preserving pancreaticoduodenectomy; DPS, distal pancreatectomy with splenectomy; DPNS, distal pancreatectomy without splenectomy; TP, total pancreatectomy; IOPN, intraductal oncocytic papillary neoplasm. *log rank test. †Continuous data split by median and recurrence rates compared using log rank test.

probit regression and patients in the treatment group were matched 1 : 1 using nearest-neighbour matching with patients in the control group. Groups were matched for age, sex, Charlson Co-morbidity Index score (split by median), differentiation, perineural invasion, lymphovascular invasion, AJCC stage, and R status. The caliper was used to impose a maximum distance in propensity scores between possible matches and was set to 0.05. Variable balancing was assessed using a standardized mean difference (SMD; less than 0.1 considered balanced) and the distance variance ratio (value close to 1.0 considered balanced). Matching without replacement was used to optimise precision, taken that matching was satisfactory. Jitter and Love plots were used to show the distribution of propensity scores and SMD respectively of unmatched and matched cohorts. KM curves were plotted for the treatment and control groups for each outcome (recurrence and OS) and the log rank test was used to compare groups. In the matched cohort, Cox proportional hazards models were developed to identify the effect of adjuvant chemotherapy on

outcome. PSM was performed using the MatchIt package and graphs were displayed using the Cobalt package in R Studio 2022.02.1[®]. The PSM analysis was repeated in patients who received more contemporary chemotherapy regimens, specifically gemcitabine–capecitabine (GEM-CAP) or folinic acid, fluorouracil, irinotecan and oxaliplatin (FOLFIRINOX). The overall cohort was investigated to determine whether any particular high-risk subgroup (for example, poor differentiation, higher AJCC stage) would benefit from the administration of adjuvant chemotherapy. Treatment and control groups were compared using KM analysis and the log rank test.

Results

Study population

Of 459 patients who underwent pancreatic resection 275 (59.9%) received adjuvant chemotherapy. Chemotherapy regimens included gemcitabine (141, 51.3%), GEM-CAP (60, 21.8%), FOLFIRINOX (22, 8.0%), and other (52, 18.9%). Before 2017, 149 of

Table 2 Clinicopathological factors associated with adjuvant chemotherapy

	Adjuvant chemotherapy (n = 275)	No adjuvant chemotherapy (n = 184)	Univariable P	OR in multivariable analysis*
Age (years), median (i.q.r.)	70 (11.0)	71 (12.5)	0.546	
Male sex	152 (55.3)	98 (53.3)	0.671	
Median Charlson Co-morbidity score	4	4	0.324	
Borderline resectable	15 (5.5)	6 (3.3)	0.270	
Operation			0.167	
Whipple's	103 (37.5)	69 (37.5)		
PPPD	63 (22.9)	27 (14.7)		
DPS	39 (14.2)	33 (17.9)		
DPNS	20 (7.3)	10 (5.4)		
TP	50 (18.2)	45 (24.5)		
Multivisceral resection	41 (14.9)	28 (15.2)	0.928	
Duct location			0.785	
Main duct	161 (58.5)	105 (57.1)		
Side	36 (13.1)	27 (14.7)		
Mixed	62 (22.5)	37 (20.1)		
Tumour location			0.274	
Head	186 (67.6)	106 (57.6)		
Body	28 (10.2)	24 (13.0)		
Tail	39 (14.2)	30 (16.3)		
Tumour size (mm), median (i.q.r.)	23 (18)	22.5 (20)	0.565	
Cyst size (mm), median (i.q.r.)	30 (23.5)	35 (22.8)	0.035	0.94 (0.85, 1.04)
Precursor epithelium			0.706	
Gastric	51 (18.5)	35 (19.0)		
Intestinal	55 (20.0)	42 (22.8)		
Pancreatobiliary	90 (32.7)	55 (29.9)		
Differentiation			0.019	
Well	42 (15.3)	40 (21.7)		1.00 (reference)
Moderately	137 (49.8)	94 (51.1)		1.11 (0.98, 1.25)
Poor	83 (30.2)	35 (19.0)		1.70 (1.07, 2.69)
Invasive component			0.047	
Ductal	213 (77.5)	130 (70.7)		1.00 (reference)
Colloid	34 (12.4)	35 (19.0)		0.93 (0.80, 1.08)
Lymphovascular invasion	151 (54.9)	85 (46.2)	0.067	1.03 (0.91, 1.17)
Perineural invasion	166 (60.4)	90 (48.9)	< 0.001	1.07 (0.96, 1.19)
R1 resection	112 (40.7)	65 (35.3)	0.244	
N1 or N2	150 (54.5)	67 (36.4)	< 0.001	1.15 (0.96, 1.38)
AJCC stage			0.008	
Ia	44 (16.0)	47 (25.5)		1.00 (reference)
Ib	27 (9.8)	25 (13.6)		1.10 (0.91, 1.33)
IIa	37 (13.5)	32 (17.4)		1.06 (0.89, 1.26)
IIb	134 (48.7)	66 (35.9)		1.10 (0.88, 1.37)
III	33 (12.0)	14 (7.6)		1.18 (0.92, 1.51)
Complication CD II	83 (30.2)	46 (25.0)	0.226	
Complication CD ≥ III	27 (9.8)	43 (23.4)	< 0.001	1.02 (0.90, 1.15)

Values are n (%) unless otherwise indicated; *values in parentheses are 95% confidence intervals. PPPD, pylorus-preserving pancreaticoduodenectomy; DPS, distal pancreatectomy with splenectomy; DPNS, distal pancreatectomy without splenectomy; TP, total pancreatectomy; CD, Clavien-Dindo.

268 patients (55.6%) received adjuvant chemotherapy, and the most common regimen was gemcitabine (108, 72.5%) followed by GEM-CAP (13, 8.7%). From 2017, 126 of 191 patients (66.0%) received chemotherapy, and the most common regimens were GEM-CAP (47, 37.3%) followed by gemcitabine (33, 26.2%) and FOLFIRINOX (18, 14.3%). The proportion of patients receiving GEM-CAP and FOLFIRINOX increased in the cohort from 2017 onwards ($P < 0.001$). Gemcitabine was used less frequently compared with before 2017 ($P < 0.001$). The median number of chemotherapy cycles was 6 (i.q.r. 4–6). The proportion of patients who received adjuvant chemotherapy did not differ among the 18 pancreatic centres (Table S1).

Ten patients (2.2%) received adjuvant radiotherapy, nine of whom also received chemotherapy. Seventeen patients (3.7%) also received neoadjuvant chemotherapy with a median of 3 cycles. Of these, 13 also received adjuvant chemotherapy. The most common neoadjuvant chemotherapy regimen was FOLFIRINOX in eight patients. Only 1 of these 17 patients had a borderline resectable tumour.

Recurrence and survival

After a median follow-up of 78 months, the overall recurrence rate was 45.5% (209 of 459). Median time to recurrence was 33 months. Eighty-three patients (18.1%) developed locoregional recurrence after a median follow up of 34 months, including 39 (47.0%) who also developed systemic recurrence. Of these 83 patients, 51 (61.4%) had peripancreatic recurrence and 31 (37.3%) had lymph node recurrence. Some 164 patients (35.7%) developed systemic recurrence after a median follow up of 31 months. Sites of systemic recurrence were liver (66 of 164, 40.2%), lung (58 of 164, 35.4%), peritoneum (48 of 164, 29.3%), and other sites (33 of 164, 20.1%). Recurrence was proven histologically in 40 patients (19.1%). The diagnosis in the other patients was based on imaging. Clinicopathological features of the overall cohort and prognostic factors for recurrence are reported in Table 1.

Single-site recurrence occurred locoregionally in 44 patients (9.6%), in the liver in 38 (8.3%), lung in 30 (6.5%), peritoneum in 18 (3.9%), and at other sites in 15 (3.3%). Multisite recurrence occurred in 64 patients (13.9%).




Fig. 1 Kaplan-Meier survival curves for patients who did or did not receive adjuvant chemotherapy in the matched cohorts

a Overall recurrence, **b** disease-free survival, **c** overall survival, **d** locoregional recurrence, and **e** systemic recurrence since resection: **a** $P = 0.713$, **b** $P = 0.284$, **c** $P = 0.455$, **d** $P = 0.283$, **e** $P = 0.592$ (log rank test).

Median OS for the entire cohort was 39 months, and median DFS was 23 months.

Factors associated with receipt of adjuvant chemotherapy

In univariable analysis, poor differentiation, larger cyst size, ductal invasive component, lymph node positivity, perineural invasion, and higher AJCC stage were positively associated with receipt of adjuvant chemotherapy (Table 2). Conversely, colloid invasive component and complication with a Clavien-Dindo

grade of III or higher were negatively associated with adjuvant chemotherapy receipt. In multivariable analysis, only poor differentiation (OR 1.70; $P = 0.023$) was positively associated with the use of adjuvant chemotherapy (Table 2).

Impact of adjuvant chemotherapy on recurrence and survival

Thirty-one patients (6.8%) died in the first 6 months and were excluded from the landmark analysis. In the unmatched cohorts, 139 of 270 patients who received adjuvant

Table 3 Association between adjuvant chemotherapy and recurrence in high-risk subgroups

	Recurrence rate		P*
	Adjuvant chemotherapy	No adjuvant chemotherapy	
Poor differentiation	53 of 80 (66)	15 of 23 (65)	0.977
Lymphovascular invasion	83 of 146 (58.7)	34 of 63 (53)	0.622
Perineural invasion	96 of 162 (59.3)	36 of 68 (51)	0.673
R1	69 of 108 (63.9)	24 of 46 (52)	0.821
N1 or N2	101 of 149 (67.8)	27 of 49 (55)	0.790
AJCC stage			
Ia	11 of 43 (26)	4 of 47 (8)	0.036†
Ib	7 of 27 (26)	6 of 25 (24)	0.890
IIa	15 of 37 (41)	14 of 32 (42)	0.189
IIb	86 of 132 (65.2)	30 of 50 (60)	0.421
III	20 of 32 (63)	4 of 9 (44)	0.700

Values are n (%). *log rank test. †indicates higher rate of recurrence in adjuvant chemotherapy group.

chemotherapy (51.5%) developed recurrence compared with 58 of 158 (36.7%) who did not receive adjuvant chemotherapy. Among those receiving adjuvant chemotherapy, the 1-, 2-, and 5-year recurrence rates were 20.9% (56 of 268), 42.4% (111 of 262), and 50.3% (86 of 171) respectively. In the group without adjuvant chemotherapy, the rates were 16.5% (26 of 158), 29.1% (44 of 151), and 40.5% (45 of 111).

Before matching, the OS rate was 52.6% (142 of 270) in the adjuvant chemotherapy group compared with 53.8% (85 of 158) in the group without adjuvant chemotherapy. In the adjuvant chemotherapy group, 1-, 2-, and 5-year survival rates were 92.5% (248 of 268), 72.1% (189 of 262), and 51.5% (88 of 171) respectively. In the group without adjuvant chemotherapy the corresponding rates were 87.3% (138 of 158), 74.2% (112 of 151), and 50.5% (56 of 111).

In the adjuvant chemotherapy group, the DFS rate was 41.5% (112 of 270) compared with 50.0% (79 of 158) in the group without adjuvant chemotherapy. In the adjuvant chemotherapy group, 1-, 2-, and 5-year DFS rates were 77.6% (208 of 268), 53.8% (141 of 262), and 42.7% (73 of 171) respectively. Corresponding DFS rates in the group without adjuvant chemotherapy were 78.5% (124 of 158), 63.6% (96 of 151), and 44.1% (49 of 111).

Propensity score-matched analysis

In the PSM postlandmark analysis, 114 patients in the adjuvant chemotherapy group were matched with 114 in the group without adjuvant chemotherapy (Table S2). For all co-variables in the matched cohort, the SMD between treatment and control groups was less than 0.10. The variance ratio of the distance between propensity scores in the matched cohort was 1.01, demonstrating balanced co-variables. Figure S1 illustrates the co-variate balance and Fig. S2 the distribution of propensity scores in the unmatched and matched cohorts.

Univariable analysis in the matched cohort revealed that adjuvant chemotherapy was not associated with overall recurrence (Fig. 1a), DFS (Fig. 1b), OS (Fig. 1c), locoregional recurrence (Fig. 1d) or systemic recurrence (Fig. 1e). Adjuvant chemotherapy was not associated with reduced site-specific recurrence (liver $P=0.275$; lung $P=0.533$; peritoneum $P=0.524$; other sites $P=0.893$).

In the multivariable analysis of the matched cohort, adjuvant chemotherapy was not independently associated with reduced

overall recurrence (HR 0.89, 95% c.i. 0.57 to 1.40; $P=0.584$) (Fig. S3), DFS (HR 0.86, 0.59 to 1.30; $P=0.456$) (Fig. S4) or OS (HR 0.77, 0.50 to 1.20; $P=0.236$) (Fig. S5). In the multivariable analysis, older age (age 60–70 years: HR 4.63, $P=0.005$; age 71–80 years: HR 4.47, $P=0.010$; age over 80 years: HR 5.71; $P=0.007$); higher AJCC stage (IIb: HR 2.74; $P=0.008$), and poor differentiation (HR 2.64; $P=0.008$) were positively associated with overall recurrence.

Subgroup analyses

Table 3 shows the results of the subgroup analysis undertaken to determine whether any particular high-risk group defined by certain characteristics (for example N1–N2 status or AJCC stage) would benefit from adjuvant chemotherapy. In none of these groups were the recurrence rate lower among patients who received adjuvant chemotherapy compared with those who did not.

Type of adjuvant chemotherapy

The recurrence rate was 56.5% (median time to recurrence 32 months) for patients who received gemcitabine, 53.4% for those treated with GEM-CAP (median time to recurrence 23 months), and 54.5% in the FOLFIRINOX group (median time to recurrence 19 months). No particular adjuvant chemotherapy regimen (gemcitabine versus GEM-CAP versus FOLFIRINOX) was associated with a lower recurrence rate, superior DFS or superior OS (Fig. 2).


A separate PSM analysis was carried out for patients who received more contemporary chemotherapy regimens, specifically GEM-CAP or FOLFIRINOX. Fifty-five treated patients (42 GEM-CAP, 13 FOLFIRINOX) were matched with 55 controls (Table S3). In this matched cohort, adjuvant chemotherapy was not associated with reduced overall ($P=0.752$), locoregional ($P=0.346$), or systemic ($P=0.694$) recurrence, or improved DFS ($P=0.863$) or OS ($P=0.823$) (log rank test).

Discussion

In this study, adjuvant chemotherapy did not influence recurrence rates, recurrence patterns or survival in patients with adenocarcinoma arising from IPMN. Among patients who received adjuvant chemotherapy, no particular regimen conferred superior outcomes.


Pancreatic resection followed by adjuvant chemotherapy for PDAC is considered the standard of care, with proven survival benefit in RCTs. Currently, a similar adjuvant treatment strategy is generally considered for patients with adenocarcinoma arising from IPMN, despite a lack of high-level evidence. The data that support adjuvant chemotherapy for adenocarcinoma arising from IPMN are extrapolated from smaller series or multi-institutional databases^{20–24}. Two large institutional data sets published in the past few years have shown a survival benefit in patients with positive lymph node metastasis but not in patients with node-negative disease^{20,23}. A recent systematic review⁶ including 11 studies and comprising of 3393 patients showed that adjuvant chemotherapy was associated with improved OS in patients with node-positive adenocarcinomas arising from IPMN. Improved OS after adjuvant chemotherapy was also demonstrated in patients with stage III–IV disease, tumour size over 2 cm, node-positive status, grade 3 tumour differentiation, positive margin status, tubular carcinoma subtype, and presence of perineural or lymphovascular invasion. These findings were, however, limited by marked heterogeneity, lack of a consistent TNM staging system, and lack of data on chemotherapy regimens that prevented quantitative analysis.

a Recurrence




No. at risk				
FFX	22	15	7	3
GEM	138	100	60	43
GEM-CAP	58	38	19	13

b Disease-free survival



No. at risk				
FFX	22	15	7	3
GEM	138	100	60	43
GEM-CAP	58	38	19	13

c Overall survival



No. at risk				
FFX	22	18	9	4
GEM	138	118	79	52
GEM-CAP	56	46	30	20

Fig. 2 Recurrence, disease-free survival, and overall survival according to type of chemotherapy

a Recurrence **b** disease-free survival, and **c** overall survival. FFX, FOLFIRINOX; GEM, gemcitabine; GEM-CAP, gemcitabine + capecitabine. **a** $P = 0.623$, **b** $P = 0.491$, **c** $P = 0.634$ (log rank test).

In the present study, the recurrence rates were higher in the adjuvant chemotherapy group than in the group without chemotherapy. This was likely related to the preponderance of adjuvant chemotherapy administration among those with adverse tumour factors such as poor differentiation, lymph node positivity, perineural invasion, and higher AJCC stage. After adjusting for confounding variables, there was no reduction in recurrence rate or survival benefit with adjuvant chemotherapy. In further analysis, there was no demonstrated benefit specifically in high-risk groups. Moreover, the type of adjuvant chemotherapy regimen did not affect the risk of recurrence, patterns of recurrence, both locoregional and systemic recurrence, or survival. These findings call into question the benefit of PDAC-derived adjuvant chemotherapy regimens for adenocarcinoma arising from IPMN, even in high-risk groups.

The results of the survival analysis were similar to those of a recent study from Heidelberg, among others^{12,23}. Kaiser *et al.*¹² reported a trend towards better median OS and 5-year survival in patients not receiving chemotherapy with AJCC stage I-IIA disease, and comparable survival with and without chemotherapy in AJCC stage IIB-IV disease.

Since the ESPAC-4 trial¹⁴, adjuvant GEM-CAP has been used routinely in the adjuvant setting and, more recently, FOLFIRINOX²⁵. Interestingly, the ESPAC-4 trial¹⁴ did not specifically include patients with adenocarcinoma arising from IPMN, similar to the more recent ESPAC-5F trial²⁶, which explored the benefits of neoadjuvant treatment in pancreatic adenocarcinoma. In the present study, the rates of GEM-CAP and FOLFIRINOX use increased significantly from 2017 (year of ESPAC-4 trial), with a concomitant reduction in the use of GEM. No particular regimen offered lower rates of recurrence or survival benefit, but it must be noted that the number of patients receiving FOLFIRINOX as monotherapy was small. There is currently no convincing guidance on the type of adjuvant chemotherapy to be used in this group and perhaps this is reflected in the rate of adjuvant chemotherapy across the cohort. The Fukuoka consensus statement²⁷ and the American College of Gastroenterology clinical guidelines²⁸ both make no recommendations on the role of adjuvant chemotherapy, and the recent European guidelines²⁹ recommend adjuvant chemotherapy for adenocarcinoma arising from IPMN with or without nodal disease in the absence of high-level evidence.

There are differences in tumour biology between adenocarcinoma arising from IPMN and PDAC which could explain the difference in response to adjuvant chemotherapy³⁰. Kato *et al.*³¹ performed organoid analyses and identified a distinct set of genetic mutations in adenocarcinoma arising from IPMN compared with PDAC. As well as differing chromatin profiles, they also found the MNX1-HNF1B axis to be critical in regulation of genes in IPMN lineages, unlike PDAC. Gentiluomo *et al.*³² aimed to investigate whether the PDAC susceptibility polymorphisms would also be responsible for the progression to malignancy in IPMNs. In a sample of 345 patients, there was no commonality between 30 susceptibility polymorphisms, only a non-genetic link likely mediated by chronic inflammation through common risk factors (such as smoking and obesity)³³. These inherent differences help explain the relatively indolent behaviour of adenocarcinoma arising from IPMN, reflected in higher survival rates and later recurrence after surgery^{15,34}. These biological differences could explain why adjuvant chemotherapy may not confer any benefit in terms of recurrence or survival^{19,20,35–38}.

Limitations of the present study include the retrospective nature of the study. This introduced selection bias for consideration of adjuvant chemotherapy. Some clinicopathological features may not have been accounted for in the PSM analysis. A significant proportion of patients received mono-gemcitabine therapy, a now outdated adjuvant chemotherapy regimen. Only a small proportion of patients received FOLFIRINOX, so few conclusions can be drawn regarding the potential effect of adjuvant FOLFIRINOX in these patients.

Funding

The authors have no funding to declare.

Author contributions

James Lucocq (CRediT contribution not specified), Jake Hawkyard (Conceptualization, Data curation, Formal analysis), Beate Haugk (Data curation), Omar Mownah (Data curation), Krishna Menon (Conceptualization, Data curation), Takaki Furukawa (Data curation, Resources), Yosuke Inoue (Conceptualization, Data curation), Yuki Hirose (CRediT contribution not specified), Naoki Sasahira (Data curation), Michael Feretis (Data curation), Anita Balakrishnan (Data curation), Carlo Ceresa (Data curation), Brian Davidson (Conceptualization, Data curation), Rupaly Pande (Data curation), Bobby Dasari (Data curation), Lulu Tanno (Data curation), Dimitrios Karavias (Data curation), Jack Helliwell (Conceptualization, Data curation), Alistair Young (CRediT contribution not specified), Quentin Nunes (Data curation), Tomas Urbonas (Data curation), Michael Silva (Data curation), Alex Gordon-Weeks (Data curation), Jenifer Barrie (Data curation), Dhanny Gomez (Data curation), Stijn Van Laarhoven (Data curation), Francis Robertson (CRediT contribution not specified), Hossam Nawara (Data curation), Joseph Doyle (Data curation), Ricky Bhogal (Data curation), Ewen Harrison (Data curation, Supervision, Writing—review & editing), Marcus Roalsø (Data curation), Deborah Ciprani (Data curation, Writing—review & editing), Somaiah Aroori (Data curation), Chathura Ratnayake (Data curation), Jonathan Koea (Data curation), Gabriele Capurso (Data curation, Writing—review & editing), Ruben Bellotti (Data curation, Writing—review & editing), Stefan Stättner (Data curation, Writing—review & editing), Tareq Alsaoudi (CRediT contribution not specified), Neil Bhardwaj (Data curation), Srujan Rajesh (Data curation), Fraser Jeffery (Data curation), Saxon Connor (Data curation), Andrew Cameron (Data curation), Nigel Jamieson (Data curation, Writing—review & editing), Amy Sheen (Data curation), Anubhav Mittal (Data curation), Jaswinder Samra (Data curation), Anthony Gill (Data curation), Keith Roberts (Data curation, Supervision, Writing—review & editing), Kjetil Søreide (Data curation, Supervision, Writing—review & editing), and Sanjay Pandanaboyana (Data curation, Investigation, Methodology, Project administration, Supervision, Validation, Writing—original draft, Writing—review & editing).

Disclosure

The authors declare no conflict of interest.

Supplementary material

Supplementary material is available at BJS online.

Data availability

Data will be available upon reasonable request.

References

- Muraki T, Jang KT, Reid MD, Pehlivanoglu B, Memis B, Basturk O et al. Pancreatic ductal adenocarcinomas associated with intraductal papillary mucinous neoplasms (IPMNs) versus pseudo-IPMNs: relative frequency, clinicopathologic characteristics and differential diagnosis. *Mod Pathol* 2022;**35**: 96–105
- Neoptolemos JP, Stocken DD, Friess H, Bassi C, Dunn JA, Hickey H et al. A randomized trial of chemoradiotherapy and chemotherapy after resection of pancreatic cancer. *N Engl J Med* 2004;**350**:1200–1210
- Seufferlein T, Bachet JB, Van Cutsem E, Rougier P; ESMO Guidelines Working Group. Pancreatic adenocarcinoma: ESMO–ESDO clinical practice guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2012; **23**(Suppl 7):vii33–vii40
- Neoptolemos JP, Stocken DD, Bassi C, Ghaneh P, Cunningham D, Goldstein D et al. Adjuvant chemotherapy with fluorouracil plus folinic acid vs gemcitabine following pancreatic cancer resection: a randomized controlled trial. *JAMA* 2010;**304**: 1073–1081
- Oettle H, Post S, Neuhaus P, Gellert K, Langrehr J, Ridwelski K et al. Adjuvant chemotherapy with gemcitabine vs observation in patients undergoing curative-intent resection of pancreatic cancer: a randomized controlled trial. *JAMA* 2007;**297**:267–277
- Chong E, Ratnayake B, Dasari BVM, Loveday BPT, Siriwardena AK, Pandanaboyana S. Adjuvant chemotherapy in the treatment of intraductal papillary mucinous neoplasms of the pancreas: systematic review and meta-analysis. *World J Surg* 2022;**46**:223–234
- Versteijne E, van Dam JL, Suker M, Janssen QP, Groothuis K, Akkermans-Vogelaar JM et al. Neoadjuvant chemoradiotherapy versus upfront surgery for resectable and borderline resectable pancreatic cancer: long-term results of the Dutch randomized PREOPANC trial. *J Clin Oncol* 2022;**40**: 1220–1230
- Motoi F, Kosuge T, Ueno H, Yamaue H, Satoi S, Sho M et al. Randomized phase II/III trial of neoadjuvant chemotherapy with gemcitabine and S-1 versus upfront surgery for resectable pancreatic cancer (Prep-02/JSAPO5). *Jpn J Clin Oncol* 2019;**49**: 190–194
- Abrams RA, Winter KA, Safran H, Goodman KA, Regine WF, Berger AC et al. Results of the NRG Oncology/RTOG 0848 adjuvant chemotherapy question—erlotinib + gemcitabine for resected cancer of the pancreatic head: a phase II randomized clinical trial. *Am J Clin Oncol* 2020;**43**:173–179
- Chen LT. Chemo-radiotherapy in adjuvant therapy of curatively resected pancreatic cancer: lesions from TCOG T3207 study. *Ann Oncol* 2019;**30**:vi75
- Tempero M, O'Reilly E, Van Cutsem E, Berlin J, Philip P, Goldstein D et al. Phase 3 APACT trial of adjuvant nab-paclitaxel plus gemcitabine (nab-P + Gem) vs gemcitabine (Gem) alone in patients with resected pancreatic cancer (PC): updated 5-year overall survival. *Ann Oncol* 2021;**32**:S226
- Kaiser J, Scheifele C, Hinz U, Leonhardt CS, Hank T, Koenig AK et al. IPMN-associated pancreatic cancer: survival, prognostic staging and impact of adjuvant chemotherapy. *Eur J Surg Oncol* 2022;**48**:1309–1320
- Turrini O, Waters JA, Schnelldorfer T, Lillemoe KD, Yiannoutsos CT, Farnell MB et al. Invasive intraductal papillary mucinous neoplasm: predictors of survival and role of adjuvant therapy. *HPB (Oxford)* 2010;**12**:447–455
- Neoptolemos JP, Palmer DH, Ghaneh P, Psarelli EE, Valle JW, Halloran CM et al. Comparison of adjuvant gemcitabine and capecitabine with gemcitabine monotherapy in patients with resected pancreatic cancer (ESPAC-4): a multicentre, open-label, randomised, phase 3 trial. *Lancet* 2017;**389**: 1011–1024
- Lucocq J, Hawkyard J, Robertson FP, Haugk B, Lye J, Parkinson D et al. Risk of recurrence after surgical resection for adenocarcinoma arising from intraductal papillary mucinous neoplasia (IPMN) with patterns of distribution and treatment: an international, multicentre, observational study. *Ann Surg* 2023; DOI: 10.1097/SLA.0000000000006144 [Epub ahead of print]
- von Elm E, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP et al. The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *J Clin Epidemiol* 2008;**61**:344–349
- Nagtegaal ID, Odze RD, Klimstra D, Paradis V, Rugge M, Schirmacher P et al. The 2019 WHO classification of tumours of the digestive system. *Histopathology* 2020;**76**:182–188
- Amin MB; American Joint Committee on Cancer. *AJCC Cancer Staging Manual* (8th edn). New York: Springer, 2017
- Takeda Y, Imamura H, Yoshimoto J, Fukumura Y, Yoshioka R, Mise Y et al. Survival comparison of invasive intraductal papillary mucinous neoplasm versus pancreatic ductal adenocarcinoma. *Surgery* 2022;**172**:335–342
- McMillan MT, Lewis RS, Drebin JA, Teitelbaum UR, Lee MK, Roses RE et al. The efficacy of adjuvant therapy for pancreatic invasive intraductal papillary mucinous neoplasm (IPMN). *Cancer* 2016;**122**:521–533
- Rodrigues C, Hank T, Qadan M, Ciprari D, Mino-Kenudson M, Weekes CD et al. Impact of adjuvant therapy in patients with invasive intraductal papillary mucinous neoplasms of the pancreas. *Pancreatology* 2020;**20**:722–728
- Hughes DL, Hughes I, Silva MA. Determining the role of adjuvant therapy in invasive intraductal papillary mucinous neoplasms; a systematic review and meta-analysis. *Eur J Surg Oncol* 2022;**48**: 1567–1575
- Mungo B, Croce C, Oba A, Ahrendt S, Gleisner A, Friedman C et al. Controversial role of adjuvant therapy in node negative invasive intraductal papillary mucinous neoplasm. *Ann Surg Oncol* 2021; **28**:1533–1542
- Swartz MJ, Hsu CC, Pawlik TM, Winter J, Hruban RH, Guler M et al. Adjuvant chemoradiotherapy after pancreatic resection for invasive carcinoma associated with intraductal papillary mucinous neoplasm of the pancreas. *Int J Radiat Oncol Biol Phys* 2010;**76**:839–844
- Conroy T, Hammel P, Hebbar M, Ben Abdelghani M, Wei AC, Raoul JL et al. FOLFIRINOX or gemcitabine as adjuvant therapy for pancreatic cancer. *N Engl J Med* 2018;**379**:2395–2406
- Ghaneh P, Palmer D, Cicconi S, Jackson R, Halloran CM, Rawcliffe C et al. Immediate surgery compared with short-course neoadjuvant gemcitabine plus capecitabine, FOLFIRINOX, or chemoradiotherapy in patients with borderline resectable pancreatic cancer (ESPAC5): a four-arm, multicentre, randomised, phase 2 trial. *Lancet Gastroenterol Hepatol* 2023;**8**:157–168
- Tanaka M, Fernández-Del Castillo C, Kamisawa T, Jang JY, Levy P, Ohtsuka T et al. Revisions of international consensus Fukuoka

- guidelines for the management of IPMN of the pancreas. *Pancreatology* 2017;**17**:738–753
28. Vege SS, Ziring B, Jain R, Moayyedi P; Clinical Guidelines Committee; American Gastroenterology Association. American Gastroenterological Association institute guideline on the diagnosis and management of asymptomatic neoplastic pancreatic cysts. *Gastroenterology* 2015;**148**:819–822
 29. European Study Group on Cystic Tumours of the Pancreas. European evidence-based guidelines on pancreatic cystic neoplasms. *Gut* 2018;**67**:789–804
 30. Adsay NV, Merati K, Nassar H, Shia J, Sarkar F, Pierson CR et al. Pathogenesis of colloid (pure mucinous) carcinoma of exocrine organs: coupling of gel-forming mucin (MUC2) production with altered cell polarity and abnormal cell–stroma interaction may be the key factor in the morphogenesis and indolent behavior of colloid carcinoma in the breast and pancreas. *Am J Surg Pathol* 2003;**27**:571–578
 31. Kato H, Tateishi K, Fujiwara H, Nakatsuka T, Yamamoto K, Kudo Y et al. MNX1–HNF1B axis is indispensable for intraductal papillary mucinous neoplasm lineages. *Gastroenterology* 2022;**162**:1272–1287.e16
 32. Gentiluomo M, Corradi C, Arcidiacono PG, Crippa S, Falconi M, Belfiori G et al. Role of pancreatic ductal adenocarcinoma risk factors in intraductal papillary mucinous neoplasm progression. *Front Oncol* 2023;**13**:1172606
 33. Liffers ST, Godfrey L, Frohn L, Haeberle L, Yavas A, Vesce R et al. Molecular heterogeneity and commonalities in pancreatic cancer precursors with gastric and intestinal phenotype. *Gut* 2023;**72**:522–534
 34. Lucocq J, Halle-Smith J, Haugk B, Joseph N, Hawkyard J, Lye J et al. Long-term outcomes following resection of adenocarcinoma arising from intraductal papillary mucinous neoplasm (A-IPMN) versus pancreatic ductal adenocarcinoma (PDAC): a propensity-score matched analysis. *Ann Surg* 2024; DOI:10.1097/SLA.0000000000006272 [Epub ahead of print]
 35. Gavazzi F, Capretti G, Giordano L, Ridolfi C, Spaggiari P, Sollai M et al. Pancreatic ductal adenocarcinoma and invasive intraductal papillary mucinous tumor: different prognostic factors for different overall survival. *Dig Liver Dis* 2022;**54**: 826–833
 36. Yamada S, Fujii T, Hirakawa A, Takami H, Suenaga M, Hayashi M et al. Comparison of the survival outcomes of pancreatic cancer and intraductal papillary mucinous neoplasms. *Pancreas* 2018;**47**:974–979
 37. Yamada S, Fujii T, Hirakawa A, Takami H, Suenaga M, Hayashi M et al. Prognosis of malignant intraductal papillary mucinous tumours of the pancreas after surgical resection. Comparison with pancreatic ductal adenocarcinoma. *Gut* 2002;**51**:717–722
 38. Schnelldorfer T, Sarr MG, Nagorney DM, Zhang L, Smyrk TC, Qin R et al. Experience with 208 resections for intraductal papillary mucinous neoplasm of the pancreas. *Arch Surg* 2008;**143**: 639–646