

Capacity for the management of kidney failure in the International Society of Nephrology Western Europe region: Report from the 2023 ISN Global Kidney Health Atlas (ISN-GKHA)

Maria Pippias PhD, Bristol Medical School: Population Health Sciences, University of Bristol, Bristol, UK; Renal Unit, North Bristol NHS Trust, Bristol, UK

Gaetano Alfano MD, Nephrology Dialysis and Transplant Unit, Azienda Ospedaliero - Universitaria di Modena, Modena, Italy

Dearbhla M. Kelly DPhil, Wolfson Centre for the Prevention of Stroke and Dementia, University of Oxford, John Radcliffe Hospital, Oxford, UK; Department of Intensive Care Medicine, John Radcliffe Hospital, Oxford, UK

Maria Jose Soler MD, Vall d'Hebron University Hospital, Vall d'Hebron Institute of Research, CSUR National Unit of Expertise for Complex Glomerular Diseases of Spain, Barcelona, Spain

Letizia De Chiara PhD, Department of Experimental and Clinical Biomedical Sciences "Mario Serio", University of Florence, Florence, Italy

Timothy O Olanrewaju MD, Division of Nephrology, Department of Medicine, College of Health Sciences, University of Ilorin, Ilorin, Nigeria; Julius Global Health, Julius Center for Health Sciences and Primary Care, University Medical Center Utrecht, Utrecht University, Utrecht, The Netherlands

Silvia Arruebo, The International Society of Nephrology, Brussels, Belgium

Aminu K. Bello PhD,* Division of Nephrology and Immunology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

Fergus J. Caskey MD, Population Health Sciences, Bristol Medical School, University of Bristol, Bristol, UK

Sandrine Damster, The International Society of Nephrology, Brussels, Belgium

Jo-Ann Donner, The International Society of Nephrology, Brussels, Belgium

Vivekanand Jha MD, George Institute for Global Health, University of New South Wales (UNSW), New Delhi, India; School of Public Health, Imperial College, London, UK; Manipal Academy of Higher Education, Manipal, India

David W. Johnson PhD,* Department of Kidney and Transplant Services, Princess Alexandra Hospital, Brisbane, Queensland, Australia; Centre for Kidney Disease Research, University of Queensland at Princess Alexandra Hospital, Brisbane, Queensland, Australia; Translational Research Institute, Brisbane, Queensland, Australia; Australasian Kidney Trials Network at the University of Queensland, Brisbane, Queensland, Australia

Adeera Levin MD, Division of Nephrology, Department of Medicine, University of British Columbia, Vancouver, British Columbia, Canada

Charu Malik PhD, The International Society of Nephrology, Brussels, Belgium

Masaomi Nangaku MD, Division of Nephrology and Endocrinology, The University of Tokyo Graduate School of Medicine, Tokyo, Japan

Ikechi G. Okpechi PhD, Division of Nephrology and Immunology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada; and Division of Nephrology and Hypertension, University of Cape Town, Cape Town, South Africa; Kidney and Hypertension Research Unit, University of Cape Town, Cape Town, South Africa

Marcello Tonelli MD, Department of Medicine, University of Calgary, Calgary, Alberta; Canada and Pan-American Health Organization/World Health Organization's Collaborating Centre in Prevention and Control of Chronic Kidney Disease, University of Calgary, Calgary, Alberta, Canada

Feng Ye MSc, Division of Nephrology and Immunology, Faculty of Medicine and
Dentistry, University of Alberta, Edmonton, Alberta, Canada

Rosanna Coppo MD,⁺ Fondazione Ricerca Molinette, Regina Margherita Hospital,
Turin, Italy

Liz Lightstone PhD,⁺ Centre for Inflammatory Disease, Department of Immunology
and Inflammation, Imperial College London, London, UK

and the Regional Board and ISN-GKHA Team Authors (Appendix 1)

* Co-chairs for International Society of Nephrology Global Kidney Health Atlas (ISN-
GKHA)

⁺ Co-senior authors

Corresponding author:

Maria Pippias: Renal Unit, Southmead Hospital, Southmead Road, Bristol BS10
5NB, UK. Tel: 01174147696, maria.pippias@bristol.ac.uk ORCID: 0000-0001-7469-
4955

Word count without abstract, references, tables, and figures: 4100

Running title: ISN-GKHA – Western Europe

Keywords (3-6): chronic kidney disease, end-stage kidney disease, dialysis, kidney
transplantation, kidney registries, Europe.

Funding Source: This work was supported by the International Society of
Nephrology (Grant RES0033080 to the University of Alberta).

Abstract (248 words)

Western Europe boasts advanced healthcare systems, robust kidney care guidelines, and a well-established healthcare workforce. Despite this, significant disparities in kidney replacement therapy incidence, prevalence, and transplant access exist. This paper presents the third International Society of Nephrology Global Kidney Health Atlas's findings on kidney care availability, accessibility, affordability, and quality in 22 Western European countries, representing 99% of the region's population. The known chronic kidney disease (CKD) prevalence across Western Europe averages 10.6%, slightly above the global median. Cardiovascular diseases account for a substantial portion of CKD-related deaths. Kidney failure incidence varies. Government health expenditure differs, however, most countries offer government-funded acute kidney injury, dialysis, and kidney transplantation care. Hemodialysis and peritoneal dialysis are universally available, with variations in the number of dialysis centers. Kidney transplantation is available in all countries (except for three microstates), with variable transplant center prevalence. Conservative kidney management is increasingly accessible. The region's kidney care workforce is substantial, exceeding global averages, however, workforce shortages are reported. Barriers to optimal kidney care include limited workforce capacity, lack of surveillance mechanisms, and suboptimal integration into national non-communicable disease strategies. Policy recognition of CKD as a health priority varies across countries. While Western Europe exhibits strong kidney care infrastructure, opportunities for improvement exist, particularly in CKD prevention, surveillance, awareness, and policy implementation. Efforts to improve CKD care should include automated detection, educational support, and enhanced workflows.

Based on these findings, healthcare professionals, stakeholders, and policymakers are called to act to enhance kidney care across the region.

Introduction

Western Europe has established functional healthcare systems, produced numerous kidney guidelines, and achieved a higher prevalence of healthcare professionals, greater availability of medicines, and more widespread universal healthcare coverage compared to other world regions.¹ For almost 60 years, the European Renal Association (ERA) Registry has been collecting data on kidney replacement therapy (KRT) modalities [i.e., hemodialysis (HD), peritoneal dialysis (PD), and transplantation] via national and regional kidney registries from 36 countries in Europe.² Despite this high level of kidney care and KRT surveillance systems, the ERA Registry and the International Society of Nephrology Global Kidney Health Atlas (ISN-GKHA) have highlighted significant disparities in kidney care across Western European countries.^{1,3-7} The incidence and prevalence of treated kidney failure, and access to kidney transplantation vary considerably across the region without clear explanation.^{4,7} For people living with kidney failure in Western Europe, HD remains the dominant dialysis modality, with significant variations across countries.^{8,9} Home-therapies, such as home HD and PD, are underutilized with significant prevalence variations across countries,⁸ and access to kidney transplantation is not equitable across Western Europe.³

Since the 2019 ISN-GKHA, kidney medicine across Europe has faced new challenges, including recovery from the COVID-19 pandemic, the war in Ukraine,^{10,11} the Turkey–Syria earthquake and rising global temperatures,¹² with heatwaves seen

across Europe. These challenges have had direct effects on the care of people with kidney disease, though the degree to which they have affected kidney care needs further clarification.

The ISN-GKHA is in its third global iteration. The aim of ISN-GKHA is to understand, compare and monitor how different countries around the world detect, treat, monitor, and advocate for people with kidney disease.¹³ In this second Western Europe-specific report, data from the 2023 ISN-GKHA is presented on the availability, accessibility, affordability, and quality of kidney failure care across the ISN Western Europe region. The methodology for this report is described in detail elsewhere.¹⁴

Results

Results of this study are presented in tables and figures and broadly summarized into two categories: desk research (Table 1,^{15,16} Table 2,^{17,18} Table 3,¹⁹⁻²⁶ Table 4,²⁷⁻⁴⁵ and Supplementary Table S1,⁴⁶) and survey administration (Tables 5-7, Figures 1-4, and Supplementary Tables S2-10).

Study setting

For this report, the ISN's regional classification of countries in Western Europe was utilized, which includes countries bordering the Mediterranean Sea. Western Europe is one of the richest regions in the world, although variability among countries is observed. Luxembourg, Ireland, Liechtenstein, Switzerland, Norway, and Denmark all appear in the top ten countries by gross domestic product (GDP) per capita (Table 1).¹⁵ Greenland and Faroe Islands are considered as their own countries by the World Bank and the data analysis was done on this basis, however

they are currently part of the Kingdom of Denmark. The Social Progress Index (SPI) measures the extent to which countries provide for the social and environmental needs of their citizens. Fifteen of the top twenty countries in the SPI list are in Western Europe,⁴⁷ whilst 14 of the top 20 world happiness report 2023 are Western European countries.⁴⁸

The current status of kidney care in the ISN Western Europe

Western Europe is the world region with the highest prevalence of medical doctors [42.3 [(interquartile range, IQR: 34.9-50.5)] per million population (pmp) and nurses [117.8 pmp (IQR: 88.5-167.8)], compared to the median reported worldwide workforce [doctors, 17.7 pmp (IQR: 3.8-32.87), nurses 36.2 pmp (IQR: 14.4-66.6)] (Supplementary Table S1).⁴⁶ According to the ERA 2020 annual report the incidence of KRT for the treatment of kidney failure is falling in some Western European countries, whilst the prevalence continues to grow, currently standing at 1,173 pmp.⁸ Individuals aged over 65 years make up almost half (45%) of the prevalent KRT population.⁸ Diabetes, hypertension, and renovascular disease account for 26% of the causes of kidney failure in the prevalent population with glomerulonephritis/sclerosis accounting for 18%.⁸ Almost half of the Western European 2020 prevalent KRT population (47%) were living with a kidney transplant.⁸ Five-year adjusted patient survival is 46.8% [95% confidence interval, (95%CI): 46.5-47.1] on dialysis, 92.1% (95%CI: 91.9-92.4) after a first deceased donor transplant and 95.0% (95%CI: 94.6-95.4) after a first living donor transplant.⁸

Literature review data for countries in the ISN Western Europe region

The burden of chronic kidney disease (CKD) and kidney failure

The prevalence of CKD, defined as stages G1-5/A1-3, ranged from 7.9% (95%CI: 7.4-8.6) in Iceland to 14.7% (95%CI: 12.7-16.9) in Greece with a median of 10.6% (IQR: 9.6-11.6) (Table 2).¹⁷ This was slightly higher than the median for all regions at 9.5% (IQR: 5.9-11.7) (Table 2).¹⁷ The proportion of deaths attributed to CKD was less than 2% in the UK (1.3% [95%CI: 1.1-1.3]), Finland (1.3% [95%CI: 1.1-1.4]), Iceland (1.6% [95%CI: 1.3-1.8]) and Norway (1.9% [95%CI: 1.6-2.0]), whereas it was over 4% in Austria (4.1% [95%CI: 3.5-4.5]) and Israel 5.8% [(95%CI: 5.0-6.4)], (Table 2).¹⁷ The median percentage of deaths worldwide attributed to CKD was 2.4% (IQR: 1.6-3.9) (Table 2).¹⁷

Risk factors associated with CKD also varied substantially across Europe with the proportion of obesity ranging from a fifth of the adult population in Switzerland (21.2% [95%CI: 17.3-25.3]) to almost a third in Malta (31.0% [95%CI: 25.1-37.0]). Hypertension ranged from a sixth of the United Kingdom's (UK's) adult population (15.2% [95%CI: 12.2-18.3]) to a quarter of Portugal's adult population (24.4% [95%CI: 18.2-31.2]). One in ten adults were reported to smoke in Sweden (11.4% [95%CI: 10.7-12.2]) whereas this was almost one in three adults in Greece (31.2% [95%CI: 28.9-33.5]) (Table 2).¹⁸

The incidence of the population on KRT failure (dialysis and kidney transplantation) ranged from less than 100 pmp in Ireland, Finland, and Switzerland (88 pmp, 94 pmp and 99 pmp, respectively) to over 250 pmp in Portugal and Greece (260 pmp and 269 pmp, respectively). Similarly, the prevalence of treated kidney failure ranged from less than 900 pmp in Luxembourg, Israel, and Iceland (522 pmp, 755 pmp, and 810 pmp, respectively) to over 1,400 pmp in Greece and Portugal (1,413 pmp and 2,008 pmp, respectively) (Table 3).¹⁹⁻²⁶ Whilst the median incidence of the population

on KRT in the ISN Western Europe region (135.0 pmp [IQR: 106-170]) was lower than the overall global median (145.5 pmp [IQR:107.0-212.5]), the median prevalence of kidney failure was higher (1,034.0 pmp [IQR: 957-1,337] for Western Europe and 822.8 pmp [IQR: 556.0-1114.0] worldwide).¹⁹⁻²⁶

Most individuals commenced KRT on chronic HD, however the incidence differed considerably from less than 70 pmp in Denmark, Sweden, Finland, and Iceland (61.8 pmp, 68.4 pmp, 69.2 pmp, and 69.3 pmp, respectively) to over 140 pmp in France, Belgium, and Greece (146.1 pmp, 175.8 pmp and 255.1 pmp, respectively). Much smaller proportions commenced KRT on PD, with an incidence of less than 15 pmp in Austria, Greece, and Switzerland (10.6 pmp, 12.7 pmp, and 12.9 pmp). Only Sweden and Denmark had an incidence of greater than 35 pmp (35.1 pmp and 36.1 pmp respectively). Very few individuals received pre-emptive kidney transplants, which were less than 5 pmp in Greece, Belgium, and Finland (1.1 pmp, 2.9 pmp and 4.3 pmp, respectively). Pre-emptive kidney transplantation was highest in the UK, Iceland, and the Netherlands (12.8 pmp, 13.9 pmp and 16.8 pmp, respectively) (Table 3).¹⁹⁻²⁶

The prevalence of chronic HD was lowest in the Nordic countries (Iceland, Norway, Finland, Sweden) and the Netherlands (196.9 pmp, 252.2 pmp, 306.8 pmp, 316.1 pmp, 322.7 pmp, respectively) and highest in Belgium, Germany, Portugal, and Greece (708.7 pmp, 768.1 pmp, 871.3 pmp, 1,100.5 pmp, respectively). The prevalence of chronic PD was only 4.7 pmp in Luxembourg, and less than 40 pmp in Germany, Ireland, and Switzerland (38.8 pmp, 39.2 pmp, and 39.3 pmp, respectively), whereas it was over 80 pmp in Sweden, Denmark, and Italy (85.1 pmp,

87.2 pmp, 89.3 pmp, respectively). Three countries had prevalences of individuals living with a kidney transplant of less than 500 pmp, Greece, Italy, and Israel (248.9 pmp, 476.0 pmp, 492.0 pmp, respectively), and three countries had prevalences of over 700 pmp Portugal, UK, and Spain (707.0 pmp, 735.0 pmp, and 751.2 pmp, respectively). Whilst the incidence of living donor kidney transplantation was less than 15 pmp for most countries, it was 26.3 pmp in the Netherlands and 37.2 pmp in Israel (Table 3).¹⁹⁻²⁶

Overview of GDP and government health expenditure

The total health expenditure as a percentage of GDP ranged from 5.4%, 6.7% and 6.7% in Luxembourg, Andorra, and Ireland, respectively, to 11.1%, 11.3% and 11.7% in France, Switzerland, and Germany, respectively, with an average of 9.3%. Total health spending per person ranged six-fold from US\$1,599 in Greece to US\$9,801 in Switzerland, with a median of US\$5,088 (IQR:3,259-5,851) (Table 1).¹⁵ This was considerably higher than the overall health spending of all regions of US\$353 (IQR:76-1,270) (Table 1).¹⁵ Government health spending per person ranged seven-fold from US\$909 in Greece to US\$6,300 in Norway, with a median of US\$3,798 (IQR: 2,470 – 4,701) (Table 1).¹⁶ This resulted in an out-of-pocket health spending per person ranging from US\$358 in Andorra to US\$2,532 in Switzerland with a median spend of US\$637 (IQR: 589-800) per person (Table 1).¹⁶ The demographics of the countries in the ISN Western Europe region can be found in Tables 1-3.¹⁵⁻²⁶

Cost of KRT

The annual cost of HD varied considerably across the region from US\$24,419 in Iceland to US\$94,085 in the UK. Likewise, the annual cost of PD ranged from US\$18,959 in Ireland to US\$83,070 in the UK. The first year after kidney transplantation was cheapest in the UK (US\$19,255) and most expensive in Germany (US\$126,993). Beyond the first year, the differences in the costs of kidney transplantation were striking, ranging from US\$578 in Finland to US\$41,254 in Norway (Table 4).²⁷⁻⁴⁵

Survey response data for the ISN Western Europe region

Characteristics of participating countries

Responses to the survey were received from 22 of 29 countries (75.9%) in the ISN Western Europe region: Andorra, Austria, Belgium, Denmark, Finland, France, Germany, Greece, Iceland, Ireland, Israel, Italy, Liechtenstein, Luxembourg, Malta, the Netherlands, Norway, Portugal, Spain, Sweden, Switzerland, and the UK (Figure 1). These 22 countries had a combined population of 439.5 million people which represented 99% of the region's population and covered 3.8 million square kilometers. All countries were classified as high income by the World Bank, with a GDP based on purchasing power parity ranging from US\$3.3 billion (2021 United States dollars), US\$5 billion, and US\$21.5 billion, in Andorra, Liechtenstein and Iceland, respectively, to US\$3.34 trillion, US\$3.42 trillion, and US\$4.82 trillion in the UK, France, and Germany, respectively (Table 1).^{15,49}

Health financing for kidney care

In eighteen countries (82%), acute kidney injury (AKI), dialysis, and kidney transplantation were publicly funded by the government and free at the point of

delivery (Table 5). Liechtenstein relied solely on a private (health insurance) scheme, whereas transplantation in Ireland was publicly funded by the government but included some fees. Sixteen countries (73%) had healthcare systems that were publicly funded by the government and free at the point of delivery for individuals with non-KRT CKD. Two countries (9%) were publicly funded by the government but included some fees. Two countries (9%) had a mixture of public and private funding systems and only one country (Liechtenstein) had a solely private and out-of-pocket funding system for non-KRT CKD. This was considerably different from the rest of the world where only 27% of non-KRT CKD was publicly funded from the government and 37% was a mixture of public and private financing. Within the ISN Western Europe region, 19 (86%) countries had government publicly funded and free at the point of delivery chronic HD, whereas this was only 45% of countries worldwide (Supplementary Table S2).

In the 2023 ISN-GKHA, funding of medications for CKD were free at the point of delivery in Austria, Denmark, Italy, and Malta (18%), whereas funding for medications for dialysis were free at the point of delivery in Austria, Denmark, Greece, Italy, Israel, Malta, and Switzerland (32%) (Table 5). Medications for people with CKD and for people receiving dialysis were publicly funded by the government but with some fees in 55% and 45% of countries, respectively. These were much higher rates of government funding of medications than the worldwide median (33% and 28%, respectively). Likewise, 72% of medications for kidney transplant recipients were either publicly funded and free at the point of delivery or with some fees (Table 5), which was higher than the worldwide median of 52%.

Availability of services for the delivery of kidney care

HD and PD were available in all countries of the ISN Western Europe region. The median number of chronic HD centers was 7.7 pmp though this ranged from 1.0 pmp in the UK to 19.0 pmp in Greece (and 25.2 pmp in the microstate of Liechtenstein), (Table 6, Figure 2). Home HD for adults and pediatrics was available in 77% of countries (generally not available in Iceland and Israel and not available in Greece, Luxembourg, or Malta). This was higher than the worldwide average of 19%. Chronic PD was available in all countries (2.7 pmp), however acute PD was only available in 15 countries (68%) (Supplementary Table S3).

Kidney transplantation, both from living and deceased donors, was available in 18 countries (but not in the microstates and small countries of Andorra, Liechtenstein, or Luxembourg) (Table 6, Figure 2). These 18 countries all had national waiting lists except Spain which only had regional waiting lists. The availability of kidney transplantation centers for the region's countries was 0.69 pmp, ranging from 0.12 pmp in Switzerland to 0.85 pmp in Spain. Kidney transplantation centers were more common in Malta (2.2 pmp) and Iceland (2.8 pmp), however both countries had populations of fewer than 500,000. Pediatric transplantation was available in 17 countries (77%).

Conservative kidney management (CKM), defined as a planned, holistic, patient-centered care for people living with CKD stage 5 which was a shared decision between the healthcare worker and the patient, was generally available in 21 (95%) countries. This is higher than the worldwide median of 45% (Supplementary Table S4).

Health workforce for the delivery of kidney care

In all countries bar two microstates (91%), nephrologists were primarily responsible for the medical care of individuals with kidney failure. This was slightly higher than the overall global average of 87%. The median number of nephrologists in the ISN Western Europe region was 25.0 (IQR: 16.4-33.9) pmp, which was over double the median number worldwide (11.8 pmp [IQR: 1.8-24.8]). Within the region, the median prevalence of nephrologists ranged from 11.8 pmp in Ireland to 53.0 pmp in Spain (excluding the microstates). Likewise, the median prevalence of nephrology trainees ranged from 0.59 pmp in Germany to 49.1 pmp in Italy (Table 6). The median prevalence of pediatric nephrologists was 1.6 pmp (IQR: 0.96-2.2), which was over double the global median prevalence of pediatric nephrologists (0.69 pmp [IQR:0.03-1.8]).

There were widespread shortages of healthcare providers for kidney failure across the region (Figure 3). Only 10 countries (45%) reported sufficient nephrologists, and only 5 countries (23%) reported sufficient pediatric nephrologists. Sixteen countries (73%) reported a shortage of transplant surgeons and hemodialysis access surgeons, and 17 countries (77%) reported a shortage of surgeons for PD access. Thirteen countries (59%) required additional counsellors or psychologists and 13 countries (59%) required additional dialysis nurses.

Health information systems, early identification mechanisms for kidney disease, advocacy, and policy for kidney disease

Availability of nephrologists (59%), patient knowledge and attitude (55%), physician availability, access, knowledge, and attitude (50%), and a lack of political will and enabling policies (45%) were the top four cited barriers to optimal kidney care in the ISN Western Europe region (Supplementary Table S5).

Country level national health policies for kidney care were variable in the region with fourteen countries (64%) stating that their country had a national strategy for non-communicable diseases (NCD) (Supplementary Table S6). Only ten countries (45%) stated that their country had a national strategy for improving the care of people with CKD with half saying that this strategy was incorporated into a NCD strategy (Supplementary Table S6). Only five countries (23%) felt that CKD was a recognized health priority by their country's government which was almost half that of the worldwide opinion (48%). This was the same for AKI (23%) though the worldwide opinion was also lower at 19%. The number of countries that felt their governments recognized kidney failure and/or KRT as a health priority was higher at 64% but this still left eight countries (36%) believing it was not recognized as a health priority.

Dialysis and transplant registries were available in all countries except for the microstates of Andorra and Liechtenstein. CKD registries were less prevalent, only being available in six countries (27%), whereas AKI and CKM registries were only available in two countries (9%) (Table 7).

Causes of hospitalization and death among people living with kidney failure on dialysis

Seventy-three percent of countries reported a mortality of less than 10% in the first year of HD and PD. The commonest cause of death was cardiovascular disease (Supplementary Tables S7-S8). Approximately half of the countries reported less than 30% of people receiving HD or PD had at least one hospitalization event in the first year of dialysis. The commonest causes of hospitalization amongst people receiving HD were cardiovascular disease (59% of countries), and access malfunction (14% of countries) (Supplementary Table S9). The commonest cases of hospitalization amongst people receiving PD were cardiovascular disease (32% of countries), and PD-related infection (23% of countries) (Supplementary Table S10).

Discussion

The aim of ISN-GKHA is to understand, compare and monitor how different countries around the world detect, treat, monitor, and advocate for people with kidney disease.¹³ In this paper, data from the third ISN-GKHA are presented with a focus on the ISN Western Europe region. Compared to the other regional median values, Western Europe is performing well, however there are many opportunities for improvement, specifically related to CKD prevention, surveillance, awareness, and policy implementation.

The median prevalence of CKD across the ISN Western Europe region was 10.6%, slightly higher than the median worldwide at 9.5%. Within the region, there was a two-fold variation between the country with the lowest prevalence, Iceland, and the country with the highest prevalence, Greece. Though caution is required when comparing CKD prevalence due to significant methodological variations in the general population sampling methods and variability in the assessment of kidney

function within studies,⁵⁰ one must still acknowledge that the data are pointing towards one in ten Western Europeans having CKD. CKD is associated with adverse health outcomes, high cardiovascular mortality, and high healthcare costs. The 2017 Global Burden of Disease study reported a 41.5% increase in the global all-age mortality from CKD between 1990 and 2017.⁵¹ The same study reported that most of the burden of CKD was concentrated in the three lowest quintiles of the socio-demographic Index. Identifying and treating CKD at the earliest stages will not only likely reduce the risk of progression to kidney failure and (cardiovascular) mortality, it is also an equity imperative.⁵¹ A recent Kidney Disease Improving Global Outcomes (KDIGO) Controversies Conference recommended that individuals with hypertension, diabetes, or cardiovascular disease should be screened for CKD.⁵² Screening the general population for CKD detection is not currently supported by evidence of benefit over costs.^{53,54} We reported high levels of obesity, hypertension, and smoking across Western Europe, all of which are known risk factors for the development of CKD. Alongside screening for CKD in high-risk populations, preventative measures based around tackling CKD risk factors, such as obesity, hypertension, and smoking, could be an effective CKD preventative measure.

As shown in this study, the ISN Western Europe region has highly developed programs to manage the care of the (relatively small) kidney failure population, with 63% of survey participants reporting that their government recognized kidney failure and KRT as a health priority. Programs to manage the care of the much larger non-KRT-dependent CKD population are less well developed.⁵⁵ Only 23% felt that CKD was a recognized health priority by their country's government, which was almost half that of the worldwide opinion (48%). Less than half of the countries surveyed

reported that their country had a national strategy for improving the care of people with CKD with half reporting that this was incorporated into a NCD strategy. Less than a quarter of countries had a national AKI strategy. Even at the European Union (EU) level, CKD was not considered in the 2022 Healthier Together EU NCD Initiative, whereas diabetes, cardiovascular disease, chronic respiratory, mental health, and neurological diseases were.⁵⁶

Bello et al. reported common barriers to the care of people with non-KRT-dependent CKD in 16 Western European countries.⁵⁵ These were: limited work force capacity, almost a total lack of disease surveillance mechanisms, absence of coordinated CKD care strategies, poor CKD care integration with other NCD control initiatives, and low awareness of the significance of CKD. They suggested potential approaches to improving CKD care by making changes to health systems and policy. These included better models of CKD care, building workforce capacity, incorporating CKD care into (inter)national NCD strategies, and population-based national CKD registries amongst others.⁵⁵

Compared to the other regions, the ISN Western Europe region has a much larger medical workforce, typically double the number of adult and pediatric nephrologists compared to the world average and higher government healthcare expenditure. Despite this, 59% of survey participants reported the insufficient availability of nephrologists as one of the main barriers to optimal kidney care. Whilst increasing the workforce is the obvious solution here, other mechanisms to streamline and improve the delivery of evidence-based CKD care may also play a role. Taylor et al. recently reported on the contexts, mechanisms, and outcomes

involved in complex interventions to improve CKD care at the primary and secondary care interface.⁵⁷ Key intervention components identified were automated identification of higher-risk individuals in primary care with management advice provided to general practitioners, educational support systems, and non-patient-facing nephrologist review. These components had the potential to promote clinician learning whilst managing individuals with CKD, promote clinician motivation to take steps toward evidence-based CKD management, and integrate dynamically with existing workflows.⁵⁷

Just under half of countries reported a lack of political will and enabling policies as a barrier to optimal kidney care in the ISN Western Europe region. Bello et al. reported low awareness of the importance of CKD among the public, policy makers and care providers, perpetuating the limitations in optimal kidney care.⁵⁵ Surveillance systems such as kidney registries provide data on disease and treatment rates which are needed to advocate for, and support the planning, delivery, and evaluation of kidney services. The data can also highlight discrepancies in available care allowing for the diversion of resources to where they are most needed.⁵⁸ This region is in a fortunate position of having dialysis and kidney transplant registries available in all countries (except for two microstates). Additionally, Europe is fortunate to have a European level registry which then collates and presents data allowing for international comparisons and benchmarking.² Within the ISN Western Europe region, CKD, AKI, and CKM registries were less prevalent, only being available in six (27%), two (9%) and two (9%) countries, respectively. Developing CKD and AKI registries will focus attention on, and effort towards prevention, whilst data on CKM, be that from patient choice or

limited resources, are necessary to advocate for this vulnerable, high-risk group. Though some efforts are being taken to develop these registries,⁵⁹ collecting both the right data and accurate data,⁶⁰ especially in the light of limited systematic early CKD detection programs, does hamper their development. The ISN's SHARing Expertise to support the set-up of Renal Registries (SharE-RR) project provides education and support to further establishment and development of kidney registries worldwide.⁵⁹

Whilst it is clear that the ISN Western Europe region is performing well in all aspects of kidney care compared with the other nine ISN regions, improvements are certainly possible and welcome. Western European kidney healthcare professionals, key stakeholders, and policy makers are called upon to prioritize CKD prevention through awareness campaigns, mitigation of CKD risk factors, and the implementation of surveillance systems, including registries for AKI and non-KRT CKD. These efforts are essential to continue the current positive trend of decreasing the incidence of KRT, a condition that is associated, in both children and adults, with dire clinical and social consequences as well as massive economic costs.⁶¹ For people reaching kidney failure, there should be a focus on patient-centered choice for treatments such as CKM and equitable access to home therapies i.e., home HD and PD. Collaborative cooperation is essential to minimize disparities between Western European countries and ensure high standard of treatment for the CKD population living in this region.

There were some limitations to the 2023 ISN-GKHA; these have been discussed.¹⁴ One such limitation specific to our region is the lack of more detailed information on paediatric kidney care. Available studies corroborate findings in adults including the availability of pediatric kidney transplantation in 93% of countries,

pediatric dialysis facilities for AKI and kidney failure in 95% of countries, and considerable variations in the availability of allied health professionals, high-tech diagnostic methods, and treatment with expensive drugs for children.⁶² However, this work is important for guiding kidney care policy in the ISN Western Europe region.

Role of the Funder/Sponsor

The ISN provided administrative support for the design and implementation of the study and data collection activities. The authors were responsible for data management, analysis, and interpretation, as well as manuscript preparation, review, and approval, and the decision to submit the manuscript for publication.

Disclosure

GA reports personal fees from Baxter, outside the submitted work. MSJ reports grants from Boehringer, Instituto Carlos III, and Marato TV3; personal fees (honoraria) from NovoNordisk, Jansen, Boehringer, Mundipharma, AstraZeneca, Ingelheim Lilly, Vifor, ICU Medical, Fresenius, and Travers Therapeutics, travel support from Travers and Vifor; patent 202131356; participation on Advisory Boards for NovoNordisk, Jansen, Boehringer, Mundipharma, AstraZeneca, Ingelheim Lilly, Vifor, ICU Medical, Bayer, GE Healthcare, and Travers Therapeutics, outside the submitted work. SA reports personal fees (salary) from The International Society of Nephrology, outside the submitted work. AKB reports other (consultancy and honoraria) from AMGEN Incorporated and Otsuka, other (consultancy) from Bayer and GSK, and grants from Canadian Institute of Health Research and Heart and Stroke Foundation of Canada, outside the submitted work; he is also Associate Editor of the Canadian Journal of Kidney Health and Disease and Co-chair of the

ISN-Global Kidney Health Atlas. SD reports personal fees (salary) from The International Society of Nephrology, outside the submitted work. JD reports personal fees (salary) from The International Society of Nephrology, outside the submitted work. VJ reports personal fees from GSK, Astra Zeneca, Baxter Healthcare, Visterra, Biocryst, Chinook, Vera, and Bayer, paid to his institution, outside the submitted work. DWJ reports consultancy fees, research grants, speaker's honoraria and travel sponsorships from Baxter Healthcare and Fresenius Medical Care, consultancy fees from Astra Zeneca, Bayer, and AWAK, speaker's honoraria from ONO and Boehringer Ingelheim & Lilly, and travel sponsorships from Ono and Amgen, outside the submitted work. He is also a current recipient of an Australian National Health and Medical Research Council Leadership Investigator Grant, outside the submitted work. CM reports personal fees (salary) from The International Society of Nephrology, outside the submitted work. MN reports grants and personal fees from KyowaKirin, Boehringer Ingelheim, Chugai, Daiichi Sankyo, Torii, JT, Mitsubishi Tanabe, grants from Takeda and Bayer, and personal fees from Astellas, Akebia, AstraZeneca, and GSK, outside the submitted work. LL reports personal fees from Alexion, Astra Zeneca, Biogen Idec, BMS, GSK, Kezar, Novartis, Pfizer, Roche, and Otsuka, outside the submitted work; and Scientific advisor and investor in Carna Health. All others have nothing to declare.

Acknowledgements

This article is published as part of a supplement supported by the International Society of Nephrology (ISN; grant RES0033080 to the University of Alberta). The ISN provided administrative support for the design and implementation of the survey and data collection activities. The authors were responsible for data

management, analysis, and interpretation, as well as manuscript preparation, review, and approval, and the decision to submit the manuscript for publication.

The authors appreciate the support from the ISN's Executive Committee, regional leadership, and Affiliated Society leaders at the regional and country levels for their help with the ISN-GKHA.

SUPPLEMENTARY MATERIAL

Supplementary File (PDF)

Table S1. Workforce (specialist physician, doctor, and nurse) prevalences in the ISN Western Europe region.

Table S2. Healthcare system funding structure for kidney replacement therapy in the ISN Western Europe region [n (100%)].

Table S3. Availability of in-center hemodialysis, home hemodialysis, and dialysis in the ISN Western Europe region [n (100%)].

Table S4. Availability of choice-restricted conservative kidney management* in the ISN Western Europe region [n (100%)].

Table S5. Barriers to optimal kidney care, overall and in the ISN Western Europe region [n (100%)].

Table S6. National health strategies for non-communicable diseases and CKD-specific policies in the ISN Western Europe region [n (100%)].

Table S7. Commonest causes of death among people receiving hemodialysis in the ISN Western Europe region [n (100%)].

Table S8. Commonest causes of death among people receiving peritoneal dialysis in the ISN Western Europe region [n (100%)].

Table S9. Commonest causes of hospitalization amongst people receiving hemodialysis in the ISN Western Europe region [n (100%)].

Table S10. Commonest causes of hospitalization amongst people receiving peritoneal dialysis in the ISN Western Europe region [n (100%)].

Figure S1. Country-level scorecard on the availability of KRT, CKM, funding for medications, registry, advocacy group, nephrologists and nephrology trainees in the ISN Western Europe region in 2019 and 2023.

References

1. Kelly DM, Anders HJ, Bello AK, et al. International Society of Nephrology Global Kidney Health Atlas: structures, organization, and services for the management of kidney failure in Western Europe. *Kidney Int Suppl* (2011). 2021;11(2):e106-e18.
2. Jager KJ, Wanner C. Fifty years of ERA-EDTA Registry-a registry in transition. *Kidney Int Suppl* (2011). 2015;5(1):12-4.
3. Boenink R, Kramer A, Masoud S, et al. International comparison and time trends of first kidney transplant recipient characteristics across Europe: an ERA Registry study. *Nephrol Dial Transplant*. Published online August 31, 2023. doi:10.1093/ndt/gfad189
4. Boenink R, Kramer A, Tuinhout RE, et al. Trends in kidney transplantation rate across Europe: study from the ERA Registry. *Nephrol Dial Transplant*. 2023;38(6):1528-39.
5. Bonthuis M, Cuperus L, Chesnaye NC, et al. Results in the ESPN/ERA-EDTA Registry suggest disparities in access to kidney transplantation but little variation in graft survival of children across Europe. *Kidney Int*. 2020;98(2):464-75.
6. Bonthuis M, Vidal E, Bjerre A, et al. Ten-year trends in epidemiology and outcomes of pediatric kidney replacement therapy in Europe: data from the ESPN/ERA-EDTA Registry. *Pediatr Nephrol*. 2021;36(8):2337-48.
7. Pippias M, Jager KJ, Kramer A, et al. The changing trends and outcomes in renal replacement therapy: data from the ERA-EDTA Registry. *Nephrol Dial Transplant*. 2016;31(5):831-41.
8. Astley ME, Boenink R, Abd ElHafeez S, et al. The ERA Registry Annual Report 2020: a summary. *Clin Kidney J*. 2023;16(8):1330-54.

9. Torreggiani M, Piccoli GB, Moio MR, et al. Choice of the Dialysis Modality: Practical Considerations. *J Clin Med*. 2023;12(9).
10. Pawlowicz-Szlarska E, Vanholder R, Sever MS, et al. Distribution, preparedness and management of Ukrainian adult refugees on dialysis-an international survey by the Renal Disaster Relief Task Force of the European Renal Association. *Nephrol Dial Transplant*. 2023;38(10):2407-15.
11. Tuglular S, Luyckx V, Vanholder R, et al. Lessons learned during the war in Ukraine: a report from the Renal Disaster Relief Task Force of the ERA. *Nephrol Dial Transplant*. 2023;38(9):1960-8.
12. Vanholder R, Agar J, Braks M, et al. The European Green Deal and nephrology: a call for action by the European Kidney Health Alliance. *Nephrol Dial Transplant*. 2023;38(5):1080-8.
13. Bello AK, Johnson DW, Feehally J, et al. Global Kidney Health Atlas (GKHA): design and methods. *Kidney Int Suppl* (2011). 2017;7(2):145-53.
14. Okpechi IG, Bello AK, Levin A, et al. An update on the variability in the organization and structures of kidney care across world region. (submitted to *Kidney Int Suppl*, this issue). 2023.
15. Central Intelligence Agency. The World Factbook. Available at: <https://www.cia.gov/the-world-factbook/>. Accessed April 8, 2022.
16. The Institute for Health Metrics and Evaluation. Global Health Data Exchange. Global Expected Health Spending 2019-2050. Available at: <https://ghdx.healthdata.org/record/ihme-data/global-expected-health-spending-2019-2050>. Accessed April 8, 2022.
17. Institute for Health Metrics and Evaluation (IHME). Global Burden of Disease Collaborative Network. Global Burden of Disease Study 2019 (GBD 2019)

- Results. 2020. Available at: <https://vizhub.healthdata.org/gbd-results/>. Accessed April 8, 2022.
18. World Health Organization. Global Health Observatory. Available at: <https://www.who.int/data/gho>. Accessed April 8, 2022.
 19. European Renal Association. ERA Registry Annual Report 2019. 2019. Available at: <https://www.era-online.org/wp-content/uploads/2022/11/ERA-Registry-Annual-Report-2019.pdf>. Accessed April 8, 2022.
 20. Finnish Registry for Kidney Diseases Report 2020. Available at: https://www.muma.fi/files/5627/Finnish_Registry_for_Kidney_Diseases_Report_2020.pdf . Accessed April 8, 2022.
 21. Jain AK, Blake P, Cordy P, et al. Global trends in rates of peritoneal dialysis. J Am Soc Nephrol. 2012;23(3):533-544.
 22. Nefrovisie, Utrecht, the Netherlands 2021. Renine Registry Annual Report 2021. Available at: https://www.nefrovisie.nl/wp-content/uploads/2022/02/Jaarrapportage_Renine_2020_web.pdf. Accessed April 8, 2022.
 23. The Norwegian Renal Registry. Annual Report 2020. Available at: https://www.nephro.no/nnr/AARSRAPPORT_NNR_2020.pdf. Accessed April 8, 2022.
 24. The Renal Association. UK Renal Registry 23rd Annual Report. Available at: https://ukkidney.org/sites/renal.org/files/23rd_UKRR_ANNUAL_REPORT.pdf. Accessed April 8, 2023.
 25. United States Renal Data System. 2019 USRDS Annual Data Report: Epidemiology of kidney disease in the United States. National Institutes of Health, National Institute of Diabetes and Digestive and Kidney Diseases,

- Bethesda, MD, 2019. Available at: <https://www.niddk.nih.gov/about-niddk/strategic-plans-reports/usrds/prior-data-reports/2019>. Accessed April 8, 2022.
26. WHO-ONT Global Observatory on Donation and Transplantation. Available at: <http://www.transplant-observatory.org/data-charts-and-tables/>. Accessed April 8, 2022.
 27. Cavallo MC, Sepe V, Conte F, et al. Cost-effectiveness of kidney transplantation from DCD in Italy. *Transplant Proc* 2014; 46(10):3289-96.
 28. Ferguson TW, Harper GD, Milad JE, et al. Cost of the quantia SC+ hemodialysis system for self-care in the United Kingdom. *Hemodial Int*, 2022; 26(3):287-294.
 29. Häckl D, Kossack N, Schoenfelder T. Prävalenz, Kosten der Versorgung und Formen des dialysepflichtigen chronischen Nierenversagens in Deutschland: Vergleich der Dialyseversorgung innerhalb und außerhalb stationärer Pflegeeinrichtungen [Prevalence, Costs of Medical Treatment and Modalities of Dialysis-dependent Chronic Renal Failure in Germany: Comparison of Dialysis Care of Nursing Home Residents and in Outpatient Units]. *Gesundheitswesen* 2021; 83(10):818-28.
 30. Haller M, Gutjahr G, Kramar R, et al. Cost-effectiveness analysis of renal replacement therapy in Austria. *Nephrol Dial Transplant*. 2011; 26(9):2988-95.
 31. Helanterä I, Isola T, Lehtonen TK, et al. Association of Clinical Factors with the Costs of Kidney Transplantation in the Current Era. *Ann Transplant*. 2019; 24:393-400.

32. Jakobsen A, Albrechtsen D, Sødal G, et al. Kostnader ved uremikeromsorg. Hva koster nyretransplantasjon? [Costs of care in uremia. How much does kidney transplantation cost?]. Tidsskr Nor Laegeforen 1990; 110(3):338-41.
33. Jarl J, Desatnik P, Peetz Hansson U, et al. Do kidney transplantations save money? A study using a before-after design and multiple register-based data from Sweden. Clin Kidney J. 2018; 11(2):283-88.
34. Jensen CE, Sørensen P, Petersen KD. In Denmark kidney transplantation is more cost-effective than dialysis. Dan Med J. 2014; 61(3):A4796.
35. Jürgensen JS, Ikenberg R, Greiner RA, et al. Cost-effectiveness of modern mTOR inhibitor based immunosuppression compared to the standard of care after renal transplantation in Germany. Eur J Health Econ 2015; 16(4):377-90.
36. Kerr M, Bray B, Medcalf J, et al. Estimating the financial cost of chronic kidney disease to the NHS in England. Nephrol Dial Transplant. 2012; 27 Suppl 3(Suppl 3):iii73-80.
37. Li B, Cairns JA, Fotheringham J, et al. Understanding cost of care for patients on renal replacement therapy: looking beyond fixed tariffs. Nephrol Dial Transplant. 2015; 30(10):1726–34.
38. Mohnen SM, van Oosten MJM, Los J, et al. Healthcare costs of patients on different renal replacement modalities - Analysis of Dutch health insurance claims data. PLoS One. 2019; 14(8):e0220800.
39. Roberts G, Holmes J, Williams G, et al. Current costs of dialysis modalities: A comprehensive analysis within the United Kingdom. Perit Dial Int. 2022; 42(6):578-84.

40. Rocha MJ, Ferreira S, Martins LS, et al. Cost analysis of renal replacement therapy by transplant in a system of bundled payment of dialysis. *Clin Transplant*. 2012; 26(4):529-31.
41. Sandoz MS, Ess SM, Keusch GW, et al. Prevalence and direct medical costs of end-stage renal disease in patients with type 2 diabetes mellitus in Switzerland for 2001. *Swiss Med Wkly*. 2004; 134(31-32):448-58.
42. Van Biesen W, Lameire N, Peeters P, et al. Belgium's mixed private/public health care system and its impact on the cost of end-stage renal disease. *Int J Health Care Finance Econ*. 2007; 7(2-3):133-48.
43. van der Tol A, Lameire N, Morton RL, et al. An International Analysis of Dialysis Services Reimbursement. *Clin J Am Soc Nephrol*. 2019; 14(1):84-93.
44. Villa G, Rodríguez-Carmona A, Fernández-Ortiz L, et al. Cost analysis of the Spanish renal replacement therapy programme. *Nephrol Dial Transplant*. 2011; 26(11):3709-14.
45. Zambrowski JJ. Coût de la dialyse [Cost of dialysis in France]. *Nephrol Ther*. 2016; 12 Suppl 1:S95-97.
46. World Health Organization Global Health Observatory. Health workforce 2021. Available at: <https://www.who.int/data/gho/data/themes/health-workforce>. Accessed April 8, 2022.
47. Social Progressive Imperative. Global Index: Results. Available at: <https://www.socialprogress.org/index/global/results/>. Accessed October 6, 2023.
48. Sustainable Development Solutions Network. World Happiness Report. Helliwell JF, Layard R, Sachs J, et al. Published 2023. Available at: <https://worldhappiness.report/ed/2023/>. Accessed October 6, 2023.

49. The World Bank. World Development Indicators. The World by Income and Region. Available at: <https://datatopics.worldbank.org/world-development-indicators/the-world-by-income-and-region.html>. Accessed April 8, 2022.
50. Bruck K, Jager KJ, Dounousi E, et al. Methodology used in studies reporting chronic kidney disease prevalence: a systematic literature review. *Nephrol Dial Transplant*. 2015;30 Suppl 4(Suppl 4):iv6-16.
51. GBD Chronic Kidney Disease Collaboration. Global, regional, and national burden of chronic kidney disease, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet*. 2020;395(10225):709-733.
52. Shlipak MG, Tummalapalli SL, Boulware LE, et al. The case for early identification and intervention of chronic kidney disease: conclusions from a Kidney Disease: Improving Global Outcomes (KDIGO) Controversies Conference. *Kidney Int*. 2021;99(1):34-47.
53. Jain V, Sinha S, Shaw C, et al. Re-evaluating national screening for chronic kidney disease in the UK. *BMJ*. 2023;382:e074265.
54. Bochud M. On the rationale of population screening for chronic kidney disease: a public health perspective. *Public Health Rev*. 2015;36:11.
55. Bello AK, Levin A, Manns BJ, et al. Effective CKD care in European countries: challenges and opportunities for health policy. *Am J Kidney Dis*. 2015;65(1):15-25.
56. Directorate-General for Health and Food Safety. EU Non-communicable diseases (NCD) Initiative: Guidance document. Published 2022. Available at: https://health.ec.europa.eu/publications/eu-non-communicable-diseases-ncds-initiative-guidance-document_en. Accessed October 6, 2023.

57. Taylor DM, Nimmo AM, Caskey FJ, et al. Complex Interventions Across Primary and Secondary Care to Optimize Population Kidney Health: A Systematic Review and Realist Synthesis to Understand Contexts, Mechanisms, and Outcomes. *Clin J Am Soc Nephrol*. 2023;18(5):563-72.
58. Alwan A, Maclean DR, Riley LM, et al. Monitoring and surveillance of chronic non-communicable diseases: progress and capacity in high-burden countries. *Lancet*. 2010;376(9755):1861-8.
59. Hole BD, Evans KM, Pyart R, et al. International collaborative efforts to establish kidney health surveillance systems. *Kidney Int*. 2020;98(4):812-6.
60. Nistor I, Bolignano D, Haller MC, et al. Why creating standardized core outcome sets for chronic kidney disease will improve clinical practice. *Nephrol Dial Transplant*. 2017;32(8):1268-73.
61. Kidney Research UK. Kidney disease: A UK public health emergency. The health economics of kidney disease to 2033. Published 2023. Available at: https://www.kidneyresearchuk.org/wp-content/uploads/2023/06/Economics-of-Kidney-Disease-full-report_accessible.pdf. Accessed October 6, 2023.
62. Prikhodina L, Ehrich J, Shroff R, et al. The European Society for Paediatric Nephrology study of pediatric renal care in Europe: comparative analysis 1998-2017. *Pediatr Nephrol*. 2020 Jan;35(1):103-111.

Tables

Table 1. Demographics and health finance in the ISN Western Europe region.^{15,16}

Country	Area (sq km)	Total population (2022)	GDP (PPP) (\$ billion)	Total health expenditures (% of GDP)	Total Health Spending per person*	Government Health Spending per person*	Out-of-pocket Health Spending per person*
Overall, median [IQR]	130,483,015	7,802,702,984	133.8 [39.7 - 545.0]	6.2 [4.3-8.2]	353 [76 -1270]	216 [23 - 908]	92 [29 - 273]
ISN Western Europe region, median [IQR]	3,801,668	439,465,153	529.3 [351.7 - 1523.9]	9.5 [8.2 - 10.5]	5088 [3259 - 5851]	3798 [2470 - 4701]	637 [589 - 800]
Andorra	468	85,560	3.3	6.7	2,875	1,993	358
Austria	83,871	8,913,088	523.3	10.4	5,721	4,361	907
Belgium	30,528	11,847,338	682.9	10.7	5,088	3,998	848
Denmark	43,094	5,920,767	378.6	10	6,300	5,392	767
Finland	357,022	5,601,547	304.8	9.2	4,553	3,666	755
France	643,801	68,305,148	3,424.2	11.1	4,838	3,629	403
Germany	357,022	84,316,622	4,815.5	11.7	5,838	4,701	627
Greece	131,957	10,533,871	333.7	7.8	1,599	909	514
Iceland	572	357,603	21.5	8.6	5,670	4,779	800
Ireland	70,723	5,275,004	535.3	6.7	6,128	4,690	637
Israel	20,770	8,914,885	409.4	7.5	3,259	2,241	627
Italy	2,586	61,095,551	2,713.3	8.7	3,205	2,491	640
Liechtenstein	160	39,711	5.0	NA	-	-	-
Luxembourg	2,586	650,364	86.1	5.4	6,782	5,976	609
Malta	316	464,186	25.2	8.2	3,553	2,470	1,013
Netherlands	41,543	17,400,824	1,118.1	10.1	5,621	3,798	535
Norway	323,802	5,553,840	428.3	10.5	7,254	6,300	929
Portugal	450,295	10,242,081	369.6	9.5	2,230	1,446	589
Spain	505,370	47,163,418	1,929.8	9.1	3,001	2,246	556
Sweden	450,295	10,483,647	617.9	10.9	5,851	5,046	740

Switzerland	41,277	8,508,698	672.5	11.3	9,801	3,290	2,532
United Kingdom	243,610	67,791,400	3,344.5	10.2	4,903	4,093	625

ISN, International Society of Nephrology; sq, square; km, kilometers; GDP, Gross Domestic Product; PPP, Purchasing Power Parity; IQR, Interquartile Range

*US\$ 2021

Table 2. Burden of CKD and risk factors in the ISN Western Europe region.¹⁷⁻¹⁸

	CKD Prevalence % (95% CI)	Death attributed to CKD % (95% CI)	DALYS attributed to CKD % (95% CI)	Obesity % (95% CI)	Increased BP % (95% CI)	Smoking % (95% CI)
Overall, median [IQR]	9.54 [5.87 - 11.73]	2.44 [1.58 - 3.88]	1.47 [1.07 - 2.30]	21.9 [7.8 - 26.6]	24.8 [20.8 - 28.5]	13.1 [7.8 - 20.1]
ISN Western Europe region, median [IQR]	10.63 [9.59 - 11.57]	2.59 [2.11 - 3.42]	1.32 [1.09 - 1.59]	24.5 [23.1 - 26.9]	19.4 [18.7 - 20.6]	18.8 [17.8 - 21.2]
Andorra	9.17 (8.48 - 9.80)	2.59 (2.19 - 2.94)	1.19 (1.01 - 1.36)	28.0 (22.0 - 34.0)	18.7 (13.3 - 24.8)	20.6 (18.1 - 23.0)
Austria	11.57 (10.73 - 12.38)	4.08 (3.47 - 4.51)	1.85 (1.61 - 2.09)	21.9 (17.5 - 26.8)	21.0 (15.7 - 27.1)	25.2 (22.8 - 27.8)
Belgium	11.00 (10.28 - 11.70)	2.6 (2.2 - 2.92)	1.33 (1.15 - 1.49)	24.5 (20.7 - 28.5)	17.5 (12.9 - 23.0)	18.7 (17.0 - 20.4)
Denmark	10.63 (9.86 - 11.37)	2.05 (1.77 - 2.26)	1.23 (1.08 - 1.38)	21.3 (17.5 - 25.3)	20.6 (15.9 - 25.7)	17.5 (16.0 - 19.1)
Finland	10.22 (9.50 - 10.89)	1.26 (1.06 - 1.4)	0.77 (0.68 - 0.86)	24.9 (21.2 - 28.7)	19.4 (15.2 - 24.1)	16.8 (15.2 - 18.6)
France	10.49 (9.77 - 11.28)	2.11 (1.71 - 2.42)	1.09 (0.94 - 1.24)	23.2 (18.8 - 27.8)	22.0 (16.6 - 27.8)	21.4 (19.3 - 23.4)
Germany	12.27 (11.57 - 12.96)	3.67 (3.15 - 4.06)	1.8 (1.57 - 2.02)	25.7 (21.9 - 29.8)	19.9 (14.7 - 25.1)	21.2 (19.2 - 23.3)
Greece	14.67 (12.73 - 16.91)	3.58 (3.13 - 3.97)	2.1 (1.85 - 2.36)	27.4 (22.5 - 32.7)	19.1 (13.8 - 25.3)	31.2 (28.9 - 33.5)
Iceland	7.99 (7.44 - 8.56)	1.58 (1.32 - 1.78)	0.81 (0.7 - 0.92)	23.1 (19.1 - 27.4)	19.7 (14.1 - 25.9)	14.7 (12.9 - 16.5)
Ireland	9.59 (9.17 - 10.04)	2.16 (1.85 - 2.43)	1.08 (0.94 - 1.23)	26.9 (22.3 - 31.7)	19.7 (14.7 - 25.3)	21.6 (19.2 - 24.2)
Israel	8.45 (7.86 - 9.05)	5.77 (4.95 - 6.37)	2.45 (2.1 - 2.83)	26.7 (21.9 - 31.6)	16.6 (11.7 - 22.4)	17.8 (15.9 - 20.0)
Italy	12.17 (11.30 - 13.06)	2.57 (2.16 - 2.81)	1.39 (1.21 - 1.55)	22.9 (19.3 - 26.8)	21.2 (16.2 - 26.3)	19.4 (17.7 - 21.4)
Liechtenstein	-	-	-	-	-	-
Luxembourg	9.84 (9.16 - 10.53)	2.87 (2.43 - 3.22)	1.32 (1.14 - 1.49)	24.2 (19.4 - 29.4)	21.9 (16.3 - 27.9)	20.9 (18.4 - 23.6)
Malta	11.91 (11.03 - 12.77)	2.79 (2.39 - 3.12)	1.59 (1.38 - 1.8)	31.0 (25.1 - 37.0)	19.4 (13.8 - 25.5)	18.8 (16.6 - 20.9)
Netherlands	11.26 (10.47 - 12.05)	2.41 (2.06 - 2.69)	1.24 (1.08 - 1.4)	23.1 (19.3 - 27.0)	18.7 (14.4 - 23.7)	17.9 (16.2 - 19.5)
Norway	8.82 (8.20 - 9.41)	1.87 (1.61 - 2.03)	0.97 (0.84 - 1.08)	25.0 (20.9 - 29.2)	19.7 (14.7 - 25.0)	15.3 (13.6 - 17.2)
Portugal	11.58 (10.63 - 12.55)	3.5 (2.98 - 3.87)	1.8 (1.56 - 2.04)	23.2 (18.8 - 27.8)	24.4 (18.2 - 31.2)	17.8 (15.9 - 19.8)
Spain	10.33 (9.57 - 11.10)	3.42 (2.82 - 3.9)	1.52 (1.3 - 1.74)	27.1 (23.2 - 31.1)	19.2 (14.5 - 24.6)	21.8 (19.7 - 24.0)
Sweden	11.14 (10.30 - 11.95)	2.13 (1.8 - 2.35)	1.13 (0.98 - 1.28)	22.1 (18.3 - 26.0)	19.3 (14.7 - 24.3)	11.4 (10.7 - 12.2)

Switzerland	11.44 (10.63 - 12.20)	3.18 (2.63 - 3.62)	1.36 (1.15 - 1.54)	21.2 (17.3 - 25.3)	18.0 (13.5 - 23.2)	19.1 (17.1 - 21.3)
United Kingdom	9.07 (8.43 - 9.72)	1.25 (1.1 - 1.34)	0.82 (0.72 - 0.9)	29.5 (26.6 - 32.5)	15.2 (12.2 - 18.3)	18.7 (17.0 - 20.4)

ISN, International Society of Nephrology; CKD, chronic kidney disease; CL, confidence interval; BP, blood pressure; IQR, Interquartile Range; DALYs, disability-adjusted life years

All presented as percentages with 95% confidence intervals (95%CI).

Table 3. Kidney replacement therapy statistics in the ISN Western Europe region.¹⁹⁻²⁶

Country	Treated KF		Chronic dialysis (HD+PD)		Chronic Hemodialysis		Chronic PD		Kidney Transplants Overall		Incidence of deceased donor transplants	Incidence of living donor transplants	Incidence of pre-emptive transplants
	Incidence	Prevalence	Incidence	Prevalence	Incidence	Prevalence	Incidence	Prevalence	Incidence	Prevalence			
Andorra	-	-	-	-	-	-	-	-	-	-	-	-	-
Austria	135	1044	128.8	536.2	118.2	495.8	10.6	40.4	34	507.1	29.33	4.67	6.7
Belgium	197	1337	194.2	766.15	175.85	708.65	18.35	57.5	35.95	569.65	30.86	5.09	2.9
Denmark	108	952	97.9	434	61.8	346.8	36.1	87.2	43.45	517.7	31.72	11.72	10
Finland	94	957	91.7	361	69.2	306.8	22.5	54.2	48.73	601	40.36	8.36	4.3
France	170	1376	162.7	746.5	146.1	700.7	16.6	45.8	49.72	622.3	42.05	7.68	7.1
Germany	-	1114	-	808	-	768.1	-	38.8	23.74	-	18.08	5.66	-
Greece	269	1413	267.8	1164.4	255.1	1100.5	12.7	63.9	16.54	248.9	7.98	8.56	1.1
Iceland	108	810	91.5	241.3	69.3	196.9	22.2	44.4	30	565.8	23.33	6.67	13.9
Ireland	88	979	-	445	-	401.39	-	39.16	27.8	533	20.8	7	-
Israel	198	755	-	756	-	698.32	-	62.35	54.09	492	16.93	37.16	-
Italy	165	1276	-	800	-	701.8	-	89.32	31.52	476	26.83	4.69	-
Liechtenstein	-	-	-	-	-	-	-	-	-	-	-	-	-
Luxembourg	-	522	-	87	-	596.7	-	4.7	12	-	12	0	-
Malta	-	-	-	-	-	-	-	-	7.5	-	7.5	0	-
Netherlands	106	1034	103.4	379	83.1	322.7	20.3	56.3	53.26	695.2	26.98	26.28	16.8
Norway	100	1014.5	101.6	317.3	72.2	252.2	29.4	65.1	44.7	686.4	33.52	10.93	11.8
Portugal	260	2008	-	1302	-	871.3	-	48.1	42.06	707	38.53	3.53	-
Spain	152	1368	139.2	599.6	114.7	531.4	24.5	68.2	63.17	751.2	56.25	6.92	6.8
Sweden	113	996	103.5	401.2	68.4	316.1	35.1	85.1	43.63	594.7	32.06	11.57	9.7
Switzerland	99	968	93.7	447.8	80.8	408.5	12.9	39.3	41.61	519.2	27.59	14.02	5.1
United Kingdom	151	1293	102.9	414.8	81.5	489	21.4	69	42.77	735	32.02	10.75	12.8

ISN, International Society of Nephrology; KF, kidney failure; HD, hemodialysis; PD, peritoneal dialysis

Table 4. Annual cost of kidney replacement therapy in the ISN Western European region.²⁷⁻⁴⁵

	Hemodialysis	Peritoneal dialysis	Kidney Transplant (First year)	Kidney Transplant (later years)	HD/PD ratio
Austria	-	-	-	-	-
Belgium	60,853	29,147	76,552	22,635	2.09
Denmark	71,318	63,935	35,737	9,920	1.12
Finland	65,842	31,057	27,394	578	2.12
France	67,830	37,269	71,627	14,480	1.82
Germany	48,256	35,465	126,993	29,533	1.36
Greece	79,499	32,622	76,629	19,924	2.44
Iceland	24,419	27,206	-	-	0.90
Ireland	81,031	18,959	-	-	4.27
Israel	66,861	40,809	-	-	1.64
Italy	62,082	72,627	-	-	0.85
Liechtenstein	32,029	21,076	96,430	14,091	1.52
Luxembourg	-	-	-	-	-
Malta	-	-	-	-	-
Netherlands	-	-	-	-	-
Norway	99,419	66,655	118,595	41,254	1.49
Portugal	71,791	50,534	41,108	25,242	1.42
Spain	30,361	30,361	116,952	12,379	1.00
Sweden	26,937	59,854	64,822	10,630	0.45
Switzerland	80,520	-	94,144	18,054	-
United Kingdom	94,085	83,070	19,255	-	1.13

ISN, International Society of Nephrology; HD, hemodialysis; PD, peritoneal dialysis

Costs in \$US in 2021

Table 5. Funding structures for kidney care in the ISN Western Europe region.

	Acute Kidney Injury			CKD no dialysis					Dialysis					Transplantation				
Country	Publicly funded by govt; free at the point of delivery	Solely private through health insurance providers	Multiple systems	Publicly funded by govt; free at the point of delivery	Publicly funded by govt but with some fees at the point of delivery	Mix of public and private funding systems	Solely private through health insurance providers	Multiple systems	Publicly funded by govt; free at the point of delivery	Publicly funded by govt but with some fees at the point of delivery	Mix of public and private funding systems	Solely private through health insurance providers	Multiple systems	Publicly funded by govt; free at the point of delivery	Publicly funded by govt but with some fees at the point of delivery	Mix of public and private funding systems	Solely private through health insurance providers	Multiple systems
Andorra			X					X									X	
Austria	X			X					X				X					X
Belgium	X				X					X				X				
Denmark	X			X					X				X					X
Finland	X				X					X				X				
France	X					X					X				X			
Germany	X				X					X				X				
Greece	X				X				X				X					X
Iceland	X				X					X			X					X
Ireland	X					X					X				X			
Israel	X				X				X				X					X
Italy	X			X					X				X					X
Liechtenstein		X					X					X				X		
Luxembourg	X				X					X			X					X
Malta	X			X					X				X					X
Netherlands	X						X					X				X		
Norway	X				X					X				X				
Portugal	X				X					X				X				
Spain	X				X					X				X				
Sweden	X				X					X				X				
Switzerland	X						X		X							X		
United Kingdom	X				X					X				X				

CKD, chronic kidney disease; ISN, International Society of Nephrology

Table 6. Availability of KRT centers and nephrology workforce in the ISN Western Europe region.

	Chronic HD Centers PMP	Chronic PD Centers PMP	Transplant Centers PMP	Nephrologists PMP	Nephrology trainees PMP
Andorra	11.69	11.69		0.00	0.00
Austria	5.61	1.12	0.45	33.66	5.61
Belgium	8.44	-	0.59	21.10	4.22
Denmark	2.70	2.70	0.51	25.33	4.05
Finland	5.18	3.93	0.18	19.64	8.93
France	9.31	2.20	0.69	17.57	5.86
Germany	8.66	2.73	0.55	27.01	0.59
Greece	18.99	2.85	0.57	36.07	9.49
Iceland	13.98	2.80	2.80	30.76	0.00
Ireland	2.46	1.90	0.19	11.75	9.48
Israel	7.74	2.24	0.67	24.68	3.93
Italy	9.00	3.27	0.65	-	49.10
Liechtenstein	25.18	25.18	NA	100.73	0.00
Luxembourg	7.69	1.54	NA	12.30	6.15
Malta	4.31	2.15	2.15	-	17.23
Netherlands	3.62	3.74	0.46	15.17	2.30
Norway	4.50	4.68	0.18	27.37	9.00
Portugal	12.89	1.95	0.78	34.17	14.65
Spain	4.24	1.70	0.85	53.01	10.60
Sweden	6.68	5.25	0.38	23.85	6.68
Switzerland	10.58	7.05	0.12	41.13	5.88
United Kingdom	1.00	1.00	0.34	12.13	6.15

KRT, kidney replacement therapy; ISN, International Society of Nephrology; HD, hemodialysis; PD, peritoneal dialysis; pmp, per million population

Table 7. Availability of CKD, dialysis, kidney transplant, AKI, and CKM registries in the ISN Western Europe region.

Country	CKD	Dialysis	Transplant	AKI	CKM
Andorra					
Austria		X	X		
Belgium		X	X		
Denmark		X	X		
Finland		X	X		
France	X	X	X		
Germany		X	X		
Greece		X	X		
Iceland	X	X	X		X
Ireland		X	X		
Israel		X	X		
Italy		X	X		
Liechtenstein					
Luxembourg	X	X	X		
Malta		X	X		
Netherlands		X	X		
Norway	X	X	X		
Portugal		X	X		
Spain		X	X		
Sweden	X	X	X	X	X
Switzerland		X	X		
United Kingdom	X	X	X	X	

ISN, International Society of Nephrology; CKD, chronic kidney disease; AKI, acute kidney injury; CKM, conservative kidney management

Figure Legends

Figure 1. Countries of the ISN Western Europe region.

ISN, International Society of Nephrology

Figure 2. Availability of KRT centres in ISN Western Europe region: chronic hemodialysis (A), peritoneal dialysis (B), and kidney transplantation (C).

ISN, International Society of Nephrology

Figure 3. Workforce shortages for medical kidney care in the ISN Western Europe region.

ISN, International Society of Nephrology; HD, hemodialysis; PD, peritoneal dialysis

Appendix 1. Regional Board and GKHA Team Authors

Atefeh Amouzegar, Division of Nephrology, Department of Medicine, Firoozgar
Clinical Research Development Center, Iran University of Medical Sciences,
Tehran, Iran

Hans-Joachim Anders, Department of Medicine IV, Hospital of Ludwig Maximilian
University of Munich, Munich, Germany

Jyoti Baharani, Department of Renal Medicine, University Hospitals Birmingham,
Heartlands Hospital, Birmingham, United Kingdom

Debasish Banerjee, Renal and Transplantation Unit, St. George's University
Hospitals National Health Service Foundation Trust, Molecular and Clinical
Sciences Research Institute, St. George's University of London, London, UK

Boris Bikbov, Scientific-Tools.Org, Global Health Department, Bergamo, Italy

Edwina A. Brown, Imperial College Renal and Transplant Center, Hammersmith
Hospital, London, UK

Yeoungjee Cho, Department of Kidney and Transplant Services, Princess Alexandra
Hospital, Brisbane, Queensland, Australia; Australasian Kidney Trials Network
at the University of Queensland, Brisbane, Queensland, Australia

Kathleen Claes, Department of Nephrology, Dialysis, and Renal Transplantation,
University Hospitals Leuven, Leuven, Belgium; Department of Microbiology,
Immunology, and Transplantation, Nephrology and Renal Transplantation
Research Group, KU Leuven, Leuven, Belgium

Naomi Clyne, Department of Nephrology, Clinical Sciences Lund, Faculty of
Medicine, Skåne University Hospital and Lund University, Lund, Sweden

M. Razeen Davids, Division of Nephrology, Department of Medicine, Stellenbosch
University and Tygerberg Hospital, Cape Town, South Africa

Sara N. Davison, Division of Nephrology and Immunology, University of Alberta
Faculty of Medicine and Dentistry, Edmonton, Alberta, Canada

Hassane M. Diongole, Division of Nephrology, Department of Medicine, National
Hospital Zinder, Zinder, Niger; Faculty of Health Sciences, University of Zinder,
Zinder, Niger

Smita Divyaveer, Department of Nephrology, Postgraduate Institute of Medical
Education and Research, Chandigarh, India

Gavin Dreyer, Department of Nephrology, Barts Health NHS Trust, London, UK

Jan Dudley, Department of Paediatric Nephrology, Bristol Royal Hospital for
Children, Bristol, United Kingdom

Udeme E. Ekrikpo, Department of Internal Medicine, University of Uyo/University of
Uyo Teaching Hospital, Uyo, Nigeria

Isabelle Ethier, Division of Nephrology, Centre Hospitalier de l'Université de
Montréal, Montréal, Québec, Canada; Health Innovation and Evaluation hub,
Centre de Recherche du Centre Hospitalier de l'Université de Montréal,
Montréal, Québec, Canada

Rhys D. R. Evans, University College London Department of Renal Medicine, Royal
Free Hospital, London, UK

Stanley L. S. Fan, Department of Renal Medicine and Transplantation, Barts Health
NHS Trust, London, UK

Winston Wing-Shing Fung, Department of Medicine & Therapeutics, Prince of Wales
Hospital, The Chinese University of Hong Kong, Hong Kong SAR, China

Maurizio Gallieni, Nephrology and Dialysis Unit ASST Fatebenefratelli Sacco,
Milano, Italy; Specialty School of Nephrology, Department of Biomedical and
Clinical Sciences, University of Milano, Italy

Anukul Ghimire, Division of Nephrology, Department of Medicine, University of
Calgary, Calgary, Alberta, Canada

Ghenette Houston, Division of Nephrology and Immunology, Faculty of Medicine and
Dentistry, University of Alberta, Edmonton, Alberta, Canada

Htay Htay, Department of Renal Medicine, Singapore General Hospital, Singapore,
Singapore; Duke-NUS Medical School, Singapore, Singapore

Kwaifa Salihu Ibrahim, Nephrology Unit, Department of Medicine, Wuse District
Hospital, Abuja, Nigeria; Department of Internal Medicine, College of Health
Sciences, Nile University, Federal Capital Territory, Abuja, Nigeria

Georgina Irish, Australia and New Zealand Dialysis and Transplant (ANZDATA)
Registry, South Australia Health and Medical Research Institute, Adelaide,
South Australia, Australia; Faculty of Health and Medical Sciences, University of
Adelaide, Adelaide, South Australia, Australia; Central and Northern Adelaide
Renal and Transplantation Service, Royal Adelaide Hospital, South Australia
Health, Adelaide, South Australia, Australia

Kailash Jindal, Division of Nephrology and Immunology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

Arif Khwaja, Sheffield Kidney Institute, Northern General Hospital, Sheffield, UK

Rowena Lalji, Centre for Kidney Disease Research, University of Queensland, Brisbane, Queensland, Australia; Department of Nephrology, Queensland Children's Hospital, Brisbane, Queensland, Australia; Metro South and Integrated Nephrology and Transplant Services (MINTS), Princess Alexandra Hospital, Brisbane, Queensland, Australia

Vassilios Liakopoulos, 2nd Department of Nephrology, AHEPA University Hospital, Medical School, Aristotle University of Thessaloniki, Thessaloniki, Greece

Valerie A. Luyckx, Department of Public and Global Health, Epidemiology, Biostatistics and Prevention Institute, University of Zurich, Zurich, Switzerland; Renal Division, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts, USA; Department of Paediatrics and Child Health, University of Cape Town, Cape Town, South Africa

Manuel Macia, Department of Nephrology, Hospital Universitario Ntra Sra de Candelaria, Santa Cruz de Tenerife, Spain

Hans Peter Marti, Division of Nephrology, Haukeland University Hospital and University of Bergen, Bergen, Norway

Piergiorgio Messa, Department of Nephrology, Dialysis and Kidney Transplants, IRCCS Ca' Granda Foundation Maggiore Policlinico Hospital, Milan, Italy

Thomas F. Müller, Clinic of Nephrology, University Spital Zürich, Zurich, Switzerland

Aisha M. Nalado, Department of Medicine, Bayero University Kano, Kano, Nigeria

Brendon L. Neuen, Kidney Trials Unit, Royal North Shore Hospital, Sydney, New South Wales, Australia; The George Institute for Global Health, Sydney, New South Wales, Australia

Dorothea Nitsch, Department of Non-communicable Disease Epidemiology, London School of Hygiene & Tropical Medicine, London, England, UK

Fernando Nolasco, Department of Nephrology, Nova Medical School, Universidade Nova de Lisboa, Lisboa, Portugal

Rainer Oberbauer, Division of Nephrology and Dialysis, Department of Internal Medicine III, Medical University of Vienna, Vienna, Austria

Mohamed A Osman, Department of Family Medicine, University of Ottawa, Ottawa, Ontario, Canada

Aikaterini Papagianni, School of Medicine, Aristotle University of Thessaloniki, Thessaloniki, Greece

Anna Petrova, Department of Propaedeutics of Internal Medicine, Bohomolets National Medical University, Kyiv, Ukraine; Department of Nephrology, "Diavita Institute," Kyiv, Ukraine

Giorgina Barbara Piccoli, Nephrology, Centre Hospitalier Le Mans, Le Mans, France

Liam Plant, Department of Renal Medicine, Cork University Hospital and University College Cork, Cork, Ireland

Giuseppe Remuzzi, Mario Negri Institute for Pharmacological Research IRCCS, Milan, Italy

Parnian Riaz, Division of Nephrology, Department of Medicine, McMaster University,
Hamilton, Ontario, Canada

Joris J. Roelofs, Department of Pathology, Amsterdam University Medical Center,
Amsterdam, The Netherlands

Michael Rudnicki, Department of Internal Medicine IV – Nephrology and
Hypertension, Medical University Innsbruck, Innsbruck, Austria

Syed Saad, Division of Nephrology and Immunology, Faculty of Medicine and
Dentistry, University of Alberta, Edmonton, Alberta, Canada

Aminu Muhammad Sakajiki, Department of Medicine, Usmanu Danfodiyo University
and Usmanu Danfodiyo University Teaching Hospital, Sokoto, Nigeria

Johannes B. Scheppach, Department of Nephrology and Medical Intensive Care,
Charité – Universitätsmedizin Berlin, Berlin, Germany

Emily See, Royal Melbourne Hospital and Royal Children's Hospital, Melbourne,
Australia; Department of Critical Care, University of Melbourne, Melbourne,
Australia

Rukshana Shroff, Paediatric Renal Unit, University College London Great Ormond
Street Hospital and Institute of Child Health, London, United Kingdom

Marit D. Solbu, Section of Nephrology, University Hospital of North Norway, Tromsø,
Norway; Metabolic and Renal Research Group, UiT The Arctic University of
Norway, Tromsø, Norway

Stephen M. Sozio, Department of Medicine, Johns Hopkins School of Medicine,
Baltimore, Maryland, USA; Department of Epidemiology, Johns Hopkins
Bloomberg School of Public Health, Baltimore, Maryland, USA

Giovanni FM Strippoli, Division of Nephrology, Department of Precision and Regenerative Medicine and Ionian Area (DiMePre-J), University of Bari, Bari, Italy

Maarten W. Taal, Department of Renal Medicine, Royal Derby Hospital, Derby, UK; Centre for Kidney Research and Innovation, School of Medicine, University of Nottingham, Nottingham, UK

James Tataw Ashu, Department of Internal Medicine and Nephrology, Moutier hospital, Moutier, Canton of Berne, Switzerland; Department of Medicine, Service of Nephrology and Hypertension, Geneva University Hospitals, Canton of Geneva, Switzerland

Sophanny Tiv, Division of Nephrology and Immunology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada

Somkanya Tungsanga, Division of Nephrology and Immunology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, Alberta, Canada; Division of General Internal Medicine-Nephrology, Department of Medicine, Faculty of Medicine, Chulalongkorn University, Bangkok, Thailand

Jeroen B. van der Net, Department of Internal Medicine, Albert Schweitzer Hospital, Dordrecht, the Netherlands

Raymond C. Vanholder, Nephrology Section, Department of Internal Medicine and Pediatrics, University Hospital, Ghent, Belgium; European Kidney Health Alliance (EKHA), Brussels, Belgium

Andrea Viecelli, Department of Kidney and Transplant Services, Division of Medicine, Princess Alexandra Hospital, Woolloongabba, Queensland, Australia;

University of Queensland, Queensland, Australia; Australasian Kidney Trials
Network at the University of Queensland, Brisbane, Queensland, Australia

Katie Vinen, Department of Renal Medicine, King's College Hospital, Denmark Hill,
London, United Kingdom

Bruno Vogt, Department of Nephrology and Hypertension, Inselspital, Bern
University Hospital, Bern, Switzerland

Marina Wainstein, Faculty of Medicine, University of Queensland, Brisbane,
Queensland, Australia; West Moreton Kidney Health Service, Ipswich Hospital,
Brisbane, Queensland, Australia

Talia Weinstein, Department of Medicine, Tel Aviv University, Tel Aviv, Israel

David C. Wheeler, Department of Renal Medicine, University College London,
London

Emily K. Yeung, Department of Nephrology, Monash Health, Clayton, Victoria,
Australia

Deenaz Zaidi, Division of Nephrology and Immunology, Faculty of Medicine and
Dentistry, University of Alberta, Edmonton, Alberta, Canada

Regional Board and GKHA Team Author Disclosures

HJA reports personal fees from GSK, Novartis, Bayer, AstraZeneca, Boehringer-
Ingelheim, Roche, Aptarion, and Otsuka, outside the submitted work. DB reports
grants from AstraZeneca ESR, speaker fees from CSL Vifor and Bayer, outside the
submitted work. EAB reports grants from Kidney Research UK, personal fees from
Baxter Healthcare, Fresenius Medical Care, and Vifor, outside the submitted work.

YC reports grants and other from Baxter Healthcare, outside the submitted work. KC reports other from Alexion, other from Astra Zeneca, other from Astellas, other from Vifor Pharma, other from Fresenius Kabi, other from Sanofi, non-financial support from GSK, outside the submitted work. MRD reports personal fees (consultancy) from National Renal Care, outside the submitted work, and is the Chair of the African Renal Registry and Co-chair of the South African Renal Registry. SND reports research funding from Canadian Institutes of Health Research, Alberta Innovates, and Alberta Health services. GD reports personal fees (consulting) from Boehringer Ingelheim, outside the submitted work. IE reports grants from Fonds de Recherche du Québec - Santé, outside the submitted work. HH reports personal fees from AWAK technology and Baxter Healthcare, and non-financial support from Mologic company, outside the submitted work. VL reports personal fees from Astra Zeneca, Sanofi, GSK, Alektor, Menarini, Astellas, and grants from Baxter healthcare, outside the submitted work. VAL reports royalties from Elsevier, consulting fees from the World Health Organization, travel support from the European Renal Association and International Society of Nephrology, and leadership role in Advocacy Working Group of the International Society of Nephrology, outside the submitted work. PM reports unpaid testimony for AIFA, travel support from his organization, and leadership role in the Italian Society of Nephrology, outside the submitted work. BLN reports personal fees (advisory boards, speaker honoraria) from AstraZeneca and Boehringer and Ingelheim, personal fees (advisory boards) from Alexion, Bayer and Cambridge Healthcare Research, and personal fees (speaker honoraria) from Cornerstone Medical Education, Medscape and The Limbic, outside the submitted work, with all fees paid to The George Institute for Global Health, outside the submitted work. DN reports grants from the UK Health Foundation, NIHR, MRC, and

GSK, and leadership role in the UKKA, outside the submitted work. RO reports personal fees from Sanofi, CSL Vifor, and Hansa, and other from Alexion, outside the submitted work. AP reports personal fees from Amgen Hellas E.P.E, Astellas Pharmaceuticals A.E.B.E, Baxter Hellas E.P.E., Genesis Pharma S.A., GlaxoSmithKline A.E.B.E., and Menarini Hellas A.E., outside the submitted work. GPB reports personal fees (consultancy honoraria) and other (travel grants) from Fresenius Kabi, outside the submitted work. LP reports personal fees from AstraZeneca, A. Menarini, Boehringer-Ingelheim, Servier, MSD, Novartis, Novo Nordisk, and Vifor Pharma and is President of the Irish Nephrology Society, outside the submitted work. JJR reports personal fees from Novartis, outside the submitted work. MDS reports personal fees from AstraZeneca, personal fees from Boehringer-Ingelheim, personal fees from ViforPharma, personal fees from Baxter, outside the submitted work. MWT reports royalties from Elsevier, grants from Fresenius Medical Care, personal fees from Bayer, personal fees from Boehringer Ingelheim, outside the submitted work; and is Chair of the International Network of CKD Cohorts. ST reports Fellowship grants from the International Society of Nephrology-Salmasi Family and the Kidney Foundation of Thailand, outside the submitted work. RCV reports personal fees from R AstraZeneca, Glaxo Smith Kline, Fresenius Kabi, Novartis, Kibow, Baxter, Nipro, Fresenius Medical Care and Nextkidney, outside the submitted work. TW reports consulting fees from GSK, Astellas, Bayer, Boehringer Ingelheim, AstraZeneca, persone fees from GSK, Bayer, AstraZeneca, outside the submitted work; TW also participates on advisory boards for Bayer, Boehringer Ingelheim, and AstraZeneca, outside the submitted work. DCW reports personal fees (consutancy) from AstraZeneca, and personal fees (honoraria) from Astellas, Bayer, Boehringer Ingelheim, Eledon, Galderma, GSK, Jansen, Mundipharma, MSD,

Menorini, ProKidney, Pharmacosmos, Tricida, and Vifor, outside the submitted work.

All other authors have nothing to declare.