Is it the place or the people in the places? Exploration of why young people in deprived coastal communities of England have worse mental health than their peers inland

Emily T Murray, Avril Keating, Cara Booker, Claire Cameron, Sam Whewall, Stephen Jivraj

PII: S2666-5581(25)00072-7

DOI: https://doi.org/10.1016/j.wss.2025.100307

Reference: WSS 100307

To appear in: Wellbeing, Space and Society

Received date: 10 June 2025
Revised date: 29 September 2025
Accepted date: 1 October 2025



Please cite this article as: Emily T Murray, Avril Keating, Cara Booker, Claire Cameron, Sam Whewall, Stephen Jivraj, Is it the place or the people in the places? Exploration of why young people in deprived coastal communities of England have worse mental health than their peers inland, *Wellbeing, Space and Society* (2025), doi: https://doi.org/10.1016/j.wss.2025.100307

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2025 Published by Elsevier Ltd.

This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/)

Highlights:

- English adolescents' mental health is lower in coastal, vs inland, deprived areas.
- We used twelve waves of a nationally representative study of adolescents living in England from 2009-2020.
- We linked individual and environmental measures to where respondents lived during adolescence.
- We then followed them up to 11 years later to assess their mental health as young adults.
- Keys drivers of the coastal mental health gap were household socio-economic circumstances.
- Investments should be made in deprived coastal areas to tackle the drivers of poor mental health.

Is it the place or the people in the places? Exploration of why young people in deprived coastal communities of England have worse mental health than their peers inland.

Emily T Murray ¹, Avril Keating ², Cara Booker ³, Claire Cameron ⁴, Sam Whewall ² and Stephen Jivraj ⁵

- 1 Institute of Public Health and Wellbeing, University of Essex, Colchester, CO4 3SQ, UK
- 2 Institute of Education, University College London; London, WC1B 7HB, UK
- 3 Institute of Social and Economic Research, University of Essex; Colchester
- 4 Social Research Institute, University College London, London
- 5 Department of Epidemiology and Public Health, University College London; London

Correspondence to:

Emily T Murray, Centre for Coastal Communities, Institute of Public Health and Wellbeing, University of Essex, Colchester, CO4 3SQ, UK

Email: emily.murray@essex.ac.uk; ORCID: 0000-0001-6297-6920

Declarations of interest: none

Funder: United Kingdom Economic and Social Research Council

Word count: 4900 (references: 53, tables: 4, figures: 0, supplementary tables: 4, supplementary figures: 0)

ABSTRACT (250 words)

Previous research has shown that English adolescents who lived in the most deprived coastal neighbourhoods had worse mental health up to 11 years later than if they had lived in equivalent inland neighbourhoods. We used the same twelve waves (2009-2022) of Understanding Society, to examine whether this association was explained by the places the study members lived (31 objectively measured built, social, economic and educational indicators linked via residential lower-super output areas) or their collective individual socio-demographics when they were adolescents (aged 10-15yrs). Coastal youth (n=764) were exposed to worse average levels of sixteen environmental measures and better average levels for five environmental measures, than their peers inland (n=4,157). The concentration of area deprivation was also greater for coastal youth compared with their inland peers. When longitudinal models were fitted between environmental measures and SF-12 mental functioning scores (MCS) during adulthood (age 16+), only local crime and higher education participation were independently associated with MCS [Top 20% vs Bottom 20% (95% Confidence interval): -1.20 (-2.38, -0.03) and Middle 20% vs Worse 20%: 1.07 (0.09, 2.05)] after adjustment for socio-demographics. As well, the amplified effect of area deprivation on MCS in coastal, compared to inland, areas was reduced the most by adjustment for individual socio-demographics [interaction term coastal*Top20% deprived area: -5.1 (-8.1, -2.2) to -4.3 (-7.0, -1.6)], rather than the two environmental measures [further reduced to -3.9 (-6.7,-1.1)]. Results from this paper suggest policies to improve young adult's mental health in England should target the socioeconomic circumstances of households in the most deprived coastal areas.

INTRODUCTION

As of 2023, one in five children and young people in England had a probable mental health disorder. This includes 20.3% of 8–16-year-olds, 23.3% of 17–19-year-olds and 21.7% of 20–25-year-olds (1). These percentages have doubled since 2010, with adolescents and young people showing the sharpest increases across multiple national data sets (2). Correspondingly, National Health Service mental health referrals for children and young people have increased by 50% between 2020 and 2023 (3).

However, the occurrence of poor mental health is not distributed equally across England. There is no publicly available small area-level data on the prevalence of mental health in only young people. There are data on the prevalence (4) of depression and mental health service demand (5) for all English adults 18 years and older, with both data sets clearly showing a higher prevalence in urban centres and specific coastal areas. The United Kingdom's (UK's) Chief Medical Officer (CMO) report from 2021 also highlighted that hospital admissions for self-harm were 35% higher for 10- to 24-year-olds living in coastal, compared to inland, Lower Super Output Areas (LSOAs) (6). When they used GP-level Quality Outcomes Framework (QOF) data from 2014/15 to 2018/19, the coastal excess youth mental health gap was only seen in LSOAs categorised as 7 and 9 of area deprivation (10 being the most deprived). Recent analysis using longitudinal population-level survey data, and self-reported levels of mental functioning and psychological distress, expanded on this finding by showing that the coastal mental health gap only occurred for young adults who had lived in the top 20% most deprived areas of England, not all coastal areas (7). Given that some of the highest rates of suicide in England are in coastal communities, and that more than two-thirds of mental illness begins by age 25 years (8), it is vital we identify the cause of these elevated levels of poor mental health in coastal youth.

Why young people in the most deprived coastal communities would have worse mental health than their peers in equally deprived inland areas has not been investigated previously. One explanation is that even within the most deprived areas, the environmental factors that cause or mitigate mental health are different in coastal compared to inland areas. The way that area deprivation is constructed is to create one index from multiple indicators that represent material and social disadvantage of residents of small geographical areas. The CMO report uses The English Indices of Deprivation 2019, which uses 39 separate indicators across seven distinct domains (9). One domain is employment deprivation, which is based on the proportion of residents claiming benefits (e.g., jobseeker's allowance) in August (10). Many coastal communities share a reliance on a seasonal economy, so a data collection at peak tourist season may not accurately reflect coastal residents' year-round employment conditions. Similar arguments could be made for other indicators. As well, there are other indicators that have been shown to be related to mental health and potentially more common in coastal youth but are not included in any of the indexes. For example, social isolation is a growing public health concern that is known to influence mental health (11) and coastal youth are more prone to social isolation from residing in areas with a high retiree populations (6, 12), geographic isolation from other communities (13) and/or public social spaces perceived as only for tourists (14).

Another explanation for why young people in coastal deprived areas would have worse mental health than their peer's inland is the high concentration of poverty in these places. Coastal towns were once thriving centres of commerce, but many have experienced stark economic declines in recent decades (15, 16). These economic declines have led to a higher proportion of the population being in poor health, which reduces incentives for business owners to invest in the area, which leads skilled workers to seek

employment elsewhere; creating a negative feedback loop of economic and health decline (12). Thus, it is not something about the environments of deprived coastal places that is directly affecting the health of residents (i.e. contextual effects), but more that there are higher concentrations of people in these places who are struggling socioeconomically, and that these individual circumstances are collectively driving poor health outcomes (i.e. compositional effect) (17). There are also indications that levels of deprivation in coastal communities are more severe in coastal than inland areas, with coastal LSOAs making up most of the top 20 rankings of deprivation in the UK (18); but this has not been tested previously.

Therefore, the aim of this study is to examine environmental characteristics – here economic, social, educational or built – or individual socio-economic circumstances during adolescence explain the coastal youth mental health gap in the most deprived small areas of England. In order to do so, we test whether: i) average levels of individual and environmental characteristics differ between coastal and inland areas (i.e., exposure) ii) are these differences seen within sub-categories of area deprivation (i.e., exposure)? iii) are environmental characteristics associated with mental health in young adulthood (i.e., outcome)? and iv) determine how much of the difference in mental health between deprived coastal youth and deprived inland youth is explained by examined individual and environmental characteristics.

METHODS

Data source:

The UK Household Longitudinal Study (UKHLS), also known as *Understanding Society*, is a comprehensive and nationally representative study that tracks the lives of individuals and households across the United Kingdom. The study was launched in 2009-2010 with

an initial sample of approximately 40,000 households and has followed up all household members fourteen times, the latest wave in 2021-23 (19).

For this analysis, we focused on data from the youth self-completion questionnaires (ages 10-15) and tracked participants over time to capture their responses to the adult self-completion questionnaires (age 16+). Data was obtained through the UK Data Service https://ukdataservice.ac.uk/), the main survey from database SN6614 (20) and geographic codes SN7248 (21). The baseline for each individual was the wave when they completed the youth questionnaire at age 15 years. In cases where a participant did not complete the questionnaire at age 15 years, we used the wave closest to that age as their baseline.

Response rates among eligible youth varied by wave, with the highest being 82% in wave seven and the lowest 58% in wave 11 (22).

Measures

Exposure: Coastal community status during adolescence:

Coastal community status was assigned based on the lower-super output area (LSOA) identifier for each youth respondent at their baseline adolescent wave. Each respondent's baseline wave was when they had completed a youth questionnaire at the age of 15 years. If the respondent did not complete a questionnaire at the age of 15 years, the questionnaire where the respondent was closest in age to 15 years was used. To maintain temporal order, coastal residence past age 15 was not included in the exposure variable.

Each respondent's usual residence was recorded at every wave, and staff from the Institute for Social and Economic Research (ISER) provided the LSOA identifier for the 2011 Census for all waves (21). In 2011, England contained 32,844 LSOAs, each representing between 400 and 1,200 households, with a typical population ranging from

1,000 to 3,000 individuals (23). These LSOA identifiers were then used to link each youth respondent's UKHLS data (20) to coastal community status, as defined in the Chief Medical Officer of England's 2021 report. Briefly, "coastal" LSOAs were those that included or were within 500 meters of built-up areas near the "Mean High Water Mark" (excluding tidal rivers). All other LSOAs in England were classified as "inland." A more detailed definition can be found in the report (6).

Outcome: Mental health during young adulthood:

For this analysis, we used the Mental Component Summary (MCS) score from the 12-item Short-Form Survey (SF-12) as a proxy for mental health. In UKHLS, a measure of psychological distress is also calculated through the General Health Questionnaire (GHQ-12) (24), but MCS scores showed a stronger association with coastal community status in previous analysis (7). In the SF-12, six mental health-related questions were asked about mental well-being in the last four weeks. Answers to these items were converted to a single score by the Ware et al (2002) method (25) calibrated against population norms, by the UKHLS research team. The MCS scores range from 0 (poorest mental health) to 100 (best mental health). Both outcomes were assessed for each respondent at all study waves completed when they were aged 16+ years. The number of possible study waves completed varied by respondent, with a maximum of 12 waves.

Covariates:

Based on prior research, we included a range of covariates that could be possible alternative explanations for why associations would be seen between coastal community residence and mental health.

Individual level covariates included age at health measurement and six covariates measured in adolescence: gender (derived from self-reported male or female at all surveys completed), ethnicity (eighteen categories derived from multiple sources such as self-reported as an adult, self-reported as a youth, reported by a household member, and ethnic group of biological parents then collapsed into white or non-white), household income (gross monthly income imputed by the ISER team, then adjusted for household size and composition through the OECD-modified equivalence scale (26), consumer price index inflation (27) at that wave and logged) and household tenure (collapsed into three categories of homeowner, social renter, or private renter/other). The adolescent measurement corresponds to when coastal community residence was measured for each respondent, ranging from age 10 to 15 years.

Environmental measures were initially considered that could proxy one of the 15 potential pathways Galster theorizes neighbourhoods may affect health of residents (28). Priority was given to measures that had previously been seen in the literature to be associated with mental health (29-33) and to be a particular issue in English coastal communities (6, 12, 13, 16). These mechanisms, or pathways, are outlined in Table 1, with the measure identified (if any), a measure of the description, data source and timing of data collection. If multiple time periods were available, the data collection closest to the 2011 census was chosen for consistency.

Effect modifier: Area deprivation was previously identified as an effect modifier of the relationship between coastal community status and mental health (7). We used the same measure of area deprivation, the 2011 Townsend Index (z-score summary derived from four census variables: unemployment, non-car ownership, non-home ownership, and overcrowding), at the LSOA-level, when respondents were adolescents. In this analysis,

we used the Index that has been divided into quintiles based on the distribution of LSOAs in England (34).

Statistical analyses:

Initially, comparisons were made across all covariates between adolescent coastal community and inland status using analysis of variance for continuous variables and the chi-square statistic for categorical variables. Additionally, the same comparisons were made within each quintile of area deprivation.

All analysis was conducted on longitudinal panel data where each study member had to have a fixed coastal status, environmental data and all covariates at age 15 years (or closest age available) with at least one mental health outcome measure, and corresponding age at outcome, allowed to vary over the eleven follow-up waves. All included study members were matched to an LSOA and all environmental variables. Any missing data remaining for covariates were handled through complete case analysis. To identify the design of the data set, the syyset command in STATA was specified with household number designated as the sampling unit. To account for attrition in the sample over time, we applied the appropriate sample weights from the wave 12 longitudinal weights (35), along with the stratum identifier variable. The weight used, m_indscus_lw, was created by staff at the ISER to match the specific sample: 'm' indicates that it refers to the last wave of the analysis, 'ind' identifies individual respondents, 'sc' denotes the self-completion aspect of the questionnaire, 'us' refers to the GPS sample, and 'lw' signifies the longitudinal weight applied to the data. The STATA xtset command was then used to specify data to be panel data with more than one wave of data per person.

The main analysis included a series of regression models to examine which environmental variables explain why coastal youth in the most deprived coastal communities had worse mental health on average than their peers in equally deprived communities. All models were fitted using Random-effects linear regression models by GLS with SF-12 MCS scores at the individual and study wave [STATA xi:streg]. To account for clustering of similar individuals within LSOAs, a VCE cluster option was used with LSOA as the cluster variable.

The first series of models aimed to assess which adolescent environmental variables were independently associated with mental health in young adulthood. Initially, each adolescent environmental measure was fitted separately to assess associations with MCS scores after adjustment for age at MCS score assessment. Following, each separate model was additionally adjusted for adolescent individual covariates. Lastly, independent environmental associations were assessed by fitting all adolescent environmental measures jointly with each other and with all individual covariates. The latter model was required as multiple environmental variables were moderately correlated (see Supplementary Table 2).

The second series of models aimed to assess whether environmental measures explained the deprived coastal youth mental health gap. First, MCS scores were regressed on coastal community status, Townsend Index quartiles, a coastal by Townsend interaction term and age at MCS assessment. Second, the individual covariates were added to the model to assess whether associations could be explained by the differences in individual attributes of residents that live in coastal, compared to inland, communities. Lastly, environmental measures identified as having an independent association with MCS scores in previous analysis were singularly added to the age- and individual covariate-adjusted models.

RESULTS

Of the 14,746 youths who self-completed a questionnaire at least once, a total of 4,921 youth (18,324 observations, mean=3.7, range 1-11) lived in England at the age of geographic linkage to determine coastal residence, completed at least one adult (age 16+) questionnaire during the follow-up period, and had data on both health outcomes and covariates. In a previous publication (7), we showed that included participants, compared to excluded, tended to have completed questionnaires at later waves, been older at MCS assessment, more female, of White ethnicity, and lived in households where the home was owned. In Supplementary Table 2, we additionally show that excluded study members lived in LSOAs with slightly worse economic environments, more crime deprivation, more urban, slightly less travel times to most destinations examined and slightly higher levels of air pollution (No2 and PM10).

Do young people who live in coastal, compared to inland, areas experience different environmental conditions during adolescence?

Table 1 shows the environmental characteristics of respondents by the coastal or inland categorisation. Across the economic, social, educational and built environment domains, coastal adolescent respondents lived in LSOAs with less favourable average levels for sixteen examined factors, compared to their inland peers. This included lower levels of homeownership, NVQ level 3+ qualifications, residents aged 10-19 years, KS4 test scores, participation in higher education, travel time to gambling and tobacco shops, green space within 900m and skilled occupations; as well as higher levels of economic inactivity, urbanicity, mean annual sulphur dioxide and travel times to GPs, further education, large employment centres, job centres, and leisure. For seven environmental variables, coastal respondents lived in LSOAs with more favourable average levels than respondents in inland LSOAs with regards deprivation, crime deprivation, proportion households

overcrowded, walking distances to food stores and levels of air pollution (annual nitrogen dioxide and particulate matter 10).

Are environmental conditions different for coastal and inland adolescents even if they live in similarly deprived areas?

Even when the sample was split further by area deprivation categories (see Table 2), there were consistent differences in environmental variables between coastal and inland LSOAs. For the UKHLS young adults who lived within the most deprived quartile of English LSOAs during adolescence, those in coastal areas had less favourable levels of most social, economic, and educational environmental measures than inland areas in the same quintile categorisation of deprivation. This included a lower proportion of adults obtaining Level 3 qualifications, more adults in lower skilled occupations, higher average crime deprivation, higher unemployment, lower average GCSE attainment scores and lower prevalence of progress to higher education. The largest differences were in the educational measures with, for example, 21% of 18-19-year-olds in the most deprived coastal MSOAs participated in higher education, while 35.8% of their inland peers in the most deprived category participated. For built environment measures, within deprivation-category differences between coastal and inland LSOAs were significant, but small (average ~1 minute difference). Similar patterns were seen when all English LSOAs were examined, not just those where UKHLS respondents resided (See supplementary Table 2). This indicates there is no geographic selection bias in the sampling strategy of the UKHLS by whether an LSOA is coastal or not.

Are adolescent environmental conditions associated with young adult mental health up to eleven year later?

Table 4 shows adjusted associations of adolescent environmental variables with SF-12 mental functioning component (MCS) scores. Of the 36 environmental measures analysed, only ten were associated with MCS scores after adjustment for age at the time of MCS score measurement: unemployment, overcrowding, no car, economic inactivity, proportion 10–19-year-olds, POLAR quintile 3, travel time to large employment centres and job centres and levels of NO2 and PM10 (table 4, model 1). Of these, only three measures (no car, age 10-19 years and POLAR quintile 3) remained after adjustment for individual socio-demographic variables (table 4, model 2). However, an additional four measures (crime quintile 5, distance to further education, and travel time to pubs/ leisure) showed significant associations at the 0.05 level after adjustment for socio-demographics (model 2). When all significant measures from model 2 were fitted jointly (model 3), only the highest crime quartile [-1.20 (95% CI: -2.38, -0.03) and the middle quartile of participation in higher education [1.07 (0.09, 2.05)] showed independent associations with MCS scores.

Are adolescent environmental conditions, or individual socio-demographics, explanations for the coastal mental health gap?

Table 5 shows associations between coastal residence, area deprivation (measured by Townsend index) and coastal*Townsend quintiles before and after adjustment for the individual socio-demographic and environmental variables. In age-adjusted models, respondents who had lived in one of the most deprived coastal communities in adolescence, their mean MCS scores in young adulthood were -5.1 points lower (95% CI: -8.1, -2.2), compared to the least deprived inland communities (table 4, model 1). The separate addition of socio-demographic covariates and higher education progression reduced this amplification effect to -4.3 (-7.0, -1.4) and -4.7 (-7.7, -1.7), respectively, while adjustment for local crime levels (and an interaction between coastal and crime)

increased the association to -6.1 (-9.7, -2.4). Inclusion of both environmental variables improved model fit slightly (R² 0.0182 to 0.0198 and 0.0196). The final model inclusive of all individual and environmental variables, showed that adolescents residing in the top 20% most deprived coastal communities had MCS scores on average -3.9 points (95% CI: -6.7, -1.1) lower than peers in the least 20% deprived inland communities; almost equivalent to the effect of bereavement in the year immediately following a significant loss (36)

DISCUSSION

Using a nationally representative sample of adolescents who lived in England from 2009 to 2020, and up to eleven years of follow-up, we show for the first time that adolescents living in coastal neighbourhoods are being exposed to a whole host of adverse environmental exposures across multiple domains. This remained the case even when comparisons were made between coastal and inland adolescents living in the most deprived fifth of neighbourhoods. We were able to follow these young people into early adulthood, showing that their mental functioning was related to the level of crime and higher education participation in their adolescent neighbourhoods. However, being able to also examine the demographic and socioeconomic circumstances of their adolescent households showed us that it was these individual factors, particularly household income and private renting, rather than the environmental measures, that explained more of the difference (16% versus 7.8%) in mental functioning between deprived coastal and inland youth: our main focus of enquiry. Further enquiry is needed to identify reasons for the remaining mental health gap.

The finding that youth respondents who lived in coastal neighbourhoods were exposed to more adverse levels of a host of environmental variables, than their inland peers, is consistent with numerous reports on English coastal communities (6, 12, 13, 16). There is a common story, which our data supports, that many coastal communities share characteristics of struggling labour markets, lower skilled populations, lower educational attainment and lower homeownership (15). However, in terms of distances to local healthharming and health-promoting infrastructure, the built environments where coastal adolescents lived ~10 years ago were not hugely different to where their peers lived inland. As well, there was a positive story that coastal youth were exposed to appreciably lower levels of outdoor air pollution, namely No2 and PM10; which has been linked to increased risk of depressive symptoms and incidence of suicide (37, 38). In addition to the above reports, we concernedly show that even within the top 20% most deprived neighbourhoods in England, coastal youth were living in areas with worse economic, social and educational environments than their equivalent peers inland, showing that deprived coastal areas should be prioritised for national and local government regeneration initiatives.

Our third finding that local area crime and participation in higher education (POLAR) were independently associated with mental health is half supported by previous studies. There is a large literature showing robust evidence that higher levels of crime are related to higher levels of depression and psychological distress, even after adjustment for individual socio-demographics (39). This is seen to occur through direct impact on mental health of being a victim or witness of a crime, or indirectly though feeling that particular places are unsafe creating a biological stress response and/or change behaviour(s) that effect health (e.g., less physical activity or social engagement in unsafe areas) (40). In our study, the main effect of local area crime was robust to adjustment for coastal residence,

area deprivation and all individual socio-demographics. There was also an indication of an amplification of the relationship between local crime rates and mental functioning for coastal, compared to inland, residents; but this was explained by socio-demographics of youth participants. This is likely due to negative confounding from some households with high income and private renting living in LSOAs with high levels of crime, which are both strongly related to youth mental health.

For the POLAR measure, as far as we are aware this is the first study to show that young people in the middle quintile of the POLAR measure had slightly better mental functioning scores than those in areas where the highest proportion of young people participate in higher education. The hypothesized mechanism for why neighbourhood higher education progression was linked to individual mental health was one of social contagion. That seeing other young people in your neighbourhood progressing to higher education would build confidence in your own ability to progress and do well academically (41, 42). However, as this association did not increase with additional levels of participation, and most of this association was reduced by adjustment for individual socio-demographics, it is likely that this finding is explained by the individual attributes of study youth who resided in these neighbourhoods.

Our results showing that only two built environmental measures - distance to pubs and leisure centres — were related to mental functioning in young adulthood is somewhat consistent with a similar aged New Zealand study (31). They showed in a cross-sectional design that young people aged 10-24 years who lived in 'health-constraining' environments had higher odds of any diagnosed mental health condition. Why we did not see similar findings across our other 'health-constraining' environmental measures could be due to reverse temporality issues in the NZ study. We used twelve waves of longitudinal data in our analysis, so can be sure of temporal order of relationships. Our

associations were also entirely explained by adjustment for individual sociodemographics, suggesting that at least in England, built environment associations with
mental health are due to clustering of similar kinds of people residing within the same
neighbourhoods, rather than infrastructure. Surprisingly, we did not see an association
between availability of green space and youth mental health, which has been seen in some
previous studies (43-45). However, the overall literature is mixed with one systematic
review suggesting positive findings were due to selection bias and residual confounding
(46). The null finding could be due to measurement error. We assessed green space
through the average total green space area available to members of each LSOA in a range
of a 900-meter buffer, which does not take into consideration accessibility, quality or
exposure of green space of individuals (47). It may also be that green space closer to the
time of mental health diagnosis is more important, compared to our study assessing
adolescent exposure up to eleven years previously.

It was therefore not too surprising that the coastal amplification of the area deprivation and mental functioning relationship was hardly explained by the inclusion of local higher education participation or local crime rates. Model fit was slightly improved by inclusion of the two environmental measures. However, our results indicate that a large part of the explanation is the economic and social challenges that people face who live in these communities. Household incomes and private renting are key factors. Many current deprived coastal towns have experienced stark economic declines in recent decades due to the decline of seaside resorts, fishing, and shipbuilding industries (48). This has created a negative feedback loop of highly skilled and healthy individuals leaving these areas for better employment opportunities (12, 14) leaving behind the sicker and more socio-demographically challenged (49). There is however no research on how much internal migration contributes to the relationship between coastal residence and health

outcomes. It is also worth noting that this pattern where individual measures explain more of a health inequality than area-level measures are common in the area-effects literature (50). It does not mean that where people live is not important for their health. The collective interactions, behaviours and decision-making of people who live in neighbourhoods helps to create the 'context' for their neighbours and future residents (51). It should also be pointed out that even after we adjusted for all potentially explanatory factors in our study, there was still a sizable difference in mental functioning between deprived coastal and inland young adults. Further work is needed to identify other potential explanations.

One such reason could be due to one of the limitations of our study, in that we only assessed environmental measures at one time point, during adolescence. Previous research has shown that the residential environment has a stronger association with health outcomes at the time of measurement (52). However, since two-thirds of mental illness begin before the age of 25 years (8), we thought it important to examine environmental exposures around the likely time of mental illness development. Research has also shown that people with mental health problems are more likely to move to deprived neighbourhoods (53). Therefore, by focusing on the relationship of adolescent environments with young adult mental health, we reduce issues from reverse temporality and health selection bias. The reason that none of the environmental measures substantially explained differences in mental functioning between deprived coastal and inland youth could be that we did not include the right environmental measure, but it could also be due to data availability issues of most environmental measures not being measured at exactly the time of adolescent residence or exposures occurred elsewhere, such as school environment (i.e., measurement error). As in any longitudinal study, attrition occurred over follow-up, particularly the transition from the youth to adult

surveys (7). However, attrition bias is most likely to create underestimates of neighbourhood effects on health over time, given that cohort members residing in the most deprived neighbourhoods, with the worse mental functioning, would be the most likely to leave this study (54).

The major strengths of our study were the ability draw upon a nationally representative, longitudinal, sample of youth that were linked to granular administrative data. This allowed us to be confident that results are generalisable to all English youth during the study period, that we have a comprehensive picture of the environments that these youth resided during their adolescence and that results are unlikely to be the result of reverse causality. The latter point buoyed by previous results showing that only 2% of the sample moved between coastal and inland communities between adolescence and young adulthood (7). Couple this with the comprehensive socio-economic data collected on UKHLS youth participants, and the rest of their household, we are confident results are robust.

In conclusion, findings from this study suggest that strategies to improve mental health of English youth needs to pay particular attention to the socio-demographics of young people, and their families, in deprived coastal communities. In particular, improvements in household income and private renting should be targeted. Suicide rates and deaths of despair, which include suicide but also those attributable to alcohol or drugs, are significantly higher in coastal compared to inland local authorities (55). Rates of deaths of despair are highest at mid-life (55), with the suicide rate in the UK the highest it has been in over 25 years (56). But given the increasing prevalence of youth mental disorders in England (1), and that more than two-thirds of mental illness begins by age 25 years (8), resources should be targeted in today's youth to prevent a tsunami of future mental ill health and suicides.

Funding

The research leading to these results has received funding from two Economic and Social Research Council (ESRC) grants, a part of the United Kingdom Research and Innovation (UKRI). The first grant funding the Understanding Society Fellowship Programme (ES/S007253/1) (PI: Michaela Benzeval) and the second grant funding research project 'Coastal youth: Exploring the impact of coastal towns on young people's life chances' (ES/X001202/1)(PI: Avril Keating). The views stated in this work are of the authors only.

Funder

United Kingdom Economic and Social Research Council.

REFERENCES

- 1. England N. Mental Health of Children and Young People in England, 2023 wave 4 follow up to the 2017 survey. 2023.
- 2. Fonagy P. Health Foundation. 2025. [cited 2025]. Available from: https://www.health.org.uk/features-and-opinion/blogs/understanding-the-crisis-in-young-people-s-mental-health.
- 3. Comissioner Cs. Children's mental health services 2022-23. London; 2024.
- 4. Tsimpida D, Tsakiridi A, Daras K, Corcoran R, Gabbay M. Unravelling the dynamics of mental health inequalities in England: A 12-year nationwide longitudinal spatial analysis of recorded depression prevalence. SSM Popul Health. 2024;26:101669.
- 5. Daras K, Barr B. Small Area Mental Health Index (SAMHI) < VERSION 5.00> [Open Dataset], Place-based Logitudinal Data Resource. University of Liverpool2020.
- 6. Officer CM. Chief Medical Officer's annual report 2021: health in coastal communities. 2021 21/07/2021.
- 7. Murray ET, Keating A, Cameron C, Benchekroun R, Whewall S, Booker C, et al. Residence in coastal communities in adolescence and health in young adulthood: An 11-year follow-up of English UKHLS youth questionnaire respondents. Health Place. 2024;87:103239.
- 8. Solmi M, Radua J, Olivola M, Croce E, Soardo L, Salazar de Pablo G, et al. Age at onset of mental disorders worldwide: large-scale meta-analysis of 192 epidemiological studies. Mol Psychiatry. 2022;27(1):281–95.

- 9. Ministry of Housing CLGt. English indices of deprivation 2019: Statistical release main findings. 2019. p. 31.
- 10. Ministry of Housing CLGt. English indices of deprivation 2019: technical report. 2019. p. 117.
- 11. Brandt L, Liu S, Heim C, Heinz A. The effects of social isolation stress and discrimination on mental health. Transl Psychiatry. 2022;12(1):398.
- 12. Emmins N, Leckle C, Munro R, Pragnell M. Communities on the edge levelling up England's coastal communities. Chelmsford, Essex: Pragmatix Advisory; 2023.
- 13. Communities HoLSCoRSTa. The future of seaside towns. 2019 04/04/2019. Contract No.: HL Paper 320.
- 14. Wenham A. "Wish you were here"? Geographies of exclusion: young people, coastal towns and marginality. Journal of Youth Studies. 2019;23(1):44–60.
- 15. Committee CaLG. Coastal Towns Second Report. London: The Stationary Office Limited; 2007 7/03/2007.
- 16. de Graaf K, McKenzie K, Asthana S, Agarwal S, Smith R. On the Waterfront. 2025 03/2025. Report No.: 978-0-9934156-7-8.
- 17. Leyland AH, Groenewegen PP. Multilevel Modelling for Public Health and Health Services Research: Health in Context. Cham (CH)2020.
- 18. English indices of deprivation 2019. In: Ministry of Housing CLGt, editor.: ONS; 2019.
- 19. Lynn P. Sample design for Understanding Society. Understanding Society Working Paper Series2009.
- 20. , University of Essex IfSaER. Understanding Society: Waves 1-12, 2009-2021 and Harmonised BHPS: Waves 1-18, 1991-2009. In: Society U, editor.
- 21. , University of Essex IfSaER. Understanding Society: Waves 1-12, 2009-2021: Special Licence Access, Census 2011 Lower Layer Super Output Areas. In: Study UKHL, editor.: UK Data Service; 2022.
- 22. Society U. Response rates 2025 [Available from:
- https://www.understandingsociety.ac.uk/documentation/mainstage/user-guides/main-survey-user-guide/response-rates/.
- 23. ONS. 2011 Census: Population and Household Estimates for Small Areas in England and Wales, March 2011. 2012.
- 24. Griffith GJ, Jones K. Understanding the population structure of the GHQ-12: Methodological considerations in dimensionally complex measurement outcomes. Soc Sci Med. 2019;243:112638.
- 25. Ware JE, KOSINSKI M, TURNER-BOWKER M, Gandel B. How to Score Version 2 of the SF-12 Health Survey. With a Supplement Documenting Version 12002.
- 26. Förster MF, D'Ercole MM. 2 The OECD Approach to Measuring Income Distribution and Poverty In: Besharov DJ, Couch KA, editors. Book cover for Counting the Poor: New Thinking About European Poverty Measures and Lessons for the United States Counting the Poor: New Thinking About European Poverty Measures and Lessons for the United States: Oxford University Press; 2012. p. 27–58.
- 27. team CPI. CPI INDEX 00: ALL ITEMS 2015=100. In: ONS, editor. 2025.
- 28. Galster GC. The Mechanism(s) of Neighbourhood Effects: Theory, Evidence, and Policy Implications. In: van Ham M, Manley D, Bailey N, Simpson L, Maclennan D, editors. Neighbourhood Effects Research: New Perspectives: Springer; 2011. p. 23–56.
- 29. Putra I, McInerney AM, Robinson E, Deschenes SS. Neighbourhood characteristics and socioeconomic inequalities in child mental health: Cross-sectional and longitudinal findings from the Growing Up in Ireland study. Health Place. 2024;86:103180.
- 30. Laporte D, Chilman N, Morgan C, Schofield P, Wykes T, Das-Munshi J. The association between area-level factors and mortality in severe mental illnesses: A systematic review. Schizophr Res. 2024;264:95–104.

- 31. Hobbs M, Bowden N, Marek L, Wiki J, Kokaua J, Theodore R, et al. The environment a young person grows up in is associated with their
- mental health: A nationwide geospatial study using the integrated data infrastructure, New Zealand. Social Science and Medicine. 2023;326(115893).
- 32. Deng B, Boden J, Ye N, Morgenroth J, Campbell M, Eggleton P, et al. Life in green: Associations between greenspace availability and mental health over the lifecourse A 40-year prospective birth cohort study. Environ Int. 2025;195:109223.
- 33. Visser K, Bolt G, Finkenauer C, Jonker M, Weinberg D, Stevens G. Neighbourhood deprivation effects on young people's mental health and well-being: A systematic review of the literature. Soc Sci Med. 2021;270:113542.
- 34. 2011 UK Townsend Deprivation Scores. In: Agency OfNSNRoSNISaR, editor.: UK Data Service; 2011.
- 35. Society U. Selecting the correct weight for your analysis 2025 [Available from: https://www.understandingsociety.ac.uk/documentation/mainstage/user-guides/main-survey-user-guide/selecting-the-correct-weight-for-your-analysis/.
- 36. Pennington B, Hernandez Alava M, Strong M. How Does Bereavement Affect the Health-Related Quality of Life of Household Members Who Do and Do Not Provide Unpaid Care? Difference-in-Differences Analyses Using the UK Household Longitudinal Survey. Pharmacoeconomics. 2025;43(4):389–402.
- 37. Hobbs M, Deng B, Woodward L, Marek L, McLeod G, Sturman A, et al. Childhood air pollution exposure is related to cognitive, educational and mental health outcomes in childhood and adolescence: A longitudinal birth cohort study. Environ Res. 2025;274:121148.
- 38. King JD, Zhang S, Cohen A. Air pollution and mental health: associations, mechanisms and methods. Curr Opin Psychiatry. 2022;35(3):192–9.
- 39. Baranyi G, Di Marco MH, Russ TC, Dibben C, Pearce J. The impact of neighbourhood crime on mental health: A systematic review and meta-analysis. Soc Sci Med. 2021;282:114106.
- 40. Lorenc T, Clayton S, Neary D, Whitehead M, Petticrew M, Thomson H, et al. Crime, fear of crime, environment, and mental health and wellbeing: mapping review of theories and causal pathways. Health Place. 2012;18(4):757–65.
- 41. Hitlin S, Johnson MK. Reconceptualizing Agency within the Life Course: The Power of Looking Ahead. AJS. 2015;120(5):1429–72.
- 42. Keating A, Melis G. Youth Attitudes Towards Their Future: the Role of Resources, Agency and Individualism in the UK. Journal of Applied Youth Studies. 2021;5:1–18.
- 43. Madzia J, Ryan P, Yolton K, Percy Z, Newman N, LeMasters G, et al. Residential Greenspace Association with Childhood Behavioral Outcomes. J Pediatr. 2019;207:233–40.
- 44. Mavoa S, Davern M, Breed M, Hahs A. Higher levels of greenness and biodiversity associate with greater subjective wellbeing in adults living in Melbourne, Australia. Health Place. 2019;57:321–9.
- 45. Van Aart CJC, Michels N, Sioen I, De Decker A, Bijnens EM, Janssen BG, et al. Residential landscape as a predictor of psychosocial stress in the life course from childhood to adolescence. Environ Int. 2018;120:456–63.
- 46. Fleckney P, Bentley R. The urban public realm and adolescent mental health and wellbeing: A systematic review. Soc Sci Med. 2021;284:114242.
- 47. Zhang Y, Wu T, Yu H, Fu J, Xu J, Liu L, et al. Green spaces exposure and the risk of common psychiatric disorders: A meta-analysis. SSM Popul Health. 2024;25:101630.
- 48. Communities AC. Coastal Inquiry: Update report 2024. Local Government Association Coastal Special Interest Group; 2024.
- 49. Fiorentino S. Coastal towns as 'left-behind places': economy, environment and planning. Cambridge Journal of Regions, Economy and Society. 2024;17(1):103–16.

- 50. Oakes JM, Andrade KE, Biyoow IM, Cowan LT. Twenty Years of Neighborhood Effect Research: An Assessment. Curr Epidemiol Rep. 2015;2(1):80–7.
- 51. Cummins S, Curtis S, Diez-Roux AV, Macintyre S. Understanding and representing 'place' in health research: a relational approach. Soc Sci Med. 2007;65(9):1825–38.
- 52. Jivraj S, Norman P, Nicholas O, Murray ET. Are there sensitive neighbourhood effect periods during the life course on midlife health and wellbeing? Health Place. 2019;57:147–56.
- 53. Tunstall H, Mitchell R, Pearce J, Shortt N. The general and mental health of movers to more- and less-disadvantaged socio-economic and physical environments within the UK. Soc Sci Med. 2014;118:97–107.
- 54. Cabrera-Alvarez P, Lynn P. Trends in panel attrition in Understanding Society: Waves 1 to 13. Understanding Society Working Paper Series2023.
- 55. Camacho C, Webb RT, Bower P, Munford L. Risk factors for deaths of despair in England: An ecological study of local authority mortality data. Soc Sci Med. 2024;342:116560.
- 56. Group PHM. Suicides in England and Wales: 2023 registrations. ONS; 2025.

Table 1. Descri	Table 1. Description of potential environmental mechanisms with the proxy							
measurement i	measurement identified (if one), a description of the measure (including							
source and tim	source and time period) and relevance to coastal communities and mental							
health.								
		Measure(s)	Description of	Source	Coastal communit	Linke d to		
Domain	Mechanism	identified	measure	(collection	y issue?	mental		
		luchtificu	measure	period)		health		
						?		
Social- Interactive Mechanisms	Social Contagion	Educationa 1 attainment and progressio n	• Average Key Stage 4 (GCSE or equivalent) total scores for all eligible pupils in a middle	 National Pupil Databas e (2011) Office for Students . Started a course 	Some	Y		

			layer super	between		
			output area	2009-10		
			(MSOA).	and		
			• POLAR4	2013-		
			assigns a	14.		
			quintile to			
			each			
			MSOA			
			based on			
			how many	C.		
			young			
			people from			
			that MSOA			
			started a			
			higher			
			education			
		_4	course.			
			Proportion			
			residents			
			with Level 3			
		0,	qualification			
		Neighbours	S			
		with	(requirement			
		qualification	for entry to	Derived		
Co	llective	s, skilled	higher	from UK	g .	3. 7
Soci	alization	occupations	education or	Census	Some	Y
		and	skilled	(2011)		
		economically	employment)			
		active				
			• Proportion			
			occupied			
			persons in the			
			LSOA with			

		classes 1, 2 or 3 / 5 classes. • Proportion of usual residents			
		aged 16 to 74 in the LSOA who were economically active on census day.	Ö		
Social Networks	Social isolation	Proportion of population in an LSOA that is aged 10-19 years.	UK Census (2011)	Y	Y
Social cohesion and control	None	-	-	N	Y
Competition	Housing availability & affordability	Median price paid for lower layer super output areas	Office for National Statistics (2009- 2021)	Some	Y
Relative Deprivation	Townsend Index	z-score summary derived from four census variables: % unemployed, % non-car	Derived from UK Census (2011)	Some	Y

			Τ			
			owners, %			
			non-			
			homeowners,			
			and %			
			overcrowded			
			households in			
			each LSOA.			
	Parental	None	-	-	Y	Y
	Mediation		Rate of			
			recorded crime	8		
			in an area for	To do . C		
			four major	Index of		
	T	T 1	crime types	Multiple		
	Exposure to	Local area	representing	Deprivatio	Some	Y
	Violence	crime	the risk of	n Crime		
			personal and	score		
			material	(2015)		
			victimisation			
Environment			at a small area			
al		<u>U</u>	level.			
Mechanisms		•		• ONS		
			Rural/urban	postcode		
		Urbanicity	classificatio	directory		
	Physical	and	n	(2011)	Some	Y
-	Surroundings	population	Persons per	• Derived	201110	•
		density	hectare	from		
			nectare	Census		
				(2011)		
			Annual	Access to		
	Toxic	Air pollution		Healthy	Some	Y
	Exposure	7 m ponduon	mean	Assets &	Some	1
			nitrogen	Hazards		

			dioxide	(AHAH)		
			(µgm3)	dataset,		
			• Annual	version 2		
				(2015)		
			mean particulate	(2013)		
			Matter 10			
			(µgm3)			
			Annual			
			mean			
			Sulphur	<u> </u>		
			Dioxide			
			(µgm3)	\cup		
			Average travel			
		Distance to	time in	Department		
Geographical	Spatial	employment	minutes to	for	Y	Y
Mechanisms	Mismatch	centres	employment	Transport		
			centre by	(2015)		
			walking			
			• Average			
			travel time in			
		O	minutes to			
			nearest			
			hospital by	Department		
		Distance to	walking	for		
	Public	hospitals,	• Average	Transport -		
-	Services	doctors'	travel time in	journey	Y	Y
	Services	offices and	minutes to	time		
		job centres	nearest GP	statistics		
			by walking	(2015)		
			• Average			
			travel time in			
			minutes to			
			employment			
						I

			center by			
			walking			
Institutional	Stigmatizatio					
Mechanisms	n	None	-	-	Y	Y
	Local Institutional Resources	Distance to further education and green space	Total green space areas available to each postcode in a range of a 900-meter buffer (passive) before creating LSOA level averages.	Access to Healthy Assets & Hazards (AHAH) dataset, version 2 (2015)	Y	Y
	Local Market Actors	Distance to pharmacies, food stores, gambling shops, fast food, pubs, leisure and tobacco shops	Mean travel times in minutes by walking (exception pharmacies is cycling).	Department for Transport - journey time statistics (2015) for pharmacies and food stores. Other measures from Access to Healthy Assets & Hazards (AHAH) dataset,	Some	Y

	version 2	
	(2015)	

Table 2: Distribution of environment variables for analysis sample: all, Inland community, Coastal community, UKHLS youth sample, 2009-2021 (n=4.921)

	Total	Inland (4,157)	Coastal	p-value
	(n=4,921)	<u> </u>	(n=764)	
Townsend index:				
Quintile 1 (ref – least deprived)	31.5	31.9	29.5	
Quintile 2	18.7	17.7	24.5	
Quintile 3	15.5	14.5	20.8	
Quintile 4	17.7	17.7	17.8	
Quintile 5 (most deprived)	16.6	18.3	7.0	< 0.001
Economic Environment:				
Mean % unemployed 2011 (SD)	4.9 (2.7)	4.8 (2.6)	5.0 (2.8)	0.182
Mean % overcrowded 2011 (SD)	3.3 (4.5)	3.6 (4.8)	1.4 (1.2)	< 0.001
Mean % no car 2011 (SD)	26.6 (17.2)	26.6 (17.6)	26.4 (14.7)	0.692
Mean % non-homeowner 2011 (SD)	37.5 (21.2)	37.9 (21.4)	35.1 (20.1)	0.001
Mean % economically inactive 2011 (SD)	30.8 (7.1)	30.7 (7.2)	31.6 (6.6)	0.001
Social Environment:				
Mean % qualifications level 3+ (SD)	37.3 (12.4)	38.0 (12.6)	33.5 (11.1)	< 0.001
Mean % adults low skilled occupations (SD)	16.4 (6.8)	16.0 (6.7)	18.5 (7.1)	< 0.001
Index of deprivation Crime quintile, 2015				
Quintile 1 (ref – least deprived)	19.4	20.0	16.4	
Quintile 2	17.4	16.6	21.6	
Quintile 3	20.5	20.8	19.0	
Quintile 4	19.5	19.8	18.1	
Quintile 5 (most deprived)	23.3	23.0	25.0	0.002
Mean % age 10-19 years (SD)	12.8 (2.9)	12.9 (2.9)	12.4 (2.9)	< 0.001
Educational Environment: a				
% 5+ A*-C grades, English & maths	58.5 (13.1)	59.2 (13.0)	54.6 (13.1)	< 0.001
Mean capped point score for MSOA (SD)	339 (24)	340 (24)	334 (26)	< 0.001
POLAR participation rates	38.7 (16.0)	40.2 (16.1)	30.4 (12.6)	< 0.001
POLAR progress higher education quintiles:				
Quintile 1 (ref – least progression)	16.1	13.1	32.6	
Quintile 2	19.6	18.5	25.4	
Quintile 3	21.4	21.4	21.5	
Quintile 4	22.8	24.3	14.9	
Quintile 5 (most progression)	20.1	22.7	5.6	< 0.001
Built Environment:				
Urban, %	83.0	81.9	89.1	< 0.001
Geometric mean persons per hectare	3.2 (1.5)	3.2 (1.6)	3.2 (1.4)	0.627
(SD)				

Geometric mean travel times in minutes (SD):				
Pharmacy, cycle	7.8 (1.7)	7.9 (1.7)	7.8 (1.5)	0.502
Food stores, walk	7.7 (1.7)	7.8 (1.7)	6.9 (1.6)	< 0.001
Hospital, walk	33.6 (1.6)	33.8 (1.6)	32.3 (1.7)	0.015
GP, walk	9.3 (1.7)	9.2 (1.7)	9.7 (1.5)	0.008
Further Education, walk	17.3 (1.7)	17.2 (1.7)	18.1 (1.5)	0.006
Large employment centres, walk	27.6 (1.8)	27.0 (1.8)	31.2 (1.8)	< 0.001
Nearest rail station, public transport	54.4 (1.4)	53.1 (1.4)	61.9 (1.4)	< 0.001
AHAH measures (SD)				
Geometric mean minutes to gambling shops	1.3 (2.5)	1.3 (2.6)	1.2 (2.2)	0.001
Geometric mean minutes to fast food shops	1.2 (2.7)	1.2 (2.8)	1.2 (2.7)	0.662
Geometric mean minutes to pubs	1.0 (2.3)	1.0 (2.3)	1.0 (2.2)	0.236
Geometric mean minutes to leisure	2.3 (2.5)	2.2 (2.5)	2.4 (2.3)	0.017
Geometric mean green space within 900m	2.1 (2.2)	2.1 (2.3)	1.8 (1.9)	<0.001
Geometric mean minutes to tobacco shops	3.0 (2.4)	3.0 (2.4)	2.7 (2.5)	0.001
Mean annual nitrogen dioxide (μgm³)	12.3 (4.1)	12.6 (4.3)	10.9 (2.8)	<0.001
Mean annual Particulate Matter 10 (μgm³)	13.4 (1.7)	13.6 (1.7)	12.7 (1.5)	<0.001
Mean annual Sulphur Dioxide (μgm³)	1.28 (0.31)	1.27 (0.27)	1.33 (0.45)	<0.002

Table 3: Average individual and environmental variables for UKHLS youth sample members who lived in Coastal lower-super output areas (difference Inland - Coastal areas) by Area deprivation quartiles, 2009-2021 (n=4,921)

	Least	Quartile 2	Quartile 3	Most	Most
	deprived	(n=922)	(n=763)	deprived	deprived
	Quartile			quartiles (4)	quartile (5)
	(1)			(n=870)	(n=817)
	(n=1,459)				
Individual variables:					
Mean wave at baseline	5.3 (-	5.2 (-0.1)	4.8 (0.6)*	4.9 (0.4)	5.5 (-0.3)
	0.01)*				
Mean age at baseline	14.8 (0.0)	14.8 (0.1)	14.7 (0.0)	14.8 (-0.1)	14.8 (-0.1)
Male, %	49.8 (3.1)	51.3 (-1.1)	49.7 (3.5)	55.6 (-3.3)	49.1 (3.3)
Non-white ethnicity, %	4.9 (2.7)	4.8 (8.7)**	11.3	10.3 (-	15.8
			(20.5)**	44.2)**	(65.7)**
Tenure, %					
Homeowner	86.2 (1.6)*	75.9 (0.8)	66.0 (-2.8)	48.5 (7.6)	35.1 (3.1)
Social renter	2.2 (2.9)	13.4 (-1.1)	18.9 (7.1)	33.8 (2.1)	49.1 (3.9)
Private renter/Other	11.6 (-4.5)	10.7 (0.3)	15.1 (3)	17.1 (-5.5)	15.8 (-7.0)
Median adjusted* household income	404 (12)	349 (10)	298 (22)	258 (2)	210 (2)
per week					
Built Environment:				-	

Urban, %	78.7 (- 17.8)**	86.1 (- 11.3)**	94.3 (-2.0)	100.0 (-0.2)	100.0 (0.0)
Geometric mean persons per	10.4 (-	22.9 (-	39.9 (-	50.6 (0.9)**	43.3
hectare	1.5)**	1.3)*	1.2)*		(0.5)**
Geometric mean travel times in minutes:					
Pharmacy, cycle	9.9 (0.9)**	7.6 (0.9)*	6.9 (1.0)	6.7 (-1.1)*	6.1 (- 1.2)**
Food stores, walk	9.2 (0.8)**	6.6 (0.8)**	6.0 (0.9)*	5.7 (0.9)	6.1 (-1.1)
Hospital, walk	37.3 (0.8)**	32.5 (0.9)**	29.5 (0.9)	28.6 (-1.0)	30.7 (- 1.3)**
GP, walk	12.1 (0.9)*	9.8 (0.9)*	8.8 (1.0)	8.8 (-1.3)**	6.7 (- 1.2)**
Further Education, walk	21.3 (0.9)*	18.2 (1.0)	16.5 (-1.0)	16.7 (-1.1)**	15.1 (- 1.3)**
Large employment centres, walk	37.3 (-1.0)	31.9 (-1.0)	31.8 (- 1.3)**	27.5 (-1.2)**	18.3 (-1.0)
Job Centre, public transport	4.2 (0.0)	4.2 (-0.5)*	4.1 (- 0.1)**	4.1 (-0.2)**	4.0 (- 0.5)**
Nearest rail station, public	65.8 (-1.0)	64.3 (-	59.2 (-	31.3.7 (-	30.7 (-
transport		1.1)*	1.1)**	1.2)**	1.5)**
Geometric mean distance to:					
Gambling shops	2.0 (0.7)**	1.2 (0.8)**	0.9 (0.9)*	0.8 (-0.9)	0.7 (-1.2)*
Fast food shops	2.1 (0.8)*	1.2 (0.8)**	0.9 (0.9)	0.8 (1.0)	0.7 (1.5)**
Pubs	1.3 (0.8)**	1.0 (0.9)	0.8 (-1.0)	0.8 (0.9)	0.9 (- 1.4)**
Leisure	3.6 (0.9)*	2.4 (0.9)*	0.9 (0.9)	1.9 (-1.2)**	1.7 (- 1.6)**
Tobacco shops	4.3 (0.8)**	2.7 (0.8)**	2.4 (-1.0)	1.8 (0.8)**	1.7 (1.0)
Green space within 900m	0.6 (0.2)*	0.7 (0.1)	0.5 (0.3)**	0.6 (0.2)**	0.6 (0.1)
Mean annual nitrogen dioxide	10.2 (-	10.6 (0.0)	11.0	14.1 (2.5)**	12.6
(μgm³)	0.4)*		(1.3)**		(5.2)**
Mean annual Particulate Matter	12.6	13.0 (0.2)	12.5	12.6 (1.2)**	12.7
10 (μgm³)	(0.3)**	()	(0.9)**	(0 .)	(2.3)**
Mean annual Sulphur Dioxide	1.3 (-	1.3 (0.0)	1.3 (0.0)	1.4 (0.1)	1.7 (-
(μgm³) Social Environment:	0.1)**				0.4)**
Mean % qualifications level 3+	40.9	35.2	31.8	25.4 (-4.8)**	21.9 (-
Weart 70 qualifications level 5.	(4.4)**	(3.8)**	(5.4)**	23.4 (4.0)	10.4)**
Mean % adults low skilled	12.6	16.2 (0.4)	20.4	25.6 (4.6)**	27.4
occupations	(1.4)**	- (- ,	(2.8)**	(- ,	(9.1)**
Index of deprivation Crime	,		` ,		` ,
quintile, 2015					
Quintile 1 (ref – least	43.6 (-6.6)	9.6 (-10.4)	5.0 (2.9)	0.74 (-0.1)	0.00 (-0.1)
deprived)					
Quintile 2	33.3 (5.5)	31.0 (3.7)	17.6 (5.5)	2.2 (-1.9)	1.8 (0.3)
Quintile 3	15.6 (-0.4)	27.3 (-4.2)	20.8 (- 16.0)	15.4 (-1.7)	8.8 (0.8)
Quintile 4	4.9 (-0.6)	18.7 (3.5)	32.7 (3.7)	26.5 (-8.3)	7.0 (20.2)
Quintile 5 (most deprived)	2.7 (2.1)**	13.4	23.9	55.2 (11.8)	82.5
		(7.4)**	(3.9)**		(21.5)*

Mean % age 10-19 years (SD)	11.5 (0.7)**	12.0 (-0.2)	12.4 (-0.1)	13.6 (0.2)	14.3 (0.1)
Economic Environment:					
Mean % unemployed 2011 (SD)	2.6 (-	3.8 (-	5.2 (-	7.6 (-0.9)**	11.1 (-
	0.2)**	0.2)**	0.4)**		2.7)**
Mean % overcrowded 2011 (SD)	0.43	0.99	1.56	2.29	3.34
	(0.06)*	(0.14)*	(0.81)**	(3.00)**	(7.57)**
Mean % no car 2011 (SD)	12.3 (-	20.3 (-	29.9	41.0 (-3.7)**	56.6 (-2.7)
	2.8)**	2.5)**	(2.8)**		
Mean % non-homeowner 2011	14.4	27.8 (1.3)	39.9 (0.5)	57.4 (-6.3)**	74.8 (-
(SD)	(2.2)**				6.0)**
Mean % economically inactive	30.5 (-	29.2 (-	30.7 (-	34.1 (0.2)	40.2 (-
2011 (SD)	3.1)**	2.2)**	1.4)*		2.7)**
Educational Environment: a					
Mean % 5+ A*-C grades, English &	64.3 (-	55.9 (-	52.4 (-	44.9 (-5.7)**	41.1 (-
maths ^b	3.4)**	4.4)**	4.9)**		12.0)**
Mean total point score for MSOA	490 (-0.7)	472 (-6.6)	469 (-4.3)	446 (-7.1)	432 (-
(SD) ^c				~	21.9)**
POLAR higher education quintiles:					
Quintile 1 (ref – least	7.6 (-4.7)	24.1 (-	35.9 (-	67.8 (-39.2)	66.7 (-
participation)		11.5)	14.0)		57.1)
Quintile 2	17.3 (-4.6)	29.4 (-4.1)	33.3 (-9.3)	21.3 (-0.5)	31.6 (-
					16.2)
Quintile 3	32.9 (-	23.5 (1.0)	18.9 (-0.8)	11.0 (5.5)	1.8 (28.1)
	13.7)				
Quintile 4	32.4 (-4.7)	18.2 (-1.3)	4.4 (10.3)	0.0 (19.8)	0.0 (37.2)
Quintile 5 (most participation)	9.8	4.8	7.6	0.0 (14.4)**	0.0 (7.9)**
	(27.7)**	(16.1)**	(13.8)**		

Table 4: Estimated mean SF-12 mental functioning component (MCS) score by 1-unit change in environmental variables, UKHLS youth sample members, 2009-2021 (n=4,921)

	Model 1: age-only	Model 2: + individual socio-demographics	Model 3: full model (fitted jointly)
Economic Environment:			_
Mean % unemployed 2011 (SD)	0.14 (0.03, 0.25)	-0.01 (-0.13, 0.11)	-
Mean % overcrowded 2011 (SD)	0.19 (0.12, 0.25)	0.03 (-0.01, 0.07)	-
Mean % no car 2011 (SD)	0.02 (0.00, 0.04)	0.10 (0.02, 0.18)	0.02 (-0.01, 0.05)
Mean % non-homeowner 2011 (SD)	0.01 (-0.002 <i>,</i> 0.02)	-0.01 (-0.03, 0.01)	-
Mean % economically inactive 2011 (SD)	0.08 (0.04, 0.12)	-0.01 (-0.02, 0.01)	-
Social Environment:			
Mean % qualifications level 3+ (SD)	-0.02 (-0.04 <i>,</i> 0.005)	-0.01 (-0.03, 0.02)	-

Mean % adults low skilled occupations (SD) Index of deprivation Crime quintile, 2015	0.01 (-0.04, 0.05)	-0.01 (-0.05, 0.04)	-
Quintile 1 (ref – least deprived)	-	-	-
Quintile 2	-0.72 (-1.63, 0.19)	-0.83 (-1.70, 0.04)	-0.70 (-1.58, 0.18)
Quintile 3	0.03 (-0.85, 0.91)	-0.31 (-1.18, 0.54)	-0.06 (-1.00, 0.88)
Quintile 4	0.57 (-0.29, 1.42)	-0.22 (-1.09, 0.65)	-0.03 (-1.03, 097)
Quintile 5 (most deprived)	-0.16 (-1.04, 0.72)	-1.33 (-2.26, - 0.41)	-1.20 (-2.38, - 0.03)
Mean % age 10-19 years (SD)	0.17 (0.08, 0.26)	0.10 (0.01, 0.19)	0.09 (-0.004,
Wedit 70 age 10 13 years (05)	0.17 (0.00) 0.120)	0.10 (0.01) 0.115)	0.19)
Educational Environment: ^a			0.20)
% 5+ A*-C grades, English & maths	0.0 (-0.02, 0.02)	0.01 (-0.01, 0.04)	-
Mean capped point score for MSOA (SD)	0.0 (-0.01, 0.01)	0.00 (0.00, 0.01)	-
POLAR participation rates	0.01 (-0.01, 0.03)	0.01 (-0.01, 0.02)	-
POLAR progress higher education quintiles:	0.01 (0.01, 0.00)	0.01 (0.01, 0.02)	
Quintile 1 (ref – least progression)	_	<u> </u>	-
Quintile 2	0.38 (-0.60, 1.36)	0.21 (-0.76, 1.17)	0.20 (-0.79, 1.20)
Quintile 3	1.51 (0.56, 2.46)	1.07 (0.12, 2.01)	1.07 (0.09, 2.05)
Quintile 4	0.85 (-0.07, 1.77)	0.57 (-0.36, 1.50)	0.65 (-0.34, 1.64)
Quintile 5 (most progression)	0.85 (-0.09, 1.79)	0.62 (-0.33, 1.57)	0.75 (-0.30, 1.80)
	0.03 (0.03) 1.73)	0.02 (0.00, 1.07)	0.75 (0.50, 1.60)
Built Environment:			
Urban, %	0.05 (-0.71, 0.82)	-0.47 (-1.21, 0.28)	-
Geometric mean persons per hectare (SD)	1.12 (0.93, 1.34)	0.86 (0.71, 1.04)	-
Geometric mean travel times in minutes (SD):			
Pharmacy, cycle	0.59 (0.34, 1.03)	1.04 (0.59, 1.85)	-
Food stores, walk	1.26 (0.74, 2.14)	1.68 (1.00, 2.84)	-
Hospital, walk	0.62 (0.36, 1.09)	1.12 (0.63, 2.00)	-
GP, walk	0.63 (0.37, 1.08)	1.14 (0.65, 2.00)	-
Further Education, walk	0.98 (0.57, 1.70)	0.77 (1.00, 3.12)	-
Large employment centres, walk	0.58 (0.36, 0.93)	0.98 (0.60, 1.61)	-
Nearest job centre, public transport	0.27 (0.12, 0.59)	0.97 (0.40, 2.36)	-
AHAH measures (SD)			
Geometric mean minutes to gambling	0.80 (0.59, 1.08)	1.17 (0.86, 1.60)	-
shops			
Geometric mean minutes to fast food	0.82 (0.62, 1.09)	1.21 (0.90, 1.63)	-
shops			
Geometric mean minutes to pubs	1.09 (0.79, 1.52)	1.46 (1.06, 2.02)	1.36 (0.88, 2.09)
Geometric mean minutes to leisure	0.93 (0.68, 1.27)	1.41 (1.01, 1.95)	1.19 (0.77, 1.86)
Mean green space within 900m	-0.07 (-0.37, 0.24)	-0.08 (-0.39, 0.22)	-
Geometric mean minutes to tobacco	-0.01 (-0.33, 0.32)	1.33 (0.96, 1.85)	-
shops			
Mean annual nitrogen dioxide (μgm³)	0.16 (0.09, 0.23)	0.05 (-0.03, 0.13)	-
Mean annual Particulate Matter 10 (μgm³)	0.18 (0.01, 0.34)	0.00 (-0.18, 0.18)	-
Mean annual Sulphur Dioxide (μgm³)	0.90 (-0.04, 1.85)	0.67 (-0.25, 1.58)	-

Table 5: Adjusted associations of SF-12 mental functioning score (95% CI) in young adulthood across 11 waves of follow-up, by environmental factors in adolescence, UKHLS youth sample, 2009-2021 (n= n=4,921, observations=18,324): Townsend index as quintiles

observations=16,524). Townse	nu muex as quint				
	Model 1:	Model 2: +	Model 3: +	Model 4: +	Model 5:
	Coastal, Area	Individual	IMD Crime &	Higher	Full model
	Deprivation,	socio-	Coastal*Crime	Education	
	Coastal*	demographics	only	Progression	
	Townsend &	only*	•	only	
	Age at	····,		····/	
	outcome				
Coastal	0.7 (-0.6, 2.0)	0.6 (-0.7, 1.9)	0.9 (-0.8, 2.6)	0.7 (-0.6,	0.7 (-0.8,
Coastai	0.7 (-0.0, 2.0)	0.0 (-0.7, 1.9)	0.9 (-0.8, 2.0)		
Townsend Index:				2.0)	1.9)
1 (least deprived)	-	-	-	-	-
2	-0.5 (-1.4, 0.4)	-0.8 (-1.6, 0.1)	-0.5 (-1.5, 0.5)	-0.5 (-1.4,	-0.7 (-1.6,
				0.4)	0.2)
3	-0.9 (-1.9, 0.1)	-1.2 (-2.2, -	-0.8 (-2.0, 0.4)	-0.8 (-1.8,	-1.0 (-2.2,
		0.2)	X	0.2)	0.1)
4	0.6 (-0.3, 1.5)	-0.2 (-1.1, 0.7)	1.1 (-0.1, 2.2)	0.8 (-0.1,	0.3 (-0.9,
				1.7)	1.5)
5 (most deprived)	1.9 (1.0, 2.8)	0.7 (-0.5, 1.9)	2.6 (1.4, 3.9)	1.9 (1.0, 2.9)	1.4 (-0.1,
			<i>,</i> ()		2.8)
Townsend*Coastal:					
1 (least deprived) x	-	-	-	-	-
coastal					
2 x coastal	-1.7 (-3.8,	-1.4 (-3.5, 0.6)	-1.7 (-4.0, 0.7)	-1.6 (-3.8,	-1.3 (-3.4,
	0.42)			0.5)	0.7)
3 x coastal	0.7 (-1.6, 3.0)	0.8 (-1.4, 3.1)	0.4 (-2.4, 3.2)	0.7 (-1.5,	0.9 (-1.3,
3 x coastai	0.7 (1.0, 3.0)	0.0 (1.4, 5.1)	0.4 (2.4, 3.2)	3.0)	3.3)
4 v coastal	-2.7 (-4.9, -0.4)	-1.7 (-4.0, 0.5)	24164	-2.5 (-4.7, -	
4 x coastal	-2.7 (-4.9, -0.4)	-1.7 (-4.0, 0.5)	-3.4 (-6.4, -		-1.6 (-3.8,
E (see a st. days six a d)	F 4 / G 4 2 2 3	42/70	0.4)	0.2)	0.7)
5 (most deprived) x	-5.1 (-8.1, -2.2)	-4.3 (-7.0, -	-6.1 (-9.7, -	-4.7 (-7.7, -	-3.9 (-6.7, -
coastal		1.6)	2.4)	1.7)	1.1)
Age at MCS assessment	-0.7 (-0.8, -0.6)	-0.7 (-0.8, -	-0.7 (-0.8, -	-0.7 (-0.8, -	-0.7 (-0.8, -
		0.6)	0.6)	0.6)	0.6)
Gender, 15y	-	-5.0 (-5.5, -	-	-	-5.0 (-5.5 <i>,</i> ·
		4.5)			4.5)
Non-white ethnicity, 15y	-	1.2 (0.5, 2.0)	-	-	1.2 (0.4, 1.9
Median adjusted* household					
income, 15y					
1 (lowest incomes)	-	-	-	-	-
2	-	0.0 (-0.7, 0.7)	-	-	0.0 (-0.7,
		, , ,			0.7)
3	-	0.3 (-0.5, 1.1)	-	-	0.3 (-0.5,
-		3.5 (3.5, 1.1)			1.1)
4	_	0.7 (-0.1, 1.5)	_	_	0.7 (-0.1,
7	_	5.7 (5.1, 1.5)	_	_	1.5)
E (highest incomes)		1 4 (0 6 2 2)			
5 (highest incomes)	-	1.4 (0.6, 2.3)	-	-	1.4 (0.6, 2.3
Tenure, 15y					
Homeowner		-	-	-	-
Social renter	-	-0.7 (-1.5, 0.1)	-	-	-0.7 (-1.5,
					0.2)

Private renter	-	-1.4 (-2.5, - 0.2)	-	-	-1.4 (-2.5, - 0.2)
Other	-	-0.4 (-2.3, 1.5)	-	-	-0.4 (-2.3, 1.4)
IoD Crime quintile (2015):					•
1 (ref – least crime)	-	-	-	-	-
2	-	-	-0.6 (-1.6, 0.4)	-	-0.7 (-1.6,
3	-	-	0.3 (-0.8, 1.4)	-	0.3) 0.3 (-0.7, 1.4)
4	-	-	-0.1 (-1.3, 1.1)	-	-0.2 (-1.4, 1.0)
5 (most crime)	-	-	-1.4 (-2.8, - 0.1)	-	-1.6 (-2.9, - 0.3)
IoD Crime*Coastal					-
1 (least crime) x coastal	-	-	- 🗙	-	-
2 x coastal	-	-	0.0 (-2.5, 2.6)	-	0.1 (-2.4,
			() `		2.5)
3 x coastal	-	-	-1.5 (-4.3, 1.3)	-	-1.6 (-4.4,
					1.2)
4 x coastal	-	-	0.7 (-2.3, 3.6)	-	1.0 (-1.9,
					3.9)
5 (most crime) x coastal	-	-	-1.3 (-2.0, 4.5)	-	-1.4 (-1.9, 4.6)
POLAR 4 quintile:					4.0)
1 (ref – most	_		-	-	_
participation)					
2	- 🔍	-	-	0.3 (-0.7,	-0.1 (-1.0,
				1.3)	0.9)
3		-	-	1.2 (0.1, 2.2)	0.6 (-0.4,
					1.6)
4		-	-	0.3 (-0.6,	0.0 (-1.0,
				1.3)	1.1)
5 (least participation)	-	-	-	0.8 (-0.2,	0.2 (-0.8,
	*			1.8)	1.2)
R-square:					
Within	0.0375	0.0369	0.0375	0.0375	0.0368
Between	0.0137	0.0914	0.0167	0.0151	0.0960
Overall	0.0182	0.0618	0.0198	0.0196	0.0644

Ethics Approval

The University of Essex Ethics Committee has approved all data collection on Understanding Society main study and innovation panel waves, including asking consent for all data linkages except to health records. Requesting consent for health record linkage was approved at <u>Wave</u> 1 by the National Research Ethics Service (NRES) Oxfordshire REC A (08/H0604/124) and at Wave 4 by NRES Southampton REC A (11/SC/0274).

More information is provided on the Understanding Society study website: https://www.understandingsociety.ac.uk/documentation/mainstage/user-guides/main-survey-user-guide/ethics/

Declaration of interests

☑ The authors declare that they have no known competing financial interests or personal relationships
that could have appeared to influence the work reported in this paper.
\mathcal{O}_{1}
☐ The authors declare the following financial interests/personal relationships which may be considered
as potential competing interests: