

Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: An updated analysis from a Phase II, randomised, double-blind, placebo-controlled trial

Jonathan A Ledermann, MD¹ Philipp Harter, MD² Charlie Gourley, PhD³ Michael Friedlander, PhD⁴ Ignace Vergote, PhD⁵ Gordon Rustin, MD⁶ Clare Scott, PhD⁷ Werner Meier, MD⁸ Ronnie Shapira-Frommer, MD⁹ Tamar Safra, MD¹⁰ Daniela Matei, MD¹¹ Anitra Fielding, MBChB¹² Stuart Spencer, MSc¹² Philip Rowe, MSc¹² Elizabeth Lowe, MD¹³ Darren Hodgson, PhD¹² Mika A Sovak, MD¹³ and Ursula Matulonis, MD¹⁴

¹UCL Cancer Institute, University College London and UCL Hospitals, London, UK; ²Kliniken Essen Mitte, Essen, Germany; ³University of Edinburgh Cancer Research UK Centre, Medical Research Council Institute of Genetics and Molecular Medicine, Western General Hospital, Edinburgh, UK; ⁴University of New South Wales Clinical School, Prince of Wales Hospital, Randwick, NSW, Australia; ⁵University of Leuven, Leuven, Belgium; ⁶Mount Vernon Hospital, Northwood, UK; ⁷Royal Melbourne Hospital, Parkville, VIC, Australia; ⁸Evangelisches Krankenhaus, Düsseldorf, Germany; ⁹Chaim Sheba Medical Center, Tel Hashomer, Israel; ¹⁰Tel Aviv Sourasky Medical Center, Tel Aviv, Israel; ¹¹Northwestern University Feinberg School of Medicine, Chicago, IL, USA; ¹²AstraZeneca, Alderley Park, Macclesfield, UK; ¹³AstraZeneca, Gaithersburg, MD, USA; ¹⁴Dana-Farber Cancer Institute, Boston, MA, USA

Article type: Original research article

Running title: Overall survival with olaparib maintenance monotherapy

Figures/tables: 5 figures/4 tables

References: 23

Corresponding author:

Professor Jonathan A Ledermann

Cancer Research UK and UCL Cancer Trials Centre

90 Tottenham Court Road

London, W1T 4TJ, UK

Phone: +44 20 7679 9898

Fax: +44 20 7679 9899

Email: j.ledermann@ucl.ac.uk

Summary

Background: In patients with platinum-sensitive recurrent serous ovarian cancer (PSR SOC), maintenance monotherapy with the PARP inhibitor olaparib (Lynparza™) significantly improves progression-free survival (PFS) versus placebo. We assessed the effect of maintenance olaparib on overall survival (OS) in patients with PSR SOC, including those with *BRCA1/2* mutations (*BRCAM*).

Methods: In this Phase II trial, patients had PSR SOC, had received ≥ 2 courses of platinum-based chemotherapy and responded to their latest regimen. Patients were randomised, using a computer-generated sequence, to oral maintenance olaparib (400 mg twice daily; capsules) or placebo by an interactive voice response system, stratified by ancestry, time to progression on penultimate platinum and response to most recent platinum. The primary endpoint was PFS. Here, we present data for OS, a secondary endpoint, from the third data analysis after >5 years' follow-up (intention-to-treat population). Randomised patients were analysed for OS; treated patients were analysed for safety. This trial is ongoing and is registered with ClinicalTrials.gov (NCT00753545).

Findings: Between 28 August 2008 and 9 February 2010, 136 patients were randomised to olaparib and 129 to placebo. 136 patients had deleterious *BRCAM*. The data cut-off for this analysis was 30 September 2015. An OS advantage was observed with maintenance olaparib versus placebo in all patients (HR 0.73 [95% CI 0.55–0.96]; median OS 29.8 vs 27.8 months) and *BRCAM* patients (HR 0.62 [0.41–0.94]; 34.9 vs 30.2 months). 11 (15%) of 74 *BRCAM* patients received maintenance olaparib for ≥ 5 years. Overall, common grade ≥ 3 adverse events (AEs) were fatigue (olaparib: 11/136 patients [8%]; placebo: 4/128 [3%]) and anaemia (olaparib: 8/136 patients [6%]; placebo: 1/128 [1%]). Serious AEs were reported in 30/136 patients (22%) on olaparib and 11/128 patients (9%) on placebo. In patients treated for ≥ 2 years, AEs included low-grade nausea (olaparib: 24/32 patients [75%]; placebo: 2/5 patients [40%]), fatigue (18/32 [56%]; 2/5 [40%]), vomiting (12/32 [38%]; 0) and anaemia (8/32 [25%]; 1/5 [20%]); generally, events were initially reported during the first 2 years of treatment.

Interpretation: An OS advantage was seen for patients with *BRCAm* PSR SOC receiving olaparib maintenance monotherapy after platinum-based chemotherapy, supporting the reported PFS benefit. Significant long-term exposure to olaparib was observed. There were no new safety signals. Taken together, these data support both the long-term clinical benefit and tolerability of maintenance olaparib in *BRCAm* patients with PSR SOC.

Funding: AstraZeneca.

Introduction

Ovarian cancer is the fifth most common type of cancer for women in developed countries.^{1,2} Approximately 70% of patients relapse within 3 years of completing first-line chemotherapy and the mean 5-year survival rate in Europe is low when compared with other tumour types (approximately 38%).^{3–5} Overall, ovarian cancer is the sixth highest cause of cancer-related deaths for women in developed countries.^{1,2}

Olaparib (Lynparza™) is an oral poly(ADP-ribose) polymerase (PARP) inhibitor that has demonstrated significant clinical activity in ovarian cancer, particularly in tumours that have mutations in *BRCA1/2* (*BRCAm*).^{6–8} Olaparib traps PARP at sites of DNA damage, blocking base-excision repair and resulting in the collapse of DNA replication forks and the accumulation of DNA double-strand breaks.⁹ Induced synthetic lethality is observed with olaparib in tumours that are deficient in homologous recombination repair (HRR) pathways, such as those with *BRCAm*.^{10,11}

Previously, we reported data from a Phase II, randomised, double-blind trial (NCT00753545, D0810C00019 [Study 19]) that demonstrated a statistically significant improvement in progression-free survival (PFS) for patients with platinum-sensitive, recurrent serous ovarian cancer (PSR SOC) who received olaparib maintenance monotherapy, compared with placebo (hazard ratio [HR] 0.35, 95% confidence interval [CI] 0.25–0.49, $P < 0.0001$).^{6,7} A pre-planned analysis of the retrospectively identified *BRCAm* subgroup showed patients with a *BRCAm* derived the greatest PFS benefit from olaparib (HR 0.18, 95% CI 0.10–0.31, $P < 0.0001$).⁷ A significant improvement in time to first subsequent therapy or death (TFST) and time to second subsequent therapy or death (TSST) was also observed with maintenance olaparib compared with placebo.⁷ Based on these data, olaparib (400 mg twice daily [bid]; capsules) was approved in the EU as maintenance therapy for patients with platinum-sensitive, relapsed, *BRCA*-mutated ovarian cancer.¹² Olaparib is also approved in the US as monotherapy for patients with germline *BRCA*-mutated advanced ovarian cancer.¹³ This indication was based on data from another Phase II study (NCT01078662, D0810C00042 [Study 42]).⁸

Two interim analyses of overall survival (OS) from Study 19 have previously been conducted, at 38% data maturity (HR 0.94, 95% CI 0.63–1.39, $P=0.75$) and 58% data maturity (HR 0.88, 95% CI 0.64–1.21, $P=0.44$) in the overall study population.^{6,7} Here, we present an updated descriptive OS analysis following the deaths of 203 (77%) of 265 patients in this study, with an additional 3 years of OS follow-up since the previous analysis. We assessed the impact of maintenance olaparib on OS in women with PSR SOC.

Methods

Study design and participants

Study 19 was a Phase II, randomised, double-blind, placebo-controlled, multicentre trial, involving 82 sites across 16 countries. The institutional review boards or independent ethics committees of all investigational sites approved the protocol. The study was conducted in accordance with the Declaration of Helsinki, Good Clinical Practice and the AstraZeneca policy on bioethics.¹⁴

Eligible patients were aged ≥ 18 years, with recurrent ovarian, fallopian tube or primary peritoneal cancer that had high-grade (grade 2 or 3) serous features or a serous component and was platinum-sensitive (no disease progression in the first 6 months after the last dose of the penultimate line of platinum-based chemotherapy). Patients must have received at least two prior courses of platinum-based chemotherapy and had to have shown a complete or partial response to their most recent regimen according to Response Evaluation Criteria in Solid Tumors (RECIST) v1.0 or Gynecological Cancer InterGroup criteria. Additional eligibility criteria have been described.⁶ All patients provided written informed consent.

Known *BRCAm* status was not required for eligibility, but was established via case report forms (CRF) documenting previous local germline *BRCA* testing, or via retrospective germline *BRCA* testing (Integrated *BRCA*Analysis assay [Myriad Genetics, Salt Lake City, UT, USA]) or tumour *BRCA* testing (next-generation sequencing [Foundation Medicine, Cambridge, MA, USA]), as described previously.⁷ Those patients whose *BRCAm* status was established during the study provided consent and samples at study entry.

Randomisation and masking

Patients were randomised (1:1) to receive olaparib or placebo within 8 weeks following completion of their most recent platinum-based regimen. An interactive voice response system (IVRS) assigned patients to their treatment, using a randomisation scheme generated by a computer program (GRand). The investigator who enrolled patients contacted an IVRS centralised randomisation office by

telephone for allocation of randomised treatment. Randomisation was stratified by ancestry (Jewish vs non-Jewish), time to progression from completion of penultimate platinum-based regimen (6–12 months vs >12 months) and response to most recent platinum-based regimen (complete vs partial response).

Treatment assignment was masked from patients and from anyone administering interventions, assessing outcomes or analysing data, by the use of unique identifiers generated during randomisation. Olaparib and placebo capsules were identical in appearance and packaging.

Procedures

Patients received oral olaparib maintenance monotherapy, at 400 mg bid (capsules; manufactured by AstraZeneca, Macclesfield, UK or Lonza, Visp, Switzerland) or matching placebo. Treatment continued until disease progression, provided that toxicities were manageable. After progression, patients could continue on study treatment if deemed appropriate by the investigator. Crossover between treatment arms within the study was not allowed. Dose modifications that were specified for toxicity management have been described previously.⁶

Tumours were assessed by computed tomography scans or magnetic resonance imaging every 12 weeks until week 60 and every 24 weeks thereafter until objective disease progression, unless patients withdrew consent. RECIST data were not collected after the primary data cut-off (DCO) of 30 June 2010. Patients were monitored for OS, with follow-up every 12 weeks after discontinuation of treatment. Safety and tolerability were monitored for patients remaining on study treatment by record of adverse events (AEs), physical examination, vital signs and laboratory findings. AEs were graded using the National Cancer Institute's Common Terminology Criteria for Adverse Events v3.0.

Outcomes

We have previously reported data for PFS, which represented the primary endpoint for this study.⁶ OS was a secondary endpoint, but represents the main outcome for

this descriptive analysis. Safety, tolerability, TFST and TSST were also assessed. Additional endpoints have been described previously.^{6,7}

Statistical analysis

Study 19 was sized to ensure a sufficient number of PFS events in the overall study population.⁷ OS was analysed on an intention-to-treat basis. The analysis set for OS included all randomised patients and the analysis sets for safety, TFST and TSST included all patients who received at least one dose of treatment. Other than for OS, no adjustments were made for multiplicity introduced by analysing multiple endpoints (TFST and TSST). No adjustments were made for analyses within the *BRCAm* or *BRCA* wild-type (*BRCAwt*) subgroups. Two previous analyses of OS have been conducted, at 38% data maturity (DCO: 31 October 2011; alpha [two-sided] = 0.1%) and 58% data maturity (DCO: 26 November 2012; alpha [two-sided] = 3%).^{6,7} The updated OS analysis described here was conducted at 77% data maturity, using an alpha (two-sided) of 0.95%. This OS analysis is considered to be descriptive and the *P*-values are nominal. Exploratory analyses of TFST and TSST were previously performed at the 2012 DCO, when these endpoints had 80% and 74% data maturity, respectively.⁷

OS, TFST and TSST were analysed using a Cox proportional hazards model, which was adjusted for treatment, ancestry (Jewish vs non-Jewish), time to progression from completion of penultimate platinum-based regimen (6–12 vs >12 months) and response to most recent platinum-based regimen (complete vs partial response). Restricted means analyses were performed for the OS data using the pseudovalues method, as previously described.¹⁵ All analyses used SAS v8.2 except the restricted means analyses, which used the Comprehensive R Archive Network “pseudo” software. This study is registered with ClinicalTrials.gov, number NCT00753545.

Role of the funding source

The corresponding author (JAL) designed the study in collaboration with the sponsor, AstraZeneca. AstraZeneca authors (AF, SS, PR, EL, DH and MAS) collected and analysed the data and had a role in data interpretation and manuscript

writing. All authors had access to the raw data. The decision to submit the manuscript for publication was made by all authors. The corresponding author (JAL) had full access to all of the data and the final responsibility to submit for publication.

Results

Patient enrolment occurred between 28 August 2008 and 9 February 2010. Of the 326 patients who enrolled, 265 met the eligibility criteria; 136 of these patients were randomised to olaparib and 129 were randomised to placebo (Figure 1). *BRCAM* status was established for 254/265 patients (96%), of whom 136 (51% of 265 patients in the overall study population) had a known or suspected deleterious *BRCAM*. Patient demographics and baseline characteristics were generally well balanced for the overall study population, *BRCAM* and *BRCAwT* subgroups (Table 1). The efficacy analysis set included all 265 randomised patients. One patient who was randomised to placebo withdrew consent and withdrew from the study without receiving treatment; therefore, the analysis sets for safety, TFST and TSST included the 264 patients who were treated.

The DCO for this updated OS analysis was 30 September 2015 (OS data maturity: 77%). At this DCO, the median follow-up for OS was 71.0 months (inter-quartile range [IQR] 67.8–72.9 months) for the overall study population (olaparib: 71.0 months [68.5–72.7] vs placebo: 70.8 months [38.2–73.0]). This represents an additional 3 years of follow-up since the previously reported OS.⁷ The Cox proportional hazards analyses indicate an OS advantage for patients who received olaparib maintenance monotherapy compared with patients who received placebo (HR 0.73, 95% CI 0.55–0.96, nominal $P=0.025$; Figure 2a), although this did not meet the required threshold for statistical significance ($P<0.0095$). The *BRCAM* subgroup data (70% OS data maturity) indicate an OS advantage for *BRCAM* patients who were treated with maintenance olaparib (HR 0.62, 95% CI 0.41–0.94, nominal $P=0.025$; Figure 2b). The OS data for the *BRCAwT* subgroup (84% OS data maturity) were: HR 0.83, 95% CI 0.55–1.24, nominal $P=0.37$ (Figure 2c).

Most patients in the *BRCAM* subgroup had germline *BRCAM* (*gBRCAM*), but 20 (15%) of 136 (olaparib: $n=10$, placebo: $n=10$) had somatic *BRCAM* (*sBRCAM*) only. We previously reported 18 *sBRCAM* patients in Study 19, based on data from tumour and blood testing, and 22 patients with tumour *BRCAM* for whom no blood testing data were available.⁷ Subsequently, we used an algorithm to distinguish *gBRCAM* and *sBRCAM* based solely on tumour sequencing data and identified the

20 *sBRCAm* patients who are discussed here: this group includes six of the 22 patients for whom blood testing data were unavailable and 14 of the original 18 *sBRCAm* patients.¹⁶ Four patients from the previously reported subgroup were therefore not included, three as a result of likely incomplete CRF-reported local blood-based *gBRCAm* tests and one as a result of discordant variant results, which revealed that the blood and tumour samples were from different individuals. Figure 3 shows the OS data for the overall, *BRCAm*, *gBRCAm* and *sBRCAm* populations. The *sBRCAm* subgroup data are not inconsistent with those from the other subgroups, but there are too few events in this group to draw conclusions. Figure 3 also shows the *BRCA1m* and *BRCA2m* OS data, and Kaplan-Meier survival curves for these two subgroups are presented in the Supplementary Material (page 2).

Formal tests of the proportionality of the hazards, using the methods of Grambsch and Therneau, indicated that there was insufficient evidence to dismiss the proportional hazards assumption in either the overall population ($P=0.19$) or the *BRCAm* subgroup ($P=0.70$).¹⁷ However, restricted means analyses were also performed, in order to enhance our understanding of average patient survival and the effect of study treatment. Table 2 shows the results from these restricted means analyses for the overall population and the *BRCAm* subgroup. These data are supportive of the OS advantage with olaparib indicated by the Cox proportional hazards analysis. In addition, the Supplementary Material (page 3) shows the restricted means data using two alternative methodologies, which gave similar estimates for the restricted mean OS. Log-rank test analyses were also consistent with the Cox proportional hazards analysis (Table 2).

Updated exploratory analyses were conducted for TFST and TSST; since the previous analysis, the data maturity had increased from 80% to 86% for TFST and from 74% to 84% for TSST.⁷ The median follow-up for TFST was 70.8 months (IQR 12.6–72.7 months) for the overall population (olaparib: 70.8 months [14.6–72.6] vs placebo: 39.0 months [4.1–74.7]); median follow-up for TSST was 70.5 months (IQR 11.2–72.8 months) for the overall population (olaparib: 70.9 months [16.4–72.6] vs placebo: 7.8 months [5.2–72.8 months]). Median TFST and TSST were significantly prolonged with olaparib compared with placebo, in the overall study population, *BRCAm* and *BRCAwt* subgroups (Figure 4).

At the DCO for this updated OS analysis, 15 (11%) of 136 patients were continuing to receive olaparib, eight of whom had a *BRCAM*. Within this group, the initial dose (olaparib 400 mg bid) was being received by nine patients (five *BRCAM*) and a reduced dose of 200 mg bid was being received by six patients (three *BRCAM*), four of whom had a dose reduction owing to AEs. One patient, with a *BRCAM*, was still receiving placebo (<1% of 129). Overall, 18 (13%) of 136 patients had received olaparib for ≥ 5 years (Table 3): 11 of these patients had a *BRCAM* (15% of 74) and seven were in the *BRCAMt* subgroup (12% of 57). Baseline characteristics for the patients who received study treatment for ≥ 5 years are listed in Table 4.

Subsequent cancer therapies had been received by 89 (65%) of 136 patients from the olaparib arm (45 of 74 *BRCAM* patients [61%]) and 111 (86%) of 129 patients from the placebo arm (55 of 62 *BRCAM* patients [89%]). From the placebo arm, 17 (13%) of 129 patients had received post-discontinuation PARP inhibitor (PARPi) treatment, of whom 14 (23%) of 62 patients had a *BRCAM*. These data include one additional patient who had received subsequent PARPi therapy since the previous DCO (26 November 2012). No patients from the olaparib arm had received subsequent PARPi treatment.

There were no new safety findings in the overall study population when compared to those that have previously been reported.^{6,7} Figure 5a shows the most common AEs reported between the start of treatment and the 2015 DCO by patients in the overall population who received treatment for ≥ 2 years. For the 32 patients who received olaparib for ≥ 2 years, 30 (94%) of 32 reported at least one AE, with 15 (47%) of 32 reporting AEs of grade ≥ 3 . For patients who received olaparib treatment for ≥ 2 years, the frequencies of previously reported common AEs, such as low-grade nausea, fatigue, vomiting and anaemia, were consistent with the frequencies that were previously reported in the overall population. In general, these AEs were initially reported during the first 2 years of treatment. Twenty-one *BRCAM* patients received olaparib for ≥ 2 years and this subgroup had a similar safety profile to the overall group of 32 patients. All five patients who received placebo for ≥ 2 years reported at least one AE; one (20%) of five reported AEs of grade ≥ 3 . Figure 5b shows the common AEs reported after 2 years by patients who received treatment for ≥ 2 years.

Twenty-three of the 32 patients who received olaparib for ≥ 2 years reported AEs after 2 years (72%), with 8 (25%) of 32 reporting AEs of grade ≥ 3 . Four of the five patients who received placebo for ≥ 2 years reported AEs after 2 years; none reported AEs of grade ≥ 3 . Fifteen of the 32 patients who received olaparib for ≥ 2 years had dose reductions (47%), eight of whom (25%) had dose reductions owing to AEs. One patient from the placebo arm (20% of five) had dose reductions, which were not related to AEs. Three of the patients who received olaparib for ≥ 2 years discontinued treatment owing to AEs, which were: pharyngitis and pancytopenia (two AEs in one patient); squamous cell carcinoma of the oral cavity and bronchiectasis (each in one patient). None of the patients who received placebo for ≥ 2 years discontinued owing to AEs.

In the overall study population, the most common grade ≥ 3 AEs were fatigue (olaparib: 11/136 patients [8%]; placebo: 4/128 [3%]) and anaemia (olaparib: 8/136 patients [6%]; placebo: 1/128 [1%]). Overall, 59 (43%) of 136 patients from the olaparib arm and 29 (23%) of 128 from the placebo arm had dose reductions. Dose reductions owing to AEs were reported in 34/136 (25%) and 5/128 (4%) patients from the olaparib and placebo arms, respectively. AEs that led to discontinuation of treatment were reported for eight (6%) of 136 patients from the olaparib arm and two (2%) of 128 patients from the placebo arm; all of these AEs were considered to be related to treatment. For the olaparib arm, in addition to the AEs that led to late discontinuation after 2 years of treatment, the other AEs resulting in discontinuation were: palpitations and myalgia (two AEs in one patient); herpes zoster, nausea, erythematous rash and haemorrhagic stroke (each in one patient). In the placebo arm, the AEs resulting in discontinuation of treatment were pruritic rash and nausea (each in one patient). Thirty (22%) of 136 patients and 11 (9%) of 128 patients reported serious AEs in the olaparib and placebo arms, respectively. There were no additional reports of AEs resulting in death at the 2015 DCO compared with the 2012 DCO, at which one patient had died solely from AEs (haemorrhagic stroke and thrombocytopenia, considered to be treatment-related). Overall, 202 patients in the safety analysis set had died at the 2015 DCO (olaparib arm: n=94; placebo arm: n=108). In the olaparib arm, 83 patients died only from the disease under investigation; one patient had AEs leading to death (haemorrhagic stroke and

thrombocytopenia); one patient died from a combination of their underlying disease and an AE (myelodysplastic syndrome); and nine patients died from other reasons (cardiac failure [n=1], euthanasia [n=1], septic shock [n=1], cerebrovascular disorder [n=1], cerebral haemorrhage [n=1] or unknown [n=4]). In the placebo arm, 99 patients died only from the disease under investigation and nine patients died from other reasons (acute renal failure and pneumonia [n=1], pulmonary embolism [n=1], cardiopulmonary failure [n=1], septic shock due to faecaloma [n=1], ovarian cancer [n=1] or unknown [n=4]). In total, three cases of myelodysplastic syndromes/acute myeloid leukaemia (MDS/AML; two in the olaparib arm and one in the placebo arm) have been reported. All three of the patients who reported MDS/AML had received two prior lines of chemotherapy. Two of these patients had received olaparib maintenance monotherapy for 57 and 10 months, respectively, and one had received placebo for 44 months.

Discussion

These updated descriptive OS analyses indicate an OS advantage for patients with PSR SOC who received olaparib maintenance monotherapy compared with placebo in Study 19. The OS data support the previously published results from Study 19, which showed that PFS, TFST and TSST are significantly prolonged with olaparib, particularly in *BRCAM* patients.^{6,7} Although a statistically significant improvement in OS was not demonstrated, we observed that olaparib had a beneficial treatment effect on OS in the overall population (HR=0.73). This was primarily driven by the treatment effect in the *BRCAM* subgroup, who received the greatest OS benefit from olaparib (HR=0.62). The Kaplan-Meier survival curves for the two treatment arms begin to separate from approximately 42 months after randomisation for the overall study population (Figure 2a). This observation may be a consequence of the heterogeneous nature of the overall population and the different treatment effect in *BRCAM* and *BRCAWt* patients. As *BRCAM* patients receive the most benefit from olaparib and have a better prognosis than *BRCAWt* patients, the proportion of *BRCAM* to *BRCAWt* patients at risk increases over time. At the tail end of the survival curve for the overall population, there are therefore relatively fewer *BRCAWt* patients at risk and so the treatment effect in *BRCAM* patients is less diluted, resulting in the observed separation. The separation of the survival curves at the tail end also suggests the observed OS advantage was influenced by a group of patients who received long-term olaparib maintenance monotherapy. Biological factors that may predict these long-term responders are being investigated.¹⁸

For the *BRCAM* subgroup, early separation of the Kaplan-Meier survival curves is evident, with maximal separation from a time point of approximately 48 months (Figure 2b). Mutations in *BRCA1* and *BRCA2* are the best characterised predictors of HRR deficiency in ovarian cancer. Our data support the proposed mechanism of action of olaparib as a synthetic lethality-inducing agent in the context of tumours with HRR deficiencies, such as *BRCAM* tumours. Ongoing translational analyses from Study 19 support the hypothesis that tumours with *sBRCAM* and *gBRCAM* are similar, both biologically and in sensitivity to olaparib.¹⁶ The OS data for *sBRCAM* patients were not inconsistent with those for *gBRCAM* patients, but the small size of the *sBRCAM* subgroup (n=20) limits the interpretation of our findings.

An exploratory restricted means analysis, using a pseudovalue methodology, showed a difference in restricted mean OS with olaparib compared with placebo of 5.2 months in the overall population and 7.5 months in the *BRCAm* subgroup. Two other methodologies were investigated for the restricted means analysis (Supplementary Material, page 3) and all analyses gave comparable results, indicating an OS advantage with maintenance olaparib versus placebo, with a greater treatment effect in the *BRCAm* subgroup. The difference in median OS with maintenance olaparib compared with placebo was 2.0 months in the overall population and 4.7 months in the *BRCAm* subgroup. This is less than the difference in restricted mean OS; the mean offers an estimate of average life expectancy, which takes account of patients who do very well on treatment, whereas the median provides a more conservative estimate that is limited to the first half of the survival observations. For example, in the *BRCAm* subgroup who received olaparib, the median OS indicates that 50% of patients lived for longer than 34.9 months, but the mean survival time was 44.3 months.

For the *BRCAwt* subgroup, some patients may have been HRR-deficient as a result of alternative factors, such as mutations in genes that encode other proteins involved in the HRR pathway, or epigenetic mechanisms, which do not yet have well-defined clinical testing strategies.^{19,20} Some separation is seen at the tail end of the *BRCAwt* survival curves for the two treatment groups (Figure 2c), suggesting that there may be a further subset of patients who receive benefit from olaparib treatment. Investigations into patients who were *BRCAwt* but deficient in other HRR genes are ongoing.²¹

Study 19 was designed to demonstrate a statistically significant difference in PFS in the patients who were randomised: a population enriched for HRR tumours as a result of high-grade serous histology and platinum sensitivity. No rules were pre-specified to control the Type 1 error rate for subgroups. The study was not designed to show a statistically significant difference in OS. However, a multiplicity strategy was pre-specified to control the error rate at 5% (two-sided) for multiple analyses of OS. Two previous OS analyses have been conducted, which did not meet statistical significance, and only 0.95% alpha (two-sided) was available to test at this updated analysis. The *P*-values did not meet this criterion for statistical significance

($P < 0.0095$) and therefore the favourable treatment effect observed for OS should only be considered descriptive and should be interpreted in the context of the clinically meaningful and statistically significant improvement in PFS. All P -values for OS are deemed nominal. The interpretation of the exploratory restricted means data is limited by the *post-hoc* nature of this analysis, as it was not pre-specified.

The updated analyses for TFST and TSST show a significant improvement in these exploratory endpoints with olaparib in the overall study population, *BRCAm* and *BRCAt* subgroups, consistent with the previous analysis.⁷ TFST is an exploratory endpoint but is clinically meaningful, as it represents the time that women are free from the next line of treatment. The updated TFST data provide a long-term view on efficacy, with the TFST Kaplan-Meier curves for the two treatment arms remaining clearly separated at a time point over 5 years from randomisation. As patients remain blinded to study treatment beyond progression, these data support an extended benefit, beyond PFS, for patients with PSR SOC receiving olaparib maintenance monotherapy. It can be challenging to understand the full therapeutic value of investigational treatments in ovarian cancer studies, as there is often a long follow-up for OS and analyses can be confounded by post-discontinuation therapy. Improvement in TSST can demonstrate continued benefit, beyond the next line of therapy, and this intermediate endpoint can therefore support other efficacy endpoints when evaluating the long-term impact of investigational treatments.²²

Crossover was not allowed in this study, but 17 patients from the placebo arm (14 *BRCAm* patients) had received post-discontinuation PARPi treatment by the 2015 DCO, via other clinical studies. This is considered to have had the potential to confound the OS data: an exploratory analysis has previously been reported for the *BRCAm* subgroup, excluding all patients from sites where at least one patient from the placebo arm received post-discontinuation PARPi therapy, and this showed a greater treatment effect than the previously published OS analysis.^{7,23}

Notably, at the DCO in 2015, there were 15 patients continuing on olaparib and one on placebo. Significant long-term exposure to maintenance olaparib was observed, with 18/136 (13%) of all patients (11/74 [15%] of *BRCAm* patients) receiving olaparib for ≥ 5 years. This observation supports the long-term benefit and tolerability of olaparib. Similar data for long-term treatment have not previously been seen in

clinical trials in recurrent ovarian cancer. Baseline data show that the majority of the 19 patients who received study treatment for ≥ 5 years had two or three prior lines of chemotherapy and a platinum-free interval >12 months.

Since the previous safety analysis, there has been an additional 3 years of follow-up, during which time no new safety signals were reported for the patients remaining on treatment and there was no change to the overall safety profile. For patients who received olaparib for ≥ 2 years, the most frequent AEs were not different to those in the overall population, specifically low-grade nausea, fatigue, constipation and vomiting, which are manageable and were generally reported prior to 2 years on treatment.⁶⁻⁸ These long-term safety findings are consistent with previous data from Study 19 and other clinical olaparib monotherapy studies. As reported in 2012, a low proportion of patients experienced AEs resulting in discontinuation of treatment.⁷

To conclude, an OS advantage is seen for patients with *BRCAm* PSR SOC treated with olaparib as maintenance therapy in Study 19. This observation is consistent with data showing a significant improvement in PFS and in the intermediate endpoints TFST and TSST with olaparib. Additionally, 11 (15%) of 74 patients with *BRCAm* continued on olaparib for ≥ 5 years, highlighting that this PARPi can significantly alter the disease course. Ongoing analyses are investigating the potential benefit of olaparib for patients who are *BRCAw*t but have other HRR deficiencies, some of whom may continue on olaparib without progression for several years. There is an extensive Phase III clinical programme for olaparib, with the SOLO2 study (NCT01874353) assessing maintenance olaparib treatment (tablets) in *BRCAm* patients with PSR SOC who have received at least two prior lines of platinum-based chemotherapy.

Research in context

Evidence before this study

We conducted searches of PubMed and the databases of the American Society of Clinical Oncology, European Society for Medical Oncology, Society of Gynecological Oncology and European Society of Gynaecological Oncology between 1 March 2015 and 1 March 2016 to identify journal publications and meeting abstracts that included the search terms “poly(ADP-ribose) polymerase inhibitor” or “PARP inhibitor” and “ovarian cancer”. No language restrictions were used. Olaparib is an oral PARP inhibitor (PARPi) that has shown significant clinical activity and tolerability in patients with recurrent ovarian cancer and is approved in the EU and the US for the treatment of *BRCA1/2*-mutated advanced ovarian cancer. Other PARPis in clinical development include rucaparib, veliparib, niraparib and talazoparib. There have been no reports of an advantage in overall survival for ovarian cancer patients treated with maintenance therapy with a PARPi compared with placebo.

Added value of this study

This is the third overall survival analysis for a Phase II, randomised trial of olaparib maintenance monotherapy in patients with platinum-sensitive recurrent serous ovarian cancer (PSR SOC) and is, to our knowledge, the first analysis to indicate a survival advantage for patients with ovarian cancer treated with a PARPi. The survival analysis was conducted after more than 5 years total follow-up, at high data maturity, with an additional 3 years of follow-up since the previous analysis. The observed survival advantage may have primarily been driven by a survival benefit in the subgroup of patients with *BRCA1/2* mutations (*BRCAm*). We believe this is also the first report of significant long-term exposure to a PARPi in recurrent ovarian cancer, with 18 (13%) of 136 patients receiving maintenance olaparib for ≥ 5 years. No new safety signals were observed and the long-term safety data were consistent with the known safety profile for olaparib monotherapy.

Implications of all of the available evidence

We have previously reported data from this Phase II study that showed a significant improvement in progression-free survival (PFS) with maintenance olaparib, with the

greatest benefit seen in patients with a *BRC*Am. Exploratory analyses have also shown a significant improvement in time to first and second subsequent therapy or death (TFST and TSST) with olaparib compared with placebo. To our knowledge, this is the first analysis to show survival data in recurrent *BRC*Am ovarian cancer that are consistent with previously reported benefits in PFS, TFST and TSST. Taken together, the available data support the long-term benefit and tolerability of maintenance olaparib in *BRC*Am patients with PSR SOC.

Contributors

JAL was responsible for the study design. JAL, PH, CG, MF, IV, GR, CS, WM, RS-F, TS, DM and UM obtained the data. AF, SS, PR, EL, DH and MAS analysed the data. All authors interpreted the data and reviewed draft and final versions of the manuscript.

Declaration of interests

JAL has participated in advisory boards and lecture symposia, and has received institutional and personal fees from AstraZeneca, personal fees from Roche and Pfizer, and institutional fees from Clovis Oncology and Merck. PH has participated in advisory boards for AstraZeneca and Roche. CG has received grants and personal fees from AstraZeneca and GlaxoSmithKline, personal fees from Roche, Nucana and Clovis Oncology, grants from Aprea and Novartis and has five patents broadly relevant to this work (issued: PCT/US2012/040805; pending: PCT/GB2013/053202, 1409479.1, 1409476.7 and 1409478.3). MF has participated in advisory boards for AstraZeneca and received personal fees from AstraZeneca, Roche and Pfizer. IV has participated in advisory boards for AstraZeneca. GR has participated in advisory boards for Oxigene and Amgen. CS has received personal fees from AstraZeneca and travel support from AstraZeneca. DM has participated as a consultant for AstraZeneca and has received research support from AstraZeneca. AF, SS, EL and DH are employees of AstraZeneca and own stock. PR and MAS are employees of AstraZeneca. UM has provided both paid and unpaid consulting to AstraZeneca. All other authors have no disclosures.

Acknowledgements

We thank all patients, their families and our co-investigators. This study was sponsored by AstraZeneca. Medical writing support during the development of this manuscript was provided by Rachel Patel, MBiochem, from Mudskipper Business Ltd and was funded by AstraZeneca.

References

1. Ferlay J, Steliarova-Foucher E, Lortet-Tieulent J, et al. Cancer incidence and mortality patterns in Europe: estimates for 40 countries in 2012. *Eur J Cancer* 2013; 49: 1374-403.
2. World Health Organization. GLOBOCAN statistics. 2012. Available at: <http://globocan.iarc.fr/> (accessed Jul 8, 2016).
3. Ledermann JA, Raja FA, Fotopoulou C, Gonzalez-Martin A, Colombo N, Sessa C. Newly diagnosed and relapsed epithelial ovarian carcinoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2013; 24(Suppl 6): vi24-vi32.
4. Kyrgiou M, Salanti G, Pavlidis N, Paraskevidis E, Ioannidis JP. Survival benefits with diverse chemotherapy regimens for ovarian cancer: meta-analysis of multiple treatments. *J Natl Cancer Inst* 2006; 98: 1655-63.
5. de Angelis R, Sant M, Coleman MP, et al. Cancer survival in Europe 1999-2007 by country and age: results of EUROCARE--5-a population-based study. *Lancet Oncol* 2014; 15: 23-34.
6. Ledermann J, Harter P, Gourley C, et al. Olaparib maintenance therapy in platinum-sensitive relapsed ovarian cancer. *N Engl J Med* 2012; 366: 1382-92.
7. Ledermann J, Harter P, Gourley C, et al. Olaparib maintenance therapy in patients with platinum-sensitive relapsed serous ovarian cancer: a preplanned retrospective analysis of outcomes by BRCA status in a randomised phase 2 trial. *Lancet Oncol* 2014; 15: 852-61.
8. Kaufman B, Shapira-Frommer R, Schmutzler RK, et al. Olaparib monotherapy in patients with advanced cancer and a germ-line *BRCA1/2* mutation. *J Clin Oncol* 2015; 33: 244-50.
9. Murai J, Huang SY, Das BB, et al. Trapping of PARP1 and PARP2 by clinical PARP inhibitors. *Cancer Res* 2012; 72: 5588-99.
10. Farmer H, McCabe N, Lord CJ, et al. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature* 2005; 434: 917-21.
11. Bryant HE, Schultz N, Thomas HD, et al. Specific killing of BRCA2-deficient tumours with inhibitors of poly(ADP-ribose) polymerase. *Nature* 2005; 434: 913-17.
12. EMA. Lynparza recommended for approval in ovarian cancer. 2014. Available at: http://www.ema.europa.eu/ema/index.jsp?curl=pages/news_and_events/new

- s/2014/10/news_detail_002196.jsp&mid=WC0b01ac058004d5c1 (accessed Jul 8, 2016).
13. FDA. Olaparib. 2014. Available at: <http://www.fda.gov/Drugs/InformationOnDrugs/ApprovedDrugs/ucm427598.htm> (accessed Jul 8, 2016).
 14. AstraZeneca. Global Policy: Bioethics. 2015. Available at: <https://www.astrazeneca.com/sustainability/responsible-research.html> (accessed Jul 8, 2016).
 15. Anderson PK, Hansen MG, Klein JP. Regression analysis of restricted mean survival time based on pseudo-observations. *Lifetime data anal* 2004; 10: 335-50.
 16. Dougherty B, Lai Z, Ledermann JA, et al. Exploratory analyses suggest ovarian tumors with somatic or germline loss of function mutations in *BRCA1* or *BRCA2* are biologically similar and sensitive to PARP inhibition. Annual Meeting of the American Association for Cancer Research. *Philadelphia, PA, USA, 18-22 April 2015*; abst 611.
 17. Grambsch P, Therneau T. Proportional hazards tests and diagnostics based on weighted residuals. *Biometrika* 1994; 81: 515-26.
 18. Lheureux S, Ledermann JA, Runswick S, et al. Genomic characterization of long-term responders to olaparib. *J Clin Oncol* 2015; 33: abst 5566.
 19. McCabe N, Turner NC, Lord CJ, et al. Deficiency in the repair of DNA damage by homologous recombination and sensitivity to poly(ADP-ribose) polymerase inhibition. *Cancer Res* 2006; 66: 8109-15.
 20. Press JZ, De Luca A, Boyd N, et al. Ovarian carcinomas with genetic and epigenetic *BRCA1* loss have distinct molecular abnormalities. *BMC Cancer* 2008; 8: 17.
 21. Hodgson DR, Dougherty B, Lai Z, et al. Candidate biomarkers of PARP inhibitor sensitivity in ovarian cancer beyond the *BRCA* genes. European Cancer Congress 2015. *Vienna, Austria, 25-29 September 2015*; abst 435.
 22. Matulonis UA, Oza AM, Ho TW, Ledermann JA. Intermediate clinical endpoints: a bridge between progression-free survival and overall survival in ovarian cancer trials. *Cancer* 2015; 121: 1737-46.
 23. Matulonis U, Harter P, Gourley C, et al. Olaparib maintenance therapy in patients with platinum-sensitive relapsed serous ovarian cancer and a *BRCA* mutation: overall survival adjusted for post-progression PARP inhibitor therapy. *Cancer* 2016; 122: 1844-52.

Figure legends

Figure 1. Enrolment, randomisation and treatment status at the third analysis of OS in Study 19.

Footnote: DCO was on 30 September 2015. *One patient was randomly assigned to the placebo arm, but withdrew consent and withdrew from the study without receiving treatment

Figure 2. OS in all patients and according to *BRCAM* status. **a)** All patients (n=265); **b)** *BRCAM* patients (n=136); **c)** *BRCAwT* patients (n=118).

Figure 3. Summary of the Cox proportional hazards analysis of OS in the overall study population and different *BRCAM* subgroups.

Figure 4. TFST and TSST in all patients and according to *BRCAM* status. **a)** TFST in all patients (n=264); **b)** TFST in *BRCAM* patients (n=136); **c)** TFST in *BRCAwT* patients (n=118); **d)** TSST in all patients (n=264); **e)** TSST in *BRCAM* patients (n=136); **f)** TSST in *BRCAwT* patients (n=118).

Figure 5. Common AEs of all grades and grade ≥ 3 in patients who received study treatment for ≥ 2 years (olaparib n=32; placebo n=5). **a)** AEs reported from the start of treatment to the 2015 DCO*

Footnote: *AEs that were reported in ≥ 8 patients are presented. Additional data for AEs in patients who received study treatment for ≥ 2 years are listed in the Supplementary Material (page 4–5)

b) AEs reported after 2 years*

Footnote: *AEs that were reported in ≥ 4 patients are presented. Additional data for AEs reported after 2 years by patients who received study treatment for ≥ 2 years are listed in the Supplementary Material (page 6–7)

Fig 1







Fig 2





Number of patients at risk:

Olaparib	74	69	65	56	50	39	33	27	27	27	25	23	11	1	0
Placebo	62	58	52	40	34	29	25	20	19	15	13	10	6	0	0



Number of patients at risk:

Olaparib	57	57	50	39	28	23	19	16	15	14	12	12	4	1	0
Placebo	61	60	56	46	37	25	17	15	12	12	11	8	3	1	0

Fig 3







Fig 4






Number of patients at risk:

Olaparib	74	61	43	30	26	25	21	20	20	19	17	16	8	1	0
Placebo	62	33	16	9	7	4	4	4	2	2	2	2	2	0	0




Number of patients at risk:

Olaparib	57	42	28	15	13	10	10	9	8	8	8	8	3	1	0
Placebo	61	37	13	6	3	1	1	1	0	0	0	0	0	0	0




Number of patients at risk:

Olaparib	136	128	96	66	48	43	38	35	30	28	27	26	12	2	0
Placebo	128	115	84	49	27	15	10	9	9	6	5	4	3	0	0



Number of patients at risk:


Olaparib	74	69	56	40	32	29	25	23	21	20	19	18	9	1	0
Placebo	62	57	42	25	18	9	6	5	5	3	3	3	3	0	0




Number of patients at risk:


Olaparib	57	56	38	25	16	14	13	12	9	8	8	8	3	1	0
Placebo	61	55	40	24	9	6	4	4	4	3	2	1	0	0	0

Fig 5






Suppl Fig 1



Number of patients at risk:

	0	6	12	18	24	30	36	42	48	54	60	66	72	78	84
Olaparib	48	44	42	38	32	24	20	14	14	14	14	13	6	1	0
Placebo	45	42	38	28	23	21	17	14	13	12	10	8	5	0	0



Number of patients at risk:

Olaparib	26	25	23	18	18	15	13	13	13	11	10	5	0	0	
Placebo	17	16	14	12	11	8	8	6	6	3	3	2	1	0	0

Table 1. Patient demographics and baseline characteristics*

	Overall study population (N=265)		<i>BRC</i> Am patients (n=136) [†]		<i>BRC</i> Awt [‡] patients (n=118) [‡]	
	Olaparib (N=136)	Placebo (N=129)	Olaparib (n=74)	Placebo (n=62)	Olaparib (n=57)	Placebo (n=61)
Age, years	58·0 (21–89)	59·0 (33–84)	57·5 (38–89)	55·0 (33–84)	62·0 (21–80)	63·0 (49–79)
Ancestry[§]						
Non-Jewish	115 (85%)	112 (87%)	60 (81%)	48 (77%)	51 (89%)	58 (95%)
Jewish	21 (15%)	17 (13%)	14 (19%)	14 (23%)	6 (11%)	3 (5%)
Number of prior lines of chemotherapy						
2	59 (43%)	63 (49%)	26 (35%)	28 (45%)	32 (56%)	35 (57%)
3	43 (32%)	34 (26%)	28 (38%)	18 (29%)	14 (25%)	14 (23%)
4	18 (13%)	19 (15%)	9 (12%)	10 (16%)	6 (11%)	9 (15%)
≥5	16 (12%)	13 (10%)	11 (15%)	6 (10%)	5 (8%)	3 (5%)
Primary tumour location						
Ovary	119 (88%)	109 (84%)	65 (88%)	54 (87%)	50 (88%)	49 (80%)
Fallopian tube or primary peritoneal	17 (13%)	20 (16%)	9 (12%)	8 (13%)	7 (12%)	12 (20%)
Time to progression after completion of penultimate platinum-based regimen						

>6 to ≤12 months	53 (39%)	54 (42%)	28 (38%)	26 (42%)	23 (40%)	24 (39%)
>12 months	83 (61%)	75 (58%)	46 (62%)	36 (58%)	34 (60%)	37 (61%)

Objective response to most recent platinum-based regimen

Complete response	57 (42%)	63 (49%)	36 (49%)	34 (55%)	20 (35%)	25 (41%)
Partial response	79 (58%)	66 (51%)	38 (51%)	28 (45%)	37 (65%)	36 (59%)

Data are median (range) or number (%). *These baseline data have also been described previously.^{6,7} †Data were not available for all randomised patients. ‡The *BRCAwt* subgroup included patients without detected *BRCAm* and patients with *BRCAm* of unknown significance. §Ancestry was self-reported

Table 2. OS using a Cox proportional hazards analysis, a restricted means analysis and a log-rank test

	Overall study population (N=265)		BRCAm patients (n=136)	
	Olaparib (N=136)	Placebo (N=129)	Olaparib (n=74)	Placebo (n=62)
Median OS (95% CI), months	29.8 (26.9–35.7)	27.8 (24.9–33.7)	34.9 (29.2–54.6)	30.2 (23.1–40.7)
Difference in median OS, months		2.0		4.7
Cox proportional hazards analysis: hazard ratio (95% CI)		0.73 (0.55–0.96)		0.62 (0.41–0.94)
Log-rank test: hazard ratio (95% CI)		0.72 (0.54–0.95)		0.61 (0.39–0.95)
Restricted mean OS, months	40.1	34.9	44.3	36.9
Difference in restricted mean OS (95% CI), months		5.2 (-0.8–11.2)		7.4 (-1.1–16.0)

Table 3. Patients receiving long-term olaparib maintenance monotherapy, by number of years of treatment received

Duration of treatment	Overall study population (N=264)		<i>BRCAm</i> patients (n=136)	
	Olaparib (N=136)	Placebo (N=128)	Olaparib (n=74)	Placebo (n=62)
≥1 year	54 (40%)	14 (11%)	34 (46%)	8 (13%)
≥2 years	32 (24%)	5 (4%)	21 (28%)	5 (8%)
≥3 years	24 (18%)	3 (2%)	16 (22%)	3 (5%)
≥4 years	20 (15%)	1 (1%)	12 (16%)	1 (2%)
≥5 years	18 (13%)	1 (1%)	11 (15%)	1 (2%)
≥6 years	7 (5%)	1 (1%)	4 (5%)	1 (2%)

Table 4. Baseline characteristics for patients receiving study treatment for ≥5 years

	Overall study population (n=19)		BRCAm patients (n=12)		BRCAwt patients (n=7)	
	Olaparib (n=18)	Placebo (n=1)	Olaparib (n=11)	Placebo (n=1)	Olaparib (n=7)	Placebo (n=0)
Number of prior lines of chemotherapy						
2	7 (39%)	1 (100%)	4 (36%)	1 (100%)	3 (43%)	0
3	7 (39%)	0	4 (36%)	0	3 (43%)	0
4	2 (11%)	0	2 (18%)	0	0	0
≥5	2 (11%)	0	1 (9%)	0	1 (14%)	0
Platinum-free interval						
6–12 months	5 (28%)	0	2 (18%)	0	3 (43%)	0
>12 months	13 (72%)	1 (100%)	9 (82%)	1 (100%)	4 (57%)	0
Objective response to most recent platinum-based regimen						
Complete response	10 (56%)	1 (100%)	5 (45%)	1 (100%)	5 (71%)	0
Partial response	8 (44%)	0	6 (55%)	0	2 (29%)	0


Overall survival in patients with platinum-sensitive recurrent serous ovarian cancer receiving olaparib maintenance monotherapy: An updated analysis from a Phase II, randomised, double-blind, placebo-controlled trial

Jonathan A Ledermann,¹ Philipp Harter,² Charlie Gourley,³ Michael Friedlander,⁴ Ignace Vergote,⁵ Gordon Rustin,⁶ Clare Scott,⁷ Werner Meier,⁸ Ronnie Shapira-Frommer,⁹ Tamar Safra,¹⁰ Daniela Matei,¹¹ Anitra Fielding,¹² Stuart Spencer,¹² Philip Rowe,¹² Elizabeth Lowe,¹³ Darren Hodgson,¹² Mika Sovak¹³ and Ursula Matulonis¹⁴

¹*UCL Cancer Institute, University College London and UCL Hospitals, London, UK;* ²*Kliniken Essen Mitte, Essen, Germany;* ³*University of Edinburgh Cancer Research UK Centre, Medical Research Council Institute of Genetics and Molecular Medicine, Western General Hospital, Edinburgh, UK;* ⁴*University of New South Wales Clinical School, Prince of Wales Hospital, Randwick, NSW, Australia;* ⁵*University of Leuven, Leuven, Belgium;* ⁶*Mount Vernon Hospital, Northwood, UK;* ⁷*Royal Melbourne Hospital, Parkville, VIC, Australia;* ⁸*Evangelisches Krankenhaus, Düsseldorf, Germany;* ⁹*Chaim Sheba Medical Center, Tel Hashomer, Israel;* ¹⁰*Tel Aviv Sourasky Medical Center, Tel Aviv, Israel;* ¹¹*Northwestern University Feinberg School of Medicine, Chicago, IL, USA;* ¹²*AstraZeneca, Macclesfield, UK;* ¹³*AstraZeneca, Gaithersburg, MD, USA;* ¹⁴*Dana-Farber Cancer Institute, Boston, MA, USA*

Figure 1. OS according to *BRCAm* status

a) *BRCA1m* patients (n=93)



b) *BRCA2m* patients (n=43)




Table 1. Restricted mean OS using three methodologies: pseudovalues, flexible parametric model and area under the Kaplan–Meier curves¹

		Overall study population (N=265)		BRCAm patients (n=136)	
		Olaparib (N=136)	Placebo (N=129)	Olaparib (n=74)	Placebo (n=62)
Pseudovalues	Restricted mean OS, months	40.1	34.9	44.3	36.9
	Difference in restricted mean OS (95% CI), months	5.2 (-0.8–11.2)		7.4 (-1.1–16.0)	
Flexible parametric model	Restricted mean OS, months	40.7	34.4	44.6	36.6
	Difference in restricted mean OS (95% CI), months	6.3 (0.3–12.1)		8.0 (-0.3–16.3)	
Area under Kaplan-Meier curves	Restricted mean OS, months	40.1	35.0	44.4	36.9
	Difference in restricted mean OS (95% CI), months	5.1 (-0.8–11.2)		7.5 (-1.1–16.0)	

Table 2. AEs of all grades (frequency >6 patients) and of grade ≥ 3 (all instances) in either treatment arm in patients who received study treatment for ≥ 2 years*

	Overall study population (n=37)				BRCAm patients (n=26)			
	All grades		Grade ≥ 3		All grades		Grade ≥ 3	
	Olaparib (n=32)	Placebo (n=5)	Olaparib (n=32)	Placebo (n=5)	Olaparib (n=21)	Placebo (n=5)	Olaparib (n=21)	Placebo (n=5)
Any AE	30 (94%)	5 (100%)	15 (47%)	1 (20%)	20 (95%)	5 (100%)	9 (43%)	1 (20%)
Nausea	24 (75%)	2 (40%)	0	0	18 (86%)	2 (40%)	0	0
Fatigue	18 (56%)	2 (40%)	3 (9%)	0	12 (57%)	2 (40%)	2 (10%)	0
Vomiting	12 (38%)	0	0	0	9 (43%)	0	0	0
Constipation	12 (38%)	1 (20%)	0	0	8 (38%)	1 (20%)	0	0
Abdominal pain	11 (34%)	1 (20%)	0	0	8 (38%)	1 (20%)	0	0
Diarrhoea	11 (34%)	1 (20%)	0	0	7 (33%)	1 (20%)	0	0
Arthralgia	11 (34%)	1 (20%)	1 (3%)	0	5 (24%)	1 (20%)	1 (5%)	0
Dizziness	10 (31%)	0	0	0	7 (33%)	0	0	0
Headache	9 (28%)	1 (20%)	0	0	7 (33%)	1 (20%)	0	0
Upper respiratory tract infection	9 (28%)	2 (40%)	0	0	6 (29%)	2 (40%)	0	0
Decreased appetite	9 (28%)	0	0	0	6 (29%)	0	0	0
Back pain	9 (28%)	2 (40%)	1 (3%)	0	6 (29%)	2 (40%)	1 (5%)	0
Pain in extremity	9 (28%)	1 (20%)	0	0	4 (19%)	1 (20%)	0	0
Anaemia	8 (25%)	0	1 (3%)	0	8 (38%)	0	1 (5%)	0
Insomnia	8 (25%)	1 (20%)	0	0	7 (33%)	1 (20%)	0	0
Pruritus	8 (25%)	1 (20%)	1 (3%)	0	6 (29%)	1 (20%)	0	0
Dysgeusia	7 (22%)	0	0	0	5 (24%)	0	0	0
Peripheral neuropathy	7 (22%)	0	0	0	5 (24%)	0	0	0
Cough	7 (22%)	2 (40%)	0	0	5 (24%)	2 (40%)	0	0

Dyspnoea	6 (19%)	1 (20%)	1 (3%)	0	3 (14%)	1 (20%)	1 (5%)	0
Myalgia	3 (9%)	1 (20%)	1 (3%)	0	2 (10%)	1 (20%)	0	0
Hypertension	3 (9%)	0	1 (3%)	0	1 (5%)	0	0	0
Pneumonia	2 (6%)	0	2 (6%)	0	2 (10%)	0	2 (10%)	0
Femur fracture	2 (6%)	0	2 (6%)	0	2 (10%)	0	2 (10%)	0
Pancytopenia	2 (6%)	0	1 (3%)	0	2 (10%)	0	1 (5%)	0
Osteoarthritis	2 (6%)	0	1 (3%)	0	2 (10%)	0	1 (5%)	0
Weight decreased	2 (6%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Anaemia megaloblastic	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Hernia pain	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Hip fracture	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Acute leukaemia	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Intraductal proliferative breast lesion	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Squamous cell carcinoma of the oral cavity	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Syncope	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Pulmonary embolism	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Neutropenia	1 (3%)	1 (20%)	1 (3%)	1 (20%)	0	1 (20%)	0	1 (20%)
Small intestinal obstruction	1 (3%)	0	1 (3%)	0	0	0	0	0
Iodine allergy	1 (3%)	0	1 (3%)	0	0	0	0	0
Blood creatine phosphokinase increased	1 (3%)	0	1 (3%)	0	0	0	0	0
Dysaesthesia	1 (3%)	0	1 (3%)	0	0	0	0	0
Confusional state	1 (3%)	0	1 (3%)	0	0	0	0	0
Urticaria	1 (3%)	0	1 (3%)	0	0	0	0	0
Myelodysplastic syndrome	0	1 (20%)	0	1 (20%)	0	1 (20%)	0	1 (20%)

*AEs are included that met the criteria stated in either treatment arm in patients who received study treatment for ≥ 2 years in the overall study population or the *BRCAM* subgroup

Table 3. AEs of all grades (frequency ≥ 3 patients) and of grade ≥ 3 (all instances) that were reported after 2 years in either treatment arm by patients who received study treatment for ≥ 2 years*

	Overall study population (n=37)				BRCAm patients (n=26)			
	All grades		Grade ≥ 3		All grades		Grade ≥ 3	
	Olaparib (n=32)	Placebo (n=5)	Olaparib (n=32)	Placebo (n=5)	Olaparib (n=21)	Placebo (n=5)	Olaparib (n=21)	Placebo (n=5)
Any AE	23 (72%)	4 (80%)	8 (25%)	0	15 (71%)	4 (80%)	6 (29%)	0
Pruritus	5 (16%)	0	1 (3%)	0	4 (19%)	0	0	0
Constipation	5 (16%)	0	0	0	4 (19%)	0	0	0
Urinary tract infection	5 (16%)	0	0	0	4 (19%)	0	0	0
Dyspnoea	5 (16%)	0	1 (3%)	0	3 (14%)	0	1 (5%)	0
Fatigue	5 (16%)	0	0	0	3 (14%)	0	0	0
Dizziness	5 (16%)	0	0	0	3 (14%)	0	0	0
Nausea	4 (13%)	0	0	0	3 (14%)	0	0	0
Pain in extremity	4 (13%)	0	0	0	2 (10%)	0	0	0
Back pain	3 (9%)	0	1 (3%)	0	3 (14%)	0	1 (5%)	0
Bone pain	3 (9%)	0	0	0	3 (14%)	0	0	0
Headache	3 (9%)	1 (20%)	0	0	2 (10%)	1 (20%)	0	0
Peripheral swelling	3 (9%)	0	0	0	2 (10%)	0	0	0
Upper respiratory tract infection	3 (9%)	0	0	0	2 (10%)	0	0	0
Cough	3 (9%)	0	0	0	2 (10%)	0	0	0
Vomiting	3 (9%)	0	0	0	1 (5%)	0	0	0
Diarrhoea	3 (9%)	0	0	0	1 (5%)	0	0	0
Sinusitis	3 (9%)	0	0	0	1 (5%)	0	0	0
Anaemia	2 (6%)	0	1 (3%)	0	2 (10%)	0	1 (5%)	0

Pancytopenia	2 (6%)	0	1 (3%)	0	2 (10%)	0	1 (5%)	0
Osteoarthritis	2 (6%)	0	1 (3%)	0	2 (10%)	0	1 (5%)	0
Arthralgia	2 (6%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Femur fracture	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Hip fracture	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Acute leukaemia	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Squamous cell carcinoma of the oral cavity	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Pulmonary embolism	1 (3%)	0	1 (3%)	0	1 (5%)	0	1 (5%)	0
Confusional state	1 (3%)	0	1 (3%)	0	0	0	0	0
Urticaria	1 (3%)	0	1 (3%)	0	0	0	0	0

*AEs that were reported after 2 years are included that met the criteria stated in either treatment arm in patients who received study treatment for ≥ 2 years in the overall study population or *BRCAM* subgroup

Table 4. Recruitment data for Study 19

Site number	Principle investigator	Number of patients recruited (N=265)
Dr. Horst Schmidt Klinik, Wiesbaden, Germany	Philipp Harter/Rita Hills/Tanja Neunhöffer	13
Western General Hospital, Edinburgh, UK	Charlie Gourley	12
University of New South Wales, Randwick, NSW, Australia	Michael Friedlander	11
Dana Faber Cancer Institute, Boston, MA, USA	Ursula Matulonis	11
UZ Gasthuisberg Dienst Gynaecologische Oncologie, Leuven, Belgium	Ignace Vergote	10
St Georges' Hospital, London, UK	Gordon Rustin	9
Massachusetts General Hospital, Boston, MA, USA	Ursula Matulonis	9
Royal Melbourne Hospital, Parkville, VIC, Australia	Clare Scott	8
Evangelisches Krankenhaus, Düsseldorf, Germany	Werner Meier	8
The Chaim Sheba Medical Center, Tel Hashomer, Israel	Ronnie Shapira-Frommer	8
Tel Aviv Sourasky Medical Center, Tel Aviv, Israel	Tamar Safra/Larisa Ryvo	6
University College London Hospitals, London, UK	Jonathan Ledermann	6
Indiana University, Indianapolis, IN, USA	Daniela Matei	6
Institutul Oncologic Prof Dr Ion Chiricuta Cluj-Napoca, Napoca, Romania	Carmen Floares	5
Netherlands Cancer Institute, Amsterdam, The Netherlands	Gabe Sonke	5
Orenburg Regional Clinical Oncological Dispensary, Orenburg, Russia	Vadim Shirinkin	5
Mount Vernon Cancer Centre, Northwood, UK	Michelle Ferguson	5
Mercy Hospital for Women, Heideberg, VIC, Australia	Linda Mileskin	4
Centre Léon Bérard, Lyon, France	Isabelle Ray-Coquard	4
Hadassah University Hospital, Jerusalem, Israel	David Edelmann	4
NYU Clinical Cancer Center, New York, NY, USA	Stephanie Blank	4
Cedars-Sinai Medical Centre, Los Angeles, CA, USA	Beth Karlan	4

Monash Medical Centre, Melbourne, VIC, Australia	Geraldine Goss	3
Sunshine Coast Cancer Centre, Nambour, QLD, Australia	Michelle Cronk	3
UZ Brussel Dienst Oncologie, Brussels, Belgium	Jacques De Greve	3
Princess Margaret Cancer Center, Toronto, ON, Canada	Amit Oza	3
Hospital Tenon, Paris, France	Frederic Selle	3
Universitätsklinikum Essen, Essen, Germany	Pauline Wimberger	3
St. Petersburg GUZ City Clinical Oncology Dispensary, St. Petersburg, Russia	Alla Lisanskaya	3
Hospital Vall d'Hebrón, Barcelona, Spain	José María Del Campo	3
Royal Marsden Hospital, London, UK	Stan Kaye	3
Clatterbridge Centre for Oncology, Wirral, UK	Rosemary Lord	3
Derzhavna Ustano va Instytut Medychnoi Radiologii im SP Grygoryeva AMN Ukrainy, Pushkinska, Ukraine	Oksana Tarasova	3
Alta Bates Summit Comprehensive Cancer Center, Berkeley, CA, USA	David Irwin	3
Centre Hospitalier Universitaire de Sherbrooke, Sherbrooke, QC, Canada	Paul Bessette	2
Insitut Bergonié, Bordeaux cedex, France	Anne Floquet	2
Universitätsfrauenklinik Ulm, Ulm, Germany	Wolfgang Janni	2
Georg-August-Universität Göttingen, Göttingen, Germany	Günter Emons	2
Western Galilee Hospital, Nahariya, Israel	Hadassah Goldberg	2
Shaare Zedek Medical Center, Jerusalem, Israel	Ora Rosengarten	2
Rambam Health Care Campus, Haifa, Israel	Amnon Amit	2
Wielkopolskie Centrum Onkologii, Poznan, Poland	Jan Kotarski	2
Wielkopolskie Centrum Onkologii, Poznan Poland	Andrzej Roszak	2
Pomeranian Medical University, Szczecin, Poland	Izabella Rzepka-Gorska	2
Szpital Kliniczny Przemienienia Pańskiego, Poznań, Poland	Janina Markowska	2
GUZ Perm Regional Dispensary, Perm, Russia	Zinaida Akishina	2
Hospital Clinico Universitario de Valencia, Valencia, Spain	Andrés Cervantes	2

Hospital Reina Sofía Avda, Córdoba, Spain	María Jesús Rubio Pérez	2
Christie Hospital, Manchester, UK	Gordon Jayson	2
Kyivska Miska Onkologichna Likarnya, Verhovynnae, Ukraine	Olena Manzhura	2
Natsionalnyi Instytut Raku, Lomonosova, Ukraine	Liusia Vorobyeva	2
Temopilskyi Derzhavnyi Medychnyi Universytet, Ternopil, Ukraine	Igor Galaychuk	2
Uzhgorodskyi Natsionalnyi Medychnyi Universytet, Uzhgorod, Ukraine	Eugen Rusyn	2
Donetskyi Oblasnyi Protypukhlynyi Tsentr, Donetsk, Ukraine	Vladimir Bondar	2
Women and Infants' Hospital, Providence, RI, USA	Paul DiSilvestro	2
Florida Cancer Specialists, West Palm Beach, FL, USA	Marilyn Raymond	2
Klinikum Rechts der Isar der Technischen, München, Germany	Barbara Schmalfeldt	2
E Wolfson Medical Center, Holon, Israel	Tally Levy	2
The Burnside War Memorial Hospital Clinical Trials Centre, Adelaide, SA, Australia	Margaret Davy	1
The Burnside War Memorial Hospital Women's Health Centre, Adelaide, SA, Australia	Margaret Davy	1
British Columbia Cancer Agency, Vancouver, BC, Canada	Paul Hoskins	1
North Estonian Medical Centre, Tallinn, Estonia	Kersti Kukk	1
Hospitel Hôtel-Dieu, Paris, France	Eric Pujade-Lauraine	1
Centre Catherine de Sienné, Nantes, France	Alain Lortholary	1
Universitätsklinikum Schleswig-Holstein, Kiel, Germany	Felix Hilpert	1
Gynäkologisch-Onkologische Praxis am Pelikanplatz, Hannover, Germany	Hans-Joachim Lück	1
Samodzielny Publiczny Szpital Kliniczny, Szczecin, Poland	Jan Lubinski	1
Ginekologiczno-Polożniczy Szpital Kliniczny, Poznań, Poland	Marek Spaczyński	1
Białostockie Centrum Onkologii, Białystok, Poland	Janusz Poznański	1
Spitalul Judetean de Urgenta, Suceava, Romania	Elena Ganea-Motan	1
Spitalul Judetean de Urgenta, Baia Mare, Romania	Dumitru Filip	1
Medical Radiological Research Centre, Obninsk, Russia	Ludmila Krikunova	1

GUZ Altaysky Regional Oncology Dispensary, Barnaul, Russia	Darya Lazare va	1
Sverdlovsk Regional Oncology Dispensary, Ekaterinburg, Russia	Valentina Popova	1
Hospital Universitario 12 de Octubre, Madrid, Spain	César Mendiola	1
Hospital Gregorio Marañón, Madrid, Spain	José Ángel Arranz Arija	1
Ninewells Hospital, Dundee, UK	Fiona Lofts	1
Odeskyi Oblasnyi Onkologichnyi Dyspanser, Odesa, Ukraine	Natalia Tavartkiladze	1
Beth Israel Deaconess Medical Center, Boston, MA, USA	Ursula Matulonis	1
Gynecologic Oncology Associates Inc, Miramar, FL, USA	Laurel King	1
UCSF Helen Diller Family Comprehensive Cancer Center, San Francisco, CA, USA	Lee May Chen	1
Faculty Hospital Bmo, Bmo, Czech Republic	Pavel Ventruba	1

References

1. Royston P, Parmar MK. The use of restricted mean survival time to estimate the treatment effect in randomized clinical trials when the proportional hazards assumption is in doubt. *Stat Med* 2011; 30: 2409-21.