

Viral Genome Sequencing Proves Nosocomial Transmission of Fatal Varicella

Daniel P. Depledge,¹ Julianne Brown,² Jasna Macanovic,³ Gill Underhill,⁴ and Judith Breuer¹

¹Division of Infection and Immunity, University College London, ²Great Ormond Street Hospital VZV Reference Laboratory, Great Ormond Street Hospital, London, ³Wessex Renal and Transplant Services, and ⁴Departments of Clinical Microbiology, Pathology Centre, Queen Alexandra Hospital, Cosham, Portsmouth, United Kingdom

We report the first use of whole viral genome sequencing to identify nosocomial transmission of varicella-zoster virus with fatal outcome. The index case patient, nursed in source isolation, developed disseminated zoster with rash present for 1 day before being transferred to the intensive care unit (ICU). Two patients who had received renal transplants while inpatients in an adjacent ward developed chickenpox and 1 died; neither patient had direct contact with the index patient.

Keywords. fatal varicella; nosocomial transmission; whole genome sequencing.

Varicella-zoster virus (VZV) causes both varicella and, after latency in sensory ganglia, herpes zoster. Universal vaccination of children has reduced the incidence of varicella and its complications in the United States, and vaccination against zoster is available for persons aged >60 years in the United States or >70 years in the United Kingdom. Neither vaccine is licensed for use in immunocompromised patients, for whom both chickenpox and herpes zoster are potentially fatal. Because transmission is thought to arise mainly through aerosolized virus from skin lesions, current advice in hospital settings is to isolate patients with diagnoses of chickenpox or widespread shingles to prevent nosocomial spread to susceptible contacts [1]. Viral genotyping provides a potential tool for linking cases where transmission is not certain [2–4]. However, where viruses are of the same clade [5], current genotyping methods cannot distinguish directly transmitted viruses from those that are unrelated or freely circulating [2]. Here we report the use of a novel method for sequencing whole viral genomes directly from residual diagnostic samples to confirm

nosocomial transmission of fatal varicella in a renal transplant recipient.

METHODS

Patient 1

A 55-year-old woman with end-stage renal failure caused by renovascular disease received a cadaveric renal transplant (MM110; donor negative/recipient positive for human cytomegalovirus [HCMV]), followed by basiliximab induction and triple therapy with tacrolimus, mycophenolate mofetil, and prednisolone. After receiving the transplant, she was an inpatient in a 4-bed bay for 9 days. Fourteen days after discharge she was readmitted into isolation with abdominal pain, a macular rash, and abnormal liver function tests. The rash vesiculated within 24 hours and a clinical diagnosis of varicella was made. She was treated with intravenous acyclovir (10 mg/kg 3 times daily), antibiotics, and cessation of immunosuppression. Three days after admission she was transferred to the ICU, where she received inotropic and ventilator support. She died 3 days later. Varicella pneumonitis was diagnosed, and it was noted that the patient had been VZV immunoglobulin (Ig) G negative before transplantation and had not received varicella vaccine. She had had no known contact with persons with varicella or zoster (Figure 1A; Supplementary Figure 1).

Patient 2

A 61-year-old male patient with advanced renal failure, due to IgA nephropathy with secondary focal segmental glomerulosclerosis, received a preemptive live donor renal transplant (MM121; donor negative/recipient negative for HCMV). He received basiliximab induction, followed by tacrolimus, mycophenolate mofetil, and prednisolone. Despite a history of chickenpox in childhood he was noted to be IgG VZV negative before transplantation. He remained on the transplant ward for 7 days. Eighteen days after discharge, he was readmitted to a different ward with a 4-day history of thoracic vesicular rash, fever, and abnormal liver function. Disseminated shingles was diagnosed, and the presence of VZV DNA was confirmed by polymerase chain reaction. The patient was immediately isolated and treated with intravenous acyclovir (10 mg/kg 3 times daily). He completed 7 days of intravenous acyclovir, followed by 1 week of oral acyclovir. Five weeks after the original presentation, he had further reactivation of VZV in the L1 dermatome and received 2 further weeks of oral acyclovir and 3 months of prophylactic treatment. He made a full recovery and retained good graft function (Figure 1A; Supplementary Figure 1). Importantly, a retrospective review indicated that disseminated shingles was a

Received 17 May 2016; accepted 22 August 2016; published online 28 August 2016.

Correspondence: D. P. Depledge, Division of Infection and Immunity, University College London, Cruciform Building, Gower Street, WC1E6BT, London, UK (d.depledge@ucl.ac.uk).

The Journal of Infectious Diseases® 2016;214:1399–402

© The Author 2016. Published by Oxford University Press for the Infectious Diseases Society of America. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited. DOI: 10.1093/infdis/jiw398




Figure 1. A, Timeline showing hospital admissions and clinical histories for patients 1, 2, and 3. B, Neighbor-joined phylogeny (100 bootstraps) comprising the 3 linked cases (patients 1, 2, and 3) and 2 unlinked cases (patients 3 and 4), together with 35 varicella-zoster virus database sequences obtained from GenBank. Abbreviations: Dx, diagnosis; ICU, intensive care unit; Pt, patient; Tx, treatment.

misdiagnosis and that recurrent varicella should have been diagnosed.

Patient 3 (Index Case Patient)

A 67-year-old man was admitted into respiratory isolation on the transplant ward, with cough and deteriorating renal function after dual cadaveric renal transplant 4 weeks previously. He was isolated immediately and treated with antibiotics. Low-grade HCMV viremia ($3.92 \times 10^2/\text{mL}$) was treated with valgancyclovir (450 mg/d). On day 10 after admission the patient was noted to have a papulovesicular rash on his right leg. A clinical diagnosis of shingles was confirmed, and treatment was started with intravenous acyclovir (10 mg/kg 3 times daily). Within 24 hours, the rash became generalized, respiratory failure developed, and the patient was transferred to the ICU. He recovered with a fully functioning graft. Because this patient was isolated and did not initially have a rash while patient had received a diagnosis of shingles, nosocomial transmission was not suspected. However, further investigation after the death of patient 1 identified a period when all 3 patients had been inpatients on the same ward, albeit physically

separated from patient 3. To investigate whether the 3 cases were linked, we genotyped and sequenced samples from all 3 patients and 2 patients with unrelated cases that occurred in the hospital at the same time (Figure 1A, Supplementary Figure 1).

Genotyping

Residual DNA samples were extracted using the QIAamp DNA Mini Kit (Qiagen) according to the manufacturer's instructions. Amplification and capillary sequencing of a 447-base pair region of open reading frame 22, containing 4 single-nucleotide polymorphisms (SNPs) that discriminate between clades 1/3, 2, 4, and 5 (SNPs 37902, 38055, 38081, and 38177) was carried out as described elsewhere [6]. In addition, the highly variable repeat region, origin of replication (OriS), which also discriminates clade 1 from other clades, was amplified and capillary sequenced using conditions published elsewhere [2].

Whole-Genome Sequencing and Assembly of VZV

Sequencing libraries were constructed as described elsewhere [7, 8] and sequenced on an Illumina MiSeq Sequencer. Data

sets were trimmed and aligned against VZV strain Dumas (NC_001348.1) using BWA (version 0.7.12) [9] and SAMTools software (version 1.0) [10]. Duplicate reads were removed, and a consensus sequence for each data set generated using in-house PERL scripts. Iterative repeat regions (R1, R2, R3, R4, and R5) and the terminal repeat region were excluded from analyses. OriS sequences were successfully captured within the read data and used to verify polymerase chain reaction–based results for all samples. Consensus sequences were aligned using MAFFT software (version 6) [11], and a neighbor-joined phylogenetic tree, based on the Tamura-Nei model, was inferred using MEGA software (version 6.06) [12]. Genome sequences are available in GenBank (accession Nos. KP771912, KP771912, and 3 more awaiting assignment by GenBank as of 16 May 2016).

RESULTS

Virus from all 3 renal patients and 2 “control” patients were typed as clade 1 by SNP genotyping, and all 5 had identical OriS sequences. Whole-genome sequencing showed the 2 putatively linked patients to be 100% identical to the index case patient, patient 3 (Figure 1B) but not the control case patients, whose sequences contained 21 (patient 4) and 11 (patient 5) SNP differences across the genome. Phylogenetic analysis confirmed the clustering of the linked cases, whereas the 2 unlinked clade 1 viruses recovered from other patients in the hospital clustered separately from the linked cases and from each other (Figure 1B).

DISCUSSION

We report the use of whole viral genome sequencing to confirm an unusual case of nosocomial VZV transmission that occurred despite source isolation of the index case patient. Although the 3 patients were in adjacent rooms for more than a week, no immediate connection between them was suspected, partly because shingles was not diagnosed in the index case patient until just before his transfer to the ICU and partly because there was no direct contact between patients. Conventional genotyping confirmed all 3 patients to be infected with clade 1 viruses, the most common genotype in European populations [7]. However, SNP genotyping failed to differentiate the putatively linked viruses from 2 epidemiologically unrelated viruses sampled from other patients at about the same time. Sequencing of highly variable regions, such as the R1 tandem repeat located in open reading frame 11 and the OriS, has also been used to investigate nosocomial transmission [2, 8]. In the cases we report, both linked case patients and controls had identical OriS sequences, and identical R1 sequences in unrelated viruses have also been observed [2, 8].

Although amplicon sequencing to track nosocomial and other transmission events is well established for small RNA viruses including human immunodeficiency virus and hepatitis C

virus [9, 13], whole-genome sequencing is required for larger DNA viruses, because these are genetically less variable. To this end, we have developed a method to sequence whole pathogen genomes directly from clinical material [10]. Our data show, unequivocally, the identity between the viruses recovered from the index case patient and 2 contact case patients. The putatively linked viruses were phylogenetically distinct from other clade 1, viruses including 1 recovered contemporaneously in the same hospital.

Although airborne and environmental transmission of virus from cases of zoster to distant contacts have previously been reported [4], it has not hitherto been possible to confirm nosocomial spread, other than in rare cases where a unique signature sequence was present in an amplicon used for SNP typing [4]. Airborne transmission despite source isolation has been described previously, particularly where viral loads in the index case patient are high (eg, in immunosuppressed patients and those with hemorrhagic varicella). Thus, the need for strict respiratory isolation of such patients, preferably in negative pressure rooms, is paramount [4].

Our case report also underlines 2 important clinical issues. First, primary varicella can be fatal in immunosuppressed patients despite appropriate early treatment and support. VZV antibody testing and immunization against VZV in patients for whom renal transplantation is planned has been shown to prevent or attenuate varicella [14]. Second, widespread rash in an immunosuppressed patient with a history of chickenpox, as in our patient 2, cannot be assumed to be disseminated shingles. Identifying cases of recurrent chickenpox due to nosocomial transmission is important for assuring infection control measures. This case report illustrates how whole-genome sequencing can contribute to this process.

In summary, whole VZV genome sequencing directly from clinical material provides a tractable method for monitoring nosocomial transmission. Applying automated methods directly to residual diagnostic samples, we can generate results within 5 days of sample receipt. The data provide support for the risk of airborne transmission of VZV, especially when the viral load is high, and underline the vulnerability of immunocompromised patients to serious infection, irrespective of varicella history. Our report highlights the ongoing need for vaccines that can prevent viral reactivation in immunocompromised patients as an important measure for control to prevent nosocomial spread.

Supplementary Data

Supplementary materials are available at <http://jid.oxfordjournals.org>. Consisting of data provided by the author to benefit the reader, the posted materials are not copyedited and are the sole responsibility of the author, so questions or comments should be addressed to the author.

Notes

Financial support. This work was supported by the MRC Centre for Medical Molecular Virology, the Medical Research Foundation (New

Investigator award to D. P. D.), the National Institute for Health Research (fellowship to J. Brown), and the NIHR Biomedical Resource Centre at University College London Hospitals NHS Foundation Trust and University College London (J. Breuer).

Potential conflicts of interest. All authors: No potential conflicts of interest. All authors have submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest. Conflicts that the editors consider relevant to the content of the manuscript have been disclosed.

References

1. Public Health England. Shingles (herpes zoster): the green book, chapter 28a. <https://www.gov.uk/government/publications/shingles-herpes-zoster-the-green-book-chapter-28a>. Accessed September 2015.
2. Groves KC, Averbeck K, Quinlivan ML, Scott FT, Breuer J. A case of nosocomial VZV transmission. *J Clin Virol* **2010**; 48:226–7.
3. Molyneux PJ, Parker S, Khan IH, Millar CG, Breuer J. Use of genomic analysis of varicella-zoster virus to investigate suspected varicella-zoster transmission within a renal unit. *J Clin Virol* **2006**; 36:76–8.
4. Lopez AS, Burnett-Hartman A, Nambiar R, et al. Transmission of a newly characterized strain of varicella-zoster virus from a patient with herpes zoster in a long-term-care facility, West Virginia, 2004. *J Infect Dis* **2008**; 197:646–53.
5. Breuer J, Grose C, Norberg P, Tipples G, Schmid DS. A proposal for a common nomenclature for viral clades that form the species varicella-zoster virus: summary of VZV Nomenclature Meeting 2008, Barts and the London School of Medicine and Dentistry, 24–25 July 2008. *J Gen Virol* **2010**; 91:821–8.
6. Parker SP, Quinlivan M, Taha Y, Breuer J. Genotyping of varicella-zoster virus and the discrimination of Oka vaccine strains by TaqMan real-time PCR. *J Clin Microbiol* **2006**; 44:3911–4.
7. Hawrami K, Breuer J. Analysis of United Kingdom wild-type strains of varicella-zoster virus: differentiation from the Oka vaccine strain. *J Med Virol* **1997**; 53:60–2.
8. Hawrami K, Harper D, Breuer J. Typing of varicella zoster virus by amplification of DNA polymorphisms. *J Virol Methods* **1996**; 57:169–74.
9. Li H, Stoddard MB, Wang S, et al. Single-genome sequencing of hepatitis C virus in donor-recipient pairs distinguishes modes and models of virus transmission and early diversification. *J Virol* **2015**; 90:152–66.
10. Depledge DP, Palser AL, Watson SJ, et al. Specific capture and whole-genome sequencing of viruses from clinical samples. *PLoS One* **2011**; 6:e27805.
11. Tamura K, Stecher G, Peterson D, Filipitski A, Kumar S. MEGA6: molecular evolutionary genetics analysis version 6.0. *Mol Biol Evol* **2013**; 30:2725–9.
12. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* **2013**; 30:772–80.
13. Ou CY, Ciesielski CA, Myers G, et al. Molecular epidemiology of HIV transmission in a dental practice. *Science* **1992**; 256:1165–71.
14. Zamora I, Simon JM, Da Silva ME, Piqueras AI. Attenuated varicella virus vaccine in children with renal transplants. *Pediatr Nephrol* **1994**; 8:190–2.