

Light activated antimicrobial agents can inactivate oral malodour causing bacteria

This content has been downloaded from IOPscience. Please scroll down to see the full text.

2016 J. Breath Res. 10 046009

(<http://iopscience.iop.org/1752-7163/10/4/046009>)

View [the table of contents for this issue](#), or go to the [journal homepage](#) for more

Download details:

IP Address: 128.41.35.117

This content was downloaded on 02/11/2016 at 17:35

Please note that [terms and conditions apply](#).

You may also be interested in:

[Quantum dots as enhancers of the efficacy of bacterial lethal photosensitization](#)

N Narband, M Mubarak, D Ready et al.

[Volatiles from oral anaerobes confounding breath biomarker discovery](#)

T Y Khalid, Saliha Saad, John Greenman et al.

[Reduction of bacterial volatile sulfur compound production by licoricidin and licorisoflavan A from licorice](#)

Shin-ichi Tanabe, Jacynthe Desjardins, Chantal Bergeron et al.

[The influence of photodynamic therapy parameters on the inactivation of Candida spp: in vitro and in vivo studies](#)

F Alves, E G Mima, L N Dovigo et al.

[Porphyrins and phthalocyanines: promising molecules for light-triggered antibacterial nanoparticles](#)

Dhananjoy Mondal and Smritilekha Bera

[\beta-Galactosidase and malodor parameters](#)

M Yoneda, Y Masuo, N Suzuki et al.



PAPER

Light activated antimicrobial agents can inactivate oral malodour causing bacteria

RECEIVED
16 August 2016ACCEPTED FOR PUBLICATION
28 September 2016PUBLISHED
18 October 2016OPEN ACCESS
1 November 2016Marika Rai¹, David Spratt¹, Paola R Gomez-Pereira², Jay Patel² and Sean P Nair¹¹ Department of Microbial Diseases, UCL Eastman Dental Institute, University College London, London, UK² Philips Research Cambridge, 101 Cambridge Science Park, Milton Rd, Cambridge CB4 0FY, UKE-mail: sean.nair@ucl.ac.uk**Keywords:** oral malodour, photodynamic therapy, methylene blue

Original content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/4.0/).

Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

**Abstract**

Oral malodour is a common condition which affects a large proportion of the population, resulting in social, emotional and psychological stress. Certain oral bacteria form a coating called a biofilm on the tongue dorsum and degrade organic compounds releasing volatile sulfur compounds that are malodorous. Current chemical treatments for oral malodour such as mouthwashes containing chlorhexidine or essential oils, are not sufficiently effective at reducing the bacterial load on the tongue. One potential alternative to current chemical treatments for oral malodour is the use of light activated antimicrobial agents (LAAAs), which display no toxicity or antimicrobial activity in the dark, but when exposed to light of a specific wavelength produce reactive oxygen species which induce damage to target cells in a process known as photodynamic inactivation.

This study aimed to determine whether oral malodour causing bacteria were susceptible to lethal photosensitization. Five bacterial species that are causative agents of oral malodour were highly sensitive to lethal photosensitization and were efficiently killed by methylene blue in conjunction with 665 nm laser light. Between 4.5–5 log₁₀ reductions in the number of viable bacteria were achieved with 20 μM methylene blue and 14.53 J cm⁻² laser light for *Porphyromonas gingivalis*, *Prevotella intermedia*, *Peptostreptococcus anaerobius* and *Solobacterium moorei*. The number of viable cells fell below the limit of detection in the case of *Fusobacterium nucleatum*.


These findings demonstrate that methylene blue in combination with 665 nm laser light is effective at killing bacteria associated with oral malodour, suggesting photodynamic therapy could be a viable treatment option for oral malodour.

1. Introduction

Upon irradiation with light of a suitable wavelength, a photoactivated antimicrobial agent, known as a photosensitizer, is able to generate reactive oxygen species which can cause damage to target cells. This process is known as photodynamic inactivation. Photodynamic therapy has been used extensively in the treatment of skin conditions, cancer, ophthalmology and more recently, in dentistry and oral infections (Diamond *et al* 1972, Johnsson *et al* 1987, Braun *et al* 2008, Trachtenberg 2008, Taylor and Gonzalez 2009, Garcez and Núñez 2015).

A photosensitizer in its ground singlet state has two electrons with opposite spins in a low energy molecular orbital. Once exposed to light of an appropriate wavelength, the photosensitizer absorbs a photon and one of the electrons transitions to a high energy excited singlet

state. This energy can then be lost through fluorescence and internal heat conversion, or can undergo a conversion to an excited triplet state via intersystem crossing (Robertson *et al* 2009). This excited triplet state is longer lived than the excited singlet state (Juzeniene and Moan 2007), and can react in one of two ways to produce reactive oxygen species. These reactive oxygen species can then cause destruction to the target area (figure 1). The type I mechanism occurs via electron or hydrogen transfer directly from the photosensitizer upon direct reaction with a substrate, such as water, which results in the formation of the superoxide anion (O₂⁻). These radicals can then react with oxygen to produce highly reactive species, such as the hydroxyl radical and hydrogen peroxide. The type II mechanism involves the photosensitizer, in its excited triplet state, reacting with molecular oxygen to produce highly reactive singlet oxygen. These reactive oxygen species can then cause rapid cytotoxic



effects to the target area and result in target cell death via damage to bacterial cell walls and cytoplasmic membranes. This in turn results in the leakage of cellular contents, damage to DNA, such as breaks in single or double stranded DNA and possible inactivation of membrane transport systems (Raghavendra *et al* 2009, Rajesh *et al* 2011, Huang *et al* 2012).

Photodynamic therapy is a promising alternative antimicrobial therapy for dental and oral conditions, due to the ease of accessibility for both photosensitizer and light application, and successful results have been seen in the treatment of dental caries (Nagata *et al* 2012), peri-implantitis (Bassetti *et al* 2013), and periodontitis (Betsy *et al* 2014).

Halitosis, also known as oral malodour, is estimated to affect approximately 8–50% of the adult population, who perceive that they suffer from oral malodour as a chronic problem (Porter and Scully 2006), with 85–90% of cases of malodour being of intra-oral origin (Quirynen *et al* 2009). It is thought that the dorso-posterior region of the tongue is the main source of malodour of an oral origin due to the unique morphology and the location, which makes the area hard to cleanse. Bacteria located on the tongue dorsum, mainly Gram-negative anaerobic bacteria, are able to putrefy organic compounds found in postnasal drip and in the tongue coating, and produce volatile sulfur compounds (VSCs), which are characteristic of oral malodour, as by products (Kazor *et al* 2003, Bollen and Beikler 2012).

Oral malodour causing bacteria are able to degrade sulfur containing amino acids such as methionine and cysteine, and non-sulfur containing amino acids such as lysine, arginine and tryptophan, and bio-transform them into VSCs such as hydrogen sulphide, methylmercaptan and dimethyl sulphide (Greenman 1999).

The bacteria which have been most commonly associated with oral malodour in the past are Gram-negative pathogens and include *Fusobacterium nucleatum*, *Porphyromonas gingivalis*, *Prevotella intermedia*, *Treponema denticola* and *Tannerella forsythensis* (Morita and Wang 2001, Porter and Scully 2006). These bacteria reside in the periodontal pockets and on the posterior of the tongue and mouth, thereby avoiding removal and cleansing (Krespi *et al* 2006). Gram-positive bacteria have also been implicated in halitosis. The Gram-positive, anaerobic coccus *Peptostreptococcus anaerobius* has been found to produce hydrogen sulphide from L-cysteine (Persson 1990), and has been isolated on the tongue dorsum of people suffering from halitosis (Tyrell *et al* 2003). *Solobacterium moorei*, a Gram-positive bacterium which has only been found to be present in the oral cavities of halitosis sufferers, has also been implicated in oral malodour, suggesting that there are distinct bacteria associated with halitosis (Haraszthy *et al* 2007). A study carried out by Tanabe and Grenier characterising the capacity of *S. moorei* to produce volatile sulphur compounds found that the bacterium was able to produce high levels of hydrogen

sulphide from cysteine. This is thought to be catalysed by the enzyme cysteine desulphydrase, the gene of which has been identified in the genome of *S. moorei* (Tanabe and Grenier 2012).

A preliminary study trialing photodynamic therapy of halitosis suggests that this treatment is effective at reducing VSC concentrations (Lopes *et al* 2014). However there is little data in the reported literature on the ability of light in combination with a photosensitizer to inactivate bacteria, in particular Gram-positive bacteria, that cause halitosis. In the study reported here we have investigated the capacity of the photosensitizer methylene blue in combination with red light to inactivate important Gram-positive and Gram-negative halitosis causing bacteria.

2. Materials and methods

2.1. Bacterial strains and growth conditions

The bacteria used in this study were *P. gingivalis* (W50), *F. nucleatum* (NCTC 10562), *Pr. intermedia* (DSMZ 20706), *Pt. anaerobius* (NCTC 11460) and *S. moorei* (DSMZ 22971). Bacteria were maintained on fastidious anaerobe agar (FAA; LabM, Bury, UK) supplemented with 5% v/v defibrinated horse blood (E&O Laboratories, Bonnybridge, Scotland) under anaerobic conditions (80% N₂; 10% H₂; 10% CO₂) at 37 °C. For the photodynamic inactivation studies bacteria were grown in liquid culture until they reached stationary phase. The Gram-negative bacteria, *P. gingivalis*, *F. nucleatum*, *Pr. intermedia* were grown anaerobically for 18 h in Brain Heart Infusion broth (BHI broth; Sigma, Dorset, UK) supplemented with 5 µg ml⁻¹ hemin (Sigma) and 0.5 µg ml⁻¹ menadione (Sigma). *Pt. anaerobius* was grown anaerobically for 22 h in Brain Heart Infusion broth supplemented with 5 µg ml⁻¹ hemin and 0.5 µg ml⁻¹ menadione. *S. moorei* was grown anaerobically for 24 h in tryptic soy broth (TSB; Sigma) supplemented with 5% v/v horse serum (Sigma).

2.2. Photosensitiser and laser

The phenothiazine dye methylene blue (Sigma, Dorset, UK) was dissolved in sterile water and filtered through a 0.22 µm syringe filter (Merck Millipore). All solutions were stored in the dark at room temperature prior to use.

A low power, non-thermal, Periwave diode laser (Ondine Biomedical Inc.) was utilised in this study (wavelength 665 nm, P_{\max} 160 mW). The laser was clamped onto a stand in such a way that the light emitting end was positioned directly above a well of a 96 well microtiter plate, at a height that allowed the laser beam to cover the entire well upon irradiation. The power delivered by the laser to the irradiated well was measured using a laser power meter (Gentec, TPM-300).

2.3. Photosensitisation procedure

Bacterial cells were pelleted from liquid culture by centrifugation (15 093 g, 3 min) and resuspended in pre-reduced PBS. Each bacterial suspension was adjusted

to an optical density at 600 nm which corresponded to approximately 1×10^8 cells ml⁻¹. The four test groups were as follows: light plus photosensitiser (L⁺S⁺), light alone (L⁺S⁻), photosensitiser alone (L⁻S⁺) and the control which was neither treated with photosensitiser nor light (L⁻S⁻). For the L⁺S⁺ group, 100 µl of methylene blue was added to an equal volume of the bacterial culture in triplicate wells of a 96 well flat bottomed plate (VWR, Leicestershire, UK) and irradiated with 665 nm laser light with an energy density of 14.53 J cm⁻², corresponding to a 30 s exposure to light. Three additional wells containing 100 µl of methylene blue and 100 µl of the bacterial suspension were kept in the dark (L⁻S⁺) to assess the toxicity of the photosensitiser alone (L⁻S⁺). Controls, L⁺S⁻ and L⁻S⁻ in triplicate, consisted of 100 µl of pre-reduced PBS added to an equal volume of the bacterial cultures. The L⁺S⁻ wells were irradiated with laser light, while the L⁻S⁻ wells, were kept in the dark. Following irradiation/dark incubation, each sample was serially diluted 10-fold in pre-reduced PBS. A 25 µl portion of each dilution was spotted and spread plated onto 5% horse blood Fastidious anaerobe agar plates in triplicate, and the plates were incubated anaerobically at 37 °C for 48–72 h. The surviving CFU/well were determined by enumerating the viable bacteria on plates. Experiments were performed on three different occasions in triplicate.

To assess the effect of light dose on lethal photosensitization, bacteria were grown as previously detailed until they had reached the stationary phase of growth. Methylene blue was diluted in PBS and added to an equal volume of bacteria in 96 well plates to give a final concentration of 10 µM. The L⁺S⁺ wells were irradiated with 665 nm laser light energy doses of 2.42 J cm⁻², 4.84 J cm⁻² or 9.69 J cm⁻², corresponding to 5, 10 or 20 s irradiation respectively. Three additional wells containing 100 µl of methylene blue and 100 µl of the bacterial suspensions were kept in the dark (L⁻S⁺) and 100 µl of PBS was also added to 100 µl of the bacterial suspensions in a further six wells, three of which were irradiated with laser light (L⁺S⁻), while the remaining three were kept in the dark (L⁻S⁻).

Following irradiation/dark incubation, each sample was serially diluted 10-fold in pre-reduced PBS. A 25 µl aliquot of each dilution was spotted and spread plated onto 5% horse blood Fastidious anaerobe agar plates in triplicate, and the plates were incubated anaerobically at 37 °C for 48–72 h. The surviving CFU/well were enumerated by viable counting. Experiments were performed on three different occasions in triplicate.

2.4. Statistics

The columns and error bars shown in the graphs represent the mean and standard deviations of three individual experiments performed in triplicate. Statistical comparisons between groups were carried out using two-way ANOVA with post-hoc Bonferroni analysis, and comparisons within groups were carried

out using one-way ANOVA with post-hoc Bonferroni analysis. A P value of less than 0.0001 was considered statistically significant for two-way ANOVA, and a P value of less than 0.05 was considered statistically significant for one-way ANOVA.

3. Results

3.1. The effect of photosensitizer dose on lethal photosensitization of oral malodour causing bacteria

The effects of increasing photosensitizer dose on the viability of oral malodour causing bacteria are shown in figure 2. When the bacteria were treated with different concentrations of methylene blue and 665 nm laser light, there was a significant ($P < 0.0001$) reduction in the viable cell counts. When *P. gingivalis* was treated with methylene blue and exposed to 14.53 J cm^{-2} of laser light, a significant ($P < 0.0001$) $3 \log_{10}$ reduction in the number of viable cells was seen with the lowest concentration of methylene blue used. When treated with the highest concentration of methylene blue, there was a $5 \log_{10}$ reduction in viable cells (figure 2(A)). Treatment of *Pr. intermedia* with methylene blue and exposure to 14.53 J cm^{-2} laser light caused a $1.6 \log_{10}$ ($P < 0.0001$) reduction in the number of viable bacteria at $2 \mu\text{M}$ methylene blue. At the highest concentration of methylene blue, there was a $4.5 \log_{10}$ reduction in viable cells (figure 2(B)). In the case of *F. nucleatum*, in the groups treated with either laser light alone (L^+S^-) or with methylene blue alone (L^-S^+) the viable counts were no different from the untreated control (L^-S^-). When exposed to $2 \mu\text{M}$ methylene blue and irradiated with 665 nm laser light, there was a $3.1 \log_{10}$ ($P < 0.0001$) reduction in the number of viable bacteria. The number of viable bacteria fell below the limit of detection with the highest concentration of methylene blue used upon exposure to 665 nm light. The bactericidal effect was dependent on photosensitizer concentration, with significantly ($P < 0.05$) more kills at 5, 10 and $20 \mu\text{M}$ methylene blue as compared to treatment with $2 \mu\text{M}$ methylene blue (figure 2(C)). Interestingly with *P. gingivalis* and *Pr. Intermedia*, a 1 and a $0.5 \log_{10}$ reduction in viable cells occurred upon irradiation with laser light alone, respectively.

The Gram-positive bacteria associated with oral malodour were also sensitive to lethal photosensitization. At the lowest concentration of methylene blue, a $2.2 \log_{10}$ ($P < 0.0001$) reduction in the number of viable *Pt. anaerobius* was achieved upon exposure to 665 nm laser light, as seen in figure 2. Treatment with $20 \mu\text{M}$ methylene blue and exposure to 14.53 J cm^{-2} laser light resulted in a $5.0 \log_{10}$ ($P < 0.0001$) reduction in viable cells (figure 2(D)).

Irradiation of *S. moorei* in the presence of $2 \mu\text{M}$ methylene blue resulted in $0.9 \log_{10}$ ($P < 0.0001$) reduction in viable bacterial cells. Treatment with the highest concentration of methylene blue and exposure

to laser light resulted in a $4.6 \log_{10}$ reduction in viable cells. As was the case with *F. nucleatum*, photodynamic inactivation of *S. moorei* was dependent on photosensitizer concentration, with a significant ($P < 0.05$) reduction in viable colony counts at 5, 10 and $20 \mu\text{M}$ methylene blue as compared to $2 \mu\text{M}$ (figure 2(E)).

3.2. The effect of light dose on lethal photosensitization of oral malodour causing bacteria

To study the effect of different light doses in combination with methylene blue on the viability of oral malodour causing bacteria, cells were exposed to light doses of either 2.42 J cm^{-2} , 4.84 J cm^{-2} or 9.68 J cm^{-2} in the presence of $10 \mu\text{M}$ methylene blue.


Upon irradiation with the lowest light dose in combination with $10 \mu\text{M}$ methylene blue, there was a significant ($P < 0.0001$) reduction in the number of viable *P. gingivalis* cells (figure 3(A)). At the lowest light dose, there was a $2.7 \log_{10}$ reduction in the number of viable cells, which increased to a $3.5 \log_{10}$ reduction in the number of viable cells at the highest light dose.

Irradiation of *Pr. intermedia* with varying light doses in the presence of $10 \mu\text{M}$ methylene blue resulted in a significant ($P < 0.0001$) reduction in viable cell counts (figure 3(B)). Exposure to a light dose of 2.42 J cm^{-2} with $10 \mu\text{M}$ methylene blue caused a $1.4 \log_{10}$ ($P < 0.0001$) reduction in cell numbers. At the longest exposure time there was a $3.1 \log_{10}$ reduction in the number of viable cells. Photodynamic inactivation of *Pr. intermedia* was dependent on light dose, with a significant ($P < 0.05$) reduction in viable cells after 10 and 20 s exposure as compared to 5 s exposure. Treatment of *F. nucleatum* cells with $10 \mu\text{M}$ methylene blue in combination with 665 nm laser light for 5 s were reduced in number by $3.6 \log_{10}$ ($P < 0.0001$). The number of viable cells fell below the limit of detection (LoD = 10 cells) after 20 s exposure (figure 3(C)). For *P. gingivalis* and *Pr. intermedia*, there was also a significant ($P < 0.0001$) reduction in viable cell numbers when treated with light alone (L^+S^-).

Upon irradiation of *Pt. anaerobius* with 2.42 J cm^{-2} laser light in the presence of methylene blue, viable cell numbers fell by $2 \log_{10}$ ($P < 0.0001$). At the highest light dose, a $2.9 \log_{10}$ reduction in the number of viable cells was observed (figure 3(D)). In the case of *S. moorei*, a significant $1 \log_{10}$ ($P < 0.0001$) reduction in the number of viable cells was observed with 2.42 J cm^{-2} of laser light and $10 \mu\text{M}$ methylene blue. With a 20 s exposure time, there was a $3.9 \log_{10}$ ($P < 0.0001$) reduction in the number of viable *S. moorei* cells.

4. Discussion

Oral malodour can arise due to a number of reasons, including transient, lifestyle, oral and non-oral causes. Whatever the cause may be, oral malodour is an extremely important condition due to the adverse impact it can have on the professional and personal



lives of sufferers. Suffering from oral malodour can lead to psychological changes, which can in turn, lead to social and personal isolation (Bollen and Beikler 2012). As such oral malodour sufferers can have a poorer quality of life compared to those that do not suffer. Current treatments include the advocacy of proper oral hygiene, mechanical removal of the tongue coating that is responsible for the malodour and the use of antimicrobial toothpastes and mouthwashes to reduce the bacterial load in the oral cavity (De Geest et al 2016). The ability of light activated antimicrobial agents (LAAAs) to cause photodynamic inactivation

of a number of bacteria has been well documented (Demidova and Hamblin 2004, Jori et al 2006). In particular, there have been numerous *in vitro* and *in vivo* studies on the use of photodynamic inactivation for the treatment of oral pathogens (Komerik et al 2003, Wilson 2004, Fimple et al 2008, Schiffner et al 2014).

In this study we found that using the phenothiazine dye methylene blue and 665 nm red laser light, lethal photosensitization of the oral malodour causing bacteria *P. gingivalis*, *Pr. intermedia*, *F. nucleatum*, *Pt. anaerobius* and *S. moorei* could be achieved. The five bacteria




Figure 3. The capacity of 665 nm laser light at doses of 2.42 J cm^{-2} , 4.84 J cm^{-2} and 9.69 J cm^{-2} (corresponding to 5, 10 and 20 s exposure) in combination with $10 \mu\text{M}$ methylene blue to cause photodynamic inactivation of oral malodour causing bacteria. An equal volume of either PBS (S^-) or methylene blue (S^+) ($10 \mu\text{M}$) was added to $100 \mu\text{l}$ of the bacterial suspensions and either kept in the dark (L^-) or exposed to 665 nm laser light (L^+) (energy doses ranging from 2.42 J cm^{-2} – 14.53 J cm^{-2}). After irradiation/dark incubation, samples were serially diluted and the surviving CFU/well were enumerated. Error bars represent the standard deviation from the mean. *** $P < 0.0001$ (two-way ANOVA). Experiments were performed three times in triplicate and the combined data are shown. (A) Lethal photosensitisation of *P. gingivalis*. (B) Lethal photosensitisation of *Pr. intermedia*. (C) Lethal photosensitisation of *F. nucleatum*. (D) Lethal photosensitisation of *Pt. anaerobius*. (E) Lethal photosensitisation of *S. moorei*.

tested in this study were all found to be susceptible to lethal photosensitisation when treated with either different concentrations of methylene blue or different light doses, with significant reduction in the number of viable cells seen for each of the bacteria.

Photodynamic inactivation was found to be dependent on photosensitiser dose in the case of *F. nucleatum* and *S. moorei*, while it was dependent on laser light dose in the case of *Pr. intermedia*. Significant kills were also demonstrated with the lowest concentration of methylene blue ($2 \mu\text{M}$) and the lowest laser light dose (2.42 J cm^{-2}) for each of the five bacteria tested. Low photosensitizer and light doses would be

beneficial in a clinical setting because the laser light dose corresponds to a short irradiation time of 5 s, and at such a low concentration of methylene blue, temporary staining of the tongue blue would be minimal. While significant kills were achieved with low energy doses, complete eradication of any of the bacteria was not achieved with low methylene blue concentrations and low light doses.

Chan and Lai demonstrated that *F. nucleatum* was highly susceptible to photoinactivation with methylene blue, achieving reductions in the number of bacteria of 1.3 logs (Chan and Lai 2003). A previous study conducted by Williams *et al* found that treatment with

20 $\mu\text{g ml}^{-1}$ of toloum chloride, also known as toluidine blue, a phenothiazine dye closely related to methylene blue, and exposure to 4.8 J cm^{-2} of 633 nm laser light resulted in a maximum mean log reduction of 2.81 in the number of *F. nucleatum* as compared to 1.14 mean logs for *Pr. intermedia* and 2.48 mean logs for *Peptostreptococcus* (now *Parvimonas*) *micros* also tested in their study (Williams et al 2006). Theirs is similar in the case of the results presented within this study with *F. nucleatum* being the most affected by photodynamic inactivation, as it was the only bacterium tested that had viable cell counts fall below the limit of detection. This demonstrates that with an increased energy dosage, higher levels of kill for *F. nucleatum* can be achieved. The results of this study are favourable however, as the concentration of methylene blue utilised in this study was much lower compared to the concentration of toloum chloride, and the 665 nm light source used in this study would have been able to penetrate more deeply as compared to the 633 nm laser device used by Williams et al therefore suggesting that our regime using methylene blue and 665 nm light produces a more promising outcome.

Energy doses in excess of 20 J cm^{-2} from a 633 nm diode laser in combination with derivatives of chlorin e6, have been demonstrated to completely eradicate cultures of *P. gingivalis* and *F. nucleatum* (Pfitzner et al 2004). In our study, the maximum energy dose tested was 14.53 J cm^{-2} which resulted in significant bacterial reduction, and in the case of *F. nucleatum*, reduced cells counts below our limit of detection. It may be possible that our regime completely eradicated *F. nucleatum* cells, and as such the findings of this study are an improvement on those already reported due to the low photosensitiser and light dose combination used in comparison to previous studies.

In the case of *P. gingivalis*, photodynamic inactivation was not dependent on the laser light dose. This is in contrast to results reported in another study using toluidine blue O (TBO) in combination with a helium/neon gas laser with a wavelength of 632.8 nm (Bhatti et al 1997), which resulted in complete bacterial kills using 2.2 J cm^{-2} of light. Conversely, an *in vivo* study investigating the killing of *P. gingivalis* also using Toluidine blue, found that photodynamic inactivation was not dependent on light dose response using a 630 nm diode laser, but was dependent on photosensitizer concentration at lower light doses. This effect was thought to be due to sufficient doses of energy being supplied to the sample in order to activate all of the photosensitiser molecules present, thereby generating enough oxygen radicals to kill the bacteria (Komerik et al 2003). It has also been found that as well as killing of *P. gingivalis* cells by photodynamic inactivation, virulence factors of this pathogen are also inactivated by the process, thereby providing another advantage of this therapy over conventional treatments (Komerik et al 2000).

Exposure to 665 nm laser light alone resulted in a significant reduction in the numbers of viable *P. gingivalis* and *Pr. intermedia* cells, albeit in a lower magnitude

compared to the L^+S^+ group, as seen in figures 2(A), (B), 3(A) and (B). *P. gingivalis* and *Pr. intermedia* are black pigmented anaerobes, both of which have been found to accumulate the endogenous porphyrins, dimeric protoporphyrin IX and monomeric protoporphyrin IX. These endogenous porphyrins have a strong absorption peak at approximately 400 nm which corresponds to blue light, and numerous studies have shown that upon irradiation with blue light, both of these pathogens can be photoinactivated in the absence of an exogenous photosensitiser (Soukos et al 2005, Hope et al 2013, 2016, AbdulAzeez et al 2014). Whilst those studies focused on blue light, which strongly activates these endogenous porphyrins, there is a smaller absorption peak at approximately 600–700 nm, corresponding to the red light which was utilized in our study. Another study utilized red light only to kill bacteria via photodynamic action, and found that following a single exposure to red light at 632.8 nm at a 100 mW cm^{-2} light intensity and with a 360 J cm^{-2} energy density, there was a pronounced photodynamic effect on *P. gingivalis* and *Pr. intermedia* (Konig et al 2000). This supports our data reported herein where there is a reduction in viable bacteria in the L^+S^- group. However although our study utilises a higher power laser, the energy dose is almost 25 times lower with a similar effect, therefore the regime utilised in this study is favourable compared to those previously reported. These results also demonstrate that using a laser light source in combination with a photosensitiser, in the case of our study methylene blue, much lower energy doses can be delivered to the target area or tissue, with significant bactericidal effects and reduced chances of causing damage to the surrounding areas.

It is interesting to note that in our study the Gram-negative bacteria were more susceptible to photodynamic inactivation than the Gram-positive bacteria, *Pt. anaerobius* and *S. moorei*. It is widely thought that Gram-negative bacteria are less susceptible to photodynamic inactivation due to structural differences in the cells as compared to Gram-positive bacteria, namely the presence of an outer membrane which serves to act as a permeability barrier (Maisch et al 2004). This apparent paradox with our findings is probably because uptake of methylene blue may be increased in Gram-negative bacteria due to its cationic nature, low molecular weight and hydrophilicity, thereby allowing the photosensitiser to pass across the porin-protein channels in the outer membrane of Gram-negative bacteria. As well as passing across the outer membrane, methylene blue is also able to localise to the anionic lipopolysaccharide found within the outer membrane and form methylene blue dimers, which themselves add to and enhance the photodynamic effect (Maisch et al 2004), and this effect could account for the differences in susceptibility seen in our study.

Photodynamic inactivation studies on *Peptostreptococcus* species have demonstrated that these bacteria are able to be killed by this technique. A study using photodynamic therapy to treat endodontic infection

found that *Peptostreptococcus* (now *Parvimonas*) *micros* was completely eradicated after photodynamic therapy with 67 μM methylene blue and exposure to 30 J cm^{-2} diode laser light (Soukos *et al* 2006). Another study carried out investigating the susceptibility of anaerobic bacteria isolated from periodontal disease to PDI with 84 μM chlorin e6 and 15 J cm^{-2} laser light resulted in a 4–6 log reduction in the 4 Peptostreptococcal species treated (Drulis-Kawa *et al* 2005). These studies highlight the susceptibility of peptostreptococci to photodynamic inactivation and as such suggest that PDI is a viable option for the eradication of these bacteria.

PDI studies on *Pt. anaerobius* are scarce, with one study achieving a 3 log kill of the bacterium using hematoporphyrin derivative (HpD) in combination with exposure to a 100 mW projector light for 20 min (Venezio 1985). In our study a significant reduction in the numbers of this bacterium were achieved using the highest concentration of methylene blue, and there was also an observed light dose response in the viability of this bacteria, however this was not statistically significant. Our study also utilised a much lower exposure time as compared to Venezio *et al*, with our maximum exposure time of 30 s resulting in a 5 \log_{10} reduction in bacteria, demonstrating that our current regime is favourable to previously reported regimes.

Our results demonstrate that a low dose of photosensitiser coupled with a low exposure time is able to cause a significant level of kill in the number of bacteria, although higher concentrations of methylene blue and longer exposure times result in higher reductions in the number of viable bacteria.

To our knowledge there have been no studies published on the use of LAAs to kill *S. moorei*, a Gram-positive bacterium implicated in oral malodour. A study by Tanabe and Grenier has found that *S. moorei* is able to produce hydrogen sulphide from cysteine, and VSC's were also produced from serum, saliva and mucin and as such *S. moorei* has been proposed to be a major player in oral malodour (Tanabe and Grenier 2012). Previous studies of photodynamic inactivation of Gram-positive bacteria have yielded successful results and significant bacterial kills. A study into the effects of photodynamic therapy of biofilms of *Enterococcus faecalis*, a Gram-positive bacterium found in the oral cavity and which is known to be resistant to some antibiotics, found that treatment with methylene blue or TBO, with 271 J cm^{-2} and 106.4 J cm^{-2} respectively, resulted in significant levels of bacterial killing (Lopez-Jimenez *et al* 2015). Another study into the photosensitisation of *E. faecalis* biofilms determined that pre-incubation for 10 min with 60 μM methylene blue with subsequent exposure to 9.6 J cm^{-2} resulted in the disruption of the biofilms, and determined that significant levels of bacterial reduction could be achieved using this photosensitiser (Garcez *et al* 2013). *Staphylococcus aureus* and *Staphylococcus epidermidis* are also Gram-positive bacteria, most commonly associated with skin infections and implant associated infections in hospitalised patients.

Biofilms of both of these organisms were treated with 40 μM TBO combined with energy doses up to 200 J cm^{-2} and it was found that a significant decrease in the bacterial biofilms could be achieved (Sharma *et al* 2008). These studies serve to show that Gram-positive have successfully been proven to be susceptible to phenothiazine photosensitizers, as used in our study.

In our study, there was a significant reduction in the number of viable cells when treated with the maximum concentration of methylene blue and exposure to laser light, with the combination of the two lower in our study than previously reported by others. The levels of bacterial reduction were slightly lower for *S. moorei* as compared to *Pt. anaerobius*, however they were still significant. When compared to the other 4 bacteria in our study, *S. moorei* reported the lowest levels of bacterial kill at the lower concentrations of methylene blue used. Significant levels of bacterial reduction were achieved for *S. moorei*, proving that a combination of methylene blue and red diode laser light is effective, however higher concentrations of methylene blue may be more beneficial in eradicating this bacteria. These results are extremely positive as they suggest that treatment with red light and methylene blue, both of which have been previously used in a clinical setting, can be potentially employed as a therapeutic method to inactivate certain oral malodour associated bacteria.

Our study has demonstrated significant kills of oral malodour associated bacteria can be achieved with 665 nm laser light and methylene blue, with both low photosensitiser concentrations and low light doses.

5. Conclusion

In conclusion, the results of this study have demonstrated the photodynamic inactivation of five important oral malodour causing pathogens, namely *P. gingivalis*, *Pr. intermedia*, *F. nucleatum*, *Pt. anaerobius* and *S. moorei*. This suggests that photodynamic inactivation is a promising treatment modality for oral malodour.

The use of PDI to reduce the bacterial load of oral malodour causing bacteria would be favourable over current treatment options such as mouthwashes which serve to mask malodour or cannot be used for prolonged periods of time due to the potential effect on taste.

Additionally, the use of the Periowave diode laser as the activating light source is also favourable as it is a low power laser. This could possibly be favourable in a clinical setting as treatment times would be short, as the relatively short exposure times resulted in a significantly high level of bacterial kill and such low photosensitiser doses would stain less intensely. Although the results of this study prove to be promising, further studies are needed in order to determine whether high levels of bacterial kill can be obtained when the bacteria are in a biofilm state, as they would be when found naturally in the tongue coating.

Acknowledgments

MR has been funded for this study by Philips Oral Healthcare: 513517, and the Engineering and Physical Sciences Research Council (EP/K504610/1): 508091.

References

- AbdulAzeez A R, Mahmood M S and Ali W M 2014 Phototoxic effect of visible blue light on aggregatibacter actinomycetemcomitans and *Porphyromonas gingivalis* with chronic periodontitis (an *in vitro* study) *J. Baghdad Coll. Dent.* **27** 144–50
- Bassetti M, Schär D, Wicki B, Eick S, Ramseier C A, Arweiler N B, Sculean A and Salvi G E 2013 Anti-infective therapy of peri-implantitis with adjunctive local drug delivery or photodynamic therapy: 12 months outcomes of a randomized controlled clinical trial *Clin. Oral Implants Res.* **25** 279–87
- Betsy J, Prasanth C S, Baiju K V, Prasanthila J and Subhash N 2014 Efficacy of antimicrobial photodynamic therapy in the management of chronic periodontitis: a randomized controlled clinical trial *J. Clin. Periodontol.* **41** 573–81
- Bhatti M, MacRobert A, Meghji S, Henderson B and Wilson M 1997 Effect of dosimetric and physiological factors on the lethal photosensitization of *Porphyromonas gingivalis* *in vitro* *Photochem. Photobiol.* **65** 1026–31
- Bollen C M L and Beikler T 2012 Halitosis: the multidisciplinary approach *Int. J. Oral Sci.* **4** 55–63
- Braun A, Dehn C, Krause F and Jepsen S 2008 Short-term clinical effects of adjunctive antimicrobial photodynamic therapy in periodontal treatment: a randomized clinical trial *J. Clin. Periodontol.* **35** 877–84
- Chan Y and Lai C H 2003 Bactericidal effects of different laser wavelengths on periodontopathic germs in photodynamic therapy *Lasers Med. Sci.* **18** 51–5
- De Geest S, Laleman I, Teughels W, Dekeyser C and Quirynen M 2016 Periodontal diseases as a source of halitosis: a review of the evidence and treatment approaches for dentists and dental hygienists *Periodontology 2000* **71** 213–27
- Demidova T N and Hamblin M R 2004 Photodynamic therapy targeted to pathogens *Int. J. Immunopathol. Pharmacol.* **17** 245
- Diamond I et al 1972 Photodynamic therapy of malignant tumours *Lancet* **2** 1175–7
- Drulis-Kawa Z, Bednarkiewicz A, Bugla-Ploskonska G, Strek W and Doroszkiwicz W 2005 The susceptibility of anaerobic bacteria isolated from periodontal diseases to photodynamic inactivation with Fotolon (chlorin e6) *Pol. J. Microbiol.* **54** 305–10
- Fimple J L, Fontana C R, Foschi F, Ruggiero K, Song X, Pagonis T C and Soukos N S 2008 Photodynamic treatment of endodontic polymicrobial infection *in vitro* *J. Endod.* **34** 728–34
- Garcez A S, Nunez S C, Azambuja Jr N, Fregnani E R, Rodriguez H M H, Hamblin M R, Suzuki H and Ribeiro M S 2013 Effects of photodynamic therapy on Gram-positive and Gram-negative bacterial biofilms by bioluminescence imaging and scanning electron microscopic analysis *Photomed. Laser Surg.* **31** 519–25
- Garcez A S and Núñez S C 2015 Bacterial reduction in root canals using antimicrobial photodynamic therapy *Lasers Dent.* **1** 133–6
- Greenman J 1999 Dental plaque revisited, oral biofilms in health and disease *Microbial Aetiology of Halitosis* ed H N Newman and M Wilson (Cardiff: Bioline Publications) pp 419–42
- Hamblin M R and Hasan T 2004 Photodynamic therapy: a new antimicrobial approach to infectious disease? *Photochem. Photobiol. Sci.* **3** 436–50
- Haraszthy V I, Zambon J J, Sreenivasan P K, Zambon M M, Gerber D, Rego R and Parker C 2007 Identification of oral bacterial species associated with halitosis *J. Am. Dent. Assoc.* **138** 1113–20
- Hope C K, Hindley J A, Khan Z, de Josselin de Jong E and Higham S M 2013 Lethal photosensitization of *Porphyromonas gingivalis* by their endogenous porphyrins under anaerobic conditions: an *in vitro* study *Photodiag. Photodyn. Ther.* **10** 677–82
- Hope C K, Strother M, Creber H K and Higham S M 2016 Lethal photosensitization of Prevotellaceae under anaerobic conditions by their endogenous porphyrins *Photodiagnosis Photodyn. Ther.* **13** 344–6
- Huang L, Xuan Y, Koide Y, Zhiyentayev T, Tanaka M and Hamblin M R 2012 Type I and Type II mechanisms of antimicrobial photodynamic therapy: an *in vitro* study on gram-negative and gram-positive bacteria *Lasers Surg. Med.* **44** 490–9
- Johnsson A, Kjeldstad B and Melo T B 1987 Fluorescence from pilosebaceous follicles *Arch. Dermatol. Res.* **279** 190–3
- Jori G, Fabris C, Soncin M, Ferro S, Coppellotti O, Dei D and Roncucci G 2006 Photodynamic therapy in the treatment of microbial infections: basic principles and perspective applications *Lasers Surg. Med.* **38** 468–81
- Juzeniene A and Moan J 2007 The history of PDT in Norway. Part one—identification of basic mechanisms of general PDT *Photodiag. Photodyn. Ther.* **4** 3–11
- Kazor C E, Mitchell P M, Lee A M, Stokes L N, Loesche W J, Dewhurst F E and Paster B J 2003 Diversity of bacterial populations on the tongue dorsa of patients with halitosis and healthy patients *J. Clin. Microbiol.* **41** 558–63
- Komerik N, Nakanishi H, MacRobert A J, Henderson B, Speight P and Wilson M 2003 *In vivo* killing of *Porphyromonas gingivalis* by toluidine blue-mediated photosensitization in an animal model *Antimicrob. Agents Chemother.* **47** 932–40
- Komerik N, Wilson M and Poole S 2000 The effects of photodynamic on two virulence factors of Gram negative bacteria *Photochem. Photobiol.* **72** 676–80
- Konig K, Teschke M, Sigusch B, Glockmann E, Eick S and Pfister W 2000 Red light kills bacteria via photodynamic action *Cell. Mol. Biol.* **46** 1297–303
- Krespi Y P, Shrome M G and Kacker A 2006 The relationship between oral malodor and volatile sulfur compound-producing bacteria *Otolaryngol. Head Neck Surg.* **135** 671–6
- Lopez-Jimenez L, Fuste E, Martinez-Garigga B, Arnabat-Dominguez J, Vinuesa T and Vinas M 2015 Effects of photodynamic therapy on *Enterococcus faecalis* biofilms *Lasers Med. Sci.* **30** 1519–26
- Lopes R G et al 2014 Photodynamic therapy as novel treatment for halitosis in adolescents: a case series study *J. Lasers Med. Sci.* **5** 146–52
- Maisch T, Szeimies R M, Jori G and Abels C 2004 Antibacterial photodynamic therapy in dermatology *Photochem. Photobiol. Sci.* **3** 907–17
- Morita M and Wang H L 2001 Association between oral malodor and adult periodontitis: a review *J. Clin. Periodontol.* **28** 813–9
- Nagata J Y, Hioka N, Kimura E, Batistela V R, Terada R S S, Graciano A X, Baesso M L and Hayacibara M F 2012 Antibacterial photodynamic therapy for dental caries: evaluation of the photosensitizers used and light source properties *Photodiag. Photodyn. Ther.* **9** 122–31
- Persson S 1990 The formation of hydrogen sulphide and methyl mercaptan by oral bacteria *Oral Microbiol. Immunol.* **5** 195–201
- Pfitzner A, Sigusch B W, Albrecht V and Glockmann E 2004 Killing of periodontopathogenic bacteria by photodynamic therapy *J. Periodontol.* **75** 1343–9
- Porter S R and Scully C 2006 Oral malodour *BMJ* **333** 632–5
- Quirynen M, Dadamio J, Van den Velde S, De Smit M, Dekeyser C, Van Tornout M and Vandekerckhove B 2009 Characteristics of 2000 patients who visited a halitosis clinic *J. Clin. Periodontol.* **36** 970–5
- Raghavendra M, Koregol A and Bhola S 2009 Photodynamic therapy: a targeted therapy in periodontics *Aust. Dent. J.* **54** S102–9
- Rajesh S, Koshi E, Philip K and Mohan A 2011 Antimicrobial photodynamic therapy: an overview *J. Indian Soc. Periodontol.* **15** 323–7
- Robertson C A, Hawkins Evans D and Abrahamse H 2009 Photodynamic therapy (PDT): a short review on cellular mechanisms and cancer research applications for PDT *J. Photochem. Photobiol. B* **96** 1–8

- Schiffner U, Cachovan G, Bastian J, Sculean A and Eick S 2014 *In vitro* activity of photoactivated disinfection using a diode laser in infected root canals *Acta Odontol. Scand.* **72** 673–80
- Sharma M, Visai L, Bragheri F, Cristiani I, Gupta P K and Speziale P 2008 Toluidine blue-mediated photodynamic effects on *Staphylococcal* biofilms *Antimicrob. Agents Chemother.* **52** 299–305
- Soukos N S et al 2006 Photodynamic therapy for endodontic disinfection *J. Endod.* **32** 979–84
- Soukos N S, Som S, Abernethy A D, Ruggiero K, Dunham J, Lee C, Doukas A G and Goodson J M 2005 Phototargeting oral black-pigmented bacteria *Antimicrob. Agents Chemother.* **49** 1391–6
- Tanabe S and Grenier D 2012 Characterization of volatile sulfur compound production by *Solobacterium moorei* *Arch. Oral Biol.* **57** 1639–43
- Taylor M N and Gonzalez M 2009 The practicalities of photodynamic therapy in acne vulgaris *Br. J. Dermatol.* **160** 1140–8
- Trachtenberg J et al 2008 Vascular-targeted photodynamic therapy (padoporphin, WST09) for recurrent prostate cancer after failure of external beam radiotherapy: a study of escalating light doses *BJU Int.* **102** 556–62
- Tyrell K L, Citron D M, Warren Y A, Nachnani S and Goldstein E J 2003 Anaerobic bacteria cultured from the tongue dorsum of subjects with oral malodor *Anaerobe* **9** 243–6
- Venezio F R et al 1985 Bactericidal effects of photoradiation therapy with hematoporphyrin derivative *J. Infect. Dis.* **151** 166–9
- Williams J A, Pearson G J and Colles M J 2006 Antibacterial action of photoactivated disinfection (PAD) used on endodontic bacteria in planktonic suspension and in artificial and human root canals *J. Dent.* **34** 363–71
- Wilson M 2004 Lethal photosensitisation of oral bacteria and its potential application in the photodynamic therapy of oral infections *Photochem. Photobiol. Sci.* **3** 412–8