Manuscript Number: MBE-15-72R3

Title: Thermodynamics-based Metabolite Sensitivity Analysis in Metabolic Networks

Article Type: Regular Article

Keywords: TMFA, TFBA, GSA, Metabolic Modeling, FBA, DoE

Corresponding Author: Dr. Vasileios Chatzimanikatis,

Corresponding Author's Institution: EPFL

First Author: Alexandros Kiparissides

Order of Authors: Alexandros Kiparissides; Vasileios Chatzimanikatis

Abstract: The increasing availability of large metabolomics datasets enhances the need for computational methodologies that can organize the data in a way that can lead to the inference of meaningful relationships. Knowledge of the metabolic state of a cell and how it responds to various stimuli and extracellular conditions can offer significant insight in the regulatory functions and how to manipulate them. Constraint based methods, such as Flux Balance Analysis (FBA) and Thermodynamics-based flux analysis (TFA), are commonly used to estimate the flow of metabolites through genome-wide metabolic networks, making it possible to identify the ranges of flux values that are consistent with the studied physiological and thermodynamic conditions. However, unless key intracellular fluxes and metabolite concentrations are known, constraint-based models lead to underdetermined problem formulations. This lack of information propagates as uncertainty in the estimation of fluxes and basic reaction properties such as the determination of reaction directionalities. Therefore, knowledge of which metabolites, if measured, would contribute the most to reducing this uncertainty can significantly improve our ability to define the internal state of the cell. In the present work we combine constraint based modeling, Design of Experiments (DoE) and Global Sensitivity Analysis (GSA) into the Thermodynamics-based Metabolite Sensitivity Analysis (TMSA) method. TMSA ranks metabolites comprising a metabolic network based on their ability to constrain the gamut of possible solutions to a limited, thermodynamically consistent set of internal states. TMSA is modular and can be applied to a single reaction, a metabolic pathway or an entire metabolic network. This is, to our knowledge, the first attempt to use metabolic modeling in order to provide a significance ranking of metabolites to guide experimental measurements.

Thermodynamics-based Metabolite Sensitivity Analysis in Metabolic Networks

A.Kiparissides^{1,2}, V. Hatzimanikatis^{1*}

1, Laboratory of Computational Systems Biotechnology
École Polytechnique Fédérale de Lausanne (EPFL)
CH-1015, Lausanne, Switzerland

2, Department of Biochemical Engineering
University College London
WC1E 6BT, London, UK

Highlights

- A method to analyze uncertainty in the thermodynamic properties of metabolic networks
- Rank metabolites on their ability to reduce thermodynamic uncertainty in a network
- Aids the design of informative experiments and the organization of existing datasets
- Modular approach, can be applied to a single reaction, pathway or entire network
- Insight on thermodynamic interactions; evaluate metabolic engineering strategies

Thermodynamics-based Metabolite Sensitivity Analysis in Metabolic Networks

A.Kiparissides^{1,2}, V. Hatzimanikatis^{1*}

1, Laboratory of Computational Systems Biotechnology
École Polytechnique Fédérale de Lausanne (EPFL)
CH-1015, Lausanne, Switzerland
2, Department of Biochemical Engineering
University College London
WC1E 6BT, London, UK

***Corresponding Author:** **Prof. Vassily Hatzimanikatis**

Laboratory of Computational Systems Biotechnology

École polytechnique fédérale de Lausanne

Lausanne, CH-1015

Switzerland

Tel: +41 (0)21 693 98 70

Email: vassily.hatzimanikatis@epfl.ch

Thermodynamics-based Metabolite Sensitivity Analysis in Metabolic Networks

A.Kiparissides^{1,2}, V. Hatzimanikatis^{1*}

1, Laboratory of Computational Systems Biotechnology
École Polytechnique Fédérale de Lausanne (EPFL)
CH-1015, Lausanne, Switzerland
2, Department of Biochemical Engineering
University College London
WC1E 6BT, London, UK

Abstract

The increasing availability of large metabolomics datasets enhances the need for computational methodologies that can organize the data in a way that can lead to the inference of meaningful relationships. Knowledge of the metabolic state of a cell and how it responds to various stimuli and extracellular conditions can offer significant insight in the regulatory functions and how to manipulate them. Constraint based methods, such as Flux Balance Analysis (FBA) and Thermodynamics-based flux analysis (TFA), are commonly used to estimate the flow of metabolites through genome-wide metabolic networks, making it possible to identify the ranges of flux values that are consistent with the studied physiological and thermodynamic conditions. However, unless key intracellular fluxes and metabolite concentrations are known, constraint-based models lead to underdetermined problem formulations. This lack of information propagates as uncertainty in the estimation of fluxes and basic reaction properties such as the determination of reaction directionalities. Therefore, knowledge of which metabolites, if measured, would contribute the most to reducing this uncertainty can significantly improve our ability to define the internal state of the cell. In the present work we combine constraint based modeling, Design of Experiments (DoE) and Global Sensitivity Analysis (GSA) into the Thermodynamics-based Metabolite Sensitivity Analysis (TMSA) method. TMSA ranks metabolites comprising a metabolic network based on their ability to constrain the gamut of possible solutions to a limited, thermodynamically consistent set of internal states. TMSA is modular and can be applied to a single reaction, a metabolic pathway or an entire metabolic network. This is, to our knowledge, the first attempt to use metabolic modeling in order to provide a significance ranking of metabolites to guide experimental measurements.

1. Introduction

The development of new modeling methods[1] has enabled the formulation of genome-scale kinetic models which are consistent with stoichiometric, thermodynamic and physiological constraints [2-5]. Constraint based methods, such as Flux Balance Analysis (FBA), are commonly used to estimate the flow of metabolites through such metabolic networks, making it possible to identify the ranges of flux values that are consistent with the studied conditions [6-8]. In the absence of explicit experimental information for key intracellular reactions, a situation commonly encountered when considering a new organism or strain, fluxes are allowed to vary between physiologically relevant bounds, often derived from literature, which leads to underdetermined problem formulations. The lack of explicit experimental information propagates as uncertainty in the estimated Gibbs free energy of reaction, $\Delta_r G'$, and consequently in the determination of reaction directionalities which in turn constrain the flux profiles and the thermodynamically feasible concentration ranges [5, 9]. This results in the existence of numerous alternative internal flux distributions that satisfy the model's constraints and can achieve the same optimum [8, 10, 11]. The size of the resulting solution space, comprising the gamut of alternate optimal flux distributions, can be viewed as a representation of the uncertainty in predicting an exact intracellular state relevant to the studied physiology [12, 13]. Therefore knowledge of which metabolites, if measured, would contribute the most to reducing this uncertainty can significantly improve our ability to define the internal state of a cell.

The space of steady state flux solutions has attracted significant scientific interest in a number of studies that either attempted to attribute biological significance to characteristics of the solution space [14, 15] or attempted to achieve more accurate flux distributions by incorporating additional physicochemical information in the form of constraints based on thermodynamic properties [16], molecular crowding considerations [17] and carbon labeling information [18-20]. However the problem of identifying network components with a significant contribution towards the observed uncertainty remains far from trivial [21]. Approaches that have been employed to study the sensitivity of FBA outputs include one-factor-a-time numerical perturbation simulations [22-24] or derivative based local measures [25, 26], which almost exclusively study the effect of the model constituents on the value of the objective function [18, 19, 21]. Whilst the proposed approaches have provided valuable qualitative information, they have not been able to

provide a functional correlation with metabolites that exert strong influence on the qualitative and quantitative behavior of a particular reaction (or a subset of reactions).

Thermodynamics-based flux analysis (TFA)[5], also referred to as thermodynamics-based metabolic flux analysis (TMFA)[16] and thermodynamics-based flux balance analysis (TFBA)[9, 11] was introduced by Hatzimanikatis and colleagues and allows integration of metabolomics data into FBA models[5, 11]. Apart from enforcing compliance with the second law of thermodynamics, TFA effectively reduces the size of the solution space by selecting only the thermodynamically feasible sets of the following biochemical and thermodynamic variables: reaction directionalities, net fluxes through a reaction, $\Delta_r G'$ and metabolite concentrations (activities). The permissible ranges of these biochemical variables can be explored within TFA using linear optimization in a methodology defined as Thermodynamic Variability Analysis (TVA) in [9, 16]. In TVA, the ranges for the activity of each metabolite, the $\Delta_r G'$ and the net flux of each reaction are estimated through their minimization and maximization subject to the thermodynamic and mass balance constraints. Consequently we can estimate whether a reaction is thermodynamically reversible under the studied physiological conditions and whether it operates near or far from thermodynamic equilibrium.

Knowledge of the metabolite concentrations for all the products and substrates of a reaction allows the exact estimation of its thermodynamic properties such as $\Delta_r G'$ and displacement from thermodynamic equilibrium (denoted as Γ [27]). The uncertainty in the exact values for metabolite concentrations, either due to experimental variability or due to lack of experimental data, propagates through the thermodynamic constraints introduced by TFA to uncertainty in the thermodynamic properties of the reactions comprising a metabolic network. In the present work we propose a new framework that combines TFA, Design of Experiments (DoE) and Global Sensitivity Analysis (GSA) methods in order to determine and quantify how variability in metabolite concentrations propagates to uncertainty in the thermodynamic properties of any reaction within a metabolic network. TMSA ranks the various metabolites according to their ability to reduce variability in the thermodynamic properties of reactions, and thus it enables the targeted reduction of uncertainty [9, 16]. The metabolites with the largest contribution to the observed uncertainty can be considered as high priority targets for experimental analysis. Consequently, TMSA can fully exploit and even upgrade the value of metabolomics data, an

important, genome-scale source of experimental information [10]. Moreover, quantification of the propagation of uncertainty from metabolite concentrations to systemic properties can be used for the design of metabolic engineering strategies and the formulation of hypotheses in many problems related to metabolism.

2. Methods

Below we present an overview of the basic concepts of the proposed methodology using a simple example before presenting each step in greater detail. Let's consider the case of a reversible reaction involving two substrates, which react towards two products:

The $\Delta_r G'$ of this reaction can be estimated[16] as:

$$\Delta_r G' = \{\Delta_f G'_C + \Delta_f G'_D - \Delta_f G'_A - \Delta_f G'_B\} + RT \ln(C \cdot D) - RT \ln(A \cdot B) \quad (2)$$

where $(\Delta_f G'_i)$ denotes the formation energy of the participating species, R is the universal gas constant and T is the absolute temperature. The formation energy terms $(\Delta_f G'_i)$ can be retrieved from relevant literature or estimated using appropriate computational methods [28]. Thus the only unknowns left in equation (2) are the concentrations of the four metabolite species participating in the reaction (A, B, C and D). We want to examine whether it is necessary to measure all four concentrations in order to define the $\Delta_r G'$ for reaction (1) within an acceptable uncertainty range. Additionally, if we can achieve the desired accuracy with fewer measurements, we want to know the order in which the participating metabolites should be measured.

Intuitively one would start by testing how different concentrations of the four participating metabolite species (A, B, C and D) affect the $\Delta_r G'$ of reaction (1). This concept can be formalized by considering it in an *uncertainty analysis* setting. In this context, the concentrations of the metabolite species can be considered as uncertain input factors (X) with a predefined range while the thermodynamic properties of reaction (1) will constitute the studied output variables (R_T). If D_T defines the variance observed in the studied output,

we are interested in estimating the fractional contributions of the variability of metabolites i towards the variance of the observed output (D_T), which are defined as *fractional variances* d_A , d_B , d_C and d_D , such that:

$$D_T = d_A + d_B + d_C + d_D \quad (3).$$

Therefore the experimental quantification of metabolite species with higher fractional variance, d_i , can be prioritized over metabolites with a lesser effect.

For single reactions or small subsets of reactions (where the total number of metabolites is $\sim < 4$) the effects of metabolite concentrations on the studied outputs can be assessed through a small set of randomized or heuristic trial and error model evaluations. However for larger subsets of reactions, pathways or genome scale models the effects of metabolite uncertainty, i.e. the fractional variances of the metabolites, propagate in a non-linear manner. Therefore we need a robust model analysis method able to identify, quantify and analyze such network-level, non-linear effects. Sensitivity Analysis, and in particular Global Sensitivity Analysis (GSA), methods are frequently the methods of choice when studying complex non-linear models due to their ability to differentiate between first and higher order (non-linear) sensitivity [29, 30]. Current state-of-the-art Global Sensitivity Analysis (GSA) methods are not directly applicable in a TFA setting due to difficulties in the implementation of commonly used numerical techniques, such as the estimation of high dimensional integrals through their Monte Carlo approximation (see Supplemental Material). In order to circumvent this limitation we developed our method based on the methodology introduced by Kiparissides et al [31] that employs Design of (*in silico*) Experiments (DoE) for the estimation of first and higher-order sensitivity measures.

Instead of measuring the effect of one-metabolite-at-a-time we use an experimental design to study how different combinations of concentration values affect the thermodynamic properties of reaction (1). In the case of the simple one-reaction example considered presently, it is possible to calculate the thermodynamic properties of reaction (1) for all the combinations defined by the full factorial. This will amount to a total of L^n evaluations of equation (2) where L is the number of concentration levels each metabolite will be considered at, while n is the number of metabolites considered. Considering, for

example, each metabolite at three distinct concentration levels will result in 3^4 possible combinations, or *in-silico* experiments, and an equal number of calculated values for the thermodynamic properties of reaction (3), or measured responses (R_T). Following the Response Surface Methodology (RSM) introduced by Box and Wilson [32], we can use a second order polynomial to describe the behavior of the response to the changes in the input factors. While a full quadratic model would be redundant in the simple case discussed, we present it for the sake of completion and continuity:

$$\begin{aligned}
 R_T = & \beta_0 + \beta_A A + \beta_B B + \beta_C C + \beta_D D \\
 & + \beta_{AB} AB + \beta_{AC} AC + \beta_{AD} AD + \beta_{BC} BC + \beta_{BD} BD + \beta_{CD} CD \\
 & + \beta_{AA} A^2 + \beta_{BB} B^2 + \beta_{CC} C^2 + \beta_{DD} D^2
 \end{aligned} \tag{4}$$

The multipliers of the first order (β_i) and quadratic terms (β_{ii}) of appropriately estimated RSM meta-models have been shown to approximate first and higher-order global sensitivity indices respectively [31]. This essentially means that we can study the multipliers of the derived RSM in order to identify which metabolite exerts a stronger influence on the behavior of the studied output R_T . For example, if β_A is bigger than the rest first order terms, then the uncertainty (variability) in the value of metabolite A contributes the most into the uncertainty (variance) of the studied output R_T . Similarly, if β_{AB} is near zero then the propagation of uncertainty (variability) in the value of metabolite A is independent of the value of metabolite B. While the example presented thus far captures the basic concepts of the proposed methodology, it does not highlight, due to its simplicity, the complexities that arise when considering even a single pair of uncertain metabolites in the context of an entire metabolic network.

Figure 1.

Figure 1 illustrates the various types of correlation that can arise between two uncertain metabolites, assuming that all other metabolites comprising the network are defined. A purely linear relationship between the metabolites and the measured response (R_T) (Figure 1a) implies no interaction between the metabolites, while a bilinear relationship implies a coupling that can have either an impeding or a compounding effect on R_T (Figures 1b and 1c respectively). Higher order interactions can arise when considering metabolites that partake and affect multiple reactions across the network such as co-factors (Figure 1d).

While constraint based TFA metabolic models are linear by nature, the cumulative effect of multiple linear constraints intersecting within a high dimensional (solution) space may give rise to piecewise-linear edges or cross-sections. During the estimation of an RSM such piecewise-linear elements are approximated by the higher order terms of the polynomial.

In the following we will show that the selection of both the experimental design levels and the studied output is not as straightforward as was presented in the simple example above. We will present the algorithm for Thermodynamics-based Metabolite Sensitivity Analysis (TMSA) in detail and discuss how to formulate the problem in the context of studying the allocation of thermodynamic uncertainty within a metabolic network. In the following we assume the existence of a thermodynamically curated metabolic model and the application of TFA. A description of TFA is given in the Supplemental Material of the present work and in [9, 16]. Metabolite concentrations are treated as uncertain input factors (X) with quantifiable uncertainty, which is defined by their thermodynamically permissible range. The algorithm consists of a several steps that are discussed in the following subsections and summarized in Figure 2.

Figure 2.

2.1 Model Calibration

Initially, all available experimental measurements are integrated in the model through the appropriate constraints. These can include measured uptake/secretion rates, estimated intracellular fluxes, and extra- and intracellular metabolite concentration values and ranges. In the absence of explicit experimental information, generalized upper and lower bounds found from relevant literature can be used. Performing a TVA analysis will determine the permissible ranges for all model variables under the set of physiological conditions considered. Reaction $\Delta_r G'$ values and metabolite concentrations that are thermodynamically feasible in a very tight range of values should be identified during this step. Such tightly constrained variables are assumed to be constant under the studied physiological conditions and are not considered further in subsequent steps.

2.2 Construction of experimental design levels

In the present context, experiments are defined as model simulations, hence the term *in silico*. Following the DoE methodology each uncertain input factor is assigned a pre-

defined amount of levels within the range of values that will be considered. In semblance to the process of designing wet-lab experiments, there are several ways of designing and selecting these levels. Ultimately, the selection of the appropriate design levels is defined by the question one wishes to investigate. Some of the possible definitions, in the context of analyzing uncertainty in the thermodynamic properties of a metabolic network, are indicatively discussed in the supplemental material (section 3). For the purposes of the case studies considered below the thermodynamically feasible range for each metabolite, defined by TVA, was discretized into a number of equal intervals. Each design level was allowed to include several of these intervals (supplemental material, section 4) and the resulting RSM provided a functional relationship describing how constraining metabolites to a certain range of concentration values affects the thermodynamic properties of the network. If we take into consideration the fact that most intracellular measurements include a considerable error range [33], this type of design level seems more realistic compared to other options (Supplemental Material). Additionally, it is not affected by bias or errors introduced through computational methods or analytical techniques.

2.3 Generation of experimental design and model evaluation

Methodical design of the conditions of the set of experiments is essential for efficient and effective information gathering. The goal is to select the levels (values) of the uncertain input factors (X), following a pre-specified design, in order to maximize the amount of information gained whilst minimizing the amount of conducted experiments. A full factorial design explores all the combinations of every factor at each possible factor level. Depending on the number of levels input factors are studied at, such an approach requires L^n experiments, where n signifies the number of factors and the base number (L) denotes the number of levels the factors are studied at. The number of total experiments scales badly with dimension. In order to reduce the number of overall experiments to be performed, one usually resorts to the use of fractional factorial designs at the cost of less accuracy in the computed RSM [34, 35]. In the context of studying metabolic networks with TFA, each experiment, e , defined by the DoE can be implemented *in silico* by adjusting the bounds of the relevant model constraints. Subsequently the studied model output for each experiment, R_T^e , can be calculated through a model evaluation or TVA.

2.4 Quantification of Thermodynamic Uncertainty in Metabolic Networks

In order to study how metabolites contribute to and affect output uncertainty in the context of a metabolic network, either qualitatively or quantitatively, it is necessary to define an appropriate metric to characterize the output uncertainty (R_T). Similar to the step described in section 2.2, many alternative performance metrics can be conceived and implemented each analyzing the thermodynamic uncertainty in metabolic networks from a different perspective (Supplemental Material, section 5).

From equation (2) we can easily deduce that allowing a metabolite concentration (M_i) to vary between M_{min} and M_{max} will impose a range on the feasible $\Delta_r G'$ values for the reactions in which (M_i) participates. The actual value of $\Delta_r G'$ (which also affects the net flux and defines Γ) under the studied physiological conditions can be anywhere within this imposed $\Delta_r G'$ range depending on the exact value of (M_i). In the context of the present work we term this range as *reaction flexibility* (F_R) and define it as:

$$F_{R,j} = \left| \Delta_r G'^{M_i, \max}_j - \Delta_r G'^{M_i, \min}_j \right| \quad (5)$$

The same rationale can be generalized to consider a subset of reactions or even an entire metabolic network by defining the *total reaction flexibility* (F_{TOT}) as:

$$F_{TOT} = \sum_{j=1}^{\#rxns} \left| \Delta_r G'^{M_i, \max}_j - \Delta_r G'^{M_i, \min}_j \right| \quad (6).$$

In the remainder, total reaction flexibility (F_{TOT} , Eq. 6) will be used as the studied experimental output (R_T) unless stated otherwise. In order to study the allocation of uncertainty within a particular pathway of interest or any subset of reactions comprising the metabolic network we need to estimate the fraction of total network uncertainty (F'_{TOT}) that is associated with the reactions of interest. For example, Equation 6 can be modified to estimate the total thermodynamic uncertainty of the TCA cycle reactions only, yielding:

$$F'_{TOT} = \sum_{j=1}^{\#TCA_rxns} \left| \Delta_r G'^{M_i, \max}_j - \Delta_r G'^{M_i, \min}_j \right| \quad (7)$$

Consequently, we can calculate an RSM that describes the behavior of F'_{TOT} as a function of the concentration of the studied metabolites. Following the same rationale, equations similar to equation 7 can be defined for any metabolic pathway or reaction of interest. Since F_{TOT} (Eq. 6) is an additive function, once a sensitivity analysis of total network flexibility (F_{TOT}) is available, additional analyses for any subset of reactions (F'_{TOT}) can be generated at no additional computational cost. Alternatively, if an analysis of total network flexibility is not available, we only need to conduct TVA for the reactions of interest during the model evaluation step which significantly reduces the computational cost.

2.5 Response Surface Model (RSM) and Sensitivity Indices

In the context of the present work we are interested to find the subset of input factors (X) with the largest impact on the studied output (R_T). This is achieved by the introduction of an intermediate step involving the approximation of the response of the original metabolic model (R_T) with a simplified meta-model in the form of a RSM. The aim herein is two-fold; the first order coefficients of the calculated RSM are used as first order sensitivity indices while the quadratic terms provide information on the quantifiable higher-order sensitivities. The estimated sensitivities can be used to rank metabolites based on their ability to reduce the thermodynamic uncertainty of the entire network. Additionally, the derived meta-model can be further analyzed with GSA in order to provide detailed global sensitivity information at a significantly reduced computational cost [31]. The complexity (and fidelity) of the calculated RSM (linear or quadratic, with or without 2-factor interaction terms) is ultimately defined by the number of metabolites considered and the number of *in silico* experiments conducted.

2.6 Quantification of Uncertainty Reduction (Q-UR).

Knowing which input factors (X) and to what extent have a significant effect on the studied output (R_T) allows for the targeted reduction of uncertainty through tailor made experiments. It is possible to simulate and characterize the effect of experimentally measuring the (r) highest-ranking metabolites on the studied output (R_T) using *in silico* studies alone and without the need for “wet-lab” measurements. To achieve this, the full factorial comprising all the L^r possible DoE level combinations needs to be explored. This will yield a set of L^r simulated values for the studied output (R_T^{Q-UR}), which can be characterized

through standard statistical metrics. The achieved level of reduction in output uncertainty can be readily estimated through a simple relationship of the form:

$$\Delta R_T = \frac{R_T^* - R_T^{Q-UR}}{R_T^*} \quad (8)$$

where R_T^{Q-UR} denotes the value of the studied output (R_T) for each Q-UR simulation and R_T^* denotes the reference value for the studied output (R_T). The reference state R_T^* is defined by allowing all *uncertain* metabolites to vary freely within their thermodynamically permissible range. A high value for ΔR_T denotes a high degree of reduction in total network uncertainty. Q-UR effectively constitutes a cost-benefit analysis that can guide the decision regarding *how many and which metabolites should be experimentally measured in order to achieve a given degree of reduction in output uncertainty*.

3. Results & Discussion

TMSA was applied on a reduced model of central carbon metabolism in *E. coli*. The model has been derived from the iJO1366 genome scale reconstruction by Palsson and coworkers [36] using the minimal set of reactions and metabolites required to link the metabolic core subsystems: glycolysis, pentose-phosphate pathway, TCA cycle, pyruvate metabolism and glyoxylate cycle[37]. The final reduced metabolic network also includes the complete electron transport chain and all anaplerotic reactions. A variety of simulations considering both aerobic and anaerobic conditions were performed and are discussed below. Experimental information for metabolite exchange rates and *E. coli* growth rates under both aerobic and anaerobic conditions (Supplemental Material) were taken from Chen et al. [18] and McCloskey et al [33]. Intracellular metabolite concentrations were obtained from McCloskey et al [33].

Generally, for any metabolic network or organism, metabolites can be divided into three categories: (a) uncertain metabolites that can be experimentally measured, (b) experimentally measured metabolites with a given measurement error and (c) uncertain metabolites that can't/won't be measured experimentally. Depending on the complexity and the novelty of the studied metabolic network various combinations of the above

categories can be expected. Taking a well-studied organism, like *E. coli*, as an example one would expect to consider metabolites belonging in each of the three categories presented above. However, when dealing with a new organism or strain information such as past experimental data and which metabolites are challenging to measure might not be readily available. In the following sections (3.1 to 3.4) we consider a number of case studies exploring the most commonly encountered scenarios and discuss how the application of the method and the interpretation of the results are affected. Table 1 provides an overview of the distribution of metabolites in each of the above categories for the case studies discussed in the remainder.

Table 1 Classification of metabolites for the case studies discussed in sections 3.1-3.4

Number of Intracellular Metabolites	Aerobic Conditions			
	3.1	3.2	3.3	3.4
in category (a)	78	57	51	51
in category (b)	-	21	21	21
in category (c)	-	-	6	6
Total	78	78	78	78
Number of Intracellular Metabolites	Anaerobic Conditions			
	3.1	3.2	3.3	3.4
in category (a)	78	60	54	54
in category (b)	-	18	18	18
in category (c)	-	-	6	6
Total	78	78	78	78

3.1 Metabolite Sensitivity Analysis with no *a priori* information

In the first case study, discussed in the present section, we emulated the analysis of a new system or organism by assuming a complete lack of intracellular experimental data. Therefore, all metabolites comprising the reduced *E. coli* network were allocated in the first category (a) of the classification presented above.

In the absence of *a priori* information metabolite concentrations were allowed to vary between widely defined upper and lower bounds in order to ensure all physiological conditions are taken into consideration [16]. TVA was performed and metabolites whose

thermodynamically permissible range was found to be smaller than a predefined tolerance (a 1mM range was used as the cutoff) were assumed to be fixed and were excluded from subsequent analysis. Moreover, due to the assumed lack of intracellular experimental information, DoE levels were defined based on a discretization of the thermodynamically permissible range for each metabolite (Supplemental Material, section 4). Subsequently, a 3-level Box-Behnken [35] experimental design to study the uncertain intracellular metabolites (X) was generated. TVA was performed for each *in silico* experiment defined by the DoE and F_{TOT} (Eq. 6) was estimated as the monitored output (R_T). Given the lack of *a priori* knowledge of the system, the choice of output metric (F_{TOT} , Eq. 6) was dictated by the need to identify metabolites that can yield information rich measurements with respect to defining the internal state of the cell. Finally, a quadratic response surface model (RSM) with interaction terms describing the functional dependence of F_{TOT} on the concentration range of the analyzed metabolites was estimated. As has been previously discussed, the RSM coefficients of the main effect terms (linear, pure quadratic) can be used as sensitivity indices [31].

Figure 3.

Analysis of the estimated coefficients revealed that higher-order sensitivity indices were able to distinguish the significance of metabolites at a higher resolution than their first order counterparts (Supplemental Figure 1). Interestingly, this effect was less pronounced under anaerobic conditions, an indication that network thermodynamics are more rigid under anoxic conditions. Pyruvate was found to have the strongest effect on F_{TOT} under both conditions (Figures 3a & b). Given the role of pyruvate in central carbon metabolism and its high connectivity (pyruvate participates in 19 out of the 146 reactions in the studied network) its classification, as the most influential metabolite is, to a certain extent, expected. In the presence of oxygen network uncertainty was found to be primarily affected by energy and redox related metabolites, while in the absence of oxygen, electron transport chain (ETC) and overflow-related metabolites were the major contributors to F_{TOT} variability (Table 2). It is therefore worth noting that the significance rankings generated for aerobic and anaerobic conditions reflect many of the key differences between fermentative and respiratory metabolism reported in relevant literature [18, 38, 39].

Subsequently we performed a Q-UR analysis, detailed above, to validate whether the estimated metabolite significance rankings provided a significant benefit in terms of

efficiently selecting metabolites that reduce uncertainty in $\Delta_r G'$. Towards this end, the potential of the 5 highest-ranking metabolites to reduce uncertainty was explored and compared against sets of 5 metabolites selected randomly. The full factorial accounting for all possible DoE level combinations of the 5 selected metabolites resulted in 32 model evaluations. The evaluation of $F_{TOT}^{Q\ UR}$ for all the combinations of the full factorial was repeated 100 times (3200 model evaluations in total) and kernel smoothed probability distribution functions of the achieved reduction in F_{TOT} (Equation 8) were evaluated (Figures 4a & b). Next, 5000 random sets of 5 metabolites to be constrained at-a-time were generated. Each metabolite within these 5000 sets was assigned a random concentration value and the resulting $F_{TOT}^{validation}$ was estimated.

Table 2 Ranking of most significant metabolites according to higher-order sensitivities when no *a priori* information is available.

Rank	Metabolites	
	Aerobic	Anaerobic
1	Pyruvate	Pyruvate
2	Phosphate	Menaquinol 8
3	NAD	D-Lactate
4	CO ₂	Menaquinone 8
5	NADH	Phosphate

Interestingly, the selection of high-ranking metabolites by the proposed method proved to be significantly more efficient in reducing F_{TOT} over an uninformed or random selection (Figures 4a & b). The benefits of the proposed method were less pronounced in the absence of oxygen which could be attributed to the fact that network thermodynamics are more constrained in anoxic conditions with $F_{TOT,Anaerobic}$ measuring up to just 65% of $F_{TOT,Aerobic}$.

Figure 4.

3.2 Metabolite Sensitivity Analysis with pre-existing experimental data

The second case study, discussed below, emulates the existence of partial experimental data by taking into account the metabolomics data from McCloskey et al [33]. Therefore, metabolites from the experimental dataset that were also part of the reduced *E. coli* network (Supplemental Material) were allocated in the second category (b) while all remaining metabolites were allocated in the first category (a) of the classification presented previously.

Experimentally measured metabolites were integrated in the model through the relevant constraints while taking into account an average experimental error as described in Supplemental Material. The concentrations for all remaining metabolites were allowed to vary between widely defined upper and lower bounds in order to ensure all physiological conditions are taken into consideration [16]. Following a similar procedure to that of section 3.1, TVA was performed and metabolites whose thermodynamically permissible range was found to be smaller than a predefined tolerance (a 1mM range was used as the cutoff) were assumed to be fixed and were excluded from subsequent analysis. The available experimental measurements alone were not enough to derive probability distribution functions for the concentrations of the measured metabolites. Therefore, DoE levels were defined based on a discretization of the thermodynamically permissible range for each metabolite (Supplemental Material). The remaining steps of the analysis (DoE generation, selection of output metric, estimation of the RSM, Q-UR and validation) followed the procedure already discussed in section 3.1.

The metabolite significance rankings were found to be significantly different from the ones computed in section 3.1, particularly in the case of aerobic conditions. Pyruvate remained in the 5 highest-ranking metabolites under both physiological conditions examined, however phosphate concentration became the highest contributor to F_{TOT} variability (Figures 3c & d). The integration of the experimental dataset during the model-preprocessing step constrained the concentration of many central carbon metabolites and as a result reduced the variability in many central carbon reactions diminishing the potential of pyruvate to strongly affect F_{TOT} (Supplemental Tables 2, 5 & 6).

Furthermore the availability of intracellular measurements for energy and redox metabolites greatly reduced the range of concentrations within which they can vary and consequently their potential effect on F_{TOT} variability. A direct comparison of the generated metabolite rankings with (Figures 3c & d, Supplemental Figure 2) and without (Figures 3a &

b, Supplemental Figure 1) the inclusion of experimental information reveals that the observed changes are not a mere reordering of the priority list (to account for the measured metabolites) but rather a complete reallocation of uncertainty that accounts for the newly integrated physiological conditions. Our Q-UR analysis (Figures 4c & d) indicated that TMSA is both sensitive to varying physiological conditions and more efficient in reducing output uncertainty making it a superior choice over previous attempts to prioritize metabolomics measurements using constraint-based models [25, 33].

3.3 Metabolite Sensitivity Analysis with excluded metabolites

In the third case study we assume that ETC metabolites, as an example, are challenging to measure experimentally due to special extraction and purification requirements prior to their quantification [38]. Therefore, metabolites of the ETC were allocated in category (c) of the classification presented above whilst all other metabolites were allocated in categories (a) and (b) accordingly as discussed above.

For the model calibration step, the procedures described in section 3.2 were repeated. However, apart from metabolites whose thermodynamically permissible range was found to be smaller than a predefined tolerance by TVA, the metabolites which are uncertain but can't be measured (category c) were also excluded from subsequent analysis. A key difference with the analysis performed in section 3.2 is that the ETC metabolites that were excluded from the analysis are still allowed to vary within their predefined bounds. The remaining steps of the analysis (DoE level selection, DoE generation, selection of output metric, estimation of the RSM, Q-UR and validation) followed the procedure already discussed in sections 3.1 and 3.2.

It is worth noting that total network flexibility (F_{TOT}) is not affected by the exclusion of the ETC metabolites from the analysis (Supplemental Table 2); we merely don't account for the uncertainty caused by the metabolites of category (c). Therefore, one would intuitively expect the significance rankings between the present case study and that of section (3.2) to differ only in the excluded metabolites. Interestingly, when we repeated the analysis the significance ranking was found to be both qualitatively and quantitatively different (Figures 3e & f, Figures 4e & f, Supplemental Figure 3). This indicates that the allocation of uncertainty in metabolic networks is not straightforward and requires the use of analysis techniques with adequate discriminating power.

The informed selection of target metabolites following the proposed method consistently provided a significant benefit over random selection (Figures 4a-f) as confirmed by the results of Q-UR. It is worth noting that in some cases, particularly in the absence of oxygen, the ranking derived from first-order sensitivities alone is either comparably or more efficient in reducing total network flexibility (F_{TOT}) compared to its higher-order counterpart (Figures 4b & f). This indicates that, under certain conditions, the dependency of F_{TOT} on the concentration of uncertain metabolites may be better represented through a linear RSM which can be attributed to several factors such as: (a) limited thermodynamic degrees of freedom (due to the inclusion of many intracellular data for example), (b) limited number of thermodynamically bidirectional reactions under the physiological conditions considered and (c) the simplicity of the metabolic network, this is particularly true when studying isolated pathways or reactions. Moreover, computational requirements for the estimation of a purely linear RSM are significantly lower making it a particularly efficient and effective screening test.

3.4 Metabolite Sensitivity Analysis on a specific pathway or reaction

Even with currently available computing power, the detailed mathematical analysis of genome scale models can become a demanding and time consuming endeavor. Moreover, working with large-scale models generates a huge amount of data that has to be subsequently “mined” and processed. However, research attention is frequently focused on particular parts of the metabolic network such as a particular metabolic pathway leading to the formation of a desired end product. Two case studies are presented in this section, one where the focus of the analysis is constrained to the reactions that participate in the tricarboxylic acid (TCA) cycle and one where the network wide effectors of a single reaction are analyzed. Both of these studies have been conducted under the premises described in section 3.3, taking into account the experimental data from McCloskey et al [33] and under the assumption that ETC metabolites can’t be measured.

In terms of algorithmic implementation, all steps starting from the calibration of the model up until the evaluation of the model for each of the *in-Silico* experiments, e , specified by the experimental design follow the exact same procedures as they have been presented in section 3.3. However, since our focus in this case study was the allocation of uncertainty

within a subset of the reactions comprising the metabolic network (i.e. only the reactions of the TCA cycle) our output metric (R_T) had to be modified.

One of the key benefits of the approach proposed in the present work is its modularity and its ability to be reformulated in ways that enable it to address a wide variety of problems. This flexibility stems from the freedom to define an appropriate performance metric (Supplemental Material). In this case, since the output metric (R_T) selected (equation 6) is an additive function, a simple re-formulation such as the one described by equation 7 allows us to focus on any possible subset of reactions, irrespective of size.

Subsequently, we estimated RSMs that describe the allocation of thermodynamic uncertainty in the reactions that comprise the TCA cycle under both aerobic and anaerobic conditions and analyzed the magnitude of first- and higher- order sensitivities. An inspection of the metabolite significance ranking generated from the higher-order sensitivities reveals that F'_{TOT} is mostly affected by metabolites that take part in the TCA (Figures 3g & h, Supplemental Figure 4). However, TCA-metabolites in the significance ranking derived from the first-order sensitivities are much less prominent even under aerobic conditions. Based on the results of Figures 3g & h alone, one could argue that a similar reduction in F'_{TOT} could be achieved just by selecting metabolites participating in the pathway of focus without the need for further analysis and computation.

In order to evaluate, whether the proposed method offered an improvement over random or empirical selection we added an additional set of random trials in our Q-UR analysis. Therefore, apart from the 5000 completely random metabolite sets we created an additional 5000 random sets of 5 TCA-only metabolites to be constrained at-a-time. Each metabolite within these 5000 sets was assigned a random concentration value and the resulting $F_{TOT}^{validation}$ was estimated. The results of our Q-UR analysis highlight that even for a small subsystem (albeit still part of a large metabolic network) comprising only 14 reactions, an informed selection of targets for metabolomics leads to a larger reduction in uncertainty (Figures 4g & h). Furthermore, comparison between the average reduction in F'_{TOT} achieved by selection of metabolites from the first and higher-order based significance rankings (Figures 4g & h) revealed only minor differences. This indicates that metabolites extrinsic to the TCA cycle (e.g. glyceraldehyde-3-phosphate, O-phospho-L-serine, ethanol etc.) could exert an equally strong influence on the thermodynamic properties of the TCA reactions.

Figure 5.

Apart from knowing which metabolites can reduce the uncertainty in a metabolic network, it is often valuable to know which reactions are expected to be affected the most as a result of the reduction in F'_{TOT} . Therefore, we examined how the concentrations of metabolites selected either randomly or from the significance rankings derived herein affect the thermodynamic properties of the reactions of the TCA cycle. We examined the degree of reduction in F_R for each reaction across all samples of the Q-UR analysis and compared the statistics between informed and random selection (Figure 5). TMSA did not only outperform random selection in terms of achieved reduction of uncertainty across all reactions but it also proved to be more consistent. In many cases the results from random selection were inconsistent to the extent that it was not possible to calculate meaningful statistical metrics (aconitase and fumarate reductase reactions; Figures 5a and b). Interestingly, in the case of 2-oxoglutarate dehydrogenase under anaerobic conditions (Figure 5b) random selection outperformed TMSA, however due to the low degree of reduction achieved by both methods it can be considered as an outlier. Overall, the results of Figure 5 highlight that random selection of metabolites is not only less efficient in reducing total uncertainty but also leads to highly variable and “information-poor” results.

Of particular interest are reactions that exhibit a shift in their thermodynamic properties in response to perturbations in metabolite concentrations. Taking the MDH reaction (Supplemental Figure 5) as an example, we observe that under aerobic conditions perturbations in the concentrations of significant metabolites may lead to $\Delta_r G'$ values near thermodynamic equilibrium. Furthermore under anaerobic conditions, such perturbations may even alter the directionality of the reaction, allowing only the backward direction to be thermodynamically feasible. By limiting the formulation of equation (7) to consider a single reaction and repeating the whole analysis, we were able to identify the metabolites which could cause the observed shifts in the properties of the reaction.

Intuitively, one would expect the substrates and products of the reaction to be found at the top of the significance rankings (Supplemental Figure 6). Interestingly, one of the reactants of the MDH reaction, malate, does not appear to be significant according to the first-order sensitivities. Fumarate, the product of the subsequent fumarase catalyzed reaction is part of the subset of experimentally measured metabolites and as such is tightly constrained within experimental error. The thermodynamic coupling between fumarate and

malate through the fumarase catalyzed reaction might therefore be limiting the thermodynamically permissible range of malate. Furthermore, the proposed methodology provides additional insight, revealing “counter-intuitive” metabolites in the lower parts of glycolysis (3-Phosphohydroxypyruvate, 3-Phospho-D-glycerate, 3-Phospho-D-glyceroyl phosphate) that can affect the thermodynamic properties of the MDH reaction (Supplemental Figures 6 & 7). Finally Q-UR, performed following the procedures highlighted in the previous sections, showed that metabolites selected through the proposed methodology outperform random or empirical selection even when considering a single reaction (Supplemental Figure 8).

4. Conclusions

Deciphering the thermodynamic properties of a metabolic network is the first step towards uniquely defining the internal state of a cell. Moreover, determining which reactions operate near thermodynamic equilibrium under a given set of environmental conditions is crucial in understanding regulation at the metabolic level. Herein we presented TMSA, a method to quantify and allocate thermodynamic uncertainty within a metabolic network. This is the first, to the extent of our knowledge, systematic method that can rank metabolites based on their ability to constrain the solution space of a metabolic network. TMSA offers a platform both for the design of informative experiments and the prioritization and organization of existing experimental datasets. Moreover, it can be used to gain insight on the thermodynamic interactions within the network leading to the identification and evaluation of metabolic engineering strategies. The approach was shown to be modular and can be applied at the single reaction, pathway and whole network level with or without taking into account pre-existing experimental information. Further studies are required to understand how the ranking of metabolites derived from reduced metabolic models transfers back to the original GEM. For example, one could ask whether metabolites found to reduce uncertainty in the thermodynamic properties of the TCA cycle using the reduced model would achieve a similar reduction when using the GEM. A systematic study aimed at understanding the transferability of metabolite sensitivity information between GEMs and their reduced counterparts is the subject of current studies for future publication.

5. References

- [1] Miskovic, L., Hatzimanikatis, V., Production of biofuels and biochemicals: in need of an ORACLE. *Trends Biotechnol* 2010, 28, 391-397.
- [2] Chakrabarti, A., Miskovic, L., Soh, K. C., Hatzimanikatis, V., Towards kinetic modeling of genome-scale metabolic networks without sacrificing stoichiometric, thermodynamic and physiological constraints. *Biotechnol J* 2013, 8, 1043-U1105.
- [3] Stanford, N. J., Lubitz, T., Smallbone, K., Klipp, E., *et al.*, Systematic Construction of Kinetic Models from Genome-Scale Metabolic Networks. *Plos One* 2013, 8.
- [4] Miskovic, L., Tokic, M., Fengos, G., Hatzimanikatis, V., Rites of passage: requirements and standards for building kinetic models of metabolic phenotypes. *Curr Opin Biotechnol* 2015, 36, 146-153.
- [5] Ataman, M., Hatzimanikatis, V., Heading in the right direction: thermodynamics-based network analysis and pathway engineering. *Curr Opin Biotechnol* 2015, 36, 176-182.
- [6] Palsson, B., *Systems biology : properties of reconstructed networks*, Cambridge University Press, New York 2006.
- [7] O'Brien, E. J., Lerman, J. A., Chang, R. L., Hyduke, D. R., Palsson, B. O., Genome-scale models of metabolism and gene expression extend and refine growth phenotype prediction. *Mol Syst Biol* 2013, 9.
- [8] Orth, J. D., Thiele, I., Palsson, B. O., What is flux balance analysis? *Nat Biotechnol* 2010, 28, 245-248.
- [9] Soh, K. C., Hatzimanikatis, V., Constraining the flux space using thermodynamics and integration of metabolomics data. *Methods Mol Biol* 2014, 1191, 49-63.
- [10] Lee, J. M., Gianchandani, E. P., Papin, J. A., Flux balance analysis in the era of metabolomics. *Brief Bioinform* 2006, 7, 140-150.
- [11] Soh, K. C., Miskovic, L., Hatzimanikatis, V., From network models to network responses: integration of thermodynamic and kinetic properties of yeast genome-scale metabolic networks. *Fems Yeast Res* 2012, 12, 129-143.
- [12] Binns, M., de Atauri, P., Vlysidis, A., Cascante, M., Theodoropoulos, C., Sampling with pooling-based flux balance analysis: optimal versus sub-optimal flux space analysis of *Actinobacillus succinogenes*. *Bmc Bioinformatics* 2015, 16.
- [13] Reed, J. L., Shrinking the Metabolic Solution Space Using Experimental Datasets. *Plos Comput Biol* 2012, 8.
- [14] Kelk, S. M., Olivier, B. G., Stougie, L., Bruggeman, F. J., Optimal flux spaces of genome-scale stoichiometric models are determined by a few subnetworks. *Sci Rep-Uk* 2012, 2.
- [15] Trinh, C. T., Wlaschin, A., Sreenc, F., Elementary mode analysis: a useful metabolic pathway analysis tool for characterizing cellular metabolism. *Appl Microbiol Biot* 2009, 81, 813-826.
- [16] Henry, C. S., Broadbelt, L. J., Hatzimanikatis, V., Thermodynamics-based metabolic flux analysis. *Biophys J* 2007, 92, 1792-1805.
- [17] Vazquez, A., Oltvai, Z. N., Molecular Crowding Defines a Common Origin for the Warburg Effect in Proliferating Cells and the Lactate Threshold in Muscle Physiology. *Plos One* 2011, 6.
- [18] Chen, X. W., Alonso, A. P., Allen, D. K., Reed, J. L., Shachar-Hill, Y., Synergy between C-13-metabolic flux analysis and flux balance analysis for understanding metabolic adaption to anaerobiosis in *E. coli*. *Metab Eng* 2011, 13, 38-48.
- [19] Masakapalli, S. K., Le Lay, P., Huddleston, J. E., Pollock, N. L., *et al.*, Subcellular Flux Analysis of Central Metabolism in a Heterotrophic Arabidopsis Cell Suspension Using Steady-State Stable Isotope Labeling. *Plant Physiol* 2010, 152, 602-619.

- [20] Quek, L. E., Dietmair, S., Kromer, J. O., Nielsen, L. K., Metabolic flux analysis in mammalian cell culture. *Metab Eng* 2010, *12*, 161-171.
- [21] Hadlich, F., Noh, K., Wiechert, W., Determination of flux directions by thermodynamic network analysis: Computing informative metabolite pools. *Math Comput Simulat* 2011, *82*, 460-470.
- [22] Edwards, J. S., Palsson, B. O., Robustness analysis of the Escherichia coli metabolic network. *Biotechnol Progr* 2000, *16*, 927-939.
- [23] Mahadevan, R., Edwards, J. S., Doyle, F. J., Dynamic flux balance analysis of diauxic growth in Escherichia coli. *Biophys J* 2002, *83*, 1331-1340.
- [24] Segre, D., Vitkup, D., Church, G. M., Analysis of optimality in natural and perturbed metabolic networks. *P Natl Acad Sci USA* 2002, *99*, 15112-15117.
- [25] Klier, C., Use of an Uncertainty Analysis for Genome-Scale Models as a Prediction Tool for Microbial Growth Processes in Subsurface Environments. *Environ Sci Technol* 2012, *46*, 2790-2798.
- [26] Hoppe, A., Hoffmann, S., Holzhutter, H. G., Including metabolite concentrations into flux balance analysis: Thermodynamic realizability as a constraint on flux distributions in metabolic networks. *Bmc Syst Biol* 2007, *1*.
- [27] Miskovic, L., Hatzimanikatis, V., Modeling of Uncertainties in Biochemical Reactions. *Biotechnol Bioeng* 2011, *108*, 413-423.
- [28] Jankowski, M. D., Henry, C. S., Broadbelt, L. J., Hatzimanikatis, V., Group contribution method for thermodynamic analysis of complex metabolic networks. *Biophys J* 2008, *95*, 1487-1499.
- [29] Saltelli, A., *Global sensitivity analysis : the primer*, John Wiley, Chichester, England ; Hoboken, NJ 2008.
- [30] Chan, K., Saltelli, A., Tarantola, S., Sensitivity analysis of model output: Variance-based methods make the difference. *Proceedings of the 1997 Winter Simulation Conference* 1997, 261-268.
- [31] Kiparissides, A., Georgakis, C., Mantalaris, A., Pistikopoulos, E. N., Design of In Silico Experiments as a Tool for Nonlinear Sensitivity Analysis of Knowledge-Driven Models. *Ind Eng Chem Res* 2014, *53*, 7517-7525.
- [32] Box, G. E. P., Wilson, K. B., On the Experimental Attainment of Optimum Conditions. *J Roy Stat Soc B* 1951, *13*, 1-45.
- [33] McCloskey, D., Gangoiti, J. A., King, Z. A., Naviaux, R. K., *et al.*, A Model-Driven Quantitative Metabolomics Analysis of Aerobic and Anaerobic Metabolism in E. coli K-12 MG1655 That Is Biochemically and Thermodynamically Consistent. *Biotechnol Bioeng* 2014, *111*, 803-815.
- [34] Box, G. E. P., Draper, N. R., *Empirical model-building and response surfaces*, Wiley, New York 1987.
- [35] Box, G. E. P., Behnken, D. W., Some New Three Level Designs for the Study of Quantitative Variables. *Technometrics* 1960, *2*, 455-475.
- [36] Orth, J. D., Conrad, T. M., Na, J., Lerman, J. A., *et al.*, A comprehensive genome-scale reconstruction of Escherichia coli metabolism-2011. *Mol Syst Biol* 2011, *7*.
- [37] Ataman, M., Hatzimanikatis, V., redGem: A method for the consistent reduction of Genome-scale Metabolic Models (GEM). *to be submitted for publication* 2016.
- [38] Bekker, M., Kramer, G., Hartog, A. F., Wagner, M. J., *et al.*, Changes in the redox state and composition of the quinone pool of Escherichia coli during aerobic batch-culture growth. *Microbiol-Sgm* 2007, *153*, 1974-1980.

[39] Ogasawara, H., Ishida, Y., Yamada, K., Yamamoto, K., Ishihama, A., PdhR (pyruvate dehydrogenase complex regulator) controls the respiratory electron transport system in *Escherichia coli*. *J Bacteriol* 2007, *189*, 5534-5541.

Figure 1: Illustrative examples describing how different types of correlations between two metabolites A and B affect the feasible solution space of a metabolic network. Red dots denote metabolite concentration bounds; Light blue shaded area defines the solution space. (a) Linear relationship ($R_T = \beta_0 + \beta_A A + \beta_B B$): metabolites A and B have an equal contribution to uncertainty ($\beta_A = \beta_B$) (b) Bilinear relationship ($R_T = \beta_0 + \beta_A A + \beta_B B + \beta_{AB} AB$): metabolites A and B affect the permissible range of each other through a negative correlation ($\beta_{AB} < 0$) (c) Bilinear relationship ($R_T = \beta_0 + \beta_A A + \beta_B B + \beta_{AB} AB$): metabolites A and B affect the permissible range of each other through a positive correlation ($\beta_{AB} > 0$) (d) Non-linear relationship ($R_T = \beta_0 + \beta_A A + \beta_B B + \beta_{AB} AB + \beta_{AA} A^2 + \beta_{BB} B^2$): metabolites A and B affect the permissible range of each other through a higher-order correlation ($\beta_{AA}, \beta_{BB} \neq 0$);

Figure 2: Diagrammatic representation of TMSA algorithm and workflow.

Figure 3: TMSA results for the case studies discussed in the present work. Values normalized by the highest sensitivity index. Results for (blue) aerobic and (red) anaerobic conditions.

(a,b): TMSA with no *a priori* information; (c,d): TMSA with pre-existing experimental data; (e,f): TMSA with pre-existing experimental data & excluded metabolites; (g,h): TMSA on a specific metabolic pathway (TCA).

Sensitivity Indices derived from the calculated RSM when considering only: First order terms (Figures a,c,e,g – left column) and Quadratic terms (Figures b,d,f,h – right column).

Figure 4: Quantification of Potential for Uncertainty Reduction (Q-UR) results for the case studies discussed in the present work. Kernel smoothed probability distribution functions of the achieved reduction in F_{TOT} estimated during the Q-UR trials along with box-plots of the raw data.

(a,b): Q-UR with no *a priori* information; (c,d): Q-UR with pre-existing experimental data; (e,f): Q-UR with pre-existing experimental data & excluded metabolites; (g,h): Q-UR on a specific metabolic pathway (TCA).

Results for Aerobic conditions (Figures a,c,e,g – left column) and Anaerobic conditions (Figures b,d,f,h – right column).

(i - blue) Top-5 metabolites from Higher-order sensitivities; (ii - red) Top-5 metabolites from First order sensitivities; (iii - black) Sets of 5 randomly selected metabolites; (Figures g,h only: iv – light blue) Sets of 5 randomly selected TCA metabolites.

Figure 5: Boxplots of the normalised reduction in F_R for each reaction of the TCA cycle: (blue - dotted) Q-UR of Top-5 metabolites selected by TMSA (Higher-order sensitivities, 3200 model evaluations); (red - solid) Q-UR of sets of 5 randomly selected metabolites (5000 model evaluations). Median and mean of the distributions are denoted by a star and vertical line respectively. Results for (A) Aerobic conditions and (B) Anaerobic conditions.

Thermodynamics-based Metabolite Sensitivity Analysis in Metabolic Networks

A.Kiparissides^{1,2}, V. Hatzimanikatis^{1*}

1, Laboratory of Computational Systems Biotechnology
École Polytechnique Fédérale de Lausanne (EPFL)
CH-1015, Lausanne, Switzerland

2, Department of Biochemical Engineering
University College London
WC1E 6BT, London, UK

Supplemental Material

1. Notes on the application of SA methods on TFBA models

A number of model analysis techniques, labeled as Sensitivity Analysis (SA), aid in examining how variability in the model parameters propagates through the mathematical model and how it affects the model's observable output. In other words, SA provides both qualitative and quantitative insight regarding the dependence of the model output to its parameters through what is known as parameter significance ranking. Global Sensitivity Analysis (GSA) methods in particular, are frequently the methods of choice when studying complex non-linear models due to their ability to differentiate between linear and higher order (non-linear) sensitivity [29,30]. Estimation of Sensitivity Indices with state-of-the-art GSA methods requires multiple model evaluations (N) where the uncertain model parameters are assigned values from one (or more) set(s) of (quasi-) randomly generated numbers. Each random trial (N_i) is defined by the vector of inputs (X_i), which contains a randomly generated value for each model parameter and the vector of outputs (Y_i), which contains the values of the observable outputs after evaluation of the model. Furthermore, the estimation of non-linear sensitivity indices requires additional model evaluations (N'), using permutations of the original random set(s) (X'_i) [29].

In general GSA methods treat model parameters as random variables within the parameter space in order to study the total output variance and how it is apportioned to the model parameters. Accordingly, the random vectors of inputs (X_i) are

generated by random sampling of the parameter space. In the context of TFBA models, metabolites concentrations have to adhere to the mass balance and thermodynamic constraints. Consequently, any set of metabolite concentrations can yield a feasible random trial (N_i) if and only if it is part of the TFBA solution space. The solution space of constraint-based models is sampled using “hit-and-run” type of samplers, which guarantee that the generated vector of inputs (X_i) satisfies all constraints. However, not all permutations (X'_i) of this original set (X_i) will result in a *feasible input vector*. Therefore, current state-of-the-art global methods are not directly applicable to a TFBA setting due to the difficulties in generating the necessary feasible sets of random trials.

In order to circumvent the above limitations related with the selection of feasible input parameter vectors we propose a new approach for the estimation of first and higher-order sensitivities based on an *in silico* Design of Experiments (DoE) method originally developed by Kiparissides et al. [31].

2. Thermodynamics-based Flux Analysis (TFA)

For a detailed description of TFA the reader is directed to the original work of Henry et al.[16]. The basic equation that defines an FBA formulation is:

$$N \times v = 0 \quad (S1)$$

where N denotes the ($m \times r$) stoichiometric matrix containing the (m) metabolites (rows) that participate in the (r) reactions (columns) of the system and v denotes the ($r \times 1$) vector of reaction fluxes. For TFA formulations additional constraints are introduced that ensure compliance with the second law of thermodynamics:

$$N \cdot v = 0,$$

$$(a) \quad 0 \leq v_i \leq z_i v_{i,\max}, \quad \{i=1, \dots, r\},$$

$$(b) \quad \Delta_r G'_i - K + K \cdot z_i < 0, \quad \left\{ i=1, \dots, r \mid \Delta_r G'_i \text{ is known} \right\},$$

$$(c) \quad \Delta_r G'_i = \Delta_r G_i^0 + RT \sum_{j=1}^m n_{i,j} \ln(x_j), \quad \left\{ i=1, \dots, r+L \mid \Delta_r G_i^0 \text{ is known} \right\},$$

$$(d) \quad \Delta_r G'_i - K \cdot y_i < 0, \quad \{i=1, \dots, r+L\},$$

$$(e) \quad y_i + \sum_{j=1}^r a_{i,j} z_j \leq \sum_{j=1}^r a_{i,j}, \quad \{i=1, \dots, r+L\},$$

(S2)

where N is the $(m \times r)$ augmented stoichiometric matrix and v is the $(r \times 1)$ flux vector. In TFA formulation each of the reversible reactions in the model are split into separate irreversible forward and backward component reactions during the formation of N . This separation is performed so that the flux through each reaction i can be constrained to be greater than or equal to zero. The binary use variable, z_i , is associated with each reaction i and is equal to one if the flux through reaction i , v_i , is positive and zero if otherwise. $V_{\max}$ in Eq.6a is the upper limit on the flux through any reaction typically set to a physiologically reasonable value such as 100 mmol/gmDW/h. Equations (S1) and (S2a) represent the mass balance constraints carried over from MFA with the only difference being the separation of reversible reactions into forward and backward component reactions[9].

The additional thermodynamic constraints of TFA begin with Eq. S2b, which ensures that the activity profiles and flux distributions generated by TFA adhere to the second law of thermodynamics; a reaction flux cannot be positive unless $\Delta_r G'_i$ is negative. K is a constant selected to be large enough that Eq. S2b is always satisfied if v_i and z_i are zero. The terms involving K in Eq. S2b ensure that the constraint is only applied to reactions with a non-zero flux. Equation S2c is the Gibbs free energy equation used to set the value of $\Delta_r G'_i$ for reaction i given the metabolite activities. In Eq. S2c, $\ln(x_j)$ is the natural logarithm of the activity of compound j and $n_{i,j}$ is the stoichiometric coefficient of compound j in reaction i .

The $\Delta_r G^\circ$ of a reaction must be known to formulate the thermodynamic constraints for a reaction. Therefore, the constraints described by Eqs. S2b and S2c are not applied to the reactions in the model with unknown $\Delta_r G^\circ$, and instead, these reactions are lumped into overall reactions with known $\Delta_r G^\circ$ and special thermodynamic constraints are applied to the reactions for which $\Delta_r G^\circ$ is unknown; these constraints are described by Eqs. S2c–e. In these equations, L is the number of lumped reactions, and a_{ij} is a coefficient equaling one if reaction j is one of the original reactions with unknown $\Delta_r G^\circ$ that makes up lumped reaction i . When Eq. S2c is applied to a lumped reaction, it sets the value for $\Delta_r G'_i$ of the lumped reaction. Equation S2d is the thermodynamic feasibility constraint for the lumped reactions similar to Eq. S2b except that the binary use variable for each lumped reaction, y_i , is set to zero when the reaction is thermodynamically feasible and one when the reaction is infeasible. Equation S2e excludes flux distributions that involve flux through the set of reactions that comprise an infeasible lumped reaction. The continuous independent variables of this optimization problem are $\ln(x_j)$, v_i , and $\Delta_r G'_i$ and the binary independent variables are z_i and y_i .

Displacement from thermodynamic equilibrium (Γ) is a dimensionless quantity with values ranging from 0 to 1, that indicates how close to or far from thermodynamic equilibrium a reaction operates [27]. It is defined as:

$$\Gamma = \frac{1}{k_{eq}} \frac{\prod_l P_l}{\prod_k S_k} \quad (\text{S3})$$

where k_{eq} denotes the equilibrium constant for the reaction. When $\Gamma \ll 1$ the reaction is far away from thermodynamic equilibrium, while when $\Gamma \approx 1$ the reaction operates near thermodynamic equilibrium.

3. Construction of experimental design levels

Following the DoE methodology each uncertain input factor is assigned a pre-defined amount of levels within the range of values that will be considered. In semblance to the process of designing wet-lab experiments, there are several ways of designing and selecting these levels. Ultimately, the selection of the appropriate design levels is defined by the question one wishes to investigate. Indicatively, and in

the context of analyzing uncertainty in the thermodynamic properties of a metabolic network, some of the possible definitions are discussed below:

- (a) **Exact Numerical Values:** Intuitively one could assign a single metabolite concentration value to each level of the experimental design. It is worth noting that such an analysis would not yield information on how uncertainty propagates through the network. Rather, the RSM computed through such a design will describe how concentration levels for each metabolite affect the thermodynamic properties of the network. Additionally, such a design will suffer from the so-called “curse of dimensionality” as the complexity of the space that needs to be sampled from the experimental design would increase exponentially with the number of metabolites considered.
- (b) **Percentile Ranges from Sampled Probability Distribution:** An alternative approach would be to consider different metabolite concentration ranges for each level of the experimental design. These ranges could correspond to specific percentiles of a probability distribution function (pdf) derived from existing experimental measurements for each metabolite. Alternatively, in the absence of explicit experimental information, the TFA solution space could be sampled using hit-and-run algorithms in order to derive approximate pdfs. However, such information is not routinely available and, in the case of using the numerical approximation, subject to bias and errors introduced by the computational methods used to derive the pdf.
- (c) **Discretization of the thermodynamically feasible range:** The thermodynamically feasible range for each metabolite, defined by TVA, can be discretized into a number of equal intervals. Each design level could include one, or more, of these intervals and the resulting RSM would provide a functional relationship describing how constraining metabolites to a certain range of concentration values affects the thermodynamic properties of the network. If we take into consideration the fact that most intracellular measurements include a considerable error range, this type of design level seems more realistic compared to option (a). Additionally, it is not affected by bias or errors introduced through computational methods or analytical techniques.

4. Implementation of TMSA by discretization of the thermodynamically permissible range for each metabolite

In the absence of *a priori* information metabolite concentrations were allowed to vary between widely defined upper and lower bounds in order to ensure all physiological conditions are taken into consideration[16]. Subsequently, TVA was used to define the thermodynamically feasible concentration range for each metabolite ($M_{i,min} - M_{i,max}$) as well as the thermodynamically feasible $\Delta_r G'$ range for each reaction ($\Delta_r G'_{j,min} - \Delta_r G'_{j,max}$) within these bounds. Finally, a 3-level Box-Behnken experimental design was generated using a commercial software package (MATLAB R2014a, The MathWorks Inc., Natick, MA, 2000). As mentioned above, intracellular metabolite concentrations were treated as variable input factors and the DoE levels were defined as follows:

- (a) The thermodynamically feasible range for each metabolite (i) was discretized into 100 equidistant intervals, numbered from $M_{i,min}$ to $M_{i,max}$. The first and last 5 intervals were treated as outliers and were not taken into further consideration. Each metabolite was assigned one of the 3 possible levels defined in the experimental design.
- (b) DoE level (1): Concentration of metabolite m was randomly assigned a value between the 5th and the 45th interval, corresponding to an arbitrarily low concentration value.
- (c) DoE level (2): Concentration of metabolite m was randomly assigned a value between the 55th and the 95th interval, corresponding to an arbitrarily high concentration value.
- (d) DoE level (3): Concentration of metabolite m was allowed to take any value between $M_{i,min}$ and $M_{i,max}$ corresponding to an unconstrained (or uncertain) metabolite.

In order to assess the sensitivity of the proposed method to the existence of partial experimental information, we integrated the metabolomics data from McCloskey et al[34], taking into account the reported experimental uncertainty, and repeated the analysis. Extracellular metabolite concentrations were not considered in the present analysis, as they are generally readily quantifiable in most present day

experimental set-ups. Analysis of the intracellular metabolite concentrations reported by McCloskey et al[34] revealed an average error of $\pm 25\%$ in the measured value (see section 6). Therefore, metabolite concentrations were allowed to vary within $\pm 25\%$ of the randomly generated values in steps (b) and (c). Consequently, a gap between the 45th and the 55th interval had to be allowed in order to accommodate for the experimental error. TVA was performed for each *in silico* experiment and the studied output response R_T was estimated. A quadratic response surface model (RSM) with interaction terms describing the functional dependence of R_T on the concentration range of the analyzed metabolites was estimated.

5. Quantification of Thermodynamic Uncertainty in Metabolic Networks

In order to study how metabolites contribute to and affect output uncertainty in the context of a metabolic network, either qualitatively or quantitatively, it is necessary to define an appropriate metric to characterize the output uncertainty (R_T). Similar to the step described in section 2.2 of the main text, many alternative performance metrics can be conceived and implemented each analyzing the thermodynamic uncertainty in metabolic networks from a different perspective. In Supplemental Table 1 we indicatively propose alternative performance metrics that can be used to address some interesting and scientifically relevant objectives using the proposed methodology.

Supplemental Table 1 Classes of problems and proposed performance metrics

Objective	Analysis Scale	Proposed metric
Identification of metabolites able to reduce the uncertainty of the thermodynamic properties of a reaction	Single Reaction	Estimation of the thermodynamically permissible range of $\Delta_r G'$ values (F_R , Eq. 5) for the reaction considered
Identification of metabolites able to move a reaction far from thermodynamic equilibrium	Reaction/Pathway	Estimation of the feasible range of values for the displacement coefficient f

Identification of metabolites able to resolve thermodynamic bottlenecks	Pathway/Genome scale	Estimation of the number of reactions near thermodynamic equilibrium
Identification of metabolites able to reduce the uncertainty of the thermodynamic properties of a metabolic network	Genome scale	Estimation of the thermodynamically permissible ranges of $\Delta_r G'$ values for all participating reactions (F_{TOT} , Eq. 6)
Identification of metabolites able to reduce uncertainty in experimental measurements	Genome scale	Estimation of the thermodynamically permissible range of values for the experimentally measured metabolites

The case studies discussed in the main text focus on just one of the objectives, namely the identification of metabolites able to reduce uncertainty in the thermodynamic properties of a metabolic network. The lack of information regarding the expected or physiological levels of intracellular metabolite concentrations for a particular strain or set of experimental conditions propagates as uncertainty in the estimated Gibbs free energy of reaction, $\Delta_r G'$, and consequently in the determination of reaction directionalities. Therefore ranking metabolites based on their ability to reduce the uncertainty of the thermodynamic properties in the reactions of the entire network can provide a guide for the systematic selection of “information rich” metabolomics measurements that can improve our ability to uniquely define the internal state of the cell. Such selection methods are of particular interest for data collection and model validation studies, especially when considering large-scale models and complex data sets.

6. Effect of experimental information on Total Network Flexibility (F_{TOT})

Supplemental Table 2 contains the values of total network flexibility (F_{TOT}) as estimated during the model calibration step for the various case studies considered in the main text. F_{TOT} values are normalized against the first case study (3.1 No *a priori* information).

Supplemental Table 2 Total network flexibility values for 3 sets of simulations. Values are normalized against the first set of simulations (no *a priori* information)

Case Study	Normalized F_{TOT}	
	Aerobic	Anaerobic
(3.1) No <i>a priori</i> information	1	1
(3.2) Intracellular data	0.79	0.845
(3.3) Intracellular data & unmeasurable metabolites	0.79	0.845

7. TCA cycle metabolites included in the reduced E.Coli model

Supplemental Table 3 contains the metabolites participating in the TCA reactions included in the reduced *E. coli* metabolic network utilized in the present study.

Supplemental Table 3 Metabolites participating in the reactions of the TCA cycle found in the reduced E.Coli metabolic network

2-Demethylmenaquinone 8	Isocitrate
2-Demethylmenaquinol 8	L-Malate
Acetate	Menaquinol 8
Acetyl-CoA	Menaquinone 8
cis-Aconitate	NAD
ADP	NADH
2-Oxoglutarate	NADP
ATP	NADPH
Citrate	Oxygen
CO ₂	Oxaloacetate
Coenzyme A	Phosphate
Fumarate	Ubiquinone-8
H ⁺	Ubiquinol-8
H ₂ O	Succinate
Hydrogen peroxide	Succinyl-CoA

8. List of TFA constraints used

Experimental information for metabolite exchange rates and *E. coli* growth rates under both aerobic and anaerobic conditions were taken from Chen et al. [18] and McCloskey et al [34] and are summarized in Supplemental Table 4.

Supplemental Table 4 List of constraints used to integrate the experimental data from [16, 34] into the *E. coli* metabolic model

Exchange rates (mmol gDW ⁻¹ h ⁻¹)	Aerobic Growth		Anaerobic Growth	
	LB	UB	LB	UB
Glucose	-9.2	-7.6	-17.5	-12.5
Acetate	3	6	7	15
CO ₂	10.7	12.9	--	--
Ethanol	--	--	6.5	16
Formate	0	0.7	16.3	25
Lactate	--	--	0.5	1.2
O ₂	-13.1	-10.7	-0.05	0
Succinate	--	--	0.75	2.2
ATPM	7.8	18	7.8	17
Biomass*	0.57	0.6	0.38	0.42

* Biomass growth rate reported in h⁻¹

Supplemental Tables 5 and 6 contain the list of constraints that were used to integrate the metabolomics data from McCloskey et al¹ into the *E. coli* metabolic model, taking into account reported measurement error.

Supplemental Table 5 Implemented Metabolite Concentrations – Aerobic Conditions¹

Metabolite Name	Concentration Values - ln(M)	
	Lower Bound	Upper Bound
Acetate[e]	-11.51	-5.40
CO ₂ [e]	-10.00	-9.00
Ethanol[e]	-18.42	-11.51
Formate[e]	-18.42	-11.51
D-Glucose[e]	-11.51	-3.82
H ⁺ [e]	-18.42	-11.51
D-Lactate[e]	-18.42	-11.51
Ammonium[e]	-4.10	-3.96
Oxygen[e]	-10.07	-8.68
Phosphate[e]	-18.42	-2.88
Sulfate[e]	-6.00	-5.81
2-Oxoglutarate[c]	-10.00	-9.67
6-Phospho-D-gluconate[c]	-11.80	-9.23
Acetyl-CoA[c]	-8.18	-7.82
ADP[c]	-10.30	-9.70
alpha-D-Ribose 5-phosphate[c]	-10.90	-10.30
AMP[c]	-12.00	-11.70
ATP[c]	-7.70	-7.58
Coenzyme A[c]	-8.27	-7.66
D-Fructose 1,6-bisphosphate[c]	-5.80	-5.06
D-Fructose 6-phosphate[c]	-8.34	-7.63
D-Glucose 6-phosphate[c]	-7.37	-6.84
D-Ribulose 5-phosphate[c]	-9.97	-9.34
L-Aspartate[c]	-8.94	-8.74
L-Glutamate[c]	-5.12	-4.91
L-Glutamine[c]	-6.74	-6.52
L-Serine[c]	-9.34	-8.75
NADH[c]	-8.91	-8.18
NAD[c]	-7.59	-7.32
NADPH[c]	-8.17	-7.67
NADP[c]	-10.30	-9.89
Phosphoenolpyruvate[c]	-10.40	-8.42

Supplemental Table 6 Implemented Metabolite Concentrations – Anaerobic Conditions¹

Metabolite Name	Concentration Values - log(M)	
	Lower Bound	Upper Bound
Acetate[e]	-7.09	-5.40
CO ₂ [e]	-10.00	-9.00
Ethanol[e]	-8.43	-5.47
Formate[e]	-9.13	-4.93
D-Glucose[e]	-11.51	-3.82
H ⁺ [e]	-18.42	-11.51
D-Lactate[e]	-11.51	-5.91
Ammonium[e]	-4.10	-3.96
Oxygen[e]	-18.42	-11.51
Phosphate[e]	-18.42	-2.88
Sulfate[e]	-6.00	-5.81
Acetyl-CoA[c]	-8.00	-7.65
ADP[c]	-9.66	-9.29
alpha-D-Ribose 5-phosphate[c]	-14.13	-10.70
AMP[c]	-11.52	-11.34
ATP[c]	-7.28	-6.91
Coenzyme A[c]	-7.87	-7.54
D-Fructose 1,6-bisphosphate[c]	-5.52	-5.06
D-Fructose 6-phosphate[c]	-8.75	-7.96
D-Glucose 6-phosphate[c]	-7.98	-6.82
D-Ribulose 5-phosphate[c]	-9.23	-8.96
L-Glutamate[c]	-5.76	-5.55
L-Glutamine[c]	-9.06	-8.64
L-Serine[c]	-9.07	-8.48
NADH[c]	-7.13	-6.77
NAD[c]	-7.26	-7.00
NADPH[c]	-8.82	-8.46
NADP[c]	-9.84	-9.50
Phosphoenolpyruvate[c]	-8.92	-7.39

Additional References

1. McCloskey, D.; Gangoiti, J. A.; King, Z. A.; Naviaux, R. K.; Barshop, B. A.; Palsson, B. O.; Feist, A. M., A model-driven quantitative metabolomics analysis of aerobic and anaerobic metabolism in E. coli K-12 MG1655 that is biochemically and thermodynamically consistent. *Biotechnol Bioeng* **2014**, *111* (4), 803-15.

9. Supplemental Figures

Supplemental Figure 1: TMSA with no a priori information. Comparison between First and Higher-order sensitivities for (a) aerobic and (b) anaerobic conditions. Values normalized by the highest sensitivity index

Supplemental Figure 2: TMSA with pre-existing experimental data. Comparison between First and (a) aerobic and (b) anaerobic conditions. Values normalized by the highest sensitivity index

Supplemental Figure 3: TMSA with pre-existing experimental data & excluded metabolites. Comp Order Sensitivity Indices for: (a) aerobic and (b) anaerobic conditions. Values normalized by the h

Supplemental Figure 4: TMSA on a specific metabolic pathway (TCA). Comparison between First for: (a) aerobic and (b) anaerobic conditions. Values normalized by the highest sensitivity index

Supplemental Figure 5: Thermodynamic Variability Analysis of the reactions of the TCA cycle for the highest reduction in F'_{TOT} : (blue) TMSA (Higher-order sensitivities); (grey) Unconstrained. Numbers in brackets order reactions from most to least affected by the TMSA selected metabolites (blue bars). Results for (a) Aerobic conditions and (b) Anaerobic conditions.

Supplemental Figure 6: TMSA on a specific reaction (MDH). Sensitivity Indices derived from the calculated RSM when considering only: (a) First order terms and (b) Quadratic terms. Values normalized by the highest sensitivity index. Results for (blue) aerobic and (red) anaerobic conditions.

Supplemental Figure 7: TMSA on a specific pathway or reaction (MDH reaction). Comparison between Sensitivity Indices for: (a) aerobic and (b) anaerobic conditions. Values normalized by the highest sensitivity index.

Supplemental Figure 8: Quantification of Potential for Uncertainty Reduction (Q-UR) on a specific reaction (MDH). Kernel smoothed probability distribution functions of the achieved reduction in F_{TOT} estimated during the Q-UR trials along with box-plots of the raw data. (i - blue) Top-5 metabolites from Higher-order sensitivities; (ii - red) Top-5 metabolites from First order sensitivities; (iii - black) Sets of 5 randomly selected metabolites. Results for (A) Aerobic conditions and (B) Anaerobic conditions.