

Longitudinal patterns in objective physical activity and sedentary time in a multi-ethnic sample of children from the UK

L. Smith¹, D. Aggio² and M. Hamer³

¹The Cambridge Centre for Sport and Exercise Sciences, Department of Life Sciences, Anglia Ruskin University, Cambridge, UK; ²UCL Department of Primary Care and Population Health, UCL Medical School, UK; UCL Physical Activity Research Group, University College London, London, UK; ³School of Sport, Exercise & Health Sciences, Loughborough University

Address for correspondence: L. Smith, The Cambridge Centre for Sport and Exercise Sciences, Department of Life Sciences, Anglia Ruskin University, Cambridge, UK, CB1 1BT. E-mail: Lee.Smith@anglia.ac.uk

Received 17 October 2016; accepted 12 April 2017

Summary

Background: Children of South Asian decent born in the UK display lower levels of physical activity than British Caucasians although no longitudinal data are available.

Objectives: We aimed to investigate change in activity levels over 1 year in a diverse ethnic sample of children residing in London, UK.

Methods: Children were categorized into ethnic groups (Caucasian/mixed, Black, South Asian). At baseline and 1-year follow-up, children's objective physical activity was monitored (Actigraph accelerometer) for at least 1 day. Mixed models were employed to investigate differences in change in activity levels between ethnic groups.

Results: A total of 281 children were included in the analyses. South Asians had a significantly greater increase in time spent sedentary at follow-up than those of a Caucasian/mixed ethnicity (B [ratio sedentary/wear time] = 0.024; 95% confidence interval 0.003, 0.046). South Asian children recorded lower moderate to vigorous physical activity at baseline (B = -6.5, 95% confidence interval, -11.1, -1.9 min d⁻¹, *p* = 0.006) although levels remained relatively stable over follow-up and changes did not differ across ethnic group.

Conclusions: In a diverse ethnic sample of children from inner city London, those of a South Asian ethnicity exhibited a significantly greater increase in sedentary time over a period of 12 months in comparison with Caucasian/mixed and Black children.

Keywords: diabetes, ethnicity, longitudinal, physical activity, sedentary.

Introduction

Diabetes prevalence is strongly associated with ethnicity. The Health Survey for England showed that all minority ethnic groups have a higher risk of diabetes compared with the general population (1). Compared with Caucasians, British South Asian adults are at an increased risk of type II diabetes (2,3). South Asians show evidence of marked insulin resistance, hypertriglyceridemia and central adiposity (4). There is increasing evidence to suggest that differences in risk of type II diabetes emerge in early life. Indeed, British South Asians are at increased risk of type II diabetes in childhood and adolescence and show evidence of insulin resistance from 10 years of age (4–6).

A large body of literature exist that shows regular participation in physical activity aids in the prevention

of both obesity and type II diabetes (7,8). Moreover, a growing body of literature is emerging that suggests increasing sedentary time may also be contributing to the diabetes epidemic (9,10), independently of physical activity. National survey data have suggested that a large proportion of children in the UK do not achieve current physical activity recommendations (11,12), and this is particularly apparent in deprived inner city areas. Recent research, although limited, suggests that differences exist in children's activity levels between ethnic groups in the UK (13,14). For example, compared with Caucasians 9–10 year olds, South Asians recorded 41 fewer accelerometer (Actigraph-GT1M; an objective measure of free-living activity) counts per minute and fewer minutes in moderate and vigorous physical activity (MVPA) and more time in sedentary activity (13). Another cross-sectional study of primary school children (14) found

that 73% of Caucasians and only 35% of South Asians achieved 60 min of MVPA daily. Both studies concluded that South Asian children participate in significantly less physical activity than Caucasians in the UK. Studies using subjective measures of physical activity have found similar findings (15). However, previous studies are all of a cross-sectional design. Therefore, it is not known how activity levels in specific ethnic populations of children change over time. It has been suggested that physical activity decreases with age throughout childhood and adolescence (16); however, UK studies demonstrating this have used samples of predominantly Caucasian children, and differences in changes in activity levels have not been investigated between children of different ethnicity. One UK study (16) monitored objective physical activity at baseline and 1-year follow-up in a sample of 844 children (baseline 9–10 years; over 90% Caucasian). It was found that physical activity decreased over 1 year with 70.4% of children meeting physical activity recommendations at baseline and 65.8% at follow-up ($p < 0.001$). Understanding how levels of activity change over time in different ethnic populations is essential for the development of targeted physical activity promotion and sedentary behaviour reduction interventions and thus aids in the prevention of obesity and type II diabetes.

The present study aims to investigate changes in activity levels over 1 year in a diverse ethnic sample (Caucasian, mixed, South Asian, Black) of inner city children residing in Camden, London, UK.

Methods

Study design

Camden Active Spaces was a school-based quasi-experimental study examining physical activity before (summer term 2014) and after (summer term 2015) playground renovation. Because no overall effects of the intervention was observed, we collapsed the sample and adjusted for intervention/control group in the present analysis. The study protocol has previously been published (17). Head teachers from each school provided explicit written consent for their school and school children to take part in the study. Parents were given the option to 'opt-out' their child (ren). Ethical approval was granted by the University College London Research Ethics Committee (4400/002).

Outcome measure: physical activity

Trained researchers fitted accelerometers (Actigraph GT3X) to children at baseline and follow-up. Children

were asked to wear the device around their waist during waking hours every day for seven consecutive days, but not during water-based activities or sleep. Devices were programmed to sample at 30 Hz. Our protocol followed methods used in the International children's Accelerometry Database study (18). Data files were re-integrated to a 60-s epoch, and non-wear time was defined as 60 min of consecutive zeros, allowing for 2 min of none zero interruptions. The first partial day of wear was excluded from our analyses in order to reduce the possibility of reactivity to wearing the device. All children with at least one school day and at least 500 min of measured monitor wear time between 07:00 AM and midnight were included. Time spent sedentary was defined as all minutes less than 100 cpm, light activity as 100 up to 3000 cpm and MVPA as more than 3000 cpm.

Children self-reported sedentary behaviour using the *validated* Girls Health Enrichment Multi-Studies physical activity survey (19). Children completed the survey with the assistance of a trained research assistant, and the meaning of each question was explained. Children were asked 'Think about whether you usually watch TV, Videos/play computer games, video games/Homework, reading. Now please check the box that best fits how much time you spent doing each activity.' Response options were as follows: none/less than 30 min/30 min–1 h/1–3 h/more than 3 h.

Exposure measure: ethnicity

Children self-reported their ethnicity (Caucasian, mixed, South Asian, Black). Ethnicity was then categorized as (Caucasian/mixed, Black, South Asian). Child self-reported ethnicity has been successfully used in previous similar studies (15).

Overweight/obesity

Trained research assistants measured participants' weight using the Tanita SC-330 Body Composition Analyser (Tanita Inc, IL, USA) and height using the Leicester Height Measure, from which body mass index (BMI) was derived in kg m^{-2} . Participants were then categorized as normal weight/overweight/obese using the BMI International Obesity TaskForce age/sex-specific cut offs (20).

Participant characteristics

A total of 450 participants from seven intervention and two control schools were recruited into the study at baseline. Valid baseline Actigraph data were provided in 396 children, and of those, 281 (71%) completed

follow-up. Reasons for drop-out included left school/absent on day of follow-up data collection ($n = 46$), insufficient wear time ($n = 28$) and failure to return device ($n = 41$). Participants that completed the baseline assessment but not follow-up were slightly older (9.8 vs. 9.0 years, $p = 0.01$) than the children in the full analytic sample but no differences in sex distribution (49.6% vs. 47.7% boys) or ethnicity (23.5% vs. 18.0% South Asian in drop-outs and final sample, respectively) were noted.

In the analytic sample, 64% were Caucasian/mixed, 18% Black and 18% South Asian. The descriptive characteristics are displayed in Table 1. Approximately 10% ($n = 27$) of the sample were categorized as obese (using BMI International Obesity TaskForce age/sex-specific cut offs; Table 1). We found no statistical differences in obesity between ethnic groups (Chi-square test, $p = 0.77$). At baseline, 77.6% of the sample recorded at least 4 days of Actigraph wear and 6.4% only 1 day of wear.

Analyses

Characteristics of the study population at baseline were analysed using descriptive statistics. Mixed models, adjusted for age, sex, the ratio of MVPA/wear time at baseline, intervention or control group (fixed effect) and school (random effect, to account for clustering at the school level) were employed to compare ratio MVPA/wear time at follow-up between ethnic groups. We also examined change in sedentary time as the outcome, but for this analysis, we adjusted for the ratio of baseline sedentary/wear time instead. Analyses were

performed using SPSS version 22 (Chicago, IL, USA), and statistical significance was set at $p < 0.05$.

Results

At baseline, South Asians recorded less time in MVPA compared with Caucasians/mixed ($B = -6.5$, 95% confidence interval [CI], $-11.1, -1.9 \text{ min d}^{-1}$, $p = 0.006$) after adjusting for age, sex, wear time and school. There were no differences in MVPA between Black and Caucasian/mixed children ($B = 3.4$, 95% CI, $-1.2, 8.0$, $p = 0.14$). There were no differences in sedentary time between South Asians and Caucasians/mixed ($B = 9.7$, 95% CI, $-7.5, 26.8 \text{ min d}^{-1}$, $p = 0.27$) or Black and Caucasian/mixed children ($B = -5.2$, 95% CI, $-22.1, 11.7 \text{ min d}^{-1}$, $p = 0.54$) after covariate adjustments.

In adjusted mixed models to examine changes over time, there were no differences in MVPA between ethnic groups (Table 2; Fig. 1). South Asians, however, had a significantly greater increase in time spent sedentary at follow-up than those of a Caucasian/mixed ethnicity ($B = 0.024$; 95% CI 0.003, 0.046) (Table 2, Fig. 2).

In further analyses, we incorporated participants' weight status (normal weight/overweight/obese; categorized using BMI International Obesity TaskForce age/sex-specific cut offs) as a covariate; this made no substantial differences to the results (results not shown). We examined changes in self-reported sedentary behaviours. At baseline, South Asian children tended to report greater TV viewing and computer usage, although at follow-up, these

Table 1 Baseline characteristics of sample

	Caucasian/Mixed($n = 179$)	Black($n = 51$)	South Asian($n = 51$)
Age	8.7 \pm 2.0	9.0 \pm 2.1	9.9 \pm 1.9
Sex (% male)	50.8	47.0	37.3
BMI (kg m^{-2})	18.1 \pm 3.8	18.2 \pm 4.1	18.2 \pm 4.0
<i>Weight Status*</i>			
Normal weight	126 (71.2)	37 (74)	40 (78.4)
Overweight	31 (17.5)	9 (18)	8 (15.7)
Obese	20 (11.3)	4 (8)	3 (5.9)
Total Actigraph wear time (min d^{-1})	741 \pm 90	775 \pm 83	777 \pm 100
Total days wear	4.6 \pm 1.7	4.7 \pm 1.4	4.8 \pm 1.8
MVPA (min d^{-1})	28 \pm 15	32 \pm 20	20 \pm 12
Light PA (min d^{-1})	363 \pm 74	375 \pm 49	355 \pm 84
Sedentary (min d^{-1})	349 \pm 86	369 \pm 81	403 \pm 99

*Categorized using BMI International Obesity TaskForce age/sex-specific cut offs.
BMI, body mass index; MVPA, moderate and vigorous physical activity; PA, physical activity.



Table 2 Mixed models examining changes in physical activity and sedentary time in relation to ethnicity

	Coefficient (95% CI) for ratio MVPA/wear time at follow-up Ref	Coefficient (95% CI) for ratio sedentary/wear time at follow-up Ref
Caucasian/Mixed		
Black	0.001 (−0.005, 0.007)	−0.019 (−0.040, 0.002)
South Asian	−0.006 (−0.013, 0.0005)	0.024 (0.003, 0.046)*

* $p < 0.05$.

†Coefficient adjusted for age, sex, and the ratio of MVPA or sedentary/wear time at baseline, intervention or control group (as fixed effects) and school (as random effect).

CI, confidence interval; MVPA, moderate and vigorous physical activity.

**Figure 1** Changes in moderate and vigorous physical activity in relation to ethnicity ($n = 281$). [Colour figure can be viewed at wileyonlinelibrary.com]**Figure 2** Changes in sedentary time in relation to ethnicity ($n = 281$). [Colour figure can be viewed at wileyonlinelibrary.com]

differences were less apparent (Table 3). South Asian children reported greater homework, and this remained evident at follow-up.

Discussion

Previous cross-sectional studies of children have reported ethnic differences in physical activity. This is the first study to examine longitudinal changes in a multi-ethnic sample of children. Our data show that children of South Asian ethnicity exhibited a significantly greater increase in sedentary time over a period of 12 months in comparison with

Caucasian/mixed children. No significant differences were observed in 12-month changes in MVPA between ethnic groups, although South Asian children displayed lower MVPA at baseline. These findings support and add to previous cross-sectional research that shows children of a South Asian ethnicity residing in the UK have the lowest levels of activity (13,14). This is of particular concern as South Asians are at a significantly increased risk of type II diabetes early in life (2,3). Indeed, low levels of physical activity and high levels of sedentary behaviour are associated with incident diabetes (8–10), possibly through promoting visceral

Table 3 Self-reported sedentary behaviours at baseline and follow-up

	Caucasian/Mixed		Black		South Asian	
	Baseline	Follow-up	Baseline	Follow-up	Baseline	Follow-up
TV viewing (min d ⁻¹)						
<30	43.5	28.5	48.8	29.2	30.6	37.5
30 to 60	35.4	38.4	26.7	33.3	32.6	33.3
>60	21.1	32.0	24.4	37.5	36.7	29.2
Computer (min d ⁻¹)						
<30	56.8	46.0	62.2	42.9	47.9	56.3
30 to 60	24.7	29.9	11.1	30.6	25.0	22.9
>60	17.9	24.1	26.7	26.5	27.1	20.8
Homework (min d ⁻¹)						
<30	45.7	41.3	42.2	46.9	26.5	29.1
30 to 60	28.4	36.8	31.1	28.6	53.1	43.8
>60	25.9	21.8	26.7	24.5	19.6	27.1

Data presented as percentages.

adiposity and impaired glucose control (21,22). However, the aetiology of increased metabolic risk in South Asians is incompletely understood.

The self-reported data on sedentary behaviours provided some context to help interpret the objective data and largely confirmed the generally higher prevalence of sedentary activities such as TV viewing, computer use and homework in the South Asian children. Nevertheless, increases in self-reported sedentary behaviours over follow-up were less marked than objectively measured sedentary time in South Asian children. This is perhaps unsurprising as many physical activity questionnaires, including ours, have not been validated to examine changes over time. In addition, self-reported measures of sedentary behaviour are often poorly correlated with objective data from motion sensors (23).

The reasons for observed differences in activity behaviour across ethnic groups in the UK are poorly understood. However, given that similar trends have been reported in adult samples (24), it is unsurprising that these results are replicated in children because parental physical activity is a strong predictor of their children's activity (25). Socio-cultural factors may play a key role (26), for example, some conventional physical activities may not fit within the health beliefs and culture of South Asian communities.

In a recent qualitative study in South Asian *adults*, a number of barriers to participation in physical activity were described and appropriate intervention content recommended (27). Key barriers included the following: issues with personal safety, cost, family and childcare commitments and the availability of free time. The study suggests that the use of the following

will be relevant to intervention development: enhancing physical activity by increasing walking; use of pedometers; multilingual spoken content and delivery; peer support, for example, bilingual community members, to facilitate engagement and motivation; undertaking activity with families or friends; and delivery in local informal settings including the home. While some of these may be appropriate for South Asian children, they are adult-specific. To the best of our knowledge, similar data for South Asian children residing in the UK do not exist.

A clear strength of the present study is the longitudinal design in an ethnically diverse sample of children residing in London, UK. We adjusted for school as a random effect, to account for clustering at the school level, thus results are unlikely to have been explained by local area factors. The 1-year interval separating baseline and follow-up data collection ensured seasonally matched data, thereby reducing the risk of artefactual changes in activity attributable to different weather conditions at two time points. We chose to use a conservative cut point to derive MVPA; thus, some moderate intensity activity could have been misclassified as light activity. Nevertheless, the relatively low levels of physical activity in the present study are consistent with data from a prior survey using self-report where only 12% of children from Camden met the physical activity guideline (28); www.camden.gov.uk/jsna.

As in previous studies (13), registered wear time differed between ethnic groups, with Caucasians/mixed recording the lowest average wear period per day. Although analyses adjust for wear time, it is unknown what activities the children undertake during non-registered wear. However, if

Caucasian/mixed children were undertaking physical activity (such as water sports) during non-registered wear periods, the ethnic differences observed may reflect a conservative estimate of the true differences. Data on individual South Asian groups (e.g. Indian/Pakistan/Bangladesh) were not collected, and thus, it was not possible to investigate differences in activity levels between groups. However, a recent study found that levels of estimated VO_{2max} were lower in South Asian children than those in white Europeans and the lower estimated VO_{2max} in South Asians, compared with white Europeans, was consistent among Indian, Pakistani and Bangladeshi children. This suggests that any differences in physical activity levels between South Asian groups are likely to be minimal (29). Given the age range of the children in the present study, some children may have been entering puberty at the time of data collection, which could potentially confound the results presented. However, data on stage of maturation were not collected. Future studies should control for such data in analyses.

Conclusion

In this ethnically diverse sample of children, those of a South Asian ethnicity exhibited a significantly greater increase in sedentary time over a period of 12 months in comparison with Caucasian/mixed children. Sedentary habits established in early life may have implications for diabetes, particularly in high-risk groups such as South Asians.

Conflict of Interest Statement

No conflict of interest was declared.

Funding source

This work was supported by The Economic and Social Research Council, UK (ES/M003795/1), and London Borough of Camden. The funders had no role in the study design; in the collection, analysis and interpretation of data; in writing of the report; or in the decision to submit the paper for publication.

Contribution Statement

L. S. conceptualized and designed the study, drafted the manuscript and approved the final version. D. A. designed the study, collected the data and approved the final manuscript. M. H. conceptualized and designed the study, drafted the manuscript and approved the final version.

Acknowledgements

We acknowledge the contributions of Georgina Knock, Abigail Barnett, Sanchia Legister, Shila Khoshkbari, Sophie Sparrow, Patrycja Cholewa, Paula Hurril, Suzie Munnery and Verena Trend to data collection, and the staff from all schools for their cooperation.

References

- Public Health England. Adult obesity and type 2 diabetes. 2014.
- McKeigue PM, Shah B, Marmot MG. Relation of central obesity and insulin resistance with high diabetes prevalence and cardiovascular risk in South Asians. *Lancet* 1991; 337: 382–386.
- Bhopal R, Unwin N, White M, *et al.* Heterogeneity of coronary heart disease risk factors in Indian, Pakistani, Bangladeshi, and European origin populations: cross sectional study. *Br Med J* 1999; 319: 215–220.
- Whincup PH, Owen CG, Orfei L, McKay C, Cook DG. Ethnic differences in risk factors for chronic disease have their origins in early life: evidence for the CHASE study. *Early Hum Dev* 2007; 83: S63.
- Whincup PH, Gilg JA, Papacosta O, *et al.* Early evidence of ethnic differences in cardiovascular risk: cross sectional comparison of British South Asian and white children. *Br Med J* 2002; 324: 635.
- Whincup PH, Gilg JA, Owen CG, Odoki K, Alberti KG, Cook DG. British South Asians aged 13–16 years have higher fasting glucose and insulin levels than Europeans. *Diabet Med* 2005; 22: 1275–1277.
- Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS Jr. Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. *N Engl J Med* 1991; 325: 147–152.
- Hu FB, Sigal RJ, Rich-Edwards JW, *et al.* Walking compared with vigorous physical activity and risk of type 2 diabetes in women: a prospective study. *JAMA* 1999; 282: 1433–1439.
- Grontved A, Hu FB. Television viewing and risk of type 2 diabetes, cardiovascular disease, and all-cause mortality: a meta-analysis. *JAMA* 2011; 305: 2448–2455.
- Smith L, Hamer M. Television viewing time and risk of incident diabetes mellitus: the English longitudinal study of ageing. *Diabet Med* 2014; 31: 1572–1576.
- Joint Health Surveys Unit. The health survey for England 2008. The Information Centre for Health and Social Care Leeds 2009. www.ic.nhs.uk/pubs/hse08physicalactivity
- Department of Health. Start active stay active: a report on physical activity for health from the four home countries' Chief Medical Officers (UK). 2011.
- Owen CG, Nightingale CM, Rudnicka AR, Cook DG, Ekelund U, Whincup PH. Ethnic and gender differences in physical activity levels among 9–10-year-old children of white European, South Asian and African-Caribbean origin:

- the Child Heart Health Study in England (CHASE Study). *Int J Epidemiol* 2009; 38: 1082–1093.
14. Eyre EL, Duncan MJ, Smith EC, Matyka KA. Objectively measured patterns of physical activity in primary school children in Coventry: the influence of ethnicity. *Diabet Med* 2013; 30: 939–945.
 15. Khunti K, Stone MA, Bankart J, *et al.* Physical activity and sedentary behaviours of South Asian and white European children in inner city secondary schools in the UK. *Fam Pract* 2007; 24: 237–244.
 16. Corder K, van Sluijs E, Ekelund U, Jones A, Griffin S. Changes in children's physical activity over 12 months: longitudinal results from the SPEEDY study. *Paediatrics* 2010; 126: 926–935.
 17. Smith L, Kipps C, Aggio D, *et al.* Camden active spaces: Does the construction of active school playgrounds influence children's physical activity levels? A longitudinal quasi-experiment protocol. *BMJ Open* 2014; 4.
 18. Sherar LB, Griew P, Esliger DW, *et al.* International children's accelerometry database (ICAD): design and methods. *BMC Public Health* 2011; 11: 485.
 19. Treuth MS, Sherwood NE, Butte NF, *et al.* Validity and reliability of activity measures in African-American girls for GEMS. *Med Sci Sports Exerc* 2003; 35: 532–539.
 20. Cole TJ, Bellizzi MC, Flegal KM, Dietz WH. Establishing a standard definition for child overweight and obesity worldwide: international survey. *BMJ* 2000; 320: 1240–1245.
 21. Murabito J, Pedley A, Massaro J, *et al.* Moderate-to-vigorous physical activity with accelerometry is associated with visceral adipose tissue in adults. *J Am Heart Assoc* 2015; 4.
 22. Dowse G, Zimmet P, Gareeboo H, *et al.* Abdominal obesity and physical inactivity as risk factors for NIDDM and Impaired Glucose Tolerance in Indian, Creole, and Chinese Mauritians. *Diabetes Care* 1999; 14: 271–282.
 23. Stamatakis E, Hamer M, Tilling K, Lawlor DA. Sedentary time in relation to cardio-metabolic risk factors: differential associations for self-report vs. accelerometry in working age adults. *Int J Epidemiol* 2012; 41: 1328–1337.
 24. Williams ED, Stamatakis E, Chandola T, Hamer M. Assessment of physical activity levels in South Asians in the UK: findings from the Health Survey for England. *J Epidemiol Community Health* 2011; 65: 517–521.
 25. Moore L, Lombardi D, White M, Campbell J, Oliveria S, Ellison C. Influence of parents' physical activity levels on activity levels of young children. *J Pediatr* 1991; 118: 215–219.
 26. Farooqi A, Naqra D, Edqra T, Khunti K. Attitudes to lifestyle factors for coronary heart disease amongst South Asians in Leicester: a focus group study. *Fam Pract* 2000; 17: 293–297.
 27. Cross-Bardell L, George T, Bhoday M, Tuomainen H, Qureshi N, Kai J. Perspective on enhancing physical activity and diet for health promotion among at-risk urban UK South Asian Communities: a qualitative study. *BMJ Open* 2016; 5.
 28. Camden Joint Strategic Needs Assessment 2014/15. Camden Clinical Commissioning Group, NHS Camden. Chapter 9: Physical Activity. 2015.
 29. Nightingale CM, Donin AS, Kerry SR, *et al.* Cross-sectional study of ethnic differences in physical fitness among children of South Asian, black African-Caribbean and white European origin: the Child Heart and Health Study in England (CHASE). *BMJ Open* 2016; 6.