

**SUPPLEMENTARY TABLE 1 - PROPORTION OF PATIENTS RECRUITED
IN EACH PARTICIPATING COUNTRY**

| | Centers (N=69) | Adult patients with a cardiomyopathy (N=3208) |
|----------------------|-----------------------|--|
| Belarus, n(%) | 1 | 21/3208 (0.7%) |
| Czech Republic, n(%) | 1 | 42/3208 (1.3%) |
| Denmark, n(%) | 2 | 82/3208 (2.6%) |
| Egypt, n(%) | 1 | 44/3208 (1.4%) |
| Finland, n(%) | 2 | 158/3208 (4.9%) |
| France, n(%) | 5 | 208/3208 (6.5%) |
| Germany, n(%) | 3 | 146/3208 (4.6%) |
| Great Britain, n(%) | 5 | 185/3208 (5.8%) |
| Greece, n(%) | 2 | 106/3208 (3.3%) |
| Hungary, n(%) | 1 | 1/3208 (0.0%) |
| Italy, n(%) | 16 | 659/3208 (20.5%) |
| Lithuania, n(%) | 2 | 118/3208 (3.7%) |
| Netherlands, n(%) | 3 | 127/3208 (4.0%) |
| Poland, n(%) | 6 | 432/3208 (13.5%) |
| Portugal, n(%) | 2 | 84/3208 (2.6%) |
| Romania, n(%) | 2 | 217/3208 (6.8%) |
| Serbia, n(%) | 2 | 135/3208 (4.2%) |
| Spain, n(%) | 13 | 443/3208 (13.8%) |