

TABLE 3. SUMMARY OF BASIC FEATURES IN ADULT PATIENTS WITH A CARDIOMYOPATHY ACCORDING TO PREDEFINED SUB-GROUPS

	Probands (N=1929)	Relatives (N=395)	Pvalue	Incident (N=1335)	Prevalent (N=1774)	Pvalue	Pilot patient (N=1115)	Long-Term patient (N=2093)	Pvalue	North (N=543)	East (N=713)	West (N=481)	South (N=1427)	Pvalue
Age at diagnosis (years),														
Median (Q1;Q3)	50.0 (38.0; 59.0)	39.0 (24.0; 50.0)	<0.001	51.0 (40.0; 60.0)	47.0 (35.0; 57.0)	<0.001	46.0 (32.0; 58.0)	49.0 (38.0; 59.0)		49.0 (38.0; 60.0)	48.0 (37.0; 58.0)	50.0 (40.0; 57.0)	49.5 (37.0; 60.0)	0.384
N= (for age at diagnosis)	1728	180		1004	969		1092	2065		272	620	201	928	
Males, n(%)	1282/1929 (66.5%)	216/395 (54.7%)	<0.001	915/1335 (68.5%)	1117/1774 (63.0%)	0.001	696/1115 (62.4%)	1393/2093 (66.6%)	0.019	349/543 (64.3%)	472/713 (66.2%)	333/481 (69.2%)	908/1427 (63.6%)	0.138
Proband (vs relative)				883/992 (89.0%)	963/1242 (77.5%)	<0.001	184/391 (47.1%)	1745/1933 (90.3%)	<0.001	262/320 (81.9%)	601/630 (95.4%)	212/250 (84.8%)	810/1080 (75.0%)	<0.001
Familial disease, n(%)	524/1510 (34.7%)	389/389 (100.0%)	<0.001	285/993 (28.7%)	655/1434 (45.7%)	<0.001	423/912 (46.4%)	534/1551 (34.4%)	<0.001	141/358 (39.4%)	132/500 (26.4%)	108/396 (27.3%)	576/1165 (49.4%)	<0.001
Incident patient (vs prevalent)	883/1846 (47.8%)	109/388 (28.1%)	<0.001				326/1111 (29.3%)	1009/1998 (50.5%)	<0.001	143/534 (26.8%)	499/641 (77.8%)	184/476 (38.7%)	465/1414 (32.9%)	<0.001
Family screening as a reason for diagnosis, n(%)	68/1808 (3.8%)	264/387 (68.2%)	<0.001	95/1269 (7.5%)	246/1676 (14.7%)	0.001	168/1042 (16.1%)	180/1978 (9.1%)	<0.001	46/520 (8.8%)	36/669 (5.4%)	31/416 (7.5%)	235/1371 (17.1%)	<0.001
Rare disease*, n(%)	62/1929 (3.2%)	16/395 (4.1%)	0.400	54/1335 (4.0%)	75/1774 (4.2%)	0.800	69/1115 (6.2%)	65/2093 (3.1%)	<0.001	12/543 (2.2%)	21/713 (2.9%)	19/481 (4.0%)	82/1427 (5.7%)	<0.001
Presence of symptoms, n(%)	1613/1928 (83.7%)	313/395 (79.2%)	0.033	1165/1332 (87.5%)	1538/1769 (86.9%)	0.668	1107/1107 (100.0%)	1675/2093 (80.0%)	<0.001	512/543 (94.3%)	627/712 (88.1%)	456/475 (96.0%)	1143/1426 (80.2%)	<0.001
MRI, n(%)	468/1929 (24.3%)	140/395 (35.4%)	<0.001	427/1335 (32.0%)	506/1774 (28.5%)	0.037	507/1115 (45.5%)	437/2093 (20.9%)	<0.001	176/543 (32.4%)	139/713 (19.5%)	153/481 (31.8%)	474/1427 (33.2%)	<0.001
Holter ECG, n(%)	1005/1929 (52.1%)	254/395 (64.3%)	<0.001	716/1335 (53.6%)	976/1774 (55.0%)	0.443	753/1115 (67.5%)	999/2093 (47.7%)	<0.001	246/543 (45.3%)	464/713 (65.1%)	176/481 (36.6%)	857/1427 (60.1%)	<0.001
Exercise test, n(%)	582/1929 (30.2%)	169/395 (42.8%)	<0.001	377/1335 (28.2%)	722/1774 (40.7%)	<0.001	564/1115 (50.6%)	546/2093 (26.1%)	<0.001	206/543 (37.9%)	146/713 (20.5%)	254/481 (52.8%)	492/1427 (34.5%)	<0.001
Genetic testing performed, n(%)	596/1763 (33.8%)	227/373 (60.9%)	<0.001	228/1222 (18.7%)	782/1644 (47.6%)	<0.001	462/1044 (44.3%)	594/1913 (31.1%)	<0.001	175/524 (33.4%)	70/672 (10.4%)	169/408 (41.4%)	642/1315 (48.8%)	<0.001
Cardioverter defibrillator implanted, n(%)	482/1929 (25.0%)	101/395 (25.6%)	0.808	223/1335 (16.7%)	596/1774 (33.6%)	<0.001	316/1115 (28.3%)	516/2093 (24.7%)	0.023	148/543 (27.3%)	122/713 (17.1%)	179/481 (37.2%)	381/1427 (26.7%)	<0.001
Pacemaker implanted, n(%)	213/1919 (11.1%)	27/387 (7.0%)	0.015	104/1327 (7.8%)	208/1743 (11.9%)	<0.001	86/1077 (8.0%)	238/2092 (11.4%)	0.003	61/536 (11.4%)	56/704 (8.0%)	65/477 (13.6%)	136/1408 (9.7%)	0.010

* Rare disease: figures on pooled Pilot + LT phase populations