

NK cells in liver disease

Victoria Male, Kerstin A. Stegmann, Nicholas J. Easom and Mala K. Maini

5 **Dr Victoria Male (PhD)**

Affiliation: Division of Infection and Immunity, University College London
Postal address: Institute of Immunity and Transplantation, UCL Medical School, Rowland Hill Street, London NW3 2PF
Telephone: 0207 794 0500 x22463

10

Dr Kerstin Stegmann (PhD), Dr Nicholas Easom (MBBChir), Prof Mala Maini (F Med Sci)

Postal address: Division of Infection and Immunity, Rayne Building, 5 University Street, WC1E 6JF
Telephone: 020 3108 2170

15

20 **Abstract**

NK cells comprise one of the most abundant immune cell populations in human liver and the nature and functions of these cells have been a focus of recent interest. Here, we consider the possible roles of NK cells in diverse liver diseases, concentrating on data from patient studies. NK cells can be protective, killing virally infected and cancerous cells in the liver and limiting fibrosis by eliminating hepatic stellate cells. However, they can also be deleterious, contributing to pathology in viral hepatitis by killing hepatocytes and downregulating virus-specific T cell responses. It has recently emerged that a large fraction of hepatic NK cells constitute a distinct liver-resident subset and we highlight the need to distinguish between circulating and liver-resident NK cells in future studies. There is also a need for further investigation into how NK cells are influenced by the liver microenvironment and what scope there is to harness their immunotherapeutic potential.

35 **Natural killer cells**

Natural killer (NK) cells are immune cells of the lymphoid lineage that recognise and kill virally infected and malignant cells. NK cells also modulate the immune response by their production of cytokines, particularly IFN γ , and by their interactions with other immune cells, notably dendritic cells and T cells. Unlike T and B cells, NK cells do not undergo antigen receptor rearrangement but instead recognise their targets using a range of germ-line encoded receptors. Therefore, in contrast to T and B cells, NK cells make a rapid and robust response without the requirement for prior sensitisation [1-3].

NK cells were first described as recognising targets that lack MHC class I: the “missing self” response. They do this by expressing a variety of inhibitory receptors with a broad specificity for MHC class I, notably the inhibitory killer immunoglobulin-like receptors (KIRs) and CD94-NKG2A. When these are not engaged, the inhibitory signal is lifted leading to target cell killing. However, some cell types constitutively express low levels of MHC class I, and some virally infected cells evade recognition by NK cells by encoding molecules that mimic MHC class I. These phenomena limit the utility of target cell recognition based entirely on “missing self”. More recently, it has become appreciated that NK cells also express a number of activating receptors whose engagement promotes killing. One such receptor is NKG2D, whose ligands, MICA, MICB and the ULPBs, are upregulated on stressed and transformed cells, providing another way for NK cells to recognise their targets. NK cells also express a group of activating receptors known as “natural cytotoxicity receptors” (NCR) which include NKp30, NKp44 and NKp46. The ligands of these receptors are not yet well-defined, but they are known to recognise virally infected and cancerous cells. Binding to antibody-coated cells via the Fc receptor CD16 allows a subset of NK cells to mediate antibody-dependent cell-mediated cytotoxicity (ADCC). The balance between ligation of activating and inhibitory receptors determines the outcome of the interaction between an NK cell and a potential target, although the precise threshold for NK cell activation can be modulated by the cytokine milieu. Selected NK cell receptors and their ligands in humans are summarised in Table I.

Having recognised a target, an NK cell may kill it either by releasing cytotoxic granules or by ligating target cell death receptors. Cytotoxic granules contain perforin and granzymes. Perforin forms a pore in the target cell membrane allowing granzymes to enter the cytosol and induce apoptosis. Therefore, the ability of NK cells to degranulate and their expression of perforin and granzymes are often used as proxy measures for their cytotoxicity. NK cells may also promote target cell death via tumour necrosis factor (TNF) receptor superfamily molecules expressed by the target cell. Some NK cells express TNF superfamily molecules, such as TRAIL or Fas ligand, which bind to these receptors, triggering apoptosis in the target cell.

The functions of circulating NK cells, then, are well-established, and the mechanisms by which they achieve target cell recognition, killing and cytokine production are well-understood. However, many non-lymphoid organs are enriched in NK cells compared to the blood. The NK populations in these organs differ phenotypically from blood NK cells, and in some cases, notably in the uterus, these organ-specific NK cells are thought to have specialist physiological functions in their home tissue [4,5]. The liver contains a large number of NK cells, and these have recently been a focus of intense research interest.

80

NK cells in the liver

NK cells have long been recognised to be extremely abundant in human liver, ordinarily representing between 30% and 50% of liver lymphocytes [6-8]. Their abundance has fueled interest in their potential role in liver disease. The specialised hepatic circulation and surrounding parenchymal and non-parenchymal cells are important local influences on the features and functions of hepatic NK cells. The extensive narrow-lumen sinusoidal vasculature is lined by a fenestrated endothelium, allowing direct contact of intravascular lymphocytes with hepatocytes and stellate cells within the space of Disse [9]. Liver NK cells were originally described as “pit cells” that were noted to be in close contact with sinusoidal endothelial cells and Kupffer cells [10]; the latter have subsequently been shown to shape NK cells by differential cytokine production following TLR signalling [11].

Human and mouse livers both contain large NK cell populations with an “immature” phenotype [7,8,12]. In mice, these “immature” NK cells are now known to represent a separate, liver-resident population that is identified by its expression of CD49a and is less cytotoxic but has a greater capacity for TNF α and GM-CSF production than circulating NK cells [13,14]. In the light of these findings, recent attempts have been made to define an equivalent NK cell population in human liver. Some human livers contain a small CD49a⁺ NK cell population [15], but these differ from CD49a⁺ NK cells in mouse liver in a number of respects and may not be equivalent [16].

100

More promisingly, all human livers seem to contain a large NK cell population that is phenotypically distinct from NK cells found in the blood [17-20] (Table II). Experiments carried out in a transplant setting have shown that these cells are unable to exit the liver and are long-lived, and are therefore considered liver-resident [19]. The liver-resident NK cells appear less mature than circulating NK cells, on the basis of their expression of the maturation marker CD56, the terminal differentiation marker CD57 and by their expression of various NK cell receptors [17-20]. They express the C-type lectin CD69, which is associated with tissue-residence [5], as well as the chemokine receptors CCR5 and CXCR6, which can mediate liver homing and residency by recognition of their ligands CCL3, CCL5 and CXCL16, which are present in hepatic sinusoids [17-19].

110

Liver-resident NK cells express lower levels of perforin and granzyme B than their circulating counterparts, are less cytotoxic and less able to produce pro-inflammatory cytokines [17-19]. This is perhaps not unexpected, since it is in line with the immunotolerant nature of the liver, but it does suggest that liver-resident NK cells do not have the same function as circulating NK cells. The large number and tissue residence of these cells perhaps point to a tissue-specific function, although it is not yet clear what this might be. There is some evidence from mice that liver-resident NK cells are able to mediate immune memory responses to haptens and viral vaccinations [21,22]. This suggests the intriguing possibility that liver-resident NK cells in humans may be “memory” cells, a function that would be consistent with their longevity [19].

115

120

125 Regardless of their function, the discovery of these liver-resident NK cells has a number of
implications for the interpretation of older studies into human NK cells in liver disease. Firstly, it
should be noted that they express CD69. In the past, this was considered a marker of immune
cell activation, but is now known to also be associated with tissue-resident cells. Therefore,
130 studies which report an increase in the proportion of CD69⁺ intrahepatic NK cells could be
interpreted either as demonstrating activation of circulating NK cells, an expansion of the
resident compartment, or even a combination of the two. Furthermore, the existence of liver-
resident cells means that the behavior of NK cells isolated from blood will not necessarily
represent the responses of all the NK cells present in the liver. Indeed, circulating and liver-
resident NK cells may later be shown to have different, even opposing, roles in liver disease.

135 **NK cells in liver fibrosis**

The liver is exceptional in its regenerative capacity, but repeated destruction and regeneration
results in fibrosis, which can progress to cirrhosis. Fibrosis occurs largely as a result of
extracellular matrix production by activated hepatic stellate cells. It is a process common to all
chronic inflammatory diseases of the liver and, as a result, any role for NK cells in fibrosis is
likely to be relevant to a broad range of liver diseases.

140 A number of studies in mice have shown that chemically-induced liver fibrosis is exacerbated in
the absence of NK cells, suggesting that NK cells control fibrosis. This is a result of the greater
susceptibility of activated hepatic stellate cells to NK cell killing, an effect that may be mediated
by their reduced expression of MHC class I [23], increased expression of ligands for NKG2D
145 [24,25], NKp46 [26] or receptors for TRAIL [27], or some combination of these.

Consistent with the findings in mice, NK cells in humans also have a role in controlling liver
fibrosis. Human peripheral blood NK cells are able to kill hepatic stellate cells in culture and this
is dependent on NKp46 [26], NKG2A [28], NKG2D, TRAIL and FasL [29]. NK cells that are
150 cytotoxic [30], and specifically more effective at killing hepatic stellate cells [29], are associated
with a lower degree of fibrosis in chronic hepatitis C. Furthermore, increased NK cell infiltration
into the liver is associated with reduced fibrosis in primary sclerosing cholangitis [31].

155 **NK cells in viral hepatitis**

One of the major causes of liver disease is viral hepatitis brought about by the unrelated
Hepatitis viruses A, B, C and E. Each of these is considered below, although the majority of NK
cell studies have focused on Hepatitis B and C [32].

160 *Hepatitis B*

In acute Hepatitis B virus (HBV) infection, there is an early expansion of NK cells [33,34] but
their antiviral potential is temporarily impaired [35], coinciding with peak viraemia and a surge in
IL-10, which has suppressive effects on NK cells. Only when viral load, and therefore IL-10
165 levels, decrease, do NK cells upregulate TRAIL and regain the ability to produce pro-
inflammatory cytokines.

Studies of NK cell frequency and function in chronic hepatitis B (CHB) have been complicated
by the heterogeneity of patients cohorts, comprised of varying proportions in the different clinical
phases. Even the nomenclature and diagnostic criteria of these four phases (currently usually
170 known as immune-tolerant, immune-active, immune-control and reactivation), based on markers
such as eAg, viral load and ALT, are controversial [36-38]. NK cell frequency in the periphery
and the liver is highest in immune-tolerant patients, while immune-active patients have a

reduction in NK cells [39,40]. There is some disagreement on the precise way in which activating and inhibitory NK cell receptors are altered in the immune-tolerant phase, but the consensus seems to be that there is a general shift towards inhibition. In immune-active patients, the consensus points to peripheral and intrahepatic NK cells expressing more activating markers such as NCRs, TRAIL, CD69 and HLA-DR, and fewer inhibitory receptors [41-44], although one study reported increased expression of the inhibitory receptor NKG2A by peripheral NK cells in immune-active patients [45]. Consistent with a generalised activation, NK cells from immune-active patients have an increased ability to degranulate and produce cytokines compared to immune-tolerant patients and healthy controls. These changes might be attributable to increased proinflammatory cytokine production in the liver of immune-active patients [40].

A broad consensus from all studies is that NK cells in CHB have a functional dichotomy, with selective impairment in cytokine production but relative preservation of cytotoxicity [42-44,46]. IFN γ is key to the non-cytolytic immune control of HBV replication without excessive liver damage, so this reduction in IFN γ production is particularly significant [47]. The selective defects in NK cell function could be recapitulated by IL-10 treatment *in vitro*, whilst blockade of IL-10 and TGF- β led to restoration of IFN γ production in both circulating and intrahepatic NK cells from patients with CHB. A role for these immunosuppressive cytokines in shaping the NK cell functional profile is in line with their preponderance in the liver and increased expression in CHB [46]. Additional mechanisms proposed to contribute to the impaired NK cell effector function in CHB are their reduced levels of activating receptors [48] and impaired activation by plasmacytoid dendritic cells [49].

Patients in the eAg-negative reactivation phase experience temporary increases in viraemia and liver damage, known as “hepatic flares”, associated with increased serum IFN α and IL-8 levels, together with enhanced NK cell activation and TRAIL expression. This cytokine combination promotes TRAIL-mediated killing by upregulating TRAIL on NK cells, and altering the balance of death-inducing versus decoy TRAIL receptors on hepatocytes. Human NK cells are able to kill both hepatoma cells and primary human hepatocytes in a partially TRAIL-dependent manner. These findings provided a paradigm in which TRAIL⁺ NK cells, which are predominantly liver-resident [18], eliminate virally-infected hepatocytes and propagate immune-mediated liver damage [50].

While NK cells can play a protective role by eliminating virally infected hepatocytes or activated hepatic stellate cells, their potential to regulate immunopathology is complicated by their interaction with antiviral T cells [51] (Figure 1). NK cells selectively kill HBV-specific CD8⁺ T cells rather than T cells of other specificities and this susceptibility is partly determined by upregulation of the death-inducing receptor TRAIL-R2 by CD8⁺ T cells during HBV-specific responses. Expression of TRAIL-R2 is greatly enriched on intrahepatic CD8⁺ T cells, to a degree determined by viral load. The *in vivo* relevance of this is underscored by the finding that antiviral T cell responses from HBV-infected livers can be rescued following overnight TRAIL blockade [52]. This finding was supported by several studies in murine models showing a vital rheostat function for NK cells in regulating T cell responses in persistent viral infections [53,54]. Another example of NK cell/T cell crosstalk in viral hepatitis is the finding that MICA is expressed on HBV-specific and intrahepatic CD4⁺ T cells. This first demonstration of uninfected human T cells expressing an NKG2D ligand was supported by *in vitro* and *ex vivo* data suggesting that this could drive NK cell cytotoxicity [55]. Further results pointing to a role for NK cells in negatively regulating HBV-specific T cells came from a study of patients treated with nucleoside/nucleotide analogs (NUCs). NUC treatment reduces NK cell activation and TRAIL

expression [46]; reversion to a quiescent NK cell phenotype is associated with restoration of HBV-specific CD4⁺ T cells [56].

225 Interferon-alpha (IFN α) is another mainstay of treatment for CHB, providing an opportunity to study the *in vivo* impact of this cytokine on NK cells. Administration of therapeutic pegylated-IFN α to cohorts with either eAg-positive or eAg-negative CHB led to an increase in serum levels of the pro-proliferative cytokine IL-15, accompanied by a cumulative expansion of CD56^{bright} NK cells expressing TRAIL, NKp46 and HLA-DR [57,58]. Whereas NUCs were unable to reverse the defective NK cell IFN γ production characteristic of CHB [44,46,56], this could be restored by pegylated IFN α [57,58]. Surprisingly, pegylated-IFN α was also able to induce sustained changes in NK cell responsiveness to sequential treatment with NUCs [58]. These prolonged *in vivo* changes in NK cells are in line with emerging data on adaptive features of NK cells [3].

235 *Hepatitis C*

An early indication of the importance of NK cells in the outcome of Hepatitis C virus (HCV) infection came from genetic studies, revealing that patients expressing the inhibitory receptor KIR2DL3 together with its ligand HLA-C1 had an increased chance of spontaneous or treatment-induced HCV clearance [59,60]. Similarly, HCV-exposed uninfected individuals were more likely to have KIR2DL3-expressing NK cells with enhanced activation and effector function [61-63]. A number of studies have reported associations between NK cell receptor expression and function with outcomes in HCV infection, including disease activity as measured by ALT and AST [43,64], viral load [43,65-67] and outcome of treatment [66,68]. Broadly speaking, higher NK cell cytotoxicity and expression of activating receptors are associated with viral clearance and good responses to treatment, while higher expression of inhibitory receptors is linked to poor responses. Therefore, activated NK cells seem to be beneficial in the control of HCV.

250 In chronic HCV, the total frequency of NK cells and the proportion of the more cytotoxic CD56^{dim} subset of cells within the NK cell compartment are reduced [30,69,70]. There is no consensus on the expression of activating and inhibitory receptors by peripheral NK cells from chronic HCV patients, although the slim majority of studies report an overall increase in activating receptors [41,43,67,69,70]. This lack of consensus could potentially be attributed to differences in patient cohorts. Hepatic NK cells in chronic HCV also seem to be broadly activated, with one study reporting that intrahepatic NK cells express increased levels of activating receptors compared to peripheral NK cells [64] while another found increased expression of activating receptors by NK cells from HCV infected liver, compared to control liver [71].

260 NK cells in chronic HCV infection display reduced cytokine production, but sustained cytotoxicity, similar to the situation in CHB. Cytotoxic function of NK cells has been reported either to be increased [43] or unaltered in HCV patients compared to healthy controls [30,70]. The HCV-infected liver is enriched in NK cells expressing high levels of NKp46, which are highly cytotoxic, produce IFN γ and block HCV replication more efficiently than NKp46^{lo} cells. The frequency of NKp46^{hi} NK cells in the liver correlates inversely with viral load and fibrosis [72], in line with NK cells having an anti-fibrotic [29] and anti-viral activity [65]. On the other hand, NK cells from HCV infected patients have an impaired ability to produce IFN γ [43,64,73].

270 IFN α is a major regulator of NK cell activity during HCV infection, and a key determinant of outcome. *In vitro*, treatment with IFN α leads to a shift towards activation with increased TRAIL expression on both peripheral and intrahepatic NK cells [64,65,68] and increased degranulation in peripheral NK cells [64]. IFN α stimulation is necessary for NK cells to effectively recognise

HCV-replicating hepatoma cells and IFN α -stimulated NK cells challenged with these targets degranulate, produce cytokines and mediate cytotoxicity via TRAIL [65,74].

275
Until recently, administration of pegylated IFN α was the standard treatment for HCV and examination of NK cell responses in patients undergoing therapy sheds light on the importance of IFN α in NK cell responses to HCV *in vivo*, with TRAIL emerging as a critical effector of virus control. NK cells in acute HCV patients have elevated TRAIL expression, which increases
280 during IFN α therapy. Patients who cleared chronic HCV infection during IFN α treatment showed a greater upregulation of TRAIL compared to patients with ongoing chronic HCV infection. Furthermore, in patients with chronic HCV infection, TRAIL upregulation during the first 48h of IFN α treatment correlated with decreases in viral load [65]. Others studies have shown an upregulation of CD69, NKp30, NKp46 and NKG2A and downregulation of CD16, perforin and
285 NKG2D during IFN α treatment [66]. There is also a decreased frequency of NK cells expressing CXCR3 and CCR7, suggesting recruitment to the liver [68]. More recently, studies of blood and liver samples from HCV patients treated with the new interferon-free regimes (direct-acting antivirals) have demonstrated normalisation of the disease-associated NK cell phenotype and function.

290
Hepatitis A and E

The role of NK cells in Hepatitis A virus (HAV) has not been investigated in detail. One study suggested NK cells have the potential to control HAV infection, although this was carried out in HAV-infected fibroblasts rather than hepatocytes [75,76]. There is controversy over changes in
295 NK cell frequency during Hepatitis A [77,78] but elevated NK cell NKG2A expression correlates with ALT, suggesting a possible role for NK cells in acute liver damage [78].

In patients with acute Hepatitis E virus (HEV) infection, NK cells produce proinflammatory cytokines in response to stimulation faster than in patients with resolved infection [79]. NCR, NKG2D and CD69 expression increase during acute and resolved HEV infection compared to healthy controls [79]. These findings suggest that NK cells are activated in HEV infection. However, NK cell degranulation is impaired [79] and NK cell numbers decrease and remain low in resolved patients [80].

305
NK cells in alcoholic and non-alcoholic fatty liver disease

Alcoholic liver disease (ALD) and non-alcoholic fatty liver disease (NAFLD) are histologically similar disorders, both of which encompass a spectrum from simple steatosis to steatohepatitis to cirrhosis and finally hepatocellular carcinoma (HCC). Hepatocyte ballooning and immune cell infiltration are features of both ALD and NAFLD and the two disorders are likely to have a
310 number of immunological characteristics in common, including the role of NK cells.

Alcoholic liver disease

The role of NK cells in ALD has not yet been extensively investigated. In mice, chronic ethanol consumption reduces NK cell number and function by inhibiting their maturation. This in turn accelerates the development of liver fibrosis in the presence of other injurious stimuli [81,82]. In human peripheral blood, a similar reduction in NK cell frequency and cytotoxic function is observed in ALD patients, compared to controls [83].

320
Non-alcoholic fatty liver disease

The role of NK cells in NAFLD has been considered in more depth. To the extent that NAFLD is regarded as the hepatic manifestation of the metabolic syndrome, NK cells in adipose tissue are likely to be implicated in NAFLD through their involvement in altering glucose homeostasis [84-

325 86]. Similar to ALD patients, NK cell frequency and cytotoxic function is decreased in the
peripheral blood of obese patients, compared to lean controls [87]. Non-alcoholic steatohepatitis
(NASH) patients have increased NK cell numbers in their liver, compared to those with simple
steatosis or healthy controls, suggesting that the decrease in peripheral blood NK cell frequency
may be a result of their recruitment to the liver [88]. Notably, this study identified NK cells using
330 CD57, demonstrating that the increase is specifically in the CD57⁺ liver-infiltrating, and not the
CD57⁻ liver-resident, population, and supporting the idea that these NK cells have been
recruited from the blood. The study also reported increased expression of NKG2D ligands and
apoptotic hepatocytes in NASH patients, suggesting increased NKG2D-dependent activation of
these NK cells [88].

335 An increase in NK cells in the liver during NASH was also observed in a mouse model of the
disease, and again this was attributable to an increase in liver-infiltrating, rather than liver-
resident NK cells [89]. Intriguingly, in this study depletion of NK cells aggravated liver disease by
allowing macrophages to become polarised to an M2 phenotype. Therefore, although a number
of associations between NK cells and NAFLD have been reported, it is not yet clear if NK cells
340 are helpful or deleterious in the disease.

NK cells in autoimmune diseases of the liver

345 NK cells are implicated to some extent in the pathology of all three main autoimmune diseases
of the liver: autoimmune hepatitis (AIH), primary biliary cirrhosis (PBC, formerly primary biliary
cirrhosis), and primary sclerosing cholangitis (PSC).

Autoimmune hepatitis

350 AIH is characterised by hepatocyte necrosis associated with T cell and plasma cell infiltrates.
NK cells have been observed to interact with hepatocytes in AIH [90]. Different HLA alleles have
been associated with AIH in different ethnic groups [91,92], and recently in a Sardinian cohort
the activating receptor KIR2DS1 has been shown to be a risk factor for AIH, while presence of
the inhibitory receptor-ligand pairs KIR2DL3/HLA-C1 and KIR3DL1/HLA-Bw4 were protective
[93], implicating NK cell activation in disease.

355

Primary biliary cirrhosis

In PBC there is destruction of the small intrahepatic bile ducts, with contributions from both
adaptive and innate cells. NK cells are more numerous in both liver and blood in PBC, with
increased perforin expression and cytotoxicity [94]. This is likely to reflect recruitment of
360 peripheral blood NK cells to the liver via CXCR3 and CX3CR1 [95,96]. *In vitro*, NK cells can be
induced to kill autologous biliary epithelial cells via TRAIL, by cross-talk with TLR-3 and -4
stimulated macrophages [97]. Further, IFN γ produced by NK cells after co-culture with biliary
epithelial cells can induce cytotoxicity in autoreactive CD4⁺ T cells [98].

Primary sclerosing cholangitis

365 PSC is an autoimmune disorder with inflammation of all parts of the biliary tree and associated
T cell infiltrates. It is associated with increased NK cell numbers in the peripheral blood but not
the liver [99] and these NK cells are functionally impaired due to the action of TNF α [100]. A
number of lines of immunogenetic evidence point to the involvement of NK cell receptors and
370 ligands, and by extension NK cells themselves, in the progression of PSC. The development of
the disease is associated with certain alleles of the NKG2D ligands MICA and MICB [101,102],
and also with the absence of HLA-Bw4 and HLA-C2, which are ligands for inhibitory KIR, an
effect that seems to require homozygosity for MICA5.1 [103]. Furthermore, two SNPs in the

375 NKG2D gene are associated with development of cholangiocarcinoma in the context of PSC [104].

NK cells in liver cancer

380 Primary liver cancer is the third commonest cause of cancer death worldwide and the liver is also a major site of metastatic spread for many other malignancies. One major role of NK cells throughout the body is tumour surveillance and NK cells have the potential to play crucial roles in the immune response to both primary and metastatic liver cancers.

Hepatocellular carcinoma

385 HCC arises from hepatocytes in the context of pre-existing liver disease, most commonly viral hepatitis, ALD or NAFLD in the developed world, and is strongly associated with aflatoxin exposure in the developing world. Apart from HBV, which can be directly carcinogenic, HCC arises only in cirrhotic livers and is associated with inflammation in all settings, with systemic inflammation carrying a poor prognosis [105].

390 As in other malignancies, NK cells are functionally impaired in HCC. Tumour-infiltrating NK cells have reduced expression of granzymes and perforin and reduced cytotoxic potential compared to healthy controls, and are also reduced as a proportion of infiltrating lymphocytes compared with non-tumour liver tissue [106]. Tumour environments can subvert NK cell immunity through a variety of mechanisms, including increased numbers of immunosuppressive cells such as regulatory T cells and myeloid-derived suppressor cells and increased expression of immunoregulatory cytokines such as TGF β [107]. Changes in NK receptors or their ligands within the tumour milieu can also downregulate immunity. For example, loss of expression of the NKG2D ligands ULBP1 and MICA by HCC is associated with a poor prognosis [108,109]. The SNP at rs2596542 is associated with MICA shedding in HCC. Among hepatitis C patients, those with the A/A genotype, which is associated with low soluble MICA, are at higher risk of HCC, whereas among hepatitis B patients, those with the GG genotype are at higher risk [110-112]. The differential effects of this SNP may be due to differences in induction and processing of MICA between the two viruses. The high-affinity allele MICA-129Met is associated with an increased risk of HCC in the context of HBV [112], providing further evidence to suggest that NKG2D may play a role in both HCC surveillance and control.

410 Conversely, there is some immunogenetic evidence to suggest that high basal NK cell activity may be deleterious. KIR-HLA interactions which allow the development of highly responsive ("licensed") NK cells and activating KIR in the presence of appropriate HLA ligands are associated with increased progression to HCC in the context of HBV [113]. Similarly, different murine models have pointed to tumorigenic and well as protective roles for the NKG2D pathway in HCC [114]. Therefore it remains to be seen whether increasing NK cell activation and/or promoting NKG2D will be helpful or harmful in HCC.

415 Attempts to overcome tumour evasion strategies and harness NK cells for immunotherapy of HCC are now being tested in the clinic. One strategy is enhancing the function of endogenous NK cells; an example is the use of an antibody that blocks engagement of the inhibitory receptors KIR2DL-1/2/3, which is currently being tried, in combination with PD-1 blockade, in patients with advanced solid tumours (NCT01714739). Alternatively, adoptive cell therapy can be carried out, using infusions of autologous or allogeneic NK cells, expanded, activated or genetically modified to optimise their anti-tumour capacity [107]. Since NK cells may function better to prevent metastases than to penetrate established solid tumours, several ongoing trials are using allogeneic NK adoptive immunotherapy once the HCC has been resected or

425 explanted (NCT01147380, NCT02008929). Previous *in vitro* studies suggested NK cells
extracted from donor liver perfusates may be a promising source for such adoptive therapy
because of an enhanced capacity to kill HCC following IL-2 activation [7]. However, peripheral
NK cells can also be rendered effective by expansion using the NK cell-stimulating cell line
430 K562-mb15-41BBL or by transduction with the chimeric receptor NKG2D-CD3z-DAP10 [115].

430 *Cholangiocarcinoma*

Cholangiocarcinoma is a primary malignancy of the biliary tree, and is closely associated with
inflammation of the biliary system, primarily due to PBC, PSC or infection with liver flukes. In
mice, cytokine induced killer cells and IFN γ are associated with control of subcutaneously
435 injected cholangiocarcinoma cells [116,117]. In humans, two SNPs in the NKG2D gene are
associated with the development of cholangiocarcinoma in the context of PSC [104] but it
remains to be seen whether this effect is NK cell-mediated.

Liver metastases

440 In most major malignancies including those of the lung, breast and GI tract, spread to the liver is
common. In mice, NK cells are well-known to have a role in the control of tumour metastases
and intrahepatic NK cells control metastases via TRAIL [118,119]. The role of NK cells in
metastases to the liver in humans is best studied in colorectal cancer, where tumours resected
445 from the liver are available for investigation. In one study, infiltrating T cells, but not NK cells,
were associated with a good prognosis [120], and in other studies tumour infiltrating CD8⁺ T
cells have been shown to contribute to good prognosis [121,122]. In many solid tumours, NK
cells become functionally impaired and unable to lyse tumour cells. These autologous NK cells
may represent an effector population that could be recruited therapeutically to target metastases
450 *in vivo*, although cytokine infusion approaches have so far been limited by clinical toxicity [123].
Combining cytokine activation of NK cells with ADCC using monoclonal antibodies that engage
CD16, such as trastuzumab for HER2+ metastatic breast cancer, may be a more promising
approach [124]. Lenalidomide, which decreases the activation threshold of NK cells via an effect
on actin remodelling [125], did not induce clinical response in metastatic renal carcinoma
455 including in patients with liver metastases as monotherapy, but could be combined with ADCC-
inducing antibodies, as in various haematological malignancies [126].

NK cells in liver transplantation

460 Liver transplantation is the only definitive treatment for patients suffering from end-stage liver
disease, but its success is heavily dependent on a number of immune-mediated factors, which
may involve the action of NK cells. One early complication of transplantation is ischaemia-
reperfusion (IR) injury, in which hypoxic organ damage is exacerbated following the re-
establishment of blood supply and oxygen delivery. Acute rejection occurs days to weeks after a
transplant and is primarily mediated by alloreactive T cells, whereas chronic rejection can occur
465 up to years later, is characterised by occlusion of the blood vessels leading to ischaemia, and is
a result of mixed mechanisms, including low-grade T cell responses and anti-graft antibody.

The role of NK cells in IR injury during liver transplant has not yet been investigated in humans,
but there is evidence from mouse models that NK cells are involved in exacerbating this
470 disease. Treatment of mice with the NK cell depleting antibody NK1.1 reduces hepatic IR injury
[127] and CD39-deficient mice, in which NK cells are sub-functional, are less susceptible to
hepatic IR injury than wild-type controls [128].

475 NK cell-mediated skin graft rejection occurs in mice [129] so it is possible that some component
of liver transplant rejection may be NK cell-mediated. In rats, recipient NK cells are rapidly

recruited to the allogeneic transplanted liver. These cells are the main producers of IFN γ in the transplanted liver, and antibody depletion of NK cells can prolong the survival of the allograft [130]. During human liver transplantation, recipient NK cells are also rapidly recruited to the graft [19]. A number of immunogenetic studies have attempted to determine whether recipient NK cells may be involved in rejection of the transplanted liver, with mixed results. One study of 100 transplants showed that matching of the main HLA ligand recognised by NK cells, HLA-C, between donor and recipient, reduced the likelihood of acute rejection, with recipients who expressed no HLA-C group 2 alleles further protected from acute rejection [131]. However, later, larger studies were unable to replicate these results [132,133]. A study of HLA-C effect on chronic rejection of 595 liver transplants showed that the presence of at least one HLA-C group 2 allele in the donor was protective [132], but again this finding could not be replicated in a later study [133]. It is unclear, then, if NK cells are important mediators of either acute or chronic rejection in liver transplantation.

490

Future directions

NK cells are one of the most prevalent immune cell types in the human liver and are implicated as players in many liver diseases (Figure 2). However, in the healthy liver and most disease settings, their dominant roles remain to be clarified. Murine models are an important adjunct, but not a substitute, for studies in humans, especially given some notable differences in NK cells between mouse and man [16]. Researchers in this field are fortunate to have access to human liver tissue from diagnostic biopsies, along with transplant perfusates, metastatic resections and cirrhotic explants, giving more access to NK cells at the site of disease than is possible for most human organs. Nevertheless, the demanding logistics of obtaining such samples has meant that the majority of studies are conducted on peripheral blood; this is becoming more of an issue as non-invasive diagnostic methods start to replace liver biopsies. Recent studies revealing large liver-resident NK cell populations that do not recirculate [19] underscore the importance of continued access to liver sampling, exemplified by a recent study tracking the kinetics of NK cell responses by an alternative approach of repeated fine needle aspiration [134].

505

Future studies will need to analyse liver-resident and non-resident NK subsets separately, since prior analyses of the whole intrahepatic fraction may have masked important disease-associated changes. Another factor that may have confounded interpretation of results in previous studies is CMV serostatus; it is now clear that prior CMV infection is a major driver of NK cell adaptive differentiation [135] that needs to be controlled for in studies of diverse infections and liver diseases. In mice, the liver is a site where “memory” NK cells accumulate [21,22]. Whether this is the case in humans remains to be established but elegant work in HCV has already revealed another “adaptive” feature of NK cells: their capacity for mediating fine specificity through altered KIR binding to subtle variants of peptide/MHC class I complexes, impairing NK cell inhibition [136,137]. It will be interesting to consider precise NK cell target selectivity in other settings.

520

In addition to analysing freshly isolated hepatic NK cells, enhanced technology for multiparameter *in situ* immunostaining will facilitate improved studies of their anatomical localisation and interactions in liver sections. More in-depth mechanistic studies of how NK cells kill their targets in the liver, and how other liver cell types influence hepatic NK cells, will take advantage of emerging 3D models. Such scaffold systems constitute exciting progress in the attempt to recapitulate the complexity of the liver microanatomy and to allow analysis of NK cell cross-talk with, for example, primary hepatocytes and hepatic stellate cells, rather than immortalised cell lines [138]. Ongoing studies are starting to address the profound influence of local factors such as hypoxia and nutrient metabolism on NK cells in the unique liver niche.

525

Initial work suggests that exposure of NK cells to hypoxic conditions mimicking those present in the liver (which receives more than 70% of its blood supply from the portal vein) has divergent effects on function: while low oxygen tensions prevent activating receptors driving target cell killing, they maintain ADCC by NK cells [139]. As such, it may be necessary to revisit ideas about NK cell function in the liver that are based on experiments carried out at atmospheric oxygen tensions. Hypoxia may also interact with the unique cytokine milieu of the liver, since TGF β is able to inhibit the mTOR-dependent metabolic reprogramming central to IL-15-induced NK cell responses [140-142]. It will also be necessary to examine interactions between hypoxia and infection. Co-culturing NK cells from HCV patients with HCV-replicating hepatocytes or with stellate cells revealed that hypoxia constrained their antiviral capacity but not their anti-fibrotic activity [143] and one of the metabolic adaptations of NK cells exposed to hypoxia, HIF-1 α -induced glycolytic gene transcription [144], appears to be defective in HCV [143].

Studies addressing these novel aspects of NK cell biology in state-of-the art *in vitro* and *in vivo* models will enhance our understanding of their diverse roles in liver disease and the scope for immunotherapeutic modification. Meanwhile NK cells are already reaching the clinic for primary and secondary liver cancers and these trials will provide powerful insights into their potential as therapeutic tools.

530

535

540

545

Acknowledgements

Victoria Male is supported by a Royal Society/Wellcome Trust Sir Henry Dale Fellowship WT 105677. Mala Maini is supported by a Wellcome Trust Investigator Award.

550

References

1. Vivier E, Tomasello E, Baratin M, Walzer T, Ugolini S. Functions of natural killer cells. *Nat Immunol.* 2008; 9:503-510.
2. Lanier LL. Up on the tightrope: natural killer cell activation and inhibition. *Nat Immunol.* 2008; 9:495-502.
3. Vivier E, Raulet DH, Moretta A et al. Innate of adaptive immunity? The example of natural killer cells. *Science* 2011; 331:44-49.
4. Sojka DK, Tian Z, Yokoyama W. Tissue-resident natural killer cells and their potential diversity. *Semin Immunol.* 2014; 26:127-131
5. Melsen JE, Lugthart G, Lankester AC, Schilham MW. Human circulating and tissue-resident CD56(bright) natural killer cell populations. *Front Immunol.* 2016; 7:262
6. Doherty DG, Norris S, Madrigal-Estebas L, et al. The human liver contains multiple populations of NK cells, T cells, and CD3+CD56+ natural T cells with distinct cytotoxic activities and Th1, Th2, and Th0 cytokine secretion patterns. *J Immunol.* 1999; 163:2314-2321.
7. Ishiyama K, Ohdan H, Ohira M, Mitsuta H, Arihiro K, Asahara T. Difference in cytotoxicity against hepatocellular carcinoma between liver and periphery natural killer cells in humans. *Hepatology.* 2006; 43:362-372.
8. Burt BM, Plitas G, Zhao Z, et al. The lytic potential of human liver NK cells is restricted by their limited expression of inhibitory killer Ig-like receptors. *J Immunol.* 2009; 183:1789-1796.
9. Guidotti LG, Inverso D, Sironi L, et al. Immunosurveillance of the liver by intravascular effector CD8(+) T cells. *Cell* 2015; 161:486-500.
10. Wisse E, van't Noorende JM, van der Meulen J, Daems WT. The pit cell: description of a new type of cell occurring in rat liver sinusoids and peripheral blood. *Cell Tissue Res.* 1976; 173:423-435.
11. Tu Z, Bozorgzadeh A, Pierce RH, Kurtis J, Crispe IN, Orloff MS. TLR-dependent cross-talk between human Kupffer cells and NK cells. *J Exp Med.* 2008; 205:233-244.
12. Gordon SM, Chaix J, Rupp LJ, et al. The transcription factors T-bet and Eomes control key checkpoints of natural killer cell maturation. *Immunity.* 2012; 36:55-67.
13. Daussy C, Faure F, Mayol K, et al. T-bet and Eomes instruct the development of two distinct natural killer cell lineages in the liver and in the bone marrow. *J Exp Med.* 2014; 211:563-577.
14. Sojka DK, Plougastel-Douglas B, Yang L, et al. Tissue-resident natural killer (NK) cells are cell lineages distinct from thymic and conventional splenic NK cells. *Elife.* 2014;3:e01659.
15. Marquardt N, Béziat V, Nyström S, et al. Cutting edge: identification and characterization of human intrahepatic CD49a+ NK cells. *J Immunol.* 2015; 194:2467-2471.
16. Male V. Liver-resident NK cells: the human factor. *Trends Immunol.* 2017; doi 10.1016/j.it.2017.02.008. [Epub ahead of print]
17. Hudspeth K, Donadon M, Cimino M, et al. Human liver-resident CD56(bright)/CD16(neg) NK cells are retained within hepatic sinusoids via the engagement of CCR5 and CXCR6 pathways. *J Autoimmun.* 2016; 66:40-50.
18. Stegmann KA, Robertson F, Hansi N, et al. CXCR6 marks a novel subset of T-bet(lo)Eomes(hi) natural killer cells residing in human liver. *Sci Rep.* 2016; 6:26157.

19. Cuff AO, Robertson FP, Stegmann KA, et al. Eomeshi NK cells in human liver are long-lived and do not recirculate but can be replenished from the circulation. *J Immunol.* 2016; 197: 4283-4291.
- 600 20. Harmon C, Robinson MW, Fahey R, et al. Tissue-resident Eomes(hi) T-bet(lo) CD56(bright) NK cells with reduced proinflammatory potential are enriched in the adult human liver. *Eur J Immunol.* 2016; 46:2111-2120.
21. Paust S, Gill HS, Wang BZ, et al. Critical role for the chemokine receptor CXCR6 in NK cell-mediated antigen-specific memory of haptens and viruses. *Nat Immunol.* 2010; 605 11:1127-1135
22. Peng H, Jiang X, Chen Y, et al. Liver-resident NK cells confer adaptive immunity in skin-contact inflammation. *J Clin Invest.* 2013; 123:1444-1456.
23. Radaeva S, Sun R, Jaruga B, Nguyen VT, Tian Z, Gao B. Natural killer cells ameliorate liver fibrosis by killing activated stellate cells in NKG2D-dependent and tumor necrosis factor-related apoptosis-inducing ligand-dependent manners. *Gastroenterology.* 2006; 610 130:435-52.
24. Radaeva S, Wang L, Radaev S, Jeong WI, Park O, Gao B. Retinoic acid signaling sensitizes hepatic stellate cells to NK cell killing via upregulation of NK cell activating ligand RAE1. *Am J Physiol Gastrointest Liver Physiol.* 2007; 293:G809-816.
- 615 25. Krizhanovsky V, Yon M, Dickins RA, et al. Senescence of activated stellate cells limits liver fibrosis. *Cell.* 2008; 134:657-667.
26. Gur C, Doron S, Kfir-Erenfeld S, et al. NKp46-mediated killing of human and mouse hepatic stellate cells attenuates liver fibrosis. *Gut.* 2012; 61:885-893.
27. Melhem A, Muhanna N, Bishara A, et al. Anti-fibrotic activity of NK cells in experimental liver injury through killing of activated HSC. *J Hepatol.* 2006; 45:60-71.
- 620 28. Muhanna N, Abu Tair L, Doron S, et al. Amelioration of hepatic fibrosis by NK cell activation. *Gut.* 2011; 60:90-98.
29. Glässner A, Eisenhardt M, Krämer B, et al. NK cells from HCV-infected patients effectively induce apoptosis of activated primary human hepatic stellate cells in a TRAIL-, FasL- and NKG2D-dependent manner. *Lab Invest.* 2012; 92:967-977.
- 625 30. Morishima C, Paschal DM, Wang CC, et al. Decreased NK cell frequency in chronic hepatitis C does not affect ex vivo cytolytic killing. *Hepatology.* 2006; 43:573-580.
31. Berglin L, Bergquist A, Johansson H, et al. In Situ Characterization of Intrahepatic Non-Parenchymal Cells in PSC Reveals Phenotypic Patterns Associated with Disease Severity. *PLoS ONE.* 2014; 9:e105375
- 630 32. Park SH, Rehermann B. Immune responses to HCV and other hepatitis viruses. *Immunity.* 2014; 40:13-24
33. Webster GJ, Reignat S, Maini MK, et al. Incubation phase of acute hepatitis B in man: dynamic of cellular immune mechanisms. *Hepatology.* 2000; 32:1117-1124
- 635 34. Fisicaro P, Valdatta C, Boni C, et al. Early kinetics of innate and adaptive immune responses during hepatitis B virus infection. *Gut.* 2009; 58:974-982.
35. Dunn C, Peppas P, Khanna P, et al. Temporal analysis of early immune responses in patients with acute hepatitis B virus infection. *Gastroenterology.* 2009; 137:1289-300.
- 640 36. Bertolotti A, Maini MK, Ferrari C. The host-pathogen interaction during HBV infection: immunological controversies. *Antivir Ther.* 2010; Suppl 3:15-24.

37. Rehermann B. Pathogenesis of chronic viral hepatitis: differential roles of T cells and NK cells. *Nat Med.* 2013; 19:859-868
38. Bertoletti A, Kennedy PT. The immune tolerant phase of chronic HBV infection: new perspectives on an old concept. *Cell Mol Immunol.* 2015; 12:258-263
- 645 39. Sprengers D, van der Molen RG, Kusters JG, et al. Different composition of intrahepatic lymphocytes in the immune-tolerance and immune-clearance phase of chronic hepatitis B. *J Med Virol.* 2006; 78:561–568.
40. Zhang Z, Zhang S, Zou Z, et al. Hypercytolytic activity of hepatic natural killer cells correlates with liver injury in chronic hepatitis B patients. *Hepatology*, 2011; 53:73–85.
- 650 41. Nattermann J, Feldmann G, Ahlenstiel G, et al. Surface expression and cytolytic function of natural killer cell receptors is altered in chronic hepatitis C. *Gut.* 2006; 55:869–877.
42. Li Y, Wang JJ, Gao S, et al. Decreased peripheral natural killer cells activity in the immune activated stage of chronic hepatitis B. *PLoS ONE*, 2014; 9:e86927.
- 655 43. Oliviero B, Varchetta S, Paudice E, et al. Natural killer cell functional dichotomy in chronic hepatitis B and chronic hepatitis C virus infections. *Gastroenterology.* 2009; 137:1151-1160.
44. Tjwa ETTL, van Oord GW, Hegmans JP, Janssen HLA, Woltman AM. Viral load reduction improves activation and function of natural killer cells in patients with chronic hepatitis B. *J Hepatol.* 2011 54:209–218.
- 660 45. Li F, Wei H, Wei H, et al. Blocking the natural killer cell inhibitory receptor NKG2A increases activity of human natural killer cells and clears hepatitis B virus infection in mice. *Gastroenterology* 2013 144:392-401.
46. Peppia D, Micco L, Javaid A, et al. Blockade of Immunosuppressive Cytokines Restores NK Cell Antiviral Function in Chronic Hepatitis B Virus Infection. *PLoS Pathog.* 2010; 6:e1001227–10.
- 665 47. Guidotti LG, Chisari FV. Immunobiology and pathogenesis of viral hepatitis. *Annu Rev Pathol.* 2006; 1:23-61
48. Sun C, Fu B, Gao Y, et al. TGF- β 1 down-regulation of NKG2D/DAP10 and 2B4/SAP expression on human NK cells contributes to HBV persistence. *PLoS Pathog.* 2012; 8: e1002594.
- 670 49. Shi CC, Tjwa ET, Biestra PJ, et al. Hepatitis B virus suppresses the functional interaction between natural killer cells and plasmacytoid dendritic cells. *J Viral Hepat.* 2012; 19:e26-33.
- 675 50. Dunn C, Brunetto M, Reynolds G, et al. Cytokines induced during chronic hepatitis B virus infection promote a pathway for NK cell-mediated liver damage. *J Exp Med.* 2007; 204: 667–680.
51. Maini MK, Peppia D. NK cells: a double-edged sword in chronic hepatitis B virus infection. *Front Immunol.* 2013; 4:57.
- 680 52. Peppia D, Gill US, Reynolds G, et al. Up-regulation of a death receptor renders antiviral T cells susceptible to NK cell-mediated deletion. *J Exp Med.* 2013; 210: 99–114.
53. Chrome SQ, Lang PA, Lang KS, Ohashi PS. Natural killer cells regulate diverse T cell responses. *Trends Immunol.* 2013; 34:342-349.
- 685 54. Waggoner SN, Reighard SD, Gyurova IE, et al. Roles of natural killer cells in antiviral immunity. *Curr Opin Virol.* 2016; 16:15-23.

55. Huang WC, Easom, NJ, Tang XZ, et al. T Cells Infiltrating Diseased Liver Express Ligands for the NKG2D Stress Surveillance System. *J Immunol.* 2016; 160:1313.
56. Boni C, Lampertico P, Talamona L, et al. Natural killer cell phenotype modulation and natural killer/T-cell interplay in nucleos(t)ide analogue-treated hepatitis e antigen-negative patients with chronic hepatitis B. *Hepatology.* 2015; 62:1697-1709.
- 690 57. Micco L, Peppia D, Loggi E, et al. Differential boosting of innate and adaptive antiviral responses during pegylated-interferon-alpha therapy of chronic hepatitis B. *J Hepatol.* 2013; 58:225–233.
58. Gill US, Peppia D, Micco L, et al. Interferon Alpha Induces Sustained Changes in NK Cell Responsiveness to Hepatitis B Viral Load Suppression In Vivo. *PLoS Pathog.* 2016; 12:e1005788.
- 695 59. Khakoo SI, Thio CL, Martin MP, et al. HLA and NK cell inhibitory receptor genes in resolving hepatitis C virus infection. *Science.* 2004; 305:872-874.
60. Knapp S, Warshaw U, Ho KM, et al. A polymorphism in IL28B distinguishes exposed, uninfected individuals from spontaneous resolvers of HCV infection. *Gastroenterology.* 2011; 141:320-325
- 700 61. Golden-Mason L, Cox AL, Randall JA, Cheng L, Rosen HR. Increased natural killer cell cytotoxicity and NKp30 expression protects against hepatitis C virus infection in high-risk individuals and inhibits replication in vitro. *Hepatology.* 2010. 52:1581-1589.
- 705 62. Sugden PB, Cameron B, Mina M, et al. Protection against hepatitis C infection via NK cells in highly-exposed uninfected injecting drug users. *J Hepatol.* 2014; 61:738-745.
63. Werner JM, Heller T, Gordon AM, et al. Innate immune responses in hepatitis C virus-exposed healthcare workers who do not develop acute infection. *Hepatology.* 2013; 58:1621-1631.
- 710 64. Ahlenstiel G, Titerence RH, Koh C, et al. Natural killer cells are polarized toward cytotoxicity in chronic hepatitis C in an interferon-alfa-dependent manner. *Gastroenterology.* 2010; 138:325–335.
65. Stegmann KA., Björkström NK, Veber H, et al. Interferon-alpha-induced TRAIL on natural killer cells is associated with control of hepatitis C virus infection. *Gastroenterology.* 2010; 138:1885–1897.
- 715 66. Oliviero B, Mele D, Degasperis E, et al. Natural killer cell dynamic profile is associated with treatment outcome in patients with chronic HCV infection. *J Hepatol.* 2013; 59:38–44.
67. Golden-Mason L, Bambha KM, Cheng L, et al. Natural killer inhibitory receptor expression associated with treatment failure and interleukin-28B genotype in patients with chronic hepatitis C. *Hepatology.* 2011; 54:1559–1569.
- 720 68. Ahlenstiel G, Edlich B, Hogdal LJ, et al. Early changes in natural killer cell function indicate virologic response to interferon therapy for hepatitis C. *Gastroenterology.* 2011; 141:1231–1239
- 725 69. Bonorino P, Ramzan M, Camous X, et al. Fine characterization of intrahepatic NK cells expressing natural killer receptors in chronic hepatitis B and C. *J Hepatol.* 2009; 51:458–467.
70. De Maria A, Fogli M, Mazza S, et al. Increased natural cytotoxicity receptor expression and relevant IL-10 production in NK cells from chronically infected viremic HCV patients. *Eur J Immunol.* 2007; 37:445–455.
- 730 71. Varchetta S, Mele D, Mantovani S, et al. Impaired intrahepatic natural killer cell

- cytotoxic function in chronic hepatitis C virus infection. *Hepatology*. 2012; 56:841–849.
72. Krämer B, Körner C, Kebschull M, et al. Natural killer p46High expression defines a natural killer cell subset that is potentially involved in control of hepatitis C virus replication and modulation of liver fibrosis. *Hepatology*. 2012; 56:1201–1213.
- 735
73. Edlich B, Ahlenstiel G, Zabaleta Azpiroz A, et al. Early changes in interferon signaling define natural killer cell response and refractoriness to interferon-based therapy of hepatitis C patients. *Hepatology*. 2012; 55:39–48.
74. Stegmann KA, Björkström NK, Ciesek S, et al. Interferon α -stimulated natural killer cells from patients with acute hepatitis C virus (HCV) infection recognize HCV-infected and uninfected hepatoma cells via DNAX accessory molecule-1. *J Infect Dis*. 2012; 205:1351–1362.
- 740
75. Kurane I, Binn LN, Bancroft WH, Ennis FA. Human lymphocyte responses to hepatitis A virus-infected cells: interferon production and lysis of infected cells. *J Immunol*. 1985; 135:2140–2144.
- 745
76. Baba M, Hasegawa H, Nakayabu M, Fukai K, Suzuki, S. Cytolytic activity of natural killer cells and lymphokine activated killer cells against hepatitis A virus infected fibroblasts. *J Clin Lab Immunol*. 1993; 40:47–60.
77. Müller C, Gödl I, Göttlicher J, Wolf HM, Eibel MM. Phenotypes of peripheral blood lymphocytes during acute hepatitis A. *Acta Paediatr Scand*. 1991; 80:931–937.
- 750
78. Cho, H. Phenotypic characteristics of natural killer cells in acute hepatitis. *J Microbiol*. 2013; 51:247-251.
79. Das R, Tripathy, A. Increased expressions of NKp44, NKp46 on NK/NKT-like cells are associated with impaired cytolytic function in self-limiting hepatitis E infection. *Med Microbiol Immunol*. 2014; 203:303–314
- 755
80. Srivastava R, Aggarwal R, Bhagat MR, Chowdhury A, Naik S. Alterations in natural killer cells and natural killer T cells during acute viral hepatitis E. *J Viral Hepat*. 2008; 15:910–916.
81. Jeong WI, Park O, Gao B. Abrogation of the antifibrotic effects of natural killer cells/interferon-gamma contributes to alcohol acceleration of liver fibrosis. *Gastroenterology*. 2008; 134:248-258.
- 760
82. Zhang F, Little A, Zhang H. Chronic alcohol consumption inhibits peripheral NK cell development and maturation by decreasing the availability of IL-15. *J Leukoc Biol*. 2016;. pii: jlb.1A0716-298RR.
- 765
83. Laso FJ, Almeida J, Torres E, Vaquero JM, Marcos M, Orfao A. Chronic alcohol consumption is associated with an increased cytotoxic profile of circulating lymphocytes that may be related with the development of liver injury. *Alcohol Clin Exp Res*. 2010; 34:876-885.
84. O'Rourke RW, Meyer KA, Neeley CK, et al. Systemic NK cell ablation attenuates intra-abdominal adipose tissue macrophage infiltration in murine obesity. *Obesity (Silver Spring)*. 2014; 22:2109-2114.
- 770
85. Wensveen FM, Jelenčić V, Valentić S, et al. NK cells link obesity-induced adipose stress to inflammation and insulin resistance. *Nat Immunol*. 2015;16:376-385.
86. Lee BC, Kim MS, Pae M, et al. Adipose Natural Killer Cells Regulate Adipose Tissue Macrophages to Promote Insulin Resistance in Obesity. *Cell Metab*. 2016; 23:685-698.
- 775
87. O'Shea D, Cawood TJ, O'Farrelly C, Lynch L. Natural killer cells in obesity: impaired function and increased susceptibility to the effects of cigarette smoke. *PLoS One*. 2010; 5:e8660.

- 780 88. Kahraman A, Schlattjan M, Kocabayoglu P, et al. Major histocompatibility complex class I-related chains A and B (MIC A/B): a novel role in nonalcoholic steatohepatitis. *Hepatology*. 2010; 51:92-102.
89. Tosello-Tramont AC, Krueger P, Narayanan S, Landes SG, Leitinger N, Hahn YS. NKp46(+) natural killer cells attenuate metabolism-induced hepatic fibrosis by regulating macrophage activation in mice. *Hepatology*. 2016; 63:799-812.
- 785 90. Kaneda K, Kurioka N, Seki S, Wake K, Yamamoto S. Pit cell-hepatocyte contact in autoimmune hepatitis. *Hepatology*. 1984; 4:955-958.
91. Donaldson PT, Doherty DG, Hayllar KM, McFarlane IG, Johnson PJ, Williams R. Susceptibility to autoimmune chronic active hepatitis: human leukocyte antigens DR4 and A1-B8-DR3 are independent risk factors. *Hepatology*. 1991; 13:701-706.
- 790 92. Muratori P, Czaja A-J, Muratori L, et al. Genetic distinctions between autoimmune hepatitis in Italy and North America. *World J Gastroenterol*. 2005; 11:1862-1866.
93. Littera R, Chessa L, Onali S, et al. Exploring the Role of Killer Cell Immunoglobulin-Like Receptors and Their HLA Class I Ligands in Autoimmune Hepatitis. *PLoS ONE*. 2016; 11:e0146086-14
- 795 94. Chuang Y-H, Lian Z-X, Tsuneyama K, et al. Increased killing activity and decreased cytokine production in NK cells in patients with primary biliary cirrhosis. *J Autoimmun*. 2006; 26:232-240.
95. Chuang Y-H, Lian Z-X, Cheng C-M, et al. Increased levels of chemokine receptor CXCR3 and chemokines IP-10 and MIG in patients with primary biliary cirrhosis and their first degree relatives. *J Autoimmun*. 2005; 25:126-132.
- 800 96. Isse K, Harada, K, Zen Y, et al. Fractalkine and CX3CR1 are involved in the recruitment of intraepithelial lymphocytes of intrahepatic bile ducts. *Hepatology*. 2005; 41:506-516.
97. Shimoda S, Harada K, Niuro H, et al. Interaction between Toll-like receptors and natural killer cells in the destruction of bile ducts in primary biliary cirrhosis. *Hepatology*. 2011; 53:1270-1281.
- 805 98. Shimoda S, Hisamoto S, Harada K, et al. Natural killer cells regulate T cell immune responses in primary biliary cirrhosis. *Hepatology*. 2015; 62:1817-1827.
99. Hata K, Zhang XR, Iwatsuki S, Van Thiel DH, Herberman RB, Whiteside TL. Isolation, phenotyping, and functional analysis of lymphocytes from human liver. *Clin Immunol Immunopathol*. 1990; 56:401-419.
- 810 100. Bo X, Broome U, Remberger M, Sumitran-Holgersson S. Tumour necrosis factor alpha impairs function of liver derived T lymphocytes and natural killer cells in patients with primary sclerosing cholangitis. *Gut*. 2001; 49:131-141.
- 815 101. Norris S, Kondeatis E, Collins R, Stephens H. Mapping MHC-encoded susceptibility and resistance in primary sclerosing cholangitis: the role of MICA polymorphism. *Gastroenterology*. 2001; 120:1475-1482.
102. Wiencke K, Spurkland A, Schrupf E, Boberg KM. Primary sclerosing cholangitis is associated to an extended B8-DR3 haplotype including particular MICA and MICB alleles. *Hepatology*. 2001; 34:625-630.
- 820 103. Karlsen TH, Boberg KM, Olsson M, et al. Particular genetic variants of ligands for natural killer cell receptors may contribute to the HLA associated risk of primary sclerosing cholangitis. *J. Hepatol*. 2007; 46:899-906.
- 825 104. Melum E, Karlsen TH, Schrupf E, et al. Cholangiocarcinoma in primary sclerosing cholangitis is associated with NKG2D polymorphisms. *Hepatology*. 2008; 47:90-96.

105. Huang J, Xu L, Luo Y, et al. The inflammation-based scores to predict prognosis of patients with hepatocellular carcinoma after hepatectomy. *Med Oncol.* 2014; 31:883-888.
- 830 106. Cai L, Zhang Z, Zhou L, et al. Functional impairment in circulating and intrahepatic NK cells and relative mechanism in hepatocellular carcinoma patients. *Clinical Immunology.* 2008; 129:428-437.
107. Guillerey C, Huntingdon ND, Smyth MJ. Targeting natural killer cells in cancer immunotherapy. *Nat Immunol.* 2016; 17:1025-1036.
- 835 108. Kamimura H, Yamagiwa S, Tsuchiya A, et al. Reduced NKG2D ligand expression in hepatocellular carcinoma correlates with early recurrence. *J Hepatol.* 2012; 56:381-388.
109. Fang L, Gong J, Wang Y, et al. MICA/B expression is inhibited by unfolded protein response and associated with poor prognosis in human hepatocellular carcinoma. *J Exp Clin Cancer Res.* 2014; 33:76.
- 840 110. Kumar V, Kato N, Urabe Y, et al. Genome-wide association study identifies a susceptibility locus for HCV-induced hepatocellular carcinoma. *Nat Genet.* 2011; 43:455-458.
- 845 111. Kumar V, Yi Lo PH, Sawai H, et al. Soluble MICA and a MICA Variation as Possible Prognostic Biomarkers for HBV-Induced Hepatocellular Carcinoma. Villa E, ed. *PLoS ONE.* 2012; 7:e44743.
112. Tong HV, Toan NL, Song LH, Bock CT, Kremsner PG, Velavan TP. Hepatitis B virus-induced hepatocellular carcinoma: functional roles of MICA variants. *J Viral Hepat.* 2013; 20:687-698.
- 850 113. Pan N, Jiang W, Sun H, et al. KIR and HLA Loci Are Associated with Hepatocellular Carcinoma Development in Patients with Hepatitis B Virus Infection: A Case-Control Study. Sandberg JK, ed. *PLoS ONE.* 2011; 6:e25682-e25688.
114. Sheppard S, Guedes J, Mroz A, et al. The immunoreceptor NKG2D promotes tumour growth in a model of hepatocellular carcinoma. *Nat Commun.* 2017; 8:13930.
- 855 115. Kamiya T, Chang YH, Campana D. Expanded and activated natural killer cells for immunotherapy of hepatocellular carcinoma. *Cancer Immunol Res.* 2016; 4:574-581.
116. Wongkajornsilp A, Somchitprasert T, Butraporn R, et al. Human Cytokine-Induced Killer Cells Specifically Infiltrated and Retarded the Growth of the Inoculated Human Cholangiocarcinoma Cells in SCID Mice. *Cancer Investigation.* 2009; 27:140-148.
- 860 117. Ahn E-Y, Pan G, Vickers SM, McDonald JM. IFN- γ upregulates apoptosis-related molecules and enhances Fas-mediated apoptosis in human cholangiocarcinoma. *Int J Cancer.* 2002; 100:445-451.
118. Takeda K, Smyth MJ, Cretney E, et al. Involvement of Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand in NK Cell-Mediated and IFN- γ -Dependent Suppression of Subcutaneous Tumor Growth. *Cellular Immunology.* 2001; 214:194-200.
- 865 119. Smyth MJ, Crowe NY, Godfrey DI. NK cells and NKT cells collaborate in host protection from methylcholanthrene-induced fibrosarcoma. *International Immunology.* 2001; 13:459-463.
120. Pugh SA, Harrison RJ, Primrose JN, Khakoo SI. T cells but not NK cells are associated with a favourable outcome for resected colorectal liver metastases. *BMC Cancer.* 2014; 14:180.
- 870 121. Galon J, Costes A, Sanchez-Cabo F, Kirilovsky A. Type, density, and location of immune cells within human colorectal tumors predict clinical outcome. *Science.* 2006; 313:1960-1964.
- 875 122. Halama N, Michel S, Kloor M, et al. Localization and Density of Immune Cells in the Invasive Margin of Human Colorectal Cancer Liver Metastases Are Prognostic for Response to Chemotherapy. *Cancer Res.* 2011; 71: 5670-5677.

123. Floros T, Tarhini AA. Anticancer Cytokines - Biology and Clinical Effects of Interferon- α 2, Interleukin (IL)-2, IL-15, IL-21, and IL-12. *Semin Oncol.* 2015; 42:539-548.
- 880 124. Parihar R, Nadella P, Lewis A, Jensen R. A phase I study of interleukin 12 with trastuzumab in patients with human epidermal growth factor receptor-2-overexpressing malignancies: analysis of sustained interferon gamma production in a subset of patients. *Clin Cancer Res.* 2004; 10:5027-5037
125. Lagrue K, Carisey A, Morgan DJ, Chopra R, Davis DM. Lenalidomide augments actin remodeling and lowers NK-cell activation thresholds. *Blood.* 2015; 126:50-60.
- 885 126. Fowler NH, Davis RE, Rawal S, et al. Safety and activity of lenalidomide and rituximab in untreated indolent lymphoma: an open-label, phase 2 trial. *Lancet Oncol.* 2014; 15:1311-1318
127. Lappas CM, Day YJ, Marshall MA, Engelhard VH, Linden J. Adenosine A2A receptor activation reduces hepatic ischemia reperfusion injury by inhibiting CD1d-dependent NKT cell activation. *J Exp Med.* 2006; 203:2639-2648.
- 890 128. Beldi G, Banz Y, Kroemer A, et al. Deletion of CD39 on natural killer cells attenuates hepatic ischemia/reperfusion injury in mice. *Hepatology.* 2010; 51:1702-1711.
129. Kroemer A, Xiao X, Degauque N, et al. The innate NK cells, allograft rejection, and a key role for IL-15. *J Immunol.* 2008; 180:7818-7826.
- 895 130. Obara H, Nagasaki K, Hsieh CL, et al. IFN-gamma, produced by NK cells that infiltrate liver allografts early after transplantation, links the innate and adaptive immune responses. *Am J Transplant.* 2005; 5:2094-2103.
131. Moya-Quiles MR, Torío A, Muro M, et al. Impact of HLA-C on acute rejection in liver transplantation. *Transplant Proc.* 2003; 35:1892-1893.
- 900 132. Hanvesakul R, Spencer N, Cook M, et al. Donor HLA-C genotype has a profound impact on the clinical outcome following liver transplantation. *Am J Transplant.* 2008; 8:1931-1941.
133. Moroso V, van der Meer A, Tilanus HW, et al. Donor and recipient HLA/KIR genotypes do not predict liver transplantation outcome. *Transpl Int.* 2011; 24:932-942.
- 905 134. Pembroke T, Gallimore A, Godkin A. Tracking the kinetics of intrahepatic immune responses by repeated fine needle aspiration of the liver. *J Immunol Methods.* 2015; 424:313-135.
135. Cerwenka A, Lanier LL. Natural killer cell memory in infection, inflammation and cancer. *Nat Rev Immunol.* 2016; 16:112-123.
- 910 136. Lunemann S, Martrus G, Hölzemer A, et al. Sequence variations in HCV core-derived epitopes alter binding of KIR2DL3 to HLA-C*03:04 and modulate NK cell function. *J Hepatol.* 2016; 65:252-258.
137. Cassidy SA, Cheent KS, Khakoo SI. Effects of Peptide on NK cell-mediated MHC I recognition. *Front Immunol.* 2014; 5:133
- 915 138. Mazza G, Rombouts K, Rennie Hall A, et al. Decellularized human liver as a natural 3D-scaffold for liver bioengineering and transplantation. *Sci Rep.* 2015; 5:13079.
139. Balsamo M, Manzini C, Pietra G, et al. Hypoxia downregulates the expression of activating receptors involved in NK-cell-mediated target cell killing without affecting ADCC. *Eur J Immunol.* 2013; 43:2756-2764
- 920 140. Gardiner CM, Finlay DK. What fuels natural killers? Metabolism and NK cell responses. *Front Immunol.* 2017; doi: 10.3389/fimmu.2017.00367
141. Viel S, Marçais, Gumaraes FS, et al. TGF- β inhibits the activation and functions of NK cells by repressing the mTOR pathway. *Sci Signal.* 2016; 9:ra19
- 925 142. Keating SE, Zaiatz-Bittencourt V, Loftus RM, et al. Metabolic Reprogramming Supports IFN- γ Production by CD56bright NK Cells. *J Immunol.* 2016; 196:2552-2560.

143. Wolter F, Glässner A, Krämer B, et al. Hypoxia impairs anti-viral activity of natural killer (NK) cells but has little effect on anti-fibrotic NK cell functions in hepatitis C virus infection. *J. Hepatol.* 2015; 63:1334-1344.
- 930 144. Velásquez SY, Killian D, Schulte J, Sticht C, Thiel M, Lindner HA. Short Term Hypoxia Synergizes with Interleukin 15 Priming in Driving Glycolytic Gene Transcription and Supports Human Natural Killer Cell Activities. *J Biol Chem.* 2016; 291:12960-12977

935

Tables

Receptor	Ligand	Activating/inhibitory
Killer Immunoglobulin-like Receptors (KIRs)	MHC class I (primarily HLA-C)	Either, depending on variant
CD94-NKG2A	MHC class I (HLA-E)	Inhibitory
CD94-NKG2C	MHC class I (HLA-E)	Activating
NKG2D	ULBPs, MICA, MICB	Activating
NCR (NKp30, NKp44, NKp46)	Not well defined	Activating
CD16	IgG-coated cells	Activating

Table I: Selected NK cell receptors and their ligands

940

	Circulating NK cells	Liver-resident NK cells
Adhesion, recruitment and residence		
CD56	Mostly low ("dim")	Mostly high ("bright")
CD57	Mostly positive	Negative
CD69	Negative	Positive
CCR5	Negative	Positive
CCR7	Positive	Negative
CXCR6	Negative	Positive
L-selectin	Positive	Negative
NK cell receptors		
CD16	Mostly positive	Negative
KIR	Mostly positive	Mostly negative
NKG2A/CD94	~50% positive	Mostly positive
NKG2D	Positive	Positive
Transcription factors		
Eomes	Low	High
T-bet	High	Low

Table II: Key phenotypic differences between circulating and liver-resident NK cells

945

Figure Legends

Figure 1: NK cell interference with the antiviral T cell response in CHB

950 During CHB infection, NK cells target HBV-specific T cells, dampening the antiviral T cell response. HBV-specific CD8⁺ T cells upregulate TRAILR2, becoming susceptible to NK cell killing via the TRAIL pathway. Meanwhile HBV-specific CD4⁺ T cells upregulate the NKG2D ligand MICA, which activates NK cells leading to degranulation and cytotoxicity.

955

Figure 2: Roles of NK cells in liver diseases

Figure 1




Figure 2

