

Article type: review

Perspective:

APOBEC mutagenesis in drug resistance and immune escape in HIV and cancer evolution

S. Venkatesan^{1,2*}, R. Rosenthal^{1*}, N. Kanu¹, N. McGranahan¹, J. Bartek^{3,4}, S. A. Quezada^{1,5}, J. Hare⁶, R. S. Harris^{7,8†}, C. Swanton^{1,2†}

*equal contribution

Affiliations:

¹CRUK Lung Cancer Centre of Excellence, UCL Cancer Institute, Paul O’Gorman Building, Huntley St., WC1E 6BT, London, UK

²Translational Cancer Therapeutics Laboratory, The Francis Crick Institute, 1 Midland Rd, NW1 1AT, London, UK

³Danish Cancer Society Research Center, Copenhagen, Denmark

⁴Science for Life Laboratory, Division of Genome Biology, Department of Biochemistry and Biophysics, Karolinska Institute, Stockholm, Sweden

⁵Cancer Immunology Unit, UCL Cancer Institute, Paul O’Gorman Building, Huntley St., WC1E 6BT, London, UK

⁶International AIDS Vaccine Initiative (IAVI), 125 Broad Street, New York, USA

⁷Masonic Cancer Center, Institute for Molecular Virology, Center for Genome Engineering, Department of Biochemistry, Molecular Biology and Biophysics, University of Minnesota, Minneapolis, USA

⁸Howard Hughes Medical Institute, University of Minnesota, Minneapolis, USA

†Correspondence to:

Prof. Reuben S. Harris, Howard Hughes Medical Institute, University of Minnesota, Minneapolis, Minnesota, USA, 55455 phone: +1 612 624 0457; fax: +1 612 625 2163; email: rsh@umn.edu ;

Prof. Charles Swanton, Translational Cancer Therapeutics Laboratory, The Francis Crick Institute, 1 Midland Road, London NW1 1AT, UK. Tel: +44-0-2072693463; Fax: +44-0-2072693463; E-mail: charles.swanton@crick.ac.uk

© The Author 2018. Published by Oxford University Press on behalf of the European Society for Medical Oncology. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted reuse, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract

The APOBEC mutational signature has only recently been detected in a multitude of cancers through next-generation sequencing. In contrast, APOBEC has been a focus of virology research for over a decade. Many lessons learnt regarding APOBEC within virology are likely to be applicable to cancer. In this review, we explore the parallels between the role of APOBEC enzymes in HIV and cancer evolution. We discuss data supporting the role of APOBEC mutagenesis in creating HIV genome heterogeneity, drug resistance, and immune escape variants. We hypothesize similar functions of APOBEC will also hold true in cancer.

Key words

APOBEC, immune escape, drug resistance, human immunodeficiency virus, intratumour heterogeneity

Key message

APOBEC mutagenesis creates diversity within viral and tumour cell populations. Sublethal APOBEC mutagenesis underlies the formation of HIV drug resistant and immune escape variants, and similar functions for APOBEC in cancer are expected. Literature suggests APOBEC inhibition is a double-edged sword, because incomplete inhibition may promote escape whereas, complete APOBEC inhibition may be attractive as adjuvant therapy.

Introduction

Apolipoprotein B mRNA-editing enzyme, catalytic polypeptide-like 3 (APOBEC3; A3) is the name of a seven-membered family of single-stranded DNA cytosine deaminases in humans. Independent approaches including analyses of next-generation sequencing data implicated APOBEC-catalyzed DNA damage and mutagenesis in breast cancer^{1,2}. Subsequent studies confirmed and extended the involvement of APOBEC in mutating the cancer genome to at least sixteen other cancer types³⁻⁵. APOBEC signature mutations (C-to-T and C-to-G in TCA and TCT trinucleotide motifs) are the most prevalent in cancer after those attributable to ageing (C-to-T in CG dinucleotide motifs, most likely due to water-mediated deamination of methyl-cytosine)⁴. Furthermore, the clinical relevance of APOBEC in cancer is underscored by associations with poor patient outcomes and treatment resistance^{6,7}, activation of oncogenic drivers⁸⁻¹⁰, tumour subclonal diversification^{9,11,12}, and increased prevalence in metastases in comparison to primary tumours¹³.

Although involvement of APOBEC mutagenesis in cancer has only recently come to light, these enzymes have been a focus of virology research for over a decade, beginning with the near simultaneous discoveries of APOBEC3G (A3G) as an HIV-1 restriction factor and as a DNA cytosine deaminase^{14,15} (reviewed elsewhere^{16,17}). We envision that many lessons learnt regarding APOBEC within virology will be applicable to oncology. For this reason, we explore the parallels between the role of APOBEC in HIV and cancer mutagenesis. We will especially focus on how APOBEC mutagenesis can promote intratumour heterogeneity, drug resistance, and immune escape.

The AID/APOBEC superfamily: a diverse set of cytosine deaminase enzymes implicated in cancer

APOBEC3 belongs to the AID/APOBEC superfamily, consisting of: activation induced deaminase (AID), APOBEC1 (A1), APOBEC2 (A2), APOBEC3A-H (A3A, A3B, A3C, A3D, A3F, A3G, and A3H) and APOBEC4 (A4). AID deaminates cytosines at the immunoglobulin locus, enabling antibody gene diversification via somatic hypermutation and class switch recombination¹⁸. A1 was identified originally as a RNA editing enzyme, deaminating apolipoprotein B mRNA at a specific position to create an early stop codon¹⁹, but it also has

robust DNA deamination activity^{14,20}. The functions of A2 and A4 are still unclear and these proteins have yet to show enzymatic activity.

In general, the A3 family members are considered part of the innate immune system, forming overlapping barriers to virus and transposon replication. Consistent with such a physiological function, A3 genes show profound copy number and amino acid variation in mammals. For instance, most humans have 7 A3 genes arranged in tandem, whereas rodents have only one at the same genomic location^{21,22}, and each A3 gene in humans as well as several other mammals manifests high levels of amino acid variation due to positive selection²³.

A3G has been studied intensely in the field of virology, as it was recognized early on to deaminate cytosines in cDNA reverse transcription intermediates of retroviruses including HIV-1^{24,25}. Reverse transcriptase places an adenine opposite to the newly created uracil nucleobase, introducing a viral genomic strand G→A mutation²⁶. This inhibits HIV replication by directly rendering the viral genome dysfunctional or by indirectly triggering viral cDNA degradation by subsequent uracil DNA glycosylase activity and endonuclease digestion^{26,27}. A3G can also directly bind to HIV-1 genomic RNA and interfere with viral cDNA synthesis²⁸. A3D, A3F, and A3H also contribute to HIV-1 mutagenesis through similar mechanisms, and it is generally accepted that different subsets of A3 family members restrict the replication of different classes of viruses and transposons (reviewed elsewhere^{16,17}).

Adding to the complexity of 7 A3 family members in humans, different subsets of A3 enzymes are expressed in different tissue types^{29,30}. Together with high levels of DNA sequence similarity (near perfect identity in many regions), determining which of these enzymes is responsible for mutagenesis of different cancer genomes has been challenging. Thus far, the greatest numbers of publications support A3B and A3A (reviewed more extensively elsewhere³¹⁻³³). However, recent reports indicate A3H is also important and, together with A3B, may account for the entire APOBEC mutation signature observed in breast and lung cancers^{34,35}.

APOBEC mutagenesis increases subclonal diversity

APOBEC mutagenesis occurs independently within single cancer cells and viruses, often resulting in strand-coordinated hypermutations (sometimes referred to as kataegis¹). Evidence for A3B upregulation has been found in over half of all cancers³ (reviewed elsewhere³¹). Additionally, our groups and others have identified APOBEC activity as contributing to branched evolution and the acquisition of subclonal mutations later in the evolutionary course of lung adenocarcinoma, estrogen receptor (ER)-negative breast cancer, head and neck squamous cell carcinoma, and esophageal adenocarcinomas^{11,35,36}. A recent analysis of the intratumour heterogeneity present in 100 TRACERx patients with untreated surgically resected primary non-small cell lung cancer (NSCLC) revealed a significant correlation between frequencies of APOBEC signature mutation and the overall number of subclonal mutations⁹. Furthermore, in 19 patients, subclonal driver events were detected as occurring in the APOBEC mutational context⁹ (Fig. 1a). Moreover, tumours with the largest number of subclonal mutations had extensive evidence of APOBEC mutational signatures, suggesting that APOBEC activity is a strong mutagenic force late in cancer evolution (Fig. 1b-c). Subclonal mutations generated from APOBEC activity could potentially drive cancer evolution by enabling the acquisition of late driver mutations. Across multiple types of cancer, there is evidence that APOBEC mutagenesis is responsible for creating driver mutations^{5,8,9,12,31}. The most striking examples are 2 helical domain hot spot mutations in

PIK3CA in papillomavirus-positive head and neck squamous cell carcinomas (E542K and E545K)⁸.

APOBEC expression has also been reported to impact responses to cancer therapy. In ER-positive breast cancer, *A3B* mRNA expression levels inversely correlated with clinical benefit to tamoxifen, and *A3B* overexpression correlated with tamoxifen resistance in mouse xenograft models, implicating the enzyme in promoting drug resistance⁶. Similarly, an enrichment of *A3B* mutations was observed in chemotherapy-resistant urothelial carcinomas³⁷.

APOBEC-induced mutagenesis is also prevalent in HIV-1 proviral DNA sequences³⁸, with some reports estimating the percentage of new mutations attributable to APOBEC activity to be as high as 98%³⁹. However, other studies have estimated lower contributions for APOBEC in HIV-1 genetic variation, and definitive work has yet to be done to dissociate the contributions of APOBEC from mistakes made by the non-proofreading viral reverse transcriptase^{38,40}. Nevertheless, APOBEC activity has been linked to early diversification in newly infected individuals as the transmitted founder virus adapts in response to the immune pressure exerted within its new host⁴¹. Thus, APOBEC mutagenesis has the capability to increase subclonal diversity in both cancer and HIV, with the potential for a profound impact on the subsequent evolution of both the tumour and the virus (2).

The fact that APOBEC increases diversity is likely to be important for the clinical course of both diseases, since genetic heterogeneity is a substrate for Darwinian evolution and is likely to confound treatments. Indeed, in some cancers, increased intratumour heterogeneity has been shown to correlate with a shorter progression free survival^{42,43}. Additionally, measures of genetic heterogeneity have been associated with poor progression free survival in cancer⁴⁴ and intermediate thresholds of copy number instability correlate with poorest clinical outcomes³¹, as they presumably allow for diversity without exceeding toxic levels of genomic instability (Fig. 2). Paralleling findings in cancer, among patients with HIV exhibiting rapid disease progression, there are fewer hypermutated viral sequences as well as fewer minimally edited sequences, suggesting that suboptimal APOBEC activity may enhance genetic diversity, allowing an increase in pathogenicity³⁹ (Fig. 2). From a clinical standpoint, this suggests that successful and complete inhibition of APOBEC enzymatic activity may provide a way to limit subclone diversification and potentially enable normal adaptive immune responses to control or clear HIV-1 infection⁴⁵. Conceivably, complete inhibition of APOBEC enzymatic activity may limit tumour evolution and boost the efficacy of a variety of targeted or immune therapies that often fail due to the acquisition of escape variants brought about by resistance mutations and/or subclonal neoantigens.

Additional lessons from HIV: APOBEC mutagenesis fuels drug resistance and immune escape

In order for HIV and cancer to replicate effectively in human hosts, these pathogenic entities must constantly avoid detection and destruction by innate and adaptive immune responses. As a consequence of this predator-prey relationship, lentiviruses and cancer have evolved elegant mechanisms to avoid or actively counteract immune responses. One of these overlapping mechanisms between HIV and cancer appears to entail co-opting APOBEC mutagenesis as a means of immune evasion.

Lentiviruses and host A3 genes have co-evolved for millions of years in non-human primate relatives⁴⁶ and more recently in humans⁴⁷. HIV-1, as well as other lentiviruses, has adapted to suppress A3-mediated innate anti-viral activity through utilizing the viral infectivity factor (Vif) as a defensive measure^{15,25}. Vif counteracts A3 functionality by polyubiquitination and

proteasomal degradation⁴⁸. Additionally, Vif has been shown to affect A3G translation through mRNA binding⁴⁹ and A3G gene transcription through heterodimerization with the transcription co-factor CBF- β ⁵⁰. These discoveries indicate that targeting the Vif-APOBEC3 interaction may be a novel avenue to combat HIV infection. However, over a decade of virology research into the A3-mediated restriction mechanism has demonstrated that interventions in this pathway may be a double-edged sword. Several studies have found that the range of APOBEC mutagenesis within HIV virions can vary largely. Lethal HIV mutagenesis will be selected against *in vivo*, and moderate APOBEC mutagenesis appears to induce sublethal variation that fuels viral heterogeneity and immune escape^{41,51-53} (Fig. 3). Similar to cancer⁵⁴, viral genetic heterogeneity together with an appropriate selective pressure can enable the emergence of resistant populations. Sublethal G→A mutations in the DNA sequence motif preferred by A3G have been observed in drug-resistant HIV variants⁵⁵⁻⁵⁸ and immune escape variants^{38,41,51,53} (Fig. 2). Mutations in the APOBEC motif are enriched in cytotoxic T lymphocyte (CTL) epitopes of HIV, and can result in diminished CD8+ T cell responses against previously antigenic epitopes, suggestive of selection as a consequence of immune escape³⁸ (Fig. 4). Finally, within cancer there is early evidence that APOBEC correlates with the overexpression of the immune checkpoint molecule PD-L1, potentially contributing to the development of immune exhaustion⁵⁹ (Fig. 4). Indeed, early clinical trial data suggest that reversing immune exhaustion with an anti-PD-L1 antibody enhances the immune response against HIV in a subset of participants⁶⁰.

Some of the evolutionary dynamics that have been identified in HIV in response to therapy resistance also appear to apply to cancer. For example, in HIV-1, the V3 loop of gp120 interacts with the host cellular coreceptor CCR5 in order to gain entry to the cell. CCR5 antagonists bind CCR5 and prevent the entry of CCR5-tropic HIV-1⁶¹. A common route of CCR5 antagonist resistance is the emergence of HIV-1 variants using CXCR4 instead of CCR5 as a coreceptor for cellular entry^{62,63}. Independent studies detected minor CCR5 antagonist-resistant variants containing resistance mutations in the APOBEC context at baseline that were rapidly selected through therapeutic selective pressure^{55,56}. Complementary to these *in vivo* studies, *in vitro* experiments point towards APOBEC mutagenesis generating subclonal resistance mutations which are under positive selection during drug exposure⁵⁸. Similarly, in cancer, minor drug resistant variants are rapidly selected upon treatment (reviewed in Schmitt et al.⁶⁴), although the contribution of APOBEC in this process remains to be quantified.

An optimal range of APOBEC mutagenesis may exist for both HIV and tumour evolution (Fig. 2). It has been shown that several Vif mutants are less potent in inhibiting A3G, with Vif-K22H incapable of fully neutralizing A3G and therefore appearing to enable sufficient levels of sublethal A3G mutagenesis for the emergence of antiretroviral resistance⁵⁷. Thus, it appears that suboptimal A3G mutagenesis is less effective in inactivating the viral genome and in contrast may promote drug resistance and immune escape^{38,45,58}. There are also parallels in the context of APOBEC mutagenesis of the cellular genome. Extensive A3B and A3A mutagenesis induces a DNA damage response (DDR) and at excessive levels it is toxic to the cell^{2,65-69}. Therefore, similar to HIV, cancer cells may have to attenuate APOBEC mutagenesis, enhance repair mechanisms, and/or dampen the DDR pathways to help ensure optimal cell survival (Fig. 2). For instance, loss of p53 enables DNA damage tolerance to A3B-mediated mutagenesis⁶⁵ and correlates positively with A3B overexpression in breast cancer². Furthermore, a recent report focusing on Y-family polymerases and PrimPol, a translesion synthesis (TLS) polymerase with re-priming properties, found that PrimPol as well as POLK and POLI may serve to limit the detrimental effects of APOBEC mutagenesis⁷⁰.

In addition to sublethal APOBEC mutagenesis driving tumour heterogeneity, APOBEC has recently also been shown to produce replication stress in cancer^{65,71,72}. This complements our previous observation that replication stress itself can drive A3B expression and activity⁷³, which suggests a positive feedforward loop involving replication stress and APOBEC mutagenesis (Fig. 5). We have previously implicated replication stress in inducing a widespread endogenous DNA damage response as a biological barrier to tumour progression⁷⁴⁻⁷⁶, as well as driving chromosome segregation errors and ensuing chromosomal instability (CIN)^{76,77}. It is therefore conceivable that oncogene-induced and APOBEC-induced replication stress itself is a driver of CIN in cancer. Furthermore, APOBEC may provide a cancer cell the necessary genomic plasticity to evade immune surveillance. Firstly, a large subclonal neoantigen burden has been associated with a poor response to immunotherapy⁷⁸ and conceivably APOBEC-induced subclonal neoantigens may contribute to this process. Although the mechanism of how subclonal neoantigens potentially confound the response to immunotherapy is still unclear, it is conceivable that subclonal neoantigens foster the outgrowth of T cells that are reactive towards subclonal neoantigens that outcompete T cells reactive towards clonal neoantigens. This may result in the outgrowth of T cells that are only reactive to a proportion of the tumour population, diminishing chances of total tumour control. Secondly, ongoing chromosome missegregations (i.e. CIN) may enable the dynamic loss of immunogenic clonal neoantigens⁷⁹. In effect, CIN-mediated loss of clonal neoantigens (provided these mutations are not required for tumour fitness) as well as the acquisition of subclonal neoantigens (due to APOBEC activity) could both aid immune evasion. Interestingly, highly aneuploid (i.e. an abnormal state of chromosomal copy number) tumours have less immune infiltration and are associated with a worse patient survival than less aneuploid tumours^{80,81}. It is yet to be quantified to which extent APOBEC-induced replication stress and potential ensuing CIN contributes to the process of immune evasion.

Therapeutic considerations

By analogy to hypermutation and hypomutation within HIV⁴⁵, there are at least 2 general strategies for targeting APOBEC in cancer. The first strategy is therapy by hypermutation, by enhancing the mutagenic effects of APOBEC to the point where cancer cells suffer catastrophic levels of DNA damage and selectively die. Indeed, recent studies have suggested that DNA damage response inhibitors, such as PARP and ATR inhibitors, may sensitize tumour cells with high levels of APOBEC to an APOBEC-dependent death^{65,71,72}.

The second strategy is therapy by hypomutation, by inhibiting APOBEC-dependent tumour evolution and potentially suppressing adverse outcomes including recurrence, metastasis, and drug resistance. Constraining cancer evolvability may be accomplished with drugs to inhibit APOBEC gene expression^{73,82} or with chemical inhibitors of DNA deaminase activity (for exemplary studies on A3G see⁸³⁻⁸⁵). Proof of principle has been achieved with a genetic knockdown of *A3B* causing an improvement in the durability of tamoxifen treatment of ER+ xenograft tumours in mice⁶.

Besides harnessing APOBEC through a cancer cell intrinsic mechanism, a complementary strategy may be provoking immune responses to cancer cells with high levels of APOBEC-induced neoantigens. It is important to consider whether neoantigens are present in the trunk of the tumour (clonal) or in the branches (subclonal), since clonal neoantigens may improve response to immune checkpoint blockade in contrast to subclonal neoantigens⁷⁸. Interestingly, one APOBEC-mediated mutational signature (signature 2) is primarily found in the branches of multiple different cancer types¹². In contrast, the other APOBEC-mediated mutational signature (signature 13) is primarily found in the trunk of bladder cancer¹². This

could potentially explain paradoxical effects of APOBEC mutagenesis on immune surveillance and patient outcome. For example, in breast cancer, A3B expression is reported to worsen overall patient survival^{6,7}. In contrast, APOBEC mutagenesis within bladder cancer has been correlated with an improved patient outcome^{86,87}, although the influence of APOBEC on survival in bladder cancer is still a matter of debate⁸⁸. Therefore, it is conceivable that patients with tumours containing extensive clonal APOBEC mutagenesis, in contrast to subclonal APOBEC mutagenesis, are more suitable for immune checkpoint blockade.

It is also important to note that several APOBEC family members including *A3G* and *A3H* are highly expressed in immune cells, including tumour infiltrating T cells, and correlated with improved outcomes^{89,90}. Thus, as studies advance, it will be crucial to not only consider APOBEC expression and mutation signature in the tumour itself, but also APOBEC expression in the larger tumour microenvironment. Immunohistochemistry approaches would be ideal to help address these relationships, but specific monoclonal antibodies for each human APOBEC family member have been challenging to develop due to extensive protein similarity.

Conclusions

Despite the numerous studies that have detected APOBEC-associated mutations by sequencing clinical cancer samples, it is still not fully clear why APOBEC mutagenesis is such a widespread and recurring mutational signature. In this review, we have postulated functions of APOBEC in cancer through exploring the past virology research focused on HIV and APOBEC. Within the field of virology, sublethal APOBEC mutagenesis of HIV virions has been linked to increasing HIV diversity and the creation of drug and immune escape variants. Conceivably, similar functions for APOBEC mutagenesis may be operating, and be selected for, during cancer evolution with clear opportunities for the development of novel therapeutic interventions.

Funding

N.M.: This work was partially funded from Cancer Research UK, Rosetrees and the University College London Hospitals Biomedical Research Centre.

J.B.: This work was partially supported by the Danish Cancer Society, Swedish Research Council, NovoNordisk Foundation (ID 16584), Danish National Research Foundation (project CARD) and Danish Council for Independent Research.

J.H.: This work was partially funded by IAVI with the generous support of USAID and other donors; a full list of IAVI donors is available at www.iavi.org. The contents of this manuscript are the responsibility of the authors and do not necessarily reflect the views of USAID or the US Government.

R.S.H.: This work was partially funded by grants from the National Institutes of Health (NIAID R37 AI064046 and NCI R21 CA206309 to RSH). R.S.H. is the Margaret Harvey Schering Land Grant Chair for Cancer Research, a Distinguished McKnight University Professor, and an Investigator of the Howard Hughes Medical Institute.

C.S.: This work was supported by Cancer Research UK (TRACERx), the Rosetrees Trust, NovoNordisk Foundation [ID 16584], EU FP7 [projects PREDICT and RESPONSIFY, ID number 259303], the Prostate Cancer Foundation, the Breast Cancer Research Foundation, the European Research Council (THESEUS), and National Institute for Health Research University College London Hospitals Biomedical Research Centre.

Competing interests statement

S.V., R.R., N.K., N.M., J.B., S.A.Q. and J.H. declare no competing interests.

R.S.H. is a co-founder, consultant, and shareholder of ApoGen Biotechnologies Inc.

C.S. is a member of the advisory board of ApoGen Biotechnologies Inc.

References

- 1 Nik-Zainal, S. *et al.* Mutational processes molding the genomes of 21 breast cancers. *Cell* **149**, 979-993, doi:10.1016/j.cell.2012.04.024 (2012).
- 2 Burns, M. B. *et al.* APOBEC3B is an enzymatic source of mutation in breast cancer. *Nature* **494**, 366-370, doi:10.1038/nature11881 nature11881 [pii] (2013).
- 3 Burns, M. B., Temiz, N. A. & Harris, R. S. Evidence for APOBEC3B mutagenesis in multiple human cancers. *Nat Genet* **45**, 977-983, doi:10.1038/ng.2701 ng.2701 [pii] (2013).
- 4 Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature* **500**, 415-421, doi:10.1038/nature12477 (2013).
- 5 Roberts, S. A. *et al.* An APOBEC cytidine deaminase mutagenesis pattern is widespread in human cancers. *Nat Genet* **45**, 970-976, doi:10.1038/ng.2702 ng.2702 [pii] (2013).
- 6 Law, E. K. *et al.* The DNA cytosine deaminase APOBEC3B promotes tamoxifen resistance in ER-positive breast cancer. *Sci Adv* **2**, e1601737, doi:10.1126/sciadv.1601737 (2016).
- 7 Sieuwerts, A. M. *et al.* Elevated APOBEC3B correlates with poor outcomes for estrogen-receptor-positive breast cancers. *Horm Cancer* **5**, 405-413, doi:10.1007/s12672-014-0196-8 (2014).
- 8 Henderson, S., Chakravarthy, A., Su, X., Boshoff, C. & Fenton, T. R. APOBEC-mediated cytosine deamination links PIK3CA helical domain mutations to human papillomavirus-driven tumor development. *Cell Rep* **7**, 1833-1841, doi:10.1016/j.celrep.2014.05.012 (2014).
- 9 Jamal-Hanjani, M. *et al.* Tracking the Evolution of Non-Small-Cell Lung Cancer. *N Engl J Med* **376**, 2109-2121, doi:10.1056/NEJMoa1616288 (2017).
- 10 Cancer Genome Atlas Research, N. *et al.* Integrated genomic and molecular characterization of cervical cancer. *Nature* **543**, 378-384, doi:10.1038/nature21386 (2017).
- 11 de Bruin, E. C. *et al.* Spatial and temporal diversity in genomic instability processes defines lung cancer evolution. *Science* **346**, 251-256, doi:10.1126/science.1253462 (2014).
- 12 McGranahan, N. *et al.* Clonal status of actionable driver events and the timing of mutational processes in cancer evolution. *Sci Transl Med* **7**, 283ra254, doi:10.1126/scitranslmed.aaa1408 (2015).
- 13 Lefebvre, C. *et al.* Mutational Profile of Metastatic Breast Cancers: A Retrospective Analysis. *PLoS Med* **13**, e1002201, doi:10.1371/journal.pmed.1002201 (2016).
- 14 Harris, R. S., Petersen-Mahrt, S. K. & Neuberger, M. S. RNA editing enzyme APOBEC1 and some of its homologs can act as DNA mutators. *Mol Cell* **10**, 1247-1253 (2002).
- 15 Sheehy, A. M., Gaddis, N. C., Choi, J. D. & Malim, M. H. Isolation of a human gene that inhibits HIV-1 infection and is suppressed by the viral Vif protein. *Nature* **418**, 646-650, doi:10.1038/nature00939 (2002).

- 16 Harris, R. S. & Dudley, J. P. APOBECs and virus restriction. *Virology* **479-480**, 131-145, doi:10.1016/j.virol.2015.03.012 (2015).
- 17 Simon, V., Bloch, N. & Landau, N. R. Intrinsic host restrictions to HIV-1 and mechanisms of viral escape. *Nat Immunol* **16**, 546-553, doi:10.1038/ni.3156 (2015).
- 18 Muramatsu, M. *et al.* Class switch recombination and hypermutation require activation-induced cytidine deaminase (AID), a potential RNA editing enzyme. *Cell* **102**, 553-563, doi:S0092-8674(00)00078-7 [pii] (2000).
- 19 Navaratnam, N. *et al.* The p27 catalytic subunit of the apolipoprotein B mRNA editing enzyme is a cytidine deaminase. *J Biol Chem* **268**, 20709-20712 (1993).
- 20 Severi, F., Chicca, A. & Conticello, S. G. Analysis of reptilian APOBEC1 suggests that RNA editing may not be its ancestral function. *Mol Biol Evol* **28**, 1125-1129, doi:10.1093/molbev/msq338 (2011).
- 21 Conticello, S. G. The AID/APOBEC family of nucleic acid mutators. *Genome Biol* **9**, 229, doi:10.1186/gb-2008-9-6-229 (2008).
- 22 LaRue, R. S. *et al.* The artiodactyl APOBEC3 innate immune repertoire shows evidence for a multi-functional domain organization that existed in the ancestor of placental mammals. *BMC Mol Biol* **9**, 104, doi:10.1186/1471-2199-9-104 (2008).
- 23 Sawyer, S. L., Emerman, M. & Malik, H. S. Ancient adaptive evolution of the primate antiviral DNA-editing enzyme APOBEC3G. *PLoS Biol* **2**, E275, doi:10.1371/journal.pbio.0020275 (2004).
- 24 Mangeat, B. *et al.* Broad antiretroviral defence by human APOBEC3G through lethal editing of nascent reverse transcripts. *Nature* **424**, 99-103, doi:10.1038/nature01709 (2003).
- 25 Zhang, H. *et al.* The cytidine deaminase CEM15 induces hypermutation in newly synthesized HIV-1 DNA. *Nature* **424**, 94-98, doi:10.1038/nature01707 (2003).
- 26 Harris, R. S. *et al.* DNA deamination mediates innate immunity to retroviral infection. *Cell* **113**, 803-809 (2003).
- 27 Weil, A. F. *et al.* Uracil DNA glycosylase initiates degradation of HIV-1 cDNA containing misincorporated dUTP and prevents viral integration. *Proc Natl Acad Sci U S A* **110**, E448-457, doi:10.1073/pnas.1219702110 (2013).
- 28 Holmes, R. K., Koning, F. A., Bishop, K. N. & Malim, M. H. APOBEC3F can inhibit the accumulation of HIV-1 reverse transcription products in the absence of hypermutation. Comparisons with APOBEC3G. *J Biol Chem* **282**, 2587-2595, doi:10.1074/jbc.M607298200 (2007).
- 29 Refsland, E. W. *et al.* Quantitative profiling of the full APOBEC3 mRNA repertoire in lymphocytes and tissues: implications for HIV-1 restriction. *Nucleic Acids Res* **38**, 4274-4284, doi:10.1093/nar/gkq174 (2010).
- 30 Koning, F. A. *et al.* Defining APOBEC3 expression patterns in human tissues and hematopoietic cell subsets. *J Virol* **83**, 9474-9485, doi:10.1128/JVI.01089-09 (2009).
- 31 Swanton, C., McGranahan, N., Starrett, G. J. & Harris, R. S. APOBEC Enzymes: Mutagenic Fuel for Cancer Evolution and Heterogeneity. *Cancer Discov* **5**, 704-712, doi:10.1158/2159-8290.CD-15-0344 (2015).

- 32 Helleday, T., Eshtad, S. & Nik-Zainal, S. Mechanisms underlying mutational signatures in human cancers. *Nat Rev Genet* **15**, 585-598, doi:10.1038/nrg3729 (2014).
- 33 Roberts, S. A. & Gordenin, D. A. Hypermutation in human cancer genomes: footprints and mechanisms. *Nat Rev Cancer* **14**, 786-800, doi:10.1038/nrc3816 (2014).
- 34 Zhu, M. *et al.* The eQTL-missense polymorphisms of APOBEC3H are associated with lung cancer risk in a Han Chinese population. *Sci Rep* **5**, 14969, doi:10.1038/srep14969 srep14969 [pii] (2015).
- 35 Starrett, G. J. *et al.* The DNA cytosine deaminase APOBEC3H haplotype I likely contributes to breast and lung cancer mutagenesis. *Nat Commun* **7**, 12918, doi:10.1038/ncomms12918 (2016).
- 36 Rosenthal, R., McGranahan, N., Herrero, J., Taylor, B. S. & Swanton, C. DeconstructSigs: delineating mutational processes in single tumors distinguishes DNA repair deficiencies and patterns of carcinoma evolution. *Genome Biol* **17**, 31, doi:10.1186/s13059-016-0893-4 (2016).
- 37 Faltas, B. M. *et al.* Clonal evolution of chemotherapy-resistant urothelial carcinoma. *Nat Genet* **48**, 1490-1499, doi:10.1038/ng.3692 (2016).
- 38 Kim, E. Y. *et al.* Human APOBEC3 induced mutation of human immunodeficiency virus type-1 contributes to adaptation and evolution in natural infection. *PLoS Pathog* **10**, e1004281, doi:10.1371/journal.ppat.1004281 (2014).
- 39 Cuevas, J. M., Geller, R., Garijo, R., Lopez-Aldeguer, J. & Sanjuan, R. Extremely High Mutation Rate of HIV-1 In Vivo. *PLoS Biol* **13**, e1002251, doi:10.1371/journal.pbio.1002251 (2015).
- 40 Delviks-Frankenberry, K. A. *et al.* Minimal Contribution of APOBEC3-Induced G-to-A Hypermutation to HIV-1 Recombination and Genetic Variation. *PLoS Pathog* **12**, e1005646, doi:10.1371/journal.ppat.1005646 (2016).
- 41 Wood, N. *et al.* HIV evolution in early infection: selection pressures, patterns of insertion and deletion, and the impact of APOBEC. *PLoS Pathog* **5**, e1000414, doi:10.1371/journal.ppat.1000414 (2009).
- 42 Birkbak, N. J. *et al.* Paradoxical relationship between chromosomal instability and survival outcome in cancer. *Cancer Res* **71**, 3447-3452, doi:10.1158/0008-5472.CAN-10-3667 0008-5472.CAN-10-3667 [pii] (2011).
- 43 Andor, N. *et al.* Pan-cancer analysis of the extent and consequences of intratumor heterogeneity. *Nat Med*, doi:10.1038/nm.3984 nm.3984 [pii] (2015).
- 44 Schwarz, R. F. *et al.* Spatial and temporal heterogeneity in high-grade serous ovarian cancer: a phylogenetic analysis. *PLoS Med* **12**, e1001789, doi:10.1371/journal.pmed.1001789 (2015).
- 45 Harris, R. S. Enhancing immunity to HIV through APOBEC. *Nat Biotechnol* **26**, 1089-1090, doi:10.1038/nbt1008-1089 (2008).
- 46 Compton, A. A. & Emerman, M. Convergence and divergence in the evolution of the APOBEC3G-Vif interaction reveal ancient origins of simian immunodeficiency viruses. *PLoS Pathog* **9**, e1003135, doi:10.1371/journal.ppat.1003135 (2013).

- 47 Nakano, Y. *et al.* HIV-1 competition experiments in humanized mice show that APOBEC3H imposes selective pressure and promotes virus adaptation. *PLoS Pathog* **13**, e1006348, doi:10.1371/journal.ppat.1006348 (2017).
- 48 Yu, X. *et al.* Induction of APOBEC3G ubiquitination and degradation by an HIV-1 Vif-Cul5-SCF complex. *Science* **302**, 1056-1060, doi:10.1126/science.1089591 1089591 [pii] (2003).
- 49 Mercenne, G. *et al.* HIV-1 Vif binds to APOBEC3G mRNA and inhibits its translation. *Nucleic Acids Res* **38**, 633-646, doi:10.1093/nar/gkp1009 (2010).
- 50 Anderson, B. D. & Harris, R. S. Transcriptional regulation of APOBEC3 antiviral immunity through the CBF-beta/RUNX axis. *Sci Adv* **1**, e1500296, doi:10.1126/sciadv.1500296 (2015).
- 51 Squires, K., Monajemi, M., Woodworth, C. F., Grant, M. D. & Larijani, M. Impact of APOBEC mutations on CD8+T cell recognition of HIV epitopes varies depending on the restricting HLA. *J Acquir Immune Defic Syndr*, doi:10.1097/QAI.0000000000000689 (2015).
- 52 Sadler, H. A., Stenglein, M. D., Harris, R. S. & Mansky, L. M. APOBEC3G contributes to HIV-1 variation through sublethal mutagenesis. *J Virol* **84**, 7396-7404, doi:10.1128/JVI.00056-10 (2010).
- 53 Monajemi, M. *et al.* Positioning of APOBEC3G/F mutational hotspots in the human immunodeficiency virus genome favors reduced recognition by CD8+ T cells. *PLoS One* **9**, e93428, doi:10.1371/journal.pone.0093428 (2014).
- 54 Greaves, M. & Maley, C. C. Clonal evolution in cancer. *Nature* **481**, 306-313, doi:10.1038/nature10762 (2012).
- 55 Tsibris, A. M. *et al.* Quantitative deep sequencing reveals dynamic HIV-1 escape and large population shifts during CCR5 antagonist therapy in vivo. *PLoS One* **4**, e5683, doi:10.1371/journal.pone.0005683 (2009).
- 56 Todesco, E. *et al.* Presence of Minority Resistant Variants After Failure of a Tenofovir, Emtricitabine, and Rilpivirine Regimen. *J Acquir Immune Defic Syndr* **72**, e43-45, doi:10.1097/QAI.0000000000000935 (2016).
- 57 Fourati, S. *et al.* Partially active HIV-1 Vif alleles facilitate viral escape from specific antiretrovirals. *AIDS* **24**, 2313-2321, doi:10.1097/QAD.0b013e32833e515a (2010).
- 58 Kim, E. Y. *et al.* Human APOBEC3G-mediated editing can promote HIV-1 sequence diversification and accelerate adaptation to selective pressure. *J Virol* **84**, 10402-10405, doi:10.1128/JVI.01223-10 (2010).
- 59 Boichard, A., Tsigelny, I. F. & Kurzrock, R. High expression of PD-1 ligands is associated with kataegis mutational signature and APOBEC3 alterations. *Oncoimmunology* **6**, e1284719, doi:10.1080/2162402X.2017.1284719 (2017).
- 60 Gay, C. L. *et al.* Clinical Trial of the Anti-PD-L1 Antibody BMS-936559 in HIV-1 Infected Participants on Suppressive Antiretroviral Therapy. *J Infect Dis*, doi:10.1093/infdis/jix191 (2017).
- 61 Strizki, J. M. *et al.* Discovery and characterization of vicriviroc (SCH 417690), a CCR5 antagonist with potent activity against human

- immunodeficiency virus type 1. *Antimicrob Agents Chemother* **49**, 4911-4919, doi:10.1128/AAC.49.12.4911-4919.2005 (2005).
- 62 Fatkenheuer, G. *et al.* Subgroup analyses of maraviroc in previously treated R5 HIV-1 infection. *N Engl J Med* **359**, 1442-1455, doi:10.1056/NEJMoa0803154 (2008).
- 63 Westby, M. *et al.* Emergence of CXCR4-using human immunodeficiency virus type 1 (HIV-1) variants in a minority of HIV-1-infected patients following treatment with the CCR5 antagonist maraviroc is from a pretreatment CXCR4-using virus reservoir. *J Virol* **80**, 4909-4920, doi:10.1128/JVI.80.10.4909-4920.2006 (2006).
- 64 Schmitt, M. W., Loeb, L. A. & Salk, J. J. The influence of subclonal resistance mutations on targeted cancer therapy. *Nat Rev Clin Oncol* **13**, 335-347, doi:10.1038/nrclinonc.2015.175 (2016).
- 65 Nikkila, J. *et al.* Elevated APOBEC3B expression drives a kataegic-like mutation signature and replication stress-related therapeutic vulnerabilities in p53-defective cells. *Br J Cancer* **117**, 113-123, doi:10.1038/bjc.2017.133 (2017).
- 66 Akre, M. K. *et al.* Mutation Processes in 293-Based Clones Overexpressing the DNA Cytosine Deaminase APOBEC3B. *PLoS One* **11**, e0155391, doi:10.1371/journal.pone.0155391 (2016).
- 67 Caval, V., Suspene, R., Shapira, M., Vartanian, J. P. & Wain-Hobson, S. A prevalent cancer susceptibility APOBEC3A hybrid allele bearing APOBEC3B 3'UTR enhances chromosomal DNA damage. *Nat Commun* **5**, 5129, doi:10.1038/ncomms6129 (2014).
- 68 Landry, S., Narvaiza, I., Linfesty, D. C. & Weitzman, M. D. APOBEC3A can activate the DNA damage response and cause cell-cycle arrest. *EMBO Rep* **12**, 444-450, doi:10.1038/embor.2011.46 (2011).
- 69 Green, A. M. *et al.* APOBEC3A damages the cellular genome during DNA replication. *Cell Cycle* **15**, 998-1008, doi:10.1080/15384101.2016.1152426 (2016).
- 70 Pilzecker, B. *et al.* PrimPol prevents APOBEC/AID family mediated DNA mutagenesis. *Nucleic Acids Res* **44**, 4734-4744, doi:10.1093/nar/gkw123 (2016).
- 71 Buisson, R., Lawrence, M. S., Benes, C. H. & Zou, L. APOBEC3A and 3B activities render cancer cells susceptible to ATR inhibition. *Cancer Res*, doi:10.1158/0008-5472.CAN-16-3389 (2017).
- 72 Green, A. M. *et al.* Cytosine deaminase APOBEC3A sensitizes leukemia cells to inhibition of the DNA replication checkpoint. *Cancer Res*, doi:10.1158/0008-5472.CAN-16-3394 (2017).
- 73 Kanu, N. *et al.* DNA replication stress mediates APOBEC3 family mutagenesis in breast cancer. *Genome Biol* **17**, 185, doi:10.1186/s13059-016-1042-9 (2016).
- 74 Bartkova, J. *et al.* DNA damage response as a candidate anti-cancer barrier in early human tumorigenesis. *Nature* **434**, 864-870, doi:10.1038/nature03482 (2005).
- 75 Bartkova, J. *et al.* Oncogene-induced senescence is part of the tumorigenesis barrier imposed by DNA damage checkpoints. *Nature* **444**, 633-637, doi:10.1038/nature05268 (2006).

- 76 Halazonetis, T. D., Gorgoulis, V. G. & Bartek, J. An oncogene-induced DNA damage model for cancer development. *Science* **319**, 1352-1355, doi:10.1126/science.1140735 (2008).
- 77 Burrell, R. A. *et al.* Replication stress links structural and numerical cancer chromosomal instability. *Nature* **494**, 492-496, doi:10.1038/nature11935 (2013).
- 78 McGranahan, N. *et al.* Clonal neoantigens elicit T cell immunoreactivity and sensitivity to immune checkpoint blockade. *Science* **351**, 1463-1469, doi:10.1126/science.aaf1490 (2016).
- 79 Anagnostou, V. *et al.* Evolution of Neoantigen Landscape during Immune Checkpoint Blockade in Non-Small Cell Lung Cancer. *Cancer Discov* **7**, 264-276, doi:10.1158/2159-8290.CD-16-0828 (2017).
- 80 Davoli, T., Uno, H., Wooten, E. C. & Elledge, S. J. Tumor aneuploidy correlates with markers of immune evasion and with reduced response to immunotherapy. *Science* **355**, doi:10.1126/science.aaf8399 (2017).
- 81 Karn, T. *et al.* Association Between Genomic Metrics and Immune Infiltration in Triple-Negative Breast Cancer. *JAMA Oncol*, doi:10.1001/jamaoncol.2017.2140 (2017).
- 82 Leonard, B. *et al.* The PKC/NF-kappaB signaling pathway induces APOBEC3B expression in multiple human cancers. *Cancer Res* **75**, 4538-4547, doi:10.1158/0008-5472.CAN-15-2171-T (2015).
- 83 Li, M. *et al.* First-in-class small molecule inhibitors of the single-strand DNA cytosine deaminase APOBEC3G. *ACS Chem Biol* **7**, 506-517, doi:10.1021/cb200440y (2012).
- 84 Olson, M. E., Li, M., Harris, R. S. & Harki, D. A. Small-molecule APOBEC3G DNA cytosine deaminase inhibitors based on a 4-amino-1,2,4-triazole-3-thiol scaffold. *ChemMedChem* **8**, 112-117, doi:10.1002/cmdc.201200411 (2013).
- 85 Olson, M. E. *et al.* Oxidative Reactivities of 2-Furylquinolines: Ubiquitous Scaffolds in Common High-Throughput Screening Libraries. *J Med Chem* **58**, 7419-7430, doi:10.1021/acs.jmedchem.5b00930 (2015).
- 86 Lerner, S. P. *et al.* Comprehensive molecular characterization and analysis of muscle-invasive urothelial carcinomas. *Journal of Clinical Oncology* **35**, 4500-4500, doi:10.1200/JCO.2017.35.15_suppl.4500 (2017).
- 87 Glaser, A. P., Fantini, D., Rimar, K. J. & Meeks, J. J. APOBEC-Mediated Mutagenesis In Urothelial Carcinoma Is Associated With Improved Survival, Mutations In DNA Damage Response Genes, And Immune Response. *bioRxiv*, 123802 (2017).
- 88 Hedegaard, J. *et al.* Comprehensive Transcriptional Analysis of Early-Stage Urothelial Carcinoma. *Cancer Cell* **30**, 27-42, doi:10.1016/j.ccell.2016.05.004 (2016).
- 89 Mullane, S. A. *et al.* Correlation of Apobec Mrna Expression with overall Survival and pd-l1 Expression in Urothelial Carcinoma. *Sci Rep* **6**, 27702, doi:10.1038/srep27702 (2016).
- 90 Leonard, B. *et al.* APOBEC3G expression correlates with T cell infiltration and improved clinical outcomes in high-grade serous ovarian carcinoma. *Clin Cancer Res*, doi:clincanres.2910.2015 [pii] 1078-0432.CCR-15-2910 [pii] 10.1158/1078-0432.CCR-15-2910 (2016).

Figure legends

Figure 1 APOBEC mutagenesis within the TRACERx 100 cohort.

(A) Phylogenetic trees of TRACERx patients harbouring a subclonal driver mutation in an APOBEC preferred motif are shown. The mutation is indicated near the clone in which it occurs. Clonal clusters are shown in blue, subclonal clusters are shown in red, and subclonal clusters containing the driver mutation in a preferred APOBEC motif are shown in yellow. (B) The total numbers of mutations in each patient are shown, with mutations in an APOBEC context shown in yellow and all other mutational contexts shown in green. (C) The total numbers of driver mutations in each patient are shown, with mutations in an APOBEC context shown in yellow and all other mutational contexts shown in green. (D) The fraction of subclonal mutations for each patient that could be attributed to APOBEC activity are shown, with mutations in an APOBEC context shown in yellow and all other mutational contexts shown in green.

Figure 2 Parallels between APOBEC mutagenesis in HIV and cancer.

(A) Within HIV, APOBEC mutagenesis is counteracted through Vif. As APOBEC mutagenesis increases, the chance of lethal mutagenesis and population variation increases. The trade-off between lethal mutagenesis and population variation creates an optimal range in which APOBEC mutagenesis increases population fitness. (B) Within cancer, the toxic effects of APOBEC mutagenesis is counteracted through DNA damage repair and DNA damage tolerance. As APOBEC mutagenesis increases, the chance of lethal mutagenesis and population variation increases. The trade-off between lethal mutagenesis and population variation creates an optimal range in which APOBEC mutagenesis increases population fitness.

Figure 3 Potential roles of APOBEC supporting the escape and progression of HIV and cancer.

(A) Sublethal APOBEC mutagenesis promotes the formation of drug escape and immune escape variants in HIV that will be selected upon exposure to treatment or the immune system. (B) Sublethal APOBEC mutagenesis promotes the formation of drug escape and immune escape variants in cancer that will be selected upon exposure to treatment or the immune system. APOBEC mutagenesis also underlies the formation of driver gene mutations and potentially also replication stress-induced genomic instability, although the latter is speculative.

Figure 4 APOBEC mutagenesis promotes HIV immune escape and potentially also that of cancer.

APOBEC mutations have been identified in CTL epitopes of HIV. APOBEC hypermutation has been linked to PD-L1 ligand overexpression and potentially contributes to immune exhaustion of tumour infiltrating lymphocytes. Studies within the HIV literature are suggestive of sublethal APOBEC mutagenesis driving the expansion of escape variants.

Figure 5 A feedforward replication stress-APOBEC loop potentially drives subclonal somatic point mutations and copy number variations.

APOBEC induces subclonal somatic point mutations and has recently been shown to induce replication stress. APOBEC-mediated replication stress could potentially contribute to CIN.




Figure 1 APOBEC mutagenesis within the TRACERx 100 cohort.

(A) Phylogenetic trees of TRACERx patients harbouring a subclonal driver mutation in an APOBEC preferred motif are shown. The mutation is indicated near the clone in which it occurs. Clonal clusters are shown in blue, subclonal clusters are shown in red, and subclonal clusters containing the driver mutation in a preferred APOBEC motif are shown in yellow. (B) The total numbers of mutations in each patient are shown, with mutations in an APOBEC context shown in yellow and all other mutational contexts shown in green. (C) The total numbers of driver mutations in each patient are shown, with mutations in an APOBEC context shown in yellow and all other mutational contexts shown in green. (D) The fraction of subclonal mutations for each patient that could be attributed to APOBEC activity are shown, with mutations in an APOBEC context shown in yellow and all other mutational contexts shown in green.

251x318mm (300 x 300 DPI)




Figure 2 Parallels between APOBEC mutagenesis in HIV and cancer. (A) Within HIV, APOBEC mutagenesis is counteracted through Vif. As APOBEC mutagenesis increases, the chance of lethal mutagenesis and population variation increases. The trade-off between lethal mutagenesis and population variation creates an optimal range in which APOBEC mutagenesis increases population fitness. (B) Within cancer, the toxic effects of APOBEC mutagenesis is counteracted through DNA damage repair and DNA damage tolerance. As APOBEC mutagenesis increases, the chance of lethal mutagenesis and population variation increases. The trade-off between lethal mutagenesis and population variation creates an optimal range in which APOBEC mutagenesis increases population fitness.

69x41mm (300 x 300 DPI)




Figure 3 Potential roles of APOBEC supporting the escape and progression of HIV and cancer. (A) Sublethal APOBEC mutagenesis promotes the formation of drug escape and immune escape variants in HIV that will be selected upon exposure to treatment or the immune system. (B) Sublethal APOBEC mutagenesis promotes the formation of drug escape and immune escape variants in cancer that will be selected upon exposure to treatment or the immune system. APOBEC mutagenesis also underlies the formation of driver gene mutations and potentially also replication stress-induced genomic instability, although the latter is speculative.

87x50mm (300 x 300 DPI)




Figure 4 APOBEC mutagenesis promotes HIV immune escape and potentially also that of cancer. APOBEC mutations have been identified in CTL epitopes of HIV. APOBEC hypermutation has been linked to PD-L1 ligand overexpression and potentially contributes to immune exhaustion of tumour infiltrating lymphocytes. Studies within the HIV literature are suggestive of sublethal APOBEC mutagenesis driving the expansion of escape variants.

93x58mm (300 x 300 DPI)




Figure 5 A feedforward replication stress-APOBEC loop potentially drives subclonal somatic points mutations and copy number variations. APOBEC induces subclonal somatic point mutations and has recently been shown to induce replication stress. APOBEC- mediated replication stress could potentially contribute to CIN.

44x13mm (300 x 300 DPI)