

Re-evaluating Microglia Expression Profiles using RiboTag and Cell isolation strategies

Zhana Haimon¹, Alon Volasky¹, Johannes Orthgiess², Sigalit Boura-Halfon¹, Diana Varol¹, Anat Shemer¹, Simon Yona¹, Binyamin Zuckerman³, Eyal David¹, Louise Chappell-Maor¹, Ingo Bechmann⁴, Martin Gericke⁴, Igor Ulitsky³ and Steffen Jung^{1*}

Departments of ¹ Immunology and ³ Biological Regulation, Weizmann Institute of Science, Rehovot 76100, Israel.

² Carl Ludwig Institute of Physiology and ⁴Institute of Anatomy, University of Leipzig, Leipzig 04103, Germany

* Corresponding author e-mail: s.jung@weizmann.ac.il (S.J.)

Abstract

Transcriptome profiling is widely used to infer functional states of specific cell types, as well as their responses to stimuli, to define contributions to physiology and pathophysiology. Focusing on microglia, the brain macrophages, we report here a side-by-side comparison of the classical cell sort-based transcriptome sequencing and the 'RiboTag' method that avoids cell retrieval from tissue context and yields translome sequencing information. Conventional whole cell microglia transcriptomes were found to be significantly tainted by artifacts induced by tissue-dissociation, cargo contaminations and transcripts sequestered from ribosomes. Conversely, our data highlight the added value of RiboTag profiling to assess the accuracy of Cre transgenic mice. Collectively, this study indicates method-based biases, alerts towards observer effects and establishes RiboTag-based translome profiling as a valuable complement to standard sort-based profiling strategies.

128 words, 150 allowed

Introduction

Cellular functions are defined by transcriptomes and proteomes. Global gene expression profiling can hence provide insights into specific contributions of distinct cell types to various physiological processes. Macrophages are myeloid immune cells that are strategically positioned to ingest and degrade dead cells, debris and foreign material, and orchestrate inflammation and immune defense. Moreover, emerging evidence supports additional critical tissue macrophage contributions to the establishment and maintenance of organ functions. Recent studies have highlighted the impact of the tissue environment on macrophage expression signatures and enhancer landscapes^{1,2}. Conversely, tissue macrophages lose distinct expression patterns once taken into culture², likely due to loss of original and exposure to novel environmental cues. Accurate expression profiling of cells in order to infer their *in vivo* functions therefore requires methods that allow efficient and rapid retrieval of phenotypically specified cells or their RNA from intact organs.

Classically, the isolation of defined cell population from their physiological tissue context involves the preparation of single cell suspensions followed by flow cytometry- or magnetic bead-based cell sorting. Depending on the cell type studied and its respective extent of tissue embedding, release of the cells can require mechanical processing and extensive enzymatic digests with prolonged incubations. Collectively, these manipulations bear the inherent risk of artifacts. Moreover, cell isolation protocols are often inefficient and likely prone to introduce bias towards subpopulations. Even optimized isolation protocols fail for instance to retrieve more than 10% of microglia cells from an intact mouse brain, estimated to comprise three million cells.

To circumvent the need for cell retrieval, alternative approaches were introduced that allow isolation of cell-specific transcriptomes by immuno-precipitation (IP) of epitope-tagged ribosomes from crude tissue extracts^{3,4}. In the 'RiboTag approach' developed by McKnight and colleagues⁴, cell type-specific expression of Cre recombinase is used to activate expression of a hemagglutinin (HA) epitope-tagged ribosomal subunit (RPL22) by deletion of a 'floxed' WT exon. IP of the tagged ribosomes from whole tissue extracts with anti-HA antibody-coupled magnetic beads enables the pull-down of cell type-specific ribosome-attached mRNA, *i.e.* the transcriptome.

Here we report the application of the RiboTag approach to the study of microglia. Specifically, we compared previously reported *Cx3cr1^{Cre}* and *Cx3cr1^{CreER}* transgenic animals⁵ for their potency and specificity to be used in expression profiling of microglia using the RiboTag strategy. Side-by-side comparison of translomes isolated by IP from crude tissue extracts and transcriptomes from sorted microglial cells highlights the advantages and disadvantages of the respective approaches. Whole cell transcriptomes were found to be contaminated by artifacts induced by tissue-dissociation, cargo contaminations and transcripts sequestered from ribosomes. Finally, we performed a translome analysis on microglia from animals exposed to acute peripheral endotoxin challenge. Collectively, our results highlight the specifics of RiboTag profiling and establish this method as a valuable complement to standard sort-based profiling strategies.

Results

Definition of cell type specificity of *CX₃CR1^{Cre}* and *CX₃CR1^{CreER}* transgenic mice

The RiboTag strategy is a two-component approach relying on the combination of a floxed *Rpl22^{HA}* allele⁴ with a cell type-specific Cre recombinase transgene. Microglia display unique high expression of CX₃CR1⁶ and transgenic mice harboring a GFP reporter gene under the promoter of this chemokine receptor have been instrumental to study microglia morphology and dynamics, as GFP expression in adult mouse brains is restricted to microglia^{6,7}. More recently, we introduced two mouse strains that display Cre recombinase activity under the control of the *Cx3cr1* promoter, either constitutively (*Cx3cr1^{Cre}* mice), or following Tamoxifen (TAM) - mediated activation of an estrogen receptor-fused latent Cre recombinase (*Cx3cr1^{CreER}* mice)^{5,8} (**Figure 1A**). To implement the RiboTag method for the study of microglia, we generated *Cx3cr1^{Cre}:Rpl22^{HA}* and *Cx3cr1^{CreER}:Rpl22^{HA}* mice. We then performed RNAseq on RNA isolated from whole brain tissue (input), RNA retrieved by a control IP (IP IgG) and by anti-HA IP (IP HA) of brain extracts from the two mouse strains (**Figure 1B**). To assess the cell type specificity of the obtained translomes, we compared them to published neuron and glia-specific gene expression signatures⁹. Translomes retrieved from *Cx3cr1^{Cre}:Rpl22^{HA}* and TAM-treated *Cx3cr1^{CreER}:Rpl22^{HA}* mice showed an enrichment for mRNAs encoding microglial proteins, such as Sall1, Csf1r, Trem2, Aif1 (Iba-1) and CD11b, that represent a small fraction in the total input, confirming

rearrangement of the 'floxed' *Rpl22^{HA}* allele in microglia (**Figure 1C**). Conversely, key astrocyte and oligodendrocyte transcripts, such as *Gfap*, *Aldh1l1*, *Aqp4* and *Mbp*, *Mog*, *Olig1*, *Plp1*, respectively, were as expected depleted in both translomes (**Figure 1C**). Surprisingly, translomes retrieved by ribosome IP from brain homogenates of *Cx3cr1^{Cre}:Rpl22^{HA}* but not TAM-treated *Cx3cr1^{CreER}:Rpl22^{HA}* mice, also exhibited a prominent neuronal signature, including mRNAs encoding Calbindin 2, CX₃CL1 and CELF (**Figure 1C**). This suggested activation of RPL22-HA expression in neuronal cells of *Cx3cr1^{Cre}:Rpl22^{HA}* mice. Analysis of *Cx3cr1^{Cre}:Rosa26^{YFP}* animals that harbor a 'floxed' reporter allele revealed prominent neuronal labeling, comparable to that recently reported for *LysM2^{Cre}* mice¹⁰ (**Figure 1D**). Moreover, immuno-histochemical analysis of brains of *Cx3cr1^{Cre}:Rpl22^{HA}* animals detected neuronal staining by anti-HA antibodies in brain sections. In contrast, RPL22-HA expression in TAM-treated *Cx3cr1^{CreER}:Rpl22^{HA}* mice was restricted to microglia, as detected by co-staining for IBA-1 (**Figure 1E**). Since *Cx3cr1^{gfp}* mice lack GFP labeling of adult neurons^{6,11}, rearrangements in *Cx3cr1^{Cre}* mice are likely due to a transient and further-to-be-defined window of *Cx3cr1* promoter activity during neuronal development. In support of this notion, one of the *Cx3cr1^{Cre}:Rpl22^{HA}* mice analyzed, also displayed astrocyte and oligodendrocyte transcripts, in line with the shared neuro-ectodermal origin of these glia cells and neurons (**Figure 1C**).

Cx3cr1^{CreER}:Rpl22^{HA} mice that were TAM-treated postnatally displayed brain macrophage-restricted activation of the RiboTag. However, some enrichment for the microglia translome was observed without TAM treatment in these mice and rare YFP⁺ cells could be detected in non-treated *Cx3cr1^{CreER}:Rosa26^{YFP}* animals (**Figure 1C, D middle panel**), corroborating reports of leakiness of the CreER system¹². However, as confirmed by flow cytometric analysis (**Figure 1F**), robust rearrangement and microglial expression of the HA epitope-tagged ribosome subunit in *Cx3cr1^{CreER}:Rpl22^{HA}* mice were dependent on the induction by the estrogen derivate, in our facility.

Collectively, these data illustrate the value of the RiboTag profiling approach to assess the accuracy of Cre transgenic mouse models, and to investigate specific cell types, including fate mapping and conditional mutagenesis.

Comparison of RiboTag profiling to cell sort-based transcriptomics

Having established the value of *Cx3cr1^{CreER}:Rpl22^{HA}* mice, we next compared translomes and transcriptomes of sorted and unsorted microglia. Specifically, individual brains of TAM-treated *Cx3cr1^{CreER}:Rpl22^{HA}* mice were divided: one hemisphere underwent direct tissue homogenization followed by IP with anti-HA (IP-HA) or an isotype control antibody to define the method-related background (IP-IgG). The second hemisphere was subjected to the classical microglia isolation protocol involving tissue digestion followed by cell sorting of microglial cells (defined as DAPI⁻ Ly6C/G (Gr1)⁻ CD11b⁺ CD45^{int}) (**Figure 2A**). A fraction of the sorted microglia was taken for direct mRNA isolation to yield the whole transcriptome (Sort); another fraction was lysed and subjected to the anti-HA IP to retrieve the translome of sorted cells (Sort-IP) (**Figure 2A**). This experimental set-up allowed comparison of translomes of sorted and unsorted microglia alongside whole transcriptomes of sorted microglia from the same brain, and thus to investigate the impact of the isolation protocol on gene expression.

Unbiased K-means clustering of the significantly differentially expressed genes between at least two sample groups (IP-HA vs IP-IgG, IP-HA vs Sort, IP-HA vs Sort-IP and Sort vs Sort-IP, fold change > 2, p < 0.05), revealed four clusters (**Figure 2B**). Cluster IV was discerned as RiboTag method-related background, since mRNA reads in the non-specific IP-IgG were higher than in the specific IP-HA and absent from sorted samples. IP specific genes were selected for being significantly higher in IP-HA than IP-IgG (fold change > 2, p < 0.05), and all genes below this threshold were removed from the analysis.

Cluster I comprised 913 mRNAs enriched in the specific IP-HA compared with IP-IgG and present in both samples of sorted microglia. This cluster includes established microglia signature genes, such as *Aif1*, *Irf8*, *Itgam*, *Sall1*, *Cx3cr1*, *Tgfb* and *Hexb* (**Supplementary Figure 2**), indicating that the retrieval methods we used are comparable. Cluster II was represented by 525 mRNAs that were highly abundant in sorted samples (both in translome and transcriptome), but not present in the direct IP-HA. Cluster III comprised 282 mRNAs prominently enriched in the direct IP-HA, but less abundant in the sorted microglia. Cluster II and III highlight differences between the retrieval methods, as well as discrepancies between transcriptomes and translomes and will be the focus of the remainder of this study.

Transcripts overrepresented in microglia translomes

Cluster III (**Figure 2B**) is defined by genes highly expressed in 'IP-HA' samples relative to 'Sort' samples and could be further subdivided according to transcript abundance in the 'Sort-IP' samples (**Figure 2C**): Cluster III-a (72 mRNAs) were low in both samples of the sorted cells ('Sort' and 'Sort-IP'); Clusters III-b and III-c (210 mRNAs) were low in the 'Sort' samples but highly expressed in the 'Sort-IP' samples (**Figure 2C**).

Cluster III-a can be largely explained by the presence of non-parenchymal macrophage mRNAs. Non-parenchymal brain macrophages, including perivascular, meningeal and choroid plexus macrophages, can be discriminated from CD11b⁺ CD45^{int} microglia as CD11b⁺ CD45^{hi} cells and therefore be excluded by fluorescence-activated cell sorting (**Figure 2D**). CX₃CR1 is expressed both in microglia and in non-parenchymal brain macrophages (**Figure 2D**)¹³; accordingly, both CD45^{int} microglia and CD45^{hi} macrophages of TAM-treated *Cx3cr1^{CreER}:Rpl22^{HA}* mice express the HA-tagged Rpl22 isoform (**Figure 2E**). Moreover, some of the non-parenchymal brain macrophages are like microglia long-lived and hence do not lose the rearranged alleles as do monocytes (Goldmann et al., 2016). Single cell transcriptomics have shown that non-parenchymal brain macrophages differ in gene expression from microglia^{13,14}. Accordingly, cluster III-a included *Cd163*, *F13a1*, *Cbr2*, *Mrc1* and *Lyve1* (**Figure 2F, 2G**).

The combined clusters III-b and III-c comprise mRNAs that are enriched in both IP-HA and Sort-IP transcriptomes over the whole transcriptomes, suggesting their functional importance for the cells at the time of isolation. These include mRNAs encoding proteins related to metabolism (*Gpx1*, *Sdhc*), vesicular transport (*Clta*, *Kdelr1*, *Ykt6*), sphingolipid metabolism (*Gm2a*, *Psap*) and lipids (*ApoE*), as well as components of the GABA-receptor signaling cascade (*Gabarap*, *Gnai2*) (**Figure 2H**). Specific functions of these genes in microglia remain to be explored. Collectively, these results highlight the value of multifaceted approach of the RiboTag and cell sorting strategy to improve cell type specificity. In addition, the RiboTag approach allows to focus specifically on genes that are actively being translated and contributing to the cellular proteome at a particular time and location.

Transcripts overrepresented in transcriptomes I

- Isolation artifacts

Cluster II (**Figure 2B**) defines 525 mRNAs that were high in Sort, but low in IP-HA. This cluster can be further subdivided according to transcript abundance in the Sort-IP samples (**Figure 3A**).

Cluster II-a comprised 190 transcripts similarly expressed in Sort-IP and Sort samples and was found to include mRNAs that are related to immune activation, such as *Cd86*, *Cd53*, *Tlr4*, and *Tlr7* (**Figure 3B**). Ingenuity Pathway Analysis (IPA) of significantly upregulated genes in Sort vs. IP-HA (>2 fold change, $p < 0.05$) showed upregulation of pathways, such as 'Production of NOS and ROS', 'Phagocytosis' and 'TLR signaling' (**Figure 3C**). Since these transcripts are high in both translomes and transcriptomes of sorted cells, we assume that they reflect cell activation resulting from the isolation process (see also reference #15¹⁵).

Commonly used macrophage isolation protocols, such as the one we applied for the microglia retrieval, include enzymatic tissue digestion at 37°C, *i.e.* a step that could cause cell activation and transcriptome alterations. Moreover, enzymes employed in these digests might contain endotoxin contaminations that could activate cells. To probe for the potential impact of these manipulations, we compared transcriptomes of sorted cells that were isolated from the same brain with or without collagenase and DNase digestion, and the RiboTag approach. Surprisingly though, both isolation procedures resulted in comparable transcriptional profiles, indicated by differential expression of 472 and 267 genes (cluster II and III, respectively), as compared to the relevant IP samples (**Supplementary Figure 3A**). Global correlation of gene expression of the samples retrieved with or without incubation was high ($r^2 = 0.99$), as compared to the correlation of sorted and IP samples ($r^2 = 0.1$) (**Supplementary Figure 3B**). Similarities were also apparent in a correlation matrix compiled from data of independent experiments (**Supplementary Figure 3C**). Collectively, these data establish that the artifact is reproducible and suggest that it is introduced by extraction of the cells from their native environment, rather than subsequent manipulation.

Transcripts overrepresented in the transcriptomes II

- Cargo contaminants and sequestered RNAs

Cluster II-b spanned 335 genes that were low in the Sort-IP compared with Sort samples (**Figure 3A**), suggesting translome-transcriptome differences.

Microglia are specialized phagocytes that like other macrophages, ingest dead cells and cell debris¹⁶. Whole cell transcriptomes hence tend to include genetic material from ingested neighboring cells. Indeed, almost half of the mRNAs in cluster II-b (157 out of 335 genes) are likely to be derived from such external source (**Figure 4A**). Examples include *Arhgap5*, *Son* and *Pisd-ps1*, which are reportedly transcribed in astrocytes and neurons⁹ (**Figure 4B**).

Long non-coding mRNAs (lncRNAs) were shown to be enriched in nuclei¹⁷, where some of them act in transcriptional regulation. As expected, representatives of these lncRNAs, such as *Malat1* and *Neat1*, were identified in the whole cell transcriptomes, while absent from translomes, and appeared in Cluster II-b (**Figure 4C**).

Gene expression is controlled at the levels of transcription and translation. The latter comprises specific mechanisms that prevent mRNAs from their integration into ribosomes, including nuclear retention and sequestration into dedicated membrane-less cytosolic ribo-nucleoprotein complexes^{18,19}. The content of these organelles, such as processing bodies (P-bodies) and stress granules is only beginning to be defined^{19,20}. However, sequestered mRNAs have been reported to be longer and to comprise extended 3'UTRs, as well as to display lower splicing efficiencies^{19,21}. When analyzed for these three parameters, mRNAs defined by Cluster II-b, showed significant presence of these hallmarks, as compared to all other clusters (**Figure 4D**). Moreover, transcripts of cluster II-b also showed significant overlap with the list of nuclear retained mRNAs reported for other cellular systems^{22,23} (**Figure 4E**). Among the protein-coding mRNAs that seem sequestered from immediate translation, we found *Fos*, *Jun*, *Egr1* and *Zfp3611* (**Figure 4F**), which are immediate early genes that have been described to be induced within minutes after activation. Of note, these mRNAs appear also in the translome of the sorted cells, suggesting that they move to the ribosomes during the isolation procedure. Collectively, discrepancies we observed between microglia translomes and whole cell transcriptomes can be explained by the inclusion of cargo-derived transcripts and mRNAs sequestered to nuclei or P-bodies (**Figure 4G**). These data highlight the value of the RiboTag approach to retrieve functionally relevant mRNAs.

RiboTag analysis of microglial response to peripheral LPS challenge

Arguably, method-related artifacts, like the ones associated with microglia isolation, could be neutralized, if controls and experimental samples are prepared using the

same approach. However, this assumes that artifacts introduced by the isolation are not affected by biological treatments and challenges. To examine this issue, we performed the RiboTag protocol on TAM-treated *Cx3cr1^{CreER}; Rpl22^{HA}* mice following an intra-peritoneal lipopolysaccharide (LPS) injection (2.5 mg/kg). Brain hemispheres of individual LPS- and PBS-treated animals were subjected to either homogenization or microglia isolation and sorting, and processed as described in **Figure 2A**. A summary heat map of the RNAseq data is shown in **Supplementary Figure 4**. Unbiased hierarchical K-means clustering analysis for significantly changed genes (>2 fold change, $p < 0.05$) between at least two samples revealed 6 clusters. To define the effect of the isolation method on microglia of LPS-treated and PBS-treated mice, we performed separate analyses for differentially expressed genes between PBS and LPS treatment in each method and then assessed the respective overlap. The majority of genes detected as up- or down-regulated by the endotoxin challenge in the IP sample were shared with the sorted samples (**Figure 5A, 5C**). Mutual genes correlated between methods, and showed a similar trend of up and downregulation (**Figure 5B, 5D**), indicating that *bona fide* LPS-induced changes are seen with both methods. Upregulated genes included *Il4ra*, *Ch25h* and *Il1b* (**Figure 5B**), while microglia signature genes, such as *Tgfb1*, *Sall1* and *Cx3cr1*, were down regulated (**Figure 5D**).

Notably, a considerable number of mRNAs changed upon LPS treatment only in the sorted samples, but not in the anti-HA IP from whole brain extract (46% of the upregulated and 71% of the downregulated genes) (**Figure 5A, 5C**). Transcripts that were detected as changed upon LPS treatment in sorted samples (both Sort and Sort-IP) but not in IP-HA included genes related to immune activation like *Il1a*, *Ccl2* and *Vcam1* that were upregulated (**Figure 7E**) and *Tlr4*, *Siglech* and *Cd48* that were downregulated (**Figure 5E**). Since they were found in Sort-IP as well, these mRNAs are likely translated in sorted, but not unsorted microglia. These data establish that the artifact introduced by the cell isolation is affected by the state of the animals the cells are retrieved from. Moreover, mRNAs that are defined as 'cargo' contamination, due to their presence in the transcriptomes but absence from IPs of the sorted cells, changed following the LPS challenge (**Figure 5F**). For example *Pisd-ps1*, *Arhgap5* and *Tia1* were downregulated and *Ranbp2*, *Tet2* and *Gas5* were upregulated upon LPS challenge (**Figure 5F**). All of these genes were reported to be more highly expressed by other brain cells, rather than microglia (**Supplementary Figure 5**)⁹,

and are absent from the translomes of either sorted or unsorted microglia in our dataset.

Collectively, these results indicate that data retrieved from sorted microglia include false information that originates from sorting-related immune activation and cargo of ingested cells. Importantly, our data establish that identical processing of control and test samples does not necessarily neutralize these artifacts, since they themselves are affected by the biological treatment.

Discussion

Here we compared experimental approaches to retrieve microglia expression signatures from brains of untreated and challenged animals. Specifically, we define strengths and weaknesses of the classical cell isolation and sort-based protocols and the RiboTag strategy⁴, that relies on poly-ribosome immuno-precipitation from crude tissue extracts. Below we discuss the pros and cons of the respective techniques (**Table 1**).

The RiboTag strategy was originally introduced by McKnight and colleagues and applied to expression profiling of neurons and Sertoli cells⁴. Cell type-specificity of the approach depends on the accuracy of the Cre-drivers that is combined with the *Rpl22^{HA}* allele. This aspect is highlighted in our study by the side-by-side comparison of CX₃CR1^{Cre} and CX₃CR1^{CreER} animals, which revealed the superiority of the inducible system to achieve brain macrophage specificity and exclude neurons. However, as reported earlier and confirmed in this study, even CX₃CR1^{CreER} mice not only target microglia, but also non-parenchymal macrophages¹³. In situations where subpopulations can be phenotypically discriminated, as in the case of CD45^{int} microglia and CD45^{hi} perivascular macrophages, sort-based approaches, potentially combined with a RiboTag analysis like is this study, can hence be advantageous. This emphasizes the need for the development of novel Cre transgenic lines targeting microglia, including combinatorial approaches, such as the split-Cre strategy²⁴, to improve the cell type- or lineage-specificity. Importantly, current tests for accuracy of Cre transgenic lines are based on their combination with reporter alleles and the analysis of resulting double transgenic animals by flow cytometry. Like in the case of CX₃CR1^{Cre} animals reported here, or the LysM^{Cre} mice, that were assumed to be myeloid cell-specific, but are now known also to target neurons²⁵, this approach is insufficient. Our results demonstrate that the RiboTag approach provides

a useful complementary method to determine Cre line accuracy, in particular for cell types such as neurons and endothelial cells that are notoriously difficult to isolate for flow cytometric analysis.

Macrophage expression signatures tend to be contaminated by material the phagocytes ingested from their surrounding. Since the RiboTag strategy only retrieves mRNAs that are associated with HA-epitope tagged host cell ribosomes, it excludes such exogenous material, and hence allows identification of *bona fide* macrophage mRNAs.

Although not directly addressed in this study, an additional benefit of the RiboTag approach is the fact, that it can be used to determine the impact of conditional mutations. Specifically, Cre-mediated rearrangements result in parallel mutagenesis and induction of the Rpl22^{HA} allele. We, for instance, recently used this approach to define the impact of an *Mecp2* deficiency on macrophages in brown adipose tissue BAT, comparing translomes retrieved from BAT of *Cx3cr1^{CreER}:Rpl22^{HA}* and *Cx3cr1^{CreER}:Rpl22^{HA}:Mecp2^{fl/y}* mice ²⁶.

Ribosome-associated mRNAs, as retrieved by the RiboTag approach are considered to reflect the translome. Interestingly though, a recent study of the microglial response to LPS using an analog experimental system, showed that mRNAs induced by the challenge can be prevented from translation by binding of Serine/Arginine-Rich Splicing Factor 3 (SRSF3) to their 3'UTRs ²⁶. Like other emerging layers of posttranscriptional expression control, this mechanism, which was revealed by a combined transcriptome and peptidome analysis on immunoprecipitated ribosomes, requires further study.

Above we listed advantages of the RiboTag approach, but an inherent weakness of this protocol is the fact that for confident assessment of gene expression in the targeted population, the method relies on an enrichment of the specific mRNA over the input, *i.e.* the whole tissue extract. The RiboTag approach hence precludes statements on the expression of genes that are equally expressed in the target cells and the surrounding tissue. For assess these co-expressed genes, sort based strategies might be superior, although these also bear caveats, as outline below.

Side-by-side comparison of the translomes isolated by IP from crude tissue extracts and from sorted microglial cells with whole cell transcriptomes revealed a number of short-comings of the latter. First, we noted a prominent activation

signature that is presumably introduced during the process of cell extraction from their tissue context. This artifact comprised pro-inflammatory genes, such as *CD86*, *Tlr4* and *Tlr7*, and will have to be considered in microglia profiling studies. Importantly, this robust and reproducible artifact could not be discerned, when control and test samples were processed similarly. Rather we found that the isolation procedure had a differential impact on microglia retrieved from either challenged or unchallenged animals.

Taken together, our study shows that cell isolation coupled with sort-based methods and the RiboTag approach both have strengths and weaknesses, which should be considered when experiments are designed and conclusions are drawn. Cell isolation bears the risk of artifacts that might significantly confound transcriptome-based studies, including single cell analysis. Our study should hence caution experimentalists and make them aware of the 'observer effect', that is well established in physics, but often less appreciated in biology.

Material and Methods

Microglia isolation protocols

Mice were anesthetized with Pental (1:2 in PBS) and were perfused with PBS. Brains were dissected, crudely chopped and incubated for 20 min at 37°C in a 1 ml HBSS solution containing 2% BSA, 1 mg/ml Collagenase D (Sigma) and 50µg/ml DNase1 (Sigma). In the middle of incubation homogenates were pipetted for further dissociation. Next the homogenates were filtered through a 150µm mesh, washed with cold FACS buffer (2% FCS, 1mM EDTA in PBS-/-) and centrifuged at 2200 RPM, at 4°C, for 5 min. For the enrichment of microglia, the cell pellet was re-suspended with 3ml of 40% percoll solution and centrifuged at 2200 RPM, no acceleration and breaks, at RT for 15 min. Next, the resuspended cell pellet was passed through 80µm mesh, washed with 5ml FACS buffer, centrifuged at 1400 RPM at 4°C for 5 min and then taken for antibody (Ab) labeling and flow cytometry analysis. For the protocol excluding the digest, brains were chopped and then filtered through a 150µm mesh. The subsequent steps were as above, but without the enzymatic digest.

Tamoxifen treatment

To induce gene recombination in CreER transgenic mice, tamoxifen (TAM) was dissolved in warm corn oil (Sigma) and administered orally via gavage for four times every other day. All animals were TAM-treated first at 4-6 weeks of age. Each oral application was of 5mg at a concentration of 0.1 mg / μ l. Mice were examined 8 weeks following treatment. For the LPS treatment, mice were injected intraperitoneally (i.p.) with a single dose of LPS [2.5 mg/kg; E. coli 0111:B4; Sigma]; controls received or the same volume of vehicle solution (PBS).

Flow cytometry and cell sorting

Antibodies against CD11b (M1/70), Ly6C/G (Gr-1) (RB6-8C5), CD45 (30-F11) purchased from Biolegend or eBioscience were used. Samples were flow sorted using AriaIII (BD Biosciences, BD Diva Software) cell sorter. Analysis was performed on Fortessa (BD Biosciences, BD Diva Software) and analyzed with FlowJo software (Treestar).

Histology

Mice were anesthetized with penthal (1:2) and were perfused with PBS. Brains and spinal cords were excised and fixed for 4 hours in 2% paraformaldehyde. Brains were incubated for 72 in 30% sucrose and spinal cords in 18% EDTA prior to OCT (TissueTek) imbedding and freezing. Post fixed and stained 20-15nm frozen sections were blocked in 2% horse serum for 2 hours and incubated overnight (O.N) in 4°C with first antibody. Sections were washed 3 times in PBSX1 0.02% TritonX (Sigma) and conjugated with 2nd antibody for 1 hour in 4°C. Before covering samples were washed 3 times and incubated for 5 minutes with Hoechst. Sections were analyzed by confocal laser scanning microscope Olympus BX51. Image acquisition was processed by Olympus image browser software. The following antibodies have been used: Mouse monoclonal anti-HA (1:200, Covance), Rabbit polyclonal anti-Iba1 (1:250, Wako), Goat polyclonal anti-GFP (1:200, abcam), Goat polyclonal anti-Chat (1:100, Millipore) and Hoechst 33342 (1:10,000, Invitrogen).

Immunofluorescence

Mouse brains were fixed in 4% para-formaldehyde and subsequently paraffin-embedded by standard routines. AT sections (7 μ m thick) were deparaffinized in xylene and rehydrated through descending grades of ethanol to water. Antigen

retrieval was performed in Tris/EDTA buffer at pH 9.5 twice for 5 minutes at 95 °C. Sections were rinsed in PBS supplemented with 0.3% Triton-X (Sigma-Aldrich, St Louis, Missouri, USA) 3 times for 5 minutes. Unspecific binding sites were blocked using 1% bovine serum albumin in PBS supplemented with 0.3% Triton-X for 30 minutes at room temperature. Primary antibodies, such as chicken anti-GFP (1:100; Abcam) and rabbit anti-Iba1 (1:200; WAKO; Richmond, VA) were incubated overnight at 4 °C. For detecting primary antibodies, fluorochrome-conjugated secondary antibodies (1:200 each; all from Invitrogen; Karlsruhe, Germany) were applied for 1 h at RT. Autofluorescence of the tissue was quenched by using pre-warmed 0.3% sudan black for 2 minutes. Nuclear counterstain was performed by using 4,6-diamidin-2-phenylindol (DAPI; 1:10,000 in PBS) for 5 minutes, followed by 3 buffer rinses. Finally, sections were embedded with Dako immunofluorescence mounting medium (DAKO, Glostrup, Denmark). Images were taken using a FV1000 confocal laser scanning microscope (Olympus, Hamburg, Germany).

RiboTag approach.

Samples were extracted from mice, flash-frozen in liquid nitrogen and stored at -80 °C until use. Samples were homogenized on ice in ice-cold homogenization buffer (50 mM Tris, pH 7.4, 100 mM KCl, 12 mM MgCl₂, 1% NP-40, 1 mM DTT, 1:100 protease inhibitor (Sigma), 200 units/ml RNasin (Promega) and 0.1 mg/ml cycloheximide (Sigma) in RNase free DDW) 10% wt/vol with a Dounce homogenizer (Sigma) until the suspension was homogeneous. To remove cell debris, 1 ml of the homogenate was transferred to an Eppendorf tube and was centrifuged at 10,000g and 4 °C for 10 min. Supernatants were transferred to a fresh Eppendorf tube on ice, then 10 µl was removed for 'input' analysis and 5 µl of anti-HA (H9658, Sigma) or 5 µl of mouse IgG (sigma) was added to the supernatant, followed by 4 h of incubation with slow rotation in a cold room at 4 °C. Meanwhile, Dynabeads Protein G (Thermo Fisher Scientific), 100 µl per sample, were equilibrated to homogenization buffer by washing three times. At the end of 4 h of incubation with antibody, beads were added to each sample, followed by incubation overnight in cold room at 4 °C. After not more than 12 h, samples were washed three times with high-salt buffer (50 mM Tris, 300 mM KCl, 12 mM MgCl₂, 1% NP-40, 1 mM DTT, 1:200 protease inhibitor, 100 units/ml RNasin and 0.1 mg/ml cycloheximide in RNase free DDW), 5 min per wash in a cold room on a rotator. At the end of the washes, beads were magnetized and excess

buffer was removed, 150µl Lysis Buffer was added to the beads and RNA was extracted with Dynabeads mRNA Direct purification kit (Thermo Fisher). RNA was eluted in 6µl H₂O and taken for RNA sequencing.

For Sort-IP samples (RiboTag IP after sorting), ~50-100*10³ cells were sorted into cold PBS, centrifuged 400g for 5 min at 4°C. Supernatant was removed and pellet was re-suspended in 1ml of lysis buffer, the rest of IP was continued as above.

RNA sequencing

RNA-Seq of populations was performed as described previously¹. In brief, 5,000 microglia cells were sorted into 50µl of Lysis Buffer (Life Technologies) and stored at -80°C. mRNA was captured with Dynabeads oligo(dT) (Life Technologies) according to manufacturer's guidelines. We used a derivation of MARS-seq²⁹. Library concentration was measured with a Qubit fluorometer (Life Technologies) and mean molecule size was determined with a 2200 TapeStation instrument. RNA-Seq libraries were sequenced using Illumina NextSeq-500.

Data Analysis

Raw reads were mapped to the genome (NCBI37/mm9) using hisat (version 0.1.6). Only reads with unique mapping were considered for further analysis. Gene expression levels were calculated using the HOMER software package (analyzeRepeats.pl rna mm9 -d <tagDir> -count exons -condenseGenes -strand + - raw)³⁰. Normalization and differential expression analysis was done using the DESeq2 R-package. Differential expressed genes were selected using a 2-fold change cutoff between at least two populations and adjusted pValue for multiple gene testing < 0.05. Gene expression matrix was clustered using k-means algorithm (matlab function kmeans) with correlation as the distance metric. Heatmaps were generated using Genee software.

QUANTIFICATION AND STATISTICAL ANALYSIS

In all experiments, data are presented as mean ± STD if not stated otherwise. Statistical tests were selected based on appropriate assumptions with respect to data distribution and variance characteristics. Statistical significance was defined as P<0.05. Sample sizes were chosen according to standard guidelines. Number of animals is indicated as 'n'. Of note, sizes of the tested animal groups were also

dictated by availability of the transgenic strains and litter sizes, allowing littermate controls. Pre-established exclusion criteria are based on IACUC guidelines. Animals of the same age, sex and genetic background were randomly assigned to treatment groups. The investigator was not blinded to the mouse group allocation. Tested samples were blindly assayed.

ACCESSION NUMBERS

The accession numbers for the RNA-seq datasets reported in this paper can be found at GEO: xxxx

SUPPLEMENTAL INFORMATION

Supplemental Information includes Supplemental Experimental Procedures and 5 figures and a table, and can be found with this article online at

AUTHOR CONTRIBUTIONS

ACKNOWLEDGEMENTS

We would like to thank all members of the Jung laboratory for helpful discussion. We further thank the staff of the Weizmann Animal facility, members of the FACS facility for expert advice and G. Friedlander for help with the bioinformatics. We would like to thank C. Glass for sharing sequencing raw data. The Jung laboratory was supported by the Israeli Science Foundation (887/11), the European Research Council (Adv ERC 340345), and the Deutsche Forschungsgemeinschaft (CRC/TRR167 'NeuroMac').

References

1. Lavin, Y. *et al.* Tissue-Resident Macrophage Enhancer Landscapes Are Shaped by the Local Microenvironment. *Cell* **159**, 1312–1326 (2014).
2. Gosselin, D. *et al.* Environment Drives Selection and Function of Enhancers Controlling Tissue-Specific Macrophage Identities. *Cell* **159**, 1327–1340 (2014).
3. Heiman, M. *et al.* A Translational Profiling Approach for the Molecular Characterization of CNS Cell Types. *Cell* **135**, 738–748 (2008).
4. Sanz, E. *et al.* Cell-type-specific isolation of ribosome-associated mRNA from complex tissues. *Proceedings of the National Academy of Sciences* **106**, 13939–

- 13944 (2009).
5. Yona, S. *et al.* Fate mapping reveals origins and dynamics of monocytes and tissue macrophages under homeostasis. *Immunity* **38**, 79–91 (2013).
 6. Jung, S. *et al.* Analysis of fractalkine receptor CX(3)CR1 function by targeted deletion and green fluorescent protein reporter gene insertion. *Molecular and Cellular Biology* **20**, 4106–4114 (2000).
 7. Davalos, D. *et al.* ATP mediates rapid microglial response to local brain injury in vivo. *Nat Neurosci* **8**, 752–758 (2005).
 8. Goldmann, T. *et al.* A new type of microglia gene targeting shows TAK1 to be pivotal in CNS autoimmune inflammation. *Nature Publishing Group* **16**, 1618–1626 (2013).
 9. Zhang, Y. *et al.* An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J. Neurosci.* **34**, 11929–11947 (2014).
 10. Orthgiess, J. *et al.* Neurons exhibit Lyz2 promoter activity in vivo: Implications for using LysM-Cre mice in myeloid cell research. *Eur. J. Immunol.* **46**, 1529–1532 (2016).
 11. Kim, K.-W. *et al.* In vivo structure/function and expression analysis of the CX3C chemokine fractalkine. **118**, e156–67 (2011).
 12. Fonseca, M. I. *et al.* Cell-specific deletion of C1qa identifies microglia as the dominant source of C1q in mouse brain. 1–15 (2017). doi:10.1186/s12974-017-0814-9
 13. Goldmann, T. *et al.* Origin, fate and dynamics of macrophages at central nervous system interfaces. *Nat Immunol* **17**, 797–805 (2016).
 14. Zeisel, A. *et al.* Cell types in the mouse cortex and hippocampus revealed by single-cell RNA-seq. *Science* 1–8 (2015). doi:10.1126/science.aaa1934
 15. van den Brink, S. C. *et al.* Single-cell sequencing reveals dissociation-induced gene expression in tissue subpopulations. *Nat Meth* **14**, 935–936 (2017).
 16. A-Gonzalez, N. *et al.* Phagocytosis imprints heterogeneity in tissue-resident macrophages. *J. Exp. Med.* **281**, jem.20161375–16 (2017).
 17. Derrien, T. *et al.* 2012. “The GENCODE v7 Catalogue of Human Long Non-Coding RNAs : Analysis of Their Structure , Evolution and Expression.” *Genome Research* **22**:1775–89.
 18. Braunschweig, Ulrich *et al.* 2014. “Widespread Intron Retention in Mammals

- Functionally Tunes Transcriptomes.” *Genome Research* 24(11):1774–86.
19. Khong, Anthony et al. 2017. “The Stress Granule Transcriptome Reveals Principles of mRNA Accumulation in Stress Granules.” *Molecular Cell* 68(4):808–820.e5. Retrieved (<https://doi.org/10.1016/j.molcel.2017.10.015>).
 20. Hubstenberger, Arnaud et al. 2017. “P-Body Purification Reveals the Condensation of Repressed mRNA Regulons.” *Molecular Cell* 68(1):144–157.e5.
 21. Tebaldi, Toma et al. 2012. “Widespread Uncoupling between Transcriptome and Translatome Variations after a Stimulus in Mammalian Cells.” *BMC Genomics* 13(1):220. Retrieved (<http://www.biomedcentral.com/1471-2164/13/220>).
 22. Bahar Halpern, Keren et al. 2015. “Nuclear Retention of mRNA in Mammalian Tissues.” *Cell Reports* 13(12):2653–62. Retrieved (<http://dx.doi.org/10.1016/j.celrep.2015.11.036>).
 23. Tan, Jennifer Y. et al. 2015. “Extensive microRNA-Mediated Crosstalk between lncRNAs and mRNAs in Mouse Embryonic Stem Cells.” *Genome Res.* 25(5):655–66.
 24. Hirrlinger, J. et al. Split-cre complementation indicates coincident activity of different genes in vivo. *PLoS ONE* 4, e4286 (2009).
 25. Orthgiess, J. et al. Neurons exhibit Lyz2 promoter activity in vivo: Implications for using LysM-Cre mice in myeloid cell research. *Eur. J. Immunol.* 46, 1529–1532 (2016).
 26. Wolf, Y. et al. Brown-adipose-tissue macrophages control tissue innervation and homeostatic energy expenditure. *Nat Immunol* 137, 495 (2017).
 27. Boutej et al. Diverging mRNA and Protein Networks in Activated Microglia Reveal SRSF3 Suppresses Translation of Highly Upregulated Innate Immune Transcripts. *Cell Rep.* 21, 3220-3233 (2017).
 28. Tilgner, Hagen et al. 2012. “Deep Sequencing of Subcellular RNA Fractions Shows Splicing to Be Predominantly Co-Transcriptional in the Human Genome but Inefficient for lncRNAs.” *Genome Research* 22(9):1616–25.
 29. Jaitin, D. A. et al. Massively parallel single-cell RNA-seq for marker-free decomposition of tissues into cell types. *Science* 343, 776–779 (2014).
 30. Heinz, S. et al. Simple Combinations of Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for Macrophage and B Cell Identities. *Molecular Cell* 38, 576–589 (2010).

FIGURE LEGENDS

Figure 1. RiboTag analysis reveals that $Cx3cr1^{Cre}$ mice but not $Cx3cr1^{CreER}$ animals display rearrangements in neurons.

(A) Scheme of CX_3CR1 -Cre and CX_3CR1 -CreER systems.

(B) Scheme describing the immuno-precipitation protocol, including brain homogenization, centrifugation to remove cell debris and incubation with magnetic beads and relevant antibodies.

(C) Heat maps of RNAseq data comparing IPs obtained from brains of $Cx3cr1^{Cre}:Rpl22^{HA}$ and $Cx3cr1^{CreER}:Rpl22^{HA}$ mice, represented by lists of genes of microglia (115), neurons (97), astrocytes (95) and oligodendrocytes (98) showing enrichment and de-enrichment of mRNAs of specific cell types in the different samples. Reference data sets⁹. Each column represents an individual mouse, n=2-3 in each group.

(D) Microscopic analysis of cortex brain sections from $Cx3cr1^{Cre}:R26-YFP$ mice (left panels) and $Cx3cr1^{CreER}:R26-YFP$ mice (TAM treated (right panels) or untreated controls (middle panels), stained for IBA-1, YFP and DAPI, showing neuronal expression of YFP in $Cx3cr1^{Cre}$ brains and microglia-restricted YFP expression in $Cx3cr1^{CreER}$ brains.

(E) Immuno-fluorescent staining of brain sections from $Cx3cr1^{Cre}:Rpl22^{HA}$ (left) and TAM-treated $Cx3cr1^{CreER}:Rpl22^{HA}$ (right) mice, stained for IBA1, HA and Hoechst, showing neuronal expression of HA in $Cx3cr1^{Cre}$ brains, and microglia-restricted HA expression in $Cx3cr1^{CreER}$ brains.

(F) Flow cytometry analysis showing HA staining in microglia ($CD11b^+ CD45^{int}$, gated on $Ly6C/G^- DAPI^-$) of $Rpl22^{HA}$ TAM-treated mice (black line), $Cx3cr1^{CreER}:Rpl22^{HA}$ mice, untreated (blue line) and TAM-treated (red line). Shaded histogram represents isotype control.

Figure 2. Comparison of cell sort-based protocol and the RiboTag method to profile microglia

(A) Scheme describing the experimental protocol comparing RiboTag and cell sort-based strategies.

(B) Heat map of RNAseq data of samples obtained in (A). Genes selected by maximum value > 100 normalized reads (3,186 out of 17,406 genes), significantly

changed (fold change >2, p-value <0.05) between IP-HA vs IP-IgG, Sort vs IP HA, Sort-IP vs IP HA and Sort vs Sort-IP, Representing 2,508 genes. n=3.

(C) Heatmap representing K-means re-clustering of genes from cluster III from Figure 3B, showing genes high in IP-HA and low in Sort samples.

(D) FACS dot plot (left panel) showing separation of microglia (CD45^{int}) from other brain macrophages (MΦ) (CD45^{hi}) by flow cytometry. Histogram (right panel) of microglia and MΦ in *Cx3cr1*^{GFP} mice indicating high CX₃CR1/GFP expression in both populations.

(E) FACS histogram of HA staining in microglia (left panel) and other brain macrophages (MΦ) (right panel) in control *Rpl22*^{HA} mice (grey) or TAM-treated *Cx3cr1*^{CreER}:*Rpl22*^{HA} mice (blue/red).

(F) Heatmap of RNA-seq data of representative non-parenchymal brain macrophages genes, showing enrichment in IP-HA, but not in sorted samples.

(G) Graphs showing normalized reads of example genes from Figure 4D. Each dot represents an individual mouse, n=3.

(H) Graphs showing normalized reads of example genes from cluster III-b and III-c in Figure 4A, showing functional genes enriched in IP and Sort-IP. Each dot represents an individual mouse, n=3.

Figure 3. Microglia isolation results in cell activation.

(A) Heatmap representing K-means re-clustering of genes from cluster II from Figure 3B, showing mRNAs high in Sort and low in IP-HA samples.

(B) Graphs showing normalized reads of example genes from cluster II-a in Figure 5A, showing high level expression of immune-activation-related genes in sorted samples. Each dot represents an individual mouse, n=3.

(C) Ingenuity Pathway Analysis of genes significantly higher (>2 fold change, p-value <0.05) in Sort compared to IP-HA, representing pathways related to immune activation.

Figure 4. Microglia transcriptomes, but not translatoemes include cargo-derived mRNAs and mRNAs sequestered in nuclei and P bodies

(A) Venn-Diagram overlapping genes of cluster II-b (Figure 2c) (yellow) with genes of non-microglial cells, as selected according to ⁹. Non-microglia genes were selected

by being with max value > 10, microglia reads < 30 and max value is > 3 fold over microglia. Genes fitting these criteria represent cargo of ingested cells.

(B) Graphs showing normalized reads of example genes from list of shared genes from Figure 6A, showing high levels cargo contamination in Sort samples, but not in IP. Each dot represents an individual mouse, n=3 (Upper panels). Expression of genes above in different brain cells; data obtained from ⁹. (Lower panels). FKPM, Fragments per kilobase of exon per million reads mapped.

(C) Graphs showing normalized reads of example genes of long non coding RNAs that reside within the nucleus and are presented only in Sort but not IP samples. Each dot represents an individual mouse, n=3.

(D) Violin plot representing splicing efficiency (left panel) and gene length (right panel) of genes in cluster II-b (orange) compared to genes not in II-b (green), showing that genes in cluster II-b are less efficiently spliced and have longer genes and longer 3'UTRs compared with the rest of genes in the dataset, suggesting nuclear retention. Splicing efficiency was computed by comparing intron-spanning and intron-crossing reads ²⁸. * = Wilcoxon test p-value < 10⁻⁶

(E) Violin plot representing cellular localization of genes in other tissues (left panel Liver, right panel – MIN6 pancreatic beta cell line) with established nuclear and cytoplasmic fractions. Genes in cluster II-b (orange) are more nuclear compared to other genes (green). * = Wilcoxon test p-value < 10⁻⁶

(F) Graphs showing normalized reads of immediate-early genes found in Sort but not in IP-HA (in cluster II-b, microglial genes), suggesting sequestration from translation in unsorted cells. Each dot represents an individual mouse, n=3.

(G) Diagram representing different states of mRNAs in the cell: nuclear retention, sequestration from translation in P-bodies or active translation in the ribosomes.

Figure 5. Analysis of microglia isolated from mice challenged with LPS

(A) Venn diagram overlapping genes upregulated by LPS treatment in IP-HA (blue) and in Sort (yellow), showing 54 genes upregulated in IP-HA only, 247 genes shared between methods and 214 genes upregulated by sorting only.

(B) Correlation analysis of 247 shared genes (from figure 7A) upregulated due to LPS in both methods, representing log2 fold change of significantly changed genes (log2 fold change>1, pValue<0.05) in LPS vs PBS in each of the methods (Left panel); Graphs showing normalized reads of example genes detected as upregulated

with LPS by both methods. Each dot represents an individual mouse, n=3-4 in each group (Right panels).

(C) Venn diagram overlapping genes downregulated by LPS treatment in IP-HA (blue) and in Sort (yellow), showing 10 genes downregulated in IP-HA only, 104 genes shared between methods and 250 genes downregulated by sorting only.

(D) Correlation analysis of 104 shared genes (from figure 5C) downregulated due to LPS in both methods, representing log₂ fold change of significantly changed genes (log₂ fold change < -1, pValue < 0.05) in LPS vs PBS in each of the methods (Left panel); Graphs showing normalized reads of example genes detected as downregulated with LPS by both methods. Each dot represents an individual mouse, n=3-4 in each group (Right panels). For IP-HA genes were selected by first being enriched over control IP-IgG, and then by LPS>PBS

(E) Graphs showing normalized reads of example genes related to immune activation that were up- (upper panels) or down- (lower panels) regulated due to LPS treatment in sorted samples only but not detected in IP-HA, showing differential susceptibility of biologically treated samples to artifacts introduced by isolation method. Each dot represents an individual mouse, n=3-4 in each group.

(F) Graphs showing normalized reads of example genes originated from ingested cargo that were up- (upper panels) or down- (lower panels) regulated due to LPS treatment in Sort only but not detected in IP-HA or Sort-IP, showing that the whole transcriptome includes LPS-dependent changes that do not originate in microglia thus introducing false information. Each dot represents an individual mouse, n=3-4 in each group.

Supplementary Figures

Supplementary Figure 1. Immunohistochemistry analysis of brains of reporter animals

Microscopic analysis of hypothalamus sections from *Cx₃cr1^{Cre}:R26-YFP* mice (left panels) and *Cx₃cr1^{CreER}:R26-YFP* mice (TAM treated (right panels) or not (middle panels), stained for IBA-1, GFP and DAPI, showing neuronal expression of GFP in *Cx₃cr1^{Cre}* brains and microglia-restricted GFP expression in *Cx₃cr1^{CreER}* brains.

Supplementary Figure 2. Microglial genes are comparable among retrieval methods.

Graphs showing normalized reads of example genes from cluster I of Figure 3B, showing microglial genes at comparable reads number, indicating that retrieval methods are comparable.

Supplementary Figure 3. Microglia activation signature is robust, reproducible and induced by cell extraction form tissue

(A) Heatmap clustered by K-means clustering, comparing direct IP to whole mRNA retrieved from sorted cells extracted with (+coll) or without (-coll) enzymatic digestion in the isolation protocol, revealing 4 clusters similar to figure 3B.

(B) Correlation plots of average of log₂ normalized reads plotting sorted cells with enzymatic digestion on the Y axis and sorted cells without enzymatic digestion on the X axis (left panel) or IP on the X axis (right panel).

(C) Correlation matrix combining two independent experiments, showing high reproducibility of the results.

Supplementary Figure 4. Analysis of microglia isolated from mice challenged with LPS

(A) Heatmap of RNA-seq data comparing RiboTag to sorting with LPS treatment, taking significantly changed genes (fold change>2, p-value<0.05).

(B) Graphs showing normalized reads of example genes taken from cluster I from figure S3A, showing similar levels of microglia signature genes among samples. Each dot represents an individual mouse, n=3-4 in each group.

Supplementary Figure 5. Expression profiles of selected genes across brain cell populations

Non-microglia mRNAs suggesting cargo contamination (Figure 5F)

Data obtained from (http://web.stanford.edu/group/barres_lab/brain_rnaseq.html)⁹.




Figure 1




Figure 3




Figure 3




Figure 4







Figure 5

Cx₃cr1^{Cre}:R26-YFP *Cx₃cr1^{CreER}:R26-YFP*
with TAM treatment






Supplementary Figure 2





Supplementary Figure 3

A



B





Supplementary Figure 5