

Published in final edited form as:

Kathmandu Univ Med J (KUMJ). 2010 ; 8(29): 62–72.

Causes of stillbirths and neonatal deaths in Dhanusha district, Nepal: A verbal autopsy study

SR Manandhar¹, A Ojha¹, DS Manandhar², B Shrestha², D Shrestha², N Saville³, AM Costello³, and D Osrin³

¹Department of Paediatrics, Kathmandu Medical College, Sinamangal

²Mother and Infant Research Activities (MIRA), Nepal

³UCL Centre for International Health and Development, Institute of Child Health, London, UK

Abstract

Background—Perinatal (stillbirths and first week neonatal deaths) and neonatal (deaths in the first 4 weeks) mortality rates remain high in developing countries like Nepal. As most births and deaths occur in the community, an option to ascertain causes of death is to conduct verbal autopsy.

Objective—The objective of this study was to classify and review the causes of stillbirths and neonatal deaths in Dhanusha district, Nepal.

Materials and Methods—Births and neonatal deaths were identified prospectively in 60 village development committees of Dhanusha district. Families were interviewed at six weeks after delivery, using a structured questionnaire. Cause of death was assigned independently by two pediatricians according to a predefined algorithm; disagreement was resolved in discussion with a consultant neonatologist.

Results—There were 25,982 deliveries in the 2 years from September 2006 to August 2008. Verbal autopsies were available for 601/813 stillbirths and 671/954 neonatal deaths. The perinatal mortality rate was 60 per 1000 births and the neonatal mortality rate 38 per 1000 live births. 84% of stillbirths were fresh and obstetric complications were the leading cause (67%). The three leading causes of neonatal death were birth asphyxia (37%), severe infection (30%) and prematurity or low birth weight (15%). Most infants were delivered at home (65%), 28% by relatives. Half of women received an injection (presumably an oxytocic) during home delivery to augment labour. Description of symptoms commensurate with birth asphyxia was commoner in the group of infants who died (41%) than in the surviving group (14%).

Conclusion—The current high rates of stillbirth and neonatal death in Dhanusha suggest that the quality of care provided during pregnancy and delivery remains sub-optimal. The high rates of stillbirth and asphyxial mortality imply that, while efforts to improve hygiene need to continue, intrapartum care is a priority. A second area for consideration is the need to reduce the uncontrolled use of oxytocic for augmentation of labour.

Keywords

Stillbirth; neonatal death; verbal autopsy; Nepal

Stillbirths and neonatal deaths remain common in many low-income countries. Nepal's most recent estimates of perinatal mortality rate (PMR: stillbirths and first week deaths) and

neonatal mortality rate (NMR: deaths in the first four weeks) are 45 per 1000 births and 33 per 1000 live births, respectively¹. About 80% of deliveries in Nepal occur at home and only 19% are conducted by health personnel¹. Over the last 15 years, Nepal's infant mortality rate (IMR: deaths in the first year) has fallen from 82 to 48 per 1000 live births¹. This is at least partly due to programmes conducted through government and other organizations, including those for acute respiratory infections and diarrhoeal diseases. There were no such programmes to reduce perinatal and neonatal deaths in Nepal, and their share of the overall figures has increased. If we are to reduce stillbirths and neonatal deaths, we need to know their causes. However, clinical registration of cause of death is available for less than a third of global newborn deaths². When families do not seek hospital care and medical records are unavailable, trying to understand the cause of death from their accounts of illness is the only option. Verbal autopsy is a method of ascribing causes of death on the basis of information on events, signs and symptoms supplied by the deceased's caretakers^{3,4,5}. Procedures for infant and child deaths have developed over the last twenty years,^{6,7,8,9,10,11,12} and verbal autopsy has been used in studies,^{13,14,15} in sample registration systems in India, and as part of the INDEPTH network of demographic surveillance sites¹⁶.

Because there is no gold standard with which to compare the findings, misclassification of causes of death is possible¹⁷. Occasionally, medical records are available and a process of triangulation can take place; or interviews with families whose children died in hospital can be compared with case notes and hospital records, even though the case-mix seen in hospitals is likely to differ from the pattern of illness in communities. There are several possible approaches to data collection. An interviewer (often with a non-medical background) may discuss the events that preceded death with a child's mother or family and record the narrative in open-ended form. Alternatively, a series of closed questions may be asked, designed to establish the presence or absence of particular signs and symptoms, or particular health care actions^{18,19}. At the analytical stage, the narrative or closed questions are read and classified by a clinician. The development of algorithms to aid this process has led to the idea of using computer analysis to classify cause of death, but this is not yet routine^{20, 21}.

Mother and Infant Research Activities (MIRA) is a non-government organisation (NGO) established to improve the health of mothers and newborn infants through research, training and advocacy. MIRA has conducted several community-based studies, including a trial of community mobilisation through women's groups in Makwanpur district, which showed a 30% reduction in neonatal mortality²². A similar study, with some modifications, is being conducted in Dhanusha district in the southern plains of Nepal. In the course of the study, stillbirths and neonatal deaths were identified in a surveillance population of around 420 000 and 1272 verbal autopsies were collected from 60 village development committees²⁴. The objective of this analysis was to classify and review the causes of stillbirths and neonatal deaths in the study area.

Materials and methods

Setting

Dhanusha is situated in the south-eastern region of Nepal and covers an area of 1180 km² with a population of around 777 000²³. In a study population of around 420 000, there are approximately 55 800 married women of reproductive age and around 14 000 annual births²⁴. The literacy rate is 56% and 89% of people live in rural areas²³. The district's PMR is 63 per 1000 births and its NMR 34 per 1000 live births²⁴.

Procedures

This was a prospective study that formed part of a major child survival study using women's groups to improve maternal and newborn health. Out of 101 village development committees (VDCs) in Dhanusha, after exclusion of 9 severely politically affected VDCs and 13 VDCs with populations of less than 4000, 60 were selected randomly for inclusion. After mapping, households were identified and data collected over two years. Each village development committee has an average population of around 8,000. In each, selected key informants provided information prospectively about every birth and death to village development committee interviewers. The interviewers verified each event and visited families at six weeks after delivery to administer a questionnaire. In the event of a stillbirth or neonatal death, they visited within three months to administer a verbal autopsy. The study was explained to participants and consent taken. We used a verbal autopsy tool which combined open-ended narrative with closed questions and was developed in light of other tools: the original WHO infant verbal autopsy²⁵, a tool updated in 2002 by the INDEPTH network, and a new tool in development (WHO/JHU/AKU/SNL 11/05/2003). Confidentiality was maintained by interviewing individuals or family groups separately and not discussing the matter openly.

Analytical methods

Questionnaires were entered in an electronic database in Microsoft Access (Microsoft Corporation). We based our classification of cause of death on the Neonatal and Intrauterine Death classification according to Etiology (NICE)²⁶, and the WHO Neonatal Child Health Epidemiology Reference Group (CHERG)²⁷. Narratives and responses to closed questions were reviewed independently by two pediatricians (SRM and AO). Cause of death was classified according to the algorithm shown in Box 1. Cases in which the two paediatricians did not agree were discussed with a consultant neonatologist (DSM) and a final diagnosis was assigned by consensus. Further analysis was carried out in SPSS 11.5 (SPSS Inc., Illinois). The findings are presented as frequencies and proportions.

Ethical issues

Study approval was obtained from the Nepal Health Research Council (NHRC), the ethical committee of Great Ormond Street Hospital and the Institute of Child Health and the Social Welfare Council of Nepal. Informed consent in local or Nepali languages was obtained from respondents prior to interview. No data used in the analysis contained the names of participants.

Results

Data were collected over two years from 1st September 2006 to 31st August 2008. There were 25 982 deliveries in this period, with 813 stillbirths and 954 neonatal deaths. Verbal autopsies were available for 601 stillbirths and 671 neonatal deaths (72% of all). Reasons for missing autopsies included political conflict and instability in the district, transportation problems and absence of male head of household to give permission for interview. Information on stillbirths, early and late neonatal deaths is presented in Table 1. The study PMR was 60 per 1000 births and the NMR 38 per 1000 live births. Deaths were slightly more common in male infants (56%) and most of the infants were born at term (76%).

The classification of cause of death is presented in Table 2. Of 601 stillbirths, 504 (84%) were fresh and 97 (16%) macerated. Obstetric complications were the leading cause (67%), followed by prematurity (10%). 10% of perinatal deaths were unclassifiable on the basis of the questionnaire. Among neonatal deaths, 529 (79%) were early and 142 (21%) late. The

three leading causes of neonatal deaths were birth asphyxia (37%), severe infection (30%) and prematurity or low birth weight (15%).

Table 3 summarises characteristics of delivery. Most of the infants (65%) were delivered at home. Many deliveries were conducted by relatives (28%), and only 26% were assisted by doctors. 20% were conducted by village practitioners who are untrained health workers. It is interesting to note that 50% of women received an injection (presumably of an oxytocic drug, Oxytocin) during home delivery to augment labour. Among neonatal deaths, 886 (74%) infants were reported by their mothers as being of normal size. In 53% of births, the umbilical cord was cut with unsterilised instruments, and dung or ash was applied to the cut end of the cord in 20% of cases. Nearly a third of infants were not fed anything within half an hour of birth and 23% were given goat's milk as the first feed.

Table 4 examines the circumstances of presumptive birth asphyxia and resuscitation. Among neonatal deaths, 41% of infants did not cry immediately after birth and some attempt at resuscitation was made in 90% of cases. Half of the infants with presumptive birth asphyxia did not cry at all. The commonest resuscitative measure was mouth-to-mouth respiration (49%). In terms of treatment in general – not only for birth asphyxia - 365 (56%) of 646 infants who died in the neonatal period did not receive any form of treatment. The main reasons given for this were poverty, unavailability of a male family member who could take a decision in time, and an assumption that infants would not survive. Of the 44% of infants who did receive treatment, 244 (88%) were seen by allopathic health care providers: 106 (38%) at hospitals, 63 (23%) at other institutions such as nursing homes and clinics, and 75 (27%) at home. 35 (12%) received treatment from providers such as traditional healers (*jhankri*) and herbalists.

Table 5 compares some of the characteristics of infants who died with those of infants who survived. Out of 25,169 surviving infants, information was available on 17,377 (69%). Migration of mothers to neighbouring districts, transportation problems and political conflicts were the main reasons for missing data. 1% of infants were preterm in the surviving group, compared to 24% in the group who died. Village practitioners delivered 6% of infants in the group who survived, and 19% of those who died. Description of symptoms of birth asphyxia was commoner in the group who died (41%) than in the surviving group (14%).

Discussion

The commonest cause ascribed to stillbirths was obstetric complications. This may reflect either delay in receiving appropriate obstetric care due to geographical difficulty or poor antepartum and intrapartum obstetric care. The fact that 504 stillbirths (84%) were fresh suggests that infants might have been viable during labour. Better intrapartum foetal monitoring with timely intervention might reduce the fresh stillbirth rate. Lack of skilled providers in the community could be one reason for the high fresh stillbirth rate. Birth asphyxia was found to be the commonest cause of neonatal death. This also indicates a lack of proper intrapartum monitoring, timely and appropriate intervention and neonatal resuscitation in the community. Indeed, the combination of fresh stillbirths and early neonatal deaths from asphyxia probably reflects the same process²⁸. In this study population, 65% of deliveries were at home and most (59%) were assisted by untrained persons. This reflects the national scenario in which 80% of deliveries are conducted by relatives at home²⁹.

A verbal autopsy analysis from Egypt found prematurity (39%), asphyxia (18%) and infections (7%) to be the commonest causes of neonatal deaths³⁰. Among perinatal deaths,

birth asphyxia was the commonest cause (44%). In a study in rural Pakistan, the main causes of neonatal death were asphyxial conditions, neonatal sepsis and conditions associated with prematurity³¹. 65% of neonatal deaths occurred at home and most of the deliveries were conducted by *dais* (untrained birth attendants). In a study from rural Maharashtra (India), Kulkarni et al found that low birth weight (36%) was the main cause of early neonatal deaths, followed by prematurity (26%)³². Prematurity was the commonest cause of stillbirth, and women under 21 years old were reported as having higher rates of premature delivery.

Few community verbal autopsy studies have been done in Nepal. In Morang district, Khanal et al found that infection (41%), birth asphyxia (37%) and prematurity or low birth weight (18.4%) were common causes of neonatal death³³. They also found that infection was seen mostly after home deliveries. Wrapping of infants with unclean cloth, cutting the umbilical cord with unsterilised instruments, and birth attendance conducted by relatives, were possible reasons for this. Our study also showed that severe infection was the second most common cause of neonatal death. In as many as 20% of cases, dung or ash was applied to the umbilical stump. A similar study from Sarlahi district found that birth asphyxia (33%), serious infection (30%) and prematurity (29%) were the commonest causes of neonatal death³⁴. This again suggests sub-optimal intrapartum and postnatal care in the community. The study also identified 30 cases of neonatal tetanus, which may have been due to inadequate coverage of tetanus toxoid immunisation and traditional cord care practices.

Our study highlights another important issue. Neonatal deaths caused by infection tend to occur later, while early deaths are more often associated with birth asphyxia³⁵. In previous studies, and in global estimates, deaths from infection, preterm and asphyxia have tended to occur at similar rates^{27,36,37}. The proportion of neonatal deaths in the first week has tended to be 60-70%. In our study, early neonatal deaths accounted for almost 80% of neonatal deaths, and asphyxia 41%. This means that asphyxial deaths are becoming the dominant problem, an interpretation that is intuitive in the sense that the numbers of late neonatal deaths are falling. An emphasis on intrapartum care is required, and this fits well with the safer motherhood agenda and recommendations. One worrying issue is the high incidence of oxytocic injection, often delivered in uncontrolled environments. This potentially harmful practice has become common and our finding of a level of 50% is a cause for concern that supports both anecdotal information about practices in the district and previous research³⁸.

Conclusion

The current high rates of stillbirth and neonatal death in Dhanusha suggest that the quality of care provided during pregnancy and delivery remains sub-optimal. The high rates of stillbirth and asphyxial mortality imply that, while efforts to improve hygiene need to continue, intrapartum care is a priority. A second area for consideration is the need to reduce the uncontrolled use of oxytocics for augmentation of labour.

Acknowledgments

Thanks are due to all the staff involved in the collection of data in the community, and to the mothers and relatives who agreed to be interviewed in a distressing situation.

References

1. Population division. Ministry of Health and Population (MOHP). Government of Nepal. Infant and child mortality. In: New ERA and Macro International Inc. , editor. Nepal Demographic and Health Survey (NDHS) 2006. Ministry of Health and Population, New ERA, and Macro International Inc; Kathmandu: 2007. p. 125

2. Lawn JE, Osrin D, Adler A, et al. Four million neonatal deaths: counting and attribution of cause of death. *Paediatr Perinat Epidemiol.* 2008; 22:410–6. [PubMed: 18782248]
3. Garenne M, Fauveau V. Potential and limits of verbal autopsies. *Bull WHO.* 2006; 84:164–5. [PubMed: 16583068]
4. Soleman N, Chandramohan D, Shibuya K. Verbal autopsy: current practices and challenges. *Bull World Health Organ.* 2006; 4(3):239–45. [PubMed: 16583084]
5. Baiden F, Bawan A, Binka R, et al. Setting international standards for verbal autopsy. *Bull World Health Organ.* 2007; 85(8):570–1. [PubMed: 17768508]
6. Datta N, Kumar V. Validation of causes of infant death in the community by verbal autopsy. *Indian J Pediatr.* 1988; 55:599–604. [PubMed: 3169930]
7. Kalter H, Gray RH, Black RE, et al. Validation of post-mortem interviews to ascertain selected causes of death in children. *Int J Epidemiol.* 1990; 19:380–6. [PubMed: 2376451]
8. Gray, R.; Smith, G.; Barrs, P. The use of verbal autopsy methods to determine selected causes of death in children. Johns Hopkins University, School of Hygiene and Public Health, Institute of International Programs; 1990. Occasional paper no. 10
9. Snow RW, Marsh K. How useful are verbal autopsies to estimate childhood causes of death? *Health Policy Plan.* 1992; 7:22–9.
10. Marsh D, Majid N, Rasmussen Z, et al. Cause-specific child mortality in a mountainous community in Pakistan by verbal autopsy. *J Pak Med Assoc.* 1993; 43:226–9. [PubMed: 8114258]
11. Nykanen M, Tamoana W, Cullinan T, et al. Verbal autopsy as a technique to establish causes of infant and child mortality. *East Africa Med J.* 1995; 72:731–4.
12. Mobley CC, Boerma JT, Titus S, et al. Validation study of a verbal autopsy method for causes of childhood mortality in Namibia. *Journal of Tropical Pediatrics.* 1996; 42:365–9. [PubMed: 9009566]
13. Baqui AH, Black RE, Arifeen SE, et al. Causes of childhood deaths in Bangladesh: results of a nationwide verbal autopsy study. *Bull World Health Organ.* 1998; 76:161–71. [PubMed: 9648357]
14. Fantahun M. Patterns of childhood mortality in three districts of north Gondar Administrative Zone. A community based study using the verbal autopsy method. *Ethiopian Med J.* 1998; 36:71–81.
15. Edmond KM, Quigley MA, Zandoh C, et al. Diagnostic accuracy of verbal autopsies in ascertaining the causes of stillbirths and neonatal deaths in rural Ghana. *Paediatr Perinat Epidemiol.* 2008; 22(5):417. [PubMed: 18782250]
16. Setel PW, Rao C, Hemed Y, et al. Core verbal autopsy procedures with comparative validation results from two countries. *PLoS Med.* 2006; 3(8):e268. [PubMed: 16942391]
17. Anker M. The effect of misclassification error on reported cause-specific mortality fractions from verbal autopsy. *Int J Epidemiol.* 1997; 26(5):1090–6. [PubMed: 9363532]
18. Quigley MA, Chandramohan D, Rodrigues LC. Diagnostic accuracy of physician review, expert algorithms and data-derived algorithms in adult verbal autopsies. *Int J Epidemiol.* 1999; 28(6): 1081–7. [PubMed: 10661651]
19. Freeman JV, Christian P, Khatry SK, et al. Evaluation of neonatal verbal autopsy using physician review versus algorithm-based cause-of-death assignment in rural Nepal. *Paediatr Perinat Epidemiol.* 2005; 19(4):323–31. [PubMed: 15958155]
20. Reeves BC, Quigley M. A review of data-derived methods for assigning causes of death from verbal autopsy data. *Int J Epidemiol.* 1997; 26(5):1080–9. [PubMed: 9363531]
21. Quigley MA, Chandramohan D, Setel P, et al. Validity of data-derived algorithms for ascertaining causes of adult death in two African sites using verbal autopsy. *Trop Med Int Health.* 2000; 5(1): 33–9. [PubMed: 10672203]
22. Manandhar DS, Osrin D, Shrestha B, et al. Effect of a participatory intervention with women's groups on birth outcomes in Nepal. *Lancet.* 2004; 364:970–9. [PubMed: 15364188]
23. Sharma, NK. Dhanusha. In: Joshi, SR., editor. District profile of Nepal 2007/2008. 1st edition. Intensive Study & Research Centre; Kathmandu: Sep 16th. 2007 p. 346-58.
24. Baseline survey MIRA (Mother and Infant Research Activities) Dhanusha. 2006. (Unpublished)

25. Anker, M.; Black, RE.; Coldham, C., et al. A standard verbal autopsy method for investigating causes of death in infants and children. WHO, Johns Hopkins School of Hygiene and Public Health, London School of Hygiene and Tropical Medicine; Geneva: 1999. WHO/ CDS/CSR/ISR/ 99.4
26. Winbo IG, Serenius FH, Dahlquist GG, et al. NICE, a new cause of death classification for stillbirths and neonatal deaths. *Int J Epidemiol.* 1998; 27(3):499–504. [PubMed: 9698143]
27. Lawn JE, Wilczynska-Ketende K, Cousens SN. Estimating the causes of 4 million neonatal deaths in the year 2000. *Int J Epidemiol.* 2006; 35(3):706–18. [PubMed: 16556647]
28. Lawn J, Shibuya K, Stein C. No cry at birth: global estimates of intrapartum stillbirths and intrapartum-related neonatal deaths. *Bull World Health Organ.* 2005; 83:409–17. [PubMed: 15976891]
29. Save the Children; Nepal: Jul. 2002 State of the world's newborns: Nepal. Saving Newborn Lives.
30. Campbell O, Gipson R, Mohandes AE, et al. The Egypt national perinatal/neonatal mortality study 2000. *J Perinatol.* 2004; 24:284–9. [PubMed: 15042109]
31. Bhutta, ZA.; Zahid, AM.; Zaidi, S., et al. A verbal autopsy survey of Perinatal Mortality in rural Pakistan. *Child Health and Nutrition Research Initiative (CHNRI); Pakistan:* 2004.
32. Kulkarni R, Chauhan S, Shah B, et al. Investigating the causes of perinatal mortality by verbal autopsy in Maharashtra, India. *Indian J Commun Med.* 2007; 32:259–63.
33. Khanal, S.; Singh, V.; Dawson, P. Verbal autopsy to ascertain the causes of death in neonates at community setting. Presented paper in the 5th Conference of Perinatal Society of Nepal (PESON); Kathmandu, Nepal. 2-3rd Dec 2005; (Unpublished)
34. Lee AC, Mullany LC, Tielsch JM, et al. Verbal autopsy methods to ascertain birth asphyxia deaths in a community based setting in southern Nepal. *Pediatrics.* 2008; 121(5):e1372–80. [PubMed: 18450880]
35. Osrin D, Vergnano S, Costello A. Serious bacterial infections in newborn infants in developing countries. *Curr Opin Infect Dis.* 2004; 17:217–24. [PubMed: 15166824]
36. Lawn JE, Cousens S, Bhutta ZA, et al. Why are 4 million newborn babies dying each year? *Lancet.* 2004; 364:399–401. [PubMed: 15288723]
37. Lawn JE, Cousens S, Zupan J, et al. 4 million neonatal deaths: When? Where? Why? *Lancet.* 2005; 365:891–900. [PubMed: 15752534]
38. Ellis M, Manandhar N, Manandhar DS, et al. Risk factors for neonatal encephalopathy in Kathmandu, Nepal, a developing country: unmatched case-controlled study. *Br Med J.* 2000; 320:1229–36. [PubMed: 10797030]

Box 1

Classification of stillbirth and neonatal death

Criteria	Classification
Infant did not breathe, cry or move at birth, weighing >1000 g or >28 weeks	Stillbirth
Baby looked swollen, puffy face with skin peeling	Macerated stillbirth
Normal looking baby with skin intact, implying death <12 hours before delivery, weighing > 1000 g	Fresh stillbirth
Gross visible malformation, such as meningomyelocele, anencephaly, gastrochisis	Lethal congenital malformation
Prolonged labour >24 hrs, bleeding per vaginam, antepartum hemorrhage, postpartum hemorrhage, cord prolapse, cord around infant's neck, pregnancy induced hypertension, eclampsia, ruptured uterus, emergency caesarean section, malpresentation	Obstetric complications
Birth weight <2500 g and age <36 weeks, or pregnancy ended early and infant was smaller than usual at birth	Prematurity
History of chronic illness like Diabetes, Tuberculosis, Rheumatic Heart Disease, HIV, Epilepsy to mother	Maternal disease
Twin or triplet	Multiple pregnancy
External trauma to mother during pregnancy	Accident or external condition
Above conditions not fulfilled and other cause discernible from history	Other
Cause could not be ascertained due to lack of information	Unclassifiable
Infant breathed, cried or moved at birth, but died before 28 complete days	Neonatal death
Death within 4 hours of birth or gestational age > 36 weeks and not able to breathe normally at birth, or not able to suckle normally at birth, convulsions or spasms, but not tetanus or fever. *Supportive criteria: history of prolonged labour, malpresentation, twin, very large baby	Birth asphyxia
Fever or cold and limp or lethargic or convulsions or bulging fontanel and refusing to feed and no localizing signs Duration > 1 day and chest indrawing or difficulty breathing Umbilical area red or discharging pus or skin red, inflamed or hot or skin rash with pustules or pus and fever > 1 day Age 3-27 days and convulsions or spasms and able to suckle and cry normally after birth and stopped suckling or crying **Supportive criteria: use of dirty instruments to cut the cord, red and discharging umbilicus, not completed tetanus toxoid immunization course	Severe infection
Birth weight <2500 g and age <36 weeks, or pregnancy ended early and infant was smaller than usual at birth	Prematurity or low birth weight
Especially in winter months and history of having been bathed immediately after birth	Hypothermia
Gross visible malformation, such as meningomyelocele, anencephaly, gastrochisis	Congenital malformation
Bruises or marks of injury on body or head at birth	Accident or external condition
Above conditions not fulfilled and due to other reasons	Others
Cause could not be ascertained due to lack of information	Unclassifiable

Table 1
Births, stillbirths and neonatal deaths in 60 village development committees of Dhanusha district, 2006-2008

Category	Frequency
Births	25 982
Stillbirths	813
Live births	25 169
Neonatal deaths	954
Early neonatal deaths	760
Late neonatal deaths	194
Perinatal deaths (stillbirths and early neonatal deaths)	1 573
Stillbirths classified on the basis of verbal autopsy	601
Neonatal deaths classified on the basis of verbal autopsy	671
Indicator	Rate
Stillbirth Rate per 1000 births	31
Neonatal Mortality Rate per 1000 live births	38
Early Neonatal Mortality Rate per 1000 live births	30
Late Neonatal Mortality Rate per 1000 live births	8
Perinatal Mortality Rate per 1000 births	60

Table 2
Causes of stillbirth and neonatal death in 60 village development committees of Dhanusha district, 2006-2008

Stillbirth	Frequency	(%)
Obstetric complications	403	(67)
Prematurity	63	(10)
Maternal specific conditions	41	(7)
Congenital malformations	16	(3)
Multiple pregnancy	10	(2)
Others	9	(1)
Unclassifiable	59	(10)
Total	601	(100)
Neonatal death		
Birth asphyxia	250	(37)
Severe infection	205	(30)
Prematurity or low birth weight	101	(15)
Hypothermia	45	(7)
Congenital malformations	6	(1)
Others	33	(5)
Unclassifiable	31	(5)
Total	671	100

Table 3

Characteristics of delivery for 1272 stillbirths and neonatal deaths in 60 village development committees of Dhanusha district, 2006-2008

Place of delivery	Frequency	(%)
Home	826	(65)
Hospital	254	(20)
Other health institution (primary health centre, private clinic)	145	(12)
On the way to hospital	30	(2)
Other	17	(1)
Total	1272	(100)
Delivery conducted by		
Relative	360	(28)
Doctor	341	(27)
Village practitioner ¹	241	(19)
Health worker ²	161	(13)
Volunteer ³	105	(8)
Other (friend, neighbor)	14	(1)
Alone	50	(4)
Total	1272	(100)
History of Injection given to mother		
Yes	628	(50)
No	640	(50)
Total	1268	(100)
Umbilical cord cutting instrument		
Unboiled instrument	345	(53)
Boiled blade	118	(18)
Other	10	(1)
Unable to recall	180	(28)
Total	653	(100)
Umbilical stump application		

Place of delivery	Frequency	(%)
Nothing applied	240	(37)
Others (dung or ash)	134	(21)
Oil	28	(4)
Antiseptic (Dettol)	23	(3)
Unable to recall	228	(35)
Total	653	(100)
Infant's first feed		
No prelacteal feeds	203	(30)
Goat's milk	155	(23)
Prelacteal (Honey, sugar, clarified butter)	112	(17)
Mother's breast milk	80	(12)
Other mother's breast milk	72	(11)
Cow or buffalo milk	9	(1)
Infant formula	3	(<1)
Unable to recall	37	(6)
Total	671	(100)

¹ Untrained health worker who conducts delivery in the community

² Health assistant, staff nurse, auxiliary health worker, community medical assistant, auxiliary nurse midwife, maternal and child health worker

³ FCHV: Female community health volunteer; TBA: Trained birth attendant

Table 4

Asphyxial conditions and resuscitation at delivery, for 671 neonatal deaths in 60 village development committees of Dhanusha district, 2006-2008

	Frequency	(%)
Infant cried immediately after birth		
Yes	395	(59)
No	276	(41)
Total	671	(100)
Time until infant cried, if not immediately		
Within 5 min	22	(8)
5 – 30 min	67	(25)
> 30 min	44	(17)
Never cried	132	(50)
Total	265	(100)
Resuscitation measures		
Mouth-to-mouth respiration	111	(45)
Tactile stimulation	77	(31)
Mouth to tube-and-mask ventilation	47	(19)
Bag-and-mask ventilation	13	(5)
Total	248	(100)

Table 5
Treatment modalities for 646 neonatal deaths in 60 village development committees of Dhanusha district, 2006 – 2008

	Frequency	(%)
Treatment received		
Yes	281	(44)
No	365	(56)
Total	646	(100)
Treatment given at:		
Hospital	106	(38)
Other health institute (private clinic, nursing home)	63	(23)
Home	75	(27)
Traditional healers (jhankri, herbalists)	35	(12)
Total	279	(100)

Table 6

Comparison of characteristics of 17 377 infants who survived and 1272 infants who were stillborn or died in the neonatal period, in 60 village development committees of Dhanusha district, 2006-2008

	Infants who survived 28 days n=17 377		Stillbirths and perinatal deaths n=1272	
	Frequency	(%)	Frequency	(%)
Infant sex				
Male	9147	(53)	713	(56)
Female	8230	(47)	559	(44)
Total	17377	(100)	1272	(100)
Maturity				
Term	17110	(99)	970	(76)
Preterm	181	(1)	302	(24)
Total	17291	(100)	1272	(100)
Place of delivery				
Home	14100	(82)	826	(65)
Hospital	2324	(14)	254	(20)
Other health Institution	602	(4)	145	(12)
On the way to hospital	107	(<1)	30	(2)
Other (including vehicle delivery)	101	(<1)	17	(1)
Total	17234	(100)	1272	(100)
Birth attendant				
Relative	5607	(34)	360	(28)
Doctor	1623	(10)	341	(27)
Village practitioner	978	(6)	241	(19)
Health worker	2104	(13)	161	(13)
Volunteer	5159	(31)	105	(8)
No attendant	500	(3)	50	(4)
Others (friends, neighbors)	460	(3)	14	(1)
Total	16431	(100)	1272	(100)
Umbilical cord cutting implement				
Unboiled instrument	10614	(66)	345	(53)
Boiled blade	4687	(29)	118	(18)
Other	54	(<1)	10	(1)

	Infants who survived 28 days n=17,377		Stillbirths and perinatal deaths n=1272	
	Frequency	(%)	Frequency	(%)
Not recalled	660	(4)	180	(28)
Total	16015	(100)	653	(100)
Infant cried immediately	9968	(86)	395	(59)
Yes	1627	(14)	276	(41)
No	11595	(100)	671	(100)
Umbilical cord application	5508	(35)	240	(37)
Applied nothing	1986	(13)	28	(4)
Oil	1435	(9)	23	(3)
Antiseptic (Dettol)	5637	(36)	134	(21)
Other (dung, ash)	1163	(7)	228	(35)
Not recalled	15729	(100)	653	(100)
First feed	0	(0)	203	(30)
No prelacteal	5255	(31)	155	(23)
Goat's milk	4175	(24)	80	(12)
Mother's breast milk	4123	(24)	72	(11)
Other mother's breast milk	3042	(18)	112	(17)
Prelacteal (Honey, sugar, clarified butter)	282	(2)	9	(1)
Cow, buffalo milk	181	(1)	3	(<1)
Infant formula	60	(<1)	37	(6)
Not recalled	17118	(100)	671	(100)
Treatment given	8869	(54)	365	(56)
Yes	7568	(46)	281	(44)
No	16437	(100)	646	(100)
Total				